



Bacterial Microcompartment-Dependent 1,2-Propanediol Utilization of *Propionibacterium freudenreichii*

Alexander Dank^{1†}, Zhe Zeng^{1†}, Sjef Boeren², Richard A. Notebaart¹, Eddy J. Smid^{1*} and Tjakko Abee^{1*}

¹ Food Microbiology, Wageningen University and Research, Wageningen, Netherlands, ² Laboratory of Biochemistry, Wageningen University and Research, Wageningen, Netherlands

OPEN ACCESS

Edited by:

Ulrike Kappler,
The University of Queensland,
Australia

Reviewed by:

Chiranjit Chowdhury,
National Chemical Laboratory (CSIR),
India
Bhawani Chamlagain,
University of Helsinki, Finland

*Correspondence:

Tjakko Abee
tjakko.abee@wur.nl
Eddy J. Smid
eddy.smid@wur.nl

[†] These authors have contributed
equally to this work and share first
authorship

Specialty section:

This article was submitted to
Microbial Physiology and Metabolism,
a section of the journal
Frontiers in Microbiology

Received: 12 March 2021

Accepted: 16 April 2021

Published: 12 May 2021

Citation:

Dank A, Zeng Z, Boeren S,
Notebaart RA, Smid EJ and Abee T
(2021) Bacterial
Microcompartment-Dependent
1,2-Propanediol Utilization of
Propionibacterium freudenreichii.
Front. Microbiol. 12:679827.
doi: 10.3389/fmicb.2021.679827

Bacterial microcompartments (BMCs) are proteinaceous prokaryotic organelles that enable the utilization of substrates such as 1,2-propanediol and ethanolamine. BMCs are mostly linked to the survival of particular pathogenic bacteria by providing a growth advantage through utilization of 1,2-propanediol and ethanolamine which are abundantly present in the human gut. Although a 1,2-propanediol utilization cluster was found in the probiotic bacterium *Propionibacterium freudenreichii*, BMC-mediated metabolism of 1,2-propanediol has not been demonstrated experimentally in *P. freudenreichii*. In this study we show that *P. freudenreichii* DSM 20271 metabolizes 1,2-propanediol in anaerobic conditions to propionate and 1-propanol. Furthermore, 1,2-propanediol induced the formation of BMCs, which were visualized by transmission electron microscopy and resembled BMCs found in other bacteria. Proteomic analysis of 1,2-propanediol grown cells compared to L-lactate grown cells showed significant upregulation of proteins involved in propanediol-utilization (*pdu*-cluster), DNA repair mechanisms and BMC shell proteins while proteins involved in oxidative phosphorylation were down-regulated. 1,2-Propanediol utilizing cells actively produced vitamin B₁₂ (cobalamin) in similar amounts as cells growing on L-lactate. The ability to metabolize 1,2-propanediol may have implications for human gut colonization and modulation, and can potentially aid in delivering propionate and vitamin B₁₂ *in situ*.

Keywords: bacterial microcompartments, *Propionibacterium freudenreichii*, vitamin B₁₂, metabolosome, propanediol utilization cluster, probiotic, intestine

INTRODUCTION

Propionibacterium freudenreichii is a Gram-positive, non-spore forming bacterium which has been linked to several potential health promoting effects, such as reducing intestinal inflammation, immunomodulation, modulation of intestinal motility and absorption, reduction of pathogen adhesion and enhancement of bifidobacteria [reviewed by Cousin et al. (2011)]. *P. freudenreichii* is able to cope with a large variety of stresses (oxidative, bile salt, and temperature; Falentin et al., 2010) and consequently has good survival capabilities in the upper intestinal tract (Huang and Adams, 2004). Survival of *P. freudenreichii* in the intestinal environment is supported by expression

of pathways involved in the metabolism of substrates present in the intestinal environment, such as propanediol (Saraoui et al., 2013).

1,2-propanediol is a major end product from anaerobic degradation of rhamnose or fucose by the human intestinal microbiota and serves as an important carbon source for 1,2-propanediol utilizing bacteria, which can metabolize 1,2-propanediol into propionate, generating ATP, and into 1-propanol for maintaining redox balance. The metabolism of 1,2-propanediol produces the toxic intermediate propionaldehyde (Sampson and Bobik, 2008). Some bacteria can protect themselves from toxic intermediates by encapsulating the enzymatic processes in self-assembling proteinaceous organelles called bacterial microcompartments (BMCs; Axen et al., 2014). BMCs are typically about 40–200 nm in diameter and are made of three types of shell proteins: hexamers, pseudo-hexamers, and pentamers (Kerfeld et al., 2018). Based on the highly conserved domain of shell proteins, Axen et al. (2014) predicted the presence of these organelles in 23 different bacterial phyla, including *Actinobacteria* which includes the species *P. freudenreichii*. However, no experimental evidence of BMC mediated utilization of 1,2-propanediol has been shown in *P. freudenreichii* and its role as carbon source is yet to be elucidated.

Bacterial microcompartment-mediated catabolism of substrates involving a toxic aldehyde intermediate is driven by three core enzymes: aldehyde dehydrogenase, alcohol dehydrogenase, and phosphotransacylase (Axen et al., 2014). The signature enzyme in the propanediol utilization (*pdu*) pathway is propanediol dehydratase, a vitamin B₁₂-dependent enzyme catalyzing the reaction of 1,2-propanediol to propionaldehyde (Cheng et al., 2011; Axen et al., 2014). Vitamin B₁₂ is produced in high quantities by *P. freudenreichii* (Burgess et al., 2009) and it also requires vitamin B₁₂ as a cofactor for a key enzyme in its characteristic Wood–Werkman cycle. It is suggested that *Actinobacteria* such as *P. freudenreichii* obtained the *pdu* cluster through horizontal gene transfer from *Clostridiales* (Ravcheev et al., 2019), in which acquisition may have been supported by the ability of *P. freudenreichii* to produce vitamin B₁₂ *de-novo* (Roessner et al., 2002). BMCs are also found in some pathogenic bacteria, such as *Salmonella enterica*, *Enterococcus faecalis*, *Listeria monocytogenes*, pathogenic *Escherichia coli*, and *Clostridium perfringens* (Kerfeld et al., 2018). BMC-mediated utilization of 1,2-propanediol increases competitive fitness of pathogens in the gut and consequently has been linked to virulence (Jakobson and Tullman-Ercek, 2016). However, symbiotic relationships depending on 1,2-propanediol metabolism have also been shown for beneficial *Lactobacillus reuteri* and *Bifidobacterium breve* (Cheng et al., 2020). The ability to degrade 1,2-propanediol may have similar implications for the bifidogenic capacity reported for *P. freudenreichii* (Kaneko et al., 1994) and consequences for gut modulation and competition with pathogenic bacteria. Furthermore, the active production of vitamin B₁₂ during metabolism of 1,2-propanediol has not been studied yet in *P. freudenreichii*.

In this study we present evidence for BMC-mediated anaerobic growth of *P. freudenreichii* on 1,2-propanediol,

evidenced by substrate utilization, propionate and 1-propanol production, and vitamin B₁₂ synthesis. Using transmission electron microscopy (TEM) and proteomics, we confirmed the presence of BMCs, Pdu BMC shell proteins and enzymes in *pdu*-induced *P. freudenreichii*.

MATERIALS AND METHODS

Strains, Culture Conditions, and Growth Measurement

Propionibacterium freudenreichii DSM 20271 was grown anaerobically (Anoxomat modified atmosphere, MART; 10% CO₂, 5% H₂, and 85% N₂) at 30°C in 50 mL tubes containing 40 mL complex media containing per liter: 10 g tryptone, 5 g yeast extract, and 5 g KH₂PO₄ supplemented with 100 mM L-lactate or 1,2-propanediol. All media was set at pH 7.0 by addition of 5 M NaOH and was filter sterilized through 0.2 μm filters into sterile flasks. OD₆₀₀ measurements and extracellular metabolite samples were taken daily for a time period of 7 days.

Analysis of Extracellular Metabolites Using High Performance Liquid Chromatography

Culture samples were taken at various time intervals and were analyzed for extracellular metabolites by high performance liquid chromatography. 1 mL culture was centrifuged at 17,000 × g for 1 min and the supernatant was collected. 0.5 mL supernatant was treated with 0.25 mL Carrez A and 0.25 mL Carrez B, vortexed and centrifuged at 17,000 × g for 2 min. 200 μL supernatant was stored in HPLC vials at –20°C upon analysis. HPLC was performed as described by Zeng et al. (2019). Quantification was performed by addition of a standard curve containing L-lactate, acetate, propionate, 1,2-propanediol, and 1-propanol.

Transmission Electron Microscopy

Propionibacterium freudenreichii cultures were grown anaerobically at 30°C in 100 mM L-lactate or 1,2-propanediol containing media. Samples were collected after 6 days of incubation (early stationary phase for 1,2-propanediol-grown cells). About 10 μg of dry cell biomass was fixed for 2 h in 2.5% (v/v) glutaraldehyde in 0.1 M sodium cacodylate buffer (pH 7.2). After rinsing in the same buffer, a post-fixation was done in 1% (w/v) OsO₄ for 1 h at room temperature. The samples were dehydrated using ethanol. The dehydrated cell pellets were then embedded in resin (Spurr HM20) for 10 h at 70°C. Thin sections (<100 nm) of polymerized resin samples were obtained with microtomes. After staining with 2% (w/v) aqueous uranyl acetate, the samples were analyzed with a Jeol 1400 plus TEM with 120 kV setting as described by Zeng et al. (2019).

Vitamin B₁₂ Quantification

After 7 days of incubation the vitamin B₁₂ (cobalamin) concentration was determined using a microbiological assay (Vitafast vitamin B₁₂ kit, R-biopharm) for *P. freudenreichii* grown in 66 mM L-lactate and 49 mM 1,2-propanediol medium.

Briefly, 1 mL of culture was disrupted by bead beating (lysing matrix B, mp-bio) 2×1 min at 4.5 m/s with 1 min on ice in between. Samples were centrifuged and diluted with water to appropriate concentrations for the test kit and were heat-extracted for 30 min at 95°C. The samples were cooled to room temperature and vitamin B₁₂ detection was performed according to the manufacturer protocol.

Proteomics

Propionibacterium freudenreichii cells were cultured in media supplemented with 100 mM L-lactate or 100 mM 1,2-propanediol for 7 days. Cell pellets were harvested by centrifugation of 1 mL of sample at $17,000 \times g$ for 1 min in table top centrifuges and cell pellets were frozen at -80 degrees. Samples were washed twice with 100 mM Tris (pH 8) and resuspended in 100 μ l 100 mM Tris. Samples were lysed by sonication for 45 s twice while cooling 1 min on ice. Protein content was determined using Pierce Coomassie protein assay and samples were diluted to 1 μ g/ μ l using Tris-HCl pH 8. Samples were prepared according to the filter assisted sample preparation protocol (FASP; Wiśniewski et al., 2009) with the following steps: reduction with 15 mM dithiothreitol, alkylation with 20 mM acrylamide, and digestion with sequencing grade trypsin overnight. Each prepared peptide sample was analyzed by injecting (18 μ l) into a nanoLC-MS/MS (Thermo nLC1000 connected to a LTQ-Orbitrap XL) as described previously (Lu et al., 2011). LCMS data with all MS/MS spectra were analyzed with the MaxQuant quantitative proteomics software package (Cox et al., 2014) as described before (Smaczniak et al., 2012; Wendrich et al., 2017).

A protein database with the protein sequence of *P. freudenreichii* DSM 20271 (ID:UP000032238) was downloaded from UniProt. Filtering and further bioinformatics and statistical analysis of the MaxQuant ProteinGroups file were performed with Perseus (Tyanova et al., 2016). Reverse hits and contaminants were filtered out. Protein groups were filtered to contain minimally two peptides for protein identification of

which at least one is unique and at least one is unmodified. Also, each group required three valid values in at least one of the two experimental groups. A volcano plot was prepared based on the Student's *t*-test difference between samples. Volcano plots were produced in Rstudio using EnhancedVolcano (Blighe, 2018). Proteins were considered to be significantly different amongst sample if $p < 0.05$ and 4-fold change difference was detected. KEGG gene set enrichment analysis was performed using ClusterProfiler (Yu et al., 2012) and 2-fold change difference amongst proteins. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Vizcaíno et al., 2016) partner repository with the dataset identifier PXD024700.


Predicting BMCs Shell Proteins

The Hidden Markov Models (HMMs) of two BMC shell protein domains listed as Pf00936 and Pf03319 were retrieved from the Pfam database to predict BMC shell proteins as described in Axen et al., 2014; Zeng et al., 2019. Shell proteins were predicted by a HMM search using the HMMER package and a local protein database of *P. freudenreichii* DSM 20271 genome [CP010341.1 (Deptula et al., 2017)]. All hits with an *e*-value less than or equal to 1e-05 that correspond to a genomic record from Genbank, RefSeq, EMBL, or DDBJ databases were accepted as BMC shell protein homologs (Supplementary File 1).

RESULTS

Growth Performance and Metabolite Production on 1,2-Propanediol and L-Lactate

Anaerobic growth of *P. freudenreichii* on 100 mM 1,2-propanediol was monitored for 6 days with daily sampling in triplicate. As a control, *P. freudenreichii* was cultured on




100 mM L-lactate. *P. freudenreichii* was able to completely metabolize 100 mM 1,2-propanediol during a period of 6 days (see **Figure 1**). This resulted in the production of 37.5 ± 0.1 mM 1-propanol and 43.2 ± 0.5 mM propionate. No production of acetate was observed, and with the current HPLC method used, approximately 28 mM of expected C₃ compounds is missing, conceivably volatile propionaldehyde. In the control cultures, 100 mM L-lactate was consumed within 2 days, resulting in 65.1 ± 0.1 mM propionate and 30.5 ± 0.1 mM acetate, close the expected molar ratio of 2:1 (Seeliger et al., 2002). Growth on 1,2-propanediol resulted in slower growth, with a stationary phase after 120 h vs 48 h in L-lactate. Our results clearly demonstrate that *P. freudenreichii* can grow on 1,2-propanediol, metabolizing it to propionate and 1-propanol.

To monitor metabolism of 1,2-propanediol in the presence of other carbon sources *P. freudenreichii* was grown in media containing both L-lactate and 1,2-propanediol. When NAD⁺/NADH pools are shared between the cytosol and the BMC as discussed by Ferlez et al. (2019), the highest ATP yield from 1,2-propanediol in the presence of L-lactate can be obtained by co-fermenting 1,2-propanediol and L-lactate solely to propionate (see **Supplementary Text 1**). The expected ATP yield with L-lactate and 1,2-propanediol as mixed substrates is higher compared to cells growing solely on L-lactate (see **Supplementary Text 1**). We found production of 67.1 ± 1.5 mM propionate, 19.0 ± 1.0 mM of acetate, and 28.3 ± 2.8 mM

1-propanol in cells growing on the mixed substrates. Assuming 1-propanol:propionate is produced in a 1:1 ratio from 1,2-propanediol and acetate:propionate is produced in a 1:2 ratio from L-lactate, 28 mM of the propionate originates from the 1,2-propanediol metabolism and 38 mM propionate originates from L-lactate metabolism. This matches the total found propionate of 67.1 ± 1.5 mM propionate in the samples. In mixed substrate conditions the amounts of total products formed indicate no apparent loss of C₃ compounds, which may indicate reduced loss of volatile propionaldehyde. Biomass formation with 1:1 L-lactate:1,2-propanediol-grown cells was found to be lower compared to 100 mM L-lactate-grown cells. These results indicate no apparent energetic benefit from mixed substrate conditions compared to mono substrate conditions, pointing toward independent pathways. Our results show mixed-substrate metabolism influences the total amount of short-chain fatty acid production, but pathway interactions are not apparent.

Vitamin B₁₂ Production

To demonstrate active vitamin B₁₂ production under 1,2-propanediol utilizing conditions, *P. freudenreichii* was grown for 7 days in either yeast extract medium supplemented with 66 mM L-lactate or 49 mM 1,2-propanediol. Biomass formation, substrate utilization, and vitamin B₁₂ production by *P. freudenreichii* was monitored after 7 days of incubation.



Incubation vessels were not opened in-between to prevent side effects by oxygen-dependent stimulation of vitamin B₁₂ production, as has been shown for *P. freudenreichii* (Quesada-Chanto et al., 1998). Again, complete utilization of L-lactate and 1,2-propanediol was found (**Supplementary File 2**).

Vitamin B₁₂ was produced both when grown on L-lactate as on 1,2-propanediol ($1.84 \pm 0.11 \mu\text{g}/\text{mmol}$ L-lactate vs $2.19 \pm 0.03 \mu\text{g}/\text{mmol}$ 1,2-propanediol). The specific vitamin B₁₂ production when grown on L-lactate and on 1,2-propanediol was $270 \pm 33 \mu\text{g}/\text{g}$ cells and $288 \pm 27 \mu\text{g}/\text{g}$ cells, respectively, (**Figure 2**). No significant difference was thus found for the production of vitamin B₁₂ per g cells when growing on either L-lactate supplemented media or 1,2-propanediol supplemented media. In complex media, 1,2-propanediol as carbon source thus supports vitamin B₁₂ production by *P. freudenreichii*.

Proteomics and Electron Microscopy of L-Lactate and 1,2-Propanediol Grown Cells


Cells growing in media supplemented with L-lactate (control) and with 1,2-propanediol were visualized using TEM. Thin sections of cells grown on 1,2-propanediol supplemented media clearly display cellular structures which were not found in the cells grown on L-lactate supplemented media (see **Figure 3**), and those structures resemble BMC structures found in other bacteria including *L. monocytogenes* (Zeng et al., 2019), *S. enterica* (Crowley et al., 2008), and *E. coli* (Toraya et al., 1979).

Proteome analysis revealed that BMC structural shell proteins (PduA, PduB, PduK, PduJ, PduN, and PduM) and enzymes involved in 1,2-propanediol utilization (PduL, PduC, PduD, PduE, PduP, PduO, and PduQ) were significantly more abundant in cells grown with 1,2-propanediol as a substrate compared

to cells grown on L-lactate (**Figure 4**). Lactate permease, succinate dehydrogenase subunits A, C1 (SdhA, SdhC1) were found to be significantly more abundant in the L-lactate-grown cells compared to 1,2-propanediol-grown cells, indicating a decrease of proteins in lactate-degradation pathways such as the Wood-Werkman cycle. Gene set enrichment analysis of KEGG pathways (see **Supplementary File 3**) revealed upregulation of the propanediol degradation pathway in 1,2-propanediol grown cells compared to L-lactate grown cells (adjusted *p*-value < 0.10). DNA repair mechanisms were also found to be activated in 1,2-propanediol-grown cells (adjusted *p*-value < 0.10). In 1,2-propanediol-grown cells significant suppression of oxidative phosphorylation was found compared to L-lactate grown cells (adjusted *p*-value < 0.01). The measured consumption of 1,2-propanediol, production of 1-propanol and propionate, visualized cellular structures, and strong upregulation of the *pdu* loci proteins provide evidence for activation of BMC-dependent 1,2-*pdu* in *P. freudenreichii* DSM 20271.

Overview of BMC-Dependent 1,2-Propanediol Metabolism Model in *P. freudenreichii*

Subsequent analysis of the *pdu* cluster in *P. freudenreichii* DSM 20271 identified two distant loci, with locus 1 starting from RM25_0852 to RM25_0857 and locus 2 starting from RM25_1258 to RM25_1273 (**Figure 5A**). Locus 1 contains four genes which are *pocR* encoding a transcriptional regulator, *pduQ* encoding 1-propanol dehydrogenase, *pduV* with unknown function and *pduU* encoding BMC shell protein. Locus 2 carries 14 genes including 6 genes encoding BMC shell proteins and 8 genes encoding enzymes for the 1,2-propanediol




degradation pathway and is not preceded by any known transcriptional regulators.

Based on the structural studies of BMCs (Sutter et al., 2017; Greening and Lithgow, 2020) and our understanding of 1,2-propanediol metabolism (Sampson and Bobik, 2008; Kerfeld et al., 2018; Zeng et al., 2019), we propose the model of BMC-dependent 1,2-propanediol metabolism in *P. freudenreichii*, with the predicted BMC shell proteins PduA, PduB, PduK, PduJ, PduM, PduN, and PduU constituting the self-organized icosahedral organelle (Kerfeld et al., 2018; Greening and Lithgow, 2020). As illustrated in **Figure 5B**, the catabolism of 1,2-propanediol starts with the conversion of 1,2-propanediol to propionaldehyde by vitamin B₁₂-dependent diol dehydratase PduCDE. The toxic propionaldehyde is then converted to propionate by the enzyme CoA-dependent propionaldehyde dehydrogenase PduP, followed by action of phosphate propanoyltransferase PduL, and potential propionate kinase located in the cytoplasm, resulting in the end product propionate and the production of ATP. The other end product is produced following conversion of propionaldehyde by propanol dehydrogenase PduQ into

1-propanol. The diol dehydratase reactivase PduGH and corrinoid adenosyltransferase PduO are linked to the supply and recycling of vitamin B₁₂.

DISCUSSION


Propionibacterium freudenreichii is commonly found in the rumen and colon of animals and in the human intestine (Bryant, 1959). In these environments fucose and rhamnose are degraded to 1,2-propanediol by the present microbiota (Xue et al., 2008). We showed 1,2-propanediol can be further metabolized into propionate and 1-propanol by *P. freudenreichii*, thereby supporting anaerobic growth. Previously, a locus containing 15 genes involved in *pdu* was detected in *P. freudenreichii* (Falentin et al., 2010). *In vivo* gene expression analysis showed the *pdu* operon to be expressed in *P. freudenreichii* cells contained in the colon environment of a pig (Saraoui et al., 2013), pointing toward *pdu* in intestinal environments. Interestingly, in this study we identified the *pdu* cluster distributed in two different loci in



P. freudenreichii DSM 20271. The presence of the *pdu* cluster seems to be species specific in propionic acid bacteria, as blasting the *pdu* cluster of *P. freudenreichii* against *Cutibacterium acnes*, *Acidipropionibacterium acidipropionici*, *A. thoenii*, and *A. jensenii* did not result significant hits for key components in the cluster (see **Supplementary File 4**). Presence of 1,2-propanediol in the medium induced expression of the two loci in *P. freudenreichii* DSM 20271 and resulted in BMC formation, 1,2-propanediol metabolism and consequently propionate and 1-propanol production. Our results show that expression of the *pdu* cluster from two different loci results in effective BMC formation and 1,2-propanediol metabolism. Indeed, BMC genes are split into two or more loci in 40% of the prokaryotic genomes containing BMCs (Abdul-Rahman et al., 2013). In *P. freudenreichii* DSM 20271, locus 1 is preceded by transcriptional activator *pocR*, which has been linked to activation of the *pdu* cluster in *Salmonella* (Chen et al., 1994). We did not find any annotated transcriptional regulator in the vicinity of locus 2. The presence of two different loci in many prokaryotic genomes suggests expression is controlled by additional regulators next to PocR. Indeed

heterologous expression of BMCs uncoupled from their cognate transcriptional regulators has been reported previously (Wilson, 2021). The transcriptional regulation and activation of the two *pdu* loci and the role of *pocR* in *P. freudenreichii* requires further attention.

Upregulation of the *pdu* cluster and DNA repair mechanisms clearly indicated the crucial role of BMCs to protect *P. freudenreichii* from the toxic intermediate propionaldehyde produced in the degradation pathway. The ability to utilize substrates producing toxic intermediates upon degradation results in a competitive advantage to other gut microbiota (Jakobson and Tullman-Ercek, 2016), as shown for ethanolamine utilization by *S. enterica* during intestinal inflammation (Thiennimitr et al., 2011). Interestingly, the presence of genes encoding metabolosomes for the utilization of ethanolamine and propanediol has been linked to pathogenicity and aids in anaerobic growth and colonization of foodborne pathogens *L. monocytogenes*, *C. perfringens*, and *S. typhimurium* (Korbel et al., 2005; De Weirdt et al., 2012). It has been suggested for beneficial bacterium *L. reuteri* that competition for 1,2-propanediol could result in decreased proliferation of pathogens



(Cheng et al., 2020). *P. freudenreichii* is considered to be non-pathogenic and has the generally recognized as safe (GRAS) status (Meile et al., 2008). Substrate competition for 1,2-propanediol by *P. freudenreichii* may thus exert similar effects as suggested for *L. reuteri* on growth of pathogenic bacteria. This is in line with reports that *P. freudenreichii* can decrease adhesion of pathogens to human intestinal mucus cells (Collado et al., 2007). Furthermore, *P. freudenreichii* stimulates the growth of beneficial bifidobacteria (Satomi et al., 1999; Hojo et al., 2002) thereby promoting a healthy gut microbiota. The role of BMC-mediated 1,2-propanediol utilization by *P. freudenreichii* and its importance for modulating gut microbiota composition, both by substrate competition and promoting other beneficial microbiota, requires further investigation.

Our study shows the potential of *P. freudenreichii* to substantially contribute to the production of propionate in the human gut. Next to 1,2-propanediol, also lactic acid (Macfarlane and Gibson, 1997) is a major end-product of microbial fermentation, which consequently can also be fermented by *P. freudenreichii* and additionally contributes to the production of propionate. In mixed substrate conditions both pathways remained active, albeit without apparent interaction based on metabolite formation. However, mixed substrate conditions decreased the loss of C₃, suggesting less loss of volatile propionaldehyde, thus potentially more efficient BMC assembly. The assembly of BMCs in mixed substrate condition requires further attention. Propionate is linked to many putative health effects [reviewed by (Hosseini et al., 2011)] and can further stimulate bifidobacteria (Kaneko et al., 1994). The production of propionate from 1,2-propanediol further supports the potential of *P. freudenreichii* as a probiotic.

The metabolism of 1,2-propanediol requires propanediol dehydratase, which is vitamin B₁₂-dependant (Axen et al., 2014). Vitamin B₁₂ is actively produced by *P. freudenreichii* as it is essential cofactor in a key enzyme in the Wood-Werkman cycle, methylmalonyl-CoA mutase. Synthesis of vitamin B₁₂ in *P. freudenreichii* follows the anaerobic pathway (Roessner et al., 2002), enabling production of vitamin B₁₂ for anaerobic metabolism of lactate. Here we show vitamin B₁₂ is produced in similar amounts in complex medium when cells metabolize 1,2-propanediol or L-lactate as carbon source. In *S. typhimurium* *pocR* mediated expression of vitamin B₁₂ is induced in the presence of propanediol (Richter-Dahlfors et al., 1994). We also identified *pocR* upstream of *pdu* loci 1 in *P. freudenreichii* DSM 20271, but as discussed before its exact regulatory role in *P. freudenreichii* remains to be elucidated. Vitamin B₁₂ production is also regulated by a vitamin B₁₂ regulated riboswitch in *P. freudenreichii*. In *Propionibacterium* strain UT1 expression of vitamin B₁₂ biosynthesis occurred at vitamin B₁₂ concentrations of 750 μM, much higher compared to the vitamin B₁₂ concentrations found in this study (~90 nM). The role of this riboswitch for vitamin B₁₂ production during metabolism of 1,2-propanediol also remains to be elucidated. Based

on our findings, we hypothesize that *P. freudenreichii* occupies a lactate and propanediol-rich niche in the gut environment. Symbiotic relationships have been shown for the production of vitamin B₁₂ (Belzer et al., 2017; Sokolovskaya et al., 2020), which may be hard-wired in the vitamin B₁₂ production of *P. freudenreichii*. As next to vitamin B₁₂-dependent lactate metabolism, BMC-mediated 1,2-propanediol metabolism supports anaerobic growth of *P. freudenreichii*, thereby contributing to *in situ* vitamin B₁₂ production in the gut.

This study presents evidence for BMC-mediated vitamin B₁₂-dependent utilization of 1,2-propanediol in *P. freudenreichii*. We have shown that 1,2-propanediol supports anaerobic growth of *P. freudenreichii*. It is conceivable that utilization of 1,2-propanediol could aid colonization of *Propionibacterium freudenreichii* in the human gut to exert beneficial effects, such as delivering vitamin B₁₂ and propionate *in situ*, but these aspects require further study.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are publicly available. This data can be found here: <https://www.ebi.ac.uk/pride/archive/projects/PXD024700>.

AUTHOR CONTRIBUTIONS

AD and ZZ designed and performed the experiments. SB, AD, and ZZ performed proteomics and analyzed data. ZZ, AD, ES, RN, and TA analyzed data. ZZ, AD, ES, and TA wrote the manuscript. All authors read, edited and approved the manuscript.

FUNDING

The authors declare that this study received funding from Arla Foods (Denmark). The funder was not involved in the study design, collection, analysis, interpretation of data, the writing of this article or the decision to submit it for publication. ZZ was supported by a grant from the China Scholarship Council.

ACKNOWLEDGMENTS

We would like to thank Wageningen Electron Microscopy Centre (WEMC) for the use of their transmission electron microscope and for their help in sample preparation.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2021.679827/full#supplementary-material>

REFERENCES

- Abdul-Rahman, F., Petit, E., and Blanchard, J. L. (2013). The distribution of polyhedral bacterial microcompartments suggests frequent horizontal transfer and operon reassembly. *J. Phylog. Evol. Biol.* 1:118.
- Axen, S. D., Erbilgin, O., and Kerfeld, C. A. (2014). A taxonomy of bacterial microcompartment loci constructed by a novel scoring method. *PLoS Comput. Biol.* 10:e1003898. doi: 10.1371/journal.pcbi.1003898
- Belzer, C., Chia, L. W., Aalvink, S., Chamlagain, B., Piironen, V., Knol, J., et al. (2017). Microbial metabolic networks at the mucus layer lead to diet-independent butyrate and Vitamin B12 production by intestinal symbionts. *mBio* 8:e00770–17. doi: 10.1128/mBio.00770-17
- Blighe, K. (2018). *EnhancedVolcano: Publication-Ready Volcano Plots With Enhanced Colouring and Labeling*. Vienna: R Foundation for Statistical Computing.
- Bryant, M. P. (1959). Bacterial species of the rumen. *Bacteriol. Rev.* 23, 125–153. doi: 10.1128/br.23.3.125-153.1959
- Burgess, C. M., Smid, E. J., and van Sinderen, D. (2009). Bacterial vitamin B2, B11 and B12 overproduction: an overview. *Int. J. Food Microbiol.* 133, 1–7. doi: 10.1016/j.ijfoodmicro.2009.04.012
- Chen, P., Andersson, D. I., and Roth, J. R. (1994). The control region of the pdu/cob regulon in *Salmonella typhimurium*. *J. Bacteriol.* 176, 5474–5482. doi: 10.1128/jb.176.17.5474-5482.1994
- Cheng, C. C., Duar, R. M., Lin, X., Perez-Munoz, M. E., Tollenaar, S., Oh, J. H., et al. (2020). Ecological importance of cross-feeding of the intermediate metabolite 1,2-propanediol between bacterial gut symbionts. *Appl. Environ. Microbiol.* 86, e190–e120. doi: 10.1128/AEM.00190-20
- Cheng, S., Sinha, S., Fan, C., Liu, Y., and Bobik, T. A. (2011). Genetic analysis of the protein shell of the microcompartments involved in coenzyme B12-dependent 1, 2-propanediol degradation by *Salmonella*. *J. Bacteriol.* 193, 1385–1392. doi: 10.1128/jb.01473-10
- Collado, M. C., Meriluoto, J., and Salminen, S. (2007). In vitro analysis of probiotic strain combinations to inhibit pathogen adhesion to human intestinal mucus. *Food Res. Int.* 40, 629–636. doi: 10.1016/j.foodres.2006.11.007
- Cousin, F. J., Mater, D. D., Foligné, B., and Jan, G. (2011). Dairy propionibacteria as human probiotics: a review of recent evidence. *Dairy Sci. Technol.* 91, 1–26.
- Cox, J., Hein, M. Y., Lubber, C. A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. *Mol. Cell. Prot.* 13, 2513–2526. doi: 10.1074/mcp.m113.031591
- Crowley, C. S., Sawaya, M. R., Bobik, T. A., and Yeates, T. O. (2008). Structure of the PduU shell protein from the Pdu microcompartment of *Salmonella*. *Structure* 16, 1324–1332. doi: 10.1016/j.str.2008.05.013
- De Weirdt, R., Crabbe, A., Roos, S., Vollenweider, S., Lacroix, C., van Pijkeren, J. P., et al. (2012). Glycerol supplementation enhances *L. reuteri*'s protective effect against *S. Typhimurium* colonization in a 3-D model of colonic epithelium. *PLoS One* 7:e37116. doi: 10.1371/journal.pone.0037116
- Deptula, P., Laine, P. K., Roberts, R. J., Smolander, O.-P., Vihinen, H., Piironen, V., et al. (2017). De novo assembly of genomes from long sequence reads reveals uncharted territories of *Propionibacterium freudenreichii*. *BMC Genomics* 18:790. doi: 10.1186/s12864-017-4165-9
- Falentin, H., Deutsch, S.-M., Jan, G., Loux, V., Thierry, A., Parayre, S., et al. (2010). The complete genome of *Propionibacterium freudenreichii* CIRM-BIA1T, a hardy actinobacterium with food and probiotic applications. *PLoS One* 5:e11748. doi: 10.1371/journal.pone.0011748
- Ferlez, B., Sutter, M., and Kerfeld, C. A. (2019). Glycyl radical enzyme-associated microcompartments: redox-replete bacterial organelles. *mBio* 10, e2327–18.
- Greening, C., and Lithgow, T. (2020). Formation and function of bacterial organelles. *Nat. Rev. Microbiol.* 18, 677–689.
- Hojó, K., Yoda, N., Tsuchita, H., Ohtsu, T., Seki, K., Taketomo, N., et al. (2002). Effect of ingested culture of *Propionibacterium freudenreichii* ET-3 on fecal microflora and stool frequency in healthy females. *Biosci. Microflora* 21, 115–120. doi: 10.12938/bifidus1996.21.115
- Hosseini, E., Grootaert, C., Verstraete, W., and Van de Wiele, T. (2011). Propionate as a health-promoting microbial metabolite in the human gut. *Nutr. Rev.* 69, 245–258. doi: 10.1111/j.1753-4887.2011.00388.x
- Huang, Y., and Adams, M. C. (2004). In vitro assessment of the upper gastrointestinal tolerance of potential probiotic dairy propionibacteria. *Int. J. Food Microbiol.* 91, 253–260. doi: 10.1016/j.ijfoodmicro.2003.07.001
- Jakobson, C. M., and Tullman-Ereck, D. (2016). Dumpster diving in the gut: bacterial microcompartments as part of a host-associated lifestyle. *PLoS Pathogens* 12:e1005558. doi: 10.1371/journal.ppat.1005558
- Kaneko, T., Mori, H., Iwata, M., and Meguro, S. (1994). Growth stimulator for bifidobacteria produced by *Propionibacterium freudenreichii* and several intestinal bacteria. *J. Dairy Sci.* 77, 393–404. doi: 10.3168/jds.s0022-0302(94)76965-4
- Kerfeld, C. A., Aussignargues, C., Zarzycki, J., Cai, F., and Sutter, M. (2018). Bacterial microcompartments. *Nat. Rev. Microbiol.* 16, 277–290.
- Korbel, J. O., Doerks, T., Jensen, L. J., Perez-Iratxeta, C., Kaczanowski, S., Hooper, S. D., et al. (2005). Systematic association of genes to phenotypes by genome and literature mining. *PLoS Biol.* 3:e134. doi: 10.1371/journal.pbio.0030134
- Lu, J., Boeren, S., De Vries, S., Van Valenberg, H., Vervoort, J., and Hettinga, K. (2011). Filter-aided sample preparation with dimethyl labeling to identify and quantify milk fat globule membrane proteins. *J. Prot.* 75, 34–43. doi: 10.1016/j.jprot.2011.07.031
- Macfarlane, G. T., and Gibson, G. R. (1997). “Carbohydrate fermentation, energy transduction and gas metabolism in the human large intestine,” in *Gastrointestinal Microbiology: Gastrointestinal Ecosystems and Fermentations*, Vol. 1, eds R. I. Mackie and B. A. White (Boston, MA: Springer US), 269–318. doi: 10.1007/978-1-4615-4111-0_9
- Meile, L., Le Blay, G., and Thierry, A. (2008). Safety assessment of dairy microorganisms: propionibacterium and Bifidobacterium. *Int. J. Food Microbiol.* 126, 316–320. doi: 10.1016/j.ijfoodmicro.2007.08.019
- Quesada-Chanto, A., Schmid-Meyer, A., Schroeder, A., Carvalho-Jonas, M., Blanco, I., and Jonas, R. (1998). Effect of oxygen supply on biomass, organic acids and vitamin B12 production by *Propionibacterium shermanii*. *World J. Microbiol. Biotechnol.* 14, 843–846.
- Ravcheev, D. A., Moussu, L., Smajic, S., and Thiele, I. (2019). Comparative genomic analysis reveals novel microcompartment-associated metabolic pathways in the human gut microbiome. *Front. Genet.* 10:636. doi: 10.3389/fgene.2019.00636
- Richter-Dahlfors, A., Ravnum, S., and Andersson, D. (1994). Vitamin B12 repression of the cob operon in *Salmonella typhimurium*: translational control of the *cbiA* gene. *Mol. Microbiol.* 13, 541–553. doi: 10.1111/j.1365-2958.1994.tb00449.x
- Roessner, C. A., Huang, K.-x., Warren, M. J., Raux, E., and Scott, A. I. (2002). Isolation and characterization of 14 additional genes specifying the anaerobic biosynthesis of cobalamin (vitamin B12) in *Propionibacterium freudenreichii* (*P. shermanii*) The GenBank accession numbers for the sequences reported in this paper are AY033235, AY033236, U13043 and U51164. *Microbiology* 148, 1845–1853. doi: 10.1099/00221287-148-6-1845
- Sampson, E. M., and Bobik, T. A. (2008). Microcompartments for B12-dependent 1, 2-propanediol degradation provide protection from DNA and cellular damage by a reactive metabolic intermediate. *J. Bacteriol.* 190, 2966–2971. doi: 10.1128/jb.01925-07
- Saroufi, T., Parayre, S., Guernec, G., Loux, V., Montfort, J., Cam, A. L., et al. (2013). A unique in vivo experimental approach reveals metabolic adaptation of the probiotic *Propionibacterium freudenreichii* to the colon environment. *BMC Genomics* 14:911. doi: 10.1186/1471-2164-14-911
- Satomi, K., Kurihara, H., Isawa, K., Mori, H., and Kaneko, T. (1999). Effects of culture-powder of *Propionibacterium freudenreichii* ET-3 on fecal microflora of normal adults. *Biosci. Microflora* 18, 27–30. doi: 10.12938/bifidus1996.18.27
- Seeliger, S., Janssen, P. H., and Schink, B. (2002). Energetics and kinetics of lactate fermentation to acetate and propionate via methylmalonyl-CoA or acrylyl-CoA. *FEMS Microbiol. Lett.* 211, 65–70. doi: 10.1111/j.1574-6968.2002.tb11204.x
- Smaczniak, C., Li, N., Boeren, S., America, T., Van Dongen, W., Goerdal, S. S., et al. (2012). Proteomics-based identification of low-abundance signaling and regulatory protein complexes in native plant tissues. *Nat. Protocols* 7:2144. doi: 10.1038/nprot.2012.129
- Sokolovskaya, O. M., Shelton, A. N., and Taga, M. E. (2020). Sharing vitamins: cobamides unveil microbial interactions. *Science* 369:eaba0165. doi: 10.1126/science.aba0165

- Sutter, M., Greber, B., Aassignargues, C., and Kerfeld, C. A. (2017). Assembly principles and structure of a 6.5-MDa bacterial microcompartment shell. *Science* 356, 1293–1297. doi: 10.1126/science.aan3289
- Thiennimitr, P., Winter, S. E., Winter, M. G., Xavier, M. N., Tolstikov, V., Huseby, D. L., et al. (2011). Intestinal inflammation allows *Salmonella* to use ethanolamine to compete with the microbiota. *Proc. Natl. Acad. Sci. U.S.A.* 108, 17480–17485. doi: 10.1073/pnas.1107857108
- Toraya, T., Honda, S., and Fukui, S. (1979). Fermentation of 1, 2-propanediol and 1, 2-ethanediol by some genera of *Enterobacteriaceae*, involving coenzyme B12-dependent diol dehydratase. *J. Bacteriol.* 139, 39–47. doi: 10.1128/jb.139.1.39-47.1979
- Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M. Y., Geiger, T., et al. (2016). The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods* 13, 731–740. doi: 10.1038/nmeth.3901
- Vizcaino, J. A., Csordas, A., del-Toro, N., Dianas, J. A., Griss, J., Lavidas, I., et al. (2016). 2016 update of the PRIDE database and related tools. *Nucleic Acids Res.* 44(D1), D447–D456.
- Wendrich, J. R., Boeren, S., Möller, B. K., Weijers, D., and De Rybel, B. (2017). *In Vivo Identification of Plant Protein Complexes Using IP-MS/MS Plant Hormones*. Berlin: Springer, 147–158.
- Wilson, J. W. (2021). Manipulating microcompartment operons to study mechanism and function. *Curr. Opin. Microbiol.* 60, 66–72. doi: 10.1016/j.mib.2021.01.014
- Wiśniewski, J. R., Zougman, A., Nagaraj, N., and Mann, M. (2009). Universal sample preparation method for proteome analysis. *Nat. Methods* 6:359. doi: 10.1038/nmeth.1322
- Xue, J., Murrieta, C. M., Rule, D. C., and Miller, K. W. (2008). Exogenous or L-rhamnose-derived 1, 2-propanediol is metabolized via a pduD-dependent pathway in *Listeria innocua*. *Appl. Environ. Microbiol.* 74, 7073–7079. doi: 10.1128/aem.01074-08
- Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics J. Integr. Biol.* 16, 284–287. doi: 10.1089/omi.2011.0118
- Zeng, Z., Smid, E. J., Boeren, S., Notebaart, R. A., and Abee, T. (2019). Bacterial microcompartment-dependent 1,2-Propanediol utilization stimulates anaerobic growth of *Listeria monocytogenes* EGDe. *Front. Microbiol.* 10:2660. doi: 10.3389/fmicb.2019.02660

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Dank, Zeng, Boeren, Notebaart, Smid and Abee. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.