



5-20-2015

Bacterial Symbionts, *Buchnera*, and Starvation on Wing Dimorphism in English Grain Aphid, *Sitobion avenae* (F.) (Homoptera: Aphididae)

Fangmei Zhang
Chinese Academy of Agricultural Sciences, China

Xiangrui Li
Chinese Academy of Agricultural Sciences, China

Yunhui Zhang
Chinese Academy of Agricultural Sciences, China

Brad Coates
US Department of Agriculture

Xuguo Joe Zhou
University of Kentucky, xuguozhou@uky.edu

See the [end of this page for additional authors](https://uknowledge.uky.edu/entomology_facpub) https://uknowledge.uky.edu/entomology_facpub



Part of the [Entomology Commons](#), and the [Physiology Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Zhang, Fangmei; Li, Xiangrui; Zhang, Yunhui; Coates, Brad; Zhou, Xuguo Joe; and Cheng, Dengfa, "Bacterial Symbionts, *Buchnera*, and Starvation on Wing Dimorphism in English Grain Aphid, *Sitobion avenae* (F.) (Homoptera: Aphididae)" (2015). *Entomology Faculty Publications*. 79.
https://uknowledge.uky.edu/entomology_facpub/79

This Article is brought to you for free and open access by the Entomology at UKnowledge. It has been accepted for inclusion in Entomology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Bacterial Symbionts, *Buchnera*, and Starvation on Wing Dimorphism in English Grain Aphid, *Sitobion avenae* (F.) (Homoptera: Aphididae)

Digital Object Identifier (DOI)

<https://doi.org/10.3389/fphys.2015.00155>

Notes/Citation Information

Published in *Frontiers in Physiology*, v. 6, article 155, p. 1-9.

© 2015 Zhang, Li, Zhang, Coates, Zhou and Cheng.

This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Authors

Fangmei Zhang, Xiangrui Li, Yunhui Zhang, Brad Coates, Xuguo Joe Zhou, and Dengfa Cheng

Bacterial symbionts, *Buchnera*, and starvation on wing dimorphism in English grain aphid, *Sitobion avenae* (F.) (Homoptera: Aphididae)

OPEN ACCESS

Edited by:

Raman Chandrasekar,
Kansas State University, USA

Reviewed by:

Raman Chandrasekar,
Kansas State University, USA
Xiaowei Yang,
Cornell University, USA
Pei Liang,
China Agricultural University, China

*Correspondence:

Xuguo "Joe" Zhou,
Department of Entomology, University
of Kentucky, S-225 Agricultural
Science Center North, Lexington,
KY 40546-0091, USA
xuguozhou@uky.edu;
Dengfa Cheng,
State Key Laboratory for Biology of
Plant Diseases and Insect Pests,
Institute of Plant Protection, Chinese
Academy of Agricultural Sciences,
Yuanmingyuan West Road,
Beijing 100193, China
dfcheng@ippcaas.cn

†These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Integrative Physiology,
a section of the journal
Frontiers in Physiology

Received: 17 March 2015

Accepted: 29 April 2015

Published: 20 May 2015

Citation:

Zhang F, Li X, Zhang Y, Coates B,
Zhou X and Cheng D (2015) Bacterial
symbionts, *Buchnera*, and starvation
on wing dimorphism in English grain
aphid, *Sitobion avenae* (F.)
(Homoptera: Aphididae).
Front. Physiol. 6:155.
doi: 10.3389/fphys.2015.00155

Fangmei Zhang^{1†}, Xiangrui Li^{1†}, Yunhui Zhang¹, Brad Coates², Xuguo "Joe" Zhou^{3*}
and Dengfa Cheng^{1*}

¹ State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China, ² Corn Insects and Crop Genetics Research Unit, United States Department of Agriculture - Agricultural Research Service, Ames, IA, USA, ³ Department of Entomology, University of Kentucky, Lexington, KY, USA

Wing dimorphism in aphids can be affected by multiple cues, including both biotic (nutrition, crowding, interspecific interactions, the presence of natural enemies, maternal and transgenerational effects, and alarm pheromone) and abiotic factors (temperature, humidity, and photoperiod). The majority of the phloem-feeding aphids carry *Buchnera*, an obligate symbiotic proteobacteria. *Buchnera* has a highly reduced genome size, but encode key enzymes in the tryptophan biosynthetic pathway and is crucial for nutritional balance, development and reproduction in aphids. In this study, we investigated the impact of two nutritional-based biotic factors, symbionts and starvation, on the wing dimorphism in the English grain aphid, *Sitobion avenae*, a devastating insect pest of cereal crops (e.g., wheat) worldwide. Elimination of *Buchnera* using the antibiotic rifampicin significantly reduced the formation of winged morphs, body mass, and fecundity in *S. avenae*. Furthermore, the absence of this primary endosymbiont may disrupt the nutrient acquisition in aphids and alter transgenerational phenotypic expression. Similarly, both survival rate and the formation of winged morphs were substantially reduced after neonatal (<24 h old) offspring were starved for a period of time. The combined results shed light on the impact of two nutritional-based biotic factors on the phenotypic plasticity in aphids. A better understanding of the wing dimorphism in aphids will provide the theoretical basis for the prediction and integrated management of these phloem-feeding insect pests.

Keywords: *Sitobion avenae*, *Buchnera*, wing dimorphism, starvation, obligate symbiont, nutrition, biotic factors, nutritional requirements

Introduction

Aphid species are soft-bodied sap-feeding insects that have developed specialized mouthparts that allow feeding on plant phloem. Although phloem contains a concentration of carbohydrates, it is nutritionally deficient in respect to nitrogenous compounds that serve as precursors to many amino acid biosynthetic pathways (Sandström and Moran, 1999). An evolutionary relationship has developed between species in insect Family Aphidoidea and the obligate intracellular

endosymbiotic proteobacteria, *Buchnera aphidicola* (Lai et al., 1994). Characteristic of several other endosymbionts *Buchnera* have a highly reduced genome content that has resulted in loss of key enzymatic components of carbohydrate and lipid metabolizing pathways (Shigenobu et al., 2000), but these losses are compensated by the host-derived cellular pathways (Douglas, 1989, 1998). Correspondingly, *Buchnera* produce many essential amino acids that are not present in the nutritionally incomplete plant phloem (Douglas, 1989, 1998). Bacterial symbionts provide a supplementary source of essential nutrients, including essential amino acids (Douglas, 1989; Wilkinson and Ishikawa, 2001; Hansen and Moran, 2011), vitamins (Dadd et al., 1967; Douglas, 1988; Nakabachi and Ishikawa, 1999), and lipids (Douglas, 1988). In addition to providing nutrients, this obligate symbiosis is crucial for aphid development and reproduction (Sasaki et al., 1991; Douglas, 1995; Hardie and Leckstein, 2007; Simon et al., 2011), behavior (Hongoh and Ishikawa, 1994), pigmentation (Tsuchida et al., 2010), and resistance against biotic and abiotic stresses (Scarborough et al., 2005). This mutualism has evolved beyond nutritional interdependence, but also includes the development of the specialized adipose cells, called bacteriocytes, that are vertically transmitted from the maternal parent to offspring (Wernegreen and Moran, 2001). Phylogenetic analysis provided evidence that this symbiotic relationship could date back to at least 200 million years (Moran and Baumann, 1994; Baumann et al., 1995).

The grain aphid, *Sitobion avenae* (F.), is a destructive pest of wheat crops worldwide. As a phloem-feeding insect, aphids reduce yield by (i) sucking sap from wheat, which affects the grain filling stage, (ii) excreting honeydew which becomes covered with sooty molds, and (iii) transmitting barley yellow dwarf virus. Similar to other Aphidoidea, *S. avenae* adults can exist in either winged or wingless forms in field conditions. This phenotypic plasticity is known to arise in response to environmental cues, including changes in nutritional availability (host plant quality), overcrowding (tactile stimulation), interspecific interactions (Braendle et al., 2006 and references within), the presence of nature enemies (Weisser et al., 1999; Mehrparvar et al., 2013), and abiotic factors, such as temperature (Parish and Bale, 1990), humidity, photoperiod (Lees, 1966; Hardie, 1987), and alarm pheromone (Hatano et al., 2010). Besides these environmental cues in a traditional phenotype-by-genotype interaction, evidence suggests that maternal and trans-generational effects, including grand-maternal phenotype, maternal phenotype, and developmental stages (Müller et al., 2001) modulate wing dimorphism in aphids. This capacity of a single genotype to produce two or more discrete phenotypic forms, termed polyphenism, is somewhat prevalent across insect Orders including hymenopteran caste systems (Wilson, 1971) and morphogenesis of Lepidoptera from larval to moth forms (Shapiro, 1976).

The function of symbionts in aphid reproduction and performance has been investigated extensively (Hardie and Leckstein, 2007), however, more limited efforts have been devoted to our understanding of the impact of *Buchnera* endosymbionts on phenotypic plasticity (polyphenism) among aphid species. The objective of this research is to study

the impacts of symbionts and starvation on *S. avenae* wing dimorphism. A better understanding of the mechanisms of wing development in aphids will provide new insights in the long term sustainable management of this key insect pest of wheat.

Materials and Methods

Aphid Colony Maintenance

Sitobion avenae (F.) adults were collected from wheat fields in Langfang (39°30'42"N, 116°36'7"E), Hebei Province, China, in 2012. Aphids were maintained on 15 cm wheat seedlings in an environmental cabinet (RXZ-380B, Nb-Jn Instrument Factory, Ningbo, China) at 20°C, 60% RH, and L:D 16:8 h photoperiod. The aphid colony was generated from a single wingless adult female. The offspring from this single clone was reared in 9 cm diameter Petri dishes individually. After three generations, aphids were subjected to the subsequent experiments. The neonatal nymphs are sensitive to environmental cues associated with wing development. To generate winged morph, more than 30 newly hatched nymphs (<24 h) were reared together in 9 cm diameter Petri dishes. In contrast, only a single nymph was kept in the Petri dishes to generate wingless morph. Selective elimination of primary endosymbionts was obtained as described below.

Rifampicin Treatment

The 24 h postnatal nymphs were maintained on artificial diets (Chen et al., 2000) supplemented with 50 µg/ml of rifampicin for 48 h (Rahbe et al., 1994; Wilkinson and Douglas, 1995; Adams et al., 1996; Wilkinson and Ishikawa, 2001). This treatment has been used to eliminate aphid-*Buchnera* endosymbiotic bacteria with minimal deleterious effects on the aphid itself (Adams and Douglas, 1997). The control aphids were maintained on regular artificial diets for 48 h. The status of *Buchnera* in aphid samples was assessed by PCR. Total DNAs were isolated from natural symbiotic (control; non-rifampicin treated) and rifampicin treated individuals using E.Z.N.A.[®] Insect DNA Kit (OMEGA, Bio-tek) according to the manufacturer's protocol. The primers for *Buchnera*, forward primer 5'-GTCGGCTCATCACATCC-3' and reverse primer 5'-TTCCGTCTGTATTATCTCCT-3', were designed based on *rrl-aroE* gene sequences obtained from the *Buchnera* genome (Accession Number: GCA_000009605.1) (Shigenobu et al., 2000) using Primer 5.0 software (Primer-E Ltd., Plymouth, UK). The PCR amplification cycles included an initial denaturation step at 95°C for 5 min, 35 cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 60 s, and a final extension step of 10 min at 72°C. Amplification products were analyzed on 1% agarose gels, stained with ethidium bromide, and visualized using the Imaging G6 System (DHS Life Science & Technology Co., Ltd., Beijing, China).

Phenotypic Impacts of Antibiotic Treatment

Rifampicin treated and symbiotic (control) aphids were transferred to wheat seedlings, with 20 nymphs per plant. A total of five replicates were carried out for each experiment. Before the experiment, the number of winged and wingless morphs was counted, and these aphids were considered the

F₁ population (namely WG-1 and WL-1). These adults were allowed to reproduce for 3 days. When their offspring developed into the fourth-instar nymphs or adults, the number of winged and wingless morphs were counted again, and this generation of aphids was defined as F₂ population (namely WG-2 and WL-2).

Phenotypic Impacts of Starvation

The phenotypic impacts of starvation on wing development were assayed using first and second instar nymphs. Neonatal (<24 h) nymphs of both winged morphs were collected and kept in groups of 20 within 9 cm diameter Petri dishes provisioned with water-saturated filter papers. Nymphs were maintained on Petri dishes for 0, 24, 48, and 72 h without food before returning to wheat seedlings. The survival rate of nymphs were documented at time points 0, 24, and 48 h after treatment. When nymphs reached the adult stage, the number of winged and wingless morphs was counted, including both F₁ and F₂ populations.

Statistical Analysis

Statistical analyses were carried out using SAS statistical software 9.2 (SAS Institute Inc., Cary, NC, USA). Data were presented as mean \pm stand error. For different treatments (antibiotics and starvation), the proportion of winged/wingless among treated and control aphids was compared and analyzed, respectively. An independent *T*-test and One-Way ANOVA for means were used to investigate the phenotypic impacts. The end point measurements included the survival rate and the percentage of winged morphs.

Results

Establishment of Rifampicin-Cured Aphid Lines

To verify the elimination of *Buchnera* after antibiotic treatment, we isolated DNAs from both symbiotic and rifampicin-cured aphids, respectively. Gel electrophoresis analysis showed that the DNAs extracted from both morphs were intact (Figure 1, Lanes 2 and 3), however, *Buchnera* was absent from rifampicin-cured aphids (Figure 1, Lane 5) compared to symbiotic controls (Figure 1, Lane 4).

Phenotypic Impact of Antibiotics Treatment on the Wing Development

F₁ Generation

The overall percentage of winged morphs was reduced in rifampicin treated aphids compared to the untreated controls (Figure 2A). Specifically, 11.67 and 45.47% of the F₁ aphids had developed wings among the *S. avenae* that were, respectively, winged and wingless before the treatment. These percentages of winged aphids were decreased to 5.25 and 25.53% in the rifampicin treatment group. For aphids derived from the parental winged group (WG-1), there was no significant difference in the wing development between the rifampicin treatment and control groups ($t = -1.60$, $P = 0.1493$), but in contrast a significant difference ($t = -3.63$, $P = 0.0176$) was observed among aphids from the initially wingless group (WL-1). Additionally, significant differences in the proportion of winged





FIGURE 1 | Electrophoretic detection of *Buchnera* among symbiotic and cured aphids. Lane 1: 100 bp ladder Marker; Lane 2: DNA of symbiotic aphids; Lane 3: DNA of rifampicin-cured aphids; Lane 4: PCR amplification of *Buchnera* in symbiotic aphids; Lane 5: PCR amplification of *Buchnera* in rifampicin-cured aphids.

and wingless individuals, were detected between WG-1 and WL-1 for both, the rifampicin ($t = -7.62$, $P = 0.0001$) and the control treatments ($t = -5.40$, $P = 0.0006$). Elimination of *Buchnera* from neonatal nymphs significantly reduced subsequent fecundity in aphids compared to control groups (Figure 2B). The number of offspring is 94.60 and 46.20 among winged and wingless *S. avenae* before the treatments. These numbers dropped to 21.60 and 13.40 after rifampicin treatment. For the initially wingless individuals (WL-1), no significant differences were detected between the number of offspring produced in the rifampicin-treated and the control groups ($t = -1.74$, $P = 0.1194$), but significant differences were detected between treatments for the initially winged individuals (WG-1) ($t = -3.98$, $P = 0.0041$).

F₂ Generation


The percentage of winged morphs was also reduced in the F₂ generation among rifampicin-treated aphids compared to the untreated controls (Figure 2C). Specifically, 54.54 and 54.11% of the F₂ aphids had wings, respectively, among winged (WG-2) and wingless *S. avenae* groups in the untreated controls



(WLG-2). These percentages dropped to 33.72 and 20.14% after rifampicin treatment for WG-2 and WLG-2, respectively. For winged individuals (WG-1), there were no significant difference in the proportion of individuals that developed wings between the rifampicin-treated and the untreated controls ($t = -1.33$, $P = 0.2200$). In contrast, significant differences were observed for the wingless group (WLG-2) ($t = -3.81$, $P = 0.0052$). The proportion of a red colored morph was not significantly different between WG-2 (48.44%) compared to WLG-2 (47.85%) ($t = 0.02$, $P = 0.9815$), although a significant difference was observed in the percentage of the red color between rifampicin-treated and untreated controls in WG-2 ($t = 2.92$, $P = 0.0433$). No significant differences in the red colored morph were detected among treatments in the WLG-2 ($t = 1.98$, $P = 0.1079$) (Figure 2D). It is worth noting that, besides the normal wingless (Figure 3A) and winged (Figure 3C) forms, a new apterous adult form in the rifampicin-treated aphids showed an intermediate winged/wingless phenotype (Figure 3B).


Between Generations

The percentage of winged morphs between generations exhibited a different trend. Specifically, the percentage of winged morphs was 5.25 and 33.7% for WG-1 and WG-2, respectively, while it was 25.53 and 12.53% in the rifampicin treated groups. Moreover, there were no significant differences in the development of wing dimorphism between the generations (WG-1 vs. WG-2: $t = -2.03$, $P = 0.1089$; WLG-1 vs. WLG-2: $t = 1.99$, $P = 0.1109$).



Phenotypic Impact of Starvation on the Wing Development

Both survival rate and the percentage of winged morphs were significantly reduced after starvation treatment to neonatal nymphs (Figures 4A,B). Starvation for 0, 24, 48, and 72 h had significantly different mean survival rates in WG-1 ($F = 341.83$, $P < 0.0001$) and WLG-1 ($F = 630.16$, $P < 0.0001$), respectively. The survival rates were consistently over 90% when aphids were starved for 24 h, with no significant difference between WG-1 and WLG-1. In contrast, survivorships in WG-1 and WLG-1 were reduced substantially to 47.5 and 66.5%, respectively, after aphids



were starved for 48 h. The survival rate decreased dramatically to 1.5 and 1.0% in, respectively, WG-1 and WL-1 after 72 h starvation, at which point the proportion of individuals that had developed wings could no longer be effectively estimated. Aphid mortality reached 100% when starved for more than 72 h. Consequently, wing development was further investigated only for aphids starved 0, 24, and 48 h.

Starvation significantly reduced the percentage of the winged morphs in F₁ *S. avenae*. Specifically, the percentage of winged aphids was drastically reduced from 18.75 to 8.33 and 1.11%, respectively, after 24 and 48 h of starvation treatments in WG-1 ($F = 8.75$, $P = 0.0045$). However, the percentage of winged morphs in WL-1 was reduced from 13.69 to 10.12 and 4.97%, respectively, after 24 and 48 h of starvation treatment, with no significant differences between different starvation times ($F = 2.20$, $P = 0.1530$) (Table 1, Figure 4B). Among the three different starvation times, there were no significant differences in the wing development between WG-1 and WL-1 (0 h: $t = 1.01$, $P = 0.3434$; 24 h: $t = -0.48$, $P = 0.6473$; 48 h: $t = -1.00$, $P = 0.3738$) (Table 1).

The starvation to neonatal nymphs affected aphid fecundity in comparison to the non-starved controls. Specifically, the number of offspring produced for the WG-2 generated increased from 44.4 to 86.4 when WG-1 parents starved 0 and 48 h were compared. The opposite effect was shown in a decrease of WL-2 offspring from 147.0 to 73.8 when contrasting those derived from WL-1 parents starved 0 and 48 h (Figure 4C). This change in reproduction (number of offspring) across time (0 vs. 48 h) was reflected as a significant difference in WG-2 ($F = 4.32$, $P = 0.0386$) and WL-2 ($F = 9.70$, $P = 0.0031$) (Table 1).

On the contrary, the proportion of the winged aphids increased as the starvation time was prolonged in WG-2 ($F = 12.60$, $P = 0.0011$) (Figure 4D). There were no significant differences in the percentage of winged individuals between controls and 24 h starvation treatments in WL-2. Starvation for 48 h led to significantly lower rate of winged offspring compared to the non-starved controls and 24 h-starved aphids ($F = 10.04$, $P = 0.0257$) (Table 1).

Discussion

Symbiotic relationships with intracellular bacteria, such as *Wolbachia* and *Buchnera*, are pronounced across Insecta, and have evolved intricate interdependencies for mutual survival and propagation (Baumann et al., 2000; Werren et al., 2008). Many insect endosymbionts have crucial roles in host survival, whereby removal of these symbionts often results in notable fitness costs. Previous studies on *Buchnera*, the primary symbionts of sap-feeding insects, has revealed several effects on host insect fitness, development and reproduction (Sasaki et al., 1990; Douglas, 1995, 1998; Febvay et al., 1995). Furthermore, detailed information available indicated the impact of aposymbionts on the performance, nutrition, metabolism, and feeding behavior of aphid (Prosser and Douglas, 1991; Wilkinson and Douglas, 1995, 1996). Moran and Yun (2015) found that aphids with *Buchnera* replacement exhibited a dramatic increase in their heat tolerance. This relationship with *Buchnera* is essential for compensation of nutritional deficiencies in plant phloem encountered by sap-feeding insects, and forms the basis of this mutualism. Thus aphids, through the mutualistic association with *Buchnera*

TABLE 1 | The number of offspring and proportion of the winged individuals derived from the winged and wingless adults.

	The proportion of F ₁		The number of offspring of F ₁		The proportion of F ₂	
	WG-1	WLG-1	WG-2	WLG-2	WG-2	WLG-2
0h	18.75 ± 4.3 ^a	13.69 ± 2.58 ^a	44.4 ± 4.87 ^a	147 ± 15.97 ^a	32.50 ± 5.2 ^a	70.54 ± 1.32 ^a
24h	8.33 ± 2.69 ^b	10.12 ± 2.64 ^a	62.8 ± 11.74 ^{ab}	145.8 ± 14.43 ^a	64.50 ± 9.84 ^b	77.66 ± 2.58 ^a
48h	1.11 ± 1.11 ^b	4.97 ± 3.55 ^a	86.4 ± 12.09 ^b	73.8 ± 9.4 ^b	81.12 ± 4.64 ^b	55.44 ± 5.36 ^b

Values are mean ± SE. Means within a column followed by different letters (a, b) are significantly different (LSD's test: $P < 0.05$).

symbionts have attained the capacity to utilize phloem sap as the sole dietary source throughout their lifecycle (Sasaki et al., 1990; Douglas and Prosser, 1992; Febvay et al., 1995). It was shown that 10 essential amino acids are scarce in the phloem sap (Sandström and Pettersson, 1994). Analysis and annotation of the *Buchnera* genome predicted the presence of most genes involved in the biosynthesis of essential amino acids and may lack certain cell wall lipid biosynthetic pathways (Shigenobu et al., 2000; Tamas et al., 2002; van Ham et al., 2003), and thus are supplied to the host through the endosymbiotic relationship (Baumann, 2005). Indeed, genome analysis reveals host–symbiont cooperation in the production of amino acids in aphids (Hansen and Moran, 2011). These prior analyses strongly suggested that that *Buchnera* are capable of synthesizing essential amino acids required that for aphid nutritional health, but may be depending on aphids in order to obtain structural lipids. Furthermore, the development and maternal transmission of specialized adipose cells, bacteriocytes, to offspring (Wernegreen and Moran, 2001) show that host cell and reproductive modifications have occurred during this intricate co-evolution. It can be argued that symbiosis resulted in the co-evolution aphids and *Buchnera* genomes in order to facilitate this partnership and this interaction could have allowed for the emergence of the aphid branch of hemipteran insects greater than 200 million years ago (Moran and Baumann, 1994; Baumann et al., 1995).

Prior studies have demonstrated that nutrient availability affects wing dimorphism in aphids and is one of the factors that influence this polyphenism, whereby nutritional deficiencies increase the proportion of apterous (winged) progeny (Johnson, 1966; Müller et al., 2001; Braendle et al., 2006). *Buchnera*, being an obligate intracellular symbiont with a relatively well-understood role in providing essential amino acids to aphid hosts, may play a role in wing dimorphism. Specifically, it was revealed that the seemingly obvious consequences of removing the *Buchnera* endosymbiont from aphids by antibiotic treatment had decreased the propensity of aphids to develop as winged individuals. Specifically, prior studies showed that feeding bean seedling treated with chlortetracycline caused a majority of *Aphis fabae* to develop as wingless or winged/wingless intermediate adult forms (Leonardo and Mondor, 2006). The primary *Buchnera* symbiont of *Acyrtosiphon pisum* has very similar effects on aphid fitness and polyphenism (Russell and Moran, 2006; Hardie and Leckstein, 2007). Data from the current study clearly demonstrated *Buchnera*-free neonatal *S. avenae* resulted in a significant reduction in the proportions of winged offspring, and, in agreement with prior studies indicated above, suggested that the primary endosymbiont can affect wing

dimorphism in Aphidoidea. Although intriguing to surmise a direct effect between *Buchnera* and wing dimorphism, a highly likely explanation for these and other results may be that the resulting lack of essential amino acids and reduced aphid vigor could be the cause of this reduction in winged forms (Hardie and Leckstein, 2007). Indeed, previous studies investigated the effect of amino acids on the wing dimorphism in aphids, and showed that the deletion of methionine, isoleucine, or histidine, respectively, could significantly increase the percentage of winged morphs in *Myzus persicae* (Dadd, 1968; Dadd and Krieger, 1968; Mittler and Kleinjan, 1970; Sutherland and Mittler, 1971). Similarly, the development of winged individuals in *A. fabae* was induced by the removal of dietary arginine, leucine, lysine, and proline, respectively (Leckstein and Liewellyn, 1973).

Our results showed a highly significant interaction between *Buchnera* and the development of winged individuals in the treatments WG-1 and WLG-1. Previous studies evidencing that the features of nutritional interactions between the aphid and its symbionts were substantially varied by aphid's species, age and morph (Douglas and Dixon, 1987). The possible explanations for this phenomenon were suggested to involve the switching off/on of the genes involved in the formation of wing tissues or wingless development (Hardie and Leckstein, 2007). Under the nutritionally adverse conditions part of mycetocytes are decreased (Hongoh and Ishikawa, 1994), causing the changes of number and size of the symbiont and further wing development changes. However, the effect of aposymbiotic on wing dimorphism in aphids was not applied to any species, such as *A. pisum* and *M. viciae* (Hardie and Leckstein, 2007). Elimination of bacteria with antibiotic rifampicin from *S. avenae* significantly reduced growth sterility, and reduced aphid mass in almost all cases, especially reproduction of F₁ of winged adults, which is consistent with previous reports (Douglas, 1995, 1998; Wilkinson and Douglas, 1995; Douglas, 1995; Hardie and Leckstein, 2007).

Moran and Jarvik (2010) observed that aphid genome contained several genes for carotenoid synthesis, which were not found in animal genomes, and might be responsible for red or green coloration of the aphids. Aphid body color can be affected by many factors such as temperature, host plant, light, symbionts, and natural enemies (Braendle et al., 2006 and references within). Intriguingly, our data showed that the number of aphids with red coloration increased significantly when the aphid were subjected to bacteria-free treatment, and cleared of the *Buchnera* endosymbiont. It represents the first known study to show this direct interaction. Previous studies showed that the coinfecting of facultative endosymbiont, *Rickettsiella* and *Hamiltonella*, could

change the body color of pea aphid (*A. pisum*) from red to green (Tsuchida et al., 2010, 2014). These changes in aphid body color can influence prey-predator interactions by changing visual cues and camouflage. Although upregulation of carotenoids may be a cause of the change in aphid coloration, further studies are required to document molecular genetic changes in this pathway at the transcriptional level and/or determine the involvement of *Buchnera* (if any) in mediating these phenotypic change.

Starvation is a common experience of natural populations which causes individuals to respond to this stressful conditions by utilizing various adaptive strategies (Karan and David, 2000). In this study, starvation was used to mimic the nutritional deficiency caused under natural conditions, and resulted in a relative decrease in the formation of winged individuals as a function of increased duration of starvation among F₁ individuals for both WG-1 and WLG-1 laboratory populations. Interestingly, both the rate of reproduction in the F₁ (as measured by the number of WG-1 and WLG-1 progeny) and the proportion of winged individuals in the F₂ generation (WG-2 and WLG-2) were significantly increased with prolonged periods of starvation. Refeeding after the starvation also led to an increase in numbers and size of mycetocytes in aphids. Prior studies showed that starved aphids exhibited a drastic decrease in the total volume of mycetocyte, which was reversed by refeeding. This observed decrease could result from host utilization of nutrients within the mycetocytes as the expense of *Buchnera* provisioning (Hongoh and Ishikawa, 1994), or could also be a consequence of reduced *Buchnera* titers as host nutrients become increasingly scarce during periods of starvation. Of greater importance, our results showed that a transgenerational effect of starvation, whereby the proportion of winged individuals produced in WG-2 and WLG-2 were directly related to the duration of starvation experienced by the maternal WG-1 and WLG-1. Many insects employ multiple reproductive strategies that enable short-term tolerance of a lack of food resource, a lack of suitable oviposition sites or other environmental stresses (Brough and Dixon, 1990; Xu et al., 2012). Some reports indicated that prolonged periods of starvation for both parent aphids and young can favored the development of wingless individuals (Johnson, 1966; Xu

et al., 2012). Thus, our laboratory experiments demonstrate that unfavorable nutritional conditions (starvation) influence wing dimorphism by increasing the proportion of mobile winged individuals in the subsequent generation, and may represent an adaptive mechanism which promotes dispersal and resulting utilization of new food resources.

Combined results from both starvation and aposymbiosis studies indicate a significant role of symbionts in the wing dimorphism in aphid, which may be related to their production of essential amino acids. Additional investigations are warranted to understand the molecular signaling and interactions between symbionts and wing dimorphism in aphids. A better understanding of the wing development in aphids will provide novel targets for the sustainable management of this phloem-feeding insect pest.

Author Contributions

Conceived and designed the experiments: XL. Performed the experiments: XL and FZ. Analyzed the data: XL, FZ, and XZ. Contributed reagents/materials/analysis tools: XL, YZ, and DC. Wrote the paper: XL, FZ, BC, and XZ. All authors read and approved the final manuscript.

Acknowledgments

The authors are grateful to anonymous reviewers and editor for their constructive criticisms. Special thanks go to Dr. Runzhi Li (Shanxi Agricultural University) for his comments on an earlier draft. This research was supported by the National Natural Science Foundation of China (Award Number: 31301659), National “Twelfth Five-Year” Plan for Science & Technology Support Development Program of China (Award Number: 2012BAD19B04), China Agriculture Research System (Award Number: CARS-3). The information reported in this paper (No. 15-08-048) is part of a project of the Kentucky Agricultural Experiment Station and is published with the approval of the Director. The granting agencies have no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Adams, D., and Douglas, A. E. (1997). How symbiotic bacteria influence plant utilization by the polyphagous aphid, *Aphis fabae*. *Oecologia* 110, 528–532. doi: 10.1007/s004420050190
- Adams, D., Wilkinson, T. L., and Douglas, A. E. (1996). The aphid-bacterial symbiosis: a comparison between pea aphids and black bean aphids. *Entomol. Exp. Appl.* 80, 275–278. doi: 10.1111/j.1570-7458.1996.tb00934.x
- Baumann, P. (2005). Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Annu. Rev. Microbiol.* 59, 155–189. doi: 10.1146/annurev.micro.59.030804.121041
- Baumann, P., Baumann, L., Lai, C. Y., Rouhbakhsh, D., Moran, N. A., and Clark, M. A. (1995). Genetics, physiology and evolutionary relationships of the genus *Buchnera*: intracellular symbionts of aphids. *Annu. Rev. Microbiol.* 49, 55–94. doi: 10.1146/annurev.mi.49.100195.000415
- Baumann, P., Moran, N. A., and Baumann, L. (2000). “Bacteriocyte-associated endosymbionts of insects,” in *The Prokaryotes*, ed M. Dworkin. New York, NY: Springer-Verlag.
- Braendle, C., Davis, G. K., Brisson, J. A., and Stern, D. L. (2006). Wing dimorphism in aphids. *Heredity* 97, 192–199. doi: 10.1038/sj.hdy.6800863
- Brough, C. N., and Dixon, A. F. G. (1990). The effects of starvation on development and reproductive potential of apterous virginoparae of vetch aphid, *Megoura viciae*. *Entomol. Exp. Appl.* 55, 41–45. doi: 10.1111/j.1570-7458.1990.tb01346.x
- Chen, J. L., Ni, H. X., Ding, H. J., and Sun, J. R. (2000). Studies on a chemically defined diet of English grain aphid, *Sitobion avenae*. *J. Integr. Agric.* 33, 54–59.
- Dadd, R. H. (1968). Dietary amino acids and wing determination in the aphid *Myzus persicae*. *Ann. Entomol. Soc. Am.* 61, 1201–1210. doi: 10.1093/aesa/61.5.1201
- Dadd, R. H., and Krieger, D. L. (1968). Dietary amino acid requirements of the aphid, *Myzus persicae*. *J. Insect Physiol.* 14, 741–764. doi: 10.1016/0022-1910(68)90186-8

- Dadd, R. H., Krieger, D. L., and Mittler, T. E. (1967). Studies on artificial feeding of the aphid *Myzus persicae* (Sulzer)-IV, requirements for water-soluble vitamins and ascorbic acid. *J. Insect Physiol.* 13, 249–272. doi: 10.1016/0022-1910(67)90152-7
- Douglas, A. E. (1988). Sulphate utilization in an aphid symbiosis. *Insect Biochem.* 18, 599–605. doi: 10.1016/0020-1790(88)90012-1
- Douglas, A. E. (1989). Mycetocyte symbiosis in insects. *Biol. Rev.* 64, 409–434. doi: 10.1111/j.1469-185X.1989.tb00682.x
- Douglas, A. E. (1995). The ecology of symbiotic micro-organisms. *Adv. Ecol. Res.* 26, 69–103. doi: 10.1016/S0065-2504(08)60064-1
- Douglas, A. E. (1998). Nutritional interactions in insect-microbial symbioses: aphids and their symbiosis bacteria *Buchnera*. *Annu. Rev. Entomol.* 43, 17–37. doi: 10.1146/annurev.ento.43.1.17
- Douglas, A. E., and Dixon, A. F. G. (1987). The mycetocyte symbiosis in aphids: variation with age and morph in virginoparae of *Megoura viciae* and *Acyrtosiphon pisum*. *J. Insect Physiol.* 33, 109–113. doi: 10.1016/0022-1910(87)90082-5
- Douglas, A. E., and Prosser, W. (1992). Synthesis of an essential amino acid tryptophan in the pea aphid (*Acyrtosiphon pisum*) symbiosis. *J. Insect Physiol.* 38, 565–568. doi: 10.1016/0022-1910(92)90107-O
- Febvay, G., Liadoze, I., Guillard, J., and Bonnet, G. (1995). Analysis of energetic necessary amino acid metabolism in *Acyrtosiphon pisum*: a multidimensional approach to amino acid metabolism on aphid. *Arch. Insect Biochem. Physiol.* 29, 45–69. doi: 10.1002/arch.940290106
- Hansen, A. K., and Moran, N. A. (2011). Aphid genome expression reveals host-symbiont cooperation in the production of amino acids. *Proc. Natl. Acad. Sci. U.S.A.* 108, 2849–2854. doi: 10.1073/pnas.1013465108
- Hardie, J. (1987). The photoperiodic control of wing development in the black bean aphid, *Aphis fabae*. *J. Insect Physiol.* 33, 543–549. doi: 10.1016/0022-1910(87)90068-0
- Hardie, J., and Leckstein, P. (2007). Antibiotics, primary symbionts and wing polyphenism in three aphid species. *Insect Biochem.* 37, 886–890. doi: 10.1016/j.ibmb.2007.05.005
- Hatano, E., Kunert, G., and Weisser, W. W. (2010). Aphid wing induction and ecological costs of alarm pheromone emission under field conditions. *PLoS ONE* 5:e11188. doi: 10.1371/journal.pone.0011188
- Hongoh, Y., and Ishikawa, H. (1994). Changes of mycetocyte symbiosis in response to flying behavior of alate aphid (*Acyrtosiphon pisum*). *Zool. Sci.* 11, 731–735.
- Johnson, B. (1966). Wing polymorphism in aphids III. The influence of the host plant. *Entomol. Exp. Appl.* 9, 213–222. doi: 10.1111/j.1570-7458.1966.tb02352.x
- Karan, D., and David, R. (2000). Cold tolerance in *Drosophila*: adaptive variations revealed by the analysis of starvation survival reaction norms. *J. Therm. Biol.* 25, 345–351. doi: 10.1016/S0306-4565(99)00106-0
- Lai, C. Y., Baumann, L., and Baumann, P. (1994). Amplification of trpEG: adaptation of *Buchnera aphidicola* to an endosymbiotic association with aphids. *Proc. Natl. Acad. Sci. U.S.A.* 91, 3819–3823. doi: 10.1073/pnas.91.9.3819
- Leckstein, P. M., and Liewellyn, M. (1973). Effect of dietary amino acids on the size and alary polymorphism of *Aphis fabae*. *J. Insect Physiol.* 19, 973–980. doi: 10.1016/0022-1910(73)90023-1
- Lees, A. D. (1966). The control of polymorphism in Aphids. *Adv. Insect Physiol.* 3, 207–277. doi: 10.1016/S0065-2806(08)60188-5
- Leonardo, T. E., and Mondor, E. B. (2006). Symbiont modifies host life-history traits that affect gene flow. *Proc. R. Soc. B.* 273, 1079–1084. doi: 10.1098/rspb.2005.3408
- Mehrpour, M., Zytynska, S. E., and Weisser, W. W. (2013). Multiple cues for winged morph production in an aphid Metacommunity. *PLoS ONE* 8:e58323. doi: 10.1371/journal.pone.0058323
- Mittler, T., and Kleinjan, J. E. (1970). Effect of artificial diet composition on wing-production by the aphid *Myzus persicae*. *J. Insect Physiol.* 16, 833–850. doi: 10.1016/0022-1910(70)90217-9
- Moran, N. A., and Jarvik, T. (2010). Lateral transfer of genes from fungi underlies carotenoid production in aphids. *Science* 328, 624–627. doi: 10.1126/science.1187113
- Moran, N. A., and Yun, Y. (2015). Experimental replacement of an obligate insect symbiont. *Proc. Natl. Acad. Sci. U.S.A.* 112, 2093–2096. doi: 10.1073/pnas.1420037112
- Moran, N., and Baumann, P. (1994). Phylogenetics of cytoplasmically inherited microorganisms of arthropods. *Trends. Ecol. Evol.* 9, 15–20. doi: 10.1016/0169-5347(94)90226-7
- Müller, C. B., Williams, I. S., and Hardie, J. (2001). The role of nutrition, crowding and interspecific interactions in the development of winged aphids. *Ecol. Entomol.* 26, 330–340. doi: 10.1046/j.1365-2311.2001.00321.x
- Nakabachi, A., and Ishikawa, H. (1999). Provision of riboflavin to the host aphid *Acyrtosiphon pisum*, by endosymbiotic bacteria, *Buchnera*. *J. Insect Physiol.* 45, 1–6. doi: 10.1016/S0022-1910(98)00104-8
- Parish, W. E. G., and Bale, J. S. (1990). Effects of short term exposure to low temperature on wing development in the grain aphid *Sitobion avenae* (F.) (Hem., Aphididae). *J. Appl. Entomol.* 109, 175–181. doi: 10.1111/j.1439-0418.1990.tb00034.x
- Prosser, W. A., and Douglas, A. E. (1991). The aposymbiotic aphid: an analysis of chlortetracycline-treated pea aphid, *Acyrtosiphon pisum*. *J. Insect Physiol.* 37, 712–719. doi: 10.1016/0022-1910(91)90104-8
- Rahbe, Y., Delobel, B., Febvay, G., and Chantegel, B. (1994). Aphid-specific triglycerides in symbiotic and aposymbiotic *Acyrtosiphon pisum*. *Insect Biochem. Mol. Biol.* 24, 95–101. doi: 10.1016/0965-1748(94)90127-9
- Russell, J. A., and Moran, N. A. (2006). Costs and benefits of symbiont infection in aphids: variation among symbionts and across temperatures. *Proc. R. Soc. B Biol. Sci.* 273, 603–610. doi: 10.1098/rspb.2005.3348
- Sandström, J., and Moran, N. A. (1999). How nutritionally imbalanced is phloem sap for aphids? *Entomol. Exp. Appl.* 91, 203–210. doi: 10.1046/j.1570-7458.1999.00485.x
- Sandström, J., and Pettersson, J. (1994). Amino acid composition of phloem sap and the relation to intraspecific variation in pea aphid *Acyrtosiphon pisum* performance. *J. Insect Physiol.* 40, 947–955. doi: 10.1016/0022-1910(94)90133-3
- Sasaki, T., Aoki, T., Hayashi, H., and Ishikawa, H. (1990). Amino acid composition of honeydew of symbiotic and aposymbiotic pea aphids *Acyrtosiphon pisum*. *J. Insect Physiol.* 36, 35–40. doi: 10.1016/0022-1910(90)90148-9
- Sasaki, T., Hayashi, H., and Ishikawa, H. (1991). Growth and reproduction of the symbiotic and aposymbiotic pea aphids *Acyrtosiphon pisum*. *J. Insect Physiol.* 31, 749–756. doi: 10.1016/0022-1910(91)90109-D
- Scarborough, C. L., Ferrari, J., and Godfray, H. C. J. (2005). Aphid protected from pathogen by endosymbiont. *Science* 310, 1781–1781. doi: 10.1126/science.1120180
- Shapiro, A. M. (1976). Seasonal polyphenism. *Evol. Biol.* 9, 259–333.
- Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y., and Ishikawa, H. (2000). Genome sequence of the endocellular bacterial symbiont of aphids *Buchnera* sp. *Nature* 407, 81–86. doi: 10.1038/35024074
- Simon, J. C., Boutin, S., Tsuchida, T., Koga, R., Le Gallic, J. F., Frantz, A., et al. (2011). Facultative symbiont infections affect aphid reproduction. *PLoS ONE* 6:e21831. doi: 10.1371/journal.pone.0021831
- Sutherland, O. R. W., and Mittler, T. E. (1971). Influence of diet composition and crowding on wing production by the aphid *Myzus persicae*. *J. Insect Physiol.* 17, 321–328. doi: 10.1016/0022-1910(71)90216-2
- Tamas, I., Klasson, L., Canbäck, B., Näslund, A. K., Eriksson, A.-S., Wernegreen, J. J., et al. (2002). 50 million years of genomic stasis in endosymbiotic bacteria. *Science* 296, 2376–2379. doi: 10.1126/science.1071278
- Tsuchida, T., Koga, R., Fujiwara, A., and Fukatsu, T. (2014). Phenotypic effect of “*Candidatus Rickettsiella viridis*”, a facultative symbiont of the pea aphid (*Acyrtosiphon pisum*), and its interaction with a coexisting symbiont. *Appl. Environ. Microbiol.* 80, 525–533. doi: 10.1128/AEM.03049-13
- Tsuchida, T., Koga, R., Horikawa, M., Tsunoda, T., Maoka, T., Matsumoto, S., et al. (2010). Symbiotic bacteria modifies aphid body color. *Science* 330, 1102–1104. doi: 10.1126/science.1195463
- van Ham, R. C. H. J., Kamerbeek, J., Palacios, C., Rausell, C., Abascal, F., Bastolla, U., et al. (2003). Reductive genome evolution in *Buchnera aphidicola*. *Proc. Natl. Acad. Sci. U.S.A.* 100, 581–586. doi: 10.1073/pnas.0235981100
- Weisser, W. W., Braendle, C., and Minoretto, N. (1999). Predator-induced morphological shift Predator-induced in the pea aphid. *Proc. R. Soc. Lond. B Biol. Sci.* 266, 1175–1181. doi: 10.1098/rspb.1999.0760
- Wernegreen, J. J., and Moran, N. A. (2001). Vertical transmission of biosynthetic plasmids in aphid endosymbionts (*Buchnera*). *J. Bacteriol.* 183, 785–790. doi: 10.1128/JB.183.2.785-790.2001

- Werren, J. H., Baldo, L., and Clark, M. E. (2008). Wolbachia: master manipulators of invertebrate biology. *Nat. Rev. Microbiol.* 6, 741–751. doi: 10.1038/nrmicro1969
- Wilkinson, T. L., and Douglas, A. E. (1995). Aphid feeding, as influenced by the disruption of the symbiotic bacteria. *J. Insect Physiol.* 41, 635–640. doi: 10.1016/0022-1910(95)00036-T
- Wilkinson, T. L., and Douglas, A. E. (1996). The impact of aposymbiosis on amino acid metabolism of pea aphids (*Acyrtosiphon pisum*). *Entomol. Exp. Appl.* 80, 279–282. doi: 10.1111/j.1570-7458.1996.tb00935.x
- Wilkinson, T. L., and Ishikawa, H. (2001). On the functional significance of symbiotic microorganisms in the Hemiptera: a comparative study of *Acyrtosiphon pisum* and *Nilaparvata lugens*. *Physiol. Entomol.* 26, 86–93. doi: 10.1046/j.1365-3032.2001.00214.x
- Wilson, E. O. (1971). *The Insect Societies*. Cambridge, MA: Harvard University Press.
- Xu, X. L., He, S. Q., and Wu, J. X. (2012). The effect of starvation and subsequent feeding on the reproductive potential of the grain aphid, *Sitibion avenae*. *Entomol. Exp. Appl.* 144, 294–300. doi: 10.1111/j.1570-7458.2012.01287.x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Zhang, Li, Zhang, Coates, Zhou and Cheng. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.