

Baryon acoustic oscillations in the Sloan Digital Sky Survey Data Release 7 galaxy sample

Will J. Percival,^{1*} Beth A. Reid,^{2,3,4} Daniel J. Eisenstein,⁵ Neta A. Bahcall,⁴ Tamas Budavari,⁶ Joshua A. Frieman,^{7,8} Masataka Fukugita,⁹ James E. Gunn,⁴ Željko Ivezić,¹⁰ Gillian R. Knapp,⁴ Richard G. Kron,¹¹ Jon Loveday,¹² Robert H. Lupton,⁴ Timothy A. McKay,¹³ Avery Meiksin,¹⁴ Robert C. Nichol,¹ Adrian C. Pope,¹⁵ David J. Schlegel,¹⁶ Donald P. Schneider,¹⁷ David N. Spergel,^{4,18} Chris Stoughton,¹⁹ Michael A. Strauss,⁴ Alexander S. Szalay,⁶ Max Tegmark,²⁰ Michael S. Vogeley,²¹ David H. Weinberg,²² Donald G. York^{11,23} and Idit Zehavi²⁴

¹*Institute of Cosmology and Gravitation, University of Portsmouth, Dennis Sciama building, Portsmouth P01 3FX*

²*Institute of Space Sciences (CSIC-IEEC), UAB, Barcelona 08193, Spain*

³*Institute for Sciences of the Cosmos (ICC), University of Barcelona, Barcelona 08028, Spain*

⁴*Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA*

⁵*Steward Observatory, University of Arizona, 933 N. Cherry Ave., Tucson, AZ 85121, USA*

⁶*Department of Physics and Astronomy, The Johns Hopkins University, 3701 San Martin Drive, Baltimore, MD 21218, USA*

⁷*Particle Astrophysics Center, Fermilab, PO Box 500, Batavia, IL 60510, USA*

⁸*Kavli Institute for Cosmological Physics, Department of Astronomy & Astrophysics, University of Chicago, Chicago, IL 60637, USA*

⁹*Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan*

¹⁰*Department of Astronomy, University of Washington Box 351580, Seattle, WA 98195, USA*

¹¹*Department of Astronomy and Astrophysics, The University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637, USA*

¹²*Astronomy Centre, University of Sussex, Falmer, Brighton BN1 9QH*

¹³*Departments of Physics and Astronomy, University of Michigan, Ann Arbor, MI 48109, USA*

¹⁴*SUPA; Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ*

¹⁵*Los Alamos National Laboratory, PO Box 1663, Los Alamos, NM 87545, USA*

¹⁶*Lawrence Berkeley National Lab, 1 Cyclotron Road, MS 50R5032, Berkeley, CA 94720, USA*

¹⁷*Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA*

¹⁸*Princeton Center for Theoretical Science, Princeton University, Jadwin Hall, Princeton, NJ 08542, USA*

¹⁹*Fermilab, PO Box 500, Batavia, IL 60510, USA*

²⁰*Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

²¹*Department of Physics, Drexel University, Philadelphia, PA 19104, USA*

²²*Department of Astronomy, The Ohio State University, Columbus, OH 43210, USA*

²³*Enrico Fermi Institute, University of Chicago, Chicago, IL 60637, USA*

²⁴*Department of Astronomy, Case Western Reserve University, Cleveland, OH 44106, USA*

Accepted 2009 October 1. Received 2009 September 23; in original form 2009 July 10

ABSTRACT

The spectroscopic Sloan Digital Sky Survey (SDSS) Data Release 7 (DR7) galaxy sample represents the final set of galaxies observed using the original SDSS target selection criteria. We analyse the clustering of galaxies within this sample, including both the luminous red galaxy and main samples, and also include the 2-degree Field Galaxy Redshift Survey data. In total, this sample comprises 893 319 galaxies over 9100 deg². Baryon acoustic oscillations (BAO) are observed in power spectra measured for different slices in redshift; this allows us to constrain the distance–redshift relation at multiple epochs. We achieve a distance measure at redshift $z = 0.275$, of $r_s(z_d)/D_V(0.275) = 0.1390 \pm 0.0037$ (2.7 per cent accuracy), where $r_s(z_d)$ is the comoving sound horizon at the baryon-drag epoch, $D_V(z) \equiv [(1+z)^2 D_A^2 cz/H(z)]^{1/3}$,

*E-mail: will.percival@port.ac.uk

$D_A(z)$ is the angular diameter distance and $H(z)$ is the Hubble parameter. We find an almost independent constraint on the ratio of distances $D_V(0.35)/D_V(0.2) = 1.736 \pm 0.065$, which is consistent at the 1.1σ level with the best-fitting Λ cold dark matter model obtained when combining our $z = 0.275$ distance constraint with the *Wilkinson Microwave Anisotropy Probe* 5-year (*WMAP5*) data. The offset is similar to that found in previous analyses of the SDSS DR5 sample, but the discrepancy is now of lower significance, a change caused by a revised error analysis and a change in the methodology adopted, as well as the addition of more data. Using *WMAP5* constraints on $\Omega_b h^2$ and $\Omega_c h^2$, and combining our BAO distance measurements with those from the Union supernova sample, places a tight constraint on $\Omega_m = 0.286 \pm 0.018$ and $H_0 = 68.2 \pm 2.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$ that is robust to allowing $\Omega_k \neq 0$ and $w \neq -1$. This result is independent of the behaviour of dark energy at redshifts greater than those probed by the BAO and supernova measurements. Combining these data sets with the full *WMAP5* likelihood constraints provides tight constraints on both $\Omega_k = -0.006 \pm 0.008$ and $w = -0.97 \pm 0.10$ for a constant dark energy equation of state.

Key words: cosmology: observations – distance scale – large-scale structure of Universe.

1 INTRODUCTION

‘What is the nature of dark energy?’ is one of the current key questions in physical science. Distinguishing between competing theories will only be achieved with precise measurements of the cosmic expansion history and the growth of structure within it. Among current measurement techniques for the cosmic expansion, baryon acoustic oscillations (BAO) appear to have the lowest level of systematic uncertainty (Albrecht et al. 2006).

BAO are a series of peaks and troughs, with a wavelength of approximately $0.06 h \text{ Mpc}^{-1}$ that are present in the power spectrum of matter fluctuations after the epoch of recombination, and on large scales. They occur because the primordial cosmological perturbations excite sound waves in the relativistic plasma of the early universe (Silk 1968; Peebles & Yu 1970; Sunyaev & Zel’dovich 1970; Bond & Efstathiou 1984, 1987; Holtzman 1989). Radiation pressure drives baryonic material away from the seed perturbations until the ionized material recombines at redshift $z \simeq 1000$. The momentum of the baryonic material means that the motion continues for a short time after recombination, until an epoch known as the baryon-drag epoch. The wavelength of the BAO is related to the comoving sound horizon at the baryon-drag epoch, which depends on the physical densities of matter $\Omega_m h^2$ and of baryons $\Omega_b h^2$ in the Universe. *Wilkinson Microwave Anisotropy Probe* 5-year (*WMAP5*) constraints on $\Omega_b h^2$ and $\Omega_m h^2$ (Komatsu et al. 2009) give that $r_s(z_d) \simeq 153.5 \text{ Mpc}$ (see Section 7 for details).

BAO occur on relatively large scales, which are still predominantly in the linear regime at present day; it is therefore expected that BAO should also be seen in the galaxy distribution (Goldberg & Strauss 1998; Meiksin, White & Peacock 1999; Seo & Eisenstein 2005; Springel et al. 2005; White 2005; Eisenstein, Seo & White 2007). We can therefore use BAO as standard rulers to constrain the expansion of the Universe if the comoving sound horizon at the baryon-drag epoch is known. The apparent size of the BAO measured from observations then leads to measurements of the Hubble parameter and the angular diameter distance (Bond & Glazebrook 2003; Seo & Eisenstein 2003; Matsubara 2004; Hu & Haiman 2006).

The acoustic signature has now been convincingly detected at low redshift (Percival et al. 2001; Cole et al. 2005; Eisenstein et al. 2005; Huetsi 2006) using the 2-degree Field Galaxy Redshift Sur-

vey (2dFGRS; Colless et al. 2003) and the Sloan Digital Sky Survey (SDSS; York et al. 2000). The detection has subsequently been refined using more data and better techniques, and is now producing competitive constraints on cosmological models. Tegmark et al. (2006) analysed the SDSS Data Release 4 (DR4; Adelman-McCarthy et al. 2006) luminous red galaxy (LRG) sample. Percival et al. (2007a,b) presented the power spectrum of the SDSS DR5 (Adelman-McCarthy et al. 2007) galaxy sample and considered the shape of the power spectrum and measured the matter density using the BAO features. Percival et al. (2007c) took this analysis a stage further by fitting the SDSS data, combined with the 2dFGRS, with models of the distance–redshift relation. Gaztanaga, Cabre & Hui (2008) and Sanchez et al. (2009) have also analysed the SDSS DR6 (Adelman-McCarthy et al. 2008) sample, obtaining cosmological constraints from the radial and spherically averaged BAO signal. In a recent analysis, Kazin et al. (2009) have calculated the correlation function of the SDSS DR7 (Abazajian et al. 2009) LRG sample, and have shown that their results agree with those presented in our paper. Two studies have also considered the clustering of the LRGs at high redshift within the SDSS survey, using photometric redshifts to estimate galaxy distances (Blake et al. 2007; Padmanabhan et al. 2007).

In this paper, we analyse the clustering of galaxies in the spectroscopic SDSS DR7 sample, including both LRG and main galaxy samples, combined with the 2dFGRS, and measure the BAO signal in a series of redshift slices. SDSS DR7 marks the final release of galaxies observed using the standard SDSS targeting algorithm, and the sample we analyse covers a solid angle of 7930 deg^2 , including a 7190 deg^2 contiguous region in the North Galactic Cap. The Baryon Oscillation Spectroscopic Survey (BOSS; Schlegel, White & Eisenstein 2009a), part of the SDSS-III project, will adopt a different targeting algorithm, focusing on galaxies and quasars at higher redshifts.

The observed amplitude of the large-scale galaxy clustering depends on both galaxy colour and luminosity (Tegmark et al. 2004; Zehavi et al. 2005; Swanson et al. 2008). Using the SDSS DR5 sample, Cresswell & Percival (2009) showed that for blue galaxies, the deviation in the shape of the galaxy power spectrum from the linear matter power spectrum at $k > 0.1 h \text{ Mpc}^{-1}$ is a strong function of luminosity, while it is almost constant for red galaxies. It is therefore difficult to extract the underlying matter power spectrum

from a galaxy power spectrum measured for a population of galaxies where the distribution of galaxy colours and luminosities changes with spatial location, such as that provided by a magnitude-limited catalogue. In contrast, the LRG population, which comprises the high-redshift part of the sample analysed here, has a simpler relation with the matter field, in that there is a single galaxy population to consider (Reid, Spergel & Bode 2009a). In a companion paper (Reid et al. 2009b), we apply a grouping algorithm to recover the halo power spectrum from the LRGs and then calibrate the relation of the halo power spectrum to the linear theory power spectrum using simulations. We are then able to extract cosmological information from the large-scale shape of the power spectrum in addition to the BAO signal, though the constraints are more tightly embedded in the assumed cosmological framework.

BAO in the galaxy power spectrum are only weakly affected by the effects of non-linear structure formation and scale-dependent galaxy bias, because they are on such large scales. The primary consequence is a damping on small scales, which can be well approximated by a Gaussian smoothing (Bharadwaj 1996; Crocce & Scoccimarro 2006, 2008; Eisenstein et al. 2007; Matsubara 2008a,b). The observed BAO, defined as the ratio of the observed power spectrum P_{obs} to a smooth fit to this power P_{nw} , $BAO_{\text{obs}} \equiv P_{\text{obs}}/P_{\text{nw}}$, are related to the original BAO in the linear matter power spectrum BAO_{lin} , defined similarly, by

$$BAO_{\text{obs}} = G_{\text{damp}} BAO_{\text{lin}} + (1 - G_{\text{damp}}), \quad (1)$$

where $G_{\text{damp}} = \exp(-\frac{1}{2}k^2 D_{\text{damp}}^2)$, and the damping scale D_{damp} is set to $10 h^{-1}$ Mpc for redshift-space power spectra at $z \simeq 0.3$ (Eisenstein et al. 2007). This damping of the linear power is a relatively benign effect as it does not affect the positions of the BAO, although it does reduce the signal available. Additional, more pernicious effects such as the mixing of modes in the power spectrum can generate shifts in the BAO position (Crocce & Scoccimarro 2008); for biased tracers, these offsets can be at the per cent level (Smith, Scoccimarro & Sheth 2007), and are therefore important as we wish to make per cent level distance measurements.

In our analysis, we measure BAO relative to a model that allows for smooth changes in the underlying shape of the power spectrum, which alleviates some of this shift. Physical models of BAO positions in observed redshift-space power spectra relative to such a fitted smooth model (Crocce & Scoccimarro 2008; Smith, Scoccimarro & Sheth 2008; Padmanabhan & White 2009; Sanchez et al. 2009) and numerical simulations (Angulo et al. 2008; Seo et al. 2008; Kim et al. 2009) suggest that we should expect residual shifts at the sub-per-cent level. These are below the precision of current experiments; for example, in this paper we present a BAO distance scale measurement with 2.7 per cent accuracy. Therefore, we adopt a procedure that allows for the damping as well as smooth changes in the underlying shape of the power spectrum, but no more. The analysis of future surveys, which will lead to tighter distance-redshift constraints, will clearly also have to allow for non-linear effects, either by physical modelling, by simulations or by using methods which attempt to reconstruct the initial fluctuation field (Eisenstein, Seo & White 2007; Seo et al. 2008; Padmanabhan, White & Cohn 2008b).

The SDSS and 2dFGRS data are discussed in Sections 2 and 3, respectively. The basic methodology, presented in Section 4, is similar to that of Percival et al. (2007c), although we have revised the calculation of the window function to increase the computational speed. We also perform an extensive test of the derived errors, running mock catalogues through our full analysis pipeline to test the confidence intervals quoted (Section 5). Results are presented

in Sections 6 and 7, tested for robustness in Section 8 and placed in a cosmological context in Section 9. A comparison with our DR5 analyses is given in Section 10 and we finish with a discussion in Section 11.

In this paper, we use the standard cosmological parameters. For flat Λ cold dark matter (Λ CDM) models, these are the Hubble constant H_0 , the densities of baryonic matter Ω_b , cold dark matter Ω_c , all matter Ω_m and dark energy Ω_Λ . Going beyond this simple class of models, we use the equation of state of the dark energy w , the curvature energy density Ω_k and total energy density Ω_{tot} . When combining with information from the cosmic microwave background (CMB), we also consider some parameters that are not constrained by the BAO: τ is the optical depth to re-ionization, n_s is the scalar spectral index and A_{05} is the amplitude of curvature perturbations at $k = 0.05 \text{ Mpc}^{-1}$.

2 THE DATA

The SDSS-I and SDSS-II projects used a 2.5 m telescope (Gunn et al. 2006), to obtain imaging data in five passbands u , g , r , i and z (Fukugita et al. 1996; Gunn et al. 1998). The images were reduced (Lupton et al. 2001; Stoughton et al. 2002; Pier et al. 2003; Ivezić et al. 2004) and calibrated (Lupton, Gunn & Szalay 1999; Hogg et al. 2001; Smith et al. 2002; Tucker et al. 2006), and galaxies were selected in two ways for follow-up spectroscopy. The main galaxy sample (Strauss et al. 2002) targeted galaxies brighter than $r = 17.77$ (approximately 90 per deg², with a weighted median redshift of $z = 0.10$). The DR7 sample (Abazajian et al. 2009) used in our analysis includes 669 905 main galaxies (Strauss et al. 2002) with a median redshift of $z = 0.12$, selected to a limiting Galactic extinction-corrected Petrosian magnitude $r < 17.77$ or $r < 17.5$ in a small subset of the early data from the survey. The effect of the inclusion of the early SDSS data is tested in Section 8.2. In addition, our sample includes 80 046 LRGs (Eisenstein et al. 2001), which form an extension of the SDSS spectroscopic survey to higher redshifts $0.2 < z < 0.5$. Of the main galaxies, 30 530 are also classified as LRGs and are intrinsically luminous with $M_{0.1,r} < -21.8$, where $M_{0.1,r}$ is the Galactic extinction and K -corrected r -band absolute galaxy magnitude. We apply this requirement to all of our LRGs, so our sample includes 110 576 LRGs in total, with a weighted median redshift of $z = 0.31$. Although the main galaxy sample contains significantly more galaxies than the LRG sample, the LRG sample covers more volume. Redshift distributions for these samples are shown in fig. 2 of Percival et al. (2007b). In our default analysis, we use SDSS Petrosian magnitudes calibrated using the ‘uber-calibration’ method (Padmanabhan et al. 2008a), although we test against data calculated using the original calibration methodology (Tucker et al. 2006). Where specified, we have K -corrected the galaxy luminosities using the methodology outlined by Blanton et al. (2003a,b). Further details of the cuts applied to the data can be found in Percival et al. (2007b).

Due to the finite size of the fibres, spectra cannot be obtained for both objects in a pair closer than 55 arcsec, within a single spectroscopic tile. Tiling (Blanton et al. 2003a) deals with this to some extent by allowing plate overlaps to provide multiple observations of crowded regions. Even so, not all galaxies in such regions which meet the target selection criteria could be observed. Zehavi et al. (2002) corrected for this undersampling by assigning the redshift of the nearest observed galaxy to a galaxy which was not observed due to crowding, and showed that this provides sufficient correction for large-scale structure studies. We apply this correction in the present

work and test it to show that our results are insensitive to this in Section 8.1.

In order to increase the volume covered at redshift $z < 0.3$, we include 143 368 galaxies from the 2dFGRS sample. These galaxies, selected to an extinction-corrected magnitude limit of approximately $b_J = 19.45$ (Colless et al. 2003) from regions of sky not covered by the SDSS sample, cover two contiguous regions totalling $\sim 1200 \text{ deg}^2$. They do not include the 2dFGRS random fields, a set of 99 random 2° fields spread over the full Southern Galactic Cap, as these would complicate the window function. The galaxies cover $0 < z < 0.3$, with a weighted median at $z = 0.17$. The redshift distribution of the sample was analysed as in Cole et al. (2005) for $0 < z < 0.3$, and we use the same synthetic catalogues to model the unclustered expected galaxy distribution within the reduced sample.

We assume that each galaxy is biased with a linear deterministic bias model and that this bias depends on $M_{0.1r}$ according to Tegmark et al. (2004) and Zehavi et al. (2005). All galaxies were weighted using this model so the fluctuation amplitudes match those of L_* galaxies, where L_* was calculated separately for the SDSS and 2dFGRS. We include an extra normalization factor to the 2dFGRS galaxy bias model to correct the relative bias of L_* galaxies in the different surveys. This was calculated by matching the normalization of the 2dFGRS and SDSS bias-corrected power spectra for $k < 0.1 h \text{ Mpc}^{-1}$. In principle, we could have added information on galaxy bias from the BAO, since the small-scale damping (see equation 1) depends on how strongly non-linear the underlying dark matter density fluctuations are. As we show in Section 8.6, this information is limited for the current data, but future surveys may be able to exploit changes in this damping as a function of galaxy properties, such as colour and luminosity.

3 SPLITTING INTO SUB-SAMPLES

In order to probe the distance–redshift relation in detail, ideally we would analyse BAO measured in many independent redshift slices. However, if the slices are too narrow in redshift, then there is insufficient signal and the BAO cannot be recovered with sufficient accuracy to give a likelihood with close to a Gaussian distribution (see the discussion in Section 5). If the slices are too wide, or too many overlapping slices are chosen, then the covariance matrix becomes close to singular, potentially leading to numerical instability. In order to balance these competing requirements, we have chosen to analyse the redshift slices presented in Table 1. The power spectra will be correlated, and these correlations, together with correlations of $P(k)$ values at different k within each redshift slice, are included in the covariance matrices in our analysis. Note that we include

Table 1. Parameters of the redshift intervals analysed.

Slice	z_{\min}	z_{\max}	N_{gal}	V_{eff}	\bar{n}
1	0.0	0.5	895 834	0.42	128.1
2	0.0	0.4	874 330	0.38	131.2
3	0.0	0.3	827 760	0.27	138.3
4	0.1	0.5	505 355	0.40	34.5
5	0.1	0.4	483 851	0.36	35.9
6	0.2	0.5	129 045	0.27	1.92
7	0.3	0.5	68 074	0.15	0.67

Note. V_{eff} is given in units of $h^{-3} \text{ Gpc}^3$, and was calculated as in equation (2) using an effective power spectrum amplitude of $\bar{P} = 10^4 h^{-3} \text{ Mpc}^3$, appropriate on scales $k \sim 0.15 h \text{ Mpc}^{-1}$ for a population with bias $b = 1.7$. The average galaxy number density in each bin \bar{n} is in units of $10^{-4} (h^{-1} \text{ Mpc})^3$.

slice 7, for which the effective volume is relatively small, because of the interesting redshift range covered.

As well as giving the redshift limits of the slices in Table 1, we also give the number of galaxies in each including both the 2dFGRS and the SDSS, and the effective volume, calculated from the integral (Feldman, Kaiser & Peacock 1994)

$$V_{\text{eff}} = \int d^3r \left[\frac{\bar{n}(\mathbf{r})\bar{P}}{1 + \bar{n}(\mathbf{r})\bar{P}} \right]^2, \quad (2)$$

where $\bar{n}(\mathbf{r})$ is the observed comoving number density of the sample at location \mathbf{r} and \bar{P} is the expected power spectrum amplitude. To calculate V_{eff} for our redshift slices, distances were calculated assuming a fiducial flat Λ CDM cosmology with $\Omega_m = 0.25$. For the numbers given in Table 1, we fix $\bar{P} = 10^4 h^{-3} \text{ Mpc}^3$, appropriate on scales $k \sim 0.15 h \text{ Mpc}^{-1}$ for a population with bias $b = 1.7$. For comparison, Eisenstein et al. (2005) analyse a sample with $V_{\text{eff}} = 0.13 h^{-3} \text{ Gpc}^3$, approximately a third of the effective volume of slice 1.

We fit models to three sets of power spectra as follows.

- (i) We fit a single power spectrum for the SDSS LRG sample covering $0.15 < z < 0.5$.
- (ii) We fit three power spectra for slices 1, 3 and 6 approximately corresponding to the procedure adopted by Percival et al. (2007c). Although we now use slices constrained by redshift rather than galaxy type, the $0 < z < 0.3$ slice is dominated by SDSS main galaxies, while the $0.2 < z < 0.5$ slice is dominated by LRGs.
- (iii) We fit six power spectra for slices 2–7, which allows a test of the distance–redshift relation at greater resolution.

We consider option (i) to tie in with the analysis presented by Reid et al. (2009b) and to demonstrate the effect of collapsing the clusters in redshift space where we try to reconstruct the halo power spectrum. Option (ii) is close to the approach of Percival et al. (2007c), where the SDSS main galaxies and 2dFGRS galaxies were analysed separately from the SDSS LRGs. Option (iii) allows us to see if there is more information available beyond measuring the distance–redshift relation at two redshifts. The slices do overlap in redshift, but we will properly take into account the covariance between the results when we fit to cosmological parameters.

4 BASIC METHODOLOGY

Power spectra were calculated for each catalogue using the Fourier method of Feldman et al. (1994), as applied by Percival et al. (2007b). In this method, a weighted galaxy over-density field is defined and Fourier transformed and then the spherically averaged power is measured. We use the luminosity-dependent galaxy weights advocated by Percival, Verde & Peacock (2004), as described in Section 2. To construct the over-density field, we need to quantify the expected spatial distribution of galaxies, in the absence of clustering. The standard method for this is to use an unclustered random catalogue, which matches the galaxy selection. To calculate this random catalogue, we fitted the redshift distributions of the galaxy samples with a spline fit (Press et al. 1992), and the angular mask was determined using a routine based on a HEALPIX (Górski et al. 2005) equal-area pixelization of the sphere (Percival et al. 2007b). Percival et al. (2007b) used a random catalogue containing 10 times as many points as galaxies. For the sparse LRGs, this approach induces significant shot noise, so we now use 100 times as many random points as LRGs. We have also increased the resolution at which the radial distribution of galaxies is quantified, now using a spline fit (Press et al. 1992) with nodes separated by

$\Delta z = 0.0025$. As an alternative to this radial selection, we could have simply adopted the redshift of a randomly chosen galaxy for each of our points in the random catalogue. In Section 8.7, we show that these two possibilities give consistent results.

Galaxy redshifts were converted to distances using a fiducial cosmology (flat Λ CDM model with $\Omega_m = 0.25$). For each distance–redshift model to be tested, we do not recalculate the power spectrum, but instead change the interpretation of the power spectrum computed assuming the fiducial Λ CDM galaxy distances. We do this through a window function, which relates the true and measured power spectra. This follows the procedure adopted by Percival et al. (2007c), but we now use a revised, computationally less intensive method for calculating the windows, as described in Appendix A.

A model of the BAO was created by fitting a linear matter power spectrum, calculated using CAMB (Lewis, Challinor & Lasenby 2000), which numerically solves the Boltzman equation describing the physical processes in the Universe before the baryon-drag epoch, with a cubic spline to remove the broad shape of the power, leaving the oscillations. The theoretical BAO were then damped with a Gaussian model as in equation (1), following the simulation results of Eisenstein et al. (2007). For our default fits, we assume that the damping scale $D_{\text{damp}} = 10 h^{-1} \text{Mpc}$ (Eisenstein et al. 2007), but we also consider fits where this scale is varied (Section 8.6). As discussed in Section 1, we do not attempt to correct for any shift induced by non-linear physics, because they are expected to be at a level below our statistical error.

The power spectrum measured from the data was fitted by a model constructed by multiplying this BAO model with a cubic spline (Press et al. 1992), which enables the power spectrum model to match the overall shape of the data power spectrum. Each power spectrum model was then convolved with a window function that corrects for both the survey geometry and the differences between our fiducial cosmological model used to convert redshift to distances and the cosmological model to be tested (see Appendix A). The free parameters of the model are the nine nodes of the cubic spline fixed empirically at $k = 0.001$, and $0.025 \leq k \leq 0.375$ with $\Delta k = 0.05$, and the parametrization of $D_V(z)$ used to calculate the correct window function. The spline nodes were refitted for every cosmology [or $D_V(z)$] tested. A power spectrum model with this spline node separation was tested by fitting many mock power spectra by Percival et al. (2007a) and was shown to match these without leaving significant residuals in the measured ‘shift’ between BAO in the model and data power spectra. This approach was also considered by Sanchez, Baugh & Angulo (2008), who found that it did not induce a bias in the recovered BAO constraints.

For a redshift survey in a thin shell, the position of the BAO approximately constrains $d_z \equiv r_s(z_d)/D_V(z)$, where $r_s(z_d)$ is the comoving sound horizon at the baryon-drag epoch, $D_V(z) \equiv [(1+z)^2 D_A^2 c z / H(z)]^{1/3}$ (Eisenstein et al. 2005; Percival et al. 2007c), D_A is the angular diameter distance and $H(z)$ is the Hubble parameter. We see that, although our power spectrum fitting procedure measures $D_V(z)$ for a fixed BAO model, we should consider the constraints as measurements of d_z , with $r_s(z_d)$ calculated for the flat Λ CDM model for which we created the BAO model, $r_s(z_d) = 111.4 h^{-1} \text{Mpc} = 154.7 \text{Mpc}$, using equation (6) of Eisenstein & Hu (1998), and assuming $h = 0.72$, $\Omega_b h^2 = 0.0223$ and $\Omega_m = 0.25$. This value of $r_s(z_d)$ is only used to index this model: as described above, the actual BAO model was calculated from a power spectrum predicted by CAMB. If the constraints provided in this paper are to be used to constrain a set of models where $r_s(z_d)$ for this fiducial model is calculated in a different way (i.e. not using equation 6 of




Figure 1. Average power spectra recovered from the LN catalogues (solid lines) compared with the data power spectra (solid circles with 1σ errors) for the six samples in Table 1. The errors on the data were calculated from the diagonal elements of the covariance matrix calculated from these LN catalogues. The power spectra have been offset by 0.5 dex for clarity, with the upper power spectrum having the correct normalization.

Eisenstein & Hu 1998), then our constraints should be adjusted to match.

The comoving distance–redshift relation is modelled as a cubic spline in the parameter $D_V(z)$. We consider models for $D_V(z)$ with two nodes at $z = 0.2$ and $z = 0.35$ or with four nodes at $z = 0.1$, $z = 0.2$, $z = 0.3$ and $z = 0.45$. Results are presented as constraints on d_z . The error between cubic spline fits to $D_V(z)$ with two nodes at $z = 0.2$ and $z = 0.35$, to the Λ CDM distance–redshift relations was shown in fig. 1 of Percival et al. (2007c), and is < 1 per cent for a flat Λ CDM cosmology with $\Omega_m = 0.25$ at $z \geq 0.15$.

Power spectra are shown in Fig. 1 for the redshift slices described in Section 3, each measured for 70 band powers equally spaced in $0.02 < k < 0.3 h \text{Mpc}^{-1}$. We see that the power spectra from the different redshift intervals are remarkably consistent, with $P(k)$ decreasing almost monotonically to small scales.

In order to calculate the covariances between the data, we have created 10 000 lognormal (LN) density fields (Coles & Jones 1991; Cole et al. 2005) from which we have drawn overlapping catalogues for each of our seven redshift slices. Catalogues were calculated on a $(512)^3$ grid with a box length of $4000 h^{-1} \text{Mpc}$. Unlike N -body simulations, these mock catalogues do not model the growth of structure, but instead return a density field with an LN distribution, similar to that seen in the real data. The window functions for these catalogues were matched to that of the 2dFGRS+SDSS catalogue with the original calibration. The input power spectrum was a cubic spline fit matched to the data power spectra (i.e. the smooth part of our standard model), multiplied by our default damped Λ CDM BAO model calculated using CAMB (Lewis et al. 2000). The LN power spectra were used to determine a covariance matrix between slices and for different band powers in each slice, assuming that the band powers were drawn from a multivariate Gaussian distribution.

The average recovered power spectra for each redshift interval are compared with the data power spectra in Fig. 1. Clearly, the general shape of the average power spectra of the LN catalogues is well matched to that recovered from the data. Using the inverse of this covariance matrix, we estimate the likelihood of each model assuming that the power spectra band powers for $0.02 < k < 0.3 h \text{ Mpc}^{-1}$ were drawn from a multivariate Gaussian distribution.

5 TESTING THE ANALYSIS METHOD WITH MOCK DATA

5.1 The model fit

We now consider using a subset of our LN catalogues to test our analysis procedure. For 1000 of the mock catalogues, we fit spline \times BAO models to extract distance constraints from the BAO, as described in Section 4. A small average shift of 1.3 per cent in the BAO scale was recovered between the power recovered from the LN catalogues and the input power spectrum used to create them. If we correct the 1000 power spectra measured from the LN catalogues by multiplying each power spectrum by the expected power divided by the average recovered power spectrum, then the average shift drops below 0.3 per cent, well within 1σ .

To test the origin of the observed 1.3 per cent shift, we have also drawn 1000 power spectrum realizations from a multivariate Gaussian distribution with covariance and mean matched to those of the data. These mock catalogues were fitted using the procedure described in Section 4. No shift in the BAO position was found from the fits to these catalogues, within the statistical limits of the analysis (~ 0.3 per cent). The distribution of recovered distance constraints was well matched to that recovered from fitting the corrected LN power spectra. Thus, the 1.3 per cent shift described above must be due to the LN procedure itself. The expected shift is dependent on the statistic used to measure the BAO position. The LN correlation function ξ_{LN} and Gaussian correlation function $\xi_G(r)$ of a field with the same power spectrum but with Gaussian statistics are related by $1 + \xi_{\text{LN}}(r) = \exp[\xi_G(r)]$. If we had used the peak in the correlation function as our standard ruler then, for the LN catalogues, we would have expected no BAO shift. However, the same is not true of our BAO \times spline model fitting procedure, which fits the BAO in the power spectrum over a range of scales.

Numerical simulations offer a better way to model the true Universe, and recent results from simulations show that we should expect a less significant shift between the BAO positions in the linear matter and galaxy power spectra than the 1.3 per cent shift found for the LN catalogues (Seo & Eisenstein 2003; Springel et al. 2005; Seo & Eisenstein 2007; Angulo et al. 2008). The exact shift required for the catalogues we analyse is not well constrained by these simulation results, and we consequently do not alter our analysis to include such a shift.

5.2 The likelihood surface

We use the Gaussian and LN power spectra samples to assess the nature of the likelihood for the BAO scale recovery. We consider fits to either three or six power spectra as described in Section 3, parametrizing $D_V(z)$ with a cubic spline with two non-zero nodes at $z = 0.2$ and $z = 0.35$, respectively. For each of the 1000 fits, we have measured the difference between the maximum likelihood value and the likelihood at the parameters of the true cosmological model. The fraction of samples with $-2 \ln \mathcal{L}/\mathcal{L}_{\text{true}} < 2.3$, 6.0, 9.3, corresponding to 68, 95 and 99 per cent confidence intervals, is given in Table 2. We find that in order to match the expected num-

Table 2. Fraction of fits to the LN power spectra in which the ratio of the likelihood maximum to the likelihood for the true cosmological model is less than the given limit. For a Gaussian likelihood, these limits correspond to 68, 95 and 99 per cent confidence intervals. We show results where we have corrected the errors as described in the text by multiplying the band-power errors by 1.14 for three redshift slices and 1.21 for six redshift slices.

$-2 \ln \mathcal{L}/\mathcal{L}_{\text{true}}$	Fraction of samples			
	Three slices		Six slices	
	Standard	Revised	Standard	Revised
< 2.3	0.579	0.666	0.551	0.667
< 6.0	0.892	0.946	0.862	0.948
< 9.3	0.966	0.983	0.955	0.981

bers of samples with likelihoods within the standard 1σ Gaussian confidence intervals, we must increase the errors on the power spectrum band powers by 14 ± 2 per cent if we fit to three power spectra. For fits to six power spectra, we must increase the errors by 21 ± 2 per cent in order to match the expected 1σ Gaussian confidence intervals. Although in this paper we do not consider fitting to a single power spectrum, we have repeated this analysis for BAO fits to the LRG sample of Reid et al. (2009b), and found that we must increase the errors on the power spectrum band powers by 10 ± 2 per cent to match the expected confidence intervals.

Because the same increase in the confidence intervals is required for both LN and Gaussian mock catalogues, this change must be caused by the methodology of fitting BAO, rather than the Gaussian to LN density field transition. In fact, we believe that it is caused by the non-Gaussian nature of the likelihood surface. We should expect the likelihood surface to be non-Gaussian to some extent in any case because there is a minimum in the likelihood where the observed and model BAO are perfectly out of phase in k -space: this represents the worst possible match between data and model. Adjusting the covariance matrix to match the distribution of best-fitting distance scales to the expected 68 per cent confidence interval does not quite match the 95 or 99 per cent confidence intervals, although it corrects for most of the difference. This shows that the confidence intervals cannot perfectly match those for a Gaussian distribution.

To test this further, we have created a set of 1000 Gaussian power spectrum realizations with errors that are 10 per cent of those in our standard sample. For these catalogues, the distribution of best-fitting $D_V(z)$ matches that expected from the likelihood distribution under Gaussian assumptions. No correction is required, and the likelihood distribution is much closer to that for a multivariate Gaussian distribution around the likelihood maximum. Thus, the requirement to increase the errors on the data disappears when we fit less noisy data, as we would expect if it is caused by fitting noisy data, which is giving a non-Gaussian likelihood surface.

The average likelihood surfaces measured from our 1000 fits to sets of three power spectra and six power spectra drawn from LN catalogues are shown in Fig. 2. We also plot the centre of the local likelihood maxima nearest to the input cosmological parameters for each model. The fractions of points within each contour are given in Table 2: the errors on the power spectrum band powers have been adjusted for each plot as described above so that ~ 68 per cent of the points lie within the $-2 \ln \mathcal{L} = 2.3$ contour.

6 RESULTS

BAO are observed in the power spectra recovered from all redshift slices of the SDSS+2dFGRS sample described in Section 3, and




Figure 2. Average likelihood contours recovered from the analysis of three power spectra (top panel) and six power spectra (bottom panel) measured from 1000 LN density fields. Contours are plotted for $-2\ln\mathcal{L} = 2.3, 6.0, 9.2$, corresponding to two-parameter confidence of 68, 95 and 99 per cent for a Gaussian distribution. Contours were calculated after increasing the errors on the power spectrum band powers as described in the text. Solid circles mark the locations of the likelihood maxima closest to the true cosmology. We have plotted the likelihood surface as a function of $D_V(z)/\text{Mpc}$, for fixed $r_s(z_d) = 154.7 \text{ Mpc}$, to show distance errors if the comoving sound horizon is known perfectly. The values of D_V for our input cosmology are shown by the vertical and horizontal solid lines.

are shown in Fig. 3, where we plot the measured power spectra divided by the spline component of the best-fitting model. In our default analysis we fit power spectra from six redshift slices as described in Section 3, using a spline for $D_V(z)$ with two nodes at $z = 0.2$ and $z = 0.35$, respectively. We assume a fixed BAO damping scale of $D_{\text{damp}} = 10 h^{-1} \text{ Mpc}$ and fit to all SDSS and non-overlapping 2dFGRS data. The effect of these assumptions is considered in Section 8. The resulting likelihood surface is shown in Fig. 4 as a function of $D_V(z) \text{ Mpc}^{-1}$, for fixed $r_s(z_d) = 154.7 \text{ Mpc}$, to show distance errors if the comoving sound horizon is known perfectly. The constraints should be considered measurements of $r_s(z_d)/D_V(z)$ (see Section 4). Fig. 4 reveals a dominant likelihood maximum close to the parameters of a ΛCDM cosmology with




Figure 3. BAO recovered from the data for each of the redshift slices (solid circles with 1σ errors). These are compared with BAO in our default ΛCDM model (solid lines).

$\Omega_m = 0.25$, $h = 0.72$ and $\Omega_b h^2 = 0.0223$. There are also weaker secondary maxima at lower $D_V(0.2)$, which are considered further in Section 8.8. The significance of the detection of BAO corresponds to $\Delta\chi^2 = 13.1$, which is approximately 3.6σ . As this is relative to an arbitrary smooth model, this test is more general, and hence the significance cannot be directly compared with results presented by Eisenstein et al. (2005).

We have matched the likelihood surface shown in Fig. 4 around the dominant maximum to a multivariate Gaussian model. Using




Figure 4. Likelihood contour plots for fits of two $D_V(z)$ cubic spline nodes at $z = 0.2$ and $z = 0.35$, calculated for our default analysis using six power spectra, uber-calibration, a fixed BAO damping scale of $D_{\text{damp}} = 10 h^{-1} \text{Mpc}$ and for all SDSS and non-overlapping 2dFGRS data. Solid contours are plotted for $-2 \ln \mathcal{L}/\mathcal{L}_{\text{true}} < 2.3, 6.0, 9.3$, which for a multivariate Gaussian distribution with two degrees of freedom correspond to 68, 95 and 99 per cent confidence intervals. Likelihoods were adjusted to match these Gaussian confidence intervals as described in Section 5. We have plotted the likelihood surface as a function of $D_V(z)/\text{Mpc}$, for fixed $r_s(z_d) = 154.7 \text{Mpc}$, to show distance errors if the comoving sound horizon is known perfectly. We also show a multivariate Gaussian fit to this likelihood surface (dashed contours). The values of D_V for a flat ΛCDM cosmology with $\Omega_m = 0.25$, $h = 0.72$ and $\Omega_b h^2 = 0.0223$ are shown by the vertical and horizontal solid lines.

this Gaussian fit, we find that the best-fitting model has

$$\begin{aligned} d_{0.2} &= 0.1905 \pm 0.0061 \quad (3.2 \text{ per cent}), \\ d_{0.35} &= 0.1097 \pm 0.0036 \quad (3.3 \text{ per cent}), \end{aligned} \quad (3)$$

where $d_z \equiv r_s(z_d)/D_V(z)$. These results are correlated with the correlation coefficient $r = 0.337$. For a cosmological distance-redshift model with \hat{d}_z the likelihood can be well approximated by a multivariate Gaussian with the covariance matrix

$$C \equiv \begin{pmatrix} \langle \Delta d_{0.2} \Delta d_{0.2} \rangle & \langle \Delta d_{0.2} \Delta d_{0.35} \rangle \\ \langle \Delta d_{0.35} \Delta d_{0.2} \rangle & \langle \Delta d_{0.35} \Delta d_{0.35} \rangle \end{pmatrix}, \quad (4)$$

where $\Delta d_z \equiv d_z - \hat{d}_z$. C has inverse:

$$C^{-1} = \begin{pmatrix} 30124 & -17227 \\ -17227 & 86977 \end{pmatrix}. \quad (5)$$

Without correcting the covariance matrix using the results from fitting to the LN power spectra as described in Section 5, the original average errors on $d_{0.2}$ and $d_{0.35}$ were 0.0051 and 0.0029, 16 and 24 per cent lower than those in equation (3), respectively. Compare with the band-power errors which were increased by 21 per cent, and we see that there is no direct relation between changes in the band-power errors and errors on d_z , because of the non-linear nature of the fit.

We diagonalize the covariance matrix of $d_{0.2}$ and $d_{0.35}$ to get quantities x and y :

$$\begin{pmatrix} x \\ y \end{pmatrix} \equiv \begin{pmatrix} 1 & 1.76 \\ -1 & 1.67 \end{pmatrix} \begin{pmatrix} d_{0.2} \\ d_{0.35} \end{pmatrix}, \quad (6)$$

which gives

$$x = 0.3836 \pm 0.0102 \quad (7)$$

$$y = -0.0073 \pm 0.0070. \quad (8)$$

The distance ratio $f \equiv D_V(0.35)/D_V(0.2)$ is given by

$$f = \frac{1.67 - 1.76y/x}{1 + y/x} \simeq 1.67 - 8.94y, \quad (9)$$

where the last approximation neglects the small variations around the best-fitting value of $x = 0.3836$; these would come to 0.002 in f , which is well within the errors. Thus, x is a measurement of distance for the concordance cosmology and y is the deviation from the concordance distance ratio: x is measured to about 2.7 per cent and y is consistent with zero to within about 1σ .

To high accuracy, the constraint x can be written as a constraint on the distance to some redshift $0.2 < z < 0.35$. In fact, $r_s(z_d)/D_V(0.275)$ predicts $x = d_{0.2} + 1.76d_{0.35}$ to a peak-to-peak precision of 0.04 per cent over the range $0.05 < \Omega_m < 1$ (assuming a flat cosmology with $w = -1$). Thus, we can quote the x measurement as a measurement of $d_{0.275}$ and the y measurement as a statistically independent measure of f .

For the best-fitting solution we have $d_{0.275} = 0.362x$, giving

$$d_{0.275} = 0.1390 \pm 0.0037 (2.7 \text{ per cent}). \quad (10)$$

We also have the statistically independent constraint

$$f \equiv D_V(0.35)/D_V(0.2) = 1.736 \pm 0.065. \quad (11)$$

$f = 1.67$ for our ΛCDM concordance cosmology, while SCDM with $\Omega_m = 1$, $\Omega_\Lambda = 0$ has $f = 1.55$, which is only 2.9σ from this result. Our constraint from the distance ratio only separates the concordance model from $\Omega_m = 1$ at 1.8σ , i.e. it is not a strong cosmological constraint, compared with the constraint on $d_{0.275}$.

7 COSMOLOGICAL INTERPRETATION

We now consider how our constraints can be mapped into the standard basis of cosmological parameters. From equation (6) of Eisenstein & Hu (1998), the sound horizon can be approximated around the *WMAP5* best-fitting location (Komatsu et al. 2009) as

$$r_s(z_d) = 153.5 \left(\frac{\Omega_b h^2}{0.02273} \right)^{-0.134} \left(\frac{\Omega_m h^2}{0.1326} \right)^{-0.255} \text{Mpc}. \quad (12)$$

Setting $r_{s,\text{fid}} = 153.5 \text{Mpc}$ and using equation (10), we have

$$\begin{aligned} D_V(0.275) &= (1104 \pm 30) [r_s(z_d)/r_{s,\text{fid}}(z_d)] \text{Mpc} \\ &= (1104 \pm 30) \left(\frac{\Omega_b h^2}{0.02273} \right)^{-0.134} \left(\frac{\Omega_m h^2}{0.1326} \right)^{-0.255} \text{Mpc}, \end{aligned} \quad (13)$$

and $f = 1.736 \pm 0.065$ as our two statistically independent constraints.

The constraint on $D_V(0.275)$, combined with a measurement of $\Omega_m h^2$ from *WMAP5* (Dunkley et al. 2009; Hinshaw et al. 2009; Komatsu et al. 2009), is enough to measure Ω_m and H_0 given information about the distance scale from $z = 0$ to $z = 0.275$. If the distance measure were at $z = 0$, then we would have a standard ruler defined by the CMB with which we could measure H_0 , and combining this with $\Omega_m h^2$ would yield Ω_m . In practice, one has to include

the small corrections to $D_V(0.275)$ that arise from the low-redshift cosmology. Noting that $D_V(0.275) = 757.4 h^{-1} \text{ Mpc}$ for a flat $\Omega_m = 0.282$ Λ CDM cosmology, we can write $h = \sqrt{\Omega_m h^2} / \sqrt{\Omega_m}$ and solve

$$\Omega_m = (0.282 \pm 0.015) \left(\frac{\Omega_m h^2}{0.1326} \right)^{0.49} \times \left(\frac{D_V(z=0.275, \Omega_m=0.282)}{D_V(z=0.275)} \right)^2, \quad (14)$$

where we have dropped the dependence of the sound horizon on $\Omega_b h^2$, which the *WMAP5* data already constrains to 0.5 per cent, five times below our statistical error.

We can perturb the ratio of distances around the best-fitting $\Omega_m = 0.282$ to give

$$\frac{D_V(z=0.275)}{D_V(z=0.275, \Omega_m=0.282)} = \left(\frac{\Omega_m}{0.282} \right)^{-0.077} [1 - 0.108\Omega_k - 0.099(1+w)]. \quad (15)$$

Using this approximation, we can manipulate equation (14) to give constraints on either Ω_m or h :

$$\Omega_m = (0.282 \pm 0.018) \left(\frac{\Omega_m h^2}{0.1326} \right)^{0.58} \times [1 + 0.25\Omega_k + 0.23(1+w)], \quad (16)$$

$$h = (0.686 \pm 0.022) \left(\frac{\Omega_m h^2}{0.1326} \right)^{0.21} \times [1 - 0.13\Omega_k - 0.12(1+w)]. \quad (17)$$

The additional uncertainty in Ω_m , ± 0.018 in equation (16) compared with ± 0.15 in equation (14), is produced by the dependence of the distance ratio on Ω_m . In equations (16) and (17), the uncertainty in the first terms is correlated so as to leave $\Omega_m h^2$ constant. One should additionally include the errors from $\Omega_m h^2$, Ω_k and w , although these are consistent between the two results.

Looking at the fractional error in Ω_m , the contribution from the uncertainty in the SDSS acoustic scale is about 6 per cent, that from the uncertainty in $\Omega_m h^2$ is about 2 per cent, that from w is about 3 per cent if the error on w is 10 per cent and that from curvature is below 1 per cent unless the cosmology is rather non-standard. Hence, our result is still limited by the SDSS-II BAO data volume and not by our knowledge of the other cosmological parameters in equation (16). Of course, these expressions only hold for mild perturbations from the concordance cosmology; for other cases, one should return to the raw distance constraints. We note that these expressions have not used the angular acoustic scale in the CMB, so they are independent of what is happening with dark energy at $z > 0.35$.

Fig. 5 shows the BAO constraints from equation (13) on Ω_m and Ω_Λ for Λ CDM cosmologies (upper panel) and on Ω_m and w for flat models where constant $w \neq -1$ is allowed (lower panel). We take a Gaussian prior of $\Omega_m h^2 = 0.1326 \pm 0.0063$ and assume that the error on $\Omega_b h^2$ is negligible as the *WMAP5* data already constrain it to 0.5 per cent (Komatsu et al. 2009). These constraints exclude the angular acoustic scale in the CMB, so they are independent of the dark energy behaviour at the redshifts beyond our sample. For comparison we plot the full *WMAP5* constraints (Komatsu et al. 2009), which include the constraints on the distance to last scattering, and constraints from the Union supernova (SN) sample (Kowalski et al. 2008), which constrain angular diameter distance ratios up to $z \sim 1$.




Figure 5. Cosmological constraints on Λ CDM cosmologies (upper panel) and flat CDM models where we allow w to vary (lower panel), from *WMAP5* (blue), Union SN (green) and our constraint on $r_s/D_V(0.275)$ (solid contours). Contours are plotted for $-2 \ln \mathcal{L}/\mathcal{L}_{\text{true}} < 2.3, 6.0$, corresponding to 68 and 95 per cent confidence intervals. The dashed lines show flat models (upper panel) and Λ models (lower panel).

Results from full likelihood fits combining these data are presented in Section 9.

8 TESTING THE ROBUSTNESS OF THE RESULTS

8.1 The effect of redshift-space distortions

We have fitted our spline \times BAO model to the observed SDSS LRG power spectrum, as calculated by Reid et al. (2009b), where the galaxy power spectrum and derived cosmological constraints are presented. Using numerical simulations, a scheme is presented in Reid et al. (2009b) to recover the halo power spectrum from the LRG distribution by only keeping a single LRG within each halo. We have fitted both the galaxy and the halo power spectra with our spline \times BAO model. The log ratio between the BAO recovered in the resulting fits is shown in Fig. 6. This shows that the cluster-collapse correction for these galaxies results in a smooth change




Figure 6. The log ratio of the BAO recovered from the SDSS LRG power spectrum to the power spectrum of the halo catalogue derived from the LRG sample as described by Reid et al. (2009b) (solid circles). For comparison we plot the BAO expected for a flat Λ CDM model with $\Omega_m = 0.25$, $h = 0.72$ and $\Omega_b h^2 = 0.0223$ (solid line), and the errors on each measurement (grey shaded region). There are no oscillatory features induced by the cluster-collapse procedure, and the scatter is well within the errors.

in the power spectrum on the scales fitted, and does not alter the position or amplitude of the BAO in a significant way.

Because of the different galaxy properties within the SDSS main galaxy sample, and the 2dFGRS, we do not attempt to correct for the more complicated distribution of galaxies within the haloes of that sample, and recover the halo power spectrum. In contrast, the halo occupation distribution of the SDSS LRGs is simple, in that there is only a single population of galaxies that are predominantly central rather than satellite galaxies in their hosting haloes (Reid et al. 2009b). But we have seen that for LRGs, the correction is smooth, and we expect this to be true for the galaxies at $z < 0.2$ as well.

8.2 Sample selection

We have run our full analysis pipeline using three sub-samples of galaxies. Results from fits to $D_V(z)$ with two nodes are shown in Fig. 7, for different catalogues, given $r_s(z_d) = 154.7$ Mpc. The best-fitting constraints for these models on d_z are given in Table 3. Our default analysis is included in panel (a) for comparison. Here, we analyse data from the SDSS and the 2dFGRS, including the early SDSS data, where we cut the sample at the extinction-corrected magnitude limit $r < 17.5$. We compare with results obtained by




Figure 7. As Fig. 4, but now considering results from four choices of catalogue: (a) all SDSS and non-overlapping 2dFGRS data, (b) excluding both the early SDSS data and 2dFGRS, (c) excluding the 2dFGRS and (d) excluding the early SDSS data.

Table 3. Measurements of $d_z \equiv r_s(z_d)/D_V(z)$ at $z = 0.2$ and $z = 0.35$ from the different analysis runs described in the captions to Figs 7 and 8.

	$d_{0.2}$	$d_{0.35}$
(a) Default	0.1905 ± 0.0061	0.1097 ± 0.0036
(b) No early SDSS, 2dFGRS	0.1923 ± 0.0072	0.1102 ± 0.0041
(c) No 2dFGRS	0.1907 ± 0.0062	0.1090 ± 0.0036
(d) No early SDSS	0.1917 ± 0.0069	0.1109 ± 0.0044
(e) Fit to three $P(k)$	0.1901 ± 0.0066	0.1080 ± 0.0043
(f) Original calibration	0.1919 ± 0.0071	0.1094 ± 0.0046
(g) Varying D_{damp}	0.1918 ± 0.0080	0.1100 ± 0.0048
(h) $\langle n(z) \rangle$ sampling galaxies	0.1890 ± 0.0068	0.1102 ± 0.0045

(b) excluding the early SDSS data and the 2dFGRS, (c) using just the SDSS data and (d) excluding the early SDSS data but including the 2dFGRS. Including the early SDSS galaxies decreases the errors at redshifts $z = 0.2$ and $z = 0.35$ by approximately 14 per cent. Including the 2dFGRS galaxies has a smaller effect, decreasing the error at $z = 0.2$ by approximately 4 per cent. The parameters of the best-fitting solutions do not move significantly with any of the sample changes: $d_{0.2}$ moves by a maximum of 0.3σ , while $d_{0.35}$ moves by a maximum of 0.2σ . The inclusion of the 2dFGRS actu-

ally moves the best-fitting solution for $D_V(0.35)/D_V(0.2)$ slightly towards that of a concordance Λ CDM model.

8.3 The number of redshift slices included

We now consider the robustness of our fit to the number of redshift slices analysed. This test was performed on the conservative data sample, excluding the early SDSS data and the 2dFGRS. In our default analysis we fit power spectra calculated for six redshift slices, and the resulting likelihood surface for the late SDSS sample is shown in panel (b) of Fig. 7. For comparison, panel (e) of Fig. 8 shows the likelihood surface calculated using power spectra from only three redshift slices (details of the slices chosen are presented in Section 3). Because we are only fitting two $D_V(z)$ nodes, these should be constrained by our reduced fit using three redshift slices. Panel (e) of Fig. 8 shows that this is true, but comparison with panel (b) of Fig. 7 shows that the constraints are tighter if we model power spectra from six redshift slices. Clearly, extra information is available from the extra redshift slices, and we therefore fit to six redshift slices for our default analysis.

It is interesting to test if there is sufficient information to constrain the shape of $D_V(z)$ beyond our simple spline model with two nodes. Results from fits allowing four $D_V(z)$ nodes are shown in Fig. 9.




Figure 8. Likelihood contour plots as in Fig. 4, for the SDSS data, excluding the early data and the 2dFGRS, but now calculated for (e) fit to three power spectra, (f) old rather than uber-calibration, (g) allowing the BAO damping scale to vary with a simple Gaussian prior $D_{\text{damp}} = 10 \pm 5 h^{-1} \text{ Mpc}$ and (h) randomized galaxy redshifts used to give the expected radial galaxy distribution.




Figure 9. Contour plots showing slices through the likelihood for four $D_V(z)$ cubic spline nodes at $z = 0.1$, $z = 0.2$, $z = 0.3$ and $z = 0.45$, calculated for our default analysis using six power spectra, uber-calibration and a fixed BAO damping scale of $D_{\text{damp}} = 10 h^{-1} \text{ Mpc}$. Shaded regions are plotted for $-2 \ln \mathcal{L}/\mathcal{L}_{\text{true}} < 2.3$, 6.0 , 9.3 , which for a multivariate Gaussian distribution with two degrees of freedom correspond to 68, 95 and 99 per cent confidence intervals. Likelihoods were adjusted to match these Gaussian confidence intervals as described in Section 5. In each panel, the nodes that are not shown were fixed at the default ΛCDM ($\Omega_m = 0.25$, $\Omega_\Lambda = 0.75$) values. We use shaded regions in this plot to show the likelihood surface, compared with the contours in Figs 7 and 8 because the likelihood surface is more complicated with four nodes, and the shading helps to distinguish peaks from troughs.

There is a clear maximum in the slices through the likelihood surface close to the ΛCDM model, but the surface is noisy, and there are secondary maxima present. There is a strong degeneracy between $D_V(0.3)$ and $D_V(0.45)$ and between $D_V(0.1)$ and $D_V(0.2)$: the data contain limited information to distinguish the shape of the distance-redshift relation between these redshifts. Consequently, we do not try to extract this information, instead concentrating on fits where there are only two nodes in $D_V(z)$.

8.4 The covariance matrix

Because we are analysing overlapping shells in redshift, the power spectra will be strongly correlated and the estimation of the covariance matrix will be in error if we do not have sufficient mock catalogues. In order to test this, we have recalculated our covariance matrix using 1/3 as many LN catalogues, and have used this matrix to recalculate the required corrections to the confidence intervals using independent sets of LN catalogues. We find consistent results in the factors required to match the confidence intervals to

those expected for a multivariate Gaussian distribution. We have also performed a full analysis using this reduced covariance matrix, and find results consistent with using our default covariance matrix.

8.5 Calibration

The likelihood surface shown in panel (f) of Fig. 8 was calculated using an SDSS galaxy sample with luminosities calibrated using the photometric calibration (Tucker et al. 2006), prior to the uber-calibration analysis (Padmanabhan et al. 2008a). This affects the calculation of the redshift completeness for any region observed and also the luminosity-dependent weights applied to the SDSS galaxies. The effect of this calibration change on our results is small, and there is no significant change between the likelihood surface in panel (f) of Fig. 8 and that in panel (b) of Fig. 7, where the uber-calibration data set was used.

8.6 BAO damping scale

Panel (g) of Fig. 8 shows the likelihood surface if we allow the BAO damping scale to be a free parameter in the fit, placing a simple Gaussian prior on its value $D_{\text{damp}} = 10 \pm 5 h^{-1} \text{ Mpc}$. This prior on the BAO damping scale is conservative. From simulations, Reid et al. (2009a) found $D_{\text{damp}} = 9.2 \pm 1 h^{-1} \text{ Mpc}$, with no variation with redshift for $0 < z < 0.5$ for halo density fields, and $D_{\text{damp}} = 9.7 \pm 1 h^{-1} \text{ Mpc}$ for density fields matched to the LRGs. The mild cosmological dependence suggested by Eisenstein et al. (2007) shows that the main cosmological dependence is through the linear growth rate; current constraints on σ_8 are much better than that required to significantly change D_{damp} , and we consider $\pm 5 h^{-1} \text{ Mpc}$ to be a conservative prior. Allowing the damping scale to vary degrades the constraint, increasing the size of the parameter confidence regions. The best-fitting solution does not move significantly, suggesting that our default assumption of a fixed damping scale is sufficiently accurate to current data precision.

8.7 Radial galaxy distribution model

Finally, analysis run (h) shows the constraints if we use a random catalogue where we randomly choose a galaxy redshift for each angular position chosen, i.e. to model the expected redshift distribution $\langle n(z) \rangle$, we sample from the galaxy redshift distribution. This test was designed to investigate the dependence of the analysis on how well we model the radial galaxy distribution. Randomly sampling galaxies to obtain this distribution perfectly matches the redshift distribution of the galaxies and that of the random catalogue used to define the survey region. In fact, we see no change in our results if we do this rather than using a smooth fit to the redshift distribution. This gives us confidence that our results are not sensitive to this modelling.

8.8 Secondary likelihood maxima

In the likelihood surfaces in Figs 7 and 8, we see secondary likelihood maxima, which appear to lie on a degeneracy stretching from $D_V(0.2) = 700 \text{ Mpc}$, $D_V(0.35) = 1500 \text{ Mpc}$ to $D_V(0.2) = 600 \text{ Mpc}$, $D_V(0.35) = 1000 \text{ Mpc}$. These minor peaks in the likelihood, which appear as isolated islands in the likelihood surface, are of lower significance than the strong peak close to the parameters of a concordance ΛCDM model. Tests have shown that the secondary peaks result from the interplay of two competing effects, which are themselves a result of using the wrong cosmology to analyse the BAO. These are as follows.

- (i) A shift in the BAO position.
- (ii) An increase in the width of the window associated with each band power, caused by BAO in different redshift shells being out of phase. This can smooth out the BAO signal.

Secondary maxima are produced where the BAO shift and the smoothing effects ‘balance’. If we redo the analysis ignoring the second effect by assuming that the window function is a δ -function centred on the peak, then these secondary maxima are removed.

8.9 Dependency on D_V

A possible concern about our method of analysis is that we assume a fiducial ΛCDM model to convert redshifts to comoving coordinate distances and measure the position of the BAO in the spherically

averaged power spectrum. If the true cosmological model has a different angular diameter distance–redshift relation $D_A(z)$ and Hubble parameter $H(z)$ than this fiducial model, then this would cause angular and radial distortions in the density field from which we estimate the power spectrum. By presenting results in terms of D_V , we remove the anisotropic information and assume that the expected BAO position for all cosmological models is solely dependent on their predicted value of D_V . This must break down for models that behave very differently from our fiducial ΛCDM model.

We now test the sensitivity of the assumption that the BAO position in the spherically averaged power spectrum only depends on D_V for cosmological models that predict significant anisotropic distortions in the density field away from our fiducial model. To do this, we compute the shifts of the BAO position expected when one measures the spherically averaged power in either real or redshift space for such models. To simplify the analysis, we assume that the BAO in the spherically averaged $P(k)$ will be shifted by the average of the shifts in k predicted over all angles, i.e. our BAO fit recovers the weighted mean shift in the 3D power. In redshift space, we also follow the distant observer approximation and assume that the angular dependence of the true 3D power spectrum is given by $(1 + \beta\mu^2)^2$, where μ is the cosine of the angle to the line of sight and $\beta = \Omega_m^{0.55}/b$. The anisotropy in the observed power spectrum caused by redshift-space distortions will act as a weight when we spherically average.

For the SDSS LRGs, which provide most of our cosmological signal, we take an effective redshift of $z = 0.35$ and assume a ΛCDM model with $\Omega_m(z=0) = 0.25$, giving $\Omega_m(z=0.35) = 0.45$. The LRGs are strongly biased and the model of Tegmark et al. (2004) gives an effective relative bias for our sample, which we correct for in the power spectrum calculation, of $\langle b/b_* \rangle = 1.9$. Matching the normalization of the measured LRG power spectrum (Reid et al. 2009b) gives that $b_* = 1.34$, assuming that the LRG clustering is constant in comoving coordinates (e.g. Percival et al. 2007b), and that $\sigma_8(\text{matter}, z=0) = 0.8$, so $\sigma_8(\text{matter}, z=0.35) = 0.68$ (Komatsu et al. 2009). This suggests that we should expect $\beta \sim 0.25$ for the LRG power spectrum, and we show contours calculated by assuming $\beta = 0.25$ in Fig. 10, which we compare with the prediction for $\beta = 1$. Note that our luminosity-dependent weighting means that we are upweighting highly biased galaxies and that our analysis will therefore have a smaller effective β than analyses without such weighting, such as the measurements presented by Cabre & Gaztanaga (2009).

Fig. 10 shows the relation between radial and angular distortions, H/H_{fid} and $D_A/D_{A,\text{fid}}$, which give rise to zero and ± 1 per cent shift in the spherically averaged power spectrum. Here, H_{fid} is the fiducial value of H and similarly for D_A . For general cosmological models, H/H_{fid} and $D_A/D_{A,\text{fid}}$ will depend on redshift, so that the final effective shift will be an average over a trajectory in this diagram which is determined by the model to be tested. Fig. 10 also shows the expected line of zero average shift we would expect if the BAO position only depends on $D_V(z)$, which would lead to behaviour such that $H(z) \propto D_A^2$. For comparison, we show the prediction for a model with increased importance of the radial distortions, with $H(z) \propto D_A$. This is included because we would expect that the redshift-space distortions will increase the importance of the radial information. However, the $H(z) \propto D_A^2$ line is a significantly better fit, even in redshift space. The $H(z) \propto D_A^2$ line does not cross the contours marking a 1 per cent average shift for our redshift-space power spectrum, showing that the assumption that the recovered BAO position only depends on D_V at most produces a 1 per cent systematic shift in the best fit for models with an anisotropy




Figure 10. The expected shift recovered from an analysis of the BAO position in a spherically averaged galaxy power spectrum, if there are radial and angular distortions induced by assuming an incorrect cosmology when analysing the data. The thick solid contour shows no residual shift, while the dotted contours show a 1 per cent shift. For comparison we plot the expected behaviour for an isotropic power spectrum $H(z) \propto D_A^2$, and for an increased importance of the radial distortion $H(z) \propto D_A$ (dashed lines). The top panel approximates redshift space, by weighting the power in the spherical average by $(1 + \beta\mu^2)^2$, with $\beta = 0.25$, matching that expected for the SDSS LRGs, while the bottom panel does not include this weighting. For comparison, the thin solid contour in the top panel marks no residual shift for data with $\beta = 1$, showing that we should expect the radial signal to increase in importance for such a sample.

distortion away from our fiducial model of up to 20 per cent in the radial direction. Such a 1 per cent systematic shift, which requires a model that is extremely discrepant from Λ CDM, is significantly below the statistical precision of our 2.7 per cent accuracy distance measurement. It is therefore a reasonable approximation to use our measurements of D_V to constrain a wide variety of cosmological models.

9 COSMOLOGICAL PARAMETER CONSTRAINTS

We now apply our full constraints to a cosmological parameter analysis. We assume that the likelihood of a model is given by a multivariate Gaussian distribution around the $D_V(z)$ measurements given

by equation (3), with the inverse covariance matrix of equation (5). Throughout this section we consider four models: a flat universe with a cosmological constant (Λ CDM), a Λ CDM universe with curvature (o Λ CDM), a flat universe with a dark energy component with a constant equation of state w (wCDM) and a wCDM universe with curvature (owCDM). This is the same model set considered by Reid et al. (2009b). We use a modified version of COSMOMC (Lewis & Bridle 2002) to perform the likelihood calculations.

9.1 SN + BAO + CMB prior likelihood fits

We first consider the constraints excluding the angular acoustic scale in the CMB, in order to consider data that are independent of the dark energy behaviour at the redshifts beyond our sample. This is important because it ensures that our results only depend on the acceleration of the Universe at late times and so do not depend on the so-called early dark energy models (Ratra & Peebles 1988; Wetterich 1988; Steinhardt, Wang & Zlatev 1999; Zlatev, Wang & Steinhardt 1999), which have non-negligible dark energy at early times. We take Gaussian priors $\Omega_c h^2 = 0.1099 \pm 0.0063$ and $\Omega_b h^2 = 0.02273 \pm 0.00061$ from the CMB; these constraints from the ratio of peak heights in the *WMAP5* data alone do not relax when Ω_k and w are allowed to vary. We also impose weak priors on $-0.3 < \Omega_k < 0.3$ and $-3 < w < 0$. The parameter constraints from the combination of Union SN (Kowalski et al. 2008) and BAO likelihoods with these priors are presented in Table 4. The best-fitting value of Ω_m ranges from 0.286 to 0.290, with the 68 per cent confidence interval, ± 0.018 , while the mean value of H_0 varies between 67.8 and 68.6 $\text{km s}^{-1} \text{Mpc}^{-1}$, and the 68 per cent confidence interval remains $\pm 2.2 \text{ km s}^{-1} \text{Mpc}^{-1}$ throughout the four models. In Section 7 we derived BAO-only constraints of ± 0.018 on Ω_m and $\pm 2.2 \text{ km s}^{-1} \text{Mpc}^{-1}$ on H_0 , for fixed $\Omega_m h^2$. If we include the 4.8 per cent error on $\Omega_m h^2$ from the *WMAP5* measurement, then we should expect these errors to increase to ± 0.019 on Ω_m and $\pm 2.3 \text{ km s}^{-1} \text{Mpc}^{-1}$ on H_0 . These agree perfectly with the COSMOMC results if we exclude the SN data, so the small difference between the errors in Table 4 and those expected is caused by the SN data helping to constrain Ω_m and H_0 slightly. Similarly, the best-fitting values of these parameters agree for COSMOMC results excluding the SN data. Comparison between Table 4 and Section 7 shows that the inclusion of the SN data is moving the best fit slightly: $+0.004$ in Ω_m and -0.5 in H_0 for the Λ CDM model. The COSMOMC analysis therefore validates the simple derivation presented in Section 7. In the space of models considered here, the BAO constraint on $D_V(0.275)$ already restricts $D_V(0.35)/D_V(0.2)$ to a much smaller region than our constraint in equation (11) allows. While the combination of these data and our priors are unable to constrain Ω_k , w is constrained at the ± 0.11 level. For the owCDM model, the weak prior on Ω_k leads to an apparent constraint on w , but these errors depend strongly on the prior.

The data are compared with the best-fitting Λ CDM model in Fig. 11. Three ways of considering the data constraints are shown in different panels. In the bottom panel we plot $D_V(z)/D_V(0.2)$, which corresponds to matching the geometry at $z = 0.2$ and $z = 0.35$ so the BAO match at these redshifts, without including information about the comoving position of the BAO. In the middle panel we plot $r_s(z_d)/D_V(z)$, where we now have to model the comoving sound horizon at the drag epoch. In the top panel, we include a constraint on the sound horizon projected at the last-scattering surface as observed in the CMB. Marginalizing over the set of flat Λ CDM models constrained only by the *WMAP5* data gives $r_s(z_d)/S_k(z_d) = 0.010824 \pm 0.000023$, where $S_k(z_d)$ is the proper distance to the

Table 4. Marginalized one-dimensional constraints (68 per cent) for BAO+SN for flat Λ CDM, Λ CDM with curvature ($\text{o}\Lambda$ CDM), flat w CDM (w CDM) and w CDM with curvature (ow CDM). The non-standard cosmological parameters are $d_{0.275} \equiv r_s(z_d)/D_V(0.275)$ and $f \equiv D_V(0.35)/D_V(0.2)$. We have assumed priors of $\Omega_c h^2 = 0.1099 \pm 0.0063$ and $\Omega_b h^2 = 0.02273 \pm 0.00061$, consistent with *WMAP5*-only fits to all of the models considered here. We also impose weak flat priors of $-0.3 < \Omega_k < 0.3$ and $-3 < w < 0$.

Parameter	Λ CDM	$\text{o}\Lambda$ CDM	w CDM	ow CDM
Ω_m	0.288 ± 0.018	0.286 ± 0.018	$0.290^{+0.018}_{-0.019}$	0.286 ± 0.018
H_0	$68.1^{+2.2}_{-2.1}$	68.6 ± 2.2	67.8 ± 2.2	68.2 ± 2.2
Ω_k	–	-0.097 ± 0.081	–	$-0.199^{+0.080}_{-0.089}$
w	–	–	-0.97 ± 0.11	$-0.838^{+0.083}_{-0.084}$
Ω_Λ	0.712 ± 0.018	$0.811^{+0.084}_{-0.085}$	$0.710^{+0.019}_{-0.018}$	$0.913^{+0.092}_{-0.082}$
$d_{0.275}$	0.1381 ± 0.0034	0.1367 ± 0.0036	0.1384 ± 0.0037	0.1386 ± 0.0037
$D_V(0.275)$	1111 ± 31	1120 ± 33	1109 ± 32	1108^{+32}_{-33}
f	1.662 ± 0.004	1.675 ± 0.011	1.659 ± 0.011	1.665 ± 0.011
Age (Gyr)	$14.02^{+0.32}_{-0.31}$	14.43 ± 0.48	13.95 ± 0.36	14.38 ± 0.44




Figure 11. The BAO constraints (solid circles with 1σ errors), compared with the best-fitting Λ CDM model. The three panels show different methods of using the data to constrain models.

baryon-drag redshift $z_d = 1020.5$, as measured by the *WMAP5* team (Komatsu et al. 2009). Ignoring the negligible error on this quantity, we combine with the BAO results to measure $S_k(z_d)/D_V(z)$. This effectively removes the dependence on the comoving sound horizon at the drag epoch, anchoring the BAO measurements at high redshift: here we have done this at the baryon-drag epoch so the CMB constraint has matched the sound horizon and projection distance.

9.2 CMB + BAO likelihood fits

We now turn to the constraints from our BAO measurement combined with the full *WMAP5* likelihood, including the constraint on $r_s(z_d)/D_A$ at the time of decoupling. While this extra constraint can break degeneracies between Ω_m , Ω_k and w inherent in our BAO constraints, the results are now sensitive to our assumption of a constant dark energy equation of state w at $z > 0.35$. Results for the four models are presented in Table 5.

For the Λ CDM model, we find $\Omega_m = 0.278 \pm 0.018$ and $H_0 = 70.1 \pm 1.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$, with errors significantly reduced compared to the *WMAP5*-alone analysis ($\Omega_m = 0.258 \pm 0.03$ and $H_0 = 70.5^{+2.6}_{-2.7} \text{ km s}^{-1} \text{ Mpc}^{-1}$). Similar limits on Ω_m were obtained by Rozo et al. (2009) who used the maxBCG cluster abundance and weak-lensing mass measurements to similarly break the tight *WMAP5* constraint on $\Omega_m h^2$.

Fig. 12 shows the impact of relaxing the flat, Λ CDM assumption. The *WMAP5* results alone tightly constrain $\Omega_m h^2$ in all of these models (dashed lines), but low-redshift information is necessary to constrain Ω_m and H_0 separately. Allowing $w \neq -1$ relaxes the constraint on Ω_m from the BAO measurement, and in addition allowing $\Omega_k \neq 0$ relaxes the constraint even further. The impact on the constraints on Ω_m and H_0 is shown in the lower right-hand panel. All of the contours lie along the banana with $\Omega_m h^2$ fixed from the CMB.

In the $\text{o}\Lambda$ CDM model, the combination of scales measured by the CMB and the BAO tightly constrains the curvature of the universe: $\Omega_k = -0.007^{+0.006}_{-0.007}$. The constraints on Ω_m and H_0 in this model are well described by equations (16) and (17), while in the w CDM cosmology they degrade because w is not well constrained by the low-redshift BAO information alone.

When the parameter space is opened to both curvature and w , the *WMAP5* data are not able to eliminate the degeneracy between Ω_m and w in the BAO constraint. The constraints relax to $\Omega_m = 0.240^{+0.044}_{-0.043}$ and $H_0 = 75.3 \pm 7.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$; $\Omega_k = -0.013 \pm 0.007$ is still well constrained but w is not (see Fig. 12). Including the constraints from the Union SN sample breaks the remaining degeneracy, and we recover the tight constraints on Ω_m and H_0 given in equations (16) and (17). These constraints, and the relative degeneracies induced and broken by different data sets, are shown in Fig. 13. For each of the four models considered, the central values for Ω_m and H_0 change only slightly when the full *WMAP5*

Table 5. Marginalized one-dimensional constraints (68 per cent) for *WMAP5*+BAO for flat Λ CDM, Λ CDM with curvature (o Λ CDM), flat wCDM (wCDM), wCDM with curvature (owCDM) and owCDM including constraints from SNe. The non-standard cosmological parameters constrained by the BAO measurements are $d_{0.275} \equiv r_s(z_d)/D_V(0.275)$ and $f \equiv D_V(0.35)/D_V(0.2)$.

Parameter	Λ CDM	o Λ CDM	wCDM	owCDM	owCDM+SN	owCDM+ H_0	owCDM+SN+ H_0
Ω_m	0.278 ± 0.018	0.283 ± 0.019	0.283 ± 0.026	$0.240^{+0.044}_{-0.043}$	0.290 ± 0.019	$0.240^{+0.025}_{-0.024}$	0.279 ± 0.016
H_0	70.1 ± 1.5	$68.3^{+2.2}_{-2.1}$	69.3 ± 3.9	75.3 ± 7.1	67.6 ± 2.2	74.8 ± 3.6	69.5 ± 2.0
Ω_k	–	$-0.007^{+0.006}_{-0.007}$	–	-0.013 ± 0.007	-0.006 ± 0.008	-0.014 ± 0.007	-0.003 ± 0.007
w	–	–	-0.97 ± 0.17	$-1.53^{+0.51}_{-0.50}$	-0.97 ± 0.10	$-1.49^{+0.32}_{-0.31}$	-1.00 ± 0.10
Ω_Λ	0.722 ± 0.018	0.724 ± 0.019	0.717 ± 0.026	0.772 ± 0.048	0.716 ± 0.019	0.773 ± 0.029	0.724 ± 0.018
$100 \Omega_b h^2$	2.267 ± 0.058	2.269 ± 0.060	2.275 ± 0.061	$2.254^{+0.062}_{-0.061}$	2.271 ± 0.061	$2.254^{+0.061}_{-0.062}$	2.284 ± 0.061
τ	0.086 ± 0.016	0.089 ± 0.017	0.087 ± 0.017	0.088 ± 0.017	0.089 ± 0.017	0.088 ± 0.017	$0.089^{+0.017}_{-0.018}$
n_s	0.961 ± 0.013	0.963 ± 0.014	0.963 ± 0.015	0.958 ± 0.014	0.963 ± 0.014	0.957 ± 0.014	0.964 ± 0.014
$\ln(10^{10} A_{05})$	$3.074^{+0.040}_{-0.039}$	3.060 ± 0.042	3.070 ± 0.041	$3.062^{+0.042}_{-0.043}$	$3.062^{+0.041}_{-0.042}$	3.062 ± 0.042	3.072 ± 0.042
$d_{0.275}$	0.1411 ± 0.0030	0.1387 ± 0.0036	$0.1404^{+0.0036}_{-0.0035}$	0.1382 ± 0.0037	0.1379 ± 0.0036	$0.1387^{+0.0036}_{-0.0037}$	$0.1402^{+0.0033}_{-0.0034}$
$D_V(0.275)$	1080 ± 18	1110^{+32}_{-31}	1089 ± 31	1111 ± 33	1115 ± 32	1107 ± 31	1091^{+27}_{-28}
f	1.6645 ± 0.0043	1.6643 ± 0.0045	1.661 ± 0.019	1.72 ± 0.056	1.660 ± 0.011	$1.7187^{+0.0337}_{-0.0334}$	1.6645 ± 0.0107
Age(Gyr)	13.73 ± 0.12	14.08 ± 0.33	$13.76^{+0.15}_{-0.14}$	14.49 ± 0.52	14.04 ± 0.36	14.48 ± 0.48	$13.86^{+0.34}_{-0.33}$
$\Omega_c h^2$	0.1139 ± 0.0041	$0.1090^{+0.0060}_{-0.0061}$	$0.1122^{+0.0068}_{-0.0069}$	$0.1107^{+0.0063}_{-0.0062}$	$0.1096^{+0.0061}_{-0.0062}$	$0.1108^{+0.0060}_{-0.0061}$	0.1115 ± 0.0061
Ω_{tot}	–	$1.007^{+0.006}_{-0.007}$	–	1.013 ± 0.007	1.006 ± 0.008	1.014 ± 0.007	1.003 ± 0.007
σ_8	0.813 ± 0.028	0.787 ± 0.037	$0.792^{+0.081}_{-0.082}$	0.907 ± 0.117	$0.780^{+0.052}_{-0.053}$	0.904 ± 0.074	$0.801^{+0.053}_{-0.052}$




Figure 12. *WMAP5*+BAO constraints on $\Omega_m h^2$, Ω_m and H_0 for Λ CDM (solid black contours), o Λ CDM (shaded green contours), wCDM (shaded red contours) and owCDM (shaded blue contours) models. Throughout, the solid contours show *WMAP5*+LRG Λ CDM constraints. The first three panels show *WMAP5*-only constraints (dashed contours) and *WMAP5*+BAO constraints (coloured contours) in the $\Omega_m h^2$ – Ω_m plane as the model is varied. In the lower right, we show all constraints from *WMAP5*+BAO for all four models in the Ω_m – H_0 plane, which lie within the tight $\Omega_m h^2 \approx 0.133 \pm 0.006$ *WMAP5*-only constraints.




Figure 13. For the owCDM model, we compare the constraints from $\text{WMAP5}+\text{BAO}$ (blue contours), $\text{WMAP5}+\text{SN}$ (green contours) and $\text{WMAP5}+\text{BAO}+\text{SN}$ (red contours). Dashed and solid contours highlight the 68 per cent confidence intervals for the $\text{WMAP5}+\text{BAO}$ and $\text{WMAP5}+\text{SN}$ models, respectively.

likelihoods are used (Table 5) instead of priors on $\Omega_b h^2$ and $\Omega_c h^2$ in combination with the Union SN sample (Table 4).

Table 5 also lists the best-fitting cosmological age (i.e. time since the big bang) for different cosmologies and data sets. While the age is very well determined for ΛCDM and wCDM , there is a degeneracy between age and curvature that increases the uncertainties and allows for an older age in $\text{o}\Lambda\text{CDM}$ and owCDM information. Adding SN and H_0 measurements reduces these uncertainties and implies a best-fitting age of $13.86^{+0.34}_{-0.33}$ Gyr.

9.3 Comparison with the Riess et al. (2009) H_0

Riess et al. (2009) recently released a new determination of the Hubble constant using a differential distance ladder: $H_0 = 74.2 \pm 3.6 \text{ km s}^{-1} \text{ Mpc}^{-1}$. This value and the values $H_0 \approx 68 \text{ km s}^{-1} \text{ Mpc}^{-1}$ determined in Table 4 using BAO, SN and a WMAP5 prior on $\Omega_c h^2$ and $\Omega_b h^2$ are within $\sim 1\sigma$ of the mean value of 70.1 determined from $\text{WMAP5}+\text{BAO}$ in a ΛCDM model. In the wCDM model, combining this new H_0 with the WMAP5 likelihood constrains $w = -1.12 \pm 0.12$. In Table 5 we show Markov Chain Monte Carlo (MCMC) results for the owCDM model for $\text{WMAP5}+\text{BAO}+H_0$

and $\text{WMAP5}+\text{BAO}+H_0+\text{SN}$.¹ In this model, the SN data are more effective than H_0 at breaking the long degeneracy in the $\text{WMAP5}+\text{BAO}$ constraints. Combining $\text{WMAP5}+\text{BAO}+\text{SN}+H_0$, the mean parameters are quite close to ΛCDM : $\Omega_k = -0.003 \pm 0.007$ and $w = -1.00 \pm 0.10$, and $\Omega_m = 0.279 \pm 0.016$ and $H_0 = 69.5 \pm 2.0 \text{ km s}^{-1} \text{ Mpc}^{-1}$ are also well constrained.

10 COMPARISON WITH DR5 ANALYSES

In Percival et al. (2007c), we presented BAO measurements calculated from fitting power spectra calculated for three samples drawn from the combined SDSS+2dFGRS catalogue, using the SDSS DR5 data. The full catalogue was split into galaxy populations, rather than redshift slices, corresponding to the SDSS LRGs, the 2dFGRS+SDSS main galaxies and the combined sample. From this, we obtained the distance constraints $r_s(z_d)/D_V(0.2) = 0.1980 \pm 0.0058$ and $r_s(z_d)/D_V(0.35) = 0.1094 \pm 0.0033$ with a correlation coefficient of 0.39, which gives a distance ratio measurement of $D_V(0.35)/D_V(0.2) = 1.812 \pm 0.062$. The concordance ΛCDM value is $D_V(0.35)/D_V(0.2) = 1.67$, measured using the Supernovae Legacy Survey (SNLS) SN data, which is discrepant with the published DR5 BAO results at the 2.4σ level. The analysis of mock catalogues presented in Section 5 showed that the cubic spline \times BAO method underestimates the true distribution of recovered distances, given noisy data, which produce a non-Gaussian likelihood surface. We should therefore increase the errors on the DR5 measurements of Percival et al. (2007c) by at least a factor of 1.14, which is the correction derived from the fits to three DR7 power spectra. If we do this, the revised DR5 constraints are $r_s(z_d)/D_V(0.2) = 0.1981 \pm 0.0071$ and $r_s(z_d)/D_V(0.35) = 0.1094 \pm 0.0040$ with a correlation coefficient of 0.38, which gives a distance ratio measurement of $D_V(0.35)/D_V(0.2) = 1.813 \pm 0.073$. The discrepancy between the old DR5 constraints and the SNLS ΛCDM value is reduced to $\lesssim 2\sigma$. Because the DR5 data were noisier than the DR7 data, we should expect the likelihood surface to be less like a Gaussian prediction, and the correction actually should be slightly larger than that for the DR7 data.

Of all the changes implemented between this DR7 analysis and the analysis of the DR5 data, it was the increase in the number of random points used to quantify the survey geometry that had the most effect when comparing different catalogues. We now find consistent results, given in Table 3, for all catalogues and analysis variations presented in Section 8. When translated into constraints on the distance ratio, for the full catalogue we find $D_V(0.35)/D_V(0.2) = 1.736 \pm 0.065$. Using only three redshift slices, we find $D_V(0.35)/D_V(0.2) = 1.765 \pm 0.079$. If the 0.5σ difference is not due to chance, the difference between these measurements could be caused by residual non-Gaussian scatter in the band powers. A scenario in which this is reduced by including fits to more redshift bins would then explain the observed trend. Excluding the 2dFGRS and early SDSS data, the constraint is reduced to $D_V(0.35)/D_V(0.2) = 1.747 \pm 0.070$, which is consistent with the tighter constraint using all of the data.

Sanchez et al. (2009), who analysed the SDSS DR6 sample, speculated that the discrepancy could be caused by the Percival et al. (2007c) analysis fixing the BAO damping scale. However, in

¹ We account for the small cosmology dependence in the H_0 constraint (seen as a slight degeneracy between H_0 and w in fig. 14 of Riess et al. 2009) by considering it as a constraint on the inverse luminosity distance at the effective redshift $z = 0.04$ (Riess, private communication).

our current analysis, if we allow the BAO damping scale D_{damp} to vary, the derived constraints on $D_V(0.35)/D_V(0.2)$ do not change significantly from that recovered in our default analysis. The mild discrepancy with Λ CDM does not appear to be caused by fixing the damping scale. The change from photometric calibration to uber-calibration has a relatively minor effect on the distance ratio, which increases to $D_V(0.35)/D_V(0.2) = 1.748 \pm 0.074$. Fig. 6 shows that the effect on the BAO of redshift-space distortions caused by the thermal motion of galaxies in clusters is similarly small. Linear redshift-space distortions propagate the apparent position of galaxies along their velocity vector in a way that simply makes the field look more evolved than it is; they do not alter the positions of the BAO.

In conclusion, the significance of the discrepancy with flat Λ CDM models is reduced because of

- (i) analysis of the non-Gaussian nature of the likelihood surface;
- (ii) analysis of more redshift slices;
- (iii) more accurate determination of the galaxy redshift distribution.

11 DISCUSSION

In this paper we have measured and analysed BAO from the SDSS DR7 sample, which represents the final data set observed using the original SDSS spectroscopic target selection algorithm. We have further developed the analysis method used by Percival et al. (2007c) to analyse the DR5 sample, including a faster method for the calculation of the window function (see Appendix A), linking the cosmological model to be tested with the power spectrum band powers measured. This has enabled us to analyse power spectra calculated for six rather than three redshift slices, which would not have been possible using the old method.

In Section 6, we have shown how the distance–redshift constraints at $z = 0.2$ and $z = 0.35$ can be decomposed into a single distance constraint at $z = 0.275$ and a ‘gradient’ around this pivot given by $D_V(0.35)/D_V(0.2)$. This allows us to easily test the consistency of the Λ CDM model without having to compare with additional data. For the best-fitting flat Λ CDM model that matches our constraint $d_{0.275} = 0.1390 \pm 0.0037$, we find that our distance–ratio measurement of $D_V(0.35)/D_V(0.2) = 1.736 \pm 0.065$ is consistent at the 1.1σ level.

Now that the SDSS-II sample is complete, the importance of including the 2dFGRS data is reduced, and the inclusion only decreases the low redshift $z = 0.2$ distance error by 4 per cent. As we showed in Section 8.2, the inclusion of the 2dFGRS galaxies does not lead to the discrepancy with the Λ CDM model; including the 2dFGRS brings our constraint slightly more into line with the predictions of Λ CDM models.

Of the cosmological parameter constraints presented in Tables 4 and 5, perhaps the most impressive are the constraints on Ω_m and H_0 . For Λ CDM models, fitting to BAO and SNe with priors on $\Omega_m h^2$ and $\Omega_b h^2$ gives H_0 to 3.2 per cent and Ω_m to 6.4 per cent. These constraints are robust to the behaviour of the Universe at high redshift, as they are based only on the distance–redshift relation at redshift $z < 0.35$: we can allow $\Omega_k \neq 0$ and $w \neq -1$ with minimal effect. This weak dependence on w and Ω_k was shown in equations (16) and (17) for the BAO data.

If we allow for the flatness constraint to be relaxed, then we obtain $\Omega_k = -0.007 \pm 0.007$ from the combination of BAO+WMAP5 data. A tight constraint was similarly obtained on $w = -0.97 \pm 0.17$ if we relax the Λ constraint. If we allow both the curvature and the

dark energy equation of state to vary, we must include more data to continue to break the degeneracy between the two parameters. We do so by including results from the Union SN data set, giving us $\Omega_k = -0.006 \pm 0.008$ and $w = -0.97 \pm 0.10$, consistent with a flat Λ CDM model. If one allows only $w \neq -1$ or $\Omega_k \neq 0$, then the combination of CMB, SN and BAO data has an internal cross-check: opening two degrees of freedom from flat Λ CDM yields results that are consistent with flat Λ CDM. We have also shown that our constraints are consistent with the recent redetermination of H_0 by Riess et al. (2009), and that combining this constraint with WMAP5, BAO and SN in a model where both curvature and w vary yields mean parameter values very close to Λ CDM.

In a companion paper (Reid et al. 2009b), we consider the LRG sample in more detail. The LRGs are distributed in haloes in a simple way and we are able to extract the halo power spectrum from the data. In addition to fitting the BAO in this power spectrum, we are able to extract limited information about the shape of the power, which gives complementary constraints. A detailed comparison between the results from our fit to the BAO in redshift slices, performed in a cosmology model-independent way and including low-redshift galaxies, and the halo power spectrum of Reid et al. (2009b) is presented in that paper, where excellent agreement is demonstrated. The data sets are correlated so they should not be used together to constrain cosmological models.

Our analysis highlights the importance of BAO as a key method for investigating cosmic acceleration and shows that the method can already provide interesting cosmological constraints. Ongoing spectroscopic surveys aiming to use BAO to analyse dark energy include the BOSS (Schlegel et al. 2009a), the Hobby-Eberly Telescope Dark Energy Experiment (HETDEX; Hill et al. 2008) and the WiggleZ survey (Glazebrook et al. 2007). There are also plans for future surveys covering significantly larger volumes of the Universe and therefore observing the BAO signal with higher precision such as the Square Kilometer Array (SKA: www.skatelescope.org), and the Joint Dark Energy Mission (JDEM: jdem.gsfc.nasa.gov) and European Space Agency Euclid satellite mission concepts, or the Big Baryon Oscillation Spectroscopic Survey (BigBOSS; Schlegel et al. 2009b). Photometric surveys such as the Dark Energy Survey (DES: www.darkenergysurvey.org), the Panoramic Survey Telescope & Rapid Response System (Pan-STARRS: pan-starrs.ifa.hawaii.edu) and the Large Synoptic Survey Telescope (LSST: www.lsst.org) will find BAO using photometric redshifts. All of these surveys will measure BAO at higher redshifts than those analysed in our paper using SDSS-II data; if dark energy does not have a simple explanation, then comparison between future high-redshift results and our current understanding of the low-redshift Universe from SDSS-II will provide an interesting test of these models.

ACKNOWLEDGMENTS

WJP is grateful for support from the UK Science and Technology Facilities Council (STFC), the Leverhulme trust and the European Research Council. DJE was supported by National Science Foundation grant AST-0707225 and NASA grant NNX07AC51G. Simulated catalogues were calculated and analysed using the COSMOS Altix 3700 supercomputer, a UK-CCC facility supported by HEFCE and STFC in cooperation with CGI/Intel. WJP would like to thank Tamara M. Davis, Ravi K. Sheth, Roman Scoccimarro, Eyal Kazin, Taka Matsubara and the referee, Fergus Simpson, for useful interactions.

The 2dFGRS was undertaken using the Two-Degree Field facility on the 3.9 m Anglo-Australian Telescope. The success of the

survey was made possible by the dedicated efforts of the staff of the Anglo-Australian Observatory, both in creating the 2dF instrument and in supporting the survey observations.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society and the Higher Education Funding Council for England. The SDSS web site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, Cambridge University, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory and the University of Washington.

REFERENCES

- Abazajian K. et al., 2009, *ApJS*, 182, 543
 Adelman-McCarthy J. et al., 2006, *ApJS*, 162, 38
 Adelman-McCarthy J. et al., 2007, *ApJS*, 172, 634
 Adelman-McCarthy J. et al., 2008, *ApJS*, 175, 297
 Albrecht A. et al., 2006, Report of the Dark Energy Task Force. preprint (arXiv:astro-ph/0609591)
 Angulo R. E., Baugh C. M., Frenk C. S., Lacey C. G., 2008, *MNRAS*, 383, 755
 Bharadwaj S., 1996, *ApJ*, 472, 1
 Blake C., Glazebrook K., 2003, *ApJ*, 594, 665
 Blake C., Collister A., Bridle S., Lahav O., 2007, *MNRAS*, 374, 1527
 Blanton M. R., Lin H., Lupton R. H., Maley F. M., Young N., Zehavi I., Loveday J., 2003a, *AJ*, 125, 2276
 Blanton M. R. et al., 2003b, *ApJ*, 592, 819
 Bond J. R., Efstathiou G., 1984, *ApJ*, 285, L45
 Bond J. R., Efstathiou G., 1987, *MNRAS*, 226, 655
 Cabre A., Gaztanaga E., 2009, *MNRAS*, 396, 1119
 Cole S. et al., 2005, *MNRAS*, 362, 505
 Coles P., Jones B., 1991, *MNRAS*, 248, 1
 Colless M. et al., 2003, preprint (arXiv:astro-ph/0306581)
 Cresswell J. G., Percival W. J., 2009, *MNRAS*, 392, 682
 Crocce M., Scoccimarro R., 2006, *Phys. Rev. D*, 73, 063519
 Crocce M., Scoccimarro R., 2008, *Phys. Rev. D*, 77, 023533
 Dunkley J. et al., 2009, *ApJS*, 180, 306
 Eisenstein D. J., Hu W., 1998, *ApJ*, 496, 605
 Eisenstein D. J. et al., 2001, *AJ*, 122, 2267
 Eisenstein D. J. et al., 2005, *ApJ*, 633, 560
 Eisenstein D. J., Seo H.-J., White M., 2007, *ApJ*, 664, 660
 Feldman H. A., Kaiser N., Peacock J. A., 1994, *MNRAS*, 426, 23
 Fukugita M., Ichikawa T., Gunn J. E., Doi M., Shimasaku K., Schneider D. P., 1996, *AJ*, 111, 1748
 Gaztanaga E., Cabre A., Hui L., 2008, preprint (arXiv:0807.3551)
 Glazebrook K. et al., 2007, in Metcalf N., Shanks T., eds, *ASP Conf. Ser.* Vol. 379, *Cosmic Frontiers*. Astron. Soc. Pac., San Francisco, p. 72
 Goldberg D. M., Strauss M. A., 1998, *ApJ*, 495, 29
 Górski K. M., Hivon E., Banday A. J., Wandelt B. D., Hansen F. K., Reinecke M., Bartelmann M., 2005, *ApJ*, 622, 759
 Gunn J. E. et al., 1998, *AJ*, 116, 3040
 Gunn J. E. et al., 2006, *AJ*, 131, 2332
 Hill G. J. et al., 2008, in Tadayuki K., Yamada T., Aoki K., eds, *ASP Conf. Ser.* Vol. 399, *Panoramic Views of Galaxy Formation and Evolution*. Astron. Soc. Pac., San Francisco, p. 115
 Hinshaw G. et al., 2009, *ApJS*, 180, 225
 Hogg D. W., Finkbeiner D. P., Schlegel D. J., Gunn J. E., 2001, *AJ*, 122, 2129
 Holtzman J. A., 1989, *ApJS*, 71, 1
 Hu W., Haiman Z., 2003, *Phys. Rev. D*, 68, 3004
 Huetsi G., 2006, *A&A*, 449, 891
 Ivezić Z. et al., 2004, *Astron. Nachr.*, 325, 583
 Kazin E. et al., 2009, *ApJ*, submitted, preprint (arXiv:0908.2598)
 Kim J., Park C., Gott R., Dubinski J., 2009, *ApJ*, 701, 1547
 Komatsu E. et al., 2009, *ApJS*, 180, 330
 Kowalski M. et al., 2008, *ApJ*, 686, 749
 Lewis A., Bridle S., 2002, *Phys. Rev. D*, 66, 103511
 Lewis A., Challinor A., Lasenby A., 2000, *ApJ*, 538, 473
 Lupton R. H., Gunn J. E., Szalay A. S., 1999, *AJ*, 118, 1406
 Lupton R., Gunn J. E., Ivezić Z., Knapp G. R., Kent S., Yasuda N., 2001, in Harnden F. R., Jr, Primini F. A., Payne H. E., eds, *ASP Conf. Ser.* Vol. 238, *Astronomical Data Analysis Software and Systems X*. Astron. Soc. Pac., San Francisco, p. 269
 Matsubara T., 2004, *ApJ*, 615, 573
 Matsubara T., 2008a, *Phys. Rev. D*, 77, 063530
 Matsubara T., 2008b, *Phys. Rev. D*, 78, 083519
 Meiksin A., White M., Peacock J. A., 1999, *MNRAS*, 304, 851
 Padmanabhan N., White M., 2009, *Phys. Rev. D*, 80, 063508
 Padmanabhan N. et al., 2007, *MNRAS*, 378, 852
 Padmanabhan N. et al., 2008a, *ApJ*, 674, 1217
 Padmanabhan N., White M., Cohn J. D., 2008b, *Phys. Rev. D*, submitted, preprint (arXiv:0812.2905)
 Peebles P. J. E., Yu J. T., 1970, *ApJ*, 162, 815
 Percival W. J. et al., 2001, *MNRAS*, 327, 1297
 Percival W. J., Verde L., Peacock J. A., 2004, *MNRAS*, 347, 645
 Percival W. J. et al., 2007a, *ApJ*, 657, 51
 Percival W. J. et al., 2007b, *ApJ*, 657, 645
 Percival W. J., Cole S., Eisenstein D., Nichol R., Peacock J. A., Pope A., Szalay A., 2007c, *MNRAS*, 381, 1053
 Pier J. R., Munn J. A., Hindsley R. B., Hennessy G. S., Kent S. M., Lupton R. H., Ivezić Z., 2003, *AJ*, 125, 1559
 Press W. H., Teukolsky S. A., Vetterling W. T., Flannery B. P., 1992, *Numerical Recipes in C. The Art of Scientific Computing*, 2nd edn. Cambridge Univ. Press, Cambridge
 Ratra B., Peebles P. J. E., 1988, *Phys. Rev. D*, 37, 3406
 Reid B. A., Spergel D. N., Bode P., 2009a, *ApJ*, 702, 249
 Reid B. et al., 2009b, *MNRAS*, submitted, preprint (arXiv:0907.1659)
 Riess A. G. et al., 2009, *ApJ*, 699, 539
 Roze E. et al., 2009, preprint (arXiv:0902.3702)
 Sanchez A. G., Baugh C. M., Angulo R., 2008, *MNRAS*, 390, 1470
 Sanchez A. G., Crocce M., Cabre A., Baugh C. M., Gaztanaga E., 2009, *MNRAS*, in press (arXiv:0901.2570)
 Schlegel D., White M., Eisenstein D. J., 2009a, preprint (arXiv:0902.4680)
 Schlegel D. et al., 2009b, preprint (arXiv:0904.0468)
 Seo H.-J., Eisenstein D. J., 2003, *ApJ*, 598, 720
 Seo H.-J., Eisenstein D. J., 2005, *ApJ*, 633, 575
 Seo H.-J., Eisenstein D. J., 2007, *ApJ*, 665, 14
 Seo H.-J., Siegel E. R., Eisenstein D. J., White M., 2008, *ApJ*, 636, 16
 Silk J., 1968, *ApJ*, 151, 459
 Smith J. A. et al., 2002, *AJ*, 123, 2121
 Smith R. E., Scoccimarro R., Sheth R. K., 2007, *Phys. Rev. D*, 75, 063512
 Smith R. E., Scoccimarro R., Sheth R. K., 2008, *Phys. Rev. D*, 77, 043525
 Springel V. et al., 2005, *Nat*, 435, 629
 Steinhart P. J., Wang L.-M., Zlatev I., 1999, *Phys. Rev. D*, 59, 123504
 Stoughton C. et al., 2002, *AJ*, 123, 485
 Strauss M. A. et al., 2002, *AJ*, 124, 1810
 Sunyaev R. A., Zel'dovich Ya. B., 1970, *Ap&SS*, 7, 3
 Swanson M. E. C., Tegmark M., Blanton M., Zehavi I., 2008, *MNRAS*, 385,

1635

- Tegmark M. et al., 2004, ApJ, 606, 702
 Tegmark M. et al., 2006, Phys. Rev. D, 74, 123507
 Tucker D. L. et al., 2006, Astron. Nachr., 327, 821
 Wetterich C., 1988, B302, 645
 White M., 2005, Astropart. Phys., 24, 334
 York D. G. et al., 2000, AJ, 120, 1579
 Zehavi I. et al., 2002, ApJ, 571, 172
 Zehavi I. et al., 2005, ApJ, 630, 1
 Zlatev I., Wang L.-M., Steinhardt P. J., 1999, Phys. Rev. Lett., 82, 896

APPENDIX A: CALCULATION OF THE WINDOW FUNCTION

In this appendix, we describe the method used to calculate the mapping between the power spectra in the ‘true’ cosmology to be tested and the measured, or observed, power spectra where a Λ CDM model was used to convert redshifts to distances. This window function includes both the effect of the survey geometry and the mapping between cosmological models. As described by Percival et al. (2007c), we should expect the observed power spectrum to be a convolution of the true power spectrum with a window function:

$$P(k)_{\text{obs}} = \int dk' W(k, k') P(k')_{\text{true}}. \quad (\text{A1})$$

The goal of this section is to introduce a fast method by which $W(k, k')$ can be calculated for any model.

In Percival et al. (2007c), this window function was calculated using Monte Carlo realizations of Gaussian density fields, created assuming the cosmological model to be tested. These fields were then distorted as if they had been analysed assuming a Λ CDM model, and the power spectrum was calculated and compared with that input. Using a large number of simple input power spectra, we were able to construct the window function from this comparison. This procedure required significant computational resources as many density fields were needed in order to accurately measure the window function, limiting the number of models that could be tested. In particular, we were only able to consider cubic spline models of $D_V(z)$ with two nodes to three power spectra. With a faster window function calculation, we can include more nodes and fit to more power spectra.

For a survey covering a thin shell, the window function relating true and observed power is an offset delta function:

$$W(k, k') = \delta_D[k/k' - \epsilon], \quad (\text{A2})$$

where $\epsilon = d_p(\text{true})/d_p(\text{obs})$ is the ratio of proper distances in the true and observed cosmologies. Here we are simply stretching the true survey prior to measuring the power spectrum.

The obvious extension to surveys over a range of redshifts is to split the sample into i redshift shells and to approximate the window function as

$$W(k, k') = \sum_i \delta_D[k/k' - \epsilon_i] w_i, \quad (\text{A3})$$

where w_i is the weighted number of galaxy pairs in redshift shell i . Because we are now considering a broad survey, this pair weight is a function of pair separation. In this paper, we bin pairs of galaxies with a comoving separation of $90 h^{-1} \text{Mpc} < d_{\Lambda\text{CDM}} < 130 h^{-1} \text{Mpc}$, where $d_{\Lambda\text{CDM}}$ is the comoving distance in the Λ CDM cosmology used to convert galaxy redshifts to distances. The bin size was chosen to approximately match the BAO scale. For the SDSS LRG, main galaxy and combined samples, the galaxy pair-weights are shown in Fig. A1. We also need to allow for differences




Figure A1. The redshift dependence of galaxy pair-weights for the SDSS DR7 LRG and main galaxy samples, and from the combination of the two. These curves were calculated assuming a flat Λ CDM cosmology with $\Omega_m = 0.25$, $h = 0.72$ and $\Omega_b h^2 = 0.0223$.

in the orientation of galaxy pairs, as the distribution of ϵ_i should allow the galaxy pairs to be of all orientations. Including radial separations introduces an asymmetric convolution for ϵ_i , and we have found that this needs to be included in order to provide approximately the correct window function shapes. Note that equation (A3) is exact when there is a perfect dilation of scale between the true and observed cosmologies: such stretching of the windows can be perfectly represented by this equation.

For each ‘true’ cosmology to be tested, we can calculate the shift in scale that stretches each pair of galaxies because we do not measure BAO using this model. We have to allow for the




Figure A2. Galaxy pair-weights for the SDSS DR7 LRG and main galaxy samples, and from the combination of the two, as a function of comoving distance shifts. These were calculated assuming that a flat Λ CDM cosmology with $\Omega_m = 0.25$, $h = 0.72$ and $\Omega_b h^2 = 0.0223$ was used to analyse the data, while the BAO are present in a true cosmological model with the distance–redshift relation defined by a cubic spline in $D_V(z)$ with nodes at $z = 0.2$ and $z = 0.35$, with amplitude as shown in the plot.




Figure A3. Window functions for three values of k , calculated for the SDSS LRG, main galaxy and combined catalogues. Dotted lines represent the windows for our fiducial Λ CDM cosmology. The solid and dashed lines show the window functions, if the true cosmology were different, but the data were analysed assuming that the fiducial Λ CDM cosmology is correct. The solid lines were calculated using the procedure outlined in this Appendix. Dashed lines were calculated using the Monte Carlo procedure of Percival et al. (2007c).

angular shift caused by a change in $D_A(z)$ and the radial shift caused by the true and observed $H(z)$ being different. An example of the weighted distribution of ‘shifts’ expected for a model cosmology defined by a cubic spline in $D_V(z)$ with two nodes at $z = 0.2$ and $z = 0.35$ is shown in Fig. A2. Here the true cosmology has a distance–redshift relation given by a spline fit to $D_V(z)$, with nodes $D_V(z = 0.2) = 550 h^{-1}$ Mpc and $D_V(z = 0.35) = 1080 h^{-1}$ Mpc. The Λ CDM values are $D_V(z = 0.2) = 568 h^{-1}$ Mpc and $D_V(z = 0.35) = 949 h^{-1}$ Mpc, so at redshift $z = 0.2$, BAO in the true cosmology are stretched to larger scales by the analysis method, while those at redshift $z = 0.35$ are compressed to smaller scales. For the SDSS main galaxies, with median redshift close to $z \simeq 0.2$, $d_{\text{true}}/d_{\text{obs}} < 1$, while for the LRGs, with median redshift $z \simeq 0.35$, $d_{\text{true}}/d_{\text{obs}} > 1$.

For each ‘true’ cosmological model, the window function relating the true and observed power spectra was calculated by convolving

the standard window function for the Λ CDM model, by the distribution of shifts such as that shown in Fig. A2. For the models shown in Fig. A2, we have calculated the window function using the approximate method outlined in this Appendix and using the Monte Carlo method described by Percival et al. (2007c). A comparison of the windows is presented in Fig. A3. Reasonable agreement is found between the different methods: it is clear that the approximate method of splitting into shells recovers the main features of the window function. The agreement is not perfect, as expected given the approximate nature of our calculation. Because we analyse the data using a Λ CDM model, the window will be correct for this model, and will only deviate if we consider significantly different distance–redshift relations.

This paper has been typeset from a \LaTeX file prepared by the author.