

Baryon Asymmetry of the Universe in the Minimal Standard Model

Glennys R. Farrar and M. E. Shaposhnikov^(a)
TH Division, CERN, CH-1211, Geneva 23, Switzerland
 (Received 23 November 1992)

We calculate the baryon asymmetry of the Universe which would arise during a first-order electroweak phase transition due to minimal-standard-model processes. It agrees in sign and magnitude with the observed baryonic excess, for reasonable Kobayashi-Maskawa parameters and m_t in the expected range, and plausible values of bubble velocity and other high temperature effects.

PACS numbers: 98.80.Cq, 12.15.Ji

The existence in the present-day Universe of an excess of matter over antimatter has long been recognized to be a fundamental problem in cosmology and has widely been considered one of the most compelling pieces of evidence that the standard model is incomplete.

In this Letter we consider the possible production of the baryon asymmetry of the Universe (BAU) as a result of purely minimal-standard-model (MSM) processes, during the electroweak (EW) phase transition. Our calculation is realistic enough to make it clear that the baryon asymmetry arising from this mechanism can be responsible for the observed baryon density to entropy ratio, $n_B/s \sim (4-10) \times 10^{-11}$, for values of the top mass and Kobayashi-Maskawa (KM) parameters in the currently favored ranges.

For an excess of baryons to develop in a Universe which initially has zero baryon number, the following conditions, first enunciated by Sakharov, must be met: (1) Some interaction of elementary particles must violate baryon number, since the net baryon number of the Universe must change over time. (2) C and CP must be violated in order that there is not a perfect equality between rates of $\Delta B \neq 0$ processes, since otherwise no asymmetry could evolve from an initially symmetric state. (3) A departure from thermal equilibrium must play an essential role, since otherwise CPT would assure compensation between processes increasing and decreasing the baryon number.

We briefly summarize several features of the standard model which are necessary to understanding how these requirements will be met.

(1) In the standard model, quarks get their masses as a result of their Yukawa couplings to the Higgs field. When the Higgs field has a nonzero vacuum expectation value (VEV), quark masses are induced which are proportional to their couplings to the Higgs field, times its VEV. There are off-diagonal Yukawa interactions, in which quarks of different generations couple to one another through the Higgs field. In general the couplings are complex, and for three generations there is one physically significant phase, δ_{CP} . A nonzero value of δ_{CP} is imagined to be responsible for the CP violation observed in the kaon system. The KM matrix describes the mixing between generations and contains the phase of the Yukawa couplings.

(2) With three generations the phase in the KM matrix could be rotated away if any pair of quarks of the same charge were degenerate in mass, or if any of the mixing angles vanished. Thus CP -violating effects are significant in particle physics only when a relevant scale is small enough to be of the order of magnitude of the splitting in mass between, e.g., m_s and m_d . The K^0 system is an example of this, where the lack of degeneracy between d and s is evidenced by $m_K \neq m_\pi$. Moreover CP violation in the MSM vanishes together with $J \equiv \sin(\theta_{12}) \sin(\theta_{13}) \sin(\theta_{23}) \sin(\delta_{CP})$, using the Particle Data Group convention for the KM mixing angles.

(3) Although the standard-model Lagrangian conserves baryon number, quantum effects produce an anomaly which leads [1] to baryon-number-violating transitions. While the rate of these transitions is negligible at $T = 0$, at high temperature their rate $\sim \exp[-\frac{\pi M_W(T)}{\alpha_w T}]$ [2]. Thus above the electroweak phase transition the baryon-number-violating transition rate is rapid compared to the expansion rate of the Universe. We will require the VEV in the low temperature phase to be large enough that baryon violation is "turned off," allowing the asymmetry which has been produced during the transition to survive. In the MSM where the only undetermined parameter of the Higgs sector is the Higgs mass, this requirement may lead (if nonperturbative thermal effects are unimportant) to an upper limit on the Higgs mass [3].

At temperatures above the electroweak phase transition, the VEV of the Higgs field vanishes [4]; it takes on a constant, nonvanishing value in the low temperature phase. We require the phase transition to be first order, although this depends on the Higgs mass and is not certain to be true. During the phase transition, "bubbles" of the new $\langle \Phi \rangle \neq 0$ phase nucleate and expand to fill the Universe. This departure from thermal equilibrium satisfies the third Sakharov requirement. A number of phenomena could produce a baryon excess during the EW phase transition. For definiteness we concentrate on a specific mechanism; see [5] for others.

As a result of their thermal motion, quarks and antiquarks in the neighborhood of the bubble wall propagate through it. Since their masses result from their interaction with the VEV, they see the bubble wall as a potential barrier and scatter from it. We model this process in

detail [5], keeping the plasma masses of the quarks and antiquarks which originate from their interactions with the gauge and Higgs particles present in the heat bath, as well as treating quantum mechanically the process of their interaction with the bubble wall of the Higgs field. As a result of the spatial variation of the effective CP -violating phase, which comes about because the physical eigenstates depend on the interplay between flavor dependent thermal effects and the masses induced by the changing Higgs VEV, there can be a difference between the reflection and transmission coefficients of quarks and antiquarks. We report below on our computation of this asymmetry, in the one-dimensional problem which results when quark motion parallel to the bubble wall is ignored. Because of the fact that Lorentz invariance is broken by the thermal medium, momentum components parallel to the wall could be dynamically important, but that issue will be left for future work.

The total baryonic current is conserved in quark scattering with the bubble wall. Nonetheless, if there is an asymmetry in reflection and transmission coefficients the wall would separate particles and antiparticles, with, e.g., quarks flowing preferentially toward the low temperature phase and antiparticles toward the high temperature side. In the high temperature phase, sphaleron transitions operate to equilibrate the antiquark excess [2], converting most of them to quarks and leptons. But as long as the VEV in the low temperature phase is sufficiently large, the sphaleron transition rate is too low to keep up with the expansion rate of the Universe, and the quark excess is preserved until now. (The idea that the BAU could result from the separation of a quantum number by the bubble wall, combined with equilibration of this quantum number in the high temperature phase due to baryon-number-violating sphaleron processes, originated with Cohen, Kaplan, and Nelson, employing a lepton-number-violating interaction with the bubble wall [6]. The idea that MSM interactions of quarks and antiquarks with the bubble wall could directly cause a separation of baryon number is due to MS [7], where the element of including thermal effects is also introduced).

As noted above, if two same-sign quarks are degenerate in mass there is no CP violation, since the phase δ_{CP} can be removed from the KM matrix by a change in the definition of the overall phases of quark fields. This fact manifests itself in the present context by a tendency for different flavor eigenstates to have canceling contributions to the baryonic asymmetry current. For instance, if the reflection probability at a given energy for a d_L is greater than that for an anti- d_L , the reflection probability of an s_L will be less than that for an \bar{s}_L by a nearly identical amount. In the limit $m_s - m_d \rightarrow 0$ the compensation is perfect, when reflection involving b 's is included. An estimate of the residual CP -violating asymmetry for

typical quark energy $\sim T$ is $(m_t^2 - m_c^2)(m_t^2 - m_u^2)(m_c^2 - m_u^2)(m_b^2 - m_s^2)(m_b^2 - m_d^2)(m_s^2 - m_d^2)/T^{12}$ times J , $\sim 10^{-21}$ [3]. For this reason it has commonly been believed that the MSM cannot be responsible for the BAU.

The new observation of the present work is that the important quark energies are *not* $\sim T$, but rather energies such that the s quark is totally reflected but the d quark is partly reflected and partly transmitted, maximizing the dynamical difference in their contributions to the asymmetry current. Taking into account thermal effects on the quark propagation (see below and Ref. [5]), one finds that the p_t for which s quarks are reflected is $p_t \sim \frac{9\alpha_W}{32} \sqrt{\frac{3\pi}{2\alpha_s}} T \sim 10^{-1}T$. The fraction of p_t 's with s but not d reflection is $\sim \frac{m_s - m_d}{T}$.

The net baryonic flux through the wall is proportional to the group velocity of the quasiparticles perpendicular to the wall ($\sim \frac{1}{3}$) and to the asymmetry in Fermi distributions on the two sides of the wall due to its motion with respect to the thermal medium [$\sim 2v \frac{\partial \omega}{\partial p_t} \frac{\omega}{T} \sim \frac{2}{3}v \sqrt{\frac{2\pi\alpha_s}{3}}$ for small wall velocity v , where $\omega = \omega(p)$ is the quasiparticle dispersion relation]. Thus putting all the factors together we expect

$$n_b/s \sim \frac{2}{9} \sqrt{\frac{2\pi\alpha_s}{3}} \left(\frac{m_s - m_d}{T} \right) \frac{J v}{N_{\text{eff}}}, \tag{1}$$

where $N_{\text{eff}} \sim 100$ is the total number of degrees of freedom. Global fits to determine KM parameters place J in the range [8] $(1.4-5.0) \times 10^{-5}$, so we expect $n_b/s \sim (2-8) \times 10^{-11}v$, which can be of the right order of magnitude to account for the BAU.

Having outlined the way in which CP violation in the quark mixing matrix can lead to a present-day baryon asymmetry, we next calculate it more quantitatively, in the $q\bar{q}$ -separation mechanism. There are two essential effects to be included: The interactions of the q 's and \bar{q} 's with the plasma of gluons, EW gauge bosons, and Higgs bosons, and the quantum mechanical scattering of the q 's and \bar{q} 's from the bubble wall in the Higgs VEV. The effects of the interactions of the q 's and \bar{q} 's with the gauge and Higgs particles in the heat bath are most efficiently taken into account by changing variables to a quasiparticle description. The propagators of the quasiparticles have been determined to one-loop accuracy, neglecting internal masses, by Klimov [9] and Weldon [10]. The quasiparticles have interesting and unfamiliar behavior, but space limitations prevent us from describing them further here. These and other details are given in a longer article [5]. Here we record only the equation of motion which these quasiparticles obey in the wall rest frame, in the limit which is relevant to total reflection of the strange quark, momentum $\ll \omega$, treating only the motion normal to the bubble wall, and in the limit of small plasma velocity with respect to the wall:

$$\left(\begin{array}{c} \omega(1 + \alpha_L + \beta_L) + i \frac{\partial}{\partial x}(1 + \alpha_L) \\ [KM_d(x)]^\dagger \end{array} \right) KM_d(x) \left(\begin{array}{c} L \\ R \end{array} \right) = 0, \tag{2}$$

where $\alpha_{L,R}$, $\beta_{L,R}$, and $M_d(x)$ are 3×3 matrices and L and R are 3-component spinors in flavor space. $M_d(x)$ is the Higgs-induced mass at T_c . E.g., for charge $-1/3$ quarks in the unbroken phase, in the gauge basis where the interactions of quarks with the W and Z are diagonal,

$$\alpha_L = -\frac{4\pi}{3} \left[\frac{\alpha_s}{6} + \frac{\alpha_w}{32} \left(3 + \frac{m_u^2 + K m_d^2 K^\dagger}{M_w^2} + \frac{\sin^2 \theta_w}{9} \right) \right] \frac{T^2}{\omega^2},$$

where m_u^2 and m_d^2 are diagonal matrices of the charge $+2/3$ and $-1/3$ masses at $T = 0$, and K is the KM matrix. For momenta small compared to ω , $\beta_{L,R} = 2\alpha_{L,R}$. More general expressions can be found in [5].

We can solve these equations analytically in two limits: no mixing and zero wall thickness with small mixing. For more realistic cases they must be solved numerically, although having the exact cases to verify the correctness and accuracy of our numerical solutions is very useful. Details of the analytical and numerical results are given in Ref. [5]. We find that when the energy is such that neither or both d and s quarks are totally reflected, the difference between reflection probabilities of quarks and antiquarks, after summing over all three flavors, is extremely small (less than our numerical integration accuracy of one part in 10^{10}). However, as expected on the basis of the heuristic discussion above, the asymmetry is substantial in the narrow energy window in which the s quark is totally reflected but the d is not.

Figure 1 shows Δ , the difference in the reflection probabilities for right chiral quarks and antiquarks incident from the unbroken phase, summed over flavors, in the interesting range of energies, taking $m_t = 150$ GeV, $m_c = 1.6$ GeV, $m_u = 0.005$ GeV, $m_b = 5$ GeV, $m_s = 0.15$ GeV, $m_d = 0.01$ GeV, $s_{12} = 0.22$, $s_{23} = 0.05$, $s_{13} = 0.007$, and $\sin(\delta_{CP}) = 1$. For the calculation of this figure the wall velocity v was zero and the wall thickness was $10/T_c$, a popular value. Taking the wall to be narrower does not significantly change the result; taking it a factor of 3 thicker increases the result by a factor of 2. This dependence on wall thickness is not surprising: even in the thin wall limit there is a nontrivial CP -conserving phase shift to interfere with the KM CP -violating phase. The asymmetry in the reflection probabilities increases when the effect of the flow of the thermal medium is included [5], so that Δ is almost a factor of 5 larger for $v = 0.25$ than it is for the $v = 0$ case shown in the figure.

The upper pair of peaks occupy the energy range in which the strange quark is totally reflected. Note that the width in energy of this region is ~ 0.1 GeV, just the mass of the strange quark at that temperature. The "notch" in the middle, of width $\sim 0.006 \sim m_d(T)$, is the region in which the down quark is also totally reflected and Glashow-Iliopoulos-Maini cancellation is perfect, as expected. The unfamiliar feature that total reflection occurs for a range of energies, rather than for all energies less than some value, results from the unusual properties of the quasiparticle dispersion relation




FIG. 1. Δ , the asymmetry in the reflection probabilities of right-chiral quarks and antiquarks incident from the unbroken phase, for zero wall velocity.

[5], but is not essential to our result. Our analytic calculation in thin-wall, small mixing approximation [5] provides an adequate description of reflection in this region. At lower energy there is another region of a different character, involving level crossing between d or s and b_R . It would not be present if mixing were absent. Its width is $\sim m_b(T) \sin(\theta_{23})$.

We have checked that the asymmetry vanishes as pairs of masses are brought together. When $m_s \rightarrow m_d$ it arises by the squeezing away of the width of the upper peaks and the diminution of the magnitude of the lower peak. When masses in the charge $+2/3$ sector are brought together, or $m_b \rightarrow m_{d,s}$, the magnitudes of the peaks decrease appropriately. We checked that the result vanishes as mixing angles are taken to zero, although in the physical range of θ_{23} the result is nonlinear, increasing by 40% as θ_{23} is changed from 0.05 to 0.06, and changing sign for $\theta_{23} \lesssim 0.03$. The m_t dependence is interesting: for low values, $m_t \lesssim 110$ GeV, the integrated asymmetry has the opposite sign as for $m_t \gtrsim 110$ GeV. It reaches its maximum value for $m_t \sim 210$ GeV, where it is more than 4 times greater than for $m_t = 150$ GeV, then decreases for larger m_t . For a more detailed discussion and additional figures see Ref. [5].

The net L baryonic current resulting from the asymmetry in reflection coefficients for R 's incident from the unbroken phase and L 's incident from the broken phase is obtained from Δ as follows [5]:

$$J_{CP}^L = \int \frac{d\omega}{2\pi} \left(n_F \left[\omega \left(1 - \frac{d\omega}{dp} v \right) \right] - n_F \left[\omega \left(1 + \frac{d\omega}{dp} v \right) \right] \right) \Delta(\omega).$$

Given the net baryonic current flowing through the bub-

ble wall, corresponding to a preferential flow of baryons in one direction and antibaryons in the other, we next wish to determine the resultant n_B/s , assuming sphaleron transitions operate on L chiral q 's and \bar{q} 's to balance the chemical potentials in the high temperature phase, but are completely suppressed in the low temperature phase. Suppose that the wall velocity is low, so that diffusion permits a back-current of baryon number to be established, which acts to replace the antibaryon number which is being destroyed in the high temperature phase at a rate Γ by the sphaleron transitions. For sufficiently low velocity the problem is essentially static and the result is [5] $n_B = 3J_{CP}^L f(\rho)$, where $\rho = \frac{3D\Gamma}{v^2}$. $f(\rho) = 1$ for $\rho \gg 1$ and $f(\rho) = \frac{4}{9}\rho$ for $\rho \ll 1$. The physical importance of ρ is clear since the typical distance from the wall, of a particle scattered at $t = 0$ into the unbroken phase, is $\sqrt{Dt} - vt$. Thus $\frac{D}{v^2}$ is the typical time in which the sphaleron transitions can act on that particle before it is enveloped by the expanding low temperature phase. The correct values to take for the diffusion length, D , and sphaleron rate and wall velocities are very uncertain, but can plausibly be such that $f(\rho) \sim 1$, though a suppression as large as 10^{-3} is also possible [5].

Since we have computed the current in one dimension, we divide by the one-dimensional entropy for the MSM particle content at the EW phase transition: $s_{1D} = \frac{73\pi T}{3}$. Taking $v = 0.25$ values for reflection probabilities and the Boltzman factors, we find

$$n_B/s \sim 4 \times 10^{-11} \left(\frac{J}{0.22 \times 0.05 \times 0.007} \right) f(\rho), \quad (3)$$

for $m_t = 150$ GeV, $\theta_{23} = 0.05$, and inverse wall thickness 10 GeV. (Until the result of the full three-dimensional calculation is known, one should take this result as an order-of-magnitude estimate.) Since the prediction increases rapidly for larger values of m_t and wall velocity, there seems to be ample margin within the favored ranges of these quantities (recent estimates place $0.1 < v < 1$; see [5] for references) to tolerate some suppression from the uncertain overall factor $Jf(\rho)$ and the difference between the one- and three-dimensional cases. Of course if there were a fourth generation with a comparable KM structure then the degeneracy between d and s would be irrelevant and would be replaced by the degeneracy between b and s , producing a large enhancement in comparison to Eq. (3).

While the sign of $\sin \delta_{CP}$ is not at present unambiguously determined [11], a positive sign is favored. In this case (3) correctly predicts a baryon, not antibaryon, excess. Changes in v and wall thickness, and changes of the poorly known m_t , θ_{23} , and θ_{13} within their favored ranges, do not change the sign of (3). Thus refinements

in the treatment of this problem may not modify the conclusion that minimal-standard-model interactions can be responsible for the BAU. The most crucial outstanding problems are those associated with our still-primitive ability to deal with the high temperature environment during the electroweak phase transition: sphaleron rate, wall velocity, quasiparticle scattering length, and propagators are obvious examples. See [5] for further discussion.

To summarize, we have argued that already known physics of the minimal standard model may explain the observed baryon asymmetry of the Universe. A quantitative calculation in a specific mechanism gives the correct sign and magnitude. The essential new ingredient is not overlooking those regimes of quark momenta in which the most degenerate pair of quarks have very different dynamical behavior. If this is the explanation for the baryon asymmetry of the Universe, then future precision comparisons between observation and theory will provide a powerful test of our understanding of the EW phase transition, as well as constrain the KM matrix and the masses and generation content of the standard model.

G.R.F. is grateful for the support and hospitality of CERN during a sabbatical visit in which most of this work was done, and M.E.S. thanks Rutgers University for its kind hospitality during the completion of the work. We have benefited from many helpful discussions with colleagues at both institutions. A. Terrano made particularly significant contributions. This research was supported in part by NSF-PHY-91-21039.

-
- (a) On leave of absence from Institute for Nuclear Research of Russian Academy of Sciences, Moscow 117312, Russia.
- [1] G. 't Hooft, Phys. Rev. D **14**, 3432 (1976).
 - [2] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, Phys. Lett. **155B**, 36 (1985).
 - [3] M. E. Shaposhnikov, Pis'ma Zh. Eksp. Teor. Fiz. **44**, 364 (1986) [JETP Lett. **44**, 465 (1986)].
 - [4] D.A. Kirzhnits and A.D. Linde, Phys. Lett. **72B**, 471 (1972).
 - [5] G. R. Farrar and M. E. Shaposhnikov, Technical Report No. RU-93-11, CERN-TH.6732/92, Rutgers/CERN, 1993.
 - [6] A. Cohen, D. Kaplan, and A. Nelson, Phys. Lett. B **245**, 561 (1990).
 - [7] M. E. Shaposhnikov, Phys. Lett. B **277**, 324 (1992); **282**, 483(E) (1992).
 - [8] Y. Grossman and Y. Nir (to be published).
 - [9] V. V. Klimov, Yad. Fiz. **33**, 1734 (1981) [Sov. J. Nucl. Phys. **33**, 934 (1981)].
 - [10] H. A. Weldon, Phys. Rev. D **26**, 2789 (1982).
 - [11] Y. Nir and H. Quinn, Phys. Rev. D **42**, 1473 (1990).