

Research Article

Bell's Inequalities, Superquantum Correlations, and String Theory

Lay Nam Chang, Zachary Lewis, Djordje Minic, Tatsu Takeuchi, and Chia-Hsiung Tze

Department of Physics, Virginia Tech, Blacksburg, VA 24061, USA

Correspondence should be addressed to Djordje Minic, dminic@vt.edu

Received 18 April 2011; Accepted 7 October 2011

Academic Editor: Yang-Hui He

Copyright © 2011 Lay Nam Chang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We offer an interpretation of superquantum correlations in terms of a “doubly” quantum theory. We argue that string theory, viewed as a quantum theory with two deformation parameters, the string tension α' , and the string coupling constant g_s , is such a superquantum theory that transgresses the usual quantum violations of Bell's inequalities. We also discuss the $\hbar \rightarrow \infty$ limit of quantum mechanics in this context. As a superquantum theory, string theory should display distinct experimentally observable supercorrelations of entangled stringy states.

1. Introduction

In this paper, we present an observation relating two fields of physics which are ostensibly quite remote, namely, the study of the foundations of quantum mechanics (QM) centered around the violation of the celebrated Bell inequalities [1–3] and string theory (ST) [4–6]. As is well known, the Bell inequalities, based on the assumption of classical local realism, are violated by the correlations of canonical QM [7–11]. This remarkable feature of QM is often called “quantum nonlocality,” though perhaps a misnomer. However, even quantum correlations, with their apparent “nonlocality,” are bounded and satisfy another inequality discovered by Cirel'son (Also spelled Tsirelson) [12]; see also [13]. The natural question that arises is as follows: do “super” quantum theories exist which predict correlations that transcend those of QM and thereby violate the Cirel'son bound? Popescu and Rohrlich have demonstrated that such “super” correlations can be consistent with relativistic causality (aka the no-signaling principle) [14]. But what theory would predict them? In the following, we give heuristic arguments which suggest that nonperturbative ST may precisely be such a “superquantum” theory.

2. Bell's Inequality, the Cirel'son Bound, and Beyond

Consider two classical variables A and B , which represent the outcomes of measurements performed on some isolated physical system by detectors 1 and 2 placed at two causally disconnected spacetime locations. Assume that the only possible values of both A and B are ± 1 . Denote the state of detector 1 by a and that of detector 2 by b . "Local realism" demands that A depend only on a and B depend only on b . They can also depend on some hidden, but shared, information, λ . The correlation between $A(a, \lambda)$ and $B(b, \lambda)$ is then

$$P(a, b) = \int d\lambda \rho(\lambda) A(a, \lambda) B(b, \lambda), \quad \int d\lambda \rho(\lambda) = 1, \quad (2.1)$$

where $\rho(\lambda)$ is the probability density of the hidden information λ . This classical correlation is bounded by the following form of Bell's inequality [1, 2] as formulated by Clauser, Horne, Shimony, and Holt (CHSH) [3]:

$$|P(a, b) + P(a, b') + P(a', b) - P(a', b')| \leq X, \quad \text{where } X = X_{\text{Bell}} = 2. \quad (2.2)$$

The quantum versions of these correlations violate this bound but are themselves bounded by a similar inequality obtained by replacing the value of X on the right-hand side with $X_{\text{QM}} = 2\sqrt{2}$. This is the famous Cirel'son bound [12, 13], the extra factor of $\sqrt{2}$ being determined by the Hilbert space structure of QM. The same Cirel'son bound has been shown to apply for quantum field theoretic (QFT) correlations also [15, 16].

Let us briefly review the simplest routes to these bounds. Following [12, 13, 17], consider 4 classical stochastic variables $A, A', B,$ and B' , each of which takes values of $+1$ or -1 . Obviously, the quantity

$$C \equiv AB + AB' + A'B - A'B' = A(B + B') + A'(B - B') \quad (2.3)$$

can be only $+2$ or -2 , and thus, the absolute value of its expectation value is bounded by 2

$$|\langle C \rangle| = |\langle AB + AB' + A'B - A'B' \rangle| \leq 2. \quad (2.4)$$

This is the classical Bell bound. For the quantum case, we replace the classical stochastic variables with hermitian operators acting on a Hilbert space. Following [12, 13], we find that if $\hat{A}^2 = \hat{A}'^2 = \hat{B}^2 = \hat{B}'^2 = 1$ and $[\hat{A}, \hat{B}] = [\hat{A}, \hat{B}'] = [\hat{A}', \hat{B}] = [\hat{A}', \hat{B}'] = 0$, then C is replaced by

$$\hat{C} = \hat{A}\hat{B} + \hat{A}\hat{B}' + \hat{A}'\hat{B} - \hat{A}'\hat{B}', \quad (2.5)$$

from which we find

$$\hat{C}^2 = 4 - [\hat{A}, \hat{A}'] \cdot [\hat{B}, \hat{B}']. \quad (2.6)$$

When the commutators are zero, we recover the classical bound of 2. If they are not, we can use the uncertainty relations $|\langle i[\hat{A}, \hat{A}'] \rangle| \leq 2\|\hat{A}\| \cdot \|\hat{A}'\|$ and $|\langle i[\hat{B}, \hat{B}'] \rangle| \leq 2\|\hat{B}\| \cdot \|\hat{B}'\|$ to obtain

$$\langle \hat{C}^2 \rangle \leq 4 + 4 \|\hat{A}\| \cdot \|\hat{A}'\| \cdot \|\hat{B}\| \cdot \|\hat{B}'\| = 8 \rightarrow |\langle \hat{C} \rangle| \leq \sqrt{\langle \hat{C}^2 \rangle} \leq 2\sqrt{2}, \quad (2.7)$$

which is the Cirel'son bound. Alternatively, we can follow [17] and let $\hat{A}|\psi\rangle = |A\rangle$, $\hat{B}|\psi\rangle = |B\rangle$, $\hat{A}'|\psi\rangle = |A'\rangle$, and $\hat{B}'|\psi\rangle = |B'\rangle$. These 4 vectors all have unit norms and

$$|\langle \hat{C} \rangle| = |\langle \psi | \hat{C} | \psi \rangle| = |\langle A | B + B' \rangle + \langle A' | B - B' \rangle| \leq \| |B\rangle + |B'\rangle \| + \| |B\rangle - |B'\rangle \|, \quad (2.8)$$

which implies

$$|\langle \hat{C} \rangle| \leq \sqrt{2(1 + \text{Re}\langle B | B' \rangle)} + \sqrt{2(1 - \text{Re}\langle B | B' \rangle)} \leq 2\sqrt{2}. \quad (2.9)$$

This second proof suggests that the Cirel'son bound is actually independent of the requirement of relativistic causality. If relativistic causality is broken, then the \hat{A} 's and \hat{B} 's will not commute. Then, \hat{C} must be symmetrized as

$$\hat{C} = \frac{1}{2} \left[(\hat{A}\hat{B} + \hat{B}\hat{A}) + (\hat{A}\hat{B}' + \hat{B}'\hat{A}) + (\hat{A}'\hat{B} + \hat{B}\hat{A}') - (\hat{A}'\hat{B}' + \hat{B}'\hat{A}') \right], \quad (2.10)$$

to make it hermitian, and its expectation value will be

$$\langle \hat{C} \rangle = \text{Re}[\langle A | B + B' \rangle + \langle A' | B - B' \rangle], \quad (2.11)$$

which is clearly subject to the same bound as before. So, it is the Hilbert space structure of QM alone which determines this bound.

Indeed, Popescu and Rohrlich have demonstrated that one can concoct superquantum correlations which violate the Cirel'son bound, while still maintaining consistency with relativistic causality [14]. However, such superquantum correlations are also bounded, the value of X in (2.2) being replaced not by $X_{\text{QM}} = 2\sqrt{2}$ but by $X = 4$

$$|P(a, b) + P(a, b') + P(a', b) - P(a', b')| \leq 4. \quad (2.12)$$

Note, though, that this is not a "bound" per se, the value of 4 being the absolute maximum that the left-hand side can possibly be, since each of the 4 terms has its absolute value bounded by one. If the four correlations represented by these 4 terms were completely independent, then, in principle, there seems to be no reason why this bound cannot be saturated.

But what type of theory would predict such correlations? It has been speculated that a specific superquantum theory could essentially be derived from the two requirements of relativistic causality and the saturation of the $X = 4$ bound, in effect elevating these

requirements to the status of “axioms” which define the theory [14]. On the other hand, it has also been proposed that relativistic causality and locality would demand the Cirel’son bound, and thus QM would be uniquely derived [18, 19]. This would imply the necessity of nonlocality to achieve $X = 4$. However, to our knowledge, no concrete realization of either of these programs has thus far emerged.

A related development has been the proof by van Dam that superquantum correlations which saturate the $X = 4$ bound can be used to render all communication complexity problems trivial [20, 21]. Subsequently, Brassard et al. discovered a protocol utilizing correlations with $X > X_{cc} = 4\sqrt{2}/3$, which solves communication complexity problems trivially in a probabilistic manner [22]. Due to this, it has been speculated that nature somehow disfavors superquantum theories and that superquantum correlations, especially those with $X > X_{cc}$, should not exist [23–26]. However, the argument obviously does not preclude the existence of superquantum theories itself.

One proposal for a superquantum theory discussed in the literature uses a formal mathematical redefinition of the norms of vectors from the usual ℓ^2 norm to the more general ℓ^p norm [27]. In a 2D vector space with basis vectors $\{\mathbf{e}_1, \mathbf{e}_2\}$, the ℓ^p norm is

$$\|\alpha\mathbf{e}_1 + \beta\mathbf{e}_2\|_p = \sqrt[p]{\alpha^p + \beta^p}. \quad (2.13)$$

If one identifies $|B\rangle = \mathbf{e}_1$ and $|B'\rangle = \mathbf{e}_2$, then

$$\||B\rangle \pm |B'\rangle\|_p = 2^{1/p}. \quad (2.14)$$

Equation (2.12) would then be saturated for the $p = 1$ case. (The ℓ^1 norm and ℓ^∞ norm are equivalent in 2D, requiring a mere 45° rotation of the coordinate axes to get from one to the other.) Unfortunately, it is unclear how one can construct a physical theory based on this proposal in which dynamical variables evolve in time while preserving total probability.

At this point, we make the very simple observation that it is the procedure of “quantization,” which takes us from classical mechanics to QM, that increases the bound from the Bell/CHSH value of 2 to the Cirel’son value of $2\sqrt{2}$. That is, “quantization” increases the bound by a factor of $\sqrt{2}$. Thus, if one could perform another step of “quantization” onto QM, would it not lead to the increase of the bound by another factor of $\sqrt{2}$, thereby take us from the Cirel’son value of $2\sqrt{2}$ to the ultimate 4? This is the main conjecture of this paper, that is, a “doubly” quantized theory would lead to the violation of the Cirel’son bound.

In the following, we will clarify which “quantization” procedure we have in mind, and how it can be applied for a second time onto QM, leading to a “doubly quantized” theory. We then argue that a physical realization of such a theory may be offered by nonperturbative open string field theory (OSFT).

3. “Double” Quantization and Open String Field Theory

Before going into the “double quantization” procedure, let us first observe that from the point of view of general mathematical deformation theory [28, 29], QM is a theory with one deformation parameter \hbar , while ST is a theory with two: the first deformation parameter of ST is the world-sheet coupling constant α' , which measures the essential nonlocality of the string, and is responsible for the organization of perturbative ST. The second deformation parameter

of ST is the string coupling constant g_s , which controls the nonperturbative aspects of ST, such as D-branes and related membrane-like solitonic excitations and the general nonperturbative string field theory (SFT) [4–6]. Therefore, ST can be expected to be more “quantum” in some sense than canonical QM, given the presence of the second deformation parameter.

Second, superquantum correlations point to a nonlocality, which is more nonlocal, so to speak, than the aforementioned “quantum nonlocality” of QM and QFT. However, QFT’s are actually local theories, and true nonlocality is expected only in theories of quantum gravity. That quantum gravity must be nonlocal stems from the requirement of diffeomorphism invariance, as has been known from the pioneering days of that field [30, 31]. Thus, quantum gravity, for which ST is a concrete example, can naturally be expected to lead to correlations more nonlocal than those in QM/QFT.

Third, the web of dualities discovered in ST [4–6], which points to the unification of QFTs in various dimensions, can themselves be considered a type of “correlation” which transcends the barriers of QFT Lagrangians and spacetime dimensions. Again, the evidence suggests “super” correlations, perhaps much more “super” than envisioned above, in the context of ST.

What follows is a heuristic attempt to make these expectations physically concrete. Our essential observation is as follows: the “quantization” procedure responsible for turning the classical Bell bound of 2 into the quantum Cirel’son bound of $2\sqrt{2}$ is given by the path integral over the classical dynamical variables, which we collectively denote as x . That is, given a classical action $S(x)$, functions of x are replaced by their expectation values defined via the path integral

$$f(x) \longrightarrow \langle f(\hat{x}) \rangle = \int Dx f(x) \exp\left[\frac{i}{\hbar} S(x)\right], \quad (3.1)$$

up to a normalization constant. In particular, the correlation between two observables $\hat{A}(a)$ and $\hat{B}(b)$ will be given by

$$\langle \hat{A}(a)\hat{B}(b) \rangle = \int Dx A(a, x)B(b, x) \exp\left[\frac{i}{\hbar} S(x)\right] \equiv A(a) \star B(b) \quad (3.2)$$

(cf. (2.1)). In a similar fashion, we can envision taking a collection of quantum operators, which we will collectively denote by $\hat{\phi}$, for which a “quantum” action $\tilde{S}(\hat{\phi})$ is given and define another path integral over the quantum operators $\hat{\phi}$

$$F(\hat{\phi}) \longrightarrow \langle\langle F(\hat{\phi}) \rangle\rangle = \int D\hat{\phi} F(\hat{\phi}) \exp\left[\frac{i}{\hbar} \tilde{S}(\hat{\phi})\right], \quad (3.3)$$

and the correlation between two “super” observables will be

$$\langle\langle \hat{A}(a)\hat{B}(b) \rangle\rangle = \int D\hat{\phi} \hat{A}(a, \hat{\phi})\hat{B}(b, \hat{\phi}) \exp\left[\frac{i}{\hbar} \tilde{S}(\hat{\phi})\right]. \quad (3.4)$$

Note that the expectation values here, denoted $\langle\langle * \rangle\rangle$, are not numbers but operators themselves. To further reduce it to a number, we must calculate its expectation value in the usual way

$$\langle\langle \hat{A}(a)\hat{B}(b) \rangle\rangle \rightarrow \langle\langle\langle \hat{A}(a)\hat{B}(b) \rangle\rangle\rangle = \left\langle \int D\hat{\phi} \hat{A}(a, \hat{\phi}) \hat{B}(b, \hat{\phi}) \exp\left[\frac{i}{\tilde{\hbar}} \tilde{S}(\hat{\phi})\right] \right\rangle, \quad (3.5)$$

which would amount to replacing all the products of operators on the right-hand side with their first-quantized expectation values, or equivalently, replacing the operators with “classical” variables except with their products defined via (3.2).

This defines our “double quantization” procedure, through which two deformation parameters, \hbar and $\tilde{\hbar}$, are introduced. We would like to emphasize that the $\hat{\phi}$ in the above expressions is already a quantum entity, depending on the first deformation parameter \hbar . Thus, the “double quantization” procedure proposed here is quite distinct from the “second quantization” procedure used in QFT, which, being a single quantization procedure of a classical field, is a misnomer to begin with. The caveats to our definition are, of course, the difficulty in precisely defining the path integral over the quantum operator $\hat{\phi}$, and thus doing any actual calculations with it, and imposing a physical interpretation on what is meant by the quantum operators themselves being probabilistically determined.

At this point, we make the observation that a “doubly quantized” theory may already be available in the form of Witten’s open string field theory (OSFT) [32]. Our “double” quantization procedure can be mapped onto ST as follows: in the first step, the classical action $S(x)$ can be identified with the world-sheet Polyakov action and the first deformation parameter \hbar with the world-sheet coupling α' [4–6]. In the second step, the quantum action $\tilde{S}(\hat{\phi})$ can be identified with Witten’s OSFT action [32] and the second deformation parameter $\tilde{\hbar}$ with the string coupling g_s .

The doubly deformed nature of the theory is explicit in the Witten action for the “classical” open string field Φ , an action of an abstract Chern-Simons type

$$S_W(\Phi) = \int \Phi \star Q_{\text{BRST}} \Phi + \Phi \star \Phi \star \Phi, \quad (3.6)$$

where Q_{BRST} is the open string theory BRST cohomology operator ($Q_{\text{BRST}}^2 = 0$) and the star product is determined via the world-sheet Polyakov action

$$S_P(X) = \frac{1}{2} \int d^2\sigma \sqrt{g} g^{ab} \partial_a X^i \partial_b X^j G_{ij} + \dots, \quad (3.7)$$

and the corresponding world-sheet path integral

$$F \star G = \int DX F(X) G(X) \exp\left[\frac{i}{\alpha'} S_P(X)\right]. \quad (3.8)$$

The fully quantum OSFT is then in principle defined by yet another path integral in the infinite dimensional space of the open string field Φ ; that is,

$$\int D\Phi \exp\left[\frac{i}{g_s} S_W(\Phi)\right], \quad (3.9)$$

with all products defined via the star-product.

In addition to its manifestly “doubly” quantized path integral, OSFT has as massless modes the ordinary photons, which are used in the experimental verification of the violation of Bell’s inequalities [7–11], and it also contains gravity (closed strings) as demanded by unitarity. (The open/closed string theory duality is nicely illustrated by the AdS/CFT duality [4–6]. It is interesting to contemplate the Bell bound and its violations, both quantum and super-quantum, in this well-defined context. Similarly, it would be interesting, even though experimentally prohibitive, to contemplate the superquantum correlations for the QCD string, perhaps in the studies of the quark-gluon plasma.) Thus, our heuristic reasoning suggests that OSFT may precisely be an example of a super-quantum theory, which violates the Cirel’son bound.

We close this section with a caveat and a speculation. In the above reasoning, the two quantizations were taken to be independent with two independent deformation parameters. In the case of OSFT, they were α' and g_s . However, from the point of view of M-theory, we would generically expect that α' and g_s are both of order one (in natural units) and that both are dynamically generated [33]. Thus, the two parameters are not completely independent, and it may not be correct to view OSFT as a fully “doubly quantized” theory. Would this mean that OSFT/M-theory correlations would not saturate the ultimate $X = 4$ bound? Would its CHSH bound be situated somewhere between $X_{QM} = 2\sqrt{2}$ and $X = 4$, perhaps below the communication complexity bound of $X_{cc} = 4\sqrt{2/3}$? If M-theory is indeed unique, it may be natural to expect that its correlations would also be unique from the point of view of communication complexity, and that they would saturate this communication complexity bound. Of course, this conjecture would be testable only in a very precise proposal for M-theory (perhaps along the lines of [34–38]).

4. The $\hbar \rightarrow \infty$ Limit

Given that a superquantum theory is supposedly more “quantum” than QM, let us now consider the the extreme quantum limit of QM, $\hbar \rightarrow \infty$. Though QM is not “doubly quantized,” could it still exhibit certain superquantum behavior in that limit? Taking a deformation parameter to infinity can be naturally performed in ST, either $\alpha' \rightarrow \infty$ or $g_s \rightarrow \infty$, and one can still retain sensible physics. Therefore, the $\hbar \rightarrow \infty$ limit of QM may also be a sensible theory but at the same time quite different from QM. After all, if the $\hbar \rightarrow 0$ limit is to recover classical mechanics, with the Bell bound of $X_{Bell} = 2$, and apparently quite different from QM, it may not be too farfetched to conjecture that the $\hbar \rightarrow \infty$ limit would flow to a superquantum theory, with the superquantum bound of $X = 4$. If this were indeed the case, it may provide us with an opportunity to explore superquantum behavior in the absence of a solution to OSFT/M-theory.

What would the $\hbar \rightarrow \infty$ limit mean from the point of view of the path integral? Given that the path-integral measure is $e^{iS/\hbar}$, in the $\hbar \rightarrow \infty$ limit this measure will be unity for any S , and all histories in the path integral contribute with equal unit weight. Similarly all phases,

measured by $e^{iS/\hbar}$, will be washed out (this immediately raises other issues, such as the meaning of quantum statistics). Because the phases are washed out, we cannot distinguish between $|B\rangle + |B'\rangle$ and $|B\rangle - |B'\rangle$ (note that $-1 = e^{i\pi}$ and that sign can be absorbed into a phase of $|B'\rangle$). This suggests that

$$\| |B\rangle \pm |B'\rangle \| = \| |B\rangle \| + \| |B'\rangle \|, \quad (4.1)$$

which, if applied to the proof of the Cirel'son bound given earlier, leads to the superquantum bound of 4. This property is similar to what was obtained by replacing the ℓ^2 norm with an ℓ^1 (or ℓ^∞) norm, *cf.* (2.14), but presumably, unlike the change of norm, this relation is independent of the choice of basis. This argument seems to suggest that the $\hbar \rightarrow \infty$ limit is indeed superquantum.

However, this observation is perhaps a bit naïve, since the proof of the Cirel'son bound itself may no longer be valid under the wash-out of all phases. Let us invoke here an optical-mechanical analogy: geometric optics is the zero wavelength limit of electromagnetism, which would correspond to the $\hbar \rightarrow 0$ limit of QM. The $\hbar \rightarrow \infty$ limit of QM would, therefore, correspond to the extreme near field limit of electromagnetism, and in that case, the superposition of waves is washed out (we thank Jean Heremans for discussions of this point). Note also that from a geometric point of view, the holomorphic sectional curvature $2/\hbar$ of the projective Hilbert space CP^N of canonical QM goes to zero as $\hbar \rightarrow \infty$, and CP^N becomes just C^N . (For a general discussion of the geometry of quantum theory and its relevance for quantum gravity and string theory, see [34–38].) From these observations, it is clear that the usual Born rule to obtain probabilities will no longer apply.

But before we ask what rule should replace that of Born, let us confront the obvious problem that in the limit $\hbar \rightarrow \infty$, only the ground state of the Hamiltonian will remain in the physical spectrum and the theory will be rendered trivial (if the system has a non-trivial topology, it could allow for degeneracies in the ground state, and thus lead to a non-trivial theory even in the $\hbar \rightarrow \infty$ limit). This can also be argued via the general Feynman-Schwinger formulation of QM [39]

$$\delta S\psi = i\hbar \delta\psi. \quad (4.2)$$

By taking the $\hbar \rightarrow \infty$ limit, we eliminate the classical part δS so that we are left only with $\delta\psi = 0$, and thus, ψ must be a constant $\psi \equiv |\psi|$, a trivial result.

Could the $\hbar \rightarrow \infty$ limit of QM be made less than trivial? Consider the corresponding $\alpha' \rightarrow \infty$ and $g_s \rightarrow \infty$ limits in ST. In the $\alpha' \rightarrow \infty$ of ST, as opposed to the usual $\alpha' \rightarrow 0$ field theory limit, one seemingly ends up with an infinite number of fields and a nontrivial higher spin theory [40, 41]. Recently, such a theory was considered from a holographically dual point of view, and the dual of such a higher spin theory in AdS space was identified to be a free field theory [42]. The $g_s \rightarrow \infty$ limit of ST appears in the context of M-theory, one of whose avatars arises in the $g_s \rightarrow \infty$ limit of type-IIA ST [4–6]. Neither the high spin theory, nor the avatars of M-theory are trivial, as the presence of the tunable second deformation parameter saves them from triviality. Thus, the introduction of a second tunable parameter into QM, for example, Newton's gravitational constant G_N , may be necessary for the limit $\hbar \rightarrow \infty$ to be nontrivial.

Another issue here is that of interpretation: in the classical ($\hbar \rightarrow 0$) case, we have one trajectory, and one event (position, for example) at one point in time. One could speculate that the superquantum ($\hbar \rightarrow \infty$) limit would correspond to the complement of all other virtual trajectories. A general linear map relating virtual and classical trajectories is presumably non-symmetric (there are in principle more possibilities than actual events). Very naïvely, one would then expect that if we impose the condition that all possible events can be “mapped” to actual events, we could end up with a symmetric linear map corresponding to quantum theory ($\hbar \sim 1$), with a natural “map” between the actual events and possibilities, presumably realized by the Born probability rule. Note that according to this scenario, the superquantum theory would correspond essentially to a theory of possibilities and without actual events, which would be an interesting lesson for the foundations of ST.

5. Possible Experimental Signatures

Finally we offer some comments on possible experimental observations of such superquantum violations of Bell’s inequalities. The usual setup involves entangled photons [7–11]. In open ST, photons are the lowest lying massless states, but there is also an entire Regge trajectory associated with them. So, the obvious experimental suggestion would be to observe entangled Reggeized photons. Such an experiment is, of course, forbidding at present, given its Planckian nature.

Superquantum correlations could also be observable in cosmology. The current understanding of the large-scale structure of the universe, that is, the distribution of galaxies and galaxy clusters, is that they are seeded by quantum fluctuations. In standard calculations, it is assumed that the quantum correlations of these fluctuations are Gaussian (non-Gaussian correlations have also been considered). If the correlations were, in fact, superquantum, however, their signature could appear as characteristic deviations from the predicted large-scale structure based on Gaussian correlations. Such superquantum correlations would presumably be generated in the quantum gravity phase, and thus should be enhanced by the expansion of the universe at the largest possible scales. It would be interesting to look for evidence of such large-scale superquantum correlations in the existing WMAP [43] or the upcoming Planck [44, 45] data.

We conclude with a few words regarding a new experimental “knob” needed to test our doubly quantized approach to superquantum correlations. In the classic experimental tests of the violation of Bell’s inequalities [7–11], such a “knob” is represented by the relative angle between polarization vectors of entangled photons. If we have another quantization, there should be, in principle, another angle-like “knob.” Thus, the usual one-dimensional data plot [7–11] should be replaced by a two dimensional surface. By cutting this surface at various values of the new, second angle, we should be able to obtain one dimensional cuts for which the value of the CHSH bound varies depending on the cut, exceeding $2\sqrt{2}$ in some cases, and perhaps not exceeding 2 in others. Thus, the second “knob” may very well allow us to interpolate between the classical, quantum, and superquantum cases. The physical meaning of such an extra “knob” is not clear at the moment. It would be natural to associate this second “knob” with the extended nature of entangled Reggeized photons. However, we must admit that the measure of such nonlocality is not as obvious as the canonical measure of polarization of entangled photons in the standard setup [7–11].

In this paper, we have obviously only scratched the surface of a possible superquantum theory, and many probing questions remain to be answered and understood. We hope to address some of them in future works.





Acknowledgments

The authors thank V. Balasubramanian, J. de Boer, J. Heremans, S. Mathur, K. Park, J. Polchinski, R. Raghavan, D. Rohrlich, V. Scarola, J. Simon, and A. Staples for helpful comments, interesting discussions, and salient questions. D. Minic acknowledges the hospitality of the Mathematics Institute at Oxford University and Merton College, Oxford, and his respective hosts, Philip Candelas and Yang-Hui He. D. Minic also thanks the Galileo Galilei Institute for Theoretical Physics, Florence, for the hospitality and the INFN for partial support. Z. Lewis, D. Minic, and T. Takeuchi are supported in part by the U.S. Department of Energy Grant no. DE-FG05-92ER40677, task A.

References

- [1] J. S. Bell, "On the Einstein Podolsky Rosen paradox," *Physics*, vol. 1, no. 3, pp. 195–200, 1964.
- [2] J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics*, Cambridge University Press, 1987.
- [3] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, "Proposed experiment to test local hidden-variable theories," *Physical Review Letters*, vol. 23, no. 15, pp. 880–884, 1969.
- [4] J. Polchinski, *String Theory*, vol. 2, Cambridge University Press, 1998.
- [5] K. Becker, M. Becker, and J. H. Schwarz, *String Theory and M-Theory: A Modern Introduction*, Cambridge University Press, 2007.
- [6] M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory*, vol. 1, Cambridge University Press, 1988, where it was suggested that string field theory can be viewed as a generalization of quantum theory.
- [7] S. J. Freedman and J. F. Clauser, "Experimental test of local hidden-variable theories," *Physical Review Letters*, vol. 28, no. 14, pp. 938–941, 1972.
- [8] J. F. Clauser and M. A. Horne, "Experimental consequences of objective local theories," *Physical Review D*, vol. 10, no. 2, pp. 526–535, 1974.
- [9] A. Aspect, P. Grangier, G. Roger et al., "Experimental tests of realistic local theories via Bell's theorem," *Physical Review Letters*, vol. 47, no. 7, pp. 460–463, 1981.
- [10] A. Aspect, P. Grangier, G. Roger et al., "Experimental realization of Einstein-Podolsky-Rosen-Bohm gedankenexperiment: a new violation of Bell's inequalities," *Physical Review Letters*, vol. 49, no. 2, pp. 91–94, 1982.
- [11] A. Aspect, J. Dalibard, G. Roger et al., "Experimental test of Bell's inequalities using time-varying analyzers," *Physical Review Letters*, vol. 49, no. 25, pp. 1804–1807, 1982.
- [12] B. S. Cirel'son, "Quantum generalizations of Bell's inequality," *Letters in Mathematical Physics*, vol. 4, no. 2, pp. 93–100, 1980.
- [13] L. J. Landau, "On the violation of Bell's inequality in quantum theory," *Physics Letters A*, vol. 120, no. 2, pp. 54–56, 1987.
- [14] S. Popescu and D. Rohrlich, "Quantum nonlocality as an axiom," *Foundations of Physics*, vol. 24, no. 3, pp. 379–385, 1994.
- [15] S. J. Summers and R. Werner, "Maximal violation of Bell's inequalities is generic in quantum field theory," *Communications in Mathematical Physics*, vol. 110, no. 2, pp. 247–259, 1987.
- [16] S. J. Summers and R. Werner, "Bell's inequalities and quantum field theory. I. General setting," *Journal of Mathematical Physics*, vol. 28, no. 10, pp. 2440–2448, 1987.
- [17] D. Dieks, "Inequalities that test locality in quantum mechanics," <http://arxiv.org/abs/quant-ph/0206172>.
- [18] Y. Aharonov and D. Rohrlich, *Quantum Paradoxes: Quantum Theory for the Perplexed*, Wiley-VCH, 2005.
- [19] D. Rohrlich, "Three attempts at two axioms for quantum mechanics," <http://arxiv.org/abs/1011.5322>.
- [20] W. van Dam, *Nonlocality and communication complexity*, D.Phil. thesis, Department of Physics, University of Oxford, 2000.
- [21] W. Van Dam, "Implausible consequences of superstrong nonlocality," <http://arxiv.org/abs/quant-ph/0501159>.
- [22] G. Brassard, H. Buhrman, N. Linden, A. A. Méthot, A. Tapp, and F. Unger, "Limit on nonlocality in any world in which communication complexity is not trivial," *Physical Review Letters*, vol. 96, no. 25, Article ID 250401, 2006.

- [23] G. Brassard, "Is information the key?" *Nature Physics*, vol. 1, pp. 2–4, 2005.
- [24] S. Popescu, "Quantum mechanics: why isn't nature more non-local?" *Nature Physics*, vol. 2, no. 8, pp. 507–508, 2006.
- [25] J. Barrett, "Information processing in generalized probabilistic theories," *Physical Review A*, vol. 75, Article ID 032304, 2007.
- [26] N. Brunner and P. Skrzypczyk, "Nonlocality distillation and postquantum theories with trivial communication complexity," *Physical Review Letters*, vol. 102, no. 16, Article ID 160403, 4 pages, 2009.
- [27] G. Ver Steeg and S. Wehner, http://arxiv.org/PS_cache/arxiv/pdf/0811/0811.3771v2.pdf.
- [28] L. D. Faddeev, "On the Relationship between Mathematics and Physics," *Asia-Pacific Physics News*, vol. 3, pp. 21–22, June-July, 1988.
- [29] L. D. Faddeev, "Frontiers in physics, high technology and mathematics," in *Proceedings of the 25th Anniversary Conference*, H. A. Cerdeira and S. Lundqvist, Eds., pp. 238–246, World Scientific, Trieste, Italy, 1989.
- [30] B. S. Dewitt, "Quantum theory of gravity. I. the canonical theory," *Physical Review*, vol. 160, no. 5, pp. 1113–1148, 1967.
- [31] W. S. Bickel, "Mean lives of some excited states in multiply ionized oxygen and neon," *Physical Review*, vol. 162, no. 1, pp. 7–11, 1967.
- [32] E. Witten, "Non-commutative geometry and string field theory," *Nuclear Physics B*, vol. 268, no. 2, pp. 253–294, 1986.
- [33] R. Dijkgraaf, "The mathematics of M-theory," in *Proceedings of the 3rd European Congress of Mathematics*, Progress in Mathematics, Birkhuser, Barcelona, Spain, 2000.
- [34] D. Minic and H. C. Tze, "Background independent quantum mechanics and gravity," *Physical Review D*, vol. 68, no. 6, Article ID 061501, 5 pages, 2003.
- [35] D. Minic and C. H. Tze, "A general theory of quantum relativity," *Physics Letters B*, vol. 581, no. 1-2, pp. 111–118, 2004.
- [36] V. Jejjala, M. Kavic, and D. Minic, "Time and M-theory," *International Journal of Modern Physics A*, vol. 22, no. 20, pp. 3317–3405, 2007.
- [37] V. Jejjala, M. Kavic, D. Minic, and C. H. Tze, http://arxiv.org/PS_cache/arxiv/pdf/0804/0804.3598v1.pdf.
- [38] V. Jejjala, M. Kavic, D. Minic, and C. H. Tze, "On the origin of time and the universe," *International Journal of Modern Physics D*, vol. 25, no. 12, pp. 2515–2523, 2010.
- [39] L. M. Brown, Ed., *Feynman's Thesis: A New Approach to Quantum Theory*, World Scientific, 2005.
- [40] M. A. Vasiliev, http://arxiv.org/PS_cache/hep-th/pdf/9910/9910096v1.pdf.
- [41] X. Bekaert, S. Cnockaert, C. Iazeolla, and M. A. Vasiliev, http://arxiv.org/PS_cache/hep-th/pdf/0503/0503128v2.pdf.
- [42] M. R. Douglas, L. Mazzucato, and S. S. Razamat, http://arxiv.org/PS_cache/arxiv/pdf/1011/1011.4926v2.pdf.
- [43] E. Komatsu, K. M. Smith, J. Dunkley et al., "Seven-year wilkinson microwave anisotropy probe (Wmap) observations: cosmological interpretation," *Astrophysical Journal Supplement Series*, vol. 192, no. 2, article 18, 2011.
- [44] G. Efstathiou, C. Lawrence, and J. Tauber, Eds., Planck Collaboration, astro-ph/0604069.
- [45] P. A. R. Ade, N. Aghanim, M. Arnaud et al., Planck Collaboration, http://arxiv.org/PS_cache/arxiv/pdf/1101/1101.2022v2.pdf.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

