

Lawrence Berkeley National Laboratory

Recent Work

Title

BELL'S THEOREM WITHOUT HIDDEN VARIABLES

Permalink

<https://escholarship.org/uc/item/6qp4f35x>

Author

Eberhard, P.H.

Publication Date

1976-05-01

00004502819

Presented at the Communication to the
Thinkshop on Physics: Experimental
Quantum Mechanics, Erice, Sicily,
April 18 - 23, 1976

LBL-4888

c.1

BELL'S THEOREM WITHOUT HIDDEN VARIABLES

P. H. Eberhard

RECEIVED
BERKELEY LABORATORY


JUN 01 1976

May 1976

LIBRARY AND
DOCUMENTS SECTION

Prepared for the U. S. Energy Research and
Development Administration under Contract W-7405-ENG-48

For Reference
Not to be taken from this room



LBL-4888
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

BELL'S THEOREM WITHOUT HIDDEN VARIABLES*

P. H. Eberhard

Physics Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

MAY 1976

*This work was done with support from the U.S. Energy Research and Development Administration. Any conclusions or opinions expressed in this report represent solely those of the author and not necessarily those of the Regents of the University of California, the Lawrence Berkeley Laboratory or the United States Energy Research and Development Administration.

BELL'S THEOREM WITHOUT HIDDEN VARIABLES

P. H. Eberhard

Introduction As stated by Prof. d'Espagnat at this Conference, hidden variables theories are not the only theories in trouble with locality. Following ideas expressed in Refs. 1 and 2, I will demonstrate the CHSH inequality³ from locality alone without using either determinism or the concept of hidden variables. Then, I will comment about the violation of this inequality by quantum theory.

Let us consider two measuring apparatus located in two different places A and B. There is a knob a on apparatus A and a knob b on apparatus B. Since A and B are separated in space, it is natural to think that what will happen in A is independent of the setting of the knob b and vice versa. The principles of relativity seem to impose this point of view if the time at which the knobs are set and the time of the measurements are so close that, in the time laps, no light signal can travel from A to B or vice versa. Then, no signal can inform a measurement apparatus of what the knob setting of the other is. However, there are cases where the predictions of quantum theory make that independence assumption impossible. If quantum theory is true, there are cases where the results of the measurements in A will depend on the setting of knob b and/or the results of the measurements in B will depend on the setting of knob a.

Definitions A simple experiment consists of the recording of N events and each one of these events involves a measurement in both apparatus A and B. There are only two possibilities for the outcome of the measurement in each apparatus

and we label these measurement results as +1 and -1. The j 'th event corresponds to a response α_j in A and β_j in B. Each term α_j and β_j is either +1 or -1. We define the correlation by a statistical mean.

$$C = \frac{1}{N} \sum_j \alpha_j \beta_j = \langle \alpha_j \beta_j \rangle \quad (1)$$

where the symbol $\langle \rangle$ around a quantity designates the statistical mean of that quantity over j . C is equal to the fraction of events where α_j and β_j have the same signs minus the fraction where they have opposite signs.

There are two positions $a^{(1)}$ and $a^{(2)}$ of the knob a and two possible positions $b^{(1)}$ and $b^{(2)}$ of the knob b . C will depend on a and b . We define

$$\left. \begin{aligned} C^{(1,1)} &= \langle \alpha_j \beta_j \rangle \text{ if } a = a^{(1)}, b = b^{(1)} \\ C^{(2,1)} &= \langle \alpha_j \beta_j \rangle \text{ if } a = a^{(2)}, b = b^{(1)} \\ C^{(1,2)} &= \langle \alpha_j \beta_j \rangle \text{ if } a = a^{(1)}, b = b^{(2)} \\ C^{(2,2)} &= \langle \alpha_j \beta_j \rangle \text{ if } a = a^{(2)}, b = b^{(2)} \end{aligned} \right\} \quad (2)$$

The Locality Assumption We first have to suppose that it makes sense to consider the different results of a future experiment for different settings of the knobs a and b , although only one setting, at most, will actually be chosen for the actual experiment.⁴ Then, for a given position of b , we can write that α_j will be equal to $\alpha_j^{(1)}$ if $a = a^{(1)}$ and α_j will be equal to $\alpha_j^{(2)}$ if $a = a^{(2)}$, for the same event j but different settings of knob a . Similarly, for a given value of a , we define the values $\beta_j^{(1)}$ and $\beta_j^{(2)}$ of β_j if $b = b^{(1)}$ or $b^{(2)}$. Let's suppose, in addition, that the measurement α_j in A will not depend on b , and the measurement β_j in B will not depend on a . Then, for each event j , we can write

$$\left. \begin{aligned} \alpha_j &= \alpha_j^{(1)} \text{ if } a = a^{(1)} \\ \alpha_j &= \alpha_j^{(2)} \text{ if } a = a^{(2)} \end{aligned} \right\} \text{ whatever } b \text{ is} \\ \left. \begin{aligned} \beta_j &= \beta_j^{(1)} \text{ if } b = b^{(1)} \\ \beta_j &= \beta_j^{(2)} \text{ if } b = b^{(2)} \end{aligned} \right\} \text{ whatever } a \text{ is} \quad (3)$$

We refer to this assumption as the locality assumption.

THE CHSH Inequality Now we can define the following mathematical expression for event #j

$$\gamma_j = \alpha_j^{(1)} \beta_j^{(1)} + \alpha_j^{(2)} \beta_j^{(1)} + \alpha_j^{(1)} \beta_j^{(2)} - \alpha_j^{(2)} \beta_j^{(2)} \quad (4)$$

Each product $\alpha_j \beta_j$ is equal to ± 1 , therefore γ_j is an even integer. Furthermore, γ_j cannot be equal to 4 because $\alpha_j^{(2)} \beta_j^{(2)}$ is equal to the product of the first three terms of the right hand side of equation (4), and it is equal to +1 if the first three terms are positive. Therefore,

$$\gamma_j \leq 2 \quad (5)$$

It follows that

$$\frac{1}{N} \sum_j \gamma_j = \langle \gamma_j \rangle = \langle \alpha_j^{(1)} \beta_j^{(1)} \rangle + \langle \alpha_j^{(2)} \beta_j^{(1)} \rangle + \langle \alpha_j^{(1)} \beta_j^{(2)} \rangle - \langle \alpha_j^{(2)} \beta_j^{(2)} \rangle \leq 2 \quad (6)$$

From equations (2), (3) and (6), we derive

$$c^{(1,1)} + c^{(2,1)} + c^{(1,2)} - c^{(2,2)} \leq 2 \quad (7)$$

Equation (7) is the CHSH inequality³ that is a generalization of Bell's inequality.⁵

Quantum Theoretical Predictions Let's consider an idealized experiment where each event consists of the detection in A and B of two photons emitted in the direction of A and B by an atom located between A and B. The detection is made with the help of a polarization analyzer based on an idealized birefringent prism so that the polarization of the photon is measured with respect to the axis of the analyzer. We count $\alpha_j = +1$ when the photon is measured with one polarization in A and $\alpha_j = -1$ with the other. We count $\beta_j = \pm 1$ similarly, depending on the polarization of the photon measured in B.

Let a and b be the angles of the analyzers in A and B. The efficiencies are all 100%. There is a 100% correlation between the planes of polarization of the two photons at emission.

Quantum theory makes a prediction for the quantity C of Eq. (1):
if the number N of events is large enough

$$C \cong \cos(2a - 2b) \quad (8)$$

in almost all cases. (The sign \cong means approximately equal.)

Then, for

$$\left. \begin{aligned} a^{(1)} &= 45^\circ \\ a^{(2)} &= 0^\circ \\ b^{(1)} &= 22.5^\circ \\ b^{(2)} &= 67.5^\circ \end{aligned} \right\} \quad (9)$$

quantum theory predicts

$$C^{(1,1)} + C^{(2,1)} + C^{(1,2)} - C^{(2,2)} \cong 2\sqrt{2} > 2 \quad (10)$$

in almost all cases. This prediction is in contradiction with (7). Therefore, the predictions of quantum theory are incompatible with our locality assumptions.

Conclusions If quantum theory is correct, it is not allowed to think about the options of the future as resulting in a sequence α_j depending on the choice made for knob a only and a sequence β_j depending on the choice of knob b alone. The α_j 's will be different for different settings of b and/or the β_j 's will be different for different settings of a .

This demonstration has been performed without invoking either the concept of hidden variables, or the concept of determinism. We used only the results α_j and β_j 's of the measurements, the hypothetical values $\alpha_j^{(1)}$, $\alpha_j^{(2)}$, $\beta_j^{(1)}$ and $\beta_j^{(2)}$ that α_j and β_j would take for different knob settings⁴ and the

locality concepts as defined by Eqs. (3). The theorem affects more theories than just the hidden variable theories.⁶

Alternate Formulation There is an alternate way to formulate this theorem without using the concept of hypothetical results of experiments that will not be performed, but involving only the results of actual experiments. Let's consider the following thought experiment. The two photon experiment with N events is repeated a lot of times in all the four combinations of a and b defined by Eq. (9). What is the probability that one can find a set of four experiment, i.e., one for each of the four knob combinations, such that Eqs. (3) are valid?

We have to select experiment for different settings of knob b such that the lists of α_j 's are identical from $j = 1$ to $j = N$. Then we have to make a similar selection for β_j and knob a . N is finite. The probability to find a set of experiments associated to such sequences $\alpha_j^{(1)}$, $\alpha_j^{(2)}$, $\beta_j^{(1)}$ and $\beta_j^{(2)}$ is very small but, because of all the possible statistical fluctuations, it is not zero. However, if in addition we require that each correlation defined by Eq. (2) differs from the expectation value predicted by quantum theory by less than the quantity $\frac{\sqrt{2}-1}{2}$, then the probability to find such a set of four experiments is an absolute zero. Indeed, any set of sequences $\alpha_j^{(1)}$, $\alpha_j^{(2)}$, $\beta_j^{(1)}$ and $\beta_j^{(2)}$ that one introduces in Eqs. (3) will generate correlations C that satisfy inequality (7), and inequality (7) cannot be satisfied if all the correlations are different from $\cos(2a - 2b)$ by less than $\frac{\sqrt{2}-1}{2}$.

If we perform a lot of experiments with the different knob settings of Eqs. (9) and look for experiments that have identical α_j 's for different settings of knob b , and the same β_j 's for different settings of knob a , the only set of experiments one may find will contain experiments whose results are very different from the prediction of quantum theory. Actually, the differences between the

statistical mean C of Eq. (1) and the quantum theoretical predictions (Eq. (8)), in absolute value, will add up to at least $2(\sqrt{2}-1)$.

Acknowledgments I am indebted to Drs. J. S. Bell, A. Berthelot, J. F. Clauser, M. A. Horne, R. R. Ross and H. P. Stapp for fruitful discussions about this subject.

REFERENCES

1. See for instance H. P. Stapp, Phys. Rev. D 3, 1303 (1971) and H. P. Stapp, Lectures on Bell's Theorem, Trieste (1975).
2. J.S. Bell, Res. Th. 2053 - CERN The Theory of Local Beables, communication at the 6th Gift conference, Jaca, 2-7 June 1975; J. F. Clauser and M. A. Horne, Phys. Rev. D 10, 526 (1974); N. Herbert Am. Jour. Phys. 43, 315 (1975).
3. CHSH stands for J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett 23, 880 (1969).
4. That consideration is related to Stapp's concept of "contrafactual definiteness" (See Ref. 1). It would not make sense to talk about non-actual experiments if, for instance, we believed that there is no free will, and the choice by an experimenter is determined anyway. Otherwise, I believe that considering different possible results of a future experiemnt, depending on one's decision, is a rather natural way of thinking.
5. J. S. Bell, Physics (N.Y.), 1, 195 (1964).
6. Note that the conventional formalism of quantum theory also involves non-local evolutions of the wave function when the wave function collapses.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720