

REVIEW ARTICLE

Benign childhood focal epilepsies: assessment of established and newly recognized syndromes

Chrysostomos P. Panayiotopoulos, Michael Michael, Sue Sanders, Thalia Valeta and Michael Koutroumanidis

Department of Clinical Neurophysiology and Epilepsies, St Thomas' Hospital, Guy's and St Thomas NHS Foundation Trust, London, UK

Correspondence to: Michael Koutroumanidis, MD, Department of Clinical Neurophysiology and Epilepsies, St Thomas' Hospital, London SE1 7EH, UK
E-mail: michael.koutroumanidis@gstt.nhs.uk

A big advance in epileptology has been the recognition of syndromes with distinct aetiology, clinical and EEG features, treatment and prognosis. A prime and common example of this is rolandic epilepsy that is well known by the general paediatricians for over 50 years, thus allowing a precise diagnosis that predicts an excellent prognosis. However, rolandic is not the only benign childhood epileptic syndrome. Converging evidence from multiple and independent clinical, EEG and magnetoencephalographic studies has documented Panayiotopoulos syndrome (PS) as a model of childhood autonomic epilepsy, which is also common and benign. Despite high prevalence, lengthy and dramatic features, PS as well as autonomic status epilepticus had eluded recognition because emetic and other ictal autonomic manifestations were dismissed as non-epileptic events of other diseases. Furthermore, PS because of frequent EEG occipital spikes has been erroneously considered as occipital epilepsy and thus confused with the idiopathic childhood occipital epilepsy of Gastaut (ICOE-G), which is another age-related but rarer and of unpredictable prognosis syndrome. Encephalitis is a common misdiagnosis for PS and migraine with visual aura for ICOE-G. Pathophysiologically, the symptomatogenic zone appears to correspond to the epileptogenic zone in rolandic epilepsy (sensory-motor symptomatology of the rolandic cortex) and the ICOE-G (occipital lobe symptomatology), while the autonomic clinical manifestations of PS are likely to be generated by variable and widely spread epileptogenic foci acting upon a temporarily hyperexcitable central autonomic network. Rolandic epilepsy, PS, ICOE-G and other possible clinical phenotypes of benign childhood focal seizures are likely to be linked together by a genetically determined, functional derangement of the systemic brain maturation that is age related (benign childhood seizure susceptibility syndrome). This is usually mild but exceptionally it may diverge to serious epileptic disorders such as epileptic encephalopathy with continuous spike and wave during sleep. Links with other benign and age-related seizures in early life such as febrile seizures, benign focal neonatal and infantile seizures is possible. Overlap with idiopathic generalized epilepsies is limited and of uncertain genetic significance. Taking all these into account, benign childhood focal seizures and related epileptic syndromes would need proper multi-disciplinary re-assessment in an evidence-based manner.

Keywords: rolandic epilepsy; Panayiotopoulos syndrome; occipital epilepsy of Gastaut; autonomic seizures; autonomic status epilepticus; ictus emeticus; visual hallucinations; childhood seizure susceptibility; migraine, cyclic vomiting syndrome

Abbreviations: AED = anti-epileptic drugs; BCSSS = benign childhood seizure susceptibility syndrome; CTS = centro-temporal spikes; CSWS = continuous spike and wave during sleep; FOS = fixation off sensitivity; GSES = giant somatosensory evoked spikes; GTCS = generalized tonic-clonic seizures; ICOE-G = idiopathic childhood occipital epilepsy of Gastaut; ILAE = International League against Epilepsy; IPS = intermittent photic stimulation; OPLS = Oro-pharyngo-laryngeal symptoms; PS = Panayiotopoulos syndrome

Received April 14, 2008. Revised June 30, 2008. Accepted July 1, 2008

Benign childhood focal seizures and related idiopathic epileptic syndromes affect approximately 22% of children with non-febrile seizures and constitute a significant part of the everyday practice of paediatricians, neurologists and electroencephalographers. They comprise three identifiable electro-clinical syndromes recognized by the International League against Epilepsy (ILAE) (Engel, 2006): rolandic epilepsy which is well known, Panayiotopoulos syndrome (PS), a common autonomic epilepsy, which is currently more readily diagnosed and the idiopathic childhood occipital epilepsy of Gastaut (ICOE-G) including the idiopathic photosensitive occipital lobe epilepsy, a less common form with uncertain prognosis. There are also reports of children with benign focal seizures of predominantly affective symptoms, and claims have been made for other clinical phenotypes associated with specific inter-ictal EEG foci, such as frontal, midline or parietal, with or without giant somatosensory evoked spikes (GSES). Neurological and mental states and brain imaging are normal, though because of their high prevalence any type of benign childhood focal seizures may incidentally occur in children with neurocognitive deficits or abnormal brain scans. The most useful diagnostic test is the EEG. In clinical practice, the combination of a normal child with infrequent seizures and an EEG showing disproportionately severe spike activity is highly suggestive of these benign childhood syndromes (Panayiotopoulos, 1999a).

All these conditions may be linked together in a broad, age-related and age-limited, benign childhood seizure susceptibility syndrome (BCSSS), which may be genetically determined (Panayiotopoulos, 1993).

This appraisal is based on over 30 years clinical research, extensive review of English and other journals including relevant book chapters and personal communications with experts. Details of original studies, numerous case histories and published reports not cited here can be found in our previous reviews (Panayiotopoulos, 1999a, 2002, 2007; Koutroumanidis, 2007).

Rolandic epilepsy (benign childhood epilepsy with centro-temporal spikes)

Rolandic epilepsy is the best known and most common benign childhood focal epilepsy ((Beaussart and Faou, 1978; Loiseau *et al.*, 1988; Bouma *et al.*, 1997; Beaussart *et al.*, 1999; Panayiotopoulos, 1999a; Dalla Bernardina *et al.*, 2005; Wirrell *et al.*, 2006; Fejerman, 2008). The age of onset ranges from 1 to 14 years with 75% starting between 7 and 10 years. There is a 1.5 male predominance, prevalence is around 15% in children aged 1–15 years with non-febrile seizures and incidence is 10–20/100 000 children aged 0–15 years (Heijbel *et al.*, 1975; Cavazzuti, 1980; Sidenvall *et al.*, 1996; Astradsson *et al.*, 1998; Berg *et al.*, 1999; Larsson and Eeg-Olofsson, 2006).

Clinical manifestations

The cardinal features of rolandic epilepsy are focal seizures consisting of unilateral facial sensory-motor symptoms (30%

of patients), oro-pharyngo-laryngeal symptoms (OPLS) (53%), speech arrest (40%) and hypersalivation (30%) (Beaussart, 1972; Loiseau and Beaussart, 1973; Lerman and Kivity, 1975; Bouma *et al.*, 1997; Panayiotopoulos, 1999a; Dalla Bernardina *et al.*, 2005; Wirrell *et al.*, 2006; Fejerman, 2008). Ictal manifestations indicative of temporal lobe involvement do not occur in rolandic epilepsy, and the term ‘centro-temporal’ refers only to the spike topography where it is partly a misnomer (see EEG section below).

Hemifacial sensory-motor seizures are mainly localized in the lower lip and may spread to the ipsilateral hand. Motor manifestations are clonic contractions sometimes concurrent with ipsilateral tonic deviation of the mouth, and sensory symptoms consist of numbness in the corner of the mouth. OPLSs are unilateral sensory-motor symptoms of numbness or paraesthesias (tingling, prickling or freezing) inside the mouth, associated with strange sounds, such as death rattle, gargling, grunting and guttural sounds.

Hypersalivation, a prominent autonomic manifestation, is often associated with hemifacial seizures, OPLSs and speech arrest. In speech arrest, the child is actually anarthric, unable to utter a single intelligible word and attempts to communicate with gestures.

Consciousness and recollection are fully retained in more than half (58%) of rolandic seizures. In the remainder, consciousness becomes impaired during the ictal progress and in one-third there is no recollection of ictal events. Progression to hemiconvulsions or generalized tonic-clonic seizures (GTCS) occurs in around half of children and hemiconvulsions may be followed by postictal Todd’s hemiparesis (Panayiotopoulos, 1999a).

Three-quarters of rolandic seizures occur during non-REM (rapid eye movement) sleep, mainly at sleep onset or just before awakening.

Rolandic seizures are usually brief lasting for 1–3 min. Focal motor, hemiconvulsive and generalized convulsive status epilepticus are rare at around 5% (Deonna *et al.*, 1986; Wirrell *et al.*, 1995; Panayiotopoulos, 1999a). Opercular status epilepticus usually occurs in children with atypical evolution (Colamaria *et al.*, 1991; Deonna *et al.*, 1993; Fejerman *et al.*, 2000) or may be induced by carbamazepine or lamotrigine (Caraballo *et al.*, 1989; Parmeggiani *et al.*, 2004). This state lasts for hours to months and consists of ongoing unilateral or bilateral contractions of the mouth, tongue or eyelids, positive or negative subtle perioral or other myoclonus, dysarthria, speech arrest, difficulties in swallowing, buccofacial apraxia and hypersalivation.

Other seizure types

Despite prominent hypersalivation, focal seizures with primarily autonomic manifestations (autonomic seizures) are not considered part of the core clinical syndrome of rolandic epilepsy. However, some children may present with independent autonomic seizures or seizures with mixed rolandic-autonomic manifestations including

emesis (see below in the relations between rolandic epilepsy and PS).

Primarily GTCS are considered part of rolandic epilepsy by the ILAE (Engel, 2006) and their occurrence can not be excluded. However, from the published ictal recordings (Watanabe, 1996; Panayiotopoulos, 1999a; Wirrell *et al.*, 2006) and the electroclinically unequivocal focal nature of rolandic epilepsy, it can be inferred that at least the majority of the GTCS follow rolandic activation, and are therefore secondarily GTCS. Short-lived initial focal symptoms may pass unnoticed in daytime GTCS and are bound to be missed in nocturnal GTCS.

Typical absence seizures are considered rare (Beydoun *et al.*, 1992; Gelisse *et al.*, 1999; Panayiotopoulos, 1999a) though a high incidence of them has also been reported (Beaumanoir *et al.*, 1974).

Electroencephalography

By definition, centro-temporal spikes (CTS) are the hallmark of benign childhood epilepsy with CTS (Fig. 1). However, although called centro-temporal, these spikes are mainly localized in the C3 and C4 (high central) or C5 and C6 (low central) supra-sylvian and not temporal electrodes (Legarda *et al.*, 1994; Panayiotopoulos, 1999a). CTS are often bilateral and typically activated by drowsiness and slow (non-REM) sleep, but not by overbreathing (Smith and Kellaway, 1964; Blom and Brorson, 1966; Clemens and Majoros, 1987). In serial EEGs of the same child, CTS may occur right or left, infrequently or frequently, and appear small or giant, alone or with spikes in other locations. Rarely, children with rolandic epilepsy may have normal EEG or CTS may appear only during non-REM sleep (3–35%) (Panayiotopoulos, 1999a). The incidence of extra-rolandic spikes in rolandic epilepsy is not precisely known but may be significant when these are sought (Drury and Beydoun, 1991).

Dipole EEG (Gregory and Wong, 1992; Tsai and Hung, 1998; Jung *et al.*, 2003), magnetoencephalography (MEG) (Minami *et al.*, 1996; Huiskamp *et al.*, 2004) and functional MRI (Boor *et al.*, 2007) studies have demonstrated that the main negative spike component of CTS is usually modelled by a single and stable tangential dipole source with the negative pole maximum in the central region and the positive pole maximum in the frontal region.

Brief 1–3 s generalized bursts of 3–5 Hz slow waves with intermixed small spikes without associated overt clinical symptoms may occur in about 4% of patients with rolandic epilepsy (Gelisse *et al.*, 1999; Panayiotopoulos, 1999a). Typical 3 Hz spike-wave discharges are probably rare (see absence seizures above).

CTS are diagnostic markers of benign rolandic epilepsy only in a suggestive clinical presentation. Their frequency, location and persistence do not determine the clinical manifestations, severity and frequency of seizures or the prognosis. It is well established that CTS are not specific to

rolandic epilepsy (Kellaway, 1980; Panayiotopoulos, 1999a) as they:

- occur in 2–3% of normal school-aged children, of whom <10% develop rolandic seizures (Gibbs and Gibbs, 1967; Petersen and Eeg-Olofsson, 1971; Cavazzuti *et al.*, 1980; Okubo *et al.*, 1994).
- are common among relatives of children with rolandic epilepsy (Bray and Wiser, 1965; Bali *et al.*, 2007)
- may occur in a variety of organic brain diseases with or without seizures, such as cerebral tumours, Rett syndrome, fragile X syndrome and focal cortical dysplasia (Kellaway, 1980; Panayiotopoulos, 1999a)
- may incidentally be found in non-epileptic children with various symptoms, such as headache, speech and behavioural and learning difficulties (Gibbs and Gibbs, 1967).

Somatosensory stimulation is common form of activation of CTS (10–20%) (De Marco and Tassinari, 1981; Panayiotopoulos, 1999a; Fonseca and Tedrus, 2000; Kubota *et al.*, 2000; Langill and Wong, 2003) and evokes GSES (Fig. 1), which correspond to mid- or long-latency somatosensory evoked potentials (Manganotti *et al.*, 1998). GSES, like spontaneous CTS, occur in children with or without seizures and disappear with age.

There have been around 20 reported ictal EEGs of rolandic seizures showing an initial paucity of spontaneous CTS before the onset of the ictal discharge, which appears in the contralateral to the clinical manifestations rolandic regions and consists of slow waves intermixed with spikes (Panayiotopoulos, 1999a; Wirrell *et al.*, 2006; Fejerman *et al.*, 2007) (Fig. 2). GTCS, when occurred, were preceded by focal clinical and EEG features. (Watanabe, 1996; Panayiotopoulos, 1999a; Wirrell *et al.*, 2006).

Aetiology

Rolandic epilepsy is genetically determined although conventional genetic influences may be less important than other mechanisms (Vadlamudi *et al.*, 2004, 2006). There is evidence of linkage with chromosome 15q14 (Neubauer *et al.*, 1998). Autosomal dominant inheritance with age-dependent penetrance refers to the EEG CTS and not to the clinical syndrome of rolandic epilepsy (Bray and Wiser, 1965; Bali *et al.*, 2007).

Siblings or parents of patients with rolandic epilepsy may rarely have the same type of seizures or other phenotypes of BCSSS, such as PS. Reported occurrence of febrile seizures ranges from 10% to 20% (probably around 18%) of patients (Kajitani *et al.*, 1992).

Pathophysiology

As indicated by the distribution of CTS, the epileptogenic zone in rolandic epilepsy involves neuronal networks within the rolandic cortex surrounding the central fissure bilaterally. This is congruent with the seizure symptomatology




Fig. 1 Interictal EEG in rolandic epilepsy (top), PS (middle) and idiopathic childhood occipital epilepsy of Gastaut (bottom). **(A)** Interictal EEG with CTS of two children with rolandic epilepsy separated by the vertical line. Left: spontaneous and GSES elicited by the patient simultaneously tapping the tip of the fingers of both hands (arrows). Right: the same EEG sample at two different montages (extreme right is laplacian montage) to show that what appears as CTS are not temporal which on this occasion are localized in the lower central electrode (C5) of the laplacian montage. **(B)** Interictal EEG in five patients with PS separated by vertical lines. Despite similar clinical features, spikes are localized in the occipital, centro-temporal and frontal regions or they are frequently multifocal and may appear as clone-like, repetitive, spike-wave complexes. Brief generalized discharges of slow wave with small spikes (extreme right) are sometimes an interictal EEG feature. **(C)** Interictal EEG in two patients with ICOE-G separated by a vertical line. Left: Classical occipital paroxysms demonstrating FOS. Right: spontaneous scattered occipital spikes and occipital photosensitivity.




Fig. 2 Ictal EEG of rolandic (**A**), autonomic (**B**) and occipital lobe (**C**) seizures. (**A**) Ictal EEG of a rolandic seizure. The onset of ictal discharge becomes apparent at the right central regions (black arrow) and the first clinical events (open arrow) occur 12 s later and consist of hemifacial motor symptoms indicated by the muscle artefacts. The seizure ended with a GTCS. (**B**) Ictal EEG of an autonomic seizure. The onset of ictal discharge becomes apparent at the right frontal regions (black arrow) and the first clinical event (open arrow) occurs 12 min later and consists of coughing. The seizure was prolonged for over 40 min terminated with intravenous benzodiazepine. Modified with permission from Koutroumanidis *et al.* (2005) where further details can be found. (**C**) Ictal EEG of an occipital seizure. The onset of ictal discharge becomes apparent at both occipital regions (black arrow) nearly simultaneously with the first clinical events (open arrow) which consist of blurring of vision. The seizure also consisted of forced eyelid closures and lasted for less than a minute.

(symptomatogenic zone) and in agreement with those described by Penfield and Rasmussen (1957) during electrical stimulation of the lower part of the pre-central and post-central gyrus in man.

The speech arrest is due to anarthria attributed to loss of the power and coordination of the musculature responsible for the articulation of words. There is no impairment of the cortical language networks. Hypersalivation most probably relates to the involvement of the superior bank of the sylvian fissure (Luders *et al.*, 1987), but defining ictal symptomatology by plotting the simple topographic coordinates of an ictal discharge can hardly explain the high prevalence of hypersalivation in benign rolandic epilepsy compared to its exceptional only occurrence in adults with symptomatic foci of similar topography. Nor can it explain the opercular status epilepticus, with the lengthy for hours speech arrest, drooling and bilateral regional twitching that is associated with diffuse or bilateral rolandic spike wave activity, but does not propagate in a conventional way and does not involve other systems like for instance the motor strip or the language function. Therefore, at variance with the symptomatic adult focal epilepsies of comparable but more discretely localized topography, rolandic epilepsy reflects an age-related maturational instability of the lower rolandic (somatosensory) cortex that represents the face and the oropharynx bilaterally (Koutroumanidis, 2007).

Evolution and prognosis

The prognosis for rolandic seizures is almost invariably excellent, with probably <2% risk of developing absence seizures and less often GTCS in adult life (Beaumanoir *et al.*, 1974; Lerman and Kivity, 1975; Blom and Heijbel, 1982; Loiseau *et al.*, 1983; Panayiotopoulos, 1999a; Datta and Sinclair, 2007; Koutroumanidis *et al.*, 2008; Caraballo *et al.*, 2008b). Remission occurs within 2–4 years from onset and before the age of 16 years. The total number of seizures is low, the majority of patients having fewer than 10 seizures; 10–20% have just a single seizure. About 10–20% may have frequent seizures, but these also remit with age.

Children with rolandic seizures may develop usually mild and reversible linguistic, cognitive and behavioural abnormalities during the active phase of the disease (Giordani *et al.*, 2006; Nicolai *et al.*, 2006; Riva *et al.*, 2007; Kossoff *et al.*, 2007; Kavros *et al.*, 2008; Perkins *et al.*, 2008). These may be worse in children with onset of seizures before 8 years of age, high rate of occurrence and multifocal EEG spikes (Bulgheroni *et al.*, 2008; Piccinelli *et al.*, 2008; Boatman *et al.*, 2008). The effect of anti-epileptic drugs (AED), the impact of stigmatizing because of epilepsy, bias in selection of the most serious cases and other factors have not been excluded in most of these studies. The development, social adaptation and occupations of adults with a previous history of rolandic seizures was found normal (Blom and Heijbel, 1982; Loiseau *et al.*, 1983).

Rarely (<1%), rolandic epilepsy may evolve to more severe syndromes with linguistic, behavioural and neuropsychological deficits, such as Landau–Kleffner syndrome, atypical focal epilepsy of childhood or epilepsy with continuous spike and wave during sleep (CSWS) (Fejerman *et al.*, 2000) as explained in the relevant section of this assessment.

Panayiotopoulos syndrome

PS is a common, childhood-related, susceptibility to autonomic seizures confirmed in long term studies of over 800 children worldwide (Panayiotopoulos, 1981, 1988, 2002; Ferrie *et al.*, 1997; Oguni *et al.*, 1999; Kivity *et al.*, 2000; Lada *et al.*, 2003; Ohtsu *et al.*, 2003, 2008; Covanis *et al.*, 2003; Caraballo *et al.*, 2007; Dura-Trave *et al.*, 2008).

The appraisal of PS here is based on in-depth analysis of our own patients and review of the related literature up to 2002 (Panayiotopoulos, 2002). Subsequent reports of around 60 publications on this and related aspects have also been thoroughly studied.

PS is defined as ‘benign age-related focal seizure disorder occurring in early and mid-childhood. It is characterized by seizures, often prolonged, with predominantly autonomic symptoms, and by an EEG that shows shifting and/or multiple foci, often with occipital predominance’ (Ferrie *et al.*, 2006). ‘Early onset benign childhood occipital epilepsy’ often used as synonymous with PS (Taylor *et al.*, 2003, 2008; Engel, 2006) does not represent the wide clinical, EEG and pathophysiological spectrum of PS, which is far beyond the occipital neocortex (Martinovic, 2007).

Onset is from age 1 to 14 years with 76% starting between 3 and 6 years. Both sexes are probably equally affected, though a female preponderance was found in some studies (Lada *et al.*, 2003; Dura-Trave *et al.*, 2008). Prevalence of PS may be high though this is practically absent in designed controlled epidemiological studies (Berg *et al.*, 1999; Jallon *et al.*, 2001; Shinnar and Pellock, 2002; Cowan, 2002; Forsgren *et al.*, 2005), which is understandable as this syndrome was only recently formally recognized, its features imitate many other conditions and often manifests with a single seizure only. In the original cohort of Panayiotopoulos (1988), prevalence was around 13% in children aged 3–6 years with one or more non-febrile seizures, and 6% in the age group 1–15 years. These figures may be higher if children who are currently considered to have atypical clinical presentation are included in the syndrome (Panayiotopoulos, 2002; Covanis, 2006). PS is the most common specific cause of non-febrile, non-convulsive status epilepticus in childhood (Okanishi *et al.*, 2008).

Clinical manifestations

The hallmark of PS is ictal autonomic aberrations that may involve any function of the autonomic system and mainly emesis (70–80% of seizures). The following description of clinical manifestations of PS are based on a synthetic analysis of available clinical historical data as perceived by patients

and witnessed by observers from our records and those provided in the literature (Panayiotopoulos, 2002). Therefore, they may not accurately represent their true prevalence and sequence in PS.

Ictal autonomic symptoms

Seizures commonly commence with autonomic manifestations (80–90%), while consciousness and speech, as a rule, are preserved. Ictus emeticus (nausea and retching) culminates in vomiting in 74–82% of seizures; in others, only nausea or retching occurs and, in a quarter, emesis may not be apparent. Emesis is usually the first apparent ictal symptom, but it may also occur long after the onset of other manifestations. Other autonomic manifestations include pallor (28%), incontinence of urine (19%) and faeces (3%), hypersalivation (10%), cyanosis (12%), mydriasis (7%) and less often miosis (2%), coughing and abnormalities of intestinal motility (3%). Breathing (7%) and cardiac irregularities may be more common than reported. Tachycardia is usually found, sometimes at the onset, in ictal EEG (Beaumanoir, 1993b; Oguni *et al.*, 1999; Koutroumanidis *et al.*, 2005; Parisi *et al.*, 2005). Cardiorespiratory arrest is rare probably occurring in 1 per 200 individuals; four cases out of around 1000 patients with PS have been reported but they all had complete recovery (Panayiotopoulos, 2002; Verrotti *et al.*, 2005; Ferrie *et al.*, 2006). Raised temperature has been documented in a few cases (2%) after seizure onset. Cephalic auras of discomfort and odd sensations or headache commonly are described with other autonomic symptoms at seizure onset.

Syncopal-like manifestations occur in at least one-fifth of seizures (Panayiotopoulos, 2002; Ferrie *et al.*, 2006, 2007; Covanis, 2006; Caraballo *et al.*, 2007). The child becomes ‘completely unresponsive and flaccid like a rag doll’, which may precede, be concurrent with other seizure symptoms or be the sole manifestation of a seizure (Oguni *et al.*, 1999; Panayiotopoulos, 2002). They may occur while the patient is standing, sitting, lying down or asleep and last from 1–2 min to half an hour.

Ictal behavioural changes

Restlessness, agitation, terror or quietness, may occur at the onset of seizures, often in combination with other autonomic manifestations.

Ictal non-autonomic symptoms

Pure autonomic seizures and pure autonomic status epilepticus appear to occur in 10% of patients. They commence and terminate solely with autonomic symptoms. However, in the majority of seizures, autonomic manifestations are followed by conventional seizure symptoms. Nearly always, the child gradually or suddenly becomes confused or unresponsive. Other non-autonomic manifestations include in order of prevalence unilateral deviation of the eyes or eyes opening (60–83%), speech arrest (8–13%), hemifacial

convulsions (6–13%), visual hallucinations (6–10%), OPLS (3%), unilateral drooping of the mouth (3%) and rarely (1%) eyelid or limb jerks, nystagmus or automatisms. The seizures may end with hemiconvulsions often with Jacksonian marching (19–30%), or generalized convulsions (21–36%).

Autonomic manifestations may not be apparent at seizure onset even in witnessed diurnal seizures (<10%). They may be absent, mild or missed in clinical observation. In these cases, eye deviation is the more common symptom. Visual symptoms are rare (1%) and not present in recurrent seizures (Ferrie *et al.*, 1997; Yalcin *et al.*, 1997; Panayiotopoulos, 2002; Caraballo *et al.*, 2007).

Duration of seizures and precipitating factors

The seizures are usually lengthy of over 6 min and almost half of them last for >30 min to many hours, thus constituting autonomic status epilepticus (Panayiotopoulos, 2002; Ferrie *et al.*, 2007). Lengthy seizures are equally common in sleep and wakefulness. Even after the most severe seizures and status, the patient is normal after a few hours’ sleep. There is no record of residual neurological abnormalities. Hemiconvulsive or convulsive status epilepticus is rare (4%).

Two-thirds of seizures start in sleep. Many seizures have been witnessed while travelling in a car, boat or aeroplane. The reason for this may be because in these circumstances children easily fall asleep, seizures are more likely to be witnessed and because travelling also precipitates motion sickness, to which children are particularly susceptible.

Intra-individual seizure variability

The same child may have brief and lengthy seizures, diurnal and nocturnal, with marked, inconspicuous, or even without any autonomic changes (Ferrie *et al.*, 1997; Oguni *et al.*, 1999; Kivity *et al.*, 2000; Panayiotopoulos, 2002; Lada *et al.*, 2003; Ohtsu *et al.*, 2003, 2008; Covanis *et al.*, 2003; Caraballo *et al.*, 2007). Even cardinal symptoms (such as vomiting or eye deviation) may be present in one but absent in another seizure. Seizures without autonomic manifestations are rare (7%) and occur in patients who also have additional autonomic seizures (Panayiotopoulos, 2002). Ictal video EEG recordings have documented that autonomic symptoms and signs may vary between seizures of the same child (Koutroumanidis *et al.*, 2005). There is no correlation between ictal semiology and topography of interictal spikes.

Aetiology

PS, like rolandic epilepsy, is probably genetically determined though conventional genetic influences may be less important than other mechanisms (Taylor *et al.*, 2008). Usually, there is no family history of similar seizures, although siblings with PS or PS and rolandic epilepsy have been reported (Ferrie *et al.*, 1997; Lada *et al.*, 2003; Covanis *et al.*,

2003; Caraballo *et al.*, 2007; Livingston *et al.*, 2008; Taylor *et al.*, 2008). There is a high prevalence of febrile seizures (about 17%) (Panayiotopoulos, 2002).

SCN1A mutations have been recently reported in a child (Grosso *et al.*, 2007) and two siblings (Livingston *et al.*, 2008) with relatively early onset of seizures, prolonged time over which many seizures have occurred and strong association with febrile precipitants even after the age of 5 years. This is an area that needs further attention but may indicate that SCN1A mutations contribute to a more severe phenotype of PS.

Pathophysiology

Autonomic symptoms of any type are often encountered in seizures, whether focal or generalized, in adults or children (Freeman, 2006; Ferrie *et al.*, 2007; Goodman *et al.*, 2008). They are generated by activation or inhibition of parts of the central autonomic network that involves the insular cortex, medial prefrontal cortex, amygdala, hypothalamus and ventrolateral medulla (Goodman *et al.*, 2008). The resultant autonomic disturbances depend on the brain areas involved in seizure onset or propagation, and appear as single or multiple symptoms some of which may be of localizing value (Elger, 2000).

In PS, the neuroanatomical and neurophysiological underpinnings of autonomic manifestations are unknown. Any hypothesis of the pathophysiology of PS should explain significant pieces of evidence that converge from clinical, EEG and magnetoencephalographic studies.

First, autonomic seizures and autonomic status epilepticus with the symptomatology and sequence as in PS appear to be specific for childhood (Panayiotopoulos, 2004; Ferrie *et al.*, 2007). For example, in adults, ictal vomiting occurs rarely, and as a rule when consciousness is impaired following other focal mainly temporal lobe symptoms, and is attributed to non-dominant mesial temporal lobe involvement (Kramer *et al.*, 1988; Schauble *et al.*, 2002; Koutroumanidis, 2003). In contrast, ictal vomiting in children is common, usually occurs when consciousness is intact without preceding focal cortical symptoms, and probably has no localizing or lateralizing value (see ictal EEG). A possible explanation for this discrepancy may relate to the fact that children are constitutionally more vulnerable to emetic disturbances as exemplified by the ‘cyclic vomiting syndrome’, a non-seizure disorder of unknown aetiology that is also specific to childhood (Li *et al.*, 1999) and associated with autonomic dysfunction (Chelimsky and Chelimsky, 2007). Thus, the preferential involvement of emetic and other autonomic manifestations in PS may be attributed to a maturation-related susceptibility of the central autonomic network (Panayiotopoulos, 2002, 2004).

Second, the epileptogenic zone in PS is wide and bilateral with multifocal pockets in cortical areas surrounding major fissures such as central, sylvian and mainly calcarine (Kanazawa *et al.*, 2005; Yoshinaga *et al.*, 2006; Saitoh *et al.*, 2007, 2008).

Third, ictal autonomic symptomatology appears to pertain to any epileptogenic cortical onset zone, be this occipital, frontotemporal or frontal (Beaumanoir, 1993b; Oguni *et al.*, 1999; Demirbilek and Dervent, 2004; Koutroumanidis *et al.*, 2005; Parisi *et al.*, 2005) (Fig. 2) and usually precede other focal cortical semiology. It is likely, that central autonomic networks have a lower threshold to epileptogenic activation than those producing focal cortical semiology (occipital, frontal, central, parietal and less often temporal). Irrespective of the localization of their onset, ictal discharges may activate the lower threshold autonomic centres (and therefore produce autonomic manifestations) commonly before other cortical regions of relatively higher threshold that generate focal cortical symptoms (sensory, motor, visual or other). Seizures remain purely autonomic if ictal neuronal activation of non-autonomic cortical areas fails to reach symptomatogenic threshold; otherwise they consist of autonomic and localization-related cortical symptoms and signs that may only rarely occur from onset. This hypothesis may explain why similar autonomic manifestations may appear from anterior or posterior, right or left brain onsets. As seizures primarily involve a particular system (the autonomic), PS may be considered as an electroclinical example of ‘system epilepsy’ (Koutroumanidis, 2007).

To explain the paradoxical discrepancy between the prolonged and ample-looking ictal discharges that have invariably featured in all published ictal recordings (Beaumanoir, 1993b; Oguni *et al.*, 1999; Vigeveno *et al.*, 2000; Demirbilek and Dervent, 2004; Koutroumanidis *et al.*, 2005; Parisi *et al.*, 2005) and the also consistent lack of conspicuous cortical (motor or sensory) manifestations for several minutes into the seizure, one may also hypothesize a suboptimal ‘strength’ of the ictal electrical activity. Despite their scalp EEG phenomenology discharges presumably fail to transform into dynamic cortico–cortical propagation and generate conventional cortical symptoms according to their distribution over the cerebrum (being though still capable of activating a more hyperexcitable autonomic network). After all, the magnitude (amplitude) of ictal discharges as they appear in scalp recordings is hardly a reliable indicator of the ‘vigour’ of the clinical manifestations: for example, dramatically appearing hypermotor frontal lobe seizures may have no scalp EEG correlates and high voltage generalized or diffuse spike wave activity (such as in atypical absence seizures of epileptic encephalopathies) may have only few and mild clinical correlates (Koutroumanidis, 2007).

Syncopal-like attacks are difficult to explain. They may be a distinct seizure-type symptom similar to atonic seizures, but on some occasions they may be due to cardiac asystole (ictal syncope) generated by the seizure discharge.

Electroencephalography

Inter-ictal EEG findings show great variability (Fig. 1) (Panayiotopoulos, 1988, 2002; Oguni *et al.*, 1999; Lada *et al.*, 2003; Ohtsu *et al.*, 2003, 2008; Covanis *et al.*, 2003;

Sanders *et al.*, 2004; Ferrie *et al.*, 2006; Caraballo *et al.*, 2007). In about 90% of cases, the EEG reveals mainly multifocal, high amplitude, sharp slow wave complexes that may appear in any area, often shifting from one region to another in the same or the contralateral hemisphere in sequential EEGs of the same child. Occipital spikes predominate but they do not occur in a third of patients. Occipital paroxysms in their classical form with fixation off sensitivity (FOS) are even rarer. Clone-like multifocal spikes are intriguing features when they occur (19%). In routine EEG, these spikes appear to occur simultaneously at various locations in one or both hemispheres but usually they are driven (secondarily activated) by a primary spike generator which is predominantly posterior, leads the other spikes by a few ms and may also be the smallest of all or inconspicuous (Panayiotopoulos 2002; Leal *et al.*, 2007, 2008). Brief generalized discharges of slow waves, intermixed with small spikes, may occur either alone (4%) or more often with focal spikes (15%). A single routine EEG may be normal in 10% of patients, and a few children have consistently normal wake EEGs before a diagnostic sleep recording. Sleep typically accentuates the spike abnormalities, and photosensitivity is practically absent.

As in benign rolandic epilepsy, the frequency, location and persistence of spikes do not determine the clinical manifestations, the duration, the severity and frequency of seizures or their prognosis. For instance, spikes may persist for many years after clinical remission or appear only once despite multiple EEGs.

The multifocal potential for epileptogenesis in PS has also been documented by EEG dipole analysis (Yoshinaga *et al.*, 2006) and magnetoencephalography, which have implicated areas along the parieto-occipital, calcarine and central sulci or in the frontal lobes (Kanazawa *et al.*, 2005; Saitoh *et al.*, 2007; Saito *et al.*, 2008).

In the few reported ictal EEGs, the discharges consist mainly of unilateral rhythmic slow activity, usually intermixed with fast rhythms and small spikes. They start in wider more often in posterior than anterior regions, quickly become diffuse and last for many minutes (Beaumanoir, 1993b; Oguni *et al.*, 1999; Demirbilek and Dervent, 2004; Koutroumanidis *et al.*, 2005; Parisi *et al.*, 2005) (Fig. 2). The first ictal clinical symptoms become apparent usually long after the onset of the electrical discharge and present as tachycardia, breathing irregularities, coughing or emesis, which would be unlikely to consider as seizure events without an EEG.

Differential diagnosis

PS is easy to diagnose because of the characteristic clustering of clinical seizure semiology, which is often supported by inter-ictal EEG findings. The main problem is to recognize emetic and other autonomic manifestations as seizure events, and not to dismiss them or erroneously consider them as unrelated to the ictus and as a feature of encephalitis,

migraine, syncope or gastroenteritis, which is the reason of the belated recognition of this common syndrome (Panayiotopoulos, 1988, 2002; Kivity and Lerman, 1992; Covanis, 2006).

A most difficult situation that demands experienced evaluation is when a child is seen at the acute stage of a seizure when symptoms may dramatically accumulate in succession and the diagnosis of true encephalitis is possible. A history of a previous similar seizure or full recovery after a few hours of sleep is reassuring and may help to avoid unnecessary investigations and promote withdrawal of any medication that may have been initiated (Kivity and Lerman, 1992; Sanders *et al.*, 2004).

Approximately 10–20% of autonomic seizures and autonomic status epilepticus in children are due to heterogeneous cerebral pathology (Panayiotopoulos, 1988, 2002). These symptomatic cases are betrayed by abnormal neurological or mental state, abnormal brain imaging and background EEG abnormalities.

PS is significantly different from the rolandic epilepsy and the ICOE-G despite some overlapping clinical and/or EEG features. These are detailed in the relevant section of this article.

Prognosis

PS is remarkably benign in terms of its evolution (Panayiotopoulos, 1988, 2002; Ferrie *et al.*, 1997; Oguni *et al.*, 1999; Kivity *et al.*, 2000; Lada *et al.*, 2003; Ohtsu *et al.*, 2003; Covanis *et al.*, 2003; Caraballo *et al.*, 2007) but autonomic seizures are of concern in the rare context of cardiorespiratory arrest though all four reported cases recovered completely (Panayiotopoulos, 2002; Verrotti *et al.*, 2005; Ferrie *et al.*, 2006, 2007). The majority of patients have a single or less than five seizures until remission. Only one quarter have multiple and sometimes very frequent and prolonged seizures that may be resistant to treatment. Remission often occurs within 1–2 years of onset but probably 10% may have more protracted active seizure periods. A fifth of patients develop rolandic and less often occipital or other seizures but these are also age-related and remit (Panayiotopoulos, 2002). Atypical evolution of PS similar to those described in rolandic epilepsy is rare, probably <3% (Caraballo *et al.*, 2001, 2007; Ferrie *et al.*, 2002; Kikumoto *et al.*, 2006).

The risk of epilepsy in adult life appears to be no higher than in the general population (Panayiotopoulos, 2002; Ferrie *et al.*, 2006; Caraballo *et al.*, 2007).

Subtle neuropsychological deficits in some children during the active phase (Germano *et al.*, 2005) may be syndrome-related but may also reflect effects of AED (most of the children were on AEDs including phenobarbital and vigabatrin) and/or other contributing factors. Prognosis of cognitive function is good even for patients with atypical evolutions (Caraballo *et al.*, 2007).

Idiopathic childhood occipital epilepsy of Gastaut

The ICOE-G is a relatively rare form of pure occipital epilepsy accounting for about 2–7% of benign childhood focal seizures (Gastaut, 1981, 1982*a, b*; Panayiotopoulos, 1981, 1999*a, b*; Beaumanoir, 1983; Gastaut and Zifkin, 1987; Gastaut *et al.*, 1992; Ferrie *et al.*, 1997; Kivity *et al.*, 2000; Covanis *et al.*, 2005; Gobbi *et al.*, 2008; Caraballo *et al.*, 2008*a*). Age at onset ranges from 3 to 15 years, but most start between 8 and 11 years. Both sexes are equally affected.

Clinical manifestations

Seizures are occipital and primarily manifest with elementary visual hallucinations, blindness or both (Gastaut, 1982*a*; Gastaut and Zifkin, 1987; Gastaut *et al.*, 1992; Panayiotopoulos, 1999*a, b*; Gobbi *et al.*, 2008; Caraballo *et al.*, 2008*a*). They are usually frequent, brief and diurnal.

Visual ictal symptoms

Elementary visual hallucinations are the most common and characteristic ictal symptom of ICOE-G. They are frequently the first and often the only seizure symptom. They develop rapidly within seconds and consist mainly of small multi-coloured circular patterns that often appear in the periphery of a visual field, becoming larger and multiplying during the course of the seizure, frequently moving towards the other side.

Ictal blindness is probably the second most common symptom after visual hallucinations. It is sudden, usually total and it is frequently the first and often the only seizure symptom in patients who may also have other visual seizures without blindness. Impairment of visual awareness is consistently reported by some patients before the appearance of visual hallucinations.

Complex visual hallucinations such as faces and figures, and visual illusions such as micropsia, palinopsia and metamorphopsia occur in <10% of patients and mainly after the appearance of elementary visual hallucinations (Gastaut and Zifkin, 1987).

Non-visual ictal occipital lobe symptoms and signs

Non-visual occipital symptoms usually appear after the elementary visual hallucinations and these in order of prevalence are deviation of the eyes, eyelid fluttering or repetitive eye closures, pain and sensory hallucinations of ocular movements (Gastaut, 1982*a*; Gastaut and Zifkin, 1987; Panayiotopoulos, 1999*a, b*; Gobbi *et al.*, 2008; Caraballo *et al.*, 2008*a*).

Deviation of the eyes, often associated with ipsilateral turning of the head, is the most common (in about 70% of cases) non-visual ictal symptom. It usually starts after the commencement of visual hallucinations and may be mild, but more often it is forceful tonic and may progress to hemiconvulsions and GTCS. Some children may have

seizures of eye deviation from the start without visual hallucinations and it is likely that these cases have a better prognosis (Beaumanoir, 1983; Ferrie *et al.*, 1997). Other ocular manifestations may include unidirectional ocular clonic seizures (oculoclonic seizures) that are rare, and eyelid fluttering or repetitive eye closures that occur in about 10% of patients, usually at a later stage when consciousness is impaired. They signal an impending secondary GTCS.

Ictal headache, or mainly orbital pain, is a common ictal symptom, and in a small number of patients it may start before the first visual or other ictal occipital symptoms.

Consciousness

Consciousness is intact during the visual symptoms (simple focal seizures), but may be disturbed or lost in the course of the seizure, usually before or at the time of eye deviation or convulsions. Syncopal-like attacks are rare (Panayiotopoulos, 2002).

Extra-occipital seizure progression

Elementary visual hallucinations or other ictal symptoms may progress to complex focal seizures (14%), hemiconvulsions (43%) or GTCS (13%) (Gastaut and Zifkin, 1987). Complex focal seizures of temporal lobe symptomatology are extremely rare and may indicate a symptomatic cause (Panayiotopoulos, 1999*b*). Ictal vomiting may occur with progression to the non-dominant temporal lobe (Guerrini *et al.*, 1995).

Post-ictal headache

Post-ictal headache, mainly diffuse, but also severe, unilateral, pulsating and indistinguishable from migraine headache, occurs in half the patients, in 10% of whom it may be associated with nausea and vomiting (Gastaut and Zifkin, 1987; Panayiotopoulos, 1999*a, b*; Caraballo *et al.*, 2008*a*). This occurs immediately, or 5–10 min after the end of the visual hallucinations. The duration and severity of the headache appears to be proportional to the duration and severity of the preceding seizure although it may also occur after brief simple visual seizures.

Seizure stereotype

For any one patient, in every seizure, the elementary visual hallucinations have a fingerprint with a stereotypic appearance regarding morphology, colours, location, movement and other characteristics. Most of patients also know at what stage of their ictal manifestations a secondary GTCS may occur.

Duration and circadian distribution

Visual seizures are usually brief, lasting from a few seconds to 1–3 min if they occur alone without other occipital or extra-occipital spreading (Gastaut, 1982*a*; Gastaut and Zifkin, 1987; Panayiotopoulos, 1999*a, b*; Covanis *et al.*, 2005; Gobbi *et al.*, 2008; Caraballo *et al.*, 2008*a*). However, a few patients with brief visual seizures may later develop lengthy visual seizures lasting for 10–20 min.

Visual seizures are predominantly diurnal and occur at any time of the day but some patients may also have infrequent seizures in sleep or on awakening.

Frequency of seizures

If untreated, the majority of patients experience frequent brief visual seizures ranging from several every day to one per week or month. However, propagation to other seizure manifestations, such as focal or generalized convulsions, is much less frequent occurring once per month, year or even rarer.

Precipitating factors and idiopathic photosensitive occipital epilepsy

This is a matter of inclusion criteria. Gastaut considered photosensitivity as part of ICOE-G (Gastaut, 1982a; Gastaut and Zifkin, 1987), while the ILAE Task Force recognizes ‘idiopathic photosensitive occipital lobe epilepsy’ as a syndrome of reflex epilepsy with age-related onset (Engel, 2001, 2006). Reflex occipital seizures induced by television, video games and intermittent photic stimulation (IPS) manifest with similar semiology as the spontaneous visual seizures (Aso *et al.*, 1987; Michelucci and Tassinari, 1993; Guerrini *et al.*, 1995, 1998; Yalcin *et al.*, 2000; Panayiotopoulos, 2007). Deviation of the eyes, epigastric discomfort and vomiting, headache and generalized convulsions may follow. Prognosis is uncertain. Some children may have only 1 or 2 seizures, but others may not remit. Interictal EEG shows spontaneous and photically induced occipital spikes (Fig. 1). CTS may coexist. Ictal EEG documented the occipital origin and the spreading of the discharges to the temporal regions (Guerrini *et al.*, 1995, 1998). There remain no other significant precipitating factors in ICOE-G if photosensitive patients are excluded. Despite FOS in EEG, only a few patients report seizure precipitation by going from bright light to darkness or by darkness itself (Beaumanoir *et al.*, 1989).

Aetiology

There is an increased family history of epilepsies (21–37%) or migraine (9–16%) (Gastaut and Zifkin, 1987; Caraballo *et al.*, 2008a) but familial ICOE-G appears to be rare (Nagendran *et al.*, 1990; Grosso *et al.*, 2008; Taylor *et al.*, 2008).

Pathophysiology

The seizures are purely of occipital lobe origin. The epileptogenic zone involves networks within the occipital lobes and this localization is congruent with the symptomatogenic zone. Elementary visual hallucinations originate from the primary visual cortex, complex visual hallucinations from the junction of the occipital with the parietal and temporal lobes, formed visual illusions from the lateral occipital–posterior temporal junction and tonic deviation of the eyes from the medial occipital cortex, above or below the calcarine sulcus. Ictal blindness may reflect bi-occipital seizure spreading but this may not explain its sudden onset, without any other preceding manifestations. From the

EEG standpoint, the occipital paroxysms are usually bilateral and synchronous because they are activated in both occipital regions by the elimination of fixation and central vision (FOS) (Panayiotopoulos, 1981) and not by thalamocortical activation proposed by Gastaut and Zifkin (1987).

The mechanisms for post-ictal headache are unknown. It is likely that the occipital seizure discharge triggers a genuine migraine headache through trigeminovascular or brain-stem mechanisms (Panayiotopoulos, 1999b, c).

Diagnostic procedures

By definition, all tests other than the EEG are normal. However, high-resolution MRI is mandatory, because symptomatic occipital epilepsy present with the same clinical EEG manifestations.

Electroencephalography

The inter-ictal EEG shows occipital paroxysms (Gastaut, 1982a; Gastaut and Zifkin, 1987), often demonstrating FOS (Panayiotopoulos, 1981) (Fig. 1). Because terminology is often unclear and FOS is not always tested, the prevalence of classical occipital paroxysms with FOS is uncertain and ranges from 100% (Gastaut and Zifkin, 1987), 88% (Caraballo *et al.*, 2008a) to 19% (Panayiotopoulos, 1999a). Some patients may have only random occipital spikes, whereas others may have occipital spikes only in sleep EEG and some may have a consistently normal EEG (Panayiotopoulos, 1999b). Centro-temporal, frontal and GSES occur together with occipital spikes in around 20% of patients (Herranz Tanarro *et al.*, 1984; Gastaut and Zifkin, 1987). IPS consistently elicits occipital spikes and/or generalized discharges in photosensitive patients.

As it happens with the rolandic spikes, occipital spikes are not pathognomonic of any particular syndrome, because they also occur in a variety of organic brain diseases with or without seizures, in children with congenital or early onset visual and ocular deficits, and even in 0.5–1.2% of normal pre-school age children (Gibbs and Gibbs, 1952, 1967; Kellaway, 1980). They are common in young children with a peak age at first discovery of 4–5 years, and ‘tend to disappear in adult life, and the subsidence of the EEG abnormality is usually accompanied by a cessation of seizures’ (Gibbs and Gibbs, 1952, 1967).

There are many reported ictal EEGs (Gastaut, 1982a; Aso *et al.*, 1987; Gastaut and Zifkin, 1987; De Romanis *et al.*, 1988, 1991; Beaumanoir, 1993a, b; Thomas *et al.*, 2003). Seizure onset is preceded by regression of occipital paroxysms, and is characterized by the sudden appearance of an occipital discharge that consists of fast rhythms, fast spikes or both and is of much lower amplitude than the occipital paroxysms. Elementary visual hallucinations relate to the initially fast spike activity and complex visual hallucinations may occur when the ictal discharge is slower. In oculoclonic seizures, spikes and spike-wave complexes are slower, and a localized ictal fast spike rhythm may occur before deviation of the eyes. Ictal EEG

Table 1 Key differences between occipital seizures and migraine

Elements of visual hallucinations	Occipital seizures	Migraine
Speed of development from onset to full image	Fast in seconds	Slow in minutes
Speed and direction of movement	Fast in seconds and usually towards the centre of the visual field and contralateral to the side of onset	Slow in minutes and usually towards the periphery of the visual field and ipsilateral to the side of onset
Quality	Usually with bright colours and circular shapes	Usually achromatic or black and white linear zigzag patterns
Duration	Usually 1–3 min	Usually over 15 min
Progression to transient neurological symptoms	Eye deviation, eyelid closures and sometimes convulsions	Scotoma, hemianopia, hemi-anaesthesia or hemi-paresis and for basilar migraine, vertigo, ataxia, bilateral weakness and dysaesthesiae

during blindness is characterized by pseudo-periodic slow waves and spikes, which differ from those seen in ictal visual hallucinations. There are usually no post-ictal abnormalities.

Differential diagnosis

The differential diagnosis of ICOE-G is mainly from symptomatic occipital epilepsy, migraine with aura, acephalgic and basilar migraine where misdiagnosis is very high (Panayiotopoulos, 1999a, b).

Patients with symptomatic occipital epilepsy may often have identical symptoms as ICOE-G with normal neuro-ophthalmological examination and routine brain imaging. Thus, high-resolution MRI is required to detect subtle lesions (Kuzniecky *et al.*, 1997). Occipital seizures of mitochondrial disorders, Lafora disease and coeliac disease should be considered (Panayiotopoulos, 1999a; Taylor *et al.*, 2003).

The differential diagnosis of ICOE-G from migraine is usually easy if all clinical elements are properly assessed and synthesized (Table 1). Contrary to visual seizures, visual aura of migraine develops slowly within minutes, lasts long for 10–20 min and consists of mainly achromatic and linear patterns (Panayiotopoulos, 1994; Russell and Olesen, 1996; Schott, 2007). Illustration of the visual symptoms of the attacks by the patient is a powerful tool in differential diagnosis and objective analysis. Orbital pain in the ictal phase of visual hallucinations is typical of occipital seizures and does not occur in migraine. However, post-attack headache is common and similar for both occipital epilepsy and migraine. Basilar migraine attacks also develop slowly within minutes, last for 30–60 min and consist of mainly bilateral impairment of vision associated with, or followed by, neurological symptoms such as vertigo, tinnitus, ataxia, bilateral weakness and dysaesthesiae, which do not occur in occipital lobe epilepsy (Panayiotopoulos, 1999c).

Mistaking visual seizures as migraine attacks may be common in publications referring to controversial diagnostic terms such as ‘migralepsy’ and ‘basilar migraine with occipital paroxysms’. A critical review of such reported cases indicates that these are likely to be genuine occipital seizures imitating migraine (Panayiotopoulos, 1999c).

Despite some overlapping features, ICOE-G is distinctive from PS (Table 2) and the differences have been statistically

validated (Panayiotopoulos, 1999a). As a rule of thumb seizure onset is primarily with visual symptoms in ICOE-G and with autonomic manifestations in PS. On the rare occasion of ICOE-G with autonomic manifestations, these always occur after occipital lobe symptomatology.

Prognosis

The prognosis of ICOE-G is unclear, although available data indicate that remission occurs in 50–60% of patients within 2–4 years of onset (Gastaut and Zifkin, 1987; Panayiotopoulos, 1999b; Caraballo *et al.*, 2008a). Seizures show a dramatically good response to carbamazepine in >90% of patients. However, 40–50% of patients may continue having visual seizures and infrequent secondarily GTCS. Rarely, atypical evolutions to epilepsy with CSWS and cognitive deterioration have been reported (Tenenbaum *et al.*, 1997). Also, rarely children with ICOE-G may manifest with typical absence seizures, which usually appear after the onset of occipital seizures (Caraballo *et al.*, 2005).

The performance scores for attention, memory and intellectual functioning were lower in patients with ICOE-G than control subjects though basic neurophysiological functions did not differ significantly (Gulgonen *et al.*, 2000).

Other phenotypes of BCSSS

There are reports of children suffering from benign childhood focal seizures with clinical EEG manifestations that cannot be classified as rolandic epilepsy, PS or ICOE-G. They may represent rare, atypical or overlapping presentations of BCSSS.

Benign childhood seizures with affective symptoms

Benign childhood epilepsy with affective symptoms, reported in less than 40 patients, is a clinical phenotype of BCSSS with features common in both PS (behavioural and autonomic symptoms) and rolandic epilepsy (speech arrest and hypersalivation) (Dalla Bernardina *et al.*, 1992, 2007). Onset is between 2 and 9 years of age and both sexes are equally affected.

Table 2 Main features of rolandic epilepsy, Panayiotopoulos syndromes and idiopathic childhood epilepsy of Gastaut

	Rolandic epilepsy	Panayiotopoulos syndrome	Idiopathic childhood occipital epilepsy of Gastaut
Prevalence amongst children aged 1–15 years	15	6	0.5–1
with non-febrile seizures (%)			
Peak age at onset (years)	7–10	3–6	8–11
Male to female ratio	1.5	1	1
Seizure characteristics			
Typical onset with	Hemifacial sensory-motor or oropharyngolaryngeal symptoms	Autonomic symptoms mainly with emesis	Visual symptoms mainly with elementary visual hallucinations
Hemifacial sensory-motor symptoms	Common and often from onset	Rare and not from onset	Rare and not from onset
Oropharyngolaryngeal symptoms	Common and often from onset	Rare and not from onset	Have not been reported
Speech arrest	Common and often at onset	Rare and not from onset	Has not been reported
Hypersalivation	Common and often at onset	Rare and not from onset	Has not been reported
Ictus emeticus	Rare and not from onset	Common and often at onset	Rare and not from onset
Autonomic disturbances other than vomiting and hypersalivation	Exceptional and not from onset	Common and often at onset	Exceptional and not from onset
Visual symptoms	Have not been reported	7% but exceptional at onset	Common and often at onset
Deviation of the eyes	Frequent during sensory-motor symptoms	Common but rarely at onset	Common but rarely at onset
Ictal behavioural changes	Exceptional and not from onset	Common and often at onset	Have not been reported
Duration for 1–3 min	As a rule	Rare	As a rule
Duration of more than 5 min	Rare	Common	Rare
Partial status epilepticus (>30 min)	Exceptional	40%	Exceptional
Total number of seizures 1–15	As a rule	As a rule	Rare
Single seizures only (%)	10–20	30	Exceptional
Frequent seizures (%)	10	10	90
Nocturnal (sleep only) (%)	70	64	Exceptional
Febrile convulsions	18	17	10
Prognosis	Excellent	Excellent	Uncertain
Remission within 1–2 years from first seizure	Common	Common	Exceptional or rare
Seizures after the age of 13 years	Rare	Exceptional	Common
Interictal EEG			
Centrottemporal spikes alone	As a rule and characteristic	Rare	Have not been reported
Occipital spikes	Have not been reported	65%	Probably 90%
Spikes in other locations	Probably uncommon	Frequent	Exceptional
Brief generalised discharged of 3–5 Hz slow waves with small spikes (%)	5	10	Exceptional
Somatosensory evoked spikes	Common	Rare	Have not been reported
Fixation-off sensitivity	Has not been reported	Rare	May be less common than reported
Photosensitivity	Has not been reported	Exceptional	Probably 20–30%
Normal EEG or focal slow after first seizure (%)	~10	~10	~10
Ictal EEG	Slow activity with spikes	Slow activity with spikes	Fast spikes and fast rhythms
Ictal onset	Rolandic regions	Anterior or posterior regions	Occipital regions

Seizures manifest with terror and screaming, autonomic disturbances (pallor, sweating, abdominal pain and hypersalivation), chewing and other automatisms, speech arrest and mild impairment of consciousness. These are usually brief, lasting 1–2 min, and frequent, occurring several times a day in wakefulness or sleep. A fifth of patients have febrile seizures and some may also have infrequent rolandic seizures. Generalized seizures have not been reported.

The inter-ictal EEG shows high-amplitude frontotemporal and parietotemporal spikes that are exaggerated by sleep. Ictal EEG discharges are mainly localized in the frontotemporal, centro-temporal or parietal regions and are stereotypical for each patient.

The response to treatment is excellent and remission occurs within 1–2 years from onset. Behavioural problems may be prominent during the active stage of the disease, but subside later with seizure remittance.

Benign childhood epilepsy with parietal spikes and frequent extreme somatosensory-evoked spikes

Benign childhood epilepsy with parietal spikes and frequent GSES (De Marco and Tassinari, 1981; Tassinari and De Marco, 1992; Fonseca and Tedrus, 2000) has been proposed as another phenotype of BCSSS. The defining features are EEG spikes in the parietal regions, which are often elicited by tactile stimulation (Fig. 1). However, GSES are not specific for any syndrome because they also occur in 10–20% of children with rolandic seizures (Fonseca and Tedrus, 2000), in a few patients with PS (Panayiotopoulos, 1999a, 2002) and in children with no seizures (Negrin and De Marco, 1977).

Versive seizures of the head and body, often without impairment of consciousness, are mainly diurnal and infrequent. Frequent seizures and focal status epilepticus are exceptional.

Remission usually occurs within 1 year from seizure onset, but EEG abnormalities may persist for longer.

Benign childhood focal seizures associated with frontal or midline spikes

Benign childhood focal seizures associated with frontal (Beaumanoir and Nahory, 1983; Martin-Santidrian *et al.*, 1998; Panayiotopoulos, 1999a) or midline spikes (Bagdord and Lee, 1993; Panayiotopoulos, 1999a) have been described and long follow-up reports have confirmed a benign course, although no systematic studies have been published. However, EEG spike foci specificity is questionable, as spike foci of various locations (including frontal and midline) are also seen in rolandic epilepsy and more frequently in PS, and midline spikes are more common in children than in adults (Kutluay *et al.*, 2001; Sanders *et al.*, 2002).

Recently, 'benign infantile focal epilepsy with midline spikes during sleep' has been described as a new syndrome of BCSSS (Bureau *et al.*, 2002; Capovilla *et al.*, 2006). Age

at onset is in the first 3 years of life and both sexes are equally affected. Seizures consist mainly of staring, motion arrest, cyanosis, loss of consciousness and stiffening of the arms. Clonic convulsions and automatisms are rare. Seizures are brief from 1 to 5 min, mainly diurnal and are generally infrequent from one to three per year. There is a strong family history of undefined types of epileptic seizures with benign epilepsies prevailing.

Inter-ictal EEG abnormalities are seen only in non-REM sleep and consist of small, mostly singular, midline spikes. The prognosis is excellent, with remission of seizures, normal development and normalization of the EEG before the age of 4 years.

Differential diagnosis between seizures and syndromes of BCSSS

The differential diagnosis between the main phenotypes of BCSSS is easy in their typical presentations (Table 2). Problems may arise in children with clinical symptoms that fall into two (or more) phenotypes or from overemphasizing on EEG localization. As in any other medical condition, a single symptom is of limited syndromic significance. The differential diagnosis requires that symptoms are meaningfully synthesized in regard to quality and quantity, chronological sequence, consistency, relation to other seizure manifestations, the circumstances of their appearance and the overall clustering of clinical EEG manifestations.

Rolandic epilepsy versus PS

Their differential diagnosis is usually easy (Table 2). However, there are some cases with overlapping features:

- One-tenth of children with PS often have typical and lengthy autonomic seizures with concurrent rolandic features such as speech arrest, hemifacial convulsions, hypersalivation and OPS but these appear after the onset of autonomic symptoms and emesis (Ferrie *et al.*, 1997; Oguni *et al.*, 1999; Kivity *et al.*, 2000; Panayiotopoulos, 2002; Lada *et al.*, 2003; Caraballo *et al.*, 2007). Conversely, these ictal symptoms occur at onset and usually without autonomic symptoms in rolandic epilepsy.
- One-tenth of children with PS develop pure rolandic seizures, either in parallel with autonomic seizures, or at a later age prior to final remission (Oguni *et al.*, 1999; Panayiotopoulos, 2002; Lada *et al.*, 2003; Caraballo *et al.*, 2007).
- The topography of interictal spikes may overlap. Covanis *et al.* (2003) studied 24 otherwise normal children with focal non-febrile seizures who had emetic manifestations in at least one seizure and CTS in at least one EEG; 21 (83%) had ictal semiology typical of PS but five also had concurrent rolandic symptoms and four later developed pure rolandic seizures. The other four children (17%) had typical

rolandic seizures with concurrent ictus emeticus. Ohtsu *et al.* (2007) found that in early-onset rolandic epilepsy vomiting usually happened in the middle of the ictus, seizures, neurocognitive and behavioural abnormalities were more frequent, while focal status epilepticus and prolonged seizures were less common than in PS.

- (d) Of siblings one may have rolandic seizures and another PS, and there is a high prevalence of febrile seizures in both (Ferrie *et al.*, 1997; Oguni *et al.*, 1999; Panayiotopoulos, 2002; Caraballo *et al.*, 2007).

Idiopathic childhood occipital epilepsy of Gastaut versus PS

The differentiation here is straightforward (Table 2) and statistically validated (Panayiotopoulos, 1999a). The seizures of ICOE-G are purely occipital and as such start and often end only with occipital lobe symptomatology. Further, seizures are mainly brief, frequent and diurnal. Rarely, seizures may be longer and also occur in sleep but these are also fundamentally different to the rolandic epilepsy or the autonomic seizures and autonomic status epilepticus of PS.

Exceptionally, ictal vomiting may occur in ICOE-G but this follows the appearance of visual symptomatology as it happens with reflex photosensitive occipital seizures (Guerrini *et al.*, 1995, 1998) and the same patient usually has frequent brief occipital seizures. Conversely, visual symptoms in PS, when present, are not the sole manifestation of a seizure or stereotypical; only exceptionally (1%) are they reported at seizure onset (Ferrie *et al.*, 1997; Yalcin *et al.*, 1997; Panayiotopoulos, 2002; Caraballo *et al.*, 2007). From the EEG standpoint, occipital spikes which characterize ICOE-G are also common in PS but these often occur with extra-occipital spikes and with shifting locations in sequential EEG (Fig. 1). Further, ictal EEG is markedly different between these syndromes (Fig. 2).

Reported difficulties in differentiating ICOE-G from PS (Genizi *et al.*, 2007; Taylor *et al.*, 2008) may arise when emphasis is unduly placed on individual symptoms that may overlap rather than a comprehensive synthetic analysis of their quality, chronological sequence and other clustering features in the respective electro-clinical phenotypes, which is the basis for precise differential diagnosis in clinical practice. If any other diagnostic approach is followed, then even non-epileptic disorders such as migraine with aura could be deemed as overlapping with ICOE-G (visual hallucinations and headache), PS (lengthy duration and vomiting) or both (age, family history of epilepsies) (Table 1). It may be because of these limitations and the retrospective character of their study that Taylor *et al.* (2008) found that only one of their 16 patients was typical in all respects of PS and that ICOE-G was as frequent as PS, which contrasts all previous prospective studies detailed in this assessment. Such a discrepancy may indicate that PS is

still unrecognized and that the study does not represent the vast majority of typical PS.

Further, the commonly quoted argument that PS is not essentially different from ICOE-G considering that ‘the younger the children are, the less likely they are to describe visual symptoms’ (Andermann and Zifkin, 1998) is not tenable: (i) more than two-thirds of children with PS are older than 4 years and therefore able to describe their visual experiences and (ii) there is no difference in seizure presentation between younger and older children with PS.

A few patients with either PS or rolandic epilepsy may later develop purely occipital seizures as of ICOE-G (Guerrini *et al.*, 1997; Parmeggiani and Guerrini, 1999; Panayiotopoulos, 2002). These cases are easy to diagnose and indicate the intimate links of these disorders within the framework of BCSSS.

Benign childhood seizure susceptibility syndrome: a unified concept of benign childhood focal seizures

Rolandic epilepsy, PS, ICOE-G and other possible clinical phenotypes of benign childhood focal seizures are likely to be linked together by a genetically determined, functional derangement of the systemic brain maturation that is mild and age related (benign childhood seizure susceptibility syndrome) (Panayiotopoulos, 1993, 2002). They have distinctive characteristics but they also share common clinical and EEG features: seizures are infrequent, usually nocturnal and remit within a few years from onset. Brief or prolonged seizures, even focal status epilepticus, may occur only once in the patient’s lifetime. Despite the distinctiveness of their core clinical and EEG features, the natural histories of these syndromes may show significant reciprocity: some children with rolandic epilepsy may present autonomic seizures referable to PS (and vice versa) before remittance, while other may have alternate autonomic and rolandic seizures. Some seizures may be of mixed character, and certainly ictal autonomic manifestations, such as hypersalivation, emesis, headache and syncopal-like attacks that are unusual in other epileptic syndromes in children or adults, are frequent in BCSSS, and may predominate. Affected siblings may have the same or another type of benign childhood focal seizures, and febrile seizures are common. EEG spikes are regional (bilateral and multifocal) than focal—and as a rule disproportionately abundant to the frequency of seizures—and there is a significant overlap of interictal topographies.

There is no reason to suggest that these syndromes differ merely because an ‘epileptogenic’ focus is slightly anterior or posterior, lateral or medial to the central regions. The relevant ictal semeiologies and EEG findings suggest that each one of these forms reflects constitutional hyperexcitability of a particular functional brain area or system: the lower rolandic (somatosensory) cortex that represents the

face and the oropharynx bilaterally in benign rolandic epilepsy, the occipital areas (cortical visual system) in ICOE-G and the central autonomic network bilaterally and diffusely in PS (Koutroumanidis, 2007). All these conditions appear to be linked together by a genetically determined, functional derangement of the systemic brain maturation that is mild and age related (Panayiotopoulos, 1993, 1999a). This derangement is often clinically silent and presents in >90% of the susceptible children only with—also age related—EEG sharp and slow waves; the remaining one-tenth of these children have infrequent focal seizures. A small number of susceptible children, with or without seizures, may also have minor and fully reversible neuropsychological symptoms that are rarely clinically overt and can be detected only by formal neuropsychological testing. Finally, in a very small number of patients (probably <1%) this disturbance of brain maturation may further evolve into a more aggressive clinical state with seizures, neuropsychological manifestations and EEG abnormalities of various combinations and severity, such as atypical benign focal epilepsy of childhood, Landau–Kleffner syndrome and epilepsy with CSWS.

This concept of BCSSS is in agreement with previously expressed views of ‘functional epilepsies of maturation’ (Sorel and Rucquoy-Ponsar, 1969), ‘multifactorial pathogenesis of epilepsies with benign focal epileptiform sharp waves’ (Doose *et al.*, 1997, 2002), ‘selective rates of maturation of the different cortical areas (Luders *et al.*, 1987)’ and more recently of possible ‘neurobiological spectrum’ between BCSSS and IGE (Taylor *et al.*, 2003, 2008).

BCSSS, febrile and other idiopathic focal seizures in neonates and infants

One of the most interesting aspects of benign childhood seizures is their striking age-related sequence that appears to reflect enhanced epileptogenicity of the developing brain, as a whole and also of its functional systems, in different stages of maturation. Benign neonatal and infantile seizures, rolandic epilepsy, PS, ICOE-G and other clinical phenotypes of BCSSS are specific to early life and do not occur in adults. This is also the case with most febrile seizures whose different genetic influences may explain their high prevalence amongst patients with BCSSS and other more severe types of epilepsy, including the febrile plus phenotypes and genotypes (Scheffer *et al.*, 2007; Harkin *et al.*, 2007). It appears that there are three main periods of age-related childhood susceptibility to benign seizures: febrile, mainly generalized, convulsions first appear in early childhood at a peak age of around 18–22 months. Rolandic epilepsy and ICOE-G manifest with purely focal seizures and occur in late childhood age. PS presents with mainly autonomic seizures and covers the intermediate early childhood period with peak at 4 or 5 years. The neonatal and early infantile periods are not immune to focal seizure susceptibility either, as indicated by the benign neonatal seizures of the first few days

of life, (Plouin and Anderson, 2005) and the benign infantile focal seizures of Watanabe and Vigevano (Vigevano *et al.*, 2008). This point is exemplified by reports of children with neonatal seizures who later developed rolandic epilepsy (Maihara *et al.*, 1999) or PS (Lada *et al.*, 2003; Panayiotopoulos, 2007).

BCSSS, Landau–Kleffner syndrome, epilepsy with CSWS and atypical benign partial epilepsy of childhood

Landau–Kleffner syndrome and epilepsy with CSWS (Beaumanoir *et al.*, 1995) are now considered by the ILAE as an entity named ‘epileptic encephalopathy of CSWS including Landau–Kleffner syndrome’ with a common pathophysiological mechanism (Engel, 2006). Epileptic encephalopathy of CSWS including Landau–Kleffner syndrome are functional disorders occurring at an age where cortical synaptogenesis with abundant axonal sprouting and elemental functional network is being established in the brain. Aggressive epileptogenic activity at this active period of brain organization is detrimental for the establishment of appropriate neuronal connections, normal brain development and functioning (Smith and Hoeppner, 2003). These disorders may constitute a rare and extreme derailment of BCSSS. EEG manifests with abundant and often continuous high amplitude sharp waves morphologically similar to the CTS and occipital paroxysms. Seizures are predominantly nocturnal and often resemble rolandic seizures. Otherwise typical rolandic epilepsy (Fejerman *et al.*, 2000), PS (Caraballo *et al.*, 2001, 2005; Ferrie *et al.*, 2002) and ICOE-G (Tenenbaum *et al.*, 1997) evolve to clinical and EEG features of epileptic encephalopathy of CSWS including Landau–Kleffner syndrome and atypical partial epilepsy of childhood (Saltik *et al.*, 2005). Atypical benign partial epilepsy of childhood (Aicardi and Chevrie, 1982) is considered either a mild form of epileptic encephalopathy of CSWS (Arzimanoglou *et al.*, 2004) or a more severe form of rolandic epilepsy (Doose *et al.*, 2001; Hahn *et al.*, 2001). The reason for this derailment of BCSSS is unknown, but may be related to location, epileptogenic threshold and other intrinsic genetic and external superimposed factors. Intense epileptic activity in the dominant temporal region would affect linguistic capabilities as in Landau–Kleffner syndrome (Morrell *et al.*, 1995). Conversely, the mainly frontal localization of CSWS primarily affects higher cognitive and executive functioning (Galanopoulou *et al.*, 2000; Smith and Hoeppner, 2003; Tassinari *et al.*, 2005).

BCSSS and idiopathic generalized epilepsies

The majority of BCSSS if properly diagnosed do not have any clinical or EEG resemblance to idiopathic generalized epilepsies, though others may disagree (Taylor *et al.*, 2003, 2008). Overlap of BCSSS with IGE is limited (Kivity *et al.*,

2000). However, a possible link, the type and extent of which should be explored further with clinical and genetic studies may be suggested by

- (a) the occurrence of EEG generalized discharges in BCSSS (though these are often markedly different from the classical generalized spike or polyspike discharges of IGE);
- (b) an undetermined but probably small proportion of patients with any type of BCSSS that may also suffer typical generalized convulsive or absence seizures either during the active phase of BCSSS or more often at a later stage and
- (c) an undetermined but probably small proportion of patients with syndromes of IGE including childhood absence epilepsy that may also have EEG focal spikes only or together with any type of seizures of BCSSS (Beaumanoir *et al.*, 1974; Taylor *et al.*, 2003, 2008; Caraballo *et al.*, 2008b).

Management of benign childhood focal seizures

Short- and long-term treatment strategies of benign childhood focal seizures are largely empirical.

In the acute stage, control of the seizure is of paramount importance. Benzodiazepines are used to terminate long-lasting seizures (>10 min) and status epilepticus. Autonomic status epilepticus needs thorough evaluation for proper diagnosis and assessment of the neurological/autonomic state of the child. Aggressive treatment should be avoided because of the risk of iatrogenic complications (Ferrie *et al.*, 2006).

Continuous anti-epileptic medication is not usually recommended. Although there may be effective AED that may prevent the occurrence of additional seizures, potential adverse effects may not commensurate with the benefit. Decisions on management must take into account the following (Bourgeois, 2000; Wirrell *et al.*, 2006; Panayiotopoulos, 2007; Fejerman, 2008):

- (a) Most children have excellent prognosis: 10–30% may have only a single seizure and 60–70% may have less than 10 in total. However, 10–20% of children may have frequent seizures, which are sometimes resistant to treatment.
- (b) Remission of benign childhood focal seizures is expected in all patients by the age of 15–16 years at the latest.
- (c) There is no evidence that the long-term prognosis is worse in untreated children (Ambrosetto and Tassinari, 1990), although they may not be protected against seizure recurrences and particularly GTCS (Peters *et al.*, 2001).
- (d) Some children become frightened, even by simple focal seizures, and some parents are unable to cope

with the possibility of another fit despite firm reassurances.

- (e) Persistence and frequency of EEG functional spikes do not predict clinical severity, frequency or degree of liability to seizures
- (f) In contrast to the other phenotypes of the BCSSS, patients with ICOE-G often suffer from frequent seizures and therefore prophylactic AED treatment may be mandatory. Secondly, GTCS are probably unavoidable without medication.

Continuous prophylaxis consists of daily monotherapy using any AED that has proven efficacy in focal seizures and minimal adverse effects in children. The 2006 ILAE treatment guidelines found that ‘no AED had level A or level B efficacy and effectiveness evidence as initial monotherapy’ in rolandic epilepsy (Glauser *et al.*, 2006). Of older AED most authorities prefer carbamazepine in USA (Wheless *et al.*, 2005) and valproic acid in Europe (Wheless *et al.*, 2007), though these may have equivalent efficacy with all others (phenobarbital, phenytoin and clonazepam) (Ferrie *et al.*, 1997; Bouma *et al.*, 2002); carbamazepine may exaggerate seizures in a minority of children with BCSSS including PS (Kikumoto *et al.*, 2006) and valproate is associated with significant adverse effects. Recently, sulthiame (available only in a few countries) has been revived as an excellent drug for the treatment of rolandic epilepsy with EEG normalization (Bast *et al.*, 2003; Ben Zeev *et al.*, 2004) but this may be associated with cognitive abnormalities (Wirrell *et al.*, 2008). Recommended newer AED include levetiracetam (Verrotti *et al.*, 2007; Kossoff *et al.*, 2007), gabapentin (Bourgeois, 2000) and lamotrigine (Wheless *et al.*, 2005). Lamotrigine on rare occasions may cause seizure exacerbation and cognitive deterioration (Cerinara *et al.*, 2004).

Opinions of when to withdraw medication differ among experts, although all agree that there is no need to continue with AED after the age 14 when most benign childhood focal seizures remit or 16 when they are practically non-existent; a common recommendation is to start slow tapering the AED after a 1–3 years seizure free period (Bourgeois, 2000).

Parental needs and support

Despite their excellent prognosis, benign childhood focal seizures usually have a dramatic impact to parents. The most dominant points of parental anxiety and concerns refer to their uncertainty of the nature, the cause and the impact of the events on their child’s development as well as the lack of appropriate information (Valeta, 2005). There is a need for supportive family management, education and specific instructions about emergency procedures for possible subsequent seizures. Education about BCSSS is the cornerstone of the optimal management. Parents should be given general information about benign childhood focal seizures and, in particular PS, in which seizures may have dramatic

features and last for many hours; the situation is frequently compounded by physicians' uncertainty over diagnosis, management and prognosis. Parents who have watched their child during a seizure may need specific psychological support to overcome anxiety and panic that may result in overprotection and interfere in turn with parent–child separation and independence (Valeta, 2007).

Conclusions and recommendations

It is apparent from this assessment that BCSSS is a formidable challenge in childhood epileptology. Despite significant progress made, we may have presented only the tip of the iceberg and it is likely that significant pieces are still missing or we may have misplaced them in the jigsaw puzzle. There are many practical, diagnostic, genetic, aetiological, electrophysiological, pathophysiological, theoretical, epidemiological, social and management issues still to be resolved. This will require a prospective multidisciplinary and evidence-based approach. Significant evidence of the processes underpinning these diseases may emerge and our old notions discarded. Furthermore, revisions should be proposed in the best spirit of unbiased work with an open-minded and dispassionate scientific decorum. Meanwhile, from the clinical point of view their phenotypic recognition and their differentiation from other epileptic and non-epileptic disorders have significant prognostic and management implications.

References

- Aicardi J, Chevrie JJ. Atypical benign partial epilepsy of childhood. *Dev Med Child Neurol* 1982; 24: 281–92.
- Ambrosetto G, Tassinari CA. Antiepileptic drug treatment of benign childhood epilepsy with rolandic spikes: is it necessary? *Epilepsia* 1990; 31: 802–5.
- Andermann F, Zifkin B. The benign occipital epilepsies of childhood: an overview of the idiopathic syndromes and of the relationship to migraine. *Epilepsia* 1998; 39 (Suppl 4): S9–23.
- Arzimanoglou A, Guerrini R, Aicardi J, editors. *Aicardi's epilepsy in children*. Philadelphia: Lippincott Williams & Wilkins; 2004.
- Aso K, Watanabe K, Negoro T, Takaesu E, Furune A, Takahashi I, et al. Visual seizures in children. *Epilepsy Res* 1987; 1: 246–53.
- Astradsson A, Olafsson E, Ludvigsson P, Bjorgvinsson H, Hauser WA. Rolandic epilepsy: an incidence study in Iceland. *Epilepsia* 1998; 39: 884–6.
- Bagdordf R, Lee SI. Midline spikes: is it another benign EEG pattern of childhood? *Epilepsia* 1993; 34: 271–4.
- Bali B, Kull LL, Strug LJ, Clarke T, Murphy PL, Akman CI, et al. Autosomal dominant inheritance of centrotemporal sharp waves in rolandic epilepsy families. *Epilepsia* 2007; 48: 2266–72.
- Bast T, Volp A, Wolf C, Rating D. The influence of sulthiame on EEG in children with benign childhood epilepsy with centrotemporal spikes (BECTS). *Epilepsia* 2003; 44: 215–20.
- Beaumanoir A. Infantile epilepsy with occipital focus and good prognosis. *Eur Neurol* 1983; 22: 43–52.
- Beaumanoir A. An EEG contribution to the study of migraine and of the association between migraine and epilepsy in childhood. In: Andermann F, Beaumanoir A, Mira L, Roger J, Tassinari CA, editors. *Occipital seizures and epilepsy in children*. London: John Libbey & company Ltd; 1993. p. 101–10.
- Beaumanoir A. Semiology of occipital seizures in infants and children. In: Andermann F, Beaumanoir A, Mira L, Roger J, Tassinari CA, editors. *Occipital seizures and epilepsies in children*. London: John Libbey and Company Ltd; 1993. p. 71–86.
- Beaumanoir A, Ballis T, Varfis G, Ansari K. Benign epilepsy of childhood with Rolandic spikes. A clinical, electroencephalographic, and telencephalographic study. *Epilepsia* 1974; 15: 301–15.
- Beaumanoir A, Bureau M, Deonna T, Mira L, Tassinari CA. Continuous spike and waves during slow sleep. Electrical status epilepticus during slow sleep. London: John Libbey & company; 1993.
- Beaumanoir A, Capizzi G, Nahory A, Yousfi Y. Scotogenic seizures. In: Beaumanoir A, Gastaut H, Roger J, editors. *Reflex seizures and reflex epilepsies*. Geneva: Medecine and Hygiene; 1989. p. 219–23.
- Beaumanoir A, Nahory A. Benign partial epilepsies: 11 cases of frontal partial epilepsy with favorable prognosis. [French]. *Revue d'Electroencephalographie et de Neurophysiologie Clinique* 1983; 13: 207–11.
- Beaussart M. Benign epilepsy of children with Rolandic (centro-temporal) paroxysmal foci. A clinical entity. Study of 221 cases. *Epilepsia* 1972; 13: 795–911.
- Beaussart M, Faou R. Evolution of epilepsy with rolandic paroxysmal foci: a study of 324 cases. *Epilepsia* 1978; 19: 337–42.
- Beaussart M, Loiseau P, Roger H. The discovery of 'benign rolandic epilepsy'. In: Berkovic SF, Genton P, Hirsch E, Picard F, editors. *Genetics of focal epilepsies*. London: John Libbey & Company; 1999. p. 3–6.
- Ben Zeev B, Waternberg N, Lerman P, Barash I, Brand N, Lerman-Sagie T. Sulthiame in childhood epilepsy. *Pediatr Int* 2004; 46: 521–4.
- Berg AT, Shinnar S, Levy SR, Testa FM. Newly diagnosed epilepsy in children: presentation at diagnosis. *Epilepsia* 1999; 40: 445–52.
- Beydoun A, Garofalo EA, Drury I. Generalized spike-waves, multiple loci, and clinical course in children with EEG features of benign epilepsy of childhood with centrotemporal spikes. *Epilepsia* 1992; 33: 1091–6.
- Blom S, Brorson LO. Central spikes or sharp waves (Rolandic spikes) in children's EEG and their clinical significance. *Acta Paediatrica Scandinavica* 1966; 55: 385–93.
- Blom S, Heijbel J. Benign epilepsy of children with centrotemporal EEG foci: a follow-up study in adulthood of patients initially studied as children. *Epilepsia* 1982; 23: 629–32.
- Boatman DF, Trescher WH, Smith C, Ewen J, Los J, Wied HM, et al. Cortical auditory dysfunction in benign rolandic epilepsy. *Epilepsia* 2008; 49: 1018–26.
- Boor R, Jacobs J, Hinzmann A, Bauermann T, Scherg M, Boor S, et al. Combined spike-related functional MRI and multiple source analysis in the non-invasive spike localization of benign rolandic epilepsy. *Clin Neurophysiol* 2007; 118: 901–9.
- Bouma PA, Bovenkerk AC, Westendorp RG, Brouwer OF. The course of benign partial epilepsy of childhood with centrotemporal spikes: a meta-analysis. *Neurology* 1997; 48: 430–7.
- Bourgeois BF. Drug treatment of benign focal epilepsies of childhood. *Epilepsia* 2000; 41: 1057–8.
- Bray PF, Wiser WC. Hereditary characteristics of familial temporal-central focal epilepsy. *Pediatric* 1965; 36: 207–21.
- Bulgheroni S, Franceschetti S, Vago C, Usilla A, Pantaleoni C, D'Arrigo S, et al. Verbal dichotic listening performance and its relationship with EEG features in benign childhood epilepsy with centrotemporal spikes. *Epilepsy Res* 2008; 79: 31–8.
- Bureau M, Cokar O, Maton B, Genton P, Dravet C. Sleep-related, low voltage Rolandic and vertex spikes: an EEG marker of benignity in infancy-onset focal epilepsies. *Epileptic Disord* 2002; 4: 15–22.
- Capovilla G, Beccaria F, Montagnini A. 'Benign focal epilepsy in infancy with vertex spikes and waves during sleep'. Delineation of the syndrome and recalling as 'benign infantile focal epilepsy with midline spikes and waves during sleep' (BIMSE). *Brain Dev* 2006; 28: 85–91.
- Caraballo R, Cersosimo R, Fejerman N. Panayiotopoulos syndrome: a prospective study of 192 patients. *Epilepsia* 2007; 48: 1054–61.
- Caraballo R, Fontana E, Michelizza B, Zullini B, Sgro V, Pajno-Ferrara F, et al. Carbamazepine inducing atypical absences, atonic seizures and continuous spike and waves during slow wave sleep. *Boll Lega It Epil* 1989; 66/67: 379–81.

- Caraballo RH, Astorino F, Cersosimo R, Soprano AM, Fejerman N. Atypical evolution in childhood epilepsy with occipital paroxysms (Panayiotopoulos type). *Epileptic Disord* 2001; 3: 157–62.
- Caraballo RH, Cersosimo RO, Fejerman N. Late-onset. 'Gastaut type', childhood occipital epilepsy: an unusual evolution. *Epileptic Disord* 2005; 7: 341–6.
- Caraballo RH, Cersosimo RO, Fejerman N. Childhood occipital epilepsy of Gastaut: a study of 33 patients. *Epilepsia* 2008; 49: 288–97.
- Caraballo RH, Fontana E, Darra F, Bongiorno L, Fiorini E, Cersosimo R, et al. Childhood absence epilepsy and electroencephalographic focal abnormalities with or without clinical manifestations. *Seizure* 2008. Advance Access published on June 3, 2008, doi:10.1016/j.seizure.2008.03.009.
- Cavazzuti GB. Epidemiology of different types of epilepsy in school age children of Modena, Italy. *Epilepsia* 1980; 21: 57–62.
- Cavazzuti GB, Cappella L, Nalin A. Longitudinal study of epileptiform EEG patterns in normal children. *Epilepsia* 1980; 21: 43–55.
- Cerminara C, Montanaro ML, Curatolo P, Seri S. Lamotrigine-induced seizure aggravation and negative myoclonus in idiopathic rolandic epilepsy. *Neurology* 2004; 63: 373–5.
- Chelmsky TC, Chelmsky GG. Autonomic abnormalities in cyclic vomiting syndrome. *J Pediatr Gastroenterol Nutr* 2007; 44: 326–30.
- Clemens B, Majoros E. Sleep studies in benign epilepsy of childhood with rolandic spikes. II. Analysis of discharge frequency and its relation to sleep dynamics. *Epilepsia* 1987; 28: 24–7.
- Colamaria V, Sgro V, Caraballo R, Simeone M, Zullini E, Fontana E, et al. Status epilepticus in benign rolandic epilepsy manifesting as anterior operculum syndrome. *Epilepsia* 1991; 32: 329–34.
- Covanis A. Panayiotopoulos syndrome: a benign childhood autonomic epilepsy frequently imitating encephalitis, syncope, migraine, sleep disorder, or gastroenteritis. *Pediatrics* 2006; 118: e1237–43.
- Covanis A, Ferrie CD, Koutroumanidis M, Oguni H, Panayiotopoulos CP. Panayiotopoulos syndrome and Gastaut type idiopathic childhood occipital epilepsy. In: Roger J, Bureau M, Dravet C, Genton P, Tassinari CA, Wolf P, editors. *Epileptic syndromes in infancy, childhood and adolescence*. (fourth edition with video). Montrouge, France: John Libbey Eurotext; 2005. p. 227–53.
- Covanis A, Lada C, Skiadas K. Children with rolandic spikes and ictus emeticus: rolandic epilepsy or Panayiotopoulos syndrome? *Epileptic Disord* 2003; 5: 139–43.
- Cowan LD. The epidemiology of the epilepsies in children. *Ment Retard Dev Disabil Res Rev* 2002; 8: 171–81.
- Dalla Bernardina B, Colamaria V, Chiamenti C, Capovilla G, Trevisan E, Tassinari CA. Benign partial epilepsy with affective symptoms ('benign psychomotor epilepsy'). In: Roger J, Bureau M, Dravet C, Dreifuss FE, Perret A, Wolf P, editors. *Epileptic syndromes in infancy, childhood and adolescence*. 2nd edn. London: John Libbey & Company Ltd; 1992. p. 219–23.
- Dalla Bernardina B, Fontana E, Darra F. Are there other types of benign focal epilepsies in childhood? In: Fejerman N, Caraballo RH, editors. *Benign focal epilepsies in infancy, childhood and adolescence*. Montrouge, France: John Libbey Eurotext; 2007. p. 169–78.
- Dalla Bernardina B, Sgro M, Fejerman N. Epilepsy with centro-temporal spikes and related syndromes. In: Roger J, Bureau M, Dravet C, Genton P, Tassinari CA, Wolf P, editors. *Epileptic syndromes in infancy, childhood and adolescence*. 4th edn. Montrouge, France: John Libbey Eurotext; 2005. p. 203–25.
- Datta A, Sinclair DB. Benign epilepsy of childhood with rolandic spikes: typical and atypical variants. *Pediatr Neurol* 2007; 36: 141–5.
- De Marco P, Tassinari CA. Extreme somatosensory evoked potential (ESEP): an EEG sign forecasting the possible occurrence of seizures in children. *Epilepsia* 1981; 22: 569–75.
- De Romanis F, Buzzi MG, Cerbo R, Feliciani M, Assenza S, Agnoli A. Migraine and epilepsy with infantile onset and electroencephalographic findings of occipital spike-wave complexes. *Headache* 1991; 31: 378–83.
- De Romanis F, Feliciani M, Cerbo R. Migraine and other clinical syndromes in children affected by EEG occipital spike-wave complexes. *Funct Neurol* 1988; 3: 187–203.
- Demirbilek V, Dervent A. Panayiotopoulos syndrome: video-EEG illustration of a typical seizure. *Epileptic Disord* 2004; 6: 121–4.
- Deonna T, Ziegler AL, Despland PA, van Melle G. Partial epilepsy in neurologically normal children: clinical syndromes and prognosis. *Epilepsia* 1986; 27: 241–7.
- Deonna TW, Roulet E, Fontan D, Marcoz JP. Speech and oromotor deficits of epileptic origin in benign partial epilepsy of childhood with rolandic spikes (BPERS). Relationship to the acquired aphasia-epilepsy syndrome. *Neuropediatrics* 1993; 24: 83–7.
- Dooze H, Brigger-Heuer B, Neubauer B. Children with focal sharp waves: clinical and genetic aspects. *Epilepsia* 1997; 38: 788–96.
- Dooze H, Hahn A, Neubauer BA, Pistohl J, Stephani U. Atypical 'benign' partial epilepsy of childhood or pseudo-lennox syndrome. Part II: family study. *Neuropediatrics* 2001; 32: 9–13.
- Dooze H, Petersen B, Neubauer BA. Occipital sharp waves in idiopathic partial epilepsies-clinical and genetic aspects. *Epilepsy Res* 2002; 48: 121–30.
- Drury I, Beydoun A. Benign partial epilepsy of childhood with monomorphic sharp waves in centrottemporal and other locations. *Epilepsia* 1991; 32: 662–7.
- Dura-Trave T, Yoldi-Petri ME, Gallinas-Victoriano F. Panayiotopoulos syndrome: epidemiological and clinical characteristics and outcome. *Eur J Neurol* 2008; 15: 336–41.
- Elger CE. Semeiology of temporal lobe seizures. In: Oxbury JM, Polkey CE, Duchowny M, editors. *Intractable focal epilepsy*. London: W.B. Saunders; 2000. p. 63–8.
- Engel J Jr. A proposed diagnostic scheme for people with epileptic seizures and with epilepsy: report of the ILAE task force on classification and terminology. *Epilepsia* 2001; 42: 796–803.
- Engel J Jr. Report of the ILAE classification core group. *Epilepsia* 2006; 47: 1558–68.
- Fejerman N. Benign childhood epilepsy with centrottemporal spikes. In: Engel J Jr, Pedley TA, editors. *Philadelphia: Lippincott Williams & Wilkins, A Wolters Kluwer Business*; 2008. p. 2369–77.
- Fejerman N, Caraballo R, Tenenbaum SN. Atypical evolutions of benign localization-related epilepsies in children: are they predictable? *Epilepsia* 2000; 41: 380–90.
- Fejerman N, Caraballo RH, editors. *Benign focal epilepsies in infancy, childhood and adolescence*. Montrouge, France: John Libbey Eurotext; 2007.
- Ferrie C, Caraballo R, Covanis A, Demirbilek V, Dervent A, Kivity S, et al. Panayiotopoulos syndrome: a consensus view. *Dev Med Child Neurol* 2006; 48: 236–40.
- Ferrie CD, Beaumanoir A, Guerrini R, Kivity S, Vigeveno F, Takaishi Y, et al. Early-onset benign occipital seizure susceptibility syndrome. *Epilepsia* 1997; 38: 285–93.
- Ferrie CD, Caraballo R, Covanis A, Demirbilek V, Dervent A, Fejerman N, et al. Autonomic status epilepticus in Panayiotopoulos syndrome and other childhood and adult epilepsies: a consensus view. *Epilepsia* 2007; 48: 1165–72.
- Ferrie CD, Koutroumanidis M, Rowlinson S, Sanders S, Panayiotopoulos CP. Atypical evolution of Panayiotopoulos syndrome: a case report [published with video-sequences]. *Epileptic Disord* 2002; 4: 35–42.
- Fonseca LC, Tedrus GM. Somatosensory evoked spikes and epileptic seizures: a study of 385 cases. *Clin Electroencephalogr* 2000; 31: 71–5.
- Forsgren L, Beghi E, Oun A, Sillanpaa M. The epidemiology of epilepsy in Europe - a systematic review. *Eur J Neurol* 2005; 12: 245–53.
- Freeman R. Cardiovascular manifestations of autonomic epilepsy. *Clin Auton Res* 2006; 16: 12–7.
- Galanopoulou AS, Bojko A, Lado F, Moshe SL. The spectrum of neuropsychiatric abnormalities associated with electrical status epilepticus in sleep. *Brain Dev* 2000; 22: 279–95.

- Gastaut H. Benign epilepsy with occipital spike waves in children. [French]. Bulletin et Memoires de l'Academie Royale de Medecine de Belgique 1981; 136: 540–55.
- Gastaut H. A new type of epilepsy: benign partial epilepsy of childhood with occipital spike-waves. Clin Electroencephalogr 1982; 13: 13–22.
- Gastaut H. The benigne epilepsy in childhood with occipital spike wave complexes. [German]. EEG-EMG Zeitschrift fur Elektroenzephalographie Elektromyographie und Verwandte Gebiete 1982; 13: 3–8.
- Gastaut H, Roger J, Bureau M. Benign epilepsy of childhood with occipital paroxysms. Up-date. In: Roger J, Bureau M, Dravet C, Dreifuss FE, Perret A, Wolf P, editors. Epileptic syndromes in infancy, childhood and adolescence. 2nd edn. London: John Libbey & Company Ltd; 1992. p. 201–17.
- Gastaut H, Zifkin BG. Benign epilepsy of childhood with occipital spike and wave complexes. In: Andermann F, Lugaesi E, editors. Migraine and epilepsy. Boston: Butterworths; 1987. p. 47–81.
- Gelisse P, Genton P, Bureau M, Dravet C, Guerrini R, Viallat D, et al. Are there generalised spike waves and typical absences in benign rolandic epilepsy? Brain Dev 1999; 21: 390–6.
- Genizi J, Zelnik N, Ravid S, Shahar E. Childhood epilepsy with occipital paroxysms: difficulties in distinct segregation into either the early-onset or late-onset epilepsy subtypes. J Child Neurol 2007; 22: 588–92.
- Germano E, Gagliano A, Magazu A, Sferro C, Calarese T, Mannarino E, et al. Benign childhood epilepsy with occipital paroxysms: neuropsychological findings. Epilepsy Res 2005; 64: 137–50.
- Gibbs FA, Gibbs EL. Atlas of electroencephalography. Vol. 2. Epilepsy. Reading, MA: Addison-Wesley; 1952.
- Gibbs FA, Gibbs EL. Medical electroencephalography. Reading, MA: Addison-Wesley Publishing Company; 1967.
- Giordani B, Caveney AF, Laughrin D, Huffman JL, Berent S, Sharma U, et al. Cognition and behavior in children with benign epilepsy with centrotemporal spikes (BECTS). Epilepsy Res 2006; 70: 89–94.
- Glauser T, Ben-Menachem E, Bourgeois B, Cnaan A, Chadwick D, Guerreiro C, et al. ILAE treatment guidelines: evidence-based analysis of antiepileptic drug efficacy and effectiveness as initial monotherapy for epileptic seizures and syndromes. Epilepsia 2006; 47: 1094–120.
- Gobbi G, Guerrini R, Grosso S. Late onset childhood occipital epilepsy (Gastaut type). In: Engel J Jr, Pedley TA, editors. Philadelphia: Lippincott Williams & Wilkins, A Wolters Kluwer Business; 2008. p. 2387–95.
- Goodman JH, Stewart M, Drislane FW. Autonomic disturbances. In: Engel J Jr, Pedley TA, editors. Philadelphia: Lippincott Williams & Wilkins, A Wolters Kluwer Business; 2008. p. 1999–2005.
- Gregory DL, Wong PK. Clinical relevance of a dipole field in rolandic spikes. Epilepsia 1992; 33: 36–44.
- Grosso S, Orrico A, Galli L, Di BR, Sorrentino V, Balestri P. SCN1A mutation associated with atypical Panayiotopoulos syndrome. Neurology 2007; 69: 609–11.
- Grosso S, Vivarelli R, Gobbi G, Bartolo RD, Berardi R, Balestri P. Late-onset childhood occipital epilepsy (Gastaut type): A family study. Eur J Paediatr Neurol 2008. Advance Access published on January 30, 2008, doi:10.1016/j.ejpn.2007.11.007.
- Guerrini R, Bonanni P, Parmeggiani L, Belmonte A. Adolescent onset of idiopathic photosensitive occipital epilepsy after remission of benign rolandic epilepsy. Epilepsia 1997; 38: 777–81.
- Guerrini R, Bonanni P, Parmeggiani L, Thomas P, Mattia D, Harvey AS, et al. Induction of partial seizures by visual stimulation. Clinical and electroencephalographic features and evoked potential studies. Adv Neurol 1998; 75: 159–78.
- Guerrini R, Dravet C, Genton P, Bureau M, Bonanni P, Ferrari AR, et al. Idiopathic photosensitive occipital lobe epilepsy. Epilepsia 1995; 36: 883–91.
- Gulgonen S, Demirbilek V, Korkmaz B, Derwent A, Townes BD. Neuropsychological functions in idiopathic occipital lobe epilepsy. Epilepsia 2000; 41: 405–11.
- Hahn A, Pistohl J, Neubauer BA, Stephani U. Atypical 'benign' partial epilepsy or pseudo-Lennox syndrome. Part I: symptomatology and long-term prognosis. Neuropediatrics 2001; 32: 1–8.
- Harkin LA, McMahon JM, Iona X, Dibbens L, Pelekanos JT, Zuberi SM, et al. The spectrum of SCN1A-related infantile epileptic encephalopathies. Brain 2007; 130: 843–52.
- Heijbel J, Blom S, Bergfors PG. Benign epilepsy of children with centrotemporal EEG foci. A study of incidence rate in outpatient care. Epilepsia 1975; 16: 657–64.
- Herranz Tanarro FJ, Saenz Lope E, Cristobal Sassot S. Occipital spike-wave with or without benign childhood epilepsy. [French]. Revue d'Electroencephalographie et de Neurophysiologie Clinique 1984; 14: 1–7.
- Huiskamp G, Van der MW, Van Huffelen A, Van Nieuwenhuizen O. High resolution spatio-temporal EEG-MEG analysis of rolandic spikes. J Clin Neurophysiol 2004; 21: 84–95.
- Jallon P, Loiseau P, Loiseau J. Newly diagnosed unprovoked epileptic seizures: presentation at diagnosis in CAROLE study. Coordination Active du Réseau Observatoire Longitudinal de l'Epilepsie. Epilepsia 2001; 42: 464–75.
- Jung KY, Kim JM, Kim DW. Patterns of interictal spike propagation across the central sulcus in benign rolandic epilepsy. Clin Electroencephalogr 2003; 34: 153–7.
- Kajitani T, Kimura T, Sumita M, Kaneko M. Relationship between benign epilepsy of children with centro-temporal EEG foci and febrile convulsions. Brain Dev 1992; 14: 230–4.
- Kanazawa O, Tohyama J, Akasaka N, Kamimura T. A magnetoencephalographic study of patients with Panayiotopoulos syndrome. Epilepsia 2005; 46: 1106–13.
- Kavros PM, Clarke T, Strug LJ, Halperin JM, Dorta NJ, Pal DK. Attention impairment in rolandic epilepsy: Systematic review. Epilepsia 2008. Advance Access published on April 11, 2008, doi:10.1111/j.1528-1167.2008.01610.x.
- Kellaway P. The incidence, significance and natural history of spike foci in children. In: Henry CE, editor. Current clinical neurophysiology. update on EEG and evoked potentials. New York: Elsevier/North Holland; 1980. p. 151–75.
- Kikumoto K, Yoshinaga H, Oka M, Ito M, Endoh F, Akiyama T, et al. EEG and seizure exacerbation induced by carbamazepine in Panayiotopoulos syndrome. Epileptic Disord 2006; 8: 53–6.
- Kivity S, Ephraim T, Weitz R, Tamir A. Childhood epilepsy with occipital paroxysms: clinical variants in 134 patients. Epilepsia 2000; 41: 1522–33.
- Kivity S, Lerman P. Stormy onset with prolonged loss of consciousness in benign childhood epilepsy with occipital paroxysms. J Neurol Neurosurg Psychiatr 1992; 55: 45–8.
- Kossoff EH, Los JG, Boatman DF. A pilot study transitioning children onto levetiracetam monotherapy to improve language dysfunction associated with benign rolandic epilepsy. Epilepsy Behav 2007; 11: 514–7.
- Koutroumanidis M. Ictal vomiting in association with left temporal lobe seizures in a left hemisphere language-dominant patient. Epilepsia 2003; 44: 1259.
- Koutroumanidis M. Panayiotopoulos syndrome: an important electro-clinical example of benign childhood system epilepsy. Epilepsia 2007; 48: 1044–53.
- Koutroumanidis M, Aggelakis K, Panayiotopoulos CP. Idiopathic epilepsy with generalized tonic clonic seizures only vs. idiopathic epilepsy with phantom absences and generalized tonic clonic seizures: one or two syndromes? Epilepsia 2008. Advance Access published on July 4, 2008, doi:10.1111/j.1528-1167.2008.01702.x.
- Koutroumanidis M, Rowlinson S, Sanders S. Recurrent autonomic status epilepticus in Panayiotopoulos syndrome: video/EEG studies. Epilepsy Behav 2005; 7: 543–7.
- Kramer RE, Luders H, Goldstick LP, Dinner DS, Morris HH, Lesser RP, et al. Ictus emeticus: an electroclinical analysis. Neurology 1988; 38: 1048–52.
- Kubota M, Takeshita K, Sakakihara Y, Yanagisawa M. Magnetoencephalographic study of giant somatosensory evoked responses in patients with rolandic epilepsy. J Child Neurol 2000; 15: 370–9.
- Kutluay E, Passaro EA, Gomez-Hassan D, Beydoun A. Seizure semiology and neuroimaging findings in patients with midline spikes. Epilepsia 2001; 42: 1563–8.

- Kuzniecky R, Gilliam F, Morawetz R, Faught E, Palmer C, Black L. Occipital lobe developmental malformations and epilepsy: clinical spectrum, treatment, and outcome. *Epilepsia* 1997; 38: 175–81.
- Lada C, Skiadas K, Theodorou V, Covanis A. A study of 43 patients with Panayiotopoulos syndrome: a common and benign childhood seizure susceptibility. *Epilepsia* 2003; 44: 81–8.
- Langill L, Wong PK. Tactile-evoked rolandic discharges: a benign finding? *Epilepsia* 2003; 44: 221–7.
- Larson K, Eeg-Olofsson O. A population based study of epilepsy in children from a Swedish county. *Eur J Paediatr Neurol* 2006; 10: 107–13.
- Leal AJ, Ferreira JC, Dias AI, Calado E. Origin of frontal lobe spikes in the early onset benign occipital lobe epilepsy (Panayiotopoulos syndrome). *Clin Neurophysiol* 2008. Advance Access published on July 11, 2008, doi:10.1016/j.clinph.2008.04.299.
- Leal AJ, Nunes S, Dias AI, Vieira JP, Moreira A, Calado E. Analysis of the generators of epileptic activity in early-onset childhood benign occipital lobe epilepsy. *Clin Neurophysiol* 2007; 118: 1341–7.
- Legarda S, Jayakar P, Duchowny M, Alvarez L, Resnick T. Benign rolandic epilepsy: high central and low central subgroups. *Epilepsia* 1994; 35: 1125–9.
- Lerman P, Kivity S. Benign focal epilepsy of childhood. A follow-up study of 100 recovered patients. *Arch Neurol* 1975; 32: 261–4.
- Li BU, Issenman RM, Sarna SK. Consensus statement—2nd International Scientific Symposium on CVS. The faculty of the 2nd International Scientific Symposium on Cyclic Vomiting Syndrome. *Dig Dis Sci* 1999; 44 (8 Suppl): 9S–11S.
- Livingston J, Cross H, Birch R, Zuberi S. Inherited SCN1A mutation in siblings with Panayiotopoulos syndrome. *Dev Med Child Neurol* 2008; 50 (Suppl 112): 3.
- Loiseau P, Duche B, Cordova S, Dartigues JF, Cohadon S. Prognosis of benign childhood epilepsy with centrotemporal spikes: a follow-up study of 168 patients. *Epilepsia* 1988; 29: 229–35.
- Loiseau P, Pestre M, Dartigues JF, Commenges D, Barberger-Gateau C, Cohadon S. Long-term prognosis in two forms of childhood epilepsy: typical absence seizures and epilepsy with rolandic (centrotemporal) EEG foci. *Ann Neurol* 1983; 13: 642–8.
- Luders HO, Lesser RP, Dinner DS, Morris III HH. Benign focal epilepsy of childhood. In: Luders H, Lesser RP, editors. *Epilepsy. Electroclinical syndromes*. Berlin, Heidelberg: Springer; 1987. p. 303–46.
- Maihara T, Tsuji M, Higuchi Y, Hattori H. Benign familial neonatal convulsions followed by benign epilepsy with centrotemporal spikes in two siblings. *Epilepsia* 1999; 40: 110–3.
- Manganotti P, Miniussi C, Santorum E, Tinazzi M, Bonato C, Marzi CA, et al. Influence of somatosensory input on paroxysmal activity in benign rolandic epilepsy with ‘extreme somatosensory evoked potentials’. *Brain* 1998; 121: 647–58.
- Martin-Santidrian MA, Garaizar C, Prats-Vinas JM. [Frontal lobe epilepsy in infancy: is there a benign partial frontal lobe epilepsy?]. *Rev Neurol* 1998; 26: 919–23.
- Martinovic Z. The new ILAE report on classification and evidence-based commentary on Panayiotopoulos syndrome and autonomic status epilepticus. *Epilepsia* 2007; 48: 1215–6.
- Michelucci R, Tassinari CA. Television-induced occipital seizures. In: Andermann F, Beaumanoir A, Mira L, Roger J, Tassinari CA, editors. *Occipital seizures and epilepsies in children*. London: John Libbey & Company Ltd; 1993. p. 141–4.
- Minami T, Gondo K, Yamamoto T, Yanai S, Tasaki K, Ueda K. Magnetoencephalographic analysis of rolandic discharges in benign childhood epilepsy. *Ann Neurol* 1996; 39: 326–34.
- Morrell F, Whisler WW, Smith MC, Hoepfner TJ, de Toledo-Morrell L, Pierre-Louis SJ, et al. Landau-Kleffner syndrome. Treatment with subpial intracortical transection. *Brain* 1995; 118: 1529–46.
- Nagendran K, Prior PF, Rossiter MA. Benign occipital epilepsy of childhood: a family study. *J R Soc Med* 1990; 83: 804–5.
- Negrin P, De Marco P. Parietal focal spikes evoked by tactile somatotopic stimulation in sixty non-epileptic children: the nocturnal sleep and clinical and EEG evolution. *Electroencephalogr Clin Neurophysiol* 1977; 43: 312–6.
- Neubauer BA, Fiedler B, Himmelein B, Kampfer F, Lassker U, Schwabe G, et al. Centrottemporal spikes in families with rolandic epilepsy: linkage to chromosome 15q14. *Neurology* 1998; 51: 1608–12.
- Nicolai J, Aldenkamp AP, Arends J, Weber JW, Vles JS. Cognitive and behavioral effects of nocturnal epileptiform discharges in children with benign childhood epilepsy with centrottemporal spikes. *Epilepsy Behav* 2006; 8: 56–70.
- Oguni H, Hayashi K, Imai K, Hirano Y, Mutoh A, Osawa M. Study on the early-onset variant of benign childhood epilepsy with occipital paroxysms otherwise described as early-onset benign occipital seizure susceptibility syndrome. *Epilepsia* 1999; 40: 1020–30.
- Ohtsu M, Oguni H, Hayashi K, Funatsuka M, Imai K, Osawa M. EEG in children with early-onset benign occipital seizure susceptibility syndrome: Panayiotopoulos syndrome. *Epilepsia* 2003; 44: 435–42.
- Ohtsu M, Oguni H, Imai K, Funatsuka M, Osawa M. Early-onset form of benign childhood epilepsy with centro-temporal EEG foci – a different nosological perspective from panayiotopoulos syndrome. *Neuropediatrics* 2008; 39: 14–9.
- Okanishi T, Maegaki Y, Ohno K, Togari H. Underlying neurologic disorders and recurrence rates of status epilepticus in childhood. *Brain Dev* 2008. Advance Access published on March 24, 2008, doi:10.1016/j.braindev.2008.02.003.
- Okubo Y, Matsuura M, Asai T, Asai K, Kato M, Kojima T, et al. Epileptiform EEG discharges in healthy children: prevalence, emotional and behavioral correlates, and genetic influences. *Epilepsia* 1994; 35: 832–41.
- Panayiotopoulos CP. Inhibitory effect of central vision on occipital lobe seizures. *Neurology* 1981; 31: 1330–3.
- Panayiotopoulos CP. Vomiting as an ictal manifestation of epileptic seizures and syndromes. *J Neurol Neurosurg Psychiatr* 1988; 51: 1448–51.
- Panayiotopoulos CP. Benign childhood partial epilepsies: benign childhood seizure susceptibility syndromes [editorial]. *J Neurol Neurosurg Psychiatr* 1993; 56: 2–5.
- Panayiotopoulos CP. Elementary visual hallucinations in migraine and epilepsy. *J Neurol Neurosurg Psychiatr* 1994; 57: 1371–4.
- Panayiotopoulos CP. Benign childhood partial seizures and related epileptic syndromes. London: John Libbey & Company Ltd; 1999.
- Panayiotopoulos CP. Elementary visual hallucinations, blindness, and headache in idiopathic occipital epilepsy: differentiation from migraine. *J Neurol Neurosurg Psychiatry* 1999; 66: 536–40.
- Panayiotopoulos CP. Visual phenomena and headache in occipital epilepsy: a review, a systematic study and differentiation from migraine. *Epileptic Disord* 1999; 1: 205–16.
- Panayiotopoulos CP. Panayiotopoulos syndrome: a common and benign childhood epileptic syndrome. London: John Libbey & Company; 2002.
- Panayiotopoulos CP. Autonomic seizures and autonomic status epilepticus peculiar to childhood: diagnosis and management. *Epilepsy Behav* 2004; 5: 286–95.
- Panayiotopoulos CP. A clinical guide to epileptic syndromes and their treatment. 2nd edn. London: Springer; 2007.
- Parisi P, Ferri R, Pagani J, Cecili M, Montemitro E, Villa MP. Ictal video-polysomnography and EEG spectral analysis in a child with severe Panayiotopoulos syndrome. *Epileptic Disord* 2005; 7: 333–9.
- Parmeggiani L, Guerrini R. Idiopathic partial epilepsy: electroclinical demonstration of a prolonged seizure with sequential rolandic and occipital involvement. Seizure spread due to regional susceptibility? *Epileptic Disord* 1999; 1: 35–40.
- Parmeggiani L, Seri S, Bonanni P, Guerrini R. Electrophysiological characterization of spontaneous and carbamazepine-induced epileptic negative myoclonus in benign childhood epilepsy with centro-temporal spikes. *Clin Neurophysiol* 2004; 115: 50–8.

- Penfield W, Rasmussen T. The cerebral cortex of man: A clinical study of localisation of function. 4th edn. New York: The Macmillan Company; 1957.
- Perkins FF Jr, Breier J, McManis MH, Castillo E, Wheless J, McGregor AL, et al. Benign rolandic epilepsy—perhaps not so benign: use of magnetic source imaging as a predictor of outcome. *J Child Neurol* 2008; 23: 389–93.
- Peters JM, Camfield CS, Camfield PR. Population study of benign rolandic epilepsy: is treatment needed? *Neurology* 2001; 57: 537–9.
- Petersen I, Eeg-Olofsson O. The development of the electroencephalogram in normal children from the age of 1 through 15 years. Non-paroxysmal activity. *Neuropadiatrie* 1971; 2: 247–304.
- Piccinelli P, Borgatti R, Aldini A, Bindelli D, Ferri M, Perna S, et al. Academic performance in children with rolandic epilepsy. *Dev Med Child Neurol* 2008; 50: 353–6.
- Plouin P, Anderson VE. Benign familial and non-familial neonatal seizures. In: Roger J, Bureau M, Dravet C, Genton P, Tassinari CA, Wolf P, editors. *Epileptic syndromes in infancy, childhood and adolescence*. 4th edn. Montrouge, France: John Libbey Eurotext Ltd; 2005. p. 3–15.
- Riva D, Vago C, Franceschetti S, Pantaleoni C, D'Arrigo S, Granata T, et al. Intellectual and language findings and their relationship to EEG characteristics in benign childhood epilepsy with centrotemporal spikes. *Epilepsy Behav* 2007; 10: 278–85.
- Russell MB, Olesen J. A nosographic analysis of the migraine aura in a general population. *Brain* 1996; 119: 355–61.
- Saito N, Kanazawa O, Tohyama J, Akasaka N, Kamimura T, Toyabe SI, et al. Brain maturation-related spike localization in Panayiotopoulos syndrome: magnetoencephalographic study. *Pediatr Neurol* 2008; 38: 104–10.
- Saitoh N, Kanazawa O, Toyama J, Akasaka N, Kamimura T. Magnetoencephalographic findings of Panayiotopoulos syndrome with frontal epileptic discharges. *Pediatric Neurol* 2007; 36: 190–4.
- Saltik S, Uluduz D, Cokar O, Demirbilek V, Dervent A. A clinical and EEG study on idiopathic partial epilepsies with evolution into ESES spectrum disorders. *Epilepsia* 2005; 46: 524–33.
- Sanders S, Rowlinson S, Koutroumanidis M, Ferrie CD, Panayiotopoulos CP. Midline spikes in children and clinical correlations. *Epilepsia* 2002; 43: 1436–9.
- Sanders S, Rowlinson S, Manidakis I, Ferrie CD, Koutroumanidis M. The contribution of the EEG technologists in the diagnosis of Panayiotopoulos syndrome (susceptibility to early onset benign childhood autonomic seizures). *Seizure* 2004; 13: 565–73.
- Schauble B, Britton JW, Mullan BP, Watson J, Sharbrough FW, Marsh WR. Ictal vomiting in association with left temporal lobe seizures in a left hemisphere language-dominant patient. *Epilepsia* 2002; 43: 1432–5.
- Scheffer IE, Harkin LA, Grinton BE, Dibbens LM, Turner SJ, Zielinski MA, et al. Temporal lobe epilepsy and GEFS+ phenotypes associated with SCN1B mutations. *Brain* 2007; 130: 100–9.
- Schott GD. Exploring the visual hallucinations of migraine aura: the tacit contribution of illustration. *Brain* 2007; 130: 1690–703.
- Shinnar S, Pellock JM. Update on the epidemiology and prognosis of pediatric epilepsy. *J Child Neurol* 2002; 17 (Suppl 1): S4–17.
- Sidenvall R, Forsgren L, Heijbel J. Prevalence and characteristics of epilepsy in children in northern Sweden. *Seizure* 1996; 5: 139–46.
- Smith JMB, Kellaway P. Central (Rolandic) foci in children: an analysis of 200 cases. *Electroencephalogr Clin Neurophysiol* 1964; 17: 460–1.
- Smith MC, Hoepfner TJ. Epileptic encephalopathy of late childhood: Landau-Kleffner syndrome and the syndrome of continuous spikes and waves during slow-wave sleep. *J Clin Neurophysiol* 2003; 20: 462–72.
- Sorel L, Rucquoy-Ponsar M. Functional epilepsy of maturation. Contribution of the EEG transverse montages in the diagnosis of this form of epilepsy. [French] *Revue Neurologique* 1969; 121: 289–97.
- Tassinari CA, Rubboli G, Volpi L, Billard C, Bureau M. Electrical status epilepticus during slow sleep (ESES or CSWS) including acquired epileptic aphasia (Landau-Kleffner syndrome). In: Roger J, Bureau M, Dravet C, Genton P, Tassinari CA, Wolf P, editors. *Epileptic syndromes in infancy, childhood and adolescence*. 4th edn. London: John Libbey & Co Ltd; 2005. p. 295–314.
- Tassinari CA, De Marco P. Benign partial epilepsy with extreme somatosensory evoked potentials. In: Roger J, Bureau M, Dravet C, Dreifuss FE, Wolf P, Perret A, editors. *Epileptic syndromes in infancy, childhood and adolescence*. 2nd edn. London: John Libbey; 1992. p. 225–9.
- Taylor I, Berkovic SF, Kivity S, Scheffer IE. Benign occipital epilepsies of childhood: clinical features and genetics. *Brain* 2008. Advance Access published on July 9, 2008; doi:10.1093/brain/awn138.
- Taylor I, Scheffer IE, Berkovic SF. Occipital epilepsies: identification of specific and newly recognized syndromes. *Brain* 2003; 126: 753–69.
- Tenembaum S, Deonna T, Fejerman N, Medina C, Ingvar-Maeder M, Gubser-Mercati D. Continuous spike-waves and dementia in childhood epilepsy with occipital paroxysms. *J Epilepsy* 1997; 10: 139–45.
- Thomas P, Arzimanoglou A, Aicardi J. Benign idiopathic occipital epilepsy: report of a case of the late (Gastaut) type. *Epileptic Disord* 2003; 5: 57–9.
- Tsai ML, Hung KL. Topographic mapping and clinical analysis of benign childhood epilepsy with centrotemporal spikes. *Brain Dev* 1998; 20: 27–32.
- Vadlamudi L, Harvey AS, Connellan MM, Milne RL, Hopper JL, Scheffer IE, et al. Is benign rolandic epilepsy genetically determined? *Ann Neurol* 2004; 56: 129–32.
- Vadlamudi L, Kjeldsen MJ, Corey LA, Solaas MH, Friis ML, Pellock JM, et al. Analyzing the etiology of benign rolandic epilepsy: a multicenter twin collaboration. *Epilepsia* 2006; 47: 550–5.
- Valeta T. Parental attitude, reaction and education in benign childhood focal seizures. In: Panayiotopoulos CP, editor. *The epilepsies: seizures, syndromes and management*. Oxford: Bladon Medical Publishing; 2005. p. 258–61.
- Valeta T. Impact of newly identified epileptic seizures in patients and family. In: Panayiotopoulos CP, editor. Vol. 3. *Newly identified epileptic seizures: diagnosis, procedures and management*. Oxford: Medicae; 2007. p. 138–44.
- Verrotti A, Coppola G, Manco R, Ciambra G, Iannetti P, Grosso S, et al. Levetiracetam monotherapy for children and adolescents with benign rolandic seizures. *Seizure* 2007; 16: 271–5.
- Verrotti A, Salladini C, Trotta D, di Corcia G, Chiarelli F. Ictal cardiorespiratory arrest in Panayiotopoulos syndrome. *Neurology* 2005; 64: 1816–7.
- Vigevano F, Lisi ML, Ricci S. Early onset benign occipital susceptibility syndrome: video-EEG documentation of an illustrative case. *Clin Neurophysiol* 2000; 111 (Suppl 2): S81–6.
- Vigevano F, Specchio N, Caraballo R, Watanabe K. Benign familial and nonfamilial seizures. In: Engel J Jr, Pedley TA, editors. Philadelphia: Lippincott Williams & Wilkins, A Wolters Kluwer Business; 2008. p. 2313–21.
- Watanabe K. Benign partial epilepsies. In: Wallace S, editor. *Epilepsy in children*. London: Chapman & Hall; 1996. p. 293–13.
- Wheless JW, Clarke DF, Carpenter D. Treatment of pediatric epilepsy: expert opinion, 2005. *J Child Neurol* 2005; 20 (Suppl 1): S1–56.
- Wheless JW, Clarke DF, Arzimanoglou A, Carpenter D. Treatment of pediatric epilepsy: European expert opinion, 2007. *Epileptic Disord* 2007; 9: 353–412.
- Wirrell E, Camfield CS, Camfield PR. Idiopathic and benign partial epilepsies of childhood. In: Wyllie E, Gupta A, Lachhwani D, editors. *The treatment of epilepsy*. 4th edn. Philadelphia: Lippincott Williams & Wilkins; 2006. p. 373–89.
- Wirrell E, Sherman EM, Vanmastrigt R, Hamiwka L. Deterioration in cognitive function in children with benign epilepsy of childhood with

- central temporal spikes treated with sulthiame. *J Child Neurol* 2008; 23: 14–21.
- Wirrell EC, Camfield PR, Gordon KE, Dooley JM, Camfield CS. Benign rolandic epilepsy: atypical features are very common. *J Child Neurol* 1995; 10: 455–8.
- Yalcin AD, Kaymaz A, Forta H. Childhood occipital epilepsy: seizure manifestations and electroencephalographic features. *Brain Dev* 1997; 19: 408–13.
- Yalcin AD, Kaymaz A, Forta H. Reflex occipital lobe epilepsy. *Seizure* 2000; 9: 436–41.
- Yoshinaga H, Koutroumanidis M, Kobayashi K, Shirasawa A, Kikumoto K, Inoue T, *et al.* EEG dipole characteristics in Panayiotopoulos syndrome. *Epilepsia* 2006; 47: 781–7.