

Besov and Triebel-Lizorkin Spaces Related to Singular Integrals with Flag Kernels

Dachun YANG

School of Mathematical Sciences
Beijing Normal University
Laboratory of Mathematics and Complex Systems
Ministry of Education
Beijing 100875 — People's Republic of China
dcyang@bnu.edu.cn

Received: August 11, 2008

Accepted: January 8, 2009

ABSTRACT

Let $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$. In this paper, the author introduces the Besov space $\dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ and the Triebel-Lizorkin space $\dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ associated to singular integrals with flag kernels. Some basic properties, including their dual spaces, some equivalent norm characterizations via Littlewood-Paley functions, lifting properties and some embedding theorems, on these spaces are given. Moreover, the author obtains the boundedness of flag singular integrals and fractional integrals on these spaces.

Key words: Besov space, Triebel-Lizorkin space, flag singular integral, flag fractional integral, Littlewood-Paley operator, dual space, lifting, embedding.

2000 Mathematics Subject Classification: 42B35, 42B20, 42B25, 46E35.

1. Introduction

In order to study the \square_b -complex on certain quadratic CR submanifolds of \mathbb{C}^n , Nagel, Ricci, and Stein [6] introduced the notion of singular integrals with flag kernels on \mathbb{R}^n . Since the flag singular integral is a special case of product singular integrals, the boundedness of flag singular integrals on $L^p(\mathbb{R}^n)$ with $p \in (1, \infty)$ is a simple corollary of the boundedness of the corresponding product singular integrals. Recently,

The author is supported by National Science Foundation for Distinguished Young Scholars (No. 10425106) of China.

Han and Lu in [3] developed a corresponding Hardy space theory for flag singular integrals on \mathbb{R}^n .

Motivated by [3, 6, 7], letting $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$, in this paper, we introduce the Besov space $B_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ and the Triebel-Lizorkin space $F_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ associated to singular integrals with flag kernels. Some basic properties, including their dual spaces, some equivalent norm characterizations via Littlewood-Paley functions, lifting properties, and some embedding theorems on these spaces are given. Moreover, we obtain the boundedness of flag singular integrals and fractional integrals on these spaces.

For the simplicity of presentation, we work on flag integrals on \mathbb{R}^2 . However, our method works for more general product Euclidean spaces.

It was proved in [5, 6] that flag kernels on \mathbb{R}^2 are closely connected with product kernels on $\mathbb{R}^2 \times \mathbb{R}$. We denote any element of $\mathbb{R}^2 \times \mathbb{R}$ by the 3-tuple $x = (x_1, x_2, x_3)$, where $(x_1, x_2) \in \mathbb{R}^2$ and $x_3 \in \mathbb{R}$. We endow \mathbb{R}^2 with the following dilation that for any $\delta > 0$ and $x = (x_1, x_2) \in \mathbb{R}^2$, $\delta x = (\delta x_1, \delta^2 x_2)$ and the norm that $\|x\| = (x_1^2 + |x_2|)^{1/2}$, which is equivalent to $|x_1| + |x_2|^{1/2}$. Obviously, the homogeneous dimension of \mathbb{R}^2 is 3.

In order to express the cancellation conditions introduced by Nagel, Ricci, and Stein in [6], we introduce the following terminology. A k -normalized bump function on \mathbb{R}^n is a C^k -function supported on the unit ball with C^k -norm bounded by 1. It was proved by Nagel, Ricci, and Stein in [6, Remark 2.1.7] that Definitions 1.1 and 1.2 given below are essentially independent of the choice of $k \in \mathbb{N}$. Hence we usually speak of normalized bump functions rather than k -normalized bump functions.

In what follows, we denote by C a positive constant which is independent of the main parameters, but it may vary from line to line. Constants with subscripts, such as C_1 , do not change in different occurrences. We also use subscripts to indicate which parameters the constant depends on. Moreover, let $\mathbb{N} \equiv \{1, 2, \dots\}$ and $\mathbb{Z}_+ \equiv \mathbb{N} \cup \{0\}$.

Definition 1.1. A product kernel on $\mathbb{R}^2 \times \mathbb{R}$, relative to the given dilations, is a distribution K on $\mathbb{R}^2 \times \mathbb{R}$ which coincides with a C^∞ function away from the coordinate subspace $x_j = 0$ for $j = 1, 2, 3$ and which satisfies:

- (i) (Differential inequalities) For each multi-index $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in (\mathbb{Z}_+)^3$, there exists a positive constant C_α so that

$$|\partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \partial_{x_3}^{\alpha_3} K(x_1, x_2, x_3)| \leq C_\alpha \| (x_1, x_2) \|^{-3-\alpha_1-2\alpha_2} |x_3|^{-1-\alpha_3}$$

away from the coordinate subspaces, where $\partial_{x_i}^{\alpha_i} = \frac{\partial^{\alpha_i}}{\partial x_i^{\alpha_i}}$.

- (ii) (Cancellation conditions)

- (a) For each multi-index $(\alpha_1, \alpha_2) \in (\mathbb{Z}_+)^2$ and any given normalized bump function φ on \mathbb{R} and any $\delta > 0$, there exists a positive constant C_{α_1, α_2}

so that, for all $(x_1, x_2) \neq (0, 0)$,

$$\left| \int_{\mathbb{R}} \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} K(x_1, x_2, x_3) \varphi(\delta x_3) dx_3 \right| \leq C_{\alpha_1, \alpha_2} \|(x_1, x_2)\|^{-3-\alpha_1-2\alpha_2}.$$

- (b) For each $\alpha_3 \in \mathbb{Z}_+$ and any given normalized bump function φ on \mathbb{R}^2 and any $\delta > 0$, there exists a positive constant C_{α_3} so that, for all $x_3 \neq 0$,

$$\left| \int_{\mathbb{R}^2} \partial_{x_3}^{\alpha_3} K(x_1, x_2, x_3) \varphi(\delta x_1, \delta^2 x_2) dx_1 dx_2 \right| \leq C_{\alpha_3} |x_3|^{-1-\alpha_3}.$$

- (c) For any given normalized bump function φ on $\mathbb{R}^2 \times \mathbb{R}$ and any $\delta_1, \delta_2 > 0$, there exists a positive constant C so that

$$\left| \int_{\mathbb{R}^2 \times \mathbb{R}} K(x_1, x_2, x_3) \varphi(\delta_1 x_1, \delta_1^2 x_2, \delta_2 x_3) dx_1 dx_2 dx_3 \right| \leq C.$$

The following definition of flag kernels is just a special case of flag kernels in [6].

Definition 1.2. A flag kernel on \mathbb{R}^2 , relative to the given dilations, is a distribution K on \mathbb{R}^2 which coincides with a C^∞ function away from the coordinate subspace $x_1 = 0$ and which satisfies:

- (i) (Differential inequalities) For each $\alpha = (\alpha_1, \alpha_2) \in (\mathbb{Z}_+)^2$, there exists a positive constant C_α so that, for all $x_1 \neq 0$,

$$|\partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} K(x_1, x_2)| \leq C_\alpha |x_1|^{-1-\alpha_1} \|(x_1, x_2)\|^{-2-2\alpha_2}.$$

- (ii) (Cancellation conditions)

- (a) For any given normalized bump function φ on \mathbb{R} , any $\alpha_1 \in \mathbb{Z}_+$, and any $\delta > 0$, there exists a positive constant C_{α_1} so that, for all $x_1 \neq 0$,

$$\left| \int_{\mathbb{R}} \partial_{x_1}^{\alpha_1} K(x_1, x_2) \varphi(\delta x_2) dx_2 \right| \leq C_{\alpha_1} |x_1|^{-1-\alpha_1}.$$

- (b) For any given normalized bump function φ on \mathbb{R} , any $\alpha_2 \in \mathbb{Z}_+$, and any $\delta > 0$, there exists a positive constant C_{α_2} so that for all $x_2 \neq 0$,

$$\left| \int_{\mathbb{R}} \partial_{x_2}^{\alpha_2} K(x_1, x_2) \varphi(\delta x_1) dx_1 \right| \leq C_{\alpha_2} |x_2|^{-1-\alpha_2}.$$

- (c) For any given normalized bump function φ on \mathbb{R}^2 , and any $\delta_1, \delta_2 > 0$, there exists a positive constant C so that

$$\left| \int_{\mathbb{R}^2} K(x_1, x_2) \varphi(\delta_1 x_1, \delta_2 x_2) dx_1 dx_2 \right| \leq C.$$

Remark 1.3. It was pointed by Nagel, Ricci, and Stein in [6] that the single normalized bump function in Definitions 1.1 and 1.2 (ii)-(c) can be replaced by the tensor product of normalized bump functions on \mathbb{R}^2 and \mathbb{R} .

The following proposition is completely similar to Proposition 3.2 and Lemma 4.5 in [5], which reveals the relation between the product kernel and the flag kernel.

Proposition 1.4. *Let K^\sharp be an integrable function on $\mathbb{R}^2 \times \mathbb{R}$ which is a product kernel as in Definition 1.1. Then, for $(x_1, x_2) \in \mathbb{R}^2$,*

$$K(x_1, x_2) = \int_{\mathbb{R}} K^\sharp(x_1, x_2 - x_3, x_3) dx_3 \tag{1}$$

is a flag singular kernel on \mathbb{R}^2 .

Conversely, given $K \in L^1(\mathbb{R}^2)$ which is a flag kernel as in Definition 1.2, then

$$K^\sharp(x_1, x_2, x_3) = \frac{1}{|x_1|^2} \chi\left(\frac{x_2}{|x_1|^2}\right) K(x_1, x_2 + x_3),$$

where χ is a non-negative smooth function supported on $[1/2, 1]$ such that $\int_{1/2}^1 \chi(t) dt = 1$, is an integrable product kernel on $\mathbb{R}^2 \times \mathbb{R}$ such that (1) holds.

The organization of this paper is as follows. In section 2, we first establish some Calderón reproducing formulae (see Lemma 2.3), whose dyadic version (see Lemma 2.4) are essentially included in [3]. However, in this paper, we use a slightly different way from [3] to define the topology of $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$; see Definition 2.1 below and Definition 1.6 in [3]. Let $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$. With a special choice of approximations of the identity associated to flag kernels (see (1.3) of [3]), we then introduce the norms of $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ in Definition 2.5, and using the Calderón reproducing formulae, we prove in Theorem 2.6 that these norms are independent of the choice of the approximations of the identity associated to flag kernels. Then we introduce the Besov space $\dot{B}_{pq}^s(\mathbb{R}^2)$ and the Triebel-Lizorkin space $\dot{F}_{pq}^s(\mathbb{R}^2)$ in Definition 2.7. Some basic properties including dual spaces of these spaces are presented in Propositions 2.9, 2.10, and 2.11. In Theorems 2.8, 2.14 and Corollary 2.22, we establish some equivalent norm characterizations of these spaces including various Littlewood-Paley functions. We remark that Corollary 2.22 clearly reveals the difference between $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ with the classical product Besov and Triebel-Lizorkin spaces in [7].

The boundedness of flag singular integrals on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ is given in Theorem 3.1 and the lifting properties of these spaces via Riesz potential operators is presented in Theorem 4.6.

Finally, in Theorems 5.1 and 5.2, we establish some embedding theorems on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$. The boundedness of flag fractional integrals is given in Theorem 5.4.

2. Besov and Triebel-Lizorkin spaces on \mathbb{R}^2

We first introduce the Calderón reproducing formulae. To this end, we need to introduce some spaces of distributions; see [3].

Definition 2.1. A Schwartz function $f \in \mathcal{S}(\mathbb{R}^2)$ is said to belong to the space of test functions, $\mathcal{S}_{\infty,F}(\mathbb{R}^2)$, related to flag singular integrals, if there exists a Schwartz function $f^\# \in \mathcal{S}(\mathbb{R}^2 \times \mathbb{R})$ such that, for all $(x_1, x_2) \in \mathbb{R}^2$,

$$f(x_1, x_2) = \int_{\mathbb{R}} f^\#(x_1, x_2 - x_3, x_3) dx_3,$$

where $f^\#$ satisfies the following conditions: for all $x_3 \in \mathbb{R}$ and $(\alpha_1, \alpha_2) \in (\mathbb{Z}_+)^2$,

$$\int_{\mathbb{R}^2} f^\#(x_1, x_2, x_3) x_1^{\alpha_1} x_2^{\alpha_2} dx_1 dx_2 = 0,$$

and for all $(x_1, x_2) \in \mathbb{R}^2$ and $\alpha_3 \in \mathbb{Z}_+$,

$$\int_{\mathbb{R}} f^\#(x_1, x_2, x_3) x_3^{\alpha_3} dx_3 = 0.$$

We endow $\mathcal{S}_{\infty,F}(\mathbb{R}^2)$ with the same topology as $\mathcal{S}(\mathbb{R}^2)$, and we denote its dual space by $\mathcal{S}_{\infty,F}(\mathbb{R}^2)'$.

Remark 2.2. It is easy to see that the space $\mathcal{S}_{\infty,F}(\mathbb{R}^2)$ is a closed subspace of $\mathcal{S}(\mathbb{R}^2)$, and if $f \in \mathcal{S}_{\infty,F}(\mathbb{R}^2)$ then, for all $\alpha_2 \in \mathbb{Z}_+$,

$$\int_{\mathbb{R}} f(x_1, x_2) x_2^{\alpha_2} dx_2 = 0.$$

Let $\mathcal{P}_{x_2}(\mathbb{R})$ be the set of all polynomials on \mathbb{R} in variable x_2 . Then, one can easily see that if $f \in \mathcal{S}_{\infty,F}(\mathbb{R}^2)'$, $h \in \mathcal{P}_{x_2}(\mathbb{R})$, and $g \in \mathcal{S}_{\infty,F}(\mathbb{R}^2)$, then $\langle f, g \rangle = \langle f, g + h \rangle$, namely, $\mathcal{S}_{\infty}(\mathbb{R}^2)' / \mathcal{P}_{x_2}(\mathbb{R}) \subset \mathcal{S}_{\infty,F}(\mathbb{R}^2)'$.

We now establish the following Calderón reproducing formulae by a method essentially similar to that of Theorem 3 in the appendix of [2] and a dual argument; see also [3].

Lemma 2.3. Let $\psi^{(1)} \in \mathcal{S}(\mathbb{R}^2)$ with $\int_{\mathbb{R}^2} \psi^{(1)}(x_1, x_2) dx_1 dx_2 = 0$ and $\psi^{(2)} \in \mathcal{S}(\mathbb{R})$ with $\int_{\mathbb{R}} \psi^{(2)}(x_3) dx_3 = 0$ satisfy the following admissible conditions: for all $(\xi_1, \xi_2) \in \mathbb{R}^2$ and $(\xi_1, \xi_2) \neq 0$,

$$\int_0^\infty |\hat{\psi}^{(1)}(t\xi_1, t^2\xi_2)|^2 \frac{dt}{t} = 1,$$

and, for all $\eta \in \mathbb{R}$ and $\eta \neq 0$,

$$\int_0^\infty |\hat{\psi}^{(2)}(t\eta)|^2 \frac{dt}{t} = 1.$$

For $t_1, t_2 > 0$ and $x_1, x_2 \in \mathbb{R}$, let

$$\psi_{t_1}^{(1)}(x_1, x_2) = \frac{1}{t_1^3} \psi^{(1)}\left(\frac{x_1}{t_1}, \frac{x_2}{t_1^2}\right), \quad \psi_{t_2}^{(2)}(x_2) = \frac{1}{t_2} \psi^{(2)}\left(\frac{x_2}{t_2}\right),$$

and

$$\psi_{t_1 t_2}(x_1, x_2) = \int_{\mathbb{R}} \psi_{t_1}^{(1)}(x_1, x_2 - x'_2) \psi_{t_2}^{(2)}(x'_2) dx'_2.$$

Then the identity

$$f(x_1, x_2) = \int_0^\infty \int_0^\infty \psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2} \tag{2}$$

holds in $L^2(\mathbb{R}^2)$, $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$, and $\mathcal{S}_{\infty, F}(\mathbb{R}^2)'$.

Proof. From the Plancherel principle and the assumptions of the lemma, it is easy to see that (2) holds in $L^2(\mathbb{R}^2)$. The fact that (2) holds in $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$ and a dual argument show that (2) also holds in $\mathcal{S}_{\infty, F}(\mathbb{R}^2)'$. Thus, to finish the proof of Lemma 2.3, we only need to show that (2) holds in $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$. To do so, for $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)$, $\epsilon_i > 0$, and $\delta_i > 0$ with $i = 1, 2$ and $(x_1, x_2) \in \mathbb{R}^2$, set

$$f_{\epsilon_1, \epsilon_2, \delta_1, \delta_2}(x_1, x_2) = \int_{\epsilon_1}^{\delta_1} \int_{\epsilon_2}^{\delta_2} \psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2}.$$

We only need to show that for any fixed $N \in \mathbb{Z}_+$ and all $(x_1, x_2) \in \mathbb{R}^2$, there exists a positive constant $C = C_{f, \psi^{(1)}, \psi^{(2)}, N}$ such that

$$|f(x_1, x_2) - f_{\epsilon_1, \epsilon_2, \delta_1, \delta_2}(x_1, x_2)| \leq C \left(\epsilon_1 + \frac{1}{\delta_1}\right) \left(\epsilon_2 + \frac{1}{\delta_2}\right) (1 + \|(x_1, x_2)\|)^{-N}. \tag{3}$$

Obviously, we may assume that $0 < \epsilon_i < 1 < \delta_i$ for $i = 1, 2$ in (3). To prove (3), we write

$$\begin{aligned} f(x_1, x_2) - f_{\epsilon_1, \epsilon_2, \delta_1, \delta_2}(x_1, x_2) &= \int_0^{\epsilon_1} \int_0^\infty \psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ &\quad + \int_{\epsilon_1}^{\delta_1} \int_0^{\epsilon_2} \dots + \int_{\epsilon_1}^{\delta_1} \int_{\delta_2}^\infty \dots + \int_{\delta_1}^\infty \int_0^\infty \dots \\ &= H_1 + H_2 + H_3. \end{aligned}$$

We only estimate H_1 . The same technique also works for H_2 and H_3 . To estimate H_1 , we further decompose it into

$$H_1 = \int_0^{\epsilon_1} \int_0^1 \psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2} + \int_0^{\epsilon_1} \int_1^{\infty} \dots = H_1^1 + H_1^2.$$

Let $\varphi^{(i)} = \psi^{(i)} * \psi^{(i)}$ for $i = 1, 2$ and $\varphi_{t_1 t_2} = \varphi_{t_1}^{(1)} * \varphi_{t_2}^{(2)}$. Then it is easy to see that $\varphi_{t_1 t_2} = \psi_{t_1 t_2} * \psi_{t_1 t_2}$. Since $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)$, by Definition 2.1, there exists a function f^\sharp as in Definition 2.1 such that, for all $(x_1, x_2) \in \mathbb{R}^2$,

$$f(x_1, x_2) = \int_{\mathbb{R}} f^\sharp(x_1, x_2 - x_3, x_3) dx_3.$$

Using this fact, we have

$$\begin{aligned} & \psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^2} \int_{\mathbb{R}} \varphi_{t_1}^{(1)}(x_1 - y_1, x_2 - y_2 - z) \varphi_{t_2}^{(2)}(z - y_3) f^\sharp(y_1, y_2, y_3) dz dy_1 dy_2 dy_3. \end{aligned}$$

By the vanishing moments of $\varphi^{(1)}$ and $\varphi^{(2)}$, we further write

$$\begin{aligned} & \psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \\ &= \int_{\mathbb{R}} \int_{|z-y_3| \leq |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| \leq \|(x_1, x_2-z)\|/2} \varphi_{t_1}^{(1)}(x_1 - y_1, x_2 - y_2 - z) \\ & \quad \times \varphi_{t_2}^{(2)}(z - y_3) \left\{ [f^\sharp(y_1, y_2, y_3) - f^\sharp(x_1, x_2 - z, y_3)] \right. \\ & \quad \left. - [f^\sharp(y_1, y_2, z) - f^\sharp(x_1, x_2 - z, z)] \right\} dy_1 dy_2 dy_3 dz \\ & + \int_{\mathbb{R}} \int_{|z-y_3| > |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| \leq \|(x_1, x_2-z)\|/2} \dots \\ & + \int_{\mathbb{R}} \int_{|z-y_3| \leq |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| > \|(x_1, x_2-z)\|/2} \dots \\ & + \int_{\mathbb{R}} \int_{|z-y_3| > |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| > \|(x_1, x_2-z)\|/2} \dots = I_1 + I_2 + I_3 + I_4. \end{aligned}$$

We denote the corresponding terms of H_1^1 to I_i , respectively, by $H_{1,i}^1$, where $i = 1, 2, 3, 4$, and by similarity we only estimate $H_{1,1}^1$ and $H_{1,2}^1$. By the mean value

theorem, we have

$$\begin{aligned}
 |I_1| &\leq C \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{\mathbb{R}^2} \left| \varphi_{t_1}^{(1)}(x_1 - y_1, x_2 - y_2 - z) \right| \\
 &\quad \times \left[\frac{|x_1 - y_1|}{(1 + \|(x_1, x_2 - z)\|)^{N+2}} + \frac{|x_2 - y_2 - z|^{1/2}}{(1 + \|(x_1, x_2 - z)\|)^{N+2}} \right] \\
 &\quad \times |\varphi_{t_2}^{(2)}(z - y_3)| |z - y_3| \frac{1}{(1 + |z|)^N} dy_1 dy_2 dy_3 dz \\
 &\leq Ct_1 t_2 \int_{\mathbb{R}} \frac{1}{(1 + \|(x_1, x_2 - z)\|)^{N+2}} \frac{1}{(1 + |z|)^N} dz \leq Ct_1 t_2 \frac{1}{(1 + \|(x_1, x_2)\|)^N}.
 \end{aligned}$$

From this, we can easily deduce a desired estimate for $H_{1,1}^1$. Similarly, we have

$$\begin{aligned}
 |I_2| &\leq Ct_1 \int_{\mathbb{R}} \int_{|z-y_3|>|z|/2} |\varphi_{t_2}^{(2)}(z - y_3)| |y_3 - z| \\
 &\quad \times \frac{1}{(1 + \|(x_1, x_2 - z)\|)^{N+2}} dy_3 dz \\
 &\leq Ct_1 t_2 \int_{\mathbb{R}} \frac{1}{(1 + \|(x_1, x_2 - z)\|)^{N+2}} \frac{1}{(1 + |z|)^N} dz \\
 &\leq Ct_1 t_2 \frac{1}{(1 + \|(x_1, x_2)\|)^N},
 \end{aligned}$$

which gives a desired estimate of $H_{1,2}^1$.

In what follows, we denote by $[a]$ the integer no more than $a \in \mathbb{R}$. To estimate H_1^2 , by the vanishing moments of $\varphi^{(1)}$ and f^\sharp , we write

$$\begin{aligned}
 &\psi_{t_1 t_2} * \psi_{t_1 t_2} * f(x_1, x_2) \\
 &= \int_{\mathbb{R}} \int_{|y_3| \leq |z|/2} \int_{\|(x_1 - y_1, x_2 - y_2 - z)\| \leq \|(x_1, x_2 - z)\|/2} \varphi_{t_1}^{(1)}(x_1 - y_1, x_2 - y_2 - z) \\
 &\quad \times \left[\varphi_{t_2}^{(2)}(z - y_3) - \sum_{\gamma=0}^{[N/2]} (-y_3)^\gamma \frac{d^\gamma}{dz^\gamma} \varphi_{t_2}^{(2)}(z) \right] \\
 &\quad \times [f^\sharp(y_1, y_2, y_3) - f^\sharp(x_1, x_2 - z, y_3)] dy_1 dy_2 dy_3 dz \\
 &+ \int_{\mathbb{R}} \int_{|y_3| > |z|/2} \int_{\|(x_1 - y_1, x_2 - y_2 - z)\| \leq \|(x_1, x_2 - z)\|/2} \dots \\
 &+ \int_{\mathbb{R}} \int_{|y_3| \leq |z|/2} \int_{\|(x_1 - y_1, x_2 - y_2 - z)\| > \|(x_1, x_2 - z)\|/2} \dots \\
 &+ \int_{\mathbb{R}} \int_{|y_3| > |z|/2} \int_{\|(x_1 - y_1, x_2 - y_2 - z)\| > \|(x_1, x_2 - z)\|/2} \dots = J_1 + J_2 + J_3 + J_4,
 \end{aligned}$$

and we also denote the corresponding terms of H_1^2 to J_i , respectively, by $H_{1,i}^2$, where $i = 1, 2, 3, 4$. By similarity, we only estimate $H_{1,1}^2$. The Taylor expansion theorem yields

$$\begin{aligned} |J_1| &\leq C \int_{\mathbb{R}} \frac{t_1}{(1 + \|(x_1, x_2 - z)\|)^{N+2}} \frac{1}{t_2} \frac{1}{(t_2 + |z|)^{\lfloor N/2 \rfloor + 1}} dz \\ &\leq C \frac{t_1}{(1 + \|(x_1, x_2)\|)^{N+2}} \frac{1}{t_2^{\lfloor N/2 \rfloor + 1}} + C \frac{t_1}{(1 + |x_1|)^N} \frac{1}{t_2} \frac{1}{(t_2 + |x_2|)^{\lfloor N/2 \rfloor + 1}}, \end{aligned}$$

which yields a desired estimate for $H_{1,1}^2$. This finishes the proof of Lemma 2.3. \square

Similarly, we have a ‘discrete’ version of Lemma 2.3 and we omit the details of its proof; see [3].

Lemma 2.4. *Let $\psi^{(1)} \in \mathcal{S}(\mathbb{R}^2)$ with $\int_{\mathbb{R}^2} \psi^{(1)}(x_1, x_2) dx_1 dx_2 = 0$ and $\psi^{(2)} \in \mathcal{S}(\mathbb{R})$ with $\int_{\mathbb{R}} \psi^{(2)}(x_3) dx_3 = 0$ satisfy the following admissible conditions that for all $(\xi_1, \xi_2) \in \mathbb{R}^2$ and $(\xi_1, \xi_2) \neq 0$,*

$$\sum_{k_1=-\infty}^{\infty} |\hat{\psi}^{(1)}(2^{-k_1} \xi_1, 2^{-2k_1} \xi_2)|^2 = 1,$$

and for all $\eta \in \mathbb{R}$ and $\eta \neq 0$, $\sum_{k_2=-\infty}^{\infty} |\hat{\psi}^{(2)}(2^{-k_2} \eta)|^2 = 1$. For all $k_1, k_2 \in \mathbb{Z}$ and $x_1, x_2 \in \mathbb{R}$, let $\psi_{k_1}^{(1)}(x_1, x_2) = 2^{3k_1} \psi^{(1)}(2^{k_1} x_1, 2^{2k_1} x_2)$, $\psi_{k_2}^{(2)}(x_2) = 2^{k_2} \psi^{(2)}(2^{k_2} x_2)$, and

$$\psi_{k_1 k_2}(x_1, x_2) = \int_{\mathbb{R}} \psi_{k_1}^{(1)}(x_1, x_2 - x'_2) \psi_{k_2}^{(2)}(x'_2) dx'_2.$$

Then the identity

$$f(x_1, x_2) = \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} \psi_{k_1 k_2} * \psi_{k_1 k_2} * f(x_1, x_2) \tag{4}$$

holds in $L^2(\mathbb{R}^2)$, $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$, and $\mathcal{S}_{\infty, F}(\mathbb{R}^2)'$.

We now introduce the norms $\|\cdot\|_{\dot{B}_{p,q}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{p,q}^s(\mathbb{R}^2)}$ for $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and using Lemma 2.3, we prove that they are independent of choices of $\psi^{(1)}$ and $\psi^{(2)}$.

Definition 2.5. Let $\psi_{t_1 t_2}$ be the same as in Lemma 2.3 and $s_1, s_2 \in \mathbb{R}$. For $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$, define

$$\|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^2)} = \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1 q} t_2^{-s_2 q} \|\psi_{t_1 t_2} * f\|_{L^p(\mathbb{R}^2)}^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q}$$

for $p, q \in [1, \infty]$, and

$$\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} = \left\| \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1q} t_2^{-s_2q} |\psi_{t_1 t_2} * f|^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)}$$

for $p \in (1, \infty)$ and $q \in (1, \infty]$, where the usual modifications are made when $p = \infty$ or $q = \infty$.

We recall the definition of the strong maximal function: for any $f \in L^1_{\text{loc}}(\mathbb{R}^2)$ and all $(x_1, x_2) \in \mathbb{R}^2$,

$$M_s(f)(x_1, x_2) = \sup_{\substack{(x_1, x_2) \in R \\ R \text{ rectangle}}} \frac{1}{|R|} \int_R |f(y_1, y_2)| dy_1 dy_2.$$

Theorem 2.6. *Let $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$. The norm $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ with $p, q \in [1, \infty]$ and the norm $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ are independent of the choices of $\psi^{(i)}$ for $i = 1, 2$.*

Proof. Let $\tilde{\psi}^{(i)}$ be some other functions satisfying the same conditions as $\psi^{(i)}$ for $i = 1, 2$. We denote the corresponding norms, respectively, by $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ and now prove that there exists a positive constant C such that, for all $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$,

$$\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \tag{5}$$

and

$$\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}. \tag{6}$$

To prove (5) and (6), by Lemma 2.3, we first need to establish a desired estimate for $\tilde{\psi}_{u_1 u_2} * \psi_{t_1 t_2}$. By its definition, it is easy to show that, for all $(x_1, x_2) \in \mathbb{R}^2$,

$$\tilde{\psi}_{u_1 u_2} * \psi_{t_1 t_2}(x_1, x_2) = (\tilde{\psi}_{u_1}^{(1)} * \psi_{t_1}^{(1)}) *_2 (\tilde{\psi}_{u_2}^{(2)} * \psi_{t_2}^{(2)})(x_1, x_2),$$

where, and in what follows, $*_2$ denotes the convolution in the second variable. We also set $a \wedge b = \min\{a, b\}$ and $a \vee b = \max\{a, b\}$ for $a, b \in \mathbb{R}$. We claim that

(i) for all $t_1, u_1 > 0$ and all $(x_1, x_2) \in \mathbb{R}^2$,

$$|(\tilde{\psi}_{u_1}^{(1)} * \psi_{t_1}^{(1)})(x_1, x_2)| \leq C \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(x_1, x_2)\|)^4}; \tag{7}$$

(ii) for all $t_2, u_2 > 0$ and all $z \in \mathbb{R}$,

$$|(\tilde{\psi}_{u_2}^{(2)} * \psi_{t_2}^{(2)})(z)| \leq C \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |z|)^2}. \tag{8}$$

By similarity, we only show (7). To this end, by the mean value theorem and some trivial computation, we can easily prove that, for all $u_1 > 0$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$|\tilde{\psi}_{u_1}^{(1)}(x_1, x_2)| \leq C \frac{u_1}{(u_1 + \|(x_1, x_2)\|)^4}, \tag{9}$$

and, for all $u_1 > 0$ and $(x_1, x_2), (y_1, y_2) \in \mathbb{R}^2$ with $\|(y_1, y_2)\| \leq (u_1 + \|(x_1, x_2)\|)/2$,

$$|\tilde{\psi}_{u_1}^{(1)}(x_1 + y_1, x_2 + y_2) - \tilde{\psi}_{u_1}^{(1)}(x_1, x_2)| \leq C \frac{\|(y_1, y_2)\|}{u_1 + \|(x_1, x_2)\|} \frac{u_1}{(u_1 + \|(x_1, x_2)\|)^4}. \tag{10}$$

The estimates (9) and (10), with $\tilde{\psi}_{u_1}^{(1)}$ and u_1 replaced respectively by $\psi_{t_1}^{(1)}$ and t_1 , also hold. We now prove (7) in the case $u_1 \geq t_1$. In this case, as

$$\int_{\mathbb{R}^2} \psi_{t_1}^{(1)}(y_1, y_2) dy_1 dy_2 = 0,$$

we can write

$$\begin{aligned} & (\tilde{\psi}_{u_1}^{(1)} * \psi_{t_1}^{(1)})(x_1, x_2) \\ &= \int_{\|(y_1, y_2)\| \leq (u_1 + \|(x_1, x_2)\|)/2} [\tilde{\psi}_{u_1}^{(1)}(x_1 - y_1, x_2 - y_2) - \tilde{\psi}_{u_1}^{(1)}(x_1, x_2)] \\ & \quad \times \psi_{t_1}^{(1)}(y_1, y_2) dy_1 dy_2 \\ & \quad + \int_{\|(y_1, y_2)\| > (u_1 + \|(x_1, x_2)\|)/2} \dots \\ &= D_1 + D_2. \end{aligned}$$

The estimate (10) yields that

$$\begin{aligned} |D_1| &\leq C \frac{t_1}{(u_1 + \|(x_1, x_2)\|)^4} \int_{\mathbb{R}^2} \left\| \left(\frac{y_1}{t_1}, \frac{y_2}{t_1} \right) \right\| |\psi_{t_1}^{(1)}(y_1, y_2)| dy_1 dy_2 \\ &\leq C \frac{t_1}{(u_1 + \|(x_1, x_2)\|)^4}, \end{aligned}$$

and the estimates (9) for $\tilde{\psi}_{u_1}^{(1)}$ and $\psi_{t_1}^{(1)}$ imply that

$$\begin{aligned} |D_2| &\leq C \int_{\|(y_1, y_2)\| > (u_1 + \|(x_1, x_2)\|)/2} |\tilde{\psi}_{u_1}^{(1)}(x_1 - y_1, x_2 - y_2)| \frac{t_1}{\|(y_1, y_2)\|^4} dy_1 dy_2 \\ & \quad + C \frac{u_1}{(u_1 + \|(x_1, x_2)\|)^4} \int_{\|(y_1, y_2)\| > u_1/2} \frac{t_1}{\|(y_1, y_2)\|^4} dy_1 dy_2 \\ &\leq C \frac{t_1}{(u_1 + \|(x_1, x_2)\|)^4}, \end{aligned}$$

which proves (7).

Let $t, s > 0$ and $(x_1, x_2) \in \mathbb{R}^2$. We now estimate

$$\begin{aligned} & \int_{\mathbb{R}} \frac{t}{(t + \|(x_1, x_2) - (0, y)\|)^4} \frac{s}{(s + |y|)^2} dy \\ & \leq \int_{|y| \leq |x_2|/2} \frac{t}{(t^2 + |x_1|^2 + |x_2 - y|)^2} \frac{s}{(s + |y|)^2} dy \\ & \quad + \int_{|x_2|/2 < |y| \leq 2|x_2|} \dots + \int_{|y| > 2|x_2|} \dots = E_1 + E_2 + E_3. \end{aligned}$$

For E_1 , in this case, we have that $|x_2 - y| \geq |x_2|/2$ and

$$E_1 \leq \frac{t}{(t^2 + |x_1|^2 + |x_2|/2)^2} \int_{\mathbb{R}} \frac{s}{(s + |y|)^2} dy \leq C \frac{t}{(t + \|(x_1, x_2)\|)^4};$$

for E_3 , by the fact that

$$t^2 + |x_1|^2 + |x_2 - y| \geq t^2 + |x_1|^2 + |x_2| \geq (t + \|(x_1, x_2)\|)^2/2,$$

we also obtain an estimate similar to E_1 . For E_2 , we have

$$E_2 \leq C \frac{t}{t^2 + |x_1|^2} \frac{s}{(s + |x_2|)^2} \int_{\mathbb{R}} \frac{1}{(1 + |y|)^2} dy \leq C \frac{t}{(t + |x_1|)^2} \frac{s}{(s + |x_2|)^2}.$$

Thus, for all $t, s > 0$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} & \int_{\mathbb{R}} \frac{t}{(t + \|(x_1, x_2) - (0, y)\|)^4} \frac{s}{(s + |y|)^2} dy \\ & \leq C \left\{ \frac{t}{(t + \|(x_1, x_2)\|)^4} + \frac{t}{(t + |x_1|)^2} \frac{s}{(s + |x_2|)^2} \right\}. \end{aligned} \tag{11}$$

Let M denote the Hardy-Littlewood maximal function on \mathbb{R}^2 . Now, the estimates (7), (8), and (11) and Lemma 2.3 yield that

$$\begin{aligned} & |\tilde{\psi}_{u_1 u_2} * f(x_1, x_2)| \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \\ & \quad \times \int_{\mathbb{R}^2} \left\{ \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(z_1, z_2)\|)^4} + \frac{u_1 \vee t_1}{(u_1 \vee t_1 + |z_1|)^2} \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |z_2|)^2} \right\} \\ & \quad \times |\psi_{t_1 t_2} * f(x_1 - z_1, x_2 - z_2)| dz_1 dz_2 \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \\ & \quad \times \{ M(\psi_{t_1 t_2} * f)(x_1, x_2) + M_s(\psi_{t_1 t_2} * f)(x_1, x_2) \} \frac{dt_1}{t_1} \frac{dt_2}{t_2}, \end{aligned} \tag{12}$$

which together with the Minkowski inequality and the $L^p(\mathbb{R}^2)$ -boundedness of M and M_s yields that, for $p \in (1, \infty)$,

$$\begin{aligned} & \|\tilde{\psi}_{u_1 u_2} * f\|_{L^p(\mathbb{R}^2)} \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \|\psi_{t_1 t_2} * f\|_{L^p(\mathbb{R}^2)} \frac{dt_1}{t_1} \frac{dt_2}{t_2}. \end{aligned} \quad (13)$$

The estimate (13) combined with the Minkowski inequality shows that

$$\begin{aligned} & \|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^2)} \\ & \leq C \left\{ \int_0^\infty \int_0^\infty \left[\int_0^{u_1} \int_0^{u_2} \left(\frac{t_1}{u_1}\right)^{s_1+1} \left(\frac{t_2}{u_2}\right)^{s_2+1} \right. \right. \\ & \quad \left. \left. \times t_1^{-s_1} t_2^{-s_2} \|\psi_{t_1 t_2} * f\|_{L^p(\mathbb{R}^2)} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right]^q \frac{du_1}{u_1} \frac{du_2}{u_2} \right\}^{1/q} \\ & + C \left\{ \int_0^\infty \int_0^\infty \left[\int_0^{u_1} \int_{u_2}^\infty \left(\frac{t_1}{u_1}\right)^{s_1+1} \left(\frac{t_2}{u_2}\right)^{s_2-1} \dots \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right]^q \frac{du_1}{u_1} \frac{du_2}{u_2} \right\}^{1/q} \\ & + C \left\{ \int_0^\infty \int_0^\infty \left[\int_{u_1}^\infty \int_0^{u_2} \left(\frac{t_1}{u_1}\right)^{s_1-1} \left(\frac{t_2}{u_2}\right)^{s_2+1} \dots \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right]^q \frac{du_1}{u_1} \frac{du_2}{u_2} \right\}^{1/q} \\ & + C \left\{ \int_0^\infty \int_0^\infty \left[\int_{u_1}^\infty \int_{u_2}^\infty \left(\frac{t_1}{u_1}\right)^{s_1-1} \left(\frac{t_2}{u_2}\right)^{s_2-1} \dots \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right]^q \frac{du_1}{u_1} \frac{du_2}{u_2} \right\}^{1/q} \\ & = F_1 + F_2 + F_3 + F_4. \end{aligned}$$

The Hölder inequality and the assumption that $s_i > -1$ for $i = 1, 2$ further imply that

$$\begin{aligned} F_1 & \leq C \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1 q} t_2^{-s_2 q} \left\| \psi_{t_1 t_2} * f \right\|_{L^p(\mathbb{R}^2)}^q \right. \\ & \quad \left. \times \left[\int_{t_1}^\infty \int_{t_2}^\infty \left(\frac{t_1}{u_1}\right)^{s_1+1} \left(\frac{t_2}{u_2}\right)^{s_2+1} \frac{du_1}{u_1} \frac{du_2}{u_2} \right] \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \\ & \leq C \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1 q} t_2^{-s_2 q} \left\| \psi_{t_1 t_2} * f \right\|_{L^p(\mathbb{R}^2)}^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} = C \|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^2)}, \end{aligned}$$

where, and in what follows, we denote by q' the conjugate index of q , namely, $1/q + 1/q' = 1$.

The same argument as for F_1 also yields the desired estimates for F_i with $i = 2, 3, 4$. This proves (5) and hence, by symmetry, the independence of the norm $\|\cdot\|_{\dot{B}_{p,q}^s(\mathbb{R}^2)}$ with respect to the choice of $\psi^{(i)}$, for $i = 1, 2$.

We now turn to the proof of (6). From the estimate (12), it follows that, for all $u_1, u_2 > 0$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} & u_1^{-s_1} u_2^{-s_2} |\tilde{\psi}_{u_1 u_2} * f(x_1, x_2)| \\ & \leq C \left\{ \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \left(\frac{t_1}{u_1} \right)^{s_1} \left(\frac{t_2}{u_2} \right)^{s_2} \right. \\ & \quad \times \left[M \left(t_1^{-s_1} t_2^{-s_2} \psi_{t_1 t_2} * f \right) (x_1, x_2) \right. \\ & \quad \left. \left. + M_s \left(t_1^{-s_1} t_2^{-s_2} \psi_{t_1 t_2} * f \right) (x_1, x_2) \right]^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q}, \end{aligned}$$

which combined with Lemma 2.3 and (10) yields that

$$\begin{aligned} \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} & \leq C \left\| \left\{ \int_0^\infty \int_0^\infty \left[M \left(t_1^{-s_1} t_2^{-s_2} \psi_{t_1 t_2} * f \right) \right. \right. \right. \\ & \quad \left. \left. + M_s \left(t_1^{-s_1} t_2^{-s_2} \psi_{t_1 t_2} * f \right) \right]^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)} \\ & \leq C \left\| \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1 q} t_2^{-s_2 q} |\psi_{t_1 t_2} * f|^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)} \\ & = C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}, \end{aligned}$$

where we have used the vector-valued inequality of Fefferman-Stein in [1]. This proves (6) and, by symmetry, the independence of the norm $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ with respect to the choice of $\psi^{(i)}$ for $i = 1, 2$. This finishes the proof of Theorem 2.6. \square

Based on Theorem 2.6, we now introduce the Besov space $\dot{B}_{pq}^s(\mathbb{R}^2)$ and the Triebel-Lizorkin space $\dot{F}_{pq}^s(\mathbb{R}^2)$ as follows.

Definition 2.7. Let $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$. The Besov space $\dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ is defined by

$$\dot{B}_{pq}^s(\mathbb{R}^2) = \{ f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)' : \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} < \infty \};$$

and the Triebel-Lizorkin space $\dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ is defined by

$$\dot{F}_{pq}^s(\mathbb{R}^2) = \{ f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)' : \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} < \infty \}.$$

Theorem 2.6 shows that the definitions of the spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ are independent of the choices of $\psi^{(i)}$ with $i = 1, 2$.

From Lemma 2.3 and Lemma 2.4, we deduce the ‘discrete’ characterization of Besov spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ and Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$ as below.

Theorem 2.8. *Let $s_1, s_2 \in (-1, 1)$, $s = (s_1, s_2)$ and all other notation be the same as in Lemma 2.4. Then $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ if and only if $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and*

$$\|f\|'_{\dot{B}_{pq}^s(\mathbb{R}^2)} = \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{k_1 s_1 q} 2^{k_2 s_2 q} \|\psi_{k_1 k_2} * f\|_{L^p(\mathbb{R}^2)}^q \right\}^{1/q} < \infty;$$

and $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ if and only if $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and

$$\|f\|'_{\dot{F}_{pq}^s(\mathbb{R}^2)} = \left\| \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{k_1 s_1 q} 2^{k_2 s_2 q} |\psi_{k_1 k_2} * f|^q \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)} < \infty.$$

Furthermore, in this case, $\|\cdot\|'_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|'_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ are equivalent to $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$, respectively.

Since the proof of Theorem 2.8 is essentially the same as that of Theorem 2.6, we omit the details.

The following properties of these spaces can easily be deduced from Theorem 2.8 and the monotonicity of the spaces l^q ; see [9, 11].

Proposition 2.9. *Let $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$. Then*

- (i) *If $p \in [1, \infty]$ and $1 \leq q_1 \leq q_2 \leq \infty$, then $\dot{B}_{pq_1}^s(\mathbb{R}^2) \subset \dot{B}_{pq_2}^s(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{B}_{pq_1}^s(\mathbb{R}^2)$,*

$$\|f\|_{\dot{B}_{pq_2}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{pq_1}^s(\mathbb{R}^2)}.$$

- (ii) *If $p \in (1, \infty)$ and $1 < q_1 \leq q_2 \leq \infty$, then $\dot{F}_{pq_1}^s(\mathbb{R}^2) \subset \dot{F}_{pq_2}^s(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{F}_{pq_1}^s(\mathbb{R}^2)$,*

$$\|f\|_{\dot{F}_{pq_2}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq_1}^s(\mathbb{R}^2)}.$$

- (iii) *If $p \in (1, \infty)$ and $q \in (1, \infty]$, then*

$$\dot{B}_{p, \min(p, q)}^s(\mathbb{R}^2) \subset \dot{F}_{pq}^s(\mathbb{R}^2) \subset \dot{B}_{p, \max(p, q)}^s(\mathbb{R}^2),$$

namely, there exists a positive constant C such that, for all $f \in \dot{B}_{p, \min(p, q)}^s(\mathbb{R}^2)$,

$$\|f\|_{\dot{B}_{p, \max(p, q)}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{p, \min(p, q)}^s(\mathbb{R}^2)}.$$

The following basic properties are useful in the study on the dual and interpolation of these function spaces; see [4, 9].

Proposition 2.10. *Let $s_1, s_2 \in (-1, 1)$ and $s = (s_1, s_2)$. Then*

- (i) *The space $\dot{B}_{pq}^s(\mathbb{R}^2)$ is a Banach space and $\mathcal{S}_{\infty, F}(\mathbb{R}^2) \subset \dot{B}_{pq}^s(\mathbb{R}^2) \subset \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ for $p, q \in [1, \infty]$. If $p, q \in [1, \infty)$, then $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$ is dense in $\dot{B}_{pq}^s(\mathbb{R}^2)$.*
- (ii) *The space $\dot{F}_{pq}^s(\mathbb{R}^2)$ is a Banach space and $\mathcal{S}_{\infty, F}(\mathbb{R}^2) \subset \dot{F}_{pq}^s(\mathbb{R}^2) \subset \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ for $p \in (1, \infty)$ and $q \in [1, \infty]$. If $p, q \in (1, \infty)$, then $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$ is dense in $\dot{F}_{pq}^s(\mathbb{R}^2)$.*

Proof. We only prove the conclusion that $\mathcal{S}_{\infty, F}(\mathbb{R}^2) \subset \dot{B}_{pq}^s(\mathbb{R}^2)$ of (i), and the conclusion that $\dot{B}_{pq}^s(\mathbb{R}^2) \subset \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ can be deduced from Lemma 2.3 and the Hölder inequality. Moreover, by a routine procedure, we can prove that $\mathcal{S}_{\infty}(\mathbb{R}^2)$ is dense in $B_{pq}^s(\mathbb{R}^2)$ for $p, q \in [1, \infty)$ and we omit the details; see [4].

Let $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)$. By Definition 2.1, there exists a Schwartz function $f^\# \in \mathcal{S}(\mathbb{R}^2 \times \mathbb{R})$ satisfying all the properties of Definition 2.1 such that, for all $(x_1, x_2) \in \mathbb{R}^2$,

$$f(x_1, x_2) = \int_{\mathbb{R}} f^\#(x_1, x_2 - x_3, x_3) dx_3.$$

Let $\psi_{t_1 t_2}$ be the same as in Lemma 2.3, and we write

$$\begin{aligned} \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} &\leq \left\{ \int_0^1 \int_0^1 t_1^{-s_1 q} t_2^{-s_2 q} \|\psi_{t_1 t_2} * f\|_{L^p(\mathbb{R}^2)}^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \\ &\quad + \left\{ \int_0^1 \int_1^\infty \dots \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} + \left\{ \int_1^\infty \int_0^1 \dots \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \\ &\quad + \left\{ \int_1^\infty \int_1^\infty \dots \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \\ &= L_1 + L_2 + L_3 + L_4. \end{aligned}$$

To estimate L_1 , for $(x_1, x_2) \in \mathbb{R}^2$ and $t_1, t_2 > 0$, by the vanishing moments of $\psi^{(1)}$ and $\psi^{(2)}$, we write

$$\begin{aligned} &\psi_{t_1 t_2} * f(x_1, x_2) \\ &= \int_{\mathbb{R}} \int_{|z-y_3| \leq |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| \leq \|(x_1, x_2-z)\|/2} \psi_{t_1}^{(1)}(x_1 - y_1, x_2 - y_2 - z) \\ &\quad \times \psi_{t_2}^{(2)}(z - y_3) \left\{ \left[f^\#(y_1, y_2, y_3) - f^\#(x_1, x_2 - z, y_3) \right] \right. \\ &\quad \left. - \left[f^\#(y_1, y_2, z) - f^\#(x_1, x_2 - z, z) \right] \right\} dy_1 dy_2 dy_3 dz + \end{aligned}$$

$$\begin{aligned}
 &+ \int_{\mathbb{R}} \int_{|z-y_3| > |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| \leq \|(x_1, x_2-z)\|/2} \dots \\
 &+ \int_{\mathbb{R}} \int_{|z-y_3| \leq |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| > \|(x_1, x_2-z)\|/2} \dots \\
 &+ \int_{\mathbb{R}} \int_{|z-y_3| > |z|/2} \int_{\|(x_1-y_1, x_2-y_2-z)\| > \|(x_1, x_2-z)\|/2} \dots
 \end{aligned}$$

Then, the same argument as that for I_i with $i = 1, 2, 3, 4$ in the proof of Lemma 2.3 yields that, for any $N \in \mathbb{Z}$, there exists a positive constant $C = C_{N,f}$ such that, for all $t_1, t_2 > 0$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$|\psi_{t_1 t_2} * f(x_1, x_2)| \leq C t_1 t_2 \frac{1}{(1 + \|(x_1, x_2)\|)^N},$$

which implies the desired estimate for L_1 . The same argument as that for J_i with $i = 1, 2, 3, 4$ in the proof of Lemma 2.3 can yield a desired estimate for L_2 . The estimates for L_3 and L_4 can be obtained in a similar way. We omit the details. This finishes the proof of Proposition 2.9. \square

Based on Proposition 2.10, we can give out the dual spaces of these spaces as below, which can be proved by an argument same as that of Theorem 7.1 in [4]. We omit the details.

Proposition 2.11. *Let $s_1, s_2 \in (-1, 1)$, $s = (s_1, s_2)$, and $-s = (-s_1, -s_2)$. Then*

- (i) $(\dot{B}_{pq}^s(\mathbb{R}^2))^* = \dot{B}_{p'q'}^{-s}(\mathbb{R}^2)$ for $p, q \in [1, \infty)$ with $1/p + 1/p' = 1/q + 1/q' = 1$. More precisely, given $g \in \dot{B}_{p'q'}^{-s}(\mathbb{R}^2)$, then $\mathcal{L}_g(f) = \langle f, g \rangle$ defines a linear functional on $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$ such that

$$|\mathcal{L}_g(f)| \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \|g\|_{\dot{B}_{p'q'}^{-s}(\mathbb{R}^2)},$$

and this linear functional can be extended to $\dot{B}_{pq}^s(\mathbb{R}^2)$ with norm at most $C \|g\|_{\dot{B}_{p'q'}^{-s}(\mathbb{R}^2)}$.

Conversely, if \mathcal{L} is a linear functional on $\dot{B}_{pq}^s(\mathbb{R}^2)$, then there exists a unique $g \in \dot{B}_{p'q'}^{-s}(\mathbb{R}^2)$ such that $\mathcal{L}_g(f) = \langle f, g \rangle$ defines a linear functional on $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$, and \mathcal{L} is the extension of \mathcal{L}_g with $\|g\|_{\dot{B}_{p'q'}^{-s}(\mathbb{R}^2)} \leq C \|\mathcal{L}\|$.

- (ii) $(\dot{F}_{pq}^s(\mathbb{R}^2))^* = \dot{F}_{p'q'}^{-s}(\mathbb{R}^2)$ for $p, q \in (1, \infty)$ with $1/p + 1/p' = 1/q + 1/q' = 1$. More precisely, given $g \in \dot{F}_{p'q'}^{-s}(\mathbb{R}^2)$, then $\mathcal{L}_g(f) = \langle f, g \rangle$ defines a linear functional on $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$ such that

$$|\mathcal{L}_g(f)| \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \|g\|_{\dot{F}_{p'q'}^{-s}(\mathbb{R}^2)},$$

and this linear functional can be extended to $\dot{F}_{pq}^s(\mathbb{R}^2)$ with norm at most $C\|g\|_{\dot{F}_{p'q'}^{-s}(\mathbb{R}^2)}$.

Conversely, if \mathcal{L} is a linear functional on $\dot{F}_{pq}^s(\mathbb{R}^2)$, then there exists a unique $g \in \dot{F}_{p'q'}^{-s}(\mathbb{R}^2)$ such that $\mathcal{L}_g(f) = \langle f, g \rangle$ defines a linear functional on $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$, and \mathcal{L} is the extension of \mathcal{L}_g with $\|g\|_{\dot{F}_{p'q'}^{-s}(\mathbb{R}^2)} \leq C\|\mathcal{L}\|$.

Remark 2.12. For $s_1, s_2 \in (-1, 1)$, $s = (s_1, s_2)$ and $p, q \in [1, \infty]$, let us now define ${}_{\circ}\dot{B}_{pq}^s(\mathbb{R}^2)$ to be the completion of $\mathcal{S}_{\infty, F}(\mathbb{R}^2)$ in $\dot{B}_{pq}^s(\mathbb{R}^2)$ endowed with the same norm as $\dot{B}_{pq}^s(\mathbb{R}^2)$. Then, in the sense of Proposition 2.11, we have

$$({}_{\circ}\dot{B}_{pq}^s(\mathbb{R}^2))^* = B_{p'q'}^{-s}(\mathbb{R}^2) \tag{14}$$

with $-s, p'$, and q' having the same meaning as in Proposition 2.11. The equality (14) is new only for the case $\max(p, q) = \infty$ in comparison with Proposition 2.11. This fact can be easily proved by combining the argument in [4, pp. 116–120] with that in [9, p. 180]; see also [10, pp. 121, 122]. We omit the details.

Now, using these properties, we establish the Lusin-area characterizations of Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$. First we introduce the following two kinds of Lusin-area functions.

Definition 2.13. Let $s_i \in \mathbb{R}$ and $\alpha_i > 0$ for $i = 1, 2$, $s = (s_1, s_2)$ and $q \in (1, \infty)$. Let $\psi_{t_1 t_2}$ for $t_i > 0$ with $i = 1, 2$ be the same as in Lemma 2.3 and $\chi = \chi_{(0,1)}$. For $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and $(x_1, x_2) \in \mathbb{R}^2$, we define

$$S_{q; \alpha_1, \alpha_2}^s(f)(x_1, x_2) = \left\{ \int_0^\infty \int_0^\infty \int_{\mathbb{R}^2} \chi\left(\frac{|x_1 - y_1|}{\alpha_1 t_1}\right) \chi\left(\frac{|x_2 - y_2|}{\alpha_2 t_2}\right) \times t_1^{-s_1 q} t_2^{-s_2 q} \left| \psi_{t_1 t_2} * f(y_1, y_2) \right|^q dy_1 dy_2 \frac{dt_1}{t_1^2} \frac{dt_2}{t_2^2} \right\}^{1/q}$$

and

$$\begin{aligned} \tilde{S}_{q; \alpha_1, \alpha_2}^s(f)(x_1, x_2) &= \left\{ \int_0^\infty \int_0^{t_1^2} \int_{\mathbb{R}^2} \chi\left(\frac{|x_1 - y_1|}{\alpha_1 t_1}\right) \chi\left(\frac{|x_2 - y_2|}{\alpha_2 t_1^2}\right) \right. \\ &\quad \left. \times t_1^{-s_1 q} t_2^{-s_2 q} \left| \psi_{t_1 t_2} * f(y_1, y_2) \right|^q dy_1 dy_2 \frac{dt_2}{t_1^2 t_2} \frac{dt_1}{t_1^2} \right\}^{1/q} \\ &+ \left\{ \int_0^\infty \int_{t_1^2}^\infty \int_{\mathbb{R}^2} \chi\left(\frac{|x_1 - y_1|}{\alpha_1 t_1}\right) \chi\left(\frac{|x_2 - y_2|}{\alpha_2 t_2}\right) \right. \\ &\quad \left. \times t_1^{-s_1 q} t_2^{-s_2 q} \left| \psi_{t_1 t_2} * f(y_1, y_2) \right|^q dy_1 dy_2 \frac{dt_2}{t_2^2} \frac{dt_1}{t_1^2} \right\}^{1/q}. \end{aligned}$$

The characterizations of Lusin-area functions of $\dot{F}_{pq}^s(\mathbb{R}^2)$ can be stated as below.

Theorem 2.14. *Let $s_1, s_2 \in (-1, 1)$, $s = (s_1, s_2)$, $\alpha_i > 0$ for $i = 1, 2$, and $p, q \in (1, \infty)$. Then, for $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$, the following three propositions are equivalent:*

- (i) $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$;
- (ii) $S_{q; \alpha_1, \alpha_2}^s(f) \in L^p(\mathbb{R}^2)$;
- (iii) $\tilde{S}_{q; \alpha_1, \alpha_2}^s(f) \in L^p(\mathbb{R}^2)$.

Furthermore, in this case,

$$\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \sim \|S_{q; \alpha_1, \alpha_2}^s(f)\|_{L^p(\mathbb{R}^2)} \sim \|\tilde{S}_{q; \alpha_1, \alpha_2}^s(f)\|_{L^p(\mathbb{R}^2)}.$$

Proof. Letting $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$, we first prove that $S_{q; \alpha_1, \alpha_2}^s(f), \tilde{S}_{q; \alpha_1, \alpha_2}^s(f) \in L^p(\mathbb{R}^2)$ and there exists a positive constant C such that, for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$,

$$\|S_{q; \alpha_1, \alpha_2}^s(f)\|_{L^p(\mathbb{R}^2)} \leq C\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \tag{15}$$

and

$$\|\tilde{S}_{q; \alpha_1, \alpha_2}^s(f)\|_{L^p(\mathbb{R}^2)} \leq C\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}. \tag{16}$$

To this end, for $(z_1, z_2) \in \mathbb{R}^2$, $t_i > 0$, $|x_i - y_i| < \alpha_i u_i$ with $u_i > 0$, $i = 1, 2$, by (11), we have

$$\begin{aligned} & \int_{\mathbb{R}} \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(y_1 - z_1, y_2 - z_2) - (0, w)\|)^4} \cdot \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |w|)^2} dw \\ & \leq C \left\{ \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(x_1 - z_1, x_2 - z_2)\|)^4} \right. \\ & \quad \left. + \frac{u_1 \vee t_1}{(u_1 \vee t_1 + |x_1 - z_1|)^2} \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |x_2 - z_2|)^2} \right\}, \tag{17} \end{aligned}$$

and, similarly, for $|x_1 - y_1| < \alpha_1 u_1$ and $|x_2 - y_2| < \alpha_2 u_1^2$, we also have

$$\begin{aligned} & \int_{\mathbb{R}} \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(y_1 - z_1, y_2 - z_2) - (0, w)\|)^4} \cdot \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |w|)^2} dw \\ & \leq C \left\{ \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(x_1 - z_1, x_2 - z_2)\|)^4} \right. \\ & \quad \left. + \frac{u_1 \vee t_1}{(u_1 \vee t_1 + |x_1 - z_1|)^2} \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |x_2 - z_2|)^2} \right\}. \tag{18} \end{aligned}$$

Now the estimates (7), (8), Lemma 2.3, and the estimates (17) and (18), respectively,

yield that

$$\begin{aligned} & \chi\left(\frac{|x_1 - y_1|}{\alpha_1 u_1}\right) \chi\left(\frac{|x_2 - y_2|}{\alpha_2 u_2}\right) |\psi_{u_1 u_2} * f(y_1, y_2)| \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \\ & \quad \times \{M(\psi_{t_1 t_2} * f)(x_1, x_2) + M_s(\psi_{t_1 t_2} * f)(x_1, x_2)\} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \end{aligned} \quad (19)$$

and

$$\begin{aligned} & \chi\left(\frac{|x_1 - y_1|}{\alpha_1 u_1}\right) \chi\left(\frac{|x_2 - y_2|}{\alpha_2 u_1^2}\right) |\psi_{u_1 u_2} * f(y_1, y_2)| \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \\ & \quad \times \{M(\psi_{t_1 t_2} * f)(x_1, x_2) + M_s(\psi_{t_1 t_2} * f)(x_1, x_2)\} \frac{dt_1}{t_1} \frac{dt_2}{t_2}. \end{aligned} \quad (20)$$

Replacing (12) with (19) and (20), respectively, and repeating the argument of (6) in Theorem 2.6, we then obtain (15) and (16).

We now show the converse of (15) and (16) and by similarity we only prove the converse of (16). To this end, letting $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ and $g \in \dot{F}_{p'q'}^{-s}(\mathbb{R}^2)$ and $\{\psi_{t_1 t_2}\}_{t_1, t_2 > 0}$ be the same as in Lemma 2.3, by (16), Proposition 2.11, and the Hölder inequality, we obtain

$$\begin{aligned} \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} &= \sup_{\|g\|_{\dot{F}_{p'q'}^{-s}(\mathbb{R}^2)} \leq 1} \left| \int_0^\infty \int_0^\infty \langle \psi_{t_1 t_2} * f, \psi_{t_1 t_2} * g \rangle \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right| \\ &\leq C \sup_{\|g\|_{\dot{F}_{p'q'}^{-s}(\mathbb{R}^2)} \leq 1} \int_{\mathbb{R}^2} \tilde{S}_{q; \alpha_1, \alpha_2}^s(f)(x_1, x_2) \tilde{S}_{q'; \alpha_1, \alpha_2}^{-s}(g)(x_1, x_2) dx_1 dx_2 \\ &\leq C \|\tilde{S}_{q; \alpha_1, \alpha_2}^s(f)\|_{L^p(\mathbb{R}^2)}, \end{aligned}$$

which establishes the equivalence of (i) and (ii) and, hence, completes the proof of Theorem 2.14. \square

Remark 2.15. From the proof of Theorem 2.14, it is easy to see that (15) and (16) also hold for $p \in (1, \infty)$ and $q = \infty$.

Let us now state the Littlewood-Paley theorem corresponding to the following g -function.

Definition 2.16. With the same notation as in Lemma 2.3, for $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$, we define its Littlewood-Paley g -function, $g(f)$, of f by

$$g(f)(x_1, x_2) = \left\{ \int_0^\infty \int_0^\infty |\psi_{t_1 t_2} * f(x_1, x_2)|^2 \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/2},$$

where $(x_1, x_2) \in \mathbb{R}^2$.

Using the boundedness of vector-valued singular integrals and a dual argument similar to the proof of Theorem 2.14, we can obtain the following result and we omit the details; see [3].

Theorem 2.17. *Let $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and $p \in (1, \infty)$. Then the norm $\|g(f)\|_{L^p(\mathbb{R}^2)}$ is equivalent to the norm $\|f\|_{L^p(\mathbb{R}^2)}$.*

From Theorem 2.17, Definition 2.7, and Theorem 2.14, it is easy to deduce the following conclusions:

Corollary 2.18. *Let $p \in (1, \infty)$. Then $\dot{F}_{p2}^0(\mathbb{R}^2) = L^p(\mathbb{R}^2)$ with an equivalent norm. Moreover, the norms $\|f\|_{L^p(\mathbb{R}^2)}$, $\|g(f)\|_{L^p(\mathbb{R}^2)}$, $\|S_{2; \alpha_1, \alpha_2}^{(0,0)}\|_{L^p(\mathbb{R}^2)}$, and $\|\tilde{S}_{2; \alpha_1, \alpha_2}^{(0,0)}\|_{L^p(\mathbb{R}^2)}$ are mutually equivalent, where $\alpha_1, \alpha_2 > 0$.*

We now establish a new characterization of Besov and Triebel-Lizorkin spaces. First, we introduce the following new Besov and Triebel-Lizorkin norms.

Definition 2.19. Let $\psi_{t_1 t_2}$ be the same as in Lemma 2.1 and $s_1, s_2 \in \mathbb{R}$. For $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$, define

$$\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} = \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1 q} (t_1^2 + t_2)^{-s_2 q} \|\psi_{t_1 t_2} * f\|_{L^p(\mathbb{R}^2)}^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q}$$

for $p, q \in [1, \infty]$, and

$$\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} = \left\| \left\{ \int_0^\infty \int_0^\infty t_1^{-s_1 q} (t_1^2 + t_2)^{-s_2 q} |\psi_{t_1 t_2} * f|^q \frac{dt_1}{t_1} \frac{dt_2}{t_2} \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)}$$

for $p \in (1, \infty)$ and $q \in (1, \infty]$, where the usual modifications are made when $p = \infty$ or $q = \infty$.

We can also show that the definitions of $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ are independent of the choices of $\psi^{(i)}$ for $i = 1, 2$ by an argument similar to that of Theorem 2.6.

Theorem 2.20. *Let $s_1, s_2 \in (-1, 1)$, $|s_1 + 2s_2| < 1$, and $s = (s_1, s_2)$. The norm $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ with $p, q \in [1, \infty]$ and the norm $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ are independent of the choices of $\psi^{(i)}$ for $i = 1, 2$.*

Proof. Let $\tilde{\psi}^{(i)}$ for $i = 1, 2$ be the same as in the proof of Theorem 2.6. We denote the corresponding norms, respectively, by $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$. To prove the theorem, by symmetry, we only need to show that there exists a positive constant C such that, for all $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$,

$$\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$$

and

$$\|f\|_{2\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C\|f\|_{1\dot{F}_{pq}^s(\mathbb{R}^2)},$$

which, with the same notation as in the proof of Theorem 2.6, are deduced from the following facts:

$$\int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_1^2+t_2}{u_1^2+u_2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \leq C \quad (21)$$

and

$$\int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_1^2+t_2}{u_1^2+u_2}\right)^{s_2} \frac{du_1}{u_1} \frac{du_2}{u_2} \leq C.$$

The proofs of both facts are similar by symmetry and we only show (21). To this end, we consider two cases.

- *Case 1:* $u_2 \leq u_1^2$. In this case, the left-hand side of (21) is equivalent to

$$\begin{aligned} & \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_1^2+t_2}{u_1^2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ &= \int_0^\infty \int_0^{t_1^2} \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_1^2+t_2}{u_1^2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ & \quad + \int_0^\infty \int_{t_1^2}^\infty \dots = P_1 + P_2. \end{aligned}$$

For P_1 , since $t_1^2+t_2 \sim t_1^2$, we then have

$$P_1 \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1+2s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \leq C,$$

by the assumption that $|s_1+2s_2| < 1$. For P_2 , since $t_1^2+t_2 \sim t_2$, we have

$$P_2 \leq C \int_0^\infty \int_{t_1^2}^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_2}{u_1^2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2}.$$

If $s_2 \leq 0$, the fact that $t_2 \geq t_1^2$ and the assumption that $|s_1+2s_2| < 1$ imply the desired estimate of P_2 ; and if $s_2 > 0$, the assumptions that $u_1^2 \geq u_2$ and $s_1, s_2 \in (-1, 1)$ also imply the desired estimate of P_2 , which completes the proof of case 1.

- *Case 2:* $u_1^2 < u_2$. In this case, the left-hand side of (21) is equivalent to

$$\begin{aligned} & \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_1+t_2}{u_2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ & \leq C \int_0^\infty \int_0^{t_1^2} \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_1^2}{u_2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2} \\ & \quad + C \int_0^\infty \int_{t_1^2}^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1}\right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2}\right) \left(\frac{t_1}{u_1}\right)^{s_1} \left(\frac{t_2}{u_2}\right)^{s_2} \frac{dt_1}{t_1} \frac{dt_2}{t_2}. \end{aligned}$$

We can obtain the desired estimate for the second term of the last expression by the assumptions that $s_1, s_2 \in (-1, 1)$. For the first term, if $s_2 \geq 0$, the fact that $u_1^2 < u_2$ and the assumption that $|s_1 + 2s_2| < 1$ can imply the desired estimate; and if $s_2 < 0$, the fact that $t_2 < t_1^2$ and the assumptions that $s_1, s_2 \in (-1, 1)$ also yield the desired estimate. This finishes the proof of (21) and hence, the proof of Theorem 2.20. \square

From Lemma 2.3 and Lemma 2.4, by an argument similar to the proof of Theorem 2.20, we also deduce the new ‘discrete’ characterization of Besov spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ and Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$ as follows. We omit the details.

Theorem 2.21. *Let $s_1, s_2 \in (-1, 1)$, $|s_1 + 2s_2| < 1$, $s = (s_1, s_2)$ and all other notation be the same as in Lemma 2.4. Then $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ if and only if $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and*

$$\begin{aligned} & \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}'' \\ & = \left\{ \sum_{k_1=-\infty}^\infty \sum_{k_2=-\infty}^\infty 2^{k_1 s_1 q} (2^{-2k_1} + 2^{-k_2})^{-s_2 q} \|\psi_{k_1 k_2} * f\|_{L^p(\mathbb{R}^2)}^q \right\}^{1/q} < \infty, \end{aligned}$$

and $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ if and only if $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ and

$$\begin{aligned} & \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}'' \\ & = \left\| \left\{ \sum_{k_1=-\infty}^\infty \sum_{k_2=-\infty}^\infty 2^{k_1 s_1 q} (2^{-2k_1} + 2^{-k_2})^{-s_2 q} |\psi_{k_1 k_2} * f|^q \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)} < \infty. \end{aligned}$$

Furthermore, in this case, $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}''$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}''$ are equivalent to $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$, respectively.

As a corollary of Theorem 2.20 and Theorem 2.6, we obtain a new characterization of Besov spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ and Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$ as below, which clearly reveals the difference between $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ with the classical product Besov spaces and Triebel-Lizorkin spaces in [7].

Corollary 2.22. *Let $s_1, s_2 \in (-1, 1)$, $|s_1 + 2s_2| < 1$, and $s = (s_1, s_2)$. Then*

- (i) *For $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$, $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ if and only if $\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} < \infty$. Moreover, in this case, $\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \sim \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$.*
- (ii) *For $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$, $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$ if and only if $\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} < \infty$. Moreover, in this case, $\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \sim \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$.*

Proof. Theorem 2.6 and Theorem 2.20 imply that the definitions of the norms $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$, $\|\cdot\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$, $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$, and $\|\cdot\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$ are independent of the choices of $\psi^{(i)}$ for $i = 1, 2$. Let $\psi^{(i)}$ for $i = 1, 2$ be the same as in Lemma 2.3 satisfying the following additional conditions:

$$\text{supp } \hat{\psi}^{(1)} \subset \{(\xi_1, \xi_2) \in \mathbb{R}^2 : 1/2 \leq \|(\xi_1, \xi_2)\| \leq 2\}$$

and

$$\text{supp } \hat{\psi}^{(2)} \subset \{\xi \in \mathbb{R} : 1/2 \leq |\xi| \leq 2\}.$$

From the definitions of $\psi_{t_1 t_2}$, the above conditions of the supports of $\psi^{(i)}$ with $i = 1, 2$, and the Plancherel principle, it is easy to deduce that $\psi_{t_1 t_2} = 0$ if $t_1^2 \leq 8t_2$. Thus, using such $\psi_{t_1 t_2}$ and noticing that $t_1^2 + t_2 \sim t_2$ if $t_1^2 \leq 8t_2$, by Definitions 2.5 and 2.19, we easily obtain that $\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \sim \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$ and $\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \sim \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}$. This finishes the proof of Corollary 2.22. \square

3. Boundedness of flag singular integrals

We now establish the boundedness on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ of flag singular integrals.

Since it is well-known that the flag singular integral is bounded on $L^p(\mathbb{R}^2)$ for $p \in (1, \infty)$, we then automatically deduce that it is also bounded on classical Besov spaces and the new Besov spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ associated with flag kernels, when $p \in (1, \infty)$. However, this is not true for Besov spaces when $p = 1$ or $p = \infty$ and Triebel-Lizorkin spaces. Moreover, our argument also gives a direct proof for the boundedness of flag singular integrals in $L^p(\mathbb{R}^2)$ with $p \in (1, \infty)$.

Theorem 3.1. *Let K be an integrable flag kernel on \mathbb{R}^2 as in Definition 1.2, and $s = (s_1, s_2)$ with $s_i \in (-1, 1)$ for $i = 1, 2$. Then*

- (i) *If $p, q \in [1, \infty]$, then there exists a positive constant C such that, for all $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$,*

$$\|K * f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}.$$
- (ii) *If $p \in (1, \infty)$ and $q \in (1, \infty]$, then there exists a positive constant C such that, for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$,*

$$\|K * f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}.$$

Before proving Theorem 3.1, we first establish several lemmas which are used in the proof of Theorem 3.1.

Lemma 3.2. *Let K be a distribution on \mathbb{R} which is a continuous function on $\Omega_1 = \mathbb{R} \setminus \{0\}$ and ψ_s be a function on \mathbb{R} for all $s > 0$. Suppose that there exists a positive constant C_K such that K satisfies the following conditions:*

- (i) For all $x_3 \in \Omega_1$, $|K(x_3)| \leq C_K \frac{1}{|x_3|}$.
- (ii) For all $x_3 \in \Omega_1$ and $x'_3 \in \mathbb{R}$ with $|x_3 - x'_3| \leq |x_3|/2$,

$$|K(x_3) - K(x'_3)| \leq C_K \frac{|x_3 - x'_3|}{|x_3|^2}.$$

- (iii) For any given bump function φ on \mathbb{R} and any $\delta > 0$,

$$\left| \int_{\mathbb{R}} K(x_3) \varphi(\delta x_3) dx_3 \right| \leq C_K.$$

Suppose also that there exists a positive constant C_ψ such that

- (iv) For all $s > 0$ and $x_3 \in \mathbb{R}$, $|\psi_s(x_3)| \leq C_\psi \frac{s}{(s+|x_3|)^2}$.
- (v) For all $s > 0$ and $x_3, x'_3 \in \mathbb{R}$ with $|x_3 - x'_3| \leq (s + |x_3|)/2$,

$$|\psi_s(x_3) - \psi_s(x'_3)| \leq C_\psi \frac{|x_3 - x'_3|}{s + |x_3|} \cdot \frac{s}{(s + |x_3|)^2}.$$

- (vi) For all $s > 0$, $\int_{\mathbb{R}} \psi_s(x_3) dx_3 = 0$.
- (vii) $\text{supp } \psi_s \subset \{x_3 \in \mathbb{R} : |x_3| \leq s\}$.

Then there exists a positive constant C such that, for all $s > 0$ and $z \in \mathbb{R}$,

$$\left| \int_{\mathbb{R}} K(z - x_3) \psi_s(x_3) dx_3 \right| \leq CC_K C_\psi \frac{s}{(s + |z|)^2}.$$

Proof. We consider two cases.

- *Case 1:* $|z| \leq 5s$. In this case, let $\theta \in C_0^\infty(\mathbb{R})$, $0 \leq \theta \leq 1$, $\text{supp } \theta \subset \{x \in \mathbb{R} : |x| \leq 2\}$, and $\theta(x) = 1$ if $|x| \leq 1$. We then define $\xi(z) = \theta(\frac{|z|}{10s})$ for $z \in \mathbb{R}$. By (vii),

we can write

$$\begin{aligned} \left| \int_{\mathbb{R}} K(z - x_3) \psi_s(x_3) dx_3 \right| &\leq \int_{|x_3 - z| \leq (s + |z|)/2} |K(z - x_3) [\psi_s(x_3) - \psi_s(z)]| \xi(x_3) dx_3 \\ &\quad + \int_{|x_3 - z| > (s + |z|)/2} \dots + |\psi_s(z)| \left| \int_{\mathbb{R}} K(z - x_3) \xi(x_3) dx_3 \right| \\ &= G_1 + G_2 + G_3. \end{aligned}$$

Notice that ξ is a normalized bump function multiplied with a normalizing constant and some dilation. The assumptions (iii) and (iv) give us the desired estimate for G_3 :

$$G_3 \leq C_K C_\psi \frac{s}{(s + |z|)^2}.$$

From (i) and (v), it follows that

$$G_1 \leq C_K C_\psi \int_{\mathbb{R}} \frac{1}{|z - x_3|} \frac{|x_3 - z|}{s + |x_3|} \frac{s}{(s + |x_3|)^2} dx_3 \leq C C_K C_\psi \frac{1}{s}.$$

The assumptions (i) and (iv) also yield that

$$\begin{aligned} G_2 &\leq \int_{|x_3 - z| > (s + |z|)/2} |K(z - x_3)| [|\psi_s(x_3)| + |\psi_s(z)|] \xi(x_3) dx_3 \\ &\leq C C_K \frac{1}{s} \left[\int_{\mathbb{R}} |\psi_s(x_3)| dx_3 + C_\psi \frac{s^2}{(s + |z|)^2} \right] \leq C C_K C_\psi \frac{s}{(s + |z|)^2}, \end{aligned}$$

which completes the proof of case 1.

- *Case 2:* $|z| > 5s$. In this case, by (vi), (vii), (ii), and (iv), we have

$$\begin{aligned} \left| \int_{\mathbb{R}} K(z - x_3) \psi_s(x_3) dx_3 \right| &= \left| \int_{\mathbb{R}} [K(z - x_3) - K(z)] \psi_s(x_3) dx_3 \right| \\ &\leq C_K \int_{\mathbb{R}} \frac{|x_3|}{|z|^2} |\psi_s(x_3)| dx_3 \leq C_K C_\psi \frac{s}{|z|^2}. \end{aligned}$$

This finishes the proof of Lemma 3.2. □

The same argument as in the proof of Lemma 3.2 gives us the following result and we omit the details.

Lemma 3.3. *Let K be a distribution on \mathbb{R}^2 which is a continuous function on $\Omega_2 = \mathbb{R}^2 \setminus \{(0, 0)\}$ and ψ_s be a function on \mathbb{R}^2 for all $s > 0$. Suppose that there exists a positive constant C_K such that K satisfies the following conditions:*

- (i) For all $(x_1, x_2) \in \Omega_2$, $|K(x_1, x_2)| \leq C_K \frac{1}{\|(x_1, x_2)\|^3}$.
- (ii) For all $(x_1, x_2) \in \Omega_2$ and $(x'_1, x'_2) \in \mathbb{R}^2$ with $\|(x_1, x_2) - (x'_1, x'_2)\| \leq \|(x_1, x_2)\|/2$,

$$|K(x_1, x_2) - K(x'_1, x'_2)| \leq C_K \frac{\|(x_1, x_2) - (x'_1, x'_2)\|}{\|(x_1, x_2)\|^4}.$$

- (iii) For any given bump function φ on \mathbb{R}^2 and any $\delta > 0$,

$$\left| \int_{\mathbb{R}^2} K(x_1, x_2) \varphi(\delta x_1, \delta^2 x_2) dx_1 dx_2 \right| \leq C_K.$$

Suppose also that there exists a positive constant C_ψ such that

- (iv) For all $s > 0$ and $(x_1, x_2) \in \mathbb{R}^2$, $|\psi_s(x_1, x_2)| \leq C_\psi \frac{s}{(s + \|(x_1, x_2)\|)^4}$.
- (v) For all $s > 0$ and $(x_1, x_2), (x'_1, x'_2) \in \mathbb{R}^2$ with $\|(x_1, x_2) - (x'_1, x'_2)\| \leq (s + \|(x_1, x_2)\|)/2$,

$$|\psi_s(x_1, x_2) - \psi_s(x'_1, x'_2)| \leq C_\psi \frac{\|(x_1, x_2) - (x'_1, x'_2)\|}{s + \|(x_1, x_2)\|} \cdot \frac{s}{(s + \|(x_1, x_2)\|)^4}.$$

- (vi) For all $s > 0$, $\int_{\mathbb{R}^2} \psi_s(x_1, x_2) dx_1 dx_2 = 0$.
- (vii) $\text{supp } \psi_s \subset \{(x_1, x_2) \in \mathbb{R}^2 : \|(x_1, x_2)\| \leq s\}$.

Then there exists a positive constant C such that, for all $s > 0$ and $z \in \mathbb{R}$,

$$\left| \int_{\mathbb{R}^2} K(x_1 - y_1, x_2 - y_2) \psi_s(y_1, y_2) dy_1 dy_2 \right| \leq C C_K C_\psi \frac{s}{(s + \|(x_1, x_2)\|)^4}.$$

We also need the following basic estimate.

Lemma 3.4. Let $\psi^{(1)} \in C_0^\infty(\mathbb{R}^2)$, $\text{supp } \psi^{(1)} \subset \{(x_1, x_2) \in \mathbb{R}^2 : \|(x_1, x_2)\| < 1\}$ and

$$\int_{\mathbb{R}^2} \psi^{(1)}(x_1, x_2) dx_1 dx_2 = 0. \tag{22}$$

For $s > 0$ and $(x_1, x_2) \in \mathbb{R}^2$, set $\psi_s^{(1)}(x_1, x_2) = \frac{1}{s^3} \psi^{(1)}(\frac{x_1}{s}, \frac{x_2}{s^2})$. Then there exists a positive constant C such that

- (i) For all $t_1, u_1 > 0$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$|(\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(x_1, x_2)| \leq C \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \frac{t_1 \vee u_1}{(t_1 \vee u_1 + \|(x_1, x_2)\|)^4}.$$

(ii) For all $t_1, u_1 > 0$ and $(x_1, x_2), (y_1, y_2) \in \mathbb{R}^2$ with $\|(y_1, y_2) - (x_1, x_2)\| \leq (t_1 \vee u_1 + \|(x_1, x_2)\|)/2$,

$$\begin{aligned} & |(\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(y_1, y_2) - (\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(x_1, x_2)| \\ & \leq C \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \frac{\|(y_1, y_2) - (x_1, x_2)\|}{t_1 \vee u_1 + \|(x_1, x_2)\|} \cdot \frac{t_1 \vee u_1}{(t_1 \vee u_1 + \|(x_1, x_2)\|)^4}. \end{aligned}$$

Proof. The same argument as in the proof of (7) also yields (i). We only show (ii) in the case $u_1 \leq t_1$. In this case, the estimate (ii) becomes that if $\|(y_1, y_2) - (x_1, x_2)\| \leq (t_1 + \|(x_1, x_2)\|)/2$ then

$$\begin{aligned} & |(\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(y_1, y_2) - (\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(x_1, x_2)| \\ & \leq C \frac{u_1}{t_1} \frac{\|(y_1, y_2) - (x_1, x_2)\|}{t_1 + \|(x_1, x_2)\|} \cdot \frac{t_1}{(t_1 + \|(x_1, x_2)\|)^4}. \end{aligned}$$

To guarantee

$$|(\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(y_1, y_2) - (\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(x_1, x_2)| \neq 0$$

when $\|(y_1, y_2) - (x_1, x_2)\| \leq (t_1 + \|(x_1, x_2)\|)/2$, we always have $\|(x_1, x_2)\| \leq Ct_1$. Thus, by (22) and the mean value theorem,

$$\begin{aligned} & |(\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(y_1, y_2) - (\psi_{t_1}^{(1)} * \psi_{u_1}^{(1)})(x_1, x_2)| \\ & = \left| \int_{\mathbb{R}^2} \left\{ [\psi_{t_1}^{(1)}(y_1 - z_1, y_2 - z_2) - \psi_{t_1}^{(1)}(x_1 - z_1, x_2 - z_2)] \right. \right. \\ & \quad \left. \left. - [\psi_{t_1}^{(1)}(y_1, y_2) - \psi_{t_1}^{(1)}(x_1, x_2)] \right\} \psi_{u_1}^{(1)}(z_1, z_2) dz_1 dz_2 \right| \\ & \leq C \frac{u_1}{t_1^5} \|(y_1, y_2) - (x_1, x_2)\| \leq C \frac{u_1}{t_1} \frac{\|(y_1, y_2) - (x_1, x_2)\|}{t_1 + \|(x_1, x_2)\|} \frac{t_1}{(t_1 + \|(x_1, x_2)\|)^4}, \end{aligned}$$

which completes the proof of Lemma 3.4. □

The same argument as in the proof of Lemma 3.4 also yields the following basic estimates and we omit the details.

Lemma 3.5. Let $\psi^{(2)} \in C_0^\infty(\mathbb{R})$, $\text{supp } \psi^{(2)} \subset (-1, 1)$, and $\int_{\mathbb{R}} \psi(x_3) dx_3 = 0$. For $s > 0$ and $x_3 \in \mathbb{R}$, set $\psi_s^{(2)}(x_3) = \frac{1}{s} \psi^{(2)}(\frac{x_3}{s})$. Then there exists a positive constant C such that

(i) For all $t_2, u_2 > 0$ and $x_3 \in \mathbb{R}$,

$$|(\psi_{t_2}^{(2)} * \psi_{u_2}^{(2)})(x_3)| \leq C \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \frac{t_2 \vee u_2}{(t_2 \vee u_2 + |x_3|)^2}.$$

(ii) For all $t_2, u_2 > 0$ and $x_3, y_3 \in \mathbb{R}$ with $|y_3 - x_3| \leq (t_2 \vee u_2 + |x_3|)/2$,

$$\begin{aligned} & |(\psi_{t_2}^{(2)} * \psi_{u_2}^{(2)})(y_3) - (\psi_{t_2}^{(2)} * \psi_{u_2}^{(2)})(x_3)| \\ & \leq C \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \frac{|y_3 - x_3|}{t_2 \vee u_2 + |x_3|} \cdot \frac{t_2 \vee u_2}{(t_2 \vee u_2 + |x_3|)^2}. \end{aligned}$$

Lemma 3.6. Let K^\sharp be a product kernel on $\mathbb{R}^2 \times \mathbb{R}$ and ψ_s be the same as in Lemma 3.2. For $(x_1, x_2) \in \mathbb{R}^2$ and $z \in \mathbb{R}$, define

$$\tilde{K}(x_1, x_2, x_3) = \int_{\mathbb{R}^2} K^\sharp(x_1, x_2, x_3 - z) \psi_s(z) dz.$$

Then \tilde{K} satisfies the same conditions of K on $(x_1, x_2) \in \mathbb{R}^2$ as in Lemma 3.3 with $C_{\tilde{K}}$ no more than $C_\psi \frac{s}{(s+|x_3|)^2}$.

Proof. For any fixed $(x_1, x_2) \in \Omega_2$, by Definition 1.1, it is easy to see that $K^\sharp(x_1, x_2, \cdot)$ satisfies all the conditions of Lemma 3.2 with $C_{K^\sharp(x_1, x_2, \cdot)} \leq C \frac{1}{\|(x_1, x_2)\|^3}$. Thus, Lemma 3.2 yields that, for all $s > 0$, $(x_1, x_2) \in \Omega_2$, and $x_3 \in \mathbb{R}$,

$$|\tilde{K}(x_1, x_2, x_3)| \leq CC_\psi \frac{s}{(s + |x_3|)^2} \frac{1}{\|(x_1, x_2)\|^3},$$

which shows that for any fixed $x_3 \in \mathbb{R}$, $\tilde{K}(\cdot, \cdot, x_3)$ satisfies Lemma 3.3 (i).

We now show that \tilde{K} satisfies Lemma 3.3 (ii). Let $\|(x_1, x_2) - (y_1, y_2)\| \leq \|(x_1, x_2)\|/2$. By the mean value theorem and Definition 1.1, we have that

$$\begin{aligned} & |K^\sharp(x_1, x_2, x_3) - K^\sharp(y_1, y_2, x_3)| \\ & \leq C \frac{1}{\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\|^4 |x_3|} \\ & \quad \times \left\{ |x_1 - y_1| + \frac{|x_2 - y_2|}{\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\|} \right\} \\ & \leq C \frac{\|(x_1, x_2) - (y_1, y_2)\|}{\|(x_1, x_2)\|^4 |x_3|}, \end{aligned}$$

where $\kappa \in (0, 1)$ and we used the fact that

$$\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\| \geq \|(x_1, x_2)\|/2.$$

Similarly, if $|x_3 - y_3| \leq |x_3|/2$, we have

$$\begin{aligned} & \left| [K^\sharp(x_1, x_2, x_3) - K^\sharp(y_1, y_2, x_3)] - [K^\sharp(x_1, x_2, y_3) - K^\sharp(y_1, y_2, y_3)] \right| \\ & \leq C \frac{|x_3 - y_3|}{\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\|^4 |(1 - \kappa_0)x_3 + \kappa_0 y_3|^2} \\ & \quad \times \left\{ |x_1 - y_1| + \frac{|x_2 - y_2|}{\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\|} \right\} \\ & \leq C \frac{\|(x_1, x_2) - (y_1, y_2)\| |x_3 - y_3|}{\|(x_1, x_2)\|^4 |x_3|^2}, \end{aligned}$$

where $\kappa, \kappa_0 \in (0, 1)$. Let φ be a normalized bump function on \mathbb{R} . The mean value theorem and Definition 1.1 further yield that, for all $\delta > 0$,

$$\begin{aligned} & \left| \int_{\mathbb{R}} [K^\sharp(x_1, x_2, x_3) - K^\sharp(y_1, y_2, x_3)] \varphi(\delta x_3) dx_3 \right| \\ & \leq C \frac{1}{\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\|^4} \\ & \quad \times \left[|x_1 - y_1| + \frac{|x_2 - y_2|}{\|((1 - \kappa)x_1 + \kappa y_1, (1 - \kappa)x_2 + \kappa y_2)\|} \right] \\ & \leq C \frac{\|(x_1, x_2) - (y_1, y_2)\|}{\|(x_1, x_2)\|^4}, \end{aligned}$$

where $\kappa \in (0, 1)$. Thus, for any fixed $(x_1, x_2) \in \Omega_2$ and any $(y_1, y_2) \in \mathbb{R}^2$ with $\|(x_1, x_2) - (y_1, y_2)\| \leq \|(x_1, x_2)\|/2$, $K^\sharp(x_1, x_2, \cdot) - K^\sharp(y_1, y_2, \cdot)$ satisfies all the conditions of Lemma 3.2 with

$$C_{K^\sharp(x_1, x_2, \cdot) - K^\sharp(y_1, y_2, \cdot)} \leq C \frac{\|(x_1, x_2) - (y_1, y_2)\|}{\|(x_1, x_2)\|^4}.$$

Lemma 3.2 yields that

$$\begin{aligned} & |\tilde{K}(x_1, x_2, x_3) - \tilde{K}(y_1, y_2, x_3)| \\ & = \left| \int_{\mathbb{R}} [K^\sharp(x_1, x_2, x_3 - z) - K^\sharp(y_1, y_2, x_3 - z)] \psi_s(z) dz \right| \\ & \leq C C_\psi \frac{s}{(s + |x_3|)^2} \cdot \frac{\|(x_1, x_2) - (y_1, y_2)\|}{\|(x_1, x_2)\|^4}, \end{aligned}$$

which shows that, for any fixed $x_3 \in \mathbb{R}$, $\tilde{K}(\cdot, \cdot, x_3)$ satisfies Lemma 3.3 (ii).

Finally, let φ be a normalized bump function on \mathbb{R}^2 and $\delta > 0$. By the mean value theorem and Definition 1.1, we can easily show that

$$\int_{\mathbb{R}^2} K^\sharp(x_1, x_2, \cdot) \varphi(\delta x_1, \delta^2 x_2) dx_1 dx_2$$

satisfies all the conditions of Lemma 3.2 and hence

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} \tilde{K}(x_1, x_2, x_3) \varphi(\delta x_1, \delta^2 x_2) dx_1 dx_2 \right| \\ &= \left| \int_{\mathbb{R}} \left[\int_{\mathbb{R}^2} K^\sharp(x_1, x_2, x_3 - z) \varphi(\delta x_1, \delta^2 x_2) dx_1 dx_2 \right] \psi_s(z) dz \right| \\ &\leq CC_\psi \frac{s}{(s + |x_3|)^2}, \end{aligned}$$

which yields that, for any fixed $x_3 \in \mathbb{R}$, $\tilde{K}(\cdot, \cdot, x_3)$ satisfies Lemma 3.3 (iii). This finishes the proof of Lemma 3.6. \square

Proof of Theorem 3.1. By Proposition 1.4, there exists an integrable product kernel K^\sharp on $\mathbb{R}^2 \times \mathbb{R}$ such that for all $(x_1, x_2) \in \mathbb{R}^2$,

$$K(x_1, x_2) = \int_{\mathbb{R}} K^\sharp(x_1, x_2 - x_3, x_3) dx_3.$$

Let $\psi_{t_1 t_2}$ be the same as in Lemma 2.3; moreover, let

$$\text{supp } \psi^{(1)} \subset \{ (x_1, x_2) : \|(x_1, x_2)\| < 1 \}$$

and $\text{supp } \psi^{(2)} \subset (-1, 1)$. By Lemma 2.3, for $u_1, u_2 > 0$ and $(x_1, x_2) \in \mathbb{R}^2$, we have

$$\begin{aligned} & (\psi_{u_1 u_2} * K * f)(x_1, x_2) \\ &= \int_0^\infty \int_0^\infty (\psi_{u_1 u_2} * K * \psi_{t_1 t_2} * \psi_{t_1 t_2} * f)(x_1, x_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2} \quad (23) \end{aligned}$$

and

$$\begin{aligned} & \psi_{u_1 u_2} * K * \psi_{t_1 t_2}(x_1, x_2) \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^2} (\psi_{u_1}^{(1)} * \psi_{t_1}^{(1)})(x_1 - y_1, x_2 - x_3 - y_2) \\ &\quad \times \left[\int_{\mathbb{R}} K^\sharp(y_1, y_2, x_3 - y_3) (\psi_{u_2}^{(2)} * \psi_{t_2}^{(2)})(y_3) dy_3 \right] dy_1 dy_2 dx_3. \end{aligned}$$

Lemma 3.5 implies that $\psi_{u_2}^{(2)} * \psi_{t_2}^{(2)}$ satisfies all the conditions of Lemma 3.2 with $s = u_2 \vee t_2$ and

$$C_{\psi_{u_2}^{(2)} * \psi_{t_2}^{(2)}} \leq C \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right),$$

which together with Lemma 3.6 shows that

$$\tilde{K}(y_1, y_2, x_3) = \int_{\mathbb{R}} K^\sharp(y_1, y_2, x_3 - y_3) (\psi_{u_2}^{(2)} * \psi_{t_2}^{(2)})(y_3) dy_3$$

satisfies all the conditions of Lemma 3.3 for any fixed $x_3 \in \mathbb{R}$ with

$$C_{\bar{K}} \leq C \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |x_3|)^2}.$$

From this, Lemma 3.5, and Lemma 3.3, it follows that

$$\begin{aligned} & |(\psi_{u_1 u_2} * K * \psi_{t_1 t_2})(x_1, x_2)| \\ & \leq C \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \int_{\mathbb{R}} \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(x_1, x_2 - x_3)\|)^4} \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |x_3|)^2} dx_3, \end{aligned}$$

which together with the estimate (11) yields that

$$\begin{aligned} & |(\psi_{u_1 u_2} * K * \psi_{t_1 t_2})(x_1, x_2)| \\ & \leq C \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \\ & \quad \times \left\{ \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(x_1, x_2)\|)^4} + \frac{u_1 \vee t_1}{(u_1 \vee t_1 + |x_1|)^2} \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |x_2|)^2} \right\}. \end{aligned}$$

Inserting the last estimate into (23) yields that

$$\begin{aligned} & |(\psi_{u_1 u_2} * K * f)(x_1, x_2)| \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \\ & \quad \times \{ M(\psi_{t_1 t_2} * f)(x_1, x_2) + M_s(\psi_{t_1 t_2} * f)(x_1, x_2) \} \frac{dt_1}{t_1} \frac{dt_2}{t_2}. \quad (24) \end{aligned}$$

Replacing (12) with (24) in the proof of Theorem 2.6 and repeating the proof there, we complete the proof of Theorem 3.1 and we omit the details. \square

4. Lifting properties

In this section, we use Theorem 3.1 in the last section to establish the lifting property of $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$. First, we introduce the following Riesz potential operators related to flag singular integrals.

Definition 4.1. Let $\{\psi_{l_1 l_2}\}_{l_1, l_2 \in \mathbb{Z}}$ be the same as in Lemma 2.4 and $(\alpha_1, \alpha_2) \in \mathbb{R} \times \mathbb{R}$. Then the Riesz potential operator $I_{(\alpha_1, \alpha_2)}$ for $f \in \mathcal{S}_{\infty, F}(\mathbb{R}^2)'$ is defined by

$$I_{(\alpha_1, \alpha_2)}(f)(x_1, x_2) = \sum_{l_1=-\infty}^\infty \sum_{l_2=-\infty}^\infty 2^{-l_1 \alpha_1} 2^{-l_2 \alpha_2} (\psi_{l_1 l_2} * f)(x_1, x_2),$$

where $(x_1, x_2) \in \mathbb{R}^2$.

From Definition 4.1, it is easy to deduce the following simple property of Riesz potential operators.

Proposition 4.2. *Let $\{\psi_{l_1}^{(1)}\}_{l_1 \in \mathbb{Z}}$ and $\{\psi_{l_2}^{(2)}\}_{l_2 \in \mathbb{Z}}$ be the same as in Lemma 2.4. For $f \in \mathcal{S}(\mathbb{R}^2)'$ and $(x_1, x_2) \in \mathbb{R}^2$, let*

$$I_{\alpha_1}^{(1)}(f)(x_1, x_2) = \sum_{l_1=-\infty}^{\infty} 2^{-l_1 \alpha_1} (\psi_{l_1}^{(1)} * f)(x_1, x_2),$$

and for $f \in \mathcal{S}(\mathbb{R})'$ and $x_3 \in \mathbb{R}$, let

$$I_{\alpha_2}^{(2)}(f)(x_3) = \sum_{l_2=-\infty}^{\infty} 2^{-l_2 \alpha_2} (\psi_{l_2}^{(2)} * f)(x_3).$$

Then

$$I_{(\alpha_1, \alpha_2)} = I_{\alpha_1}^{(1)} I_{\alpha_2}^{(2)} = I_{\alpha_2}^{(2)} I_{\alpha_1}^{(1)}.$$

One of the main theorems in this section is the following boundedness of $I_{(\alpha_1, \alpha_2)}$ on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ as below.

Proposition 4.3. *Let $|s_i| < 1$, $|\alpha_i| < 1$, $|s_i + \alpha_i| < 1$ for $i = 1, 2$, $s = (s_1, s_2)$ and $s + \alpha = (s_1 + \alpha_1, s_2 + \alpha_2)$. Then*

- (i) *If $p, q \in [1, \infty]$, $I_{(\alpha_1, \alpha_2)}$ is bounded from $\dot{B}_{pq}^s(\mathbb{R}^2)$ to $\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)$, namely, there exists a positive constant C such that for all $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$,*

$$\|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}.$$

- (ii) *If $p \in (1, \infty)$ and $q \in (1, \infty]$, $I_{(\alpha_1, \alpha_2)}$ is bounded from $\dot{F}_{pq}^s(\mathbb{R}^2)$ to $\dot{F}_{pq}^{s+\alpha}(\mathbb{R}^2)$, namely, there exists a positive constant C such that for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$,*

$$\|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{F}_{pq}^{s+\alpha}(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}.$$

To prove Proposition 4.3, we first establish the following basic estimate. In what follows, for any $a, b \in \mathbb{R}$, let $a \wedge b = \min\{a, b\}$.

Lemma 4.4. *For $i = 1, 2$, let $I_{\alpha_i}^{(i)}$ be the same as in Proposition 4.2 and $\{\tilde{\psi}_{k_i}^{(i)}\}_{k_i \in \mathbb{Z}}$ be the same as in Lemma 2.4 with $\text{supp } \tilde{\psi}^{(1)} \subset \{(x_1, x_2) \in \mathbb{R}^2 : \|(x_1, x_2)\| \leq 1\}$ and*

$$\text{supp } \tilde{\psi}^{(2)} \subset \{x_3 \in \mathbb{R} : |x_3| \leq 1\}.$$

Then,

- (i) For $|\alpha_1| < 1$ and any $\epsilon_1 > 0$, there exists a positive constant C_{ϵ_1, α_1} such that for all $k_1, j_1 \in \mathbb{Z}$ and all $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} & |\tilde{\psi}_{k_1}^{(1)} * I_{\alpha_1}^{(1)} * \tilde{\psi}_{j_1}^{(1)}(x_1, x_2)| \\ & \leq C_{\epsilon_1, \alpha_1} (1 + |k_1 - j_1|) 2^{-(k_1 \wedge j_1) \alpha_1} 2^{-|k_1 - j_1| (1 + \alpha_1 \wedge 0)} \\ & \quad \times \frac{2^{-(k_1 \wedge j_1)(1 - \alpha_1 - \epsilon_1)}}{(2^{-(k_1 \wedge j_1)} + \|(x_1, x_2)\|)^{4 - \alpha_1 - \epsilon_1}}. \end{aligned}$$

- (ii) For $|\alpha_2| < 1$ and any $\epsilon_2 > 0$, there exists a positive constant C_{ϵ_2, α_2} such that for all $k_2, j_2 \in \mathbb{Z}$ and all $x_3 \in \mathbb{R}$,

$$\begin{aligned} & |\tilde{\psi}_{k_2}^{(2)} * I_{\alpha_2}^{(2)} * \tilde{\psi}_{j_2}^{(2)}(x_3)| \\ & \leq C_{\epsilon_2, \alpha_2} (1 + |k_2 - j_2|) 2^{-(k_2 \wedge j_2) \alpha_2} 2^{-|k_2 - j_2| (1 + \alpha_2 \wedge 0)} \\ & \quad \times \frac{2^{-(k_2 \wedge j_2)(1 - \alpha_2 - \epsilon_2)}}{(2^{-(k_2 \wedge j_2)} + |x_3|)^{2 - \alpha_2 - \epsilon_2}}. \end{aligned}$$

Proof. By similarity, we only show (i). Without loss of generality, we may further assume that $j_1 \geq k_1$. In this case, we write

$$\begin{aligned} & \tilde{\psi}_{k_1}^{(1)} * I_{\alpha_1}^{(1)} * \tilde{\psi}_{j_1}^{(1)}(x_1, x_2) \\ & = \sum_{l_1=-\infty}^{k_1} 2^{-l_1 \alpha_1} \tilde{\psi}_{k_1}^{(1)} * \psi_{l_1}^{(1)} * \tilde{\psi}_{j_1}^{(1)}(x_1, x_2) + \sum_{l_1=k_1+1}^{j_1} \dots + \sum_{l_1=j_1+1}^{\infty} \dots \\ & = O_1 + O_2 + O_3. \end{aligned}$$

We now consider two cases.

- *Case 1:* $\|(x_1, x_2)\| \geq 5 \cdot 2^{-k_1}$. In this case, by the vanishing moment of $\tilde{\psi}^{(1)}$ and the mean value theorem,

$$\begin{aligned} |O_1| & = \left| \sum_{l_1=-\infty}^{k_1} 2^{-l_1 \alpha_1} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \tilde{\psi}_{k_1}^{(1)}(x_1 - y_1, x_2 - y_2) \right. \\ & \quad \left. \times [\psi_{l_1}^{(1)}(y_1 - u_1, y_2 - u_2) - \psi_{l_1}^{(1)}(y_1, y_2)] \tilde{\psi}_{j_1}^{(1)}(u_1, u_2) dy_1 dy_2 du_1 du_2 \right| \leq \end{aligned}$$

$$\begin{aligned}
 &\leq \sum_{l_1=-\infty}^{k_1} 2^{-l_1\alpha_1} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} |\tilde{\psi}_{k_1}^{(1)}(y_1, y_2)| \\
 &\quad \times \left[\left| \frac{\partial \psi_{l_1}^{(1)}}{\partial y_1}(x_1 - y_1 - \theta u_1, x_2 - y_2 - \theta u_2) \right| |u_1| \right. \\
 &\quad \left. + \left| \frac{\partial \psi_{l_1}^{(1)}}{\partial y_2}(x_1 - y_1 - \theta u_1, x_2 - y_2 - \theta u_2) \right| |u_2| \right] \\
 &\quad \times |\tilde{\psi}_{j_1}^{(1)}(u_1, u_2)| dy_1 dy_2 du_1 du_2, \tag{25}
 \end{aligned}$$

where $\theta \in (0, 1)$. The support condition of $\tilde{\psi}^{(1)}$ yields that

$$\begin{aligned}
 \|(x_1 - y_1 - \theta u_1, x_2 - y_2 - \theta u_2)\| &\geq \|(x_1, x_2)\| - \|(y_1, y_2)\| - \|(u_1, u_2)\| \\
 &\geq \|(x_1, x_2)\|/2.
 \end{aligned}$$

From this, it follows that $|O_1|$ is further controlled by

$$|O_1| \leq C \sum_{l_1=-\infty}^{k_1} 2^{l_1\epsilon_1} 2^{-j_1} \frac{1}{\|(x_1, x_2)\|^{4-\alpha_1-\epsilon_1}} \leq C 2^{-k_1\alpha_1} 2^{k_1-j_1} \frac{2^{-k_1(1-\alpha_1-\epsilon_1)}}{\|(x_1, x_2)\|^{4-\alpha_1-\epsilon_1}}.$$

Similarly, for O_2 , we have

$$\begin{aligned}
 |O_2| &= \left| \sum_{l_1=k_1+1}^{j_1} 2^{-l_1\alpha_1} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \psi_{l_1}^{(1)}(x_1 - y_1, x_2 - y_2) \right. \\
 &\quad \left. \times [\tilde{\psi}_{k_1}^{(1)}(y_1 - u_1, y_2 - u_2) - \tilde{\psi}_{k_1}^{(1)}(y_1, y_2)] \tilde{\psi}_{j_1}^{(1)}(u_1, u_2) dy_1 dy_2 du_1 du_2 \right| \\
 &\leq C 2^{k_1-j_1} \frac{1}{\|(x_1, x_2)\|^{4-\alpha_1}} \sum_{l_1=k_1+1}^{j_1} 2^{-l_1} = C 2^{-k_1\alpha_1} 2^{k_1-j_1} \frac{2^{-k_1(1-\alpha_1)}}{\|(x_1, x_2)\|^{4-\alpha_1}}, \tag{26}
 \end{aligned}$$

since

$$\|(x_1 - y_1, x_2 - y_2)\| \geq \|(x_1, x_2)\| - \|(y_1, y_2)\| \geq \|(x_1, x_2)\|/2.$$

For O_3 , by the vanishing moment of $\psi^{(1)}$ and the mean value theorem, we have

$$\begin{aligned}
 |O_3| &= \left| \sum_{l_1=j_1+1}^{\infty} 2^{-l_1\alpha_1} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} [\tilde{\psi}_{k_1}^{(1)}(u_1 - y_1, u_2 - y_2) - \tilde{\psi}_{k_1}^{(1)}(u_1, u_2)] \right. \\
 &\quad \left. \times \psi_{l_1}^{(1)}(y_1, y_2) \tilde{\psi}_{j_1}^{(1)}(x_1 - u_1, x_2 - u_2) du_1 du_2 dy_1 dy_2 \right| \\
 &\leq \sum_{l_1=j_1+1}^{\infty} 2^{-l_1\alpha_1} \int_{\mathbb{R}^2} \int_{\|(u_1, u_2)\| \leq 2^{-k_1}} |\tilde{\psi}_{k_1}^{(1)}(u_1 - y_1, u_2 - y_2) - \tilde{\psi}_{k_1}^{(1)}(u_1, u_2)| \\
 &\quad \times |\psi_{l_1}^{(1)}(y_1, y_2) \tilde{\psi}_{j_1}^{(1)}(x_1 - u_1, x_2 - u_2)| du_1 du_2 dy_1 dy_2 + \sum_{l_1=j_1+1}^{\infty} 2^{-l_1\alpha_1} \\
 &\quad \times \int_{\mathbb{R}^2} \int_{\substack{\|(u_1, u_2)\| > 2^{-k_1} \\ \|(u_1 - y_1, u_2 - y_2)\| \leq 2^{-k_1}}} |\tilde{\psi}_{k_1}^{(1)}(u_1 - y_1, u_2 - y_2) - \tilde{\psi}_{k_1}^{(1)}(u_1, u_2)| \\
 &\quad \times |\psi_{l_1}^{(1)}(y_1, y_2) \tilde{\psi}_{j_1}^{(1)}(x_1 - u_1, x_2 - u_2)| du_1 du_2 dy_1 dy_2 = O_3^1 + O_3^2. \quad (27)
 \end{aligned}$$

For O_3^1 , we have

$$\|(x_1 - u_1, x_2 - u_2)\| \geq \|(x_1, x_2)\| - \|(u_1, u_2)\| \geq \|(x_1, x_2)\|/2;$$

and, for O_3^2 , we have

$$\|(y_1, y_2)\| \geq \|(x_1, x_2)\| - \|(y_1 - u_1, y_2 - u_2)\| - \|(u_1 - x_1, u_2 - x_2)\| \geq \|(x_1, x_2)\|/2.$$

These facts, respectively, imply that

$$\begin{aligned}
 O_3^1 &\leq C \frac{1}{\|(x_1, x_2)\|^{4-\alpha_1}} 2^{k_1-j_1(1-\alpha_1)} \sum_{l_1=j_1+1}^{\infty} 2^{-l_1(1+\alpha_1)} \\
 &\leq C 2^{-k_1\alpha_1} 2^{2(k_1-j_1)} \frac{2^{-k_1(1-\alpha_1)}}{\|(x_1, x_2)\|^{4-\alpha_1}}
 \end{aligned}$$

and

$$\begin{aligned}
 O_3^2 &\leq C \frac{1}{\|(x_1, x_2)\|^{4-\alpha_1}} 2^{k_1} \sum_{l_1=j_1+1}^{\infty} 2^{-2l_1} \\
 &\leq C 2^{-k_1\alpha_1} 2^{2(k_1-j_1)} \frac{2^{-k_1(1-\alpha_1)}}{\|(x_1, x_2)\|^{4-\alpha_1}}.
 \end{aligned}$$

This finishes the proof of case 1.

- *Case 2:* $\|(x_1, x_2)\| < 5 \cdot 2^{-k_1}$. In this case, by (25), we have

$$|O_1| \leq C \sum_{l_1=-\infty}^{k_1} 2^{4l_1-l_1\alpha_1-j_1} = C 2^{-k_1\alpha_1} 2^{k_1-j_1} 2^{3k_1};$$

by (26), we obtain

$$|O_2| \leq C \sum_{l_1=k_1+1}^{j_1} 2^{4k_1-l_1\alpha_1-j_1} = \begin{cases} (j_1 - k_1)2^{k_1-j_1}2^{3k_1}, & \alpha_1 = 0, \\ C2^{-k_1\alpha_1}2^{k_1-j_1}2^{3k_1}, & \alpha_1 > 0, \\ C2^{-k_1\alpha_1}2^{(k_1-j_1)(1+\alpha_1)}2^{3k_1}, & \alpha_1 < 0; \end{cases}$$

and, finally by (27), we have

$$|O_3| \leq C \sum_{l_1=j_1+1}^{\infty} 2^{4k_1-l_1\alpha_1-l_1} = C2^{-k_1\alpha_1}2^{(k_1-j_1)(1+\alpha_1)}2^{3k_1},$$

which are desired estimates. This finishes the proof of case 2 and, hence, the proof of Lemma 4.4. \square

Proof of Proposition 4.3. Let $\{\tilde{\psi}_{k_i}^{(i)}\}_{k_i \in \mathbb{Z}}$ with $i = 1, 2$ and $\{\psi_{l_1 l_2}\}_{l_1, l_2 \in \mathbb{Z}}$ be the same, respectively, as in Lemma 4.4 and Lemma 2.4. By Theorem 2.8 and Lemma 2.4, we have

$$\begin{aligned} & \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)} \\ & \leq C \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{k_1(s_1+\alpha_1)q} 2^{k_2(s_2+\alpha_2)q} \right. \\ & \quad \left. \times \left\| \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \tilde{\psi}_{k_1 k_2} * I_{(\alpha_1, \alpha_2)} * \tilde{\psi}_{j_1 j_2} * \tilde{\psi}_{j_1 j_2} * f \right\|_{L^p(\mathbb{R}^2)}^q \right\}^{1/q}. \end{aligned}$$

Lemma 4.4 further yields that for all $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} & |\tilde{\psi}_{k_1 k_2} * I_{(\alpha_1, \alpha_2)} * \tilde{\psi}_{j_1 j_2}(x_1, x_2)| \\ & \leq C 2^{-(k_1 \wedge j_1)\alpha_1 - |k_1 - j_1|(1+\alpha_1 \wedge 0) - (k_2 \wedge j_2)\alpha_2 - |k_2 - j_2|(1+\alpha_2 \wedge 0)} \\ & \quad \times (1 + |k_1 - j_1|)(1 + |k_2 - j_2|) \\ & \quad \times \left\{ \frac{2^{-(k_1 \wedge j_1)(1-\alpha_1-\epsilon_1)}}{(2^{-(k_1 \wedge j_1)} + \|(x_1, x_2)\|)^{4-\alpha_1-\epsilon_1}} \right. \\ & \quad \left. + \frac{2^{-(k_1 \wedge j_1)(1-\alpha_1-\epsilon_1)}}{(2^{-(k_1 \wedge j_1)} + |x_1|)^{2-\alpha_1-\epsilon_1}} \frac{2^{-(k_2 \wedge j_2)(1-\alpha_2-\epsilon_2)}}{(2^{-(k_2 \wedge j_2)} + |x_2|)^{2-\alpha_2-\epsilon_2}} \right\}. \quad (28) \end{aligned}$$

From this, the Minkowski inequality, and the boundedness of M_s in $L^p(\mathbb{R}^2)$, it follows

that, for $p \in (1, \infty]$,

$$\begin{aligned} & \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)} \\ & \leq C \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{k_1(s_1+\alpha_1)q} 2^{k_2(s_2+\alpha_2)q} \right. \\ & \quad \times \left[\sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} 2^{-(k_1 \wedge j_1)\alpha_1 - |k_1 - j_1|(1+\alpha_1 \wedge 0) - (k_2 \wedge j_2)\alpha_2 - |k_2 - j_2|(1+\alpha_2 \wedge 0)} \right. \\ & \quad \left. \left. \times (1 + |k_1 - j_1|)(1 + |k_2 - j_2|) \|\tilde{\psi}_{j_1 j_2} * f\|_{L^p(\mathbb{R}^2)} \right]^q \right\}^{1/q}. \end{aligned} \tag{29}$$

If $p = 1$, by the Minkowski inequality and the Fubini theorem, we also obtain the same estimate. Now the assumptions that $|s_i| < 1$ and $|s_i + \alpha_i| < 1$ imply that

$$\sup_{k_i \in \mathbb{Z}} \sum_{j_i=-\infty}^{\infty} (1 + |k_i - j_i|) 2^{-(k_i \wedge j_i)\alpha_i - |k_i - j_i|(1+\alpha_i \wedge 0) + (k_i - j_i)s_i + k_i \alpha_i} < \infty \tag{30}$$

and

$$\sup_{j_i \in \mathbb{Z}} \sum_{k_i=-\infty}^{\infty} (1 + |k_i - j_i|) 2^{-(k_i \wedge j_i)\alpha_i - |k_i - j_i|(1+\alpha_i \wedge 0) + (k_i - j_i)s_i + k_i \alpha_i} < \infty, \tag{31}$$

where $i = 1, 2$. Combining these estimates (30) and (31) with (29) and using the Hölder inequality yield that

$$\begin{aligned} \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)} & \leq C \left\{ \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} 2^{j_1 s_1 q} 2^{j_2 s_2 q} \|\tilde{\psi}_{j_1 j_2} * f\|_{L^p(\mathbb{R}^2)}^q \right\}^{1/q} \\ & \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}, \end{aligned}$$

which completes the proof of Proposition 4.3 (i).

To prove Proposition 4.3 (ii), by Theorem 2.8, Lemma 2.4, the estimates (28), (30), and (31), and the Hölder inequality, we obtain

$$\begin{aligned} & \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{F}_{pq}^{s+\alpha}(\mathbb{R}^2)} \\ & \leq C \left\| \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{k_1(s_1+\alpha_1)q} 2^{k_2(s_2+\alpha_2)q} |\tilde{\psi}_{k_1 k_2} * I_{(\alpha_1, \alpha_2)}(f)|^q \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)} \leq \end{aligned}$$

$$\begin{aligned} &\leq C \left\| \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{k_1(s_1+\alpha_1)q} 2^{k_2(s_2+\alpha_2)q} \right. \right. \\ &\quad \times \left. \left[\sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} 2^{-(k_1 \wedge j_1)\alpha_1 - |k_1 - j_1|(1+\alpha_1 \wedge 0) - (k_2 \wedge j_2)\alpha_2 - |k_2 - j_2|(1+\alpha_2 \wedge 0)} \right. \right. \\ &\quad \left. \left. \times (1 + |k_1 - j_1|)(1 + |k_2 - j_2|) M_s(\tilde{\psi}_{j_1 j_2} * f) \right] \right\}^{1/q} \Big\|_{L^p(\mathbb{R}^2)} \\ &\leq C \left\| \left\{ \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} 2^{j_1 s_1 q} 2^{j_2 s_2 q} |\tilde{\psi}_{j_1 j_2} * f|^q \right\}^{1/q} \right\|_{L^p(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}, \end{aligned}$$

where, in the second-to-last inequality, we used the vector-valued inequality of Fefferman-Stein in [1]. This finishes the proof of Proposition 4.3. \square

We now establish the converse of Proposition 4.3.

Proposition 4.5. *Let $|s_i| < 1$, $|s_i + \alpha_i| < 1$ for $i = 1, 2$, $s = (s_1, s_2)$, and $s + \alpha = (s_1 + \alpha_1, s_2 + \alpha_2)$. Then there exist a positive constant C and $\alpha_i^0(s_1, s_2) \in (0, 1)$ such that, if $|\alpha_i| < \alpha_i^0(s_1, s_2)$ with $i = 1, 2$,*

$$\|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq C \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)}$$

for all $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$, and for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$,

$$\|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{F}_{pq}^{s+\alpha}(\mathbb{R}^2)}.$$

Proof. The key of the proof is to show that the operator $I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)}$ is invertible in $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ if α_1 and α_2 are small. To this end, we need to prove that the operator $I - I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)}$ is bounded on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ with operator norms less than 1 when α_1 and α_2 are small, where I is the identity operator. We obtain this by using Theorem 3.1. Let $\{\psi_{k_1 k_2}\}_{k_1, k_2 \in \mathbb{Z}}$ be the same as in Lemma 2.4. We write

$$I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)} = \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} 2^{-k_1 \alpha_1} 2^{-k_2 \alpha_2} \psi_{j_1 j_2} * \psi_{k_1+j_1, k_2+j_2}$$

and

$$\begin{aligned} I - I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)} &= \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} (1 - 2^{-k_1 \alpha_1} 2^{-k_2 \alpha_2}) \psi_{j_1 j_2} * \psi_{k_1+j_1, k_2+j_2}. \end{aligned}$$

We denote the kernel of $I - I_{(-\alpha_1, -\alpha_2)}I_{(\alpha_1, \alpha_2)}$ simply by $K_{(\alpha_1, \alpha_2)}$. Noticing that, for $(x_1, x_2) \in \mathbb{R}^2$,

$$\psi_{j_1 j_2} * \psi_{k_1+j_1, k_2+j_2}(x_1, x_2) = (\psi_{j_1}^{(1)} * \psi_{k_1+j_1}^{(1)}) *_2 (\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)})(x_1, x_2),$$

we then have that

$$K_{(\alpha_1, \alpha_2)}^\sharp(x_1, x_2, x_3) = \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} (1 - 2^{-k_1 \alpha_1} 2^{-k_2 \alpha_2}) \times (\psi_{j_1}^{(1)} * \psi_{k_1+j_1}^{(1)})(x_1, x_2) (\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)})(x_3),$$

where $(x_1, x_2, x_3) \in \mathbb{R}^2 \times \mathbb{R}$, is the corresponding product kernel on $\mathbb{R}^2 \times \mathbb{R}$ of $K_{(\alpha_1, \alpha_2)}$. We only need to show that $K_{(\alpha_1, \alpha_2)}^\sharp$ satisfies the conditions of Definition 1.1 with a constant no more than

$$C \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1} 2^{-k_2 \alpha_2}| 2^{-|k_1|} 2^{-|k_2|},$$

where C is a positive constant independent of α_1 and α_2 . First, we point out that by a modified argument of (7), we can easily to show that for $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{Z}_+$, $(x_1, x_2) \in \mathbb{R}^2$, and $x_3 \in \mathbb{R}$,

$$|\partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} (\psi_{j_1}^{(1)} * \psi_{k_1+j_1}^{(1)})(x_1, x_2)| \leq C_{\gamma_1, \gamma_2} 2^{-|k_1|} \frac{2^{-j_1 \wedge (k_1+j_1)}}{(2^{-j_1 \wedge (k_1+j_1)} + \|(x_1, x_2)\|)^{4+\gamma_1+2\gamma_2}} \quad (32)$$

and

$$|\partial_{x_3}^{\gamma_3} (\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)})(x_3)| \leq C_{\gamma_3} 2^{-|k_2|} \frac{2^{-j_2 \wedge (k_2+j_2)}}{(2^{-j_2 \wedge (k_2+j_2)} + |x_3|)^{2+\gamma_3}}, \quad (33)$$

by noticing that $\psi^{(i)}$ for $i = 1, 2$ are the Schwartz functions. In fact, we only need that these estimates are true for $\gamma_1 = \gamma_2 = \gamma_3 = 1$. The estimates (32) and (33) imply that for $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{Z}_+$, $(x_1, x_2) \in \mathbb{R}^2$, and $x_3 \in \mathbb{R}$,

$$\begin{aligned} & |\partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} \partial_{x_3}^{\gamma_3} K_{(\alpha_1, \alpha_2)}^\sharp(x_1, x_2, x_3)| \\ & \leq \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1} 2^{-k_2 \alpha_2}| \\ & \quad \times |\partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} (\psi_{j_1}^{(1)} * \psi_{k_1+j_1}^{(1)})(x_1, x_2) \partial_{x_3}^{\gamma_3} (\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)})(x_3)| \\ & \leq C \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1} 2^{-k_2 \alpha_2}| 2^{-|k_1| - |k_2|} \right\} \\ & \quad \times \frac{1}{\|(x_1, x_2)\|^{3+\gamma_1+2\gamma_2}} \cdot \frac{1}{|x_3|^{1+\gamma_3}}. \end{aligned} \quad (34)$$

Let φ be a normalized bump function on \mathbb{R} and $\delta > 0$. We now estimate, by (32) and (33), that, for $\gamma_1, \gamma_2 \in \mathbb{Z}_+$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} & \left| \int_{\mathbb{R}} \partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} K_{(\alpha_1, \alpha_2)}^\#(x_1, x_2, x_3) \varphi(\delta x_3) dx_3 \right| \\ & \leq \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| \\ & \quad \times \left| \int_{\mathbb{R}} \partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} \left(\psi_{j_1}^{(1)} * \psi_{k_1+j_1}^{(1)} \right) (x_1, x_2) \left(\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)} \right) (x_3) \varphi(\delta x_3) dx_3 \right| \\ & \leq C \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{\infty} \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| 2^{-|k_1|} \\ & \quad \frac{2^{-j_1 \wedge (k_1+j_1)}}{(2^{-j_1 \wedge (k_1+j_1)} + \|(x_1, x_2)\|)^{4+\gamma_1+2\gamma_2}} \\ & \quad \times \left| \int_{\mathbb{R}} \left(\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)} \right) (x_3) \varphi(\delta x_3) dx_3 \right|. \end{aligned}$$

We choose $j_2^0 \in \mathbb{Z}$ such that $\delta \leq 2^{-j_2^0 \wedge (k_2+j_2^0)} < 2\delta$. For this j_2^0 and any $\epsilon > 0$, by the vanishing moment of $\psi^{(2)} * \psi^{(2)}$, we have

$$\begin{aligned} & \sum_{j_2=-\infty}^{\infty} \left| \int_{\mathbb{R}} \left(\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)} \right) (x_3) \varphi(\delta x_3) dx_3 \right| \\ & \leq C \sum_{j_2=-\infty}^{j_2^0} 2^{-|k_2|} \int_{\mathbb{R}} \frac{2^{-j_2 \wedge (k_2+j_2)}}{(2^{-j_2 \wedge (k_2+j_2)} + |x_3|)^2} |\varphi(\delta x_3)| dx_3 \\ & \quad + \sum_{j_2=j_2^0+1}^{\infty} \left| \int_{\mathbb{R}} \left(\psi_{j_2}^{(2)} * \psi_{k_2+j_2}^{(2)} \right) (x_3) [\varphi(\delta x_3) - \varphi(0)] dx_3 \right| \\ & \leq C 2^{-|k_2|} \left\{ 1 + \delta^\epsilon \sum_{j_2=j_2^0+1}^{\infty} 2^{-j_2 \wedge (k_2+j_2)\epsilon} \right\} \leq C 2^{-|k_2|}, \end{aligned}$$

which yields that

$$\begin{aligned} & \left| \int_{\mathbb{R}} \partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} K_{(\alpha_1, \alpha_2)}^\#(x_1, x_2, x_3) \varphi(\delta x_3) dx_3 \right| \\ & \leq C \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| 2^{-|k_1| - |k_2|} \right\} \frac{1}{\|(x_1, x_2)\|^{3+\gamma_1+2\gamma_2}}, \quad (35) \end{aligned}$$

where C is a positive constant independent of $(x_1, x_2) \in \mathbb{R}^2$ and $\alpha_1, \alpha_2 \in \mathbb{Z}_+$. Similarly, we can show that for all normalized bump function φ on \mathbb{R}^2 , $\delta > 0$,

and all $x_3 \in \mathbb{R}$,

$$\left| \int_{\mathbb{R}^2} \partial_{x_3}^{\gamma_3} K_{(\alpha_1, \alpha_2)}^\sharp(x_1, x_2, x_3) \varphi(\delta x_1, \delta^2 x_2) dx_1 dx_2 \right| \leq C \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| 2^{-|k_1| - |k_2|} \right\} \frac{1}{|x_3|^{1+\gamma_3}}, \quad (36)$$

where C is a positive constant independent of $x_3 \in \mathbb{R}$ and $\alpha_1, \alpha_2 \in \mathbb{Z}_+$. The estimates (34) and (35), and the special structure of $K_{(\alpha_1, \alpha_2)}^\sharp$ imply that for all normalized bump functions φ_1 and φ_2 , respectively, on \mathbb{R}^2 and \mathbb{R} , and all $\delta_1, \delta_2 > 0$,

$$\left| \int_{\mathbb{R}^2 \times \mathbb{R}} K_{(\alpha_1, \alpha_2)}^\sharp(x_1, x_2, x_3) \varphi_1(\delta_1 x_1, \delta_1^2 x_2) \varphi_2(\delta_2 x_2) dx_1 dx_2 dx_3 \right| \leq C \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| 2^{-|k_1| - |k_2|} \right\} \quad (37)$$

with the positive constant C independent of $\alpha_1, \alpha_2 \in \mathbb{Z}_+$. Thus, the estimates (34), (35), (36), and (37) and Remark 1.3 imply that the kernel $K_{(\alpha_1, \alpha_2)}^\sharp$ is a product kernel on $\mathbb{R}^2 \times \mathbb{R}$ with a constant no more than

$$C_0 \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| 2^{-|k_1| - |k_2|} \right\}, \quad (38)$$

where C_0 is a positive constant independent of $\alpha_1, \alpha_2 \in \mathbb{Z}_+$. Now, Theorem 3.1 and its proof imply that $I - I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)}$ is bounded on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ with operator norms no more than the quantity in (38). It is easy to see that we can choose $\alpha_1^0(s_1, s_2) > 0$ and $\alpha_2^0(s_1, s_2) > 0$ so small that if $|\alpha_1| < \alpha_1^0(s_1, s_2)$ and $|\alpha_2| < \alpha_2^0(s_1, s_2)$, then

$$C_0 \left\{ \sum_{k_1=-\infty}^{\infty} \sum_{k_2=-\infty}^{\infty} |1 - 2^{-k_1 \alpha_1 - k_2 \alpha_2}| 2^{-|k_1| - |k_2|} \right\} < 1,$$

where C_0 is the same positive constant as in (38). Thus, under this restriction, we know that $(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})^{-1}$ exists and is bounded, respectively, on $\dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$. Namely, there exists a positive constant C such that

$$\|(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})^{-1}(f)\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$$

for all $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$, and for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$,

$$\|(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})^{-1}(f)\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}.$$

Combining these with Proposition 4.3 yields that, if $|\alpha_1| < \alpha_1^0(s_1, s_2)$ and $|\alpha_2| < \alpha_2^0(s_1, s_2)$, then there exists a positive constant C such that

$$\begin{aligned} \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} &= \|(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})^{-1} (I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})(f)\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \\ &\leq C \|(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})(f)\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq C \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)}, \end{aligned}$$

for all $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$, and for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$,

$$\begin{aligned} \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} &= \|(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})^{-1} (I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})(f)\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \\ &\leq C \|(I_{(-\alpha_1, -\alpha_2)} I_{(\alpha_1, \alpha_2)})(f)\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq C \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{F}_{pq}^{s+\alpha}(\mathbb{R}^2)}, \end{aligned}$$

which completes the proof of Proposition 4.5. □

Combining Proposition 4.3 with Proposition 4.5 yields the following lifting properties of $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$.

Theorem 4.6. *Let $|s_i| < 1$, $|s_i + \alpha_i| < 1$ for $i = 1, 2$, $s = (s_1, s_2)$ and $s + \alpha = (s_1 + \alpha_1, s_2 + \alpha_2)$. Then there exist a positive constant C and $\alpha_i^0(s_1, s_2) \in (0, 1)$ such that, if $|\alpha_i| < \alpha_i^0(s_1, s_2)$ with $i = 1, 2$,*

$$C^{-1} \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)} \leq \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{B}_{pq}^{s+\alpha}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{pq}^s(\mathbb{R}^2)}$$

for all $f \in \dot{B}_{pq}^s(\mathbb{R}^2)$ with $p, q \in [1, \infty]$, and for all $f \in \dot{F}_{pq}^s(\mathbb{R}^2)$ with $p \in (1, \infty)$ and $q \in (1, \infty]$,

$$C^{-1} \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)} \leq \|I_{(\alpha_1, \alpha_2)}(f)\|_{\dot{F}_{pq}^{s+\alpha}(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{pq}^s(\mathbb{R}^2)}.$$

5. Embedding theorems and fractional integrals

In this section we first present some embedding theorems for both Besov spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ and Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$. We remark that our embedding theorems for Triebel-Lizorkin spaces are not the same as those for Besov spaces, which reflects the difference between these two kinds of spaces. As an application, we then obtain another boundedness of fractional integrals $I_{(\alpha_1, \alpha_2)}$ on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$ again.

Theorem 5.1. *Let $q \in [1, \infty]$, $p_1, p_2 \in [1, \infty]$, $|s_i| < 1$, $|\bar{s}_i| < 1$, and $\bar{s}_i < s_i$ with $i = 1, 2$. Then*

- (i) *If $s_1 - 3/p_1 = \bar{s}_1 - 3/p_2$, then $\dot{B}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2) \subset \dot{B}_{p_2, q}^{(\bar{s}_1, s_2)}(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{B}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)$,*

$$\|f\|_{\dot{B}_{p_2, q}^{(\bar{s}_1, s_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)}.$$

(ii) If $\bar{s}_1 - s_1 = 1/p_2 - 1/p_1 = \bar{s}_2 - s_2$, then $\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2) \subset \dot{B}_{p_2,q}^{(\bar{s}_1,\bar{s}_2)}(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)$,

$$\|f\|_{\dot{B}_{p_2,q}^{(\bar{s}_1,\bar{s}_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)}.$$

Proof. To simplify our proof, based on Theorem 2.8 and Theorem 2.6, we may suppose that $\psi^{(i)}$ for $i = 1, 2$ in Lemma 2.4 have compact supports, namely,

$$\text{supp } \psi^{(1)} \subset \{(x_1, x_2) \in \mathbb{R}^2 : \|(x_1, x_2)\| \leq 1\}$$

and $\text{supp } \psi^{(2)} \subset \{x_3 \in \mathbb{R} : |x_3| \leq 1\}$. Let other notation be the same as in Lemma 2.4. We then easily show that for all $j_1, k_1 \in \mathbb{Z}$, $(x_1, x_2) \in \mathbb{R}^2$, and $x_3 \in \mathbb{R}$,

$$\begin{aligned} \text{supp } \psi_{j_1}^{(1)} * \psi_{k_1}^{(1)} &\subset \{(x_1, x_2) \in \mathbb{R}^2 : \|(x_1, x_2)\| \leq 2^{1-(j_1 \wedge k_1)}\}, \\ |(\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)})(x_1, x_2)| &\leq C 2^{-|j_1 - k_1|} 2^{3(j_1 \wedge k_1)} \end{aligned} \tag{39}$$

and

$$\begin{aligned} \text{supp } \psi_{j_2}^{(2)} * \psi_{k_2}^{(2)} &\subset \{x_3 \in \mathbb{R} : |x_3| \leq 2^{1-(j_2 \wedge k_2)}\}, \\ |(\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(x_3)| &\leq C 2^{-|j_2 - k_2|} 2^{j_2 \wedge k_2}. \end{aligned} \tag{40}$$

The estimates (39) and (40), Lemma 2.4, and the Hölder inequality yield that for all $j_1, j_2 \in \mathbb{Z}$ and $(x_1, x_2) \in \mathbb{R}^2$

$$\begin{aligned} |\psi_{j_1 j_2} * f(x_1, x_2)| &= \left| \sum_{k_1, k_2 \in \mathbb{Z}} \int_{\mathbb{R}^2} (\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)}) *_2 (\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(x_1 - y_1, x_2 - y_2) \right. \\ &\quad \left. \times \psi_{k_1 k_2} * f(y_1, y_2) dy_1 dy_2 \right| \\ &\leq C \sum_{k_1, k_2 \in \mathbb{Z}} 2^{-|j_1 - k_1|/p'_1 - |j_2 - k_2|/p'_1} \\ &\quad \times \left\{ \int_{\mathbb{R}^2} |(\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)}) *_2 (\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(x_1 - y_1, x_2 - y_2)| \right. \\ &\quad \left. \times |(\psi_{k_1 k_2} * f)(y_1, y_2)|^{p_1} dy_1 dy_2 \right\}^{1/p_1}. \end{aligned}$$

Noticing that $p_2 > p_1$, by the above estimate and the Minkowski inequality, we obtain

$$\begin{aligned} \|\psi_{j_1 j_2} * f\|_{L^{p_2}(\mathbb{R}^2)} &\leq C \sum_{k_1, k_2 \in \mathbb{Z}} 2^{-|j_1 - k_1|/p'_1 - |j_2 - k_2|/p'_1} \|\psi_{k_1 k_2} * f\|_{L^{p_1}(\mathbb{R}^2)} \\ &\quad \times \left\{ \int_{\mathbb{R}^2} |(\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)}) *_2 (\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(x_1, x_2)|^{p_2/p_1} dx_1 dx_2 \right\}^{p_1/p_2}. \end{aligned} \tag{41}$$

We now have two ways to estimate the last quantity in (41), which result in the conclusions (i) and (ii) of Theorem 5.1, respectively.

Proof of (i). The Minkowski inequality and the estimates (39) and (40) imply that

$$\left\{ \int_{\mathbb{R}^2} |(\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)}) *_2 (\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(x_1, x_2)|^{p_2/p_1} dx_1 dx_2 \right\}^{p_1/p_2} \leq C 2^{-|j_1-k_1|} 2^{3(j_1 \wedge k_1)(1-p_1/p_2)} 2^{-|j_2-k_2|}.$$

Inserting this in (41) leads us to

$$\|\psi_{j_1 j_2} * f\|_{L^{p_2}(\mathbb{R}^2)} \leq C \sum_{k_1, k_2 \in \mathbb{Z}} 2^{-|j_1-k_1|-|j_2-k_2|} 2^{3(j_1 \wedge k_1)(1/p_1-1/p_2)} \|\psi_{k_1 k_2} * f\|_{L^{p_1}(\mathbb{R}^2)},$$

which together with Lemma 2.4 and the assumption that $s_1 - 3/p_1 = \bar{s}_1 - 3/p_2$ yields that

$$\begin{aligned} & \|f\|_{\dot{B}_{p_2, q}^{(\bar{s}_1, s_2)}(\mathbb{R}^2)} \\ & \leq C \left\{ \sum_{j_1, j_2 \in \mathbb{Z}} 2^{j_1 \bar{s}_1 q} 2^{j_2 s_2 q} \right. \\ & \quad \times \left. \left[\sum_{k_1, k_2 \in \mathbb{Z}} 2^{-|j_1-k_1|-|j_2-k_2|} 2^{3(j_1 \wedge k_1)(1/p_1-1/p_2)} \|\psi_{k_1 k_2} * f\|_{L^{p_1}(\mathbb{R}^2)} \right]^q \right\}^{1/q} \\ & \leq C \left\{ \sum_{k_1, k_2 \in \mathbb{Z}} 2^{k_1 s_1 q} 2^{k_2 s_2 q} \|\psi_{k_1 k_2} * f\|_{L^{p_1}(\mathbb{R}^2)}^q \right\}^{1/q} \leq C \|f\|_{\dot{B}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)}, \end{aligned}$$

where in the second-to-last inequality, we used the assumptions that $\bar{s}_1 < 1$, $s_1 > -1$, and $|s_2| < 1$. This proves (i) of Theorem 5.1. ■

Proof of (ii). Again, the Minkowski inequality and the estimates (39) and (40) yield that

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^2} |(\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)}) *_2 (\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(x_1, x_2)|^{p_2/p_1} dx_1 dx_2 \right\}^{p_1/p_2} \\ & \leq \left\{ \int_{\mathbb{R}} \left[\int_{\mathbb{R}} |(\psi_{j_1}^{(1)} * \psi_{k_1}^{(1)})(x_1, x_3)| \right. \right. \\ & \quad \times \left. \left. \left\{ \int_{\mathbb{R}} |(\psi_{j_2}^{(2)} * \psi_{k_2}^{(2)})(z)|^{p_2/p_1} dz \right\}^{p_1/p_2} dx_3 \right]^{p_2/p_1} dx_1 \right\}^{p_1/p_2} \\ & \leq C 2^{(j_1 \wedge k_1)(1-p_1/p_2)-|j_1-k_1|} 2^{(j_2 \wedge k_2)(1-p_1/p_2)-|j_2-k_2|}. \end{aligned}$$

This, Lemma 2.4, and the assumptions that $\bar{s}_1 - s_1 = 1/p_2 - 1/p_1 = \bar{s}_2 - s_2$ lead us to

$$\begin{aligned} \|f\|_{\dot{B}_{p_2,q}^{(\bar{s}_1,\bar{s}_2)}(\mathbb{R}^2)} &\leq C \left\{ \sum_{j_1,j_2 \in \mathbb{Z}} 2^{j_1 \bar{s}_1 q} 2^{j_2 \bar{s}_2 q} \left[\sum_{k_1,k_2 \in \mathbb{Z}} 2^{(j_1 \wedge k_1)(1/p_1 - 1/p_2) - |j_1 - k_1|} \right. \right. \\ &\quad \left. \left. \times 2^{(j_2 \wedge k_2)(1/p_1 - 1/p_2) - |j_2 - k_2|} \|\psi_{k_1 k_2} * f\|_{L^{p_1}(\mathbb{R}^2)} \right]^q \right\}^{1/q} \\ &\leq C \left\{ \sum_{k_1,k_2 \in \mathbb{Z}} 2^{k_1 s_1 q} 2^{k_2 s_2 q} \|\psi_{k_1 k_2} * f\|_{L^{p_1}(\mathbb{R}^2)}^q \right\}^{1/q} \\ &\leq C \|f\|_{\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)}, \end{aligned}$$

where in the second-to-last inequality, we used the assumptions that $\bar{s}_i < 1$ and $s_i > -1$ for $i = 1, 2$, which proves (ii) of Theorem 5.1. This finishes the proof of Theorem 5.1. \square

We now establish an embedding theorem for Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$.

Theorem 5.2. *Let $q_i \in (1, \infty]$, $p_i \in (1, \infty)$, $|s_i| < 1$, and $|\bar{s}_i| < 1$ with $i = 1, 2$. Assume that $s_2 < \bar{s}_2$, $\bar{s}_1 + 2\bar{s}_2 < s_1 + 2s_2$, and $s_1 + 2s_2 - 3/p_1 = \bar{s}_1 + 2\bar{s}_2 - 3/p_2$. Then $\dot{F}_{p_1,q_1}^{(s_1,s_2)}(\mathbb{R}^2) \subset \dot{F}_{p_2,q_2}^{(\bar{s}_1,\bar{s}_2)}(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{F}_{p_1,q_1}^{(s_1,s_2)}(\mathbb{R}^2)$,*

$$\|f\|_{\dot{F}_{p_2,q_2}^{(\bar{s}_1,\bar{s}_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{p_1,q_1}^{(s_1,s_2)}(\mathbb{R}^2)}.$$

Proof. By Proposition 2.9 (ii), we may assume that $q_1 = \infty$ and $q_2 = 1$. Thus, to prove the theorem, we only need to show $\dot{F}_{p_1,\infty}^{(s_1,s_2)}(\mathbb{R}^2) \subset \dot{F}_{p_2,1}^{(\bar{s}_1,\bar{s}_2)}(\mathbb{R}^2)$. Let $f \in \dot{F}_{p_1,\infty}^{(s_1,s_2)}(\mathbb{R}^2)$ and $\|f\|_{\dot{F}_{p_1,\infty}^{(s_1,s_2)}(\mathbb{R}^2)} = 1$ by homogeneity. We also let $\psi^{(i)}$ for $i = 1, 2$ be the same as in the proof of Corollary 2.22. Using the same notation as in Lemma 2.4, by Lemma 2.4 and this special choice of $\psi^{(i)}$ for $i = 1, 2$, we then have that

$$f(x_1, x_2) = \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{2j_1+3} \psi_{j_1 j_2} * \psi_{j_1 j_2} * f(x_1, x_2) \tag{42}$$

holds with the same meaning as in (4). By (42) and the Hölder inequality, we have that, for $k_1, k_2 \in \mathbb{Z}$ and $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} |\psi_{k_1 k_2} * f(x_1, x_2)| &\leq \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{2j_1+3} \left\{ \int_{\mathbb{R}^2} |(\psi_{k_1 k_2} * \psi_{j_1 j_2})(y_1, y_2)|^{p'_1} dy_1 dy_2 \right\}^{1/p'_1} \\ &\quad \times \|\psi_{j_1 j_2} * f\|_{L^{p_1}(\mathbb{R}^2)}. \end{aligned} \tag{43}$$

The estimate (32) and the Minkowski inequality imply that

$$\begin{aligned} & \left\{ \int_{\mathbb{R}^2} |(\psi_{k_1 k_2} * \psi_{j_1 j_2})(y_1, y_2)|^{p'_1} dy_1 dy_2 \right\}^{1/p'_1} \\ & \leq C 2^{-|j_2-k_2|} 2^{-|j_1-k_1|} \left\{ \int_{\mathbb{R}^2} \frac{2^{-(j_1 \wedge k_1)p'_1}}{(2^{-(j_1 \wedge k_1)} + \|(y_1, y_2)\|)^{4p'_1}} dy_1 dy_2 \right\}^{1/p'_1} \\ & \leq C 2^{-|j_2-k_2|} 2^{3(j_1 \wedge k_1)/p_1 - |j_1-k_1|}. \end{aligned}$$

Combining this estimate with (43) and using the assumption that $\|f\|_{\dot{F}_{p_1, \infty}^{(s_1, s_2)}(\mathbb{R}^2)} = 1$ yield that

$$|\psi_{k_1 k_2} * f(x_1, x_2)| \leq C \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{2j_1+3} 2^{-|j_2-k_2|-j_2 s_2} 2^{3(j_1 \wedge k_1)/p_1 - |j_1-k_1| - j_1 s_1}.$$

From this and the assumptions that $\bar{s}_2 > s_2$, $s_1 + 2s_2 - 3/p_1 = \bar{s}_1 + 2\bar{s}_2 - 3/p_2$, and $|s_i| < 1$ for $i = 1, 2$, it follows that, for any fixed $N \in \mathbb{Z}$ and all $(x_1, x_2) \in \mathbb{R}^2$,

$$\begin{aligned} & \sum_{k_1=-\infty}^N \sum_{k_2=-\infty}^{2k_1+3} 2^{k_1 \bar{s}_1} 2^{k_2 \bar{s}_2} |\psi_{k_1 k_2} * f(x_1, x_2)| \\ & \leq C \sum_{k_1=-\infty}^N \sum_{k_2=-\infty}^{2k_1+3} 2^{k_1 \bar{s}_1} 2^{k_2(\bar{s}_2 - s_2)} \\ & \quad \times \sum_{j_1=-\infty}^{\infty} \sum_{j_2=-\infty}^{2j_1+3} 2^{(k_2 - j_2)s_2 - |j_2 - k_2|} 2^{3(j_1 \wedge k_1)/p_1 - |j_1 - k_1| - j_1 s_1} \\ & \leq C_0 2^{3N/p_2}, \end{aligned} \tag{44}$$

where C_0 is a positive constant independent of $N \in \mathbb{Z}$. On the other hand, for all $N \in \mathbb{Z}$ and $(x_1, x_2) \in \mathbb{R}^2$, by the assumptions that $s_2 < \bar{s}_2$ and $\bar{s}_1 + 2\bar{s}_2 < s_1 + 2s_2$, we have

$$\begin{aligned} & \sum_{k_1=N+1}^{\infty} \sum_{k_2=-\infty}^{2k_1+3} 2^{k_1 \bar{s}_1} 2^{k_2 \bar{s}_2} |\psi_{k_1 k_2} * f(x_1, x_2)| \\ & \leq \sum_{k_1=N+1}^{\infty} \sum_{k_2=-\infty}^{2k_1+3} 2^{k_1(\bar{s}_1 - s_1)} 2^{k_2(\bar{s}_2 - s_2)} \sup_{\substack{j_1 \in \mathbb{Z} \\ -\infty < j_2 \leq 2j_1 + 3}} 2^{j_1 s_1} 2^{j_2 s_2} |\psi_{j_1 j_2} * f(x_1, x_2)| \\ & \leq C_1 2^{N(\bar{s}_1 + 2\bar{s}_2 - s_1 - 2s_2)} \sup_{\substack{j_1 \in \mathbb{Z} \\ -\infty < j_2 \leq 2j_1 + 3}} 2^{j_1 s_1} 2^{j_2 s_2} |\psi_{j_1 j_2} * f(x_1, x_2)|, \end{aligned} \tag{45}$$

where the positive constant C_1 is independent of $N \in \mathbb{Z}$. The estimates (44) and (45) then imply that

$$\begin{aligned} \|f\|_{\dot{F}_{p_2,1}^{(s_1,s_2)}(\mathbb{R}^2)}^{p_2} &= p_2 \sum_{N=-\infty}^{\infty} \int_{C_0 2^{3N/p_2}}^{C_0 2^{3(N+1)/p_2}} t^{p_2-1} \\ &\quad \times \left| \left\{ (x_1, x_2) \in \mathbb{R}^2 : \sum_{k_1=-\infty}^N \sum_{k_2=-\infty}^{2k_1+3} 2^{k_1 s_1} 2^{k_2 s_2} |\psi_{k_1 k_2} * f(x_1, x_2)| \right. \right. \\ &\quad \left. \left. + \sum_{k_1=N+1}^{\infty} \sum_{k_2=-\infty}^{2k_1+3} \dots > t \right\} \right| dt \\ &\leq C p_2 \int_0^{\infty} t^{p_1-1} \left| \left\{ (x_1, x_2) \in \mathbb{R}^2 : \sup_{\substack{j_1 \in \mathbb{Z} \\ -\infty < j_2 \leq 2j_1+3}} 2^{j_1 s_1} 2^{j_2 s_2} \right. \right. \\ &\quad \left. \left. \times |\psi_{j_1 j_2} * f(x_1, x_2)| > t \right\} \right| dt \\ &= C \|f\|_{\dot{F}_{p_1,\infty}^{(s_1,s_2)}(\mathbb{R}^2)}^{p_1} = C, \end{aligned}$$

where the positive constant C is independent of f . This finishes the proof of Theorem 5.2. □

As a corollary of Proposition 4.3 and Theorems 5.1 and 5.2, we have the following boundedness of $I_{(\alpha_1,\alpha_2)}$ on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$.

Corollary 5.3. *Let $|s_i| < 1$, $|\alpha_i| < 1$, and $|s_i + \alpha_i| < 1$ for $i = 1, 2$. Then,*

- (i) *If $q \in [1, \infty]$, $\alpha_1 > 0$, $p_1 \in (1, \infty)$, and $1/p_2 = 1/p_1 - \alpha_1/3$, then $I_{(\alpha_1,\alpha_2)}$ is bounded from $\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)$ to $\dot{B}_{p_2,q}^{(s_1,s_2+\alpha_2)}(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)$,*

$$\|I_{(\alpha_1,\alpha_2)}(f)\|_{\dot{B}_{p_2,q}^{(s_1,s_2+\alpha_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)}.$$

- (ii) *If $q \in [1, \infty]$, $0 < \alpha_1 = \alpha_2 < 1$, $p_1 \in (1, \infty)$, and $1/p_2 = 1/p_1 - \alpha_1$, then $I_{(\alpha_1,\alpha_2)}$ is bounded from $\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)$ to $\dot{B}_{p_2,q}^{(s_1,s_2)}(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)$,*

$$\|I_{(\alpha_1,\alpha_2)}(f)\|_{\dot{B}_{p_2,q}^{(s_1,s_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{p_1,q}^{(s_1,s_2)}(\mathbb{R}^2)}.$$

- (iii) *If $q_1, q_2 \in (1, \infty]$, $\alpha_2 < 0$, $\alpha_1 > -2\alpha_2$, $p_1 \in (1, \infty)$, and $1/p_2 = 1/p_1 - (\alpha_1 + 2\alpha_2)/3$, then $I_{(\alpha_1,\alpha_2)}$ is bounded from $\dot{F}_{p_1,q_1}^{(s_1,s_2)}(\mathbb{R}^2)$ to $\dot{F}_{p_2,q_2}^{(s_1,s_2)}(\mathbb{R}^2)$, namely, there exists a positive constant C such that, for all $f \in \dot{F}_{p_1,q_1}^{(s_1,s_2)}(\mathbb{R}^2)$,*

$$\|I_{(\alpha_1,\alpha_2)}(f)\|_{\dot{F}_{p_2,q_2}^{(s_1,s_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{p_1,q_1}^{(s_1,s_2)}(\mathbb{R}^2)}.$$

Finally, we consider the boundedness of semi-fractional integrals determined by the kernel K_γ on $\dot{B}_{pq}^s(\mathbb{R}^2)$ and $\dot{F}_{pq}^s(\mathbb{R}^2)$.

Theorem 5.4. *Let K_0 be a distribution kernel on \mathbb{R} which coincides with a C^∞ function away from $\{0\}$ and which satisfies:*

- (i) *For any $\alpha \in \mathbb{Z}_+$, there exists a positive constant C_α such that, for all $z \in \mathbb{R} \setminus \{0\}$,*

$$|\partial_z^\alpha K_0(z)| \leq C_\alpha |z|^{-1-\alpha}.$$

- (ii) *For any given normalized bump function φ on \mathbb{R} and any $\delta > 0$, there exists a positive constant C such that*

$$\left| \int_{\mathbb{R}} K_0(z) \varphi(\delta z) dz \right| \leq C.$$

For $\gamma \in (0, 3)$, define the distribution kernel

$$K_\gamma(x_1, x_2) = (\|x_1, \cdot\|)^{\gamma-3} *_2 K_0(x_2),$$

where $(x_1, x_2) \in \mathbb{R}^2$. Let $|s_i| < 1$ for $i = 1, 2$. If $p_1 \in (1, \infty)$ and $1/p_2 = 1/p_1 - \gamma/3$, then there exists a positive constant C such that, for all $f \in \dot{B}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)$ with $q \in [1, \infty]$,

$$\|K_\gamma * f\|_{\dot{B}_{p_2, q}^{(s_1, s_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{B}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)},$$

and for all $f \in \dot{F}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)$ with $q \in (1, \infty]$,

$$\|K_\gamma * f\|_{\dot{F}_{p_2, q}^{(s_1, s_2)}(\mathbb{R}^2)} \leq C \|f\|_{\dot{F}_{p_1, q}^{(s_1, s_2)}(\mathbb{R}^2)}.$$

Proof. Let $\psi^{(i)}$ for $i = 1, 2$ be the same as in the proof of Theorem 3.1. Repeating the proof of Theorem 3.1 implies that

$$\begin{aligned} & |(\psi_{u_1 u_2} * K_\gamma * \psi_{t_1 t_2})(x_1, x_2)| \\ & \leq C \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) \\ & \quad \times \int_{\mathbb{R}} \frac{u_1 \vee t_1}{(u_1 \vee t_1 + \|(x_1, x_2 - x_3)\|)^{4-\gamma}} \frac{u_2 \vee t_2}{(u_2 \vee t_2 + |x_3|)^2} dx_3. \end{aligned}$$

Inserting this estimate into (23) yields that

$$\begin{aligned} & |(\psi_{u_1 u_2} * K_\gamma * f)(x_1, x_2)| \\ & \leq C \int_0^\infty \int_0^\infty \left(\frac{u_1}{t_1} \wedge \frac{t_1}{u_1} \right) \left(\frac{u_2}{t_2} \wedge \frac{t_2}{u_2} \right) I_\gamma \circ M_2(|(\psi_{t_1 t_2} * f)(x_1, \cdot)|)(x_2) \frac{dt_1}{t_1} \frac{dt_2}{t_2}, \end{aligned}$$

where I_γ is the standard fractional integral on \mathbb{R}^2 when \mathbb{R}^2 is regarded as a space of homogeneous type, and M_2 is the usual Hardy-Littlewood maximal function on the second variable. It is well-known that I_γ is bounded from $L^{p_1}(\mathbb{R}^2)$ into $L^{p_2}(\mathbb{R}^2)$, where $p_1 \in (1, \infty)$ and $1/p_2 = 1/p_1 - \gamma/3$; see [8]. Using this fact and noticing that I_γ is a positive operator, by some computation similar to the proof of Theorem 2.6, we complete the proof of Theorem 5.4. We leave the details to the reader. \square

Finally, we point out that using some discrete Calderón reproducing formulae as in Theorem 1.8 in [3], we can develop a theory of Besov spaces $\dot{B}_{pq}^s(\mathbb{R}^2)$ and Triebel-Lizorkin spaces $\dot{F}_{pq}^s(\mathbb{R}^2)$ for full $s \in \mathbb{R}$ and $p, q \in (0, \infty]$. However, to limit the length of this paper and to simplify its presentation, we restrict ourself to the current case.

Acknowledgements. This paper was initially written during the period when Professor Yongsheng Han from Auburn University visited Beijing Normal University in 2005. The author sincerely thanks Professor Yongsheng Han to show him many ideas which are now presented in [3] and to have many stimulating conversations on this subject at that period. He also thanks editor Jose L. Gámez for his careful reading of the manuscript and valuable remarks which made this article more readable.

References

- [1] C. Fefferman and E. M. Stein, *Some maximal inequalities*, Amer. J. Math. **93** (1971), 107–115.
- [2] M. Frazier, B. Jawerth, and G. Weiss, *Littlewood-Paley theory and the study of function spaces*, CBMS Regional Conference Series in Mathematics, vol. 79, American Mathematical Society, Providence, RI, 1991.
- [3] Y. Han and G. Lu, *Discrete Littlewood-Paley-Stein theory and multi-parameter Hardy spaces associated with flag singular integrals*, available at <http://arxiv.org/abs/0801.1701v1>.
- [4] Y. S. Han and E. T. Sawyer, *Littlewood-Paley theory on spaces of homogeneous type and the classical function spaces*, Mem. Amer. Math. Soc. **110** (1994), no. 530, 1–126.
- [5] D. Müller, F. Ricci, and E. M. Stein, *Marcinkiewicz multipliers and multi-parameter structure on Heisenberg(-type) groups*, I, Invent. Math. **119** (1995), no. 2, 199–233.
- [6] A. Nagel, F. Ricci, and E. M. Stein, *Singular integrals with flag kernels and analysis on quadratic CR manifolds*, J. Funct. Anal. **181** (2001), no. 1, 29–118.
- [7] H.-J. Schmeisser and H. Triebel, *Topics in Fourier analysis and function spaces*, A Wiley-Interscience Publication, John Wiley & Sons Ltd., Chichester, 1987.
- [8] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, vol. 30, Princeton University Press, Princeton, N.J., 1970.
- [9] H. Triebel, *Theory of function spaces*, Monographs in Mathematics, vol. 78, Birkhäuser Verlag, Basel, 1983.
- [10] ———, *Spaces of Besov-Hardy-Sobolev type*, Teubner-Texte zur Mathematik, vol. 15, BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1978.
- [11] ———, *The structure of functions*, Monographs in Mathematics, vol. 97, Birkhäuser Verlag, Basel, 2001.