

## BEST PROXIMITY POINT RESULTS VIA SIMULATION FUNCTIONS IN METRIC-LIKE SPACES

G. V. V. J. RAO<sup>1</sup>, H. K. NASHINE<sup>2\*</sup>, AND Z. KADELBURG<sup>3</sup>

ABSTRACT. In this paper, we discuss the existence of best proximity points of certain mappings via simulation functions in the frame of complete metric-like spaces. Some consequences and examples are given of the obtained results.

### 1. INTRODUCTION

Khojasteh et al. introduced in [13] the notion of simulation function in order to unify several fixed point results obtained by various authors. These functions were later utilized by Karapinar and Khojasteh in [9] to solve some problems concerning best proximity points.

On the other hand, spaces more general than metric and fixed point and related problems in them have been lately a wide field of interest of huge number of mathematicians. Among them, metric-like spaces, introduced by Amini-Harandi in [2], took a prominent place.

In this paper, we are going to extend these investigations to best proximity points of mappings acting in complete metric-like spaces, using conditions involving simulation functions. The results will be illustrated by several examples, showing the strength of these results compared with others existing in the literature.

### 2. PRELIMINARIES

Throughout the paper,  $\mathbb{R}$  and  $\mathbb{R}^+$ ,  $\mathbb{R}_0^+$  will denote the set of real numbers, the set of positive real numbers and the set of nonnegative real numbers, respectively. Also,  $\mathbb{N}_0$  and  $\mathbb{N}$  will denote the set of nonnegative, resp. positive integers.

---

*Key words and phrases.*  $\mathcal{Z}$ -contraction, best proximity point, simulation function, admissible mapping.

2010 *Mathematics Subject Classification.* Primary: 41A50. Secondary: 47H10, 54H25.

DOI 10.46793/KgJMat2003.401R

*Received:* April 14, 2018.

*Accepted:* May 22, 2018.

We shall first recall some basic definitions and some results from [1, 5, 13].

**Definition 2.1** ([13]). A simulation function is a mapping  $\zeta : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$  satisfying the following conditions:

- ( $\zeta_1$ )  $\zeta(0, 0) = 0$ ;
- ( $\zeta_2$ )  $\zeta(t, s) < s - t$  for all  $t, s > 0$ ;
- ( $\zeta_3$ ) if  $\{t_n\}$  and  $\{s_n\}$  are sequences in  $(0, \infty)$  such that  $\lim_{n \rightarrow \infty} t_n = \lim_{n \rightarrow \infty} s_n = l \in (0, \infty)$ , then  $\limsup_{n \rightarrow \infty} \zeta(t_n, s_n) < 0$ .

Note that, according to the axiom ( $\zeta_2$ ), each simulation function  $\zeta$  satisfies  $\zeta(t, t) < 0$  for all  $t > 0$ . The family of all simulation functions will be denoted by  $\mathcal{Z}$ .

*Example 2.1* (See, e.g., [1, 5, 7, 13]). For  $i = 1, 2, \dots, 6$ , define mappings  $\zeta_i : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$ , as follows.

- (i)  $\zeta_1(t, s) = \phi_1(s) - \phi_2(t)$  for all  $t, s \in \mathbb{R}_0^+$ , where  $\phi_1, \phi_2 : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  are continuous functions, with  $\phi_i(t) = 0$  if and only if  $t = 0$  and  $\phi_1(t) < t \leq \phi_2(t)$  for all  $t > 0$ .
- (ii)  $\zeta_2(t, s) = s - \frac{f(t,s)}{g(t,s)} t$  for all  $t, s \in \mathbb{R}_0^+$ , where  $f, g : \mathbb{R}_0^{+2} \rightarrow \mathbb{R}_0^+$  are two functions, continuous with respect to each variable and such that  $f(t, s) > g(t, s)$  for all  $t, s > 0$ .
- (iii)  $\zeta_3(t, s) = s - \phi(s) - t$  for all  $t, s \in \mathbb{R}_0^+$ , where  $\phi : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  is a continuous functions, with  $\phi(t) = 0$  if and only if  $t = 0$ .
- (iv) If  $\varphi : \mathbb{R}_0^+ \rightarrow [0, 1)$  is a function such that  $\limsup_{t \rightarrow r^+} \varphi(t) < 1$  for all  $r > 0$ , let

$$\zeta_4(t, s) = s\varphi(s) - t, \quad \text{for all } t, s \in \mathbb{R}_0^+.$$

- (v) If  $\eta : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  is an upper semi-continuous function such that  $\eta(t) < t$  for all  $t > 0$  and  $\eta(0) = 0$ , let

$$\zeta_5(t, s) = \eta(s) - t, \quad \text{for all } t, s \in \mathbb{R}_0^+.$$

- (vi) If  $\phi : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  is a function such that  $\int_0^\epsilon \phi(u) du > \epsilon$  for each  $\epsilon > 0$ , let

$$\zeta_6(t, s) = s - \int_0^t \phi(u) du, \quad \text{for all } t, s \in \mathbb{R}_0^+.$$

It is clear that each function  $\zeta_i$ ,  $i = 1, 2, \dots, 6$ , is a simulation function.

**Definition 2.2** ([2]). Let  $X$  be a nonempty set, and a mapping  $\sigma : X \times X \rightarrow \mathbb{R}_0^+$  is such that, for all  $x, y, z \in X$ ,

- ( $\sigma_1$ )  $\sigma(x, y) = 0$  implies  $x = y$ ;
- ( $\sigma_2$ )  $\sigma(x, y) = \sigma(y, x)$ ;
- ( $\sigma_3$ )  $\sigma(x, y) \leq \sigma(x, z) + \sigma(z, y)$ .

Then  $(X, \sigma)$  is said to be a metric-like space.

As is well known, each partial metric space is an example of a metric-like space. The converse is not true. The following example illustrates this statement.

*Example 2.2.* Take  $X = \{1, 2, 3\}$  and consider the metric-like  $\sigma : X \times X \rightarrow \mathbb{R}_0^+$  given by

$$\begin{aligned} \sigma(1, 1) &= 0, & \sigma(2, 2) &= 1, & \sigma(3, 3) &= \frac{2}{3}, \\ \sigma(2, 1) = \sigma(1, 2) &= \frac{9}{10}, & \sigma(1, 3) = \sigma(3, 1) &= \frac{7}{10}, & \sigma(2, 3) = \sigma(3, 2) &= \frac{4}{5}. \end{aligned}$$

Since  $\sigma(2, 2) \neq 0$ ,  $\sigma$  is not a metric and since  $\sigma(2, 2) > \sigma(2, 1)$ ,  $\sigma$  is not a partial metric.

Every metric-like  $\sigma$  on  $X$  generates a topology  $\tau_\sigma$  whose base is the family of all open  $\sigma$ -balls

$$\{ B_\sigma(x, \delta) : x \in X, \delta > 0 \},$$

where  $B_\sigma(x, \delta) = \{ y \in X : |\sigma(x, y) - \sigma(x, x)| < \delta \}$ , for all  $x \in X$  and  $\delta > 0$ .

**Definition 2.3** ([2]). Let  $(X, \sigma)$  be a metric-like space, let  $\{x_n\}$  be a sequence in  $X$  and  $x \in X$ . Then

- (i)  $\{x_n\}$  is said to converge to  $x$ , w.r.t.  $\tau_\sigma$ , if  $\lim_{n \rightarrow \infty} \sigma(x_n, x) = \sigma(x, x)$ ;
- (ii)  $\{x_n\}$  is called a Cauchy sequence in  $(X, \sigma)$  if  $\lim_{n, m \rightarrow \infty} \sigma(x_n, x_m)$  exists (and is finite);
- (iii)  $(X, \sigma)$  is called complete if every Cauchy sequence  $\{x_n\}$  in  $X$  converges with respect to  $\tau_\sigma$  to a point  $x \in X$  such that

$$\lim_{n, m \rightarrow \infty} \sigma(x_n, x_m) = \lim_{n \rightarrow \infty} \sigma(x_n, x) = \sigma(x, x);$$

- (iv) a function  $f : X \rightarrow X$  is continuous if for any sequence  $\{x_n\}$  in  $X$  such that  $\sigma(x_n, x) \rightarrow \sigma(x, x)$  as  $n \rightarrow \infty$ , we have  $\sigma(fx_n, fx) \rightarrow \sigma(fx, fx)$  as  $n \rightarrow \infty$ .

Note that the limit of a sequence in a metric-like space might not be unique.

**Lemma 2.1** ([11]). Let  $(X, \sigma)$  be a metric-like space. Let  $\{x_n\}$  be a sequence in  $X$  such that  $x_n \rightarrow x$  where  $x \in X$  and  $\sigma(x, x) = 0$ . Then for all  $y \in X$ , we have

$$\lim_{n \rightarrow \infty} \sigma(x_n, y) = \sigma(x, y).$$

$\Psi$  will denote the family of non-decreasing functions  $\psi : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  satisfying the following conditions:

- (i)  $\psi(t) < t$ , for any  $t \in \mathbb{R}^+$ ;
- (ii)  $\psi$  is continuous at 0.

Let  $(X, \sigma)$  be a metric-like space, and  $U$  and  $V$  be two non-empty subsets of  $X$ . Recall the following standard notation:

$$\begin{aligned} \sigma(U, V) &:= \inf\{\sigma(u, v) : u \in U, v \in V\}, \\ U_0 &:= \{u \in U : \sigma(u, v) = \sigma(U, V) \text{ for some } v \in V\}, \\ V_0 &:= \{v \in V : \sigma(u, v) = \sigma(U, V) \text{ for some } u \in U\}. \end{aligned}$$

Consider now a non-self mapping  $T : U \rightarrow V$  and the equation  $Tu = u$  ( $u \in U$ ). As is well known, a solution of this equation, if it exists, is called a fixed point of  $T$ . If such solution does not exist, an approximate solution  $u^* \in U$  have the least possible error when  $\sigma(u^*, Tu^*) = \sigma(U, V)$ . In this case,  $u^*$  is called a best proximity point of the mapping  $T : U \rightarrow V$ .

Finally, recall the following useful notions.

**Definition 2.4** ([6]). Let  $U$  and  $V$  be nonempty subsets of a metric-like space  $(X, \sigma)$ , and  $\alpha : U \times U \rightarrow \mathbb{R}_0^+$  be a function. We say that the mapping  $T$  is  $\alpha$ -proximal admissible if

$$\alpha(x, y) \geq 1 \text{ and } \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) \Rightarrow \alpha(u, v) \geq 1,$$

for all  $x, y, u, v \in X$ .

If  $\sigma(U, V) = 0$ , then  $T$  reduces from  $\alpha$ -proximal admissible to  $\alpha$ -admissible.

**Definition 2.5** ([8, 10]). Let  $T : X \rightarrow X$  be a mapping and  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  be a function. We say that the mapping  $T$  is triangular weakly- $\alpha$ -admissible if

$$\alpha(x, y) \geq 1 \text{ and } \alpha(y, z) \geq 1 \Rightarrow \alpha(x, z) \geq 1.$$

### 3. MAIN RESULTS

**Definition 3.1.** Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$ ,  $\psi \in \Psi$ ,  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  and  $\zeta \in \mathcal{Z}$ . We say that  $T : U \rightarrow V$  is an  $\alpha$ - $\psi$ - $\zeta$ -contraction if  $T$  is  $\alpha$ -proximal admissible and

$$(3.1) \quad \alpha(x, y) \geq 1 \text{ and } \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) \Rightarrow \zeta(\alpha(x, y)\sigma(u, v), \psi(\sigma(x, y))) \geq 0,$$

for all  $x, y, u, v \in U$ .

**Definition 3.2.** Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$ ,  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  and  $\zeta \in \mathcal{Z}$ . We say that  $T : U \rightarrow V$  is an  $\alpha$ - $\zeta$ -contraction if  $T$  is  $\alpha$ -proximal admissible and

$$(3.2) \quad \alpha(x, y) \geq 1 \text{ and } \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) \Rightarrow \zeta(\alpha(x, y)\sigma(u, v), \sigma(x, y)) \geq 0,$$

for all  $x, y, u, v \in U$ .

Notice that Definition 3.2 is not a special case of Definition 3.1 since the function  $\psi(t) = t$  does not belong to  $\Psi$ .

The following lemma provides a standard step in proving that the given sequence is Cauchy in a certain space.

**Lemma 3.1** (See, e.g., [14]). *Let  $(X, \sigma)$  be a metric-like space and let  $\{x_n\}$  be a sequence in  $X$  such that  $\sigma(x_{n+1}, x_n)$  is non-increasing and that  $\lim_{n \rightarrow \infty} \sigma(x_{n+1}, x_n) = 0$ . If  $\{x_n\}$  is not a Cauchy sequence, then there exist an  $\epsilon > 0$  and two sequences  $\{m_k\}$*

and  $\{n_k\}$  of positive integers such that the following four sequences tend to  $\epsilon$  when  $k \rightarrow \infty$ :

$$\sigma(x_{m_k}, x_{n_k}), \sigma(x_{m_k+1}, x_{n_k+1}), \sigma(x_{m_k-1}, x_{n_k}), \sigma(x_{m_k}, x_{n_k-1}).$$

Now we present the main results of this article.

**Theorem 3.1.** *Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$ ,  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$ ,  $\psi \in \Psi$  and  $\zeta \in \mathcal{Z}$  is non-decreasing with respect to its second argument. Suppose that  $T : U \rightarrow V$  is an  $\alpha$ - $\psi$ - $\zeta$ -contraction and*

- (1)  $T$  is triangular weakly- $\alpha$ -admissible;
- (2)  $U$  is closed with respect to the topology  $\tau_\sigma$ ;
- (3)  $T(U_0) \subset V_0$ ;
- (4) there exist  $x_0, x_1 \in U$  such that  $\sigma(x_1, Tx_0) = \sigma(U, V)$  and  $\alpha(x_0, x_1) \geq 1$ ;
- (5)  $T$  is continuous.

Then,  $T$  has a best proximity point, that is, there exists  $z \in U$  such that  $\sigma(z, Tz) = \sigma(U, V)$ .

*Proof.* Take  $x_0, x_1 \in U$  given as in (4). Taking (3) into account, we conclude that  $Tx_1 \in V_0$  which implies that there exists  $x_2 \in U$  such that  $\sigma(x_2, Tx_1) = \sigma(U, V)$ . Since  $\alpha(x_0, x_1) \geq 1$  and  $T$  is  $\alpha$ -proximal admissible, we conclude that  $\alpha(x_1, x_2) \geq 1$ . Recursively, a sequence  $\{x_n\} \subset U$  can be chosen satisfying

$$(3.3) \quad \sigma(x_{n+1}, Tx_n) = \sigma(U, V) \text{ and } \alpha(x_n, x_{n+1}) \geq 1, \quad \text{for all } n \in \mathbb{N}_0.$$

If  $x_k = x_{k+1}$  for some  $k \in \mathbb{N}_0$ , then  $\sigma(x_k, Tx_k) = \sigma(x_{k+1}, Tx_k) = \sigma(U, V)$ , meaning that  $x_k$  is the required best proximal point. Hence, we will further assume that

$$(3.4) \quad x_n \neq x_{n+1}, \quad \text{for all } n \in \mathbb{N}_0.$$

Using relations (3.3) and (3.4), we get that  $\sigma(x_n, Tx_{n-1}) = \sigma(x_{n+1}, Tx_n) = \sigma(U, V)$ , for all  $n \in \mathbb{N}$ . Furthermore, by (3.1)

$$(3.5) \quad \zeta(\alpha(x_{n-1}, x_n)\sigma(x_n, x_{n+1}), \psi(\sigma(x_{n-1}, x_n))) \geq 0, \quad \text{for all } n \in \mathbb{N},$$

since  $T : U \rightarrow V$  is an  $\alpha$ - $\psi$ - $\zeta$ -contraction. Regarding (3.4) and  $(\zeta_2)$ , the inequality (3.5) implies that

$$\sigma(x_n, x_{n+1}) \leq \alpha(x, y)\sigma(x_n, x_{n+1}) \leq \psi(\sigma(x_{n-1}, x_n)) < \sigma(x_{n-1}, x_n), \quad \text{for all } n \in \mathbb{N}.$$

Thus,  $\{\sigma(x_n, x_{n+1})\}$  is a non-increasing sequence bounded from below and there exists  $L \in \mathbb{R}_0^+$  such that  $\sigma(x_n, x_{n+1}) \rightarrow L$  as  $n \rightarrow \infty$ . We shall prove that  $L = 0$ . Suppose, on the contrary, that  $L > 0$ . Taking the upper limit in (3.5) as  $n \rightarrow \infty$ , regarding  $(\zeta_3)$ , property (i) of  $\psi \in \Psi$  and that  $\zeta$  is non-decreasing with respect to the second argument, we deduce

$$\begin{aligned} 0 &\leq \limsup_{n \rightarrow \infty} \zeta(\alpha(x_n, x_{n-1})\sigma(x_n, x_{n+1}), \psi(\sigma(x_n, x_{n-1}))) \\ &\leq \limsup_{n \rightarrow \infty} \zeta(\alpha(x_n, x_{n-1})\sigma(x_n, x_{n+1}), \sigma(x_n, x_{n-1})) < 0, \end{aligned}$$

which is a contradiction. We conclude that  $\lim_{n \rightarrow \infty} \sigma(x_n, x_{n+1}) = 0$ .

We shall now prove that the sequence  $\{x_n\}$  is Cauchy. Suppose that it is not. Then, there exist  $\epsilon > 0$  and subsequences  $\{x_{m_k}\}$  and  $\{x_{n_k}\}$  of  $\{x_n\}$ , so that  $n_k > m_k > k$  and

$$(3.6) \quad \sigma(x_{m_k}, x_{n_k}) \geq \epsilon \text{ and } \sigma(x_{m_k}, x_{n_k-1}) < \epsilon.$$

By Lemma 2.1, we have

$$\lim_{k \rightarrow \infty} \sigma(x_{m_k}, x_{n_k}) = \lim_{k \rightarrow \infty} \sigma(x_{n_k-1}, x_{m_k-1}) = \epsilon.$$

Since  $T$  is triangular weakly- $\alpha$ -admissible, from (3.3), we get that

$$\alpha(x_n, x_m) \geq 1, \quad \text{for all } n, m \in \mathbb{N}_0 \text{ with } n > m.$$

Hence,

$$(3.7) \quad \alpha(x_{m_k}, x_{n_k}) \geq 1 \text{ and } \sigma(x_{m_k}, Tx_{m_k-1}) = \sigma(x_{n_k}, Tx_{n_k-1}) = \sigma(U, V), \quad \text{for all } k \in \mathbb{N}.$$

Since  $T$  is an  $\alpha$ - $\psi$ - $\zeta$ -contraction, the obtained relations (3.7) yield the following inequality:

$$0 \leq \zeta(\alpha(x_n, x_{n-1})\sigma(x_{m_k}, x_{n_k}), \psi(\sigma(x_{m_k}, x_{n_k}))), \quad \text{for all } k \in \mathbb{N}.$$

Letting  $k \rightarrow \infty$ , using (3.6) and ( $\zeta_3$ ), and regarding properties of  $\psi \in \Psi$  and that  $\zeta$  is non-decreasing with respect to the second argument, we obtain

$$\begin{aligned} 0 &\leq \limsup_{n \rightarrow \infty} \zeta(\alpha(x_n, x_{n-1})\sigma(x_{m_k}, x_{n_k}), \psi(\sigma(x_{m_k-1}, Tx_{n_k-1}))) \\ &\leq \limsup_{n \rightarrow \infty} \zeta(\alpha(x_n, x_{n-1})\sigma(x_{m_k}, x_{n_k}), \sigma(x_{m_k-1}, Tx_{n_k-1})) < 0, \end{aligned}$$

which is a contradiction. Thus, we conclude that the sequence  $\{x_n\}$  is Cauchy in  $U$ .

Since  $U$  is a closed subset of a complete metric-like space  $(X, \sigma)$ , there exists  $z \in U$  such that

$$(3.8) \quad \lim_{n \rightarrow \infty} \sigma(x_n, z) = 0.$$

Since  $T$  is continuous, we deduce that

$$(3.9) \quad \lim_{n \rightarrow \infty} \sigma(Tx_n, Tz) = 0.$$

From (3.3), using the triangle inequality together with (3.8) and (3.9), we find that

$$\sigma(U, V) = \lim_{n \rightarrow \infty} \sigma(x_{n+1}, Tx_n) = \sigma(z, Tz).$$

Thus,  $z \in U$  is a best proximity point of the mapping  $T$ . □

The continuity hypothesis in Theorem 3.1 can be omitted if we assume the following additional condition on  $U$ :

- (P) if a sequence  $\{u_n\}$  in  $U$  converges to  $u \in U$  and is such that  $\alpha(u_n, u_{n+1}) \geq 1$  for  $n \geq 1$ , then there is a subsequence  $\{u_{n(k)}\}$  of  $\{u_n\}$  with  $\alpha(u_{n(k)}, u) \geq 1$  for all  $k$ .

**Theorem 3.2.** *Let all the conditions of Theorem 3.1 hold, except that the condition (5) is replaced by*

(5') (P) holds.

*Then T has a best proximity point.*

*Proof.* As in the proof of Theorem 3.1 we conclude that there exists a sequence  $\{x_n\}$  in  $U_0$  which converges to  $z \in U_0$ . Using (3), we note that  $Tz \in V_0$  and hence

$$\sigma(u_1, Tz) = \sigma(U, V), \quad \text{for some } u_1 \in U_0.$$

Notice that from (P), we have  $\alpha(x_{n_k}, z) \geq 1$  for all  $k \in \mathbb{N}$ . Since  $T$  is  $\alpha$ -proximal admissible and

$$(3.10) \quad \sigma(u_1, Tz) = \sigma(x_{n_k+1}, Tx_{n_k}) = \sigma(U, V),$$

we obtain that  $\alpha(x_{n_k+1}, u_1) \geq 1$  for all  $k \in \mathbb{N}$  and

$$\zeta(\alpha(x_{n_k+1}, u_1)\sigma(u_1, x_{n_k+1}), \psi(\sigma(z, x_{n_k}))) \geq 0.$$

Then,  $(\zeta_2)$  implies that

$$\sigma(u_1, x_{n_k+1}) \leq \alpha(x_{n_k+1}, u_1)\sigma(u_1, x_{n_k+1}) \leq \psi(\sigma(z, x_{n_k})) < \sigma(z, x_{n_k})$$

and so  $\lim_{k \rightarrow \infty} \sigma(u_1, x_{n_k+1}) \rightarrow 0$ . Thus,  $u_1 = z$  and by (3.10) we have  $\sigma(z, Tz) = \sigma(U, V)$ . □

**Theorem 3.3.** *Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$ ,  $\zeta \in \mathcal{Z}$  and  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$ . Suppose that  $T : U \rightarrow V$  is an  $\alpha$ - $\zeta$ -contraction and that conditions (1)-(4) of Theorem 3.1 are satisfied, as well as*

(5'')  $T$  is continuous or (P) holds.

*Then, T has a best proximity point.*

*Proof.* By following the lines in the proof of Theorem 3.1, we easily construct a sequence  $\{x_n\}$  in  $U$  which converges to some  $z \in U$ , moreover

$$(3.11) \quad \lim_{n \rightarrow \infty} \sigma(x_n, z) = 0.$$

Suppose first that  $T$  is continuous. Then

$$(3.12) \quad \lim_{n \rightarrow \infty} \sigma(Tx_n, Tz) = 0.$$

From (3.3), the triangle inequality together with (3.11) and (3.12) imply

$$\sigma(U, V) = \lim_{n \rightarrow \infty} \sigma(x_{n+1}, Tx_n) = \sigma(z, Tz).$$

In other words,  $z \in U$  is a best proximity of the mapping  $T$ .

Suppose now that (P) holds. Regarding (3), we note that  $Tz \in V_0$  and hence

$$\sigma(u_1, Tz) = \sigma(U, V), \quad \text{for some } u_1 \in U_0.$$

Notice that from (P), we have  $\alpha(x_{n_k}, z) \geq 1$  for all  $k \in \mathbb{N}$ . Since  $T$  is  $\alpha$ -proximal admissible, and

$$\sigma(u_1, Tz) = \sigma(x_{n_k+1}, Tx_{n_k}) = \sigma(U, V),$$

we get that  $\alpha(x_{n_k+1}, u_1) \geq 1$  for all  $k \in \mathbb{N}$  and

$$(3.13) \quad \zeta(\alpha(x_{n_k+1}, u_1)\sigma(u_1, x_{n_k+1}), \sigma(z, x_{n_k})) \geq 0.$$

Then,  $(\zeta_2)$  implies that  $\sigma(u_1, x_{n_k+1}) \leq \alpha(x_{n_k+1}, u_1)\sigma(u_1, x_{n_k+1}) \leq \sigma(z, x_{n_k})$  and so

$$\lim_{k \rightarrow \infty} \sigma(u_1, x_{n_k+1}) \rightarrow 0.$$

Thus,  $u_1 = z$  and by (3.13) we have  $\sigma(z, Tz) = \sigma(U, V)$  and the proof is completed.  $\square$

Notice that Theorem 3.3 cannot be obtained by combining Theorems 3.1 and 3.2, since the function  $\psi(t) = t$  does not belong to  $\Psi$ . Furthermore, in Theorems 3.1 and 3.2, we have an additional condition that  $\zeta$  is non-decreasing in its second argument.

**Definition 3.3.** Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$ ,  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  and  $\zeta \in \mathcal{Z}$ . We say that  $T : U \rightarrow V$  is a generalized  $\alpha$ - $\zeta$ -contraction if  $T$  is  $\alpha$ -proximal admissible and

$$(3.14) \quad \alpha(x, y) \geq 1 \text{ and } \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) \Rightarrow \zeta(\alpha(x, y)\sigma(u, v), r(x, y)) \geq 0,$$

for all  $x, y, u, v \in U$  with  $x \neq y$ , where

$$r(x, y) = \max \left\{ \sigma(x, y), \frac{\sigma(x, u)\sigma(y, v)}{\sigma(x, y)} \right\}.$$

**Theorem 3.4.** Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$  and  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$ ,  $\zeta \in \mathcal{Z}$ . Suppose that  $T : U \rightarrow V$  is a generalized  $\alpha$ - $\zeta$ -contraction and conditions (1)-(5) of Theorem 3.1 are satisfied. Then  $T$  has a best proximity point.

*Proof.* As in the proof of Theorem 3.1, we can construct a sequence  $\{x_n\}$  in  $X$  satisfying conditions (3.3) and (3.4). Combining these relations with (3.14), we get that  $\sigma(x_n, Tx_{n-1}) = \sigma(x_{n+1}, Tx_n) = \sigma(U, V)$  for all  $n \in \mathbb{N}$  and

$$\zeta(\alpha(x_{n-1}, x_n)\sigma(x_n, x_{n+1}), r(x_{n-1}, x_n)) \geq 0, \quad \text{for all } n \in \mathbb{N}.$$

Here,

$$\begin{aligned} r(x_{n-1}, x_n) &= \max \left\{ \frac{\sigma(x_{n-1}, x_n)\sigma(x_n, x_{n+1})}{\sigma(x_{n-1}, x_n)}, \sigma(x_{n-1}, x_n) \right\} \\ &= \max \{ \sigma(x_n, x_{n+1}), \sigma(x_{n-1}, x_n) \}. \end{aligned}$$

Suppose that for some  $n \in \mathbb{N}$

$$\max \{ \sigma(x_n, x_{n+1}), \sigma(x_{n-1}, x_n) \} = \sigma(x_n, x_{n+1}).$$

Since  $\sigma(x_n, x_{n+1}) > 0$ , using the property (2) of the simulation function, we obtain

$$\zeta(\alpha(x_{n-1}, x_n)\sigma(x_n, x_{n+1}), \sigma(x_n, x_{n+1})) < 0,$$

which is a contradiction. It follows that  $r(x_{n-1}, x_n) = \sigma(x_{n-1}, x_n)$  for all  $n \in \mathbb{N}$ , implying that

$$(3.15) \quad \zeta(\alpha(x_{n-1}, x_n)\sigma(x_n, x_{n+1}), \sigma(x_{n-1}, x_n)) \geq 0, \quad \text{for all } n \in \mathbb{N}.$$



Using  $(\zeta_2)$ , the inequality (3.15) yields that

$$\sigma(x_n, x_{n+1}) \leq \sigma(x_{n-1}, x_n), \quad \text{for all } n \in \mathbb{N}.$$

Hence,  $\{\sigma(x_n, x_{n+1})\}$  is a non-increasing sequence, bounded from below, converging to some  $L \geq 0$ . Suppose that  $L > 0$ . Taking the upper limit as  $n \rightarrow \infty$  in (3.15), using  $(\zeta_3)$ , we get

$$0 \leq \limsup_{n \rightarrow \infty} \zeta(\alpha(x_n, x_{n-1})\sigma(x_n, x_{n+1}), \psi(\sigma(x_{n-1}, x_n))) < 0,$$

which is a contradiction. Hence, we conclude that  $\lim_{n \rightarrow \infty} \sigma(x_n, x_{n+1}) = 0$ .

In order to prove that  $\{x_n\}$  is a Cauchy sequence, suppose the contrary. Then, as in the proof of Theorem 3.1, there exist  $\epsilon > 0$  and subsequences  $\{x_{m_k}\}$  and  $\{x_{n_k}\}$  of  $\{x_n\}$ , so that for  $n_k > m_k > k$  we have

$$\sigma(x_{m_k}, x_{n_k}) \geq \epsilon \text{ and } \sigma(x_{m_k}, x_{n_k-1}) < \epsilon.$$

Also, in the same way, the following inequalities hold:

$$(3.16) \quad \begin{aligned} \lim_{k \rightarrow \infty} \sigma(x_{m_k}, x_{n_k}) &= \lim_{k \rightarrow \infty} \sigma(x_{n_k-1}, x_{m_k-1}) = \epsilon, \\ \lim_{k \rightarrow \infty} \sigma(x_{m_k-1}, x_{n_k}) &= \lim_{k \rightarrow \infty} \sigma(x_{n_k-1}, x_{m_k}) = \epsilon. \end{aligned}$$

Since  $T$  is triangular weakly- $\alpha$ -admissible, we derive that

$$\alpha(x_n, x_m) \geq 1, \quad \text{for all } n, m \in \mathbb{N}_0 \text{ with } n > m.$$

Thus, we have

$$(3.17) \quad \alpha(x_{m_k}, x_{n_k}) \geq 1 \text{ and } \sigma(x_{m_k}, Tx_{m_k-1}) = \sigma(x_{n_k}, Tx_{n_k-1}) = \sigma(U, V),$$

for all  $k \in \mathbb{N}$ . Since  $T$  is a generalized  $\alpha$ - $\zeta$ -contraction, the obtained relations (3.17) imply

$$0 \leq \zeta(\alpha(x_{m_k-1}, x_{n_k-1})\sigma(x_{m_k}, x_{n_k}), r(x_{m_k-1}, x_{n_k-1})), \quad \text{for all } k \in \mathbb{N}.$$

Since

$$(3.18) \quad r(x_{m_k-1}, x_{n_k-1}) = \max \left\{ \frac{\sigma(x_{m_k-1}, x_{m_k})\sigma(x_{n_k-1}, x_{n_k})}{\sigma(x_{m_k-1}, x_{n_k-1})}, \sigma(x_{m_k-1}, x_{n_k-1}) \right\},$$

taking limits of both sides of (3.18), we conclude that  $\lim_{k \rightarrow \infty} r(x_{m_k-1}, x_{n_k-1}) = \epsilon$ . Letting  $k \rightarrow \infty$  and keeping (3.16) and  $(\zeta_3)$  in mind, we get

$$0 \leq \limsup_{n \rightarrow \infty} \zeta(\alpha(x_{m_k-1}, x_{n_k-1})\sigma(x_{m_k}, x_{n_k}), r(x_{m_k-1}, x_{n_k-1})) < 0,$$

which is a contradiction. Thus, we conclude that the sequence  $\{x_n\}$  is Cauchy in  $U$ .

The final step of the proof is the same as for Theorem 3.1. □

4. COROLLARIES AND EXAMPLES

Using Example 2.1, it is possible to get a number of consequences of our main results by choosing the simulation function  $\zeta$  and  $\alpha(x, y)$  in a proper way. We skip making such a list of corollaries since they seem clear. We just state the following one as a sample

**Corollary 4.1.** *Let  $(X, \sigma)$  be a metric-like space,  $U$  and  $V$  be two non-empty subsets of  $X$  and  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$ ,  $\psi \in \Psi$ . Suppose that  $T : U \rightarrow V$  is a given  $\alpha$ -proximal admissible mapping such that*

$$\alpha(x, y) \geq 1 \text{ and } \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) \Rightarrow \alpha(x, y)\sigma(u, v) \leq \psi(\sigma(x, y)),$$

for all  $x, y, u, v \in U$ . Suppose also

- (a)  $T$  is triangular weakly- $\alpha$ -admissible;
- (b)  $U$  is closed with respect to the topology induced by  $\tau_\sigma$ ;
- (c)  $T(U_0) \subset V_0$ ;
- (d) there exist  $x_0, x_1 \in U$  such that  $\sigma(x_1, Tx_0) = \sigma(U, V)$  and  $\alpha(x_0, x_1) \geq 1$ ;
- (e)  $T$  is continuous or (P) holds.

Then,  $T$  has a best proximity point.

In particular, if the given space  $(X, \sigma)$  is also endowed with a partial order  $\preceq$ , by taking

$$\alpha(x, y) \geq 1 \Leftrightarrow x \succeq y,$$

one can get standard variations of the given results in a partially ordered space.

The following illustrative examples show how our results can be used for certain mappings acting in metric-like spaces.

*Example 4.1.* Consider  $X = \{a, b, c, d\}$  equipped with  $\sigma : X \times X \rightarrow \mathbb{R}_0^+$  defined by

$$\begin{aligned} \sigma(a, a) &= \frac{1}{2}, & \sigma(b, b) &= 0, & \sigma(c, c) &= 2, & \sigma(d, d) &= \frac{1}{3}, & \sigma(a, b) &= 3, \\ \sigma(a, c) &= \frac{5}{2}, & \sigma(a, d) &= \frac{3}{2}, & \sigma(b, c) &= 2, & \sigma(b, d) &= \frac{3}{2}, & \sigma(c, d) &= \frac{5}{2}, \end{aligned}$$

and  $\sigma(x, y) = \sigma(y, x)$  for  $x, y \in X$ . It is clear that  $(X, \sigma)$  is a complete metric-like space. Take  $U = \{b, c\}$  and  $V = \{c, d\}$ . Consider the mapping  $T : U \rightarrow V$  defined by  $Tb = d$ , and  $Tc = c$ . Remark that  $\sigma(U, V) = \sigma(b, d) = \frac{3}{2}$ . Also,  $U_0 = \{b\}$  and  $V_0 = \{d\}$ . Note that  $T(U_0) \subseteq V_0$ . Take  $\psi(t) = \frac{5}{6}t$ , and  $\zeta(t, s) = \frac{3}{4}s - t$  for all  $t, s \geq 0$ . Define  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  by

$$\alpha(x, y) = \begin{cases} 1, & x, y \in U, \\ 0, & \text{otherwise.} \end{cases}$$

Let  $x, y, u, v \in U$  be such that

$$\alpha(x, y) \geq 1 \text{ and } \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) = \frac{3}{2}.$$

Then, necessarily, we have  $x = y = u = v = b$ . So,  $\alpha(u, v) \geq 1$ , that is,  $T$  is  $\alpha$ -proximal admissible.

We need to prove that  $T$  is an  $\alpha$ - $\psi$ - $\zeta$  contraction. By the previous conclusion, the only case to be checked is when  $x = y = u = v = b$ . Then we have

$$\zeta(\alpha(b, b)\sigma(b, b), \psi(\sigma(b, b))) = \zeta(1 \cdot 0, \psi(0)) = 0.$$

Thus, all the conditions of Theorem 3.1 are satisfied. So  $T$  has a best proximity point (which is  $z = b$ ). On the other hand, e.g., Corollary 2.2 (with  $k = 2$ ) of [4] is not applicable for the standard metric.

*Example 4.2.* Consider the set  $X = \{a, b, c, d\}$  equipped with the following complete metric-like  $\sigma$ :

$$\begin{aligned} \sigma(a, a) = \sigma(b, b) &= \frac{1}{4}, & \sigma(c, c) = \sigma(d, d) &= 2, \\ \sigma(a, b) = \sigma(c, d) &= \frac{1}{2}, & \sigma(a, c) = \sigma(b, d) &= 1, & \sigma(a, d) = \sigma(b, c) &= \frac{3}{2}, \end{aligned}$$

and  $\sigma(x, y) = \sigma(y, x)$  for all  $x, y \in X$ . Let  $U = \{a, b\}$  and  $V = \{c, d\}$ ; then  $\sigma(U, V) = 1$ ,  $U_0 = U$  and  $V_0 = V$ . Consider, further, the mappings  $T : U \rightarrow V$  given by  $Ta = c$ ,  $Tb = c$ ,  $\alpha : X \times X \rightarrow [0, +\infty)$  given by

$$\alpha(x, y) = \begin{cases} 1, & \text{if } x, y \in U, \\ 0, & \text{otherwise,} \end{cases}$$

and  $\zeta \in \mathcal{Z}$  given by  $\zeta(t, s) = s - \frac{2+t}{1+t}t$ . Let us check that the mapping  $T$  is a generalized  $\alpha$ - $\zeta$ -contraction. Let  $x, y, u, v \in U$  be such that  $x \neq y$ ,  $\alpha(x, y) \geq 1$ ,  $\sigma(u, Tx) = \sigma(v, Ty) = 1$ . Then it must be  $u = v = a$  and either  $x = a$ ,  $y = b$  or  $x = b$ ,  $y = a$ . In both cases, it is  $\alpha(u, v) \geq 1$ . In order to check condition (3.14), it is enough to consider the case  $x = a$ ,  $y = b$ ,  $u = v = a$  (the other is treated symmetrically). Then,

$$\begin{aligned} \zeta(\alpha(x, y)\sigma(u, v), r(x, y)) &= \zeta\left(1 \cdot \frac{1}{4}, \max\left\{\frac{1}{2}, \frac{\frac{1}{4} \cdot \frac{1}{2}}{\frac{1}{2}}\right\}\right) = \zeta\left(\frac{1}{4}, \frac{1}{2}\right) \\ &= \frac{1}{2} - \frac{2 + \frac{1}{4}}{1 + \frac{1}{4}} \cdot \frac{1}{4} = \frac{1}{20} > 0, \end{aligned}$$

and the condition is satisfied. All other conditions of Theorem 3.4 are fulfilled, hence, we conclude that the mapping  $T$  has a best proximity point (which is  $z = a$ ).

### 5. APPLICATION TO BEST PROXIMITY RESULTS ON A METRIC-LIKE SPACE WITH A GRAPH

Throughout this section,  $(X, \sigma)$  will denote a metric-like space and  $G = (V(G), E(G))$  will be a directed graph such that its set of vertices  $V(G) = X$  and the set of edges  $E(G)$  contains all loops, i.e.,  $\Delta := \{(x; x) : x \in X\} \subseteq E(G)$ . We need in the sequel the following hypothesis:

( $P_G$ ) if a sequence  $\{u_n\}$  in  $X$  converges to  $u \in A$  such that  $(u_n, u_{n+1}) \in E(G)$ , then there is a subsequence  $\{u_{n(k)}\}$  of  $\{u_n\}$  with  $(u_{n(k)}, u) \in E(G)$  for all  $k$ .

**Definition 5.1.** Let  $U$  and  $V$  be two non-empty subsets of  $X$  and  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$ . We say that  $T : U \rightarrow V$  is a  $G$ -proximal mapping if

$$(5.1) \quad \left. \begin{array}{l} (x, y) \in E(G), \quad \alpha(x, y) \geq 1, \\ \sigma(u, Tx) = \sigma(v, Ty) = \sigma(U, V) \end{array} \right\} \Rightarrow (u, v) \in E(G),$$

for all  $x, y, u, v \in U$ .

**Definition 5.2** ([8, 10]). Let  $U$  and  $V$  be two non-empty subsets of  $X$ , let  $T : U \rightarrow V$  be a mapping and  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  be a function. We say that  $T$  is triangular weakly- $G$ -admissible if

$$\alpha(x, y) \in E(G) \text{ and } \alpha(y, z) \in E(G) \Rightarrow \alpha(x, z) \in E(G).$$

**Corollary 5.1.** Let  $U$  and  $V$  be two non-empty subsets of  $X$  and  $\psi \in \Psi$ . Suppose that  $T : U \rightarrow V$  is a mapping such that

$$\sigma(Tx, Ty) \leq \psi(\sigma(x, y)),$$

for all  $x, y \in U$  such that  $(x, y) \in E(G)$ . Suppose also:

- (a)  $T$  is triangular weakly- $G$ -admissible;
- (b)  $T(U_0) \subset V_0$ ;
- (c) there exist  $x_0, x_1 \in U$  such that  $\sigma(x_1, Tx_0) = \sigma(U, V)$  and  $(x_0, x_1) \in E(G)$ ;
- (d)  $T$  is continuous or  $(R_G)$  holds.

Then,  $T$  has a best proximity point.

*Proof.* It suffices to consider  $\alpha : X \times X \rightarrow \mathbb{R}_0^+$  such that

$$\alpha(x, y) = \begin{cases} 1, & \text{if } (x, y) \in E(G), \\ 0, & \text{if not.} \end{cases}$$

All the hypotheses of Corollary 4.1 are satisfied. □

In this way, we can derive all results and consequences of the paper [15], extending them to partially ordered metric-like spaces. Similarly, we can extend the frame of several other existing results from, e.g., [3, 10, 12, 16].

**Acknowledgements.** The authors are thankful to the referee for his/her careful reading of the article.

## REFERENCES

- [1] H. H. Alsulami, E. Karapinar, F. Khojasteh and A. F. R.-L. de Hierro, *A proposal to the study of contractions in quasi-metric spaces*, Discrete Dyn. Nat. Soc. **2014** (2014), Article ID 269286.
- [2] A. Amini-Harandi, *Metric-like spaces, partial metric spaces and fixed points*, Fixed Point Theory Appl. **2012**(204) (2012), 10 pages.

- [3] N. Bilgili, E. Karapinar and B. Samet, *Generalized  $\alpha$ - $\psi$ -contractive mappings in quasi-metric spaces and related fixed-point theorems*, J. Inequal. Appl. **2014**(36) (2014), 15 pages.
- [4] S. Chandok and M. Postolache, *Fixed point theorem for weakly Chatterjea-type cyclic contractions*, Fixed Point Theory Appl. **2013**(28) (2013), 9 pages.
- [5] A. F. R.-L. de Hierro, E. Karapinar, C. R.-L. de Hierro and J. Martínez-Moreno, *Coincidence point theorems on metric spaces via simulation functions*, J. Comput. Appl. Math. **75** (2015), 345–355.
- [6] M. Jleli, E. Karapinar and B. Samet, *Best proximity points for generalized  $\alpha$ - $\psi$ -proximal contractive type mappings*, J. Appl. Math. **2013** (2013), Article ID 534127, 10 pages.
- [7] E. Karapinar, *Fixed points results via simulation functions*, Filomat **30**(8) (2016), 2343–2350 .
- [8] E. Karapinar, H. H. Alsulami and M. Noorwali, *Some extensions for Geraghty type contractive mappings*, J. Inequal. Appl. **2015**(303) (2015), 22 pages.
- [9] E. Karapinar and F. Khojasteh, *An approach to best proximity points via simulation functions*, J. Fixed Point Theory Appl. **19** (2017), 1983–1995.
- [10] E. Karapinar, P. Kuman and P. Salimi, *On  $\alpha$ - $\psi$ -Meir-Keeler contractive mappings*, Fixed Point Theory Appl. **2013** (2013), Article ID 94.
- [11] E. Karapinar and P. Salimi, *Dislocated metric space to metric spaces with some fixed point theorems*, Fixed Point Theory Appl. **2013** (2013), Article ID 222.
- [12] E. Karapinar and B. Samet, *Fixed point theorems for generalized  $\alpha$ - $\psi$  contractive type mappings and applications*, Abstr. Appl. Anal. **2012** (2012), 17 pages.
- [13] F. Khojasteh, S. Shukla and S. Radenović, *A new approach to the study of fixed point theorems via simulation functions*, Filomat **29** (2015), 1189–1194.
- [14] S. Radenović, Z. Kadelburg, D. Jandrlić and A. Jandrlić, *Some results on weak contraction maps*, Bull. Iranian Math. Soc. **38** (2012), 625–645.
- [15] B. Samet, *Best proximity point results in partially ordered metric spaces via simulation functions*, Fixed Point Theory Appl. **2015**(232) (2015), 15 pages.
- [16] B. Samet, C. Vetro and P. Vetro, *Fixed point theorem for  $\alpha$ - $\psi$  contractive type mappings*, Nonlinear Anal. **75** (2012), 2154–2165.

<sup>1</sup>DEPARTMENT OF MATHEMATICS,  
ICFAI UNIVERSITY,  
RAIPUR, CHHATTISGARH-490042, INDIA  
Email address: gvvjagan1@gmail.com

<sup>2</sup>DEPARTMENT OF MATHEMATICS,  
SCHOOL OF ADVANCED SCIENCES,  
VELLORE INSTITUTE OF TECHNOLOGY,  
VELLORE-632014, TN, INDIA  
Email address: drhknashine@gmail.com, hemant.nashine@vit.ac.in

<sup>3</sup>FACULTY OF MATHEMATICS,  
UNIVERSITY OF BELGRADE,  
STUDENTSKI TRG 16, 11000 BEOGRAD, SERBIA  
Email address: kadelbur@matf.bg.ac.rs

\*CORRESPONDING AUTHOR