REVIEW



### **Beyond RGD: virus interactions with integrins**

Hosni A. M. Hussein<sup>1</sup> · Lia R. Walker<sup>1</sup> · Usama M. Abdel-Raouf<sup>2</sup> · Sayed A. Desouky<sup>2</sup> · Abdel Khalek M. Montasser<sup>3</sup> · Shaw M. Akula<sup>1</sup>

Received: 27 May 2015/Accepted: 21 August 2015/Published online: 1 September 2015 © Springer-Verlag Wien 2015

Abstract Viruses successfully infect host cells by initially binding to the surfaces of the cells, followed by an intricate entry process. As multifunctional heterodimeric cell-surface receptor molecules, integrins have been shown to usefully serve as entry receptors for a plethora of viruses. However, the exact role(s) of integrins in viral pathogen internalization has yet to be elaborately described. Notably, several viruses harbor integrin-recognition motifs displayed on viral envelope/capsid-associated proteins. The most common of these motifs is the minimal peptide sequence for binding integrins, RGD (Arg-Gly-Asp), which is known for its role in virus infection via its ability to interact with over half of the more than 20 known integrins. Not all virus-integrin interactions are RGD-dependent, however. Non-RGD-binding integrins have also been shown to effectively promote virus entry and infection as well. Such virus-integrin binding is shown to facilitate adhesion, cytoskeleton rearrangement, integrin activation, and increased intracellular signaling. Also, we have attempted to discuss the role of carbohydrate moieties in virus interactions with receptor-like host cell surface integrins that drive the process of internalization. As much as possible, this article examines the published literature regarding the role of integrins in terms of virus infection and virus-encoded glycosylated proteins that mediate interactions with integrins, and it explores the idea of targeting these receptors as a therapeutic treatment option.

#### Introduction

Viruses may be small in size, but they carry enough genetic material that they are capable of inflicting some of the deadliest diseases in the world. If not for their ability to enter host cells and efficiently impair them, we would not even talk about them. An efficient pathogen is one that has evolved a robust entry mechanism for delivery of genetic material into different target host cells, which is critical for replication and sustenance. Over the years, viruses as obligate parasites have evolved successful ways to colonize host cells, using complicated but well-orchestrated mechanisms to enter cells.

The whole process of virus entry-otherwise referred to as internalization-begins with the virus binding to target cells. Binding to cells is a reversible process that does not ensure virus entry. Virus binding or attachment only ensures viral proximity to cells. However, this process is the most essential step that kick-starts the whole cascade of events, resulting in the eventual internalization of the virus. Several viruses utilize different glycosaminoglycans expressed on the target cells as binding receptors. Glycosaminoglycans serve as good receptor molecules that promote binding, as they are expressed ubiquitously in eukaryotic cells. Some of the most common binding receptors are heparan sulfate (HS) and chondroitin sulfate [4, 82, 99, 143, 167]. Virus binding to such receptors brings them closer to cells and provides the opportunity to interact with other receptor molecules that promote the actual have internalization process. Several scientists

Shaw M. Akula akulas@ecu.edu

<sup>&</sup>lt;sup>1</sup> Department of Microbiology and Immunology, Brody School of Medicine, East Carolina University, Greenville, NC 27834, USA

<sup>&</sup>lt;sup>2</sup> Faculty of Science, Al Azhar University, Assiut Branch, Assiut 71524, Egypt

<sup>&</sup>lt;sup>3</sup> Brucellosis Research Department, Animal Health Research Institute, Dokki, Egypt

hypothesized that the initial step of virus binding to cells also induces conformational changes to the glycoproteins expressed on the target cells that are critical for the virus to interact with other receptor molecules, thus promoting internalization.

The actual virus entry is complicated to the extent that a single virus may utilize different receptors to efficiently enter different target cells [6, 19]. Viruses have evolved such mechanisms to be effective pathogens. Such variations in the entry mechanisms dictate the actual entry route. For example, Epstein-Barr virus (EBV) enters lymphoblastoid cells by fusion but enters B cells via endocytosis [112]. Integrins are a family of receptor molecules that serve as entry receptors for a variety of different viruses, including foot-and-mouth disease virus (FMDV) [97], Kaposi's sarcoma-associated herpesvirus (KSHV) [5], herpes simplex virus-2 (HSV-2) [31], adenovirus [168], human papillomavirus-16 (HPV-16) [3], reovirus [40], and others. Our understanding of the role of integrins in promoting virus entry is still not complete. In this review, we have attempted to elaborate on the role of integrins in virus internalization.

#### What are integrins?

Integrins play an important role in regulating a variety of cellular functions, including cell adhesion, cell migration, and critical signaling processes. This is possible because of their ability to interact with various ligands, including extracellular matrix glycoproteins (i.e., collagens, fibronectins, laminins, etc.) and cellular receptors (i.e., vascular cell adhesion molecule-1 and intercellular cell adhesion molecules) [85, 86, 124]. Discovered over twenty years ago, integrins are a large family of transmembrane glycoproteins found in a variety of organisms ranging from sponges, corals, nematodes, and echinoderms to mammals [20].

There are about 24 integrins that have been identified. These heterodimeric receptor molecules result from different pairings among 18  $\alpha$  and 8  $\beta$  subunits [86]. Each integrin subunit has three domains: an extracellular, transmembrane, and cytoplasmic domain. The extracellular domain is the largest part, ranging from 80 to 150 kDa, while the cytoplasmic domain is a short and largely unstructured domain of 10-70 amino acid (aa) residues, with the exception of the b4 subunit, which contains >1,000 aa residues [148]. The transmembrane domains of integrins are single-spanning structures comprised of 25-29 aa residues that form  $\alpha$ -helical coiled coils that exist as either homo- or heterodimers [2].

High-resolution X-ray crystallography structural data are available for the extracellular domains of integrins [9,

98, 183], but no high-resolution experimental X-ray crystal structures are available for the transmembrane or cytoplasmic domain of any integrin heterodimer. Much of the structural data of the transmembrane and cytoplasmic domains are based purely on NMR analysis. Integrins can shift between high- and low-affinity conformations for ligand binding to transduce intracellular signals following ligand binding. In the inactive state, the extracellular domain of integrins is not bound to ligands and exists in a bent conformation. However, signals from the cell induce conformational changes that expose the external ligand-binding site, where ligands bind and transmit the signals from outside to inside the cell [148].

Although some integrins can bind their ligands in a resting state, there are other integrins whose binding to their ligand requires activation through alterations in the intracellular domains by signaling events, which subsequently lead to transmission of signals from inside to outside of the cell; this is commonly referred to as insideout signaling [37]. Ligand binding to extracellular domains of integrins leads to activation of integrins and subsequent transmission of cellular signals from outside to inside of the cell, which is known as outside-in signaling [182]. These intracellular signals are very important for cell growth, differentiation, and apoptosis. Additionally, intracellular signals lead to formation of the focal adhesion complex, which is a large and dynamic multi-protein complex that includes a vast number of intracellular proteins [185].

Along with proteoglycans, integrins form the major adhesion receptors for extracellular cellular matrix (ECM) proteins, making them important for signaling events that determine cell fate [94]. Cellular signaling processes depend on the pattern of expression and the composition of integrins, which determine the ECM type a cell can bind and initiate downstream signaling events [78]. Integrins provide a connection between the ECM proteins and the actin cytoskeleton that is crucial for regulating cytoskeletal organization and intracellular signaling pathways, all of which are a necessity for cell survival, proliferation, shape, attachment, migration, and angiogenesis [148].

As adhesion molecules, integrins mediate cell-to-cell, cell-to-ECM, and cell-to-pathogen interactions, and such adhesion is regulated through the inside-out signaling process. Integrin-induced adhesion is very important in the regulation of the immune system during leukocyte trafficking, migration, and phagocytosis [108]. Adhesion of integrins to a solid surface is the first step in cell migration and motility. The vital role of integrins in cell migration makes them essential for many important biological events, including embryonic development [58, 189], inflammatory responses [50, 55, 101], wound healing [7, 24], and tumor metastasis [81]. There are many pathogens, including

viruses [45, 89, 146] and bacteria [76, 80], that have the ability to use integrins with different mechanisms for invading cells. Through regulation of several cell functions, integrins have a role in human disease. For example, tumors, cancer, and immunodeficiency disorders are all associated with altered integrin-mediated adhesion and migration [84].

The hallmark of tumor development is cell attachment, migration, and proliferation; all of which are regulated by integrin-based cellular signaling [71, 113, 170]. Expression of particular integrins, including  $\alpha 6\beta 4$ ,  $\alpha 6\beta 1$ ,  $\alpha V\beta 5$ ,  $\alpha 2\beta 1$  and  $\alpha 3\beta 1$ , on tumor cells in the context of activated cytokine receptors or growth factor receptors leads to increased disease progression and severity [42]. Integrins integrate the extracellular and intracellular environments by binding to ligands outside the cell and cytoskeletal components and signaling molecules inside the cell [108]. The role of integrins in cancer initiation and progression makes them targets for several therapeutic agents in clinical trials of cancer therapy.

## What are the roles of integrins in virus entry of cells?

Integrins have been exploited by many pathogens, including bacteria and viruses to infect cells. Penetration of the host-cell plasma membrane is a crucial step for a successful virus infection [178]. To invade the host cell, several animal viruses physically interact with integrins to infect cells. There are many studies that demonstrate the critical role of membrane rafts in viral entry and infection [22, 153]. Interestingly, many integrins used by viruses for binding and internalization are localized to and associated with membrane rafts, which consolidate the role of integrin in virus internalization and infection [30, 70, 116, 153]. A list of viruses and the manner by which they utilize integrins to infect cells is provided in Table 1.

## **RGD-binding integrins and their effect on virus infection of cells**

Many viruses, including adenoviruses and herpesviruses, have an RGD (Arg-Gly-Asp) tripeptide motif displayed on their viral envelope glycoproteins. RGD is the minimal peptide sequence for binding integrins. As an integrinrecognition motif, RGD plays an important role in virus infection by binding one or different combinations of several integrins, which include  $\alpha V\beta 1$ ,  $\alpha V\beta 3$ ,  $\alpha V\beta 5$ ,  $\alpha 5\beta 1$ ,  $\alpha V\beta 6$ ,  $\alpha V\beta 8$ , and  $\alpha IIb\beta 3$  [28, 127, 132, 140]. RGD binding to these integrins activates cellular signals such as phosphatidylinositol-3-kinase (PI-3K) and mitogen-activate protein kinase (MAPK) pathways, which are critical for supporting virus infection of cells [5, 106].

Many serotypes of adenovirus use RGD-binding integrins to stimulate endocytosis and thereby promote virus entry [41, 106, 177]. Binding of the RGD motif on the adenovirus penton base capsid protein to integrins initiates virus internalization by stimulating endocytosis via clathrin-coated vesicles [11, 104, 150]. Philpott and colleagues demonstrated that blocking adenovirus binding to integrins using an RGD peptide resulted in a 2- to 3-fold reduction in viral DNA intake [123]. Likewise, studies by Shayakhmetov and colleagues revealed that the deletion of the RGD motif in the penton base did not affect virus attachment but significantly reduced the rate of virus internalization, specifically at the step that involves endosomal escape [140].

Several members of the family *Herpesviridae* interact with integrins in an RGD-dependent manner. For example, herpes simplex virus type 1 (HSV-1) envelope-associated gH interacts with  $\alpha V\beta3$ , which is critical for virus entry of cells [122]. KSHV or human herpesvirus 8 (HHV-8) interacts with a variety of cellular integrins, including  $\alpha 3\beta1$ ,  $\alpha V\beta3$ , and  $\alpha V\beta5$ , and activates focal adhesion kinase (FAK), Src, PI-3K, Rho GTPases, and diaphanous 2 (Dia2)-associated signaling, which is a necessity for the internalization of the virus [28, 117]. The ability of KSHV to interact with integrins is mediated by the RGD motif of envelope-associated gB [5, 61, 74]. Interestingly, the RGD motif of gB is also required to mediate attachment of cells to the endothelium [44].

RGD-binding integrins are also important for other viruses. For example, the interaction between RGD of capsid protein VP1 of coxsackievirus A9 and  $\alpha V\beta 3$  is essential for virus binding and entry into cells [77, 130, 131]. There are also other studies that demonstrate that a high-affinity interaction between RGD of coxsackievirus A9 and  $\alpha V\beta 6$  (compared to  $\alpha V\beta 3$ ) is important for cell entry and virus uncoating [77, 138, 180]. Notably, the RGD sequence is highly conserved in the VP1 protein of FMDV and mediates virus attachment to integrins, and thus internalization [47]. Synthetic RGD peptides have been shown to block FMDV attachment in a dose-dependent manner [57, 88, 119]. HIV-1 Tat protein interacts with RGD-binding  $\alpha V\beta 5$ ,  $\alpha V\beta 3$ , and  $\alpha 5\beta 1$  and initiates the integrin endocytic pathway, which is essential for entry of the virus [13, 160, 173]. HIV-1 Tat protein's interaction with RGD-binding integrin(s) is important for adhesion of target cells [12, 114]. Interactions of the more deadly Ebola virus with  $\alpha 5\beta 1$  is deemed critical for modulating infection of fibroblasts [135]. Thus, RGD interactions between viral proteins and integrins seem to regulate not only virus infection of cells but also the associated pathogenesis.

Table 1 Integrins used by different viruses and their role in virus infection

Virus	Integrins	Role of integrins	References
Human adenovirus type 2/5	ανβ3, ανβ5, ανβ1, α5β1, αLβ2, αΜβ2	Cell entry, endosome escape	[32, 68, 151, 177, 181]
Human cytomegalovirus (HCMV)	α2β1, α6β1, ανβ3	Cell entry	[53]
Kaposi's sarcoma-associated herpesvirus (KSHV)	α3β1, αVβ3, αVβ5, α9β1	Cell entry	[5, 28, 169, 174]
Epstein-Barr virus (EBV)	αVβ3, αVβ5, α5β1	Cell entry	[166]
Human immunodeficiency virus 1 (HIV-1)	α4β7, αVβ5, αVβ3, α5β1	Cell attachment	[10, 35, 114, 173]
HPS-associated hantaviruses NY-1 and Sin Nombre virus (SNV)	αVβ3, αllbβ3	Cell attachment, entry	[62, 127]
Rotavirus	α4β1, α4β7, α2β1, αVβ3, αxβ2	Cell attachment, entry	[66, 67, 69, 107]
Echovirus types 1	α2β1, αVβ3	Cell attachment, entry	[49, 92, 162]
Echovirus types 9	αVβ3	Cell attachment, entry	[150]
Foot-and-mouth disease virus (FMDV)	α5β1, αVβ3, αVβ6, αVβ8, αVβ1	Cell attachment, entry	[88, 100, 176]
Coxsackievirus A9	αVβ3, αVβ6	Cell attachment, entry	[129, 138, 150, 164]
Murine polyomavirus	α4β1	Cell entry	[25]
Vaccinia virus	β1	Cell entry	[87]
West Nile virus	ανβ3, ανβ1	Cell entry	[33, 134]
Simian virus 40	α2β1	Cell attachment, entry	[149]
Ross River (RR) virus	α1β1	Cell attachment, entry	[96]
Human papillomavirus	α6β4	Cell attachment	[37, 184]
Ebola virus	α5β1	Cell entry	[135]

### Do viruses utilize non-RGD-binding integrins to enter cells?

Not all integrins recognize and interact with the conserved RGD motif of viral proteins. There are multiple other non-RGD-binding integrins that drive virus entry and infection [105, 150]. Some of the non-RGD-binding integrins that promote entry and infection of HCMV [54], KSHV [174], simian virus 40 (SV40) [149], and Ross River virus (RRV) [96] are  $\alpha 1\beta 1$ ,  $\alpha 2\beta 1$ ,  $\alpha 4\beta 1$ ,  $\alpha 6\beta 1$ ,  $\alpha 9\beta 1$  and  $\alpha x\beta 2$ .

 $\alpha 1\beta 1$  and  $\alpha 2\beta 1$  are collagen receptors that are utilized by viruses for cell entry and infection. RRV is an alphavirus that is endemic to Australia and New Guinea and is etiologically associated with epidemic polyarthritis [75, 115]. RRV interacts with  $\alpha 1\beta 1$  to infect target cells [60, 96]. Infection of RRV has been shown to be blocked by function-blocking antibodies to  $\alpha 1\beta 1$ , soluble  $\alpha 1\beta 1$  integrin, or peptides representing the  $\alpha 1\beta 1$  integrin-binding site on collagen IV [96]. Rotavirus utilizes different non-RGD binding domains to interact with integrins. (i) The Tyr-Gly-Leu (YGL) sequence of the rotavirus spike protein, VP4, interacts with  $\alpha 4\beta 1$  and  $\alpha 4\beta 7$  and helps the rotavirus to bind and enter cells [66], and (ii) the GPR sequence in the rotavirus spike protein VP7 interacts with  $\alpha x \beta 2$  and helps the rotavirus to enter cells [64, 65]. Integrins also have a critical role to play in rotavirus pathogenesis. Integrin  $\alpha 1\beta 1$  and  $\alpha 2\beta 1$  are important receptors for enterotoxin function and pathogenesis through their interactions with rotavirus NSP4, which eventually results in diarrhea [137].

Human echovirus 1 (EV1), which belongs to the family Picornaviridae (a family of RNA viruses) is implicated in many human diseases, including meningoencephalitis and carditis. EV1 successfully infects cells by interacting with  $\alpha 2\beta 1$  integrin in an RGD-independent manner. In fact,  $\alpha 2\beta 1$ clustering on the surface of cells is the determining factor that defines the success rate of the EV1-mediated signaling pathway and virus infection of cells [15, 92, 186]. Also, HIV-1 interacts with the non-RGD-binding integrin  $\alpha 4\beta 7$  via gp120, which is critical for efficient cell-to-cell spread of the virus [10, 35, 72]. More interestingly, Cicala et al. hypothesize that gp120- $\alpha$ 4 $\beta$ 7 interactions play an important role in the very early events following sexual transmission of HIV and may have important implications in the design of vaccine strategies for the prevention of acquisition of HIV infection [36]. Stergiou et al., determined that  $\alpha 2\beta 1$  plays a crucial role in modulating SV40-induced cellular signaling and infection [149]. There is also evidence that indicates a role for  $\alpha 2\beta 1$ integrin in promoting human papillomavirus (HPV)-induced squamous epithelial dysplasia [161].

Apart from the traditional RGD motif, herpesvirus glycoprotein B (gB) possesses a disintegrin-like domain (DLD) [53, 54]. A role for DLD in HCMV and KSHV entry and infection has been described recently. HCMV interacts with  $\alpha 2\beta 1$  and  $\alpha 6\beta 1$  [53], and KSHV interacts with  $\alpha 9\beta 1$  [174] via the DLD contained within gB to successfully enter cells. The minimum component of the disintegrin module required for integrin engagement is the 12- to 13-amino-acid disintegrin loop, for which a consensus sequence has been described:  $RX_6DLXXF$  [48]. In the case of KSHV gB, the DLD sequence is  $RX_{5-7}D/ELXXF/LX_5C$  (aa 66-85; with a conservative D to E substitution). Overall, viruses seem to use non-RGD- and RGD-binding integrins to a comparable extent as a means of binding and entering cells.

#### Comparison of the roles of RGD- versus non-RGDbinding integrins in assisting virus infection of cells

Integrins play a crucial role in cellular function through interactions with a variety of ligands. Integrins are ligand specific and can be grouped into four major groups: laminin-binding integrins ( $\alpha 1\beta 1$ ,  $\alpha 2\beta 1$ ,  $\alpha 3\beta 1$ ,  $\alpha 6\beta 1$ ,  $\alpha 7\beta 1$ , and  $\alpha 6\beta 4$ ), collagen-binding integrins ( $\alpha 1\beta 1$ ,  $\alpha 2\beta 1$ ,  $\alpha 3\beta 1$ ,  $\alpha 10\beta 1$ , and  $\alpha 11\beta 1$ ), leukocyte integrins ( $\alpha L\beta 2$ ,  $\alpha M\beta 2$ ,  $\alpha X\beta 2$ , and  $\alpha D\beta 2$ ), and RGD-recognizing integrins ( $\alpha 5\beta 1$ ,  $\alpha V\beta 1$ ,  $\alpha V\beta 3$ ,  $\alpha V\beta 5$ ,  $\alpha V\beta 6$ ,  $\alpha V\beta 8$ , and  $\alpha IIb\beta 3$ ) [155]. Interestingly, RGD and non-RGD-binding integrins aid equally in the internalization of viruses (Table 2).

#### The key cellular components that are altered by interactions of integrin with viruses

Virus-integrin binding induces changes in the quaternary structure of the integrin resulting in clustering of subunits, which increases virus affinity, cytoskeletal rearrangement,

Table 2 Functions of RGD and non-RGD binding integrins

Class	Integrin	Functions in virus infection	References
RGD binding	αVβ1 αVβ3 αVβ5 α5 β1 αVβ6 αVβ8 αΠbβ3 αMβ2 αLβ2 α281	Cell attachment, entry, signaling, and endosome escape	[5, 62, 83, 100, 107, 127, 140, 150, 176]
Non- RGD binding	α1β1 α2β1 α4β1 α6β1 α9β1 αxβ2 α6β4	Cell attachment, entry, signaling, and endosome escape	[10, 25, 28, 32, 35, 37, 53, 68, 96, 169, 173]

and subsequent virus internalization [150]. These conformational changes are critical for integrins to achieve outside-in and inside-out signaling necessary for various cellular functions, including cytoskeleton activation, endocytosis, gene expression, cell motility, attachment, cell cycle, cell growth, apoptosis, and differentiation [28, 108]. On the other hand, interactions of viruses with cellular integrins induce conformational changes in the viral surface proteins, helping to expose the essential domains required for virus entry into a host cell [107].

Through integrin activation, viruses can induce FAK phosphorylation which is followed by the activation of several focal adhesion-associated signal molecules, including Src, PI-3K, Rho GTPases (RhoA, Rac, and Cdc42), Dia2, and other effector molecules, such as AKT, ezrin, protein kinase C (PKC), MAPK (MEK, ERK1/2), NF-kB, and p38MAPK [5, 28]. Focal adhesion and associated molecules play critical roles in mediating the internalization of viral DNA into target cells [1, 95].

Src is one of the cellular components that is activated immediately upon activation of FAK by virusintegrin interactions. Src-mediated tyrosine phosphorylation of clathrin regulates clathrin translocation to the plasma membrane, which is important for interactions of clathrin with a number of other essential proteins, including AP2, Eps15, and dynamin. Src-mediated tyrosine phosphorylation also plays a role in endocytosis by releasing the internalized endocytic vesicles and initiating the assembly of the plasma-membrane-associated Ras activation complex [21, 26]. PI-3K and Ras are directly responsible for activating Rho and Rab GTPases. These GTPases, along with the activated ERK1/2, are critical for the microtubule and microfilament reorganization that determines the formation of various types of endocytic vesicles and their movements, as well as acting as a force to drive the virus inside and closer to the nuclear membrane [28].

Many viruses use microtubules and cytoplasmic dynein for nuclear targeting [43]. Virus-induced activity of Rho GTPases such as RhoA and Rac1 increase the efficiency of viral trafficking along microtubules to the nucleus. Thus, inactivation of these Rho proteins affects the stability of microtubules, thereby limiting the delivery of viral nucleic acids into the nucleus [110]. Ezrin is the best-studied member of the downstream effector molecules induced by Rho GTPases, and it is a critical element for cross-linking the actin cytoskeleton with the plasma membrane and inducing the morphological changes that are commonly observed in cells being infected by viruses and other pathogens [139]. Rac, Rho, Cdc42, and Rab5 also act as switching molecules that are essential for internalization of any pathogen due to their ability to modulate actin dynamics, formation of endocytic vesicles and their fission,

cytoskeletal transport, endosome movement, fusion of endocytic vesicles, and recycling [1, 27, 139].

Adenoviruses, echoviruses, FMDV, parechoviruses, parvoviruses, rotaviruses, KSHV, hantaviruses, and others enter target cells via endocytosis by physically interacting with integrins, resulting in the activation of the FAK-Src-PI-3K signaling pathway [111, 139, 179]. PI-3K-induction in a integrin-FAK-Src-dependent manner plays an important role in virus entry and infection via activation of the Rho family of GTPases and ezrin, and mediates actin cytoskeleton reorganization. Interestingly, these events initiate a cascade of intracellular signals that eventually activate the mitogen-activated protein kinase (MAPK) pathways, which are very important in modulating a variety of cellular processes, including proliferation, differentiation, survival, and apoptosis [117, 163]. In terms of cellular machinery, actin cytoskeleton reorganization is the crux that supports integrin-associated signaling-induced virus entry. Apart from working as a structural platform stabilizing cellular signaling molecules, actin provides mechanical force for endosome formation and endocytic vesicle propulsion [139, 157]. Microtubules and microfilaments along with other cytoskeletal elements play an important role in controlling the intracellular movement of many viruses [39, 102, 103, 118, 126].

# A key role for carbohydrate moieties on the cell surface in virus-integrin interactions

Successful virus infection involves multiple steps, which include initial binding to the cell surface, internalization, replication, and egress. In the initial step of virus infection, these versatile infectious agents can bind several different cellular surface molecules, such as proteins, lipids, and carbohydrates. These molecules may function in mediating attachment (i.e., concentrating virus on the cell surface) or serve as receptors or co-receptors facilitating viral endocytosis, conformational changes, and the initiation of signaling pathways associated with infection [121, 158].

In addition to the protein receptor, which is generally dubbed the 'principal' receptor, the carbohydrate moiety of host-cell membrane proteoglycans, glycosphingolipids, and glycoproteins also serve as viral receptors. For instance, HIV-1, via its glycoprotein subunits gp41 and gp120, attached to cell-surface carbohydrates (i.e., glycosphingolipids, galactosylceramide, and heparan sulfate proteoglycans [HSPGs]) as a means of promoting actual virus binding to cells [8, 51, 171]. Similarly, several human herpesviruses, including HSV [147], KSHV [4], and CMV [93], make their initial contact with cells by binding to cellsurface HSPGs.

Expressing many glycan-binding proteins [46], viruses have been shown to bind host-cell carbohydrate attachment receptors in what is considered a charge-transfer-based lectin-sugar interaction [18]. Lectins are glycoproteins that bind sugar with high specificity [8], and these lectin-glycoconjugate interactions have been shown to involve hydrogen bonding, van der Waals interactions, and hydrophobic binding [8, 159]. Many enveloped and nonenveloped viruses have multiple lectin sites on their multimeric surface proteins; the apparent lectin clustering allows for potentially numerous interactions with cellular surface receptors and notably higher-affinity binding to the oligosaccharide attachment receptors compared to monomeric endogenous lectins [18, 79, 142, 158, 175]. The influenza virus haemagglutinin glycoprotein is one example of a common viral lectin-binding protein that binds to sialyloligosaccharide carbohydrate receptors on epithelial cell surfaces with high affinity [144, 172, 187]. Similarly, sialyloligosaccharide receptors serve as a means of cell attachment for other viruses, including human JC virus, Sendai virus, and sialyloligosaccharide-dependent strains of rotavirus and reovirus [18]. Specifically, C-type lectins are associated with HIV, SARS coronavirus, and measles virus infections. Likewise, S-type, P-type and I-type lectins are important for promoting infection of HIV, HTLV-1,VZV, and porcine reproductive and respiratory syndrome virus (PRRSV) [38, 63, 133].

In general, binding of viruses to carbohydrate moieties on the surface of cells is the key step that induces conformational changes in the viral structure that are critical for interactions with entry-promoting receptors such as integrins. Blocking this step of virus interactions with carbohydrate moieties impairs viral entry via integrins [4]. A list of viruses that utilize carbohydrate moieties to promote virus binding to cells is provided in Table 3.

#### Integrins are targeted specifically to develop therapeutic strategies to treat infections

Integrins are exciting pharmacological targets because (i) they are exposed on the cell surface and are sensitive to pharmacological blockades and (ii) they regulate the interactions of cells and precisely sense their microenvironment. Inhibitors of integrin functions have been successfully tested as drugs to treat several pathological conditions. PSK1404, a nonpeptide antagonist of  $\alpha\nu\beta3$ , inhibited osteoclast-mediated bone resorption in a cancer animal model of bone loss [188]. Volociximab, now known as M200, is a humanized monoclonal antibody that binds specifically to  $\alpha5\beta1$  integrin [14]. In a phase I trial conducted by Ricart et al., volociximab was shown to stabilize Reovirus virus

Table 3 Carbohydrate moieties that interact with viruses

Virus	Viral protein	Carbohydrate moiety	Cell-surface glycoprotein	References
KSHV	Glycoproteins B and K8.1	HS	Proteoglycans	[4, 16]
HSV-1	Glycoproteins B and C	HS	Proteoglycans	[141, 154, 165]
VZV	gB (gpII)	HS	Proteoglycans	[ <del>90</del> ]
CMV	gB and gM (gC-II)	HS	Proteoglycans	[93, 120]
HHV7	gB and gp65	HS	Proteoglycans	[136, 145]
BPV	Hemagglutinin	Sialic acid and O-linked alpha2,3 neuraminic acids	Sialylglycoproteins and glycophorin A (GPA)	[17, 91]
Human norovirus [23]		Histo-blood group antigens (HBGA) and HS	Glycosphingolipids and proteoglycans	[156, 158]
Vaccinia virus		HS	Proteoglycans	[18]
HIV-1	Subunits gp41 and gp120	Galactosylceramide	Glycosphingolipids and galactosylceramide	[51, 171]
Influenza virus	Haemagglutinin and neuraminidase	Sialyloligosaccharide	Glycoproteins or glycosphingolipids	[144, 187]
JC virus		Sialyloligosaccharide	Glycoproteins or glycosphingolipids	[152]
Sendai virus		Sialyloligosaccharide	Glycoproteins or glycosphingolipids	[59, 109]
Rotavirus virus		Sialyloligosaccharide	Glycoproteins or glycosphingolipids	[34]

Sialyloligosaccharide

2675

[29]

disease in patients with advanced solid tumors [128]. Vedolizumab, a humanized monoclonal antibody that specifically recognizes the  $\alpha 4\beta 7$  heterodimer, underwent a phase 3 trial to determine its effectiveness and safety in treating patients with ulcerative colitis [52]. In that study, vedolizumab as both an initial and maintenance therapy for patients with active ulcerative colitis was shown to be effective in achieving a response and remission [52]. Recently, natalizumab, one of the five therapeutic drugs targeting integrins, has been approved for clinic use. This engineered pan- $\alpha$ 4 antibody has been approved for recurrent multiple sclerosis (MS) patients and has been shown to yield promising results for relapsed MS patients by reducing the frequency of relapse, a unique therapeutic result [125]. Its efficacy against Crohn's disease has also been demonstrated [125].

Sigma1 protein

Though integrins have been targeted to treat cancers and other pathological disorders, we have not made a significant breakthrough in targeting integrins to treat virus infections. This does not seem encouraging, especially with many viruses having been shown to utilize integrins to enter cells. This may be due to the fact that (i) viruses utilize multiple receptor molecules to enter the same cell and that (ii) the receptors utilized by the same virus to enter cells *in vitro* and *in vivo* may differ [56]. To overcome this pitfall, we may have to decipher the key elements in the motifs on the virus that interact with integrins and conduct detailed comparative studies outlining the manner by which the virus enters cells under in vitro and in vivo conditions. This will be crucial for gaining comprehensive knowledge of the receptors utilized by viruses to infect cells. Such studies, we hope, will get us one step closer to developing treatment strategies targeting integrins to combat viral infections.

Glycoproteins or glycosphingolipids

#### **Conclusions and recommendations**

Integrins are just not receptors expressed on the surface of cells. They regulate a diverse set of cellular functions are involved in the pathology of autoimmune diseases [73] and viral infections [174]. Viruses utilize different types of integrins, which are classified primarily based on the manner in which they interact with their ligands, ECM proteins. Integrins physically recognize and interact with distinct amino acid sequences contained within the ligands or pathogens. These can be RGD or any other specific (non-RGD) sequences. Whatever the amino acid recognition sequence may be, integrins (RGD and non-RGD) seem to generally aid in virus attachment and entry into cells. The function of integrins is not limited to providing anchoring for the virus. They are also critical for preparing the cells to support a permissive infection via outside-in signaling. Expression of integrins seems to be of relevance in the initial infection as well as in the pathobiology of the virus-induced condition. Recent growth in the field of biomedical sciences has already aided in the development of therapeutics based on integrin interactions to treat various cancers and other pathological conditions. To date, such novel therapeutics to treat virus infections are still only a dream, even though multiple viruses seem to utilize integrins to enter cells. Future studies, we hope, will work toward understanding the roles of integrins in virus infection and associated pathogenesis, as such studies may result in novel treatment regimens aimed at preventing the internalization of viruses. After all, the ideal method of treating infection is to block the entry of a pathogen into cells.

#### References

- Abban CY, Meneses PI (2010) Usage of heparan sulfate, integrins, and FAK in HPV16 infection. Virology 403:1–16
- Adair BD, Yeager M (2002) Three-dimensional model of the human platelet integrin alpha IIbbeta 3 based on electron cryomicroscopy and X-ray crystallography. Proc Natl Acad Sci USA 99:14059–14064
- Aksoy P, Abban CY, Kiyashka E, Qiang W, Meneses PI (2014) HPV16 infection of HaCaTs is dependent on beta4 integrin, and alpha6 integrin processing. Virology 449:45–52
- Akula SM, Wang FZ, Vieira J, Chandran B (2001) Human herpesvirus 8 interaction with target cells involves heparan sulfate. Virology 282:245–255
- Akula SM, Pramod NP, Wang FZ, Chandran B (2002) Integrin alpha3beta1 (CD 49c/29) is a cellular receptor for Kaposi's sarcoma-associated herpesvirus (KSHV/HHV-8) entry into the target cells. Cell 108:407–419
- Akula SM, Naranatt PP, Walia NS, Wang FZ, Fegley B, Chandran B (2003) Kaposi's sarcoma-associated herpesvirus (human herpesvirus 8) infection of human fibroblast cells occurs through endocytosis. J Virol 77:7978–7990
- AlDahlawi S, Eslami A, Hakkinen L, Larjava HS (2006) The alphavbeta6 integrin plays a role in compromised epidermal wound healing. Wound Repair Regen 14:289–297
- Alfsen A, Bomsel M (2002) HIV-1 gp41 envelope residues 650-685 exposed on native virus act as a lectin to bind epithelial cell galactosyl ceramide. J Biol Chem 277:25649–25659
- Arnaout MA, Goodman SL, Xiong JP (2007) Structure and mechanics of integrin-based cell adhesion. Curr Opin Cell Biol 19:495–507
- Arthos J, Cicala C, Martinelli E, Macleod K, Van Ryk D, Wei D, Xiao Z, Veenstra TD, Conrad TP, Lempicki RA, McLaughlin S, Pascuccio M, Gopaul R, McNally J, Cruz CC, Censoplano N, Chung E, Reitano KN, Kottilil S, Goode DJ, Fauci AS (2008) HIV-1 envelope protein binds to and signals through integrin alpha4beta7, the gut mucosal homing receptor for peripheral T cells. Nat Immunol 9:301–309
- Asokan A, Hamra JB, Govindasamy L, Agbandje-McKenna M, Samulski RJ (2006) Adeno-associated virus type 2 contains an integrin alpha5beta1 binding domain essential for viral cell entry. J Virol 80:8961–8969
- 12. Barillari G, Gendelman R, Gallo RC, Ensoli B (1993) The Tat protein of human immunodeficiency virus type 1, a growth

factor for AIDS Kaposi sarcoma and cytokine-activated vascular cells, induces adhesion of the same cell types by using integrin receptors recognizing the RGD amino acid sequence. Proc Natl Acad Sci USA 90:7941–7945

- 13. Barillari G, Sgadari C, Fiorelli V, Samaniego F, Colombini S, Manzari V, Modesti A, Nair BC, Cafaro A, Sturzl M, Ensoli B (1999) The Tat protein of human immunodeficiency virus type-1 promotes vascular cell growth and locomotion by engaging the alpha5beta1 and alphavbeta3 integrins and by mobilizing sequestered basic fibroblast growth factor. Blood 94:663–672
- Barkan D, Chambers AF (2011) beta1-integrin: a potential therapeutic target in the battle against cancer recurrence. Clin Cancer Res 17:7219–7223
- Bergelson JM, Shepley MP, Chan BM, Hemler ME, Finberg RW (1992) Identification of the integrin VLA-2 as a receptor for echovirus 1. Science 255:1718–1720
- Birkmann A, Mahr K, Ensser A, Yaguboglu S, Titgemeyer F, Fleckenstein B, Neipel F (2001) Cell surface heparan sulfate is a receptor for human herpesvirus 8 and interacts with envelope glycoprotein K8.1. J Virol 75:11583–11593
- Blackburn SD, Cline SE, Hemming JP, Johnson FB (2005) Attachment of bovine parvovirus to O-linked alpha 2,3 neuraminic acid on glycophorin A. Arch Virol 150:1477–1484
- Bomsel M, Alfsen A (2003) Entry of viruses through the epithelial barrier: pathogenic trickery. Nat Rev Mol Cell Biol 4:57–68
- Bryan BA, Dyson OF, McCubrey JA, Akula SM (2005) Biology of Kaposi's sarcoma-associated herpesvirus. Front Biosci 10:2882–2891
- 20. Burke RD (1999) Invertebrate integrins: structure, function, and evolution. Int Rev Cytol 191:257–284
- Calderwood DA, Shattil SJ, Ginsberg MH (2000) Integrins and actin filaments: reciprocal regulation of cell adhesion and signaling. J Biol Chem 275:22607–22610
- 22. Campbell SM, Crowe SM, Mak J (2001) Lipid rafts and HIV-1: from viral entry to assembly of progeny virions. J Clin Virol 22:217–227
- 23. Capper D, Weissert S, Balss J, Habel A, Meyer J, Jager D, Ackermann U, Tessmer C, Korshunov A, Zentgraf H, Hartmann C, von Deimling A (2010) Characterization of R132H mutationspecific IDH1 antibody binding in brain tumors. Brain Pathol 20:245–254
- 24. Carter RT (2009) The role of integrins in corneal wound healing. Vet Ophthalmol 12(Suppl 1):2–9
- 25. Caruso M, Belloni L, Sthandier O, Amati P, Garcia MI (2003) Alpha4beta1 integrin acts as a cell receptor for murine polyomavirus at the postattachment level. J Virol 77:3913–3921
- Cavalli V, Corti M, Gruenberg J (2001) Endocytosis and signaling cascades: a close encounter. FEBS Lett 498:190–196
- Chakraborty S, Veettil MV, Chandran B (2012) Kaposi's sarcoma associated herpesvirus entry into target cells. Front Microbiol 3:6
- Chandran B (2010) Early events in Kaposi's sarcoma-associated herpesvirus infection of target cells. J Virol 84:2188–2199
- Chappell JD, Duong JL, Wright BW, Dermody TS (2000) Identification of carbohydrate-binding domains in the attachment proteins of type 1 and type 3 reoviruses. J Virol 74:8472–8479
- Chazal N, Gerlier D (2003) Virus entry, assembly, budding, and membrane rafts. Microbiol Mol Biol Rev 67:226–237 (table of contents)
- 31. Cheshenko N, Trepanier JB, Gonzalez PA, Eugenin EA, Jacobs WR Jr, Herold BC (2014) Herpes simplex virus type 2 glycoprotein H interacts with integrin alphavbeta3 to facilitate viral entry and calcium signaling in human genital tract epithelial cells. J Virol 88:10026–10038

- Chintakuntlawar AV, Zhou X, Rajaiya J, Chodosh J (2010) Viral capsid is a pathogen-associated molecular pattern in adenovirus keratitis. PLoS Pathog 6:e1000841
- 33. Chu JJ, Ng ML (2004) Interaction of West Nile virus with alpha v beta 3 integrin mediates virus entry into cells. J Biol Chem 279:54533–54541
- 34. Ciarlet M, Estes MK (2001) Interactions between rotavirus and gastrointestinal cells. Curr Opin Microbiol 4:435–441
- 35. Cicala C, Martinelli E, McNally JP, Goode DJ, Gopaul R, Hiatt J, Jelicic K, Kottilil S, Macleod K, O'Shea A, Patel N, Van Ryk D, Wei D, Pascuccio M, Yi L, McKinnon L, Izulla P, Kimani J, Kaul R, Fauci AS, Arthos J (2009) The integrin alpha4beta7 forms a complex with cell-surface CD4 and defines a T-cell subset that is highly susceptible to infection by HIV-1. Proc Natl Acad Sci USA 106:20877–20882
- Cicala C, Arthos J, Fauci AS (2011) HIV-1 envelope, integrins and co-receptor use in mucosal transmission of HIV. J Transl Med 9(Suppl 1):S2
- Cox D, Brennan M, Moran N (2010) Integrins as therapeutic targets: lessons and opportunities. Nat Rev Drug Discov 9:804–820
- Crocker PR, Redelinghuys P (2008) Siglecs as positive and negative regulators of the immune system. Biochem Soc Trans 36:1467–1471
- Cudmore S, Reckmann I, Way M (1997) Viral manipulations of the actin cytoskeleton. Trends Microbiol 5:142–148
- Danthi P, Holm GH, Stehle T, Dermody TS (2013) Reovirus receptors, cell entry, and proapoptotic signaling. Adv Exp Med Biol 790:42–71
- 41. Davison E, Kirby I, Whitehouse J, Hart I, Marshall JF, Santis G (2001) Adenovirus type 5 uptake by lung adenocarcinoma cells in culture correlates with Ad5 fibre binding is mediated by alpha(v)beta1 integrin and can be modulated by changes in beta1 integrin function. J Gene Med 3:550–559
- Desgrosellier JS, Cheresh DA (2010) Integrins in cancer: biological implications and therapeutic opportunities. Nat Rev Cancer 10:9–22
- Dohner K, Nagel CH, Sodeik B (2005) Viral stop-and-go along microtubules: taking a ride with dynein and kinesins. Trends Microbiol 13:320–327
- 44. Dyson OF, Oxendine TL, Hamden KE, Ford PW, Akula SM (2008) Differential regulation of the attachment of Kaposi's sarcoma-associated herpesvirus (KSHV)-infected human B cells to extracellular matrix by KSHV-encoded gB and cellular alphaV integrins. Cell Microbiol 10:1546–1558
- 45. Esfandiarei M, Suarez A, Amaral A, Si X, Rahmani M, Dedhar S, McManus BM (2006) Novel role for integrin-linked kinase in modulation of coxsackievirus B3 replication and virus-induced cardiomyocyte injury. Circ Res 99:354–361
- Esko JD, Sharon N (2009) Microbial lectins: hemagglutinins, adhesins, and toxins
- 47. Esmaelizad M, Jelokhani-Niaraki S, Hashemnejad K, Kamalzadeh M, Lotfi M (2011) Molecular characterization of amino acid deletion in VP1 (1D) protein and novel amino acid substitutions in 3D polymerase protein of foot and mouth disease virus subtype A/Iran87. J Vet Sci 12:363–371
- 48. Eto K, Huet C, Tarui T, Kupriyanov S, Liu HZ, Puzon-McLaughlin W, Zhang XP, Sheppard D, Engvall E, Takada Y (2002) Functional classification of ADAMs based on a conserved motif for binding to integrin alpha 9beta 1: implications for sperm-egg binding and other cell interactions. J Biol Chem 277:17804–17810
- Evans DJ, Almond JW (1998) Cell receptors for picornaviruses as determinants of cell tropism and pathogenesis. Trends Microbiol 6:198–202

- Evans R, Patzak I, Svensson L, De Filippo K, Jones K, McDowall A, Hogg N (2009) Integrins in immunity. J Cell Sci 122:215–225
- 51. Fantini J, Hammache D, Delezay O, Yahi N, Andre-Barres C, Rico-Lattes I, Lattes A (1997) Synthetic soluble analogs of galactosylceramide (GalCer) bind to the V3 domain of HIV-1 gp120 and inhibit HIV-1-induced fusion and entry. J Biol Chem 272:7245–7252
- 52. Feagan BG, Rutgeerts P, Sands BE, Hanauer S, Colombel JF, Sandborn WJ, Van Assche G, Axler J, Kim HJ, Danese S, Fox I, Milch C, Sankoh S, Wyant T, Xu J, Parikh A (2013) Vedolizumab as induction and maintenance therapy for ulcerative colitis. N Engl J Med 369:699–710
- Feire AL, Koss H, Compton T (2004) Cellular integrins function as entry receptors for human cytomegalovirus via a highly conserved disintegrin-like domain. Proc Natl Acad Sci USA 101:15470–15475
- Feire AL, Roy RM, Manley K, Compton T (2010) The glycoprotein B disintegrin-like domain binds beta 1 integrin to mediate cytomegalovirus entry. J Virol 84:10026–10037
- 55. Flick MJ, Du X, Witte DP, Jirouskova M, Soloviev DA, Busuttil SJ, Plow EF, Degen JL (2004) Leukocyte engagement of fibrin(ogen) via the integrin receptor alphaMbeta2/Mac-1 is critical for host inflammatory response in vivo. J Clin Invest 113:1596–1606
- 56. Fowler V, Bashiruddin JB, Belsham GJ, Stenfeldt C, Botner A, Knowles NJ, Bankowski B, Parida S, Barnett P (2014) Characteristics of a foot-and-mouth disease virus with a partial VP1 G-H loop deletion in experimentally infected cattle. Vet Microbiol 169:58–66
- 57. Fox G, Parry NR, Barnett PV, McGinn B, Rowlands DJ, Brown F (1989) The cell attachment site on foot-and-mouth disease virus includes the amino acid sequence RGD (arginine-glycine-aspartic acid). J Gen Virol 70(Pt 3):625–637
- 58. Francis SE, Goh KL, Hodivala-Dilke K, Bader BL, Stark M, Davidson D, Hynes RO (2002) Central roles of alpha5beta1 integrin and fibronectin in vascular development in mouse embryos and embryoid bodies. Arterioscler Thromb Vasc Biol 22:927–933
- 59. Fujioka H, Emancipator SN, Aikawa M, Huang DS, Blatnik F, Karban T, DeFife K, Mazanec MB (1998) Immunocytochemical colocalization of specific immunoglobulin A with sendai virus protein in infected polarized epithelium. J Exp Med 188:1223–1229
- Gardner H (2014) Integrin alpha1beta1. Adv Exp Med Biol 819:21–39
- 61. Garrigues HJ, Rubinchikova YE, Dipersio CM, Rose TM (2008) Integrin alphaVbeta3 Binds to the RGD motif of glycoprotein B of Kaposi's sarcoma-associated herpesvirus and functions as an RGD-dependent entry receptor. J Virol 82:1570–1580
- 62. Gavrilovskaya IN, Shepley M, Shaw R, Ginsberg MH, Mackow ER (1998) beta3 Integrins mediate the cellular entry of hantaviruses that cause respiratory failure. Proc Natl Acad Sci USA 95:7074–7079
- Geijtenbeek TB, Gringhuis SI (2009) Signalling through C-type lectin receptors: shaping immune responses. Nat Rev Immunol 9:465–479
- 64. Graham KL, Halasz P, Tan Y, Hewish MJ, Takada Y, Mackow ER, Robinson MK, Coulson BS (2003) Integrin-using rotaviruses bind alpha2beta1 integrin alpha2 I domain via VP4 DGE sequence and recognize alphaXbeta2 and alphaVbeta3 by using VP7 during cell entry. J Virol 77:9969–9978
- 65. Graham KL, Zeng W, Takada Y, Jackson DC, Coulson BS (2004) Effects on rotavirus cell binding and infection of monomeric and polymeric peptides containing alpha2beta1 and alphaxbeta2 integrin ligand sequences. J Virol 78:11786–11797

- 66. Graham KL, Fleming FE, Halasz P, Hewish MJ, Nagesha HS, Holmes IH, Takada Y, Coulson BS (2005) Rotaviruses interact with alpha4beta7 and alpha4beta1 integrins by binding the same integrin domains as natural ligands. J Gen Virol 86:3397–3408
- 67. Graham KL, Takada Y, Coulson BS (2006) Rotavirus spike protein VP5\* binds alpha2beta1 integrin on the cell surface and competes with virus for cell binding and infectivity. J Gen Virol 87:1275–1283
- 68. Granio O, Ashbourne Excoffon KJ, Henning P, Melin P, Norez C, Gonzalez G, Karp PH, Magnusson MK, Habib N, Lindholm L, Becq F, Boulanger P, Zabner J, Hong SS (2010) Adenovirus 5-fiber 35 chimeric vector mediates efficient apical correction of the cystic fibrosis transmembrane conductance regulator defect in cystic fibrosis primary airway epithelia. Hum Gene Ther 21:251–269
- 69. Guerrero CA, Mendez E, Zarate S, Isa P, Lopez S, Arias CF (2000) Integrin alpha(v)beta(3) mediates rotavirus cell entry. Proc Natl Acad Sci USA 97:14644–14649
- Guerrero CA, Zarate S, Corkidi G, Lopez S, Arias CF (2000) Biochemical characterization of rotavirus receptors in MA104 cells. J Virol 74:9362–9371
- Guo W, Giancotti FG (2004) Integrin signalling during tumour progression. Nat Rev Mol Cell Biol 5:816–826
- 72. Hait SH, Darc M, Machado ES, Soares EA, Sprinz E, Soares MA (2014) Conservation of the alpha4beta7 lymphocyte homing receptor in HIV-infected patients with distinct transmission routes and disease progression profiles. AIDS Res Hum Retroviruses 30:493–497
- Halpert G, Sredni B (2014) The effect of the novel tellurium compound AS101 on autoimmune diseases. Autoimmun Rev 13:1230–1235
- 74. Haque M, Davis DA, Wang V, Widmer I, Yarchoan R (2003) Kaposi's sarcoma-associated herpesvirus (human herpesvirus 8) contains hypoxia response elements: relevance to lytic induction by hypoxia. J Virol 77:6761–6768
- Harley D, Sleigh A, Ritchie S (2001) Ross River virus transmission, infection, and disease: a cross-disciplinary review. Clin Microbiol Rev 14:909–932 (table of contents)
- Harokopakis E, Hajishengallis G (2005) Integrin activation by bacterial fimbriae through a pathway involving CD14, Toll-like receptor 2, and phosphatidylinositol-3-kinase. Eur J Immunol 35:1201–1210
- 77. Heikkila O, Susi P, Tevaluoto T, Harma H, Marjomaki V, Hyypia T, Kiljunen S (2010) Internalization of coxsackievirus A9 is mediated by {beta}2-microglobulin, dynamin, and Arf6 but not by caveolin-1 or clathrin. J Virol 84:3666–3681
- Hemler ME, Lobb RR (1995) The leukocyte beta 1 integrins. Curr Opin Hematol 2:61–67
- Hidari KI, Abe T, Suzuki T (2013) Carbohydrate-related inhibitors of dengue virus entry. Viruses 5:605–618
- Hoffmann C, Ohlsen K, Hauck CR (2011) Integrin-mediated uptake of fibronectin-binding bacteria. Eur J Cell Biol 90:891–896
- Holly SP, Larson MK, Parise LV (2000) Multiple roles of integrins in cell motility. Exp Cell Res 261:69–74
- 82. Hsiao JC, Chung CS, Chang W (1999) Vaccinia virus envelope D8L protein binds to cell surface chondroitin sulfate and mediates the adsorption of intracellular mature virions to cells. J Virol 73:8750–8761
- Huang S, Kamata T, Takada Y, Ruggeri ZM, Nemerow GR (1996) Adenovirus interaction with distinct integrins mediates separate events in cell entry and gene delivery to hematopoietic cells. J Virol 70:4502–4508
- Huttenlocher A, Horwitz AR (2011) Integrins in cell migration. Cold Spring Harb Perspect Biol 3:a005074

- Hynes RO (1987) Integrins: a family of cell surface receptors. Cell 48:549–554
- Hynes RO (2002) Integrins: bidirectional, allosteric signaling machines. Cell 110:673–687
- Izmailyan R, Hsao JC, Chung CS, Chen CH, Hsu PW, Liao CL, Chang W (2012) Integrin beta1 mediates vaccinia virus entry through activation of PI3K/Akt signaling. J Virol 86:6677–6687
- 88. Jackson T, Blakemore W, Newman JW, Knowles NJ, Mould AP, Humphries MJ, King AM (2000) Foot-and-mouth disease virus is a ligand for the high-affinity binding conformation of integrin alpha5beta1: influence of the leucine residue within the RGDL motif on selectivity of integrin binding. J Gen Virol 81:1383–1391
- Jackson T, Clark S, Berryman S, Burman A, Cambier S, Mu D, Nishimura S, King AM (2004) Integrin alphavbeta8 functions as a receptor for foot-and-mouth disease virus: role of the betachain cytodomain in integrin-mediated infection. J Virol 78:4533–4540
- 90. Jacquet A, Haumont M, Chellun D, Massaer M, Tufaro F, Bollen A, Jacobs P (1998) The varicella zoster virus glycoprotein B (gB) plays a role in virus binding to cell surface heparan sulfate proteoglycans. Virus Res 53:197–207
- Johnson FB, Fenn LB, Owens TJ, Faucheux LJ, Blackburn SD (2004) Attachment of bovine parvovirus to sialic acids on bovine cell membranes. J Gen Virol 85:2199–2207
- 92. Jokinen J, White DJ, Salmela M, Huhtala M, Kapyla J, Sipila K, Puranen JS, Nissinen L, Kankaanpaa P, Marjomaki V, Hyypia T, Johnson MS, Heino J (2010) Molecular mechanism of alpha2beta1 integrin interaction with human echovirus 1. EMBO J 29:196–208
- Kari B, Gehrz R (1992) A human cytomegalovirus glycoprotein complex designated gC-II is a major heparin-binding component of the envelope. J Virol 66:1761–1764
- 94. Kim SH, Turnbull J, Guimond S (2011) Extracellular matrix and cell signalling: the dynamic cooperation of integrin, proteoglycan and growth factor receptor. J Endocrinol 209:139–151
- 95. Krishnan HH, Sharma-Walia N, Streblow DN, Naranatt PP, Chandran B (2006) Focal adhesion kinase is critical for entry of Kaposi's sarcoma-associated herpesvirus into target cells. J Virol 80:1167–1180
- 96. La Linn M, Eble JA, Lubken C, Slade RW, Heino J, Davies J, Suhrbier A (2005) An arthritogenic alphavirus uses the alpha1beta1 integrin collagen receptor. Virology 336:229–239
- Lawrence P, LaRocco M, Baxt B, Rieder E (2013) Examination of soluble integrin resistant mutants of foot-and-mouth disease virus. Virol J 10:2
- Lee JO, Rieu P, Arnaout MA, Liddington R (1995) Crystal structure of the A domain from the alpha subunit of integrin CR3 (CD11b/CD18). Cell 80:631–638
- Leistner CM, Gruen-Bernhard S, Glebe D (2008) Role of glycosaminoglycans for binding and infection of hepatitis B virus. Cell Microbiol 10:122–133
- 100. Leu ST, Jacques SA, Wingerd KL, Hikita ST, Tolhurst EC, Pring JL, Wiswell D, Kinney L, Goodman NL, Jackson DY, Clegg DO (2004) Integrin alpha4beta1 function is required for cell survival in developing retina. Dev Biol 276:416–430
- 101. Ley K, Laudanna C, Cybulsky MI, Nourshargh S (2007) Getting to the site of inflammation: the leukocyte adhesion cascade updated. Nat Rev Immunol 7:678–689
- 102. Li E, Stupack D, Bokoch GM, Nemerow GR (1998) Adenovirus endocytosis requires actin cytoskeleton reorganization mediated by Rho family GTPases. J Virol 72:8806–8812
- 103. Li E, Stupack D, Klemke R, Cheresh DA, Nemerow GR (1998) Adenovirus endocytosis via alpha(v) integrins requires phosphoinositide-3-OH kinase. J Virol 72:2055–2061

- 104. Li E, Brown SL, Stupack DG, Puente XS, Cheresh DA, Nemerow GR (2001) Integrin alpha(v)beta1 is an adenovirus coreceptor. J Virol 75:5405–5409
- 105. Li P, Lu Z, Bao H, Li D, King DP, Sun P, Bai X, Cao W, Gubbins S, Chen Y, Xie B, Guo J, Yin H, Liu Z (2011) In-vitro and in-vivo phenotype of type Asia 1 foot-and-mouth disease viruses utilizing two non-RGD receptor recognition sites. BMC Microbiol 11:154
- 106. Liu Q, Muruve DA (2003) Molecular basis of the inflammatory response to adenovirus vectors. Gene Ther 10:935–940
- 107. Lopez S, Arias CF (2004) Multistep entry of rotavirus into cells: a Versaillesque dance. Trends Microbiol 12:271–278
- Luo BH, Carman CV, Springer TA (2007) Structural basis of integrin regulation and signaling. Annu Rev Immunol 25:619–647
- 109. Markwell MA, Paulson JC (1980) Sendai virus utilizes specific sialyloligosaccharides as host cell receptor determinants. Proc Natl Acad Sci USA 77:5693–5697
- 110. Mathew SJ, Haubert D, Kronke M, Leptin M (2009) Looking beyond death: a morphogenetic role for the TNF signalling pathway. J Cell Sci 122:1939–1946
- 111. McPherson PS, Kay BK, Hussain NK (2001) Signaling on the endocytic pathway. Traffic 2:375–384
- 112. Miller N, Hutt-Fletcher LM (1992) Epstein–Barr virus enters B cells and epithelial cells by different routes. J Virol 66:3409–3414
- 113. Mitra SK, Schlaepfer DD (2006) Integrin-regulated FAK-Src signaling in normal and cancer cells. Curr Opin Cell Biol 18:516–523
- 114. Monini P, Cafaro A, Srivastava IK, Moretti S, Sharma VA, Andreini C, Chiozzini C, Ferrantelli F, Cossut MR, Tripiciano A, Nappi F, Longo O, Bellino S, Picconi O, Fanales-Belasio E, Borsetti A, Toschi E, Schiavoni I, Bacigalupo I, Kan E, Sernicola L, Maggiorella MT, Montin K, Porcu M, Leone P, Collacchi B, Palladino C, Ridolfi B, Falchi M, Macchia I, Ulmer JB, Butto S, Sgadari C, Magnani M, Federico MP, Titti F, Banci L, Dallocchio F, Rappuoli R, Ensoli F, Barnett SW, Garaci E, Ensoli B (2012) HIV-1 tat promotes integrin-mediated HIV transmission to dendritic cells by binding Env spikes and competes neutralization by anti-HIV antibodies. PLoS One 7:e48781
- 115. Mylonas AD, Brown AM, Carthew TL, McGrath B, Purdie DM, Pandeya N, Vecchio PC, Collins LG, Gardner ID, de Looze FJ, Reymond EJ, Suhrbier A (2002) Natural history of Ross River virus-induced epidemic polyarthritis. Med J Aust 177:356–360
- 116. Nakano T, Inoue I, Shinozaki R, Matsui M, Akatsuka T, Takahashi S, Tanaka K, Akita M, Seo M, Hokari S, Katayama S, Komoda T (2009) A possible role of lysophospholipids produced by calcium-independent phospholipase A(2) in membrane-raft budding and fission. Biochim Biophys Acta 1788:2222–2228
- 117. Naranatt PP, Akula SM, Zien CA, Krishnan HH, Chandran B (2003) Kaposi's sarcoma-associated herpesvirus induces the phosphatidylinositol 3-kinase-PKC-zeta-MEK-ERK signaling pathway in target cells early during infection: implications for infectivity. J Virol 77:1524–1539
- 118. Naranatt PP, Krishnan HH, Smith MS, Chandran B (2005) Kaposi's sarcoma-associated herpesvirus modulates microtubule dynamics via RhoA-GTP-diaphanous 2 signaling and utilizes the dynein motors to deliver its DNA to the nucleus. J Virol 79:1191–1206
- 119. Nemerow GR, Stewart PL (2001) Antibody neutralization epitopes and integrin binding sites on nonenveloped viruses. Virology 288:189–191
- Neyts J, Snoeck R, Schols D, Balzarini J, Esko JD, Van Schepdael A, De Clercq E (1992) Sulfated polymers inhibit the

interaction of human cytomegalovirus with cell surface heparan sulfate. Virology 189:48–58

- 121. Olofsson S, Bergstrom T (2005) Glycoconjugate glycans as viral receptors. Ann Med 37:154–172
- 122. Parry C, Bell S, Minson T, Browne H (2005) Herpes simplex virus type 1 glycoprotein H binds to alphavbeta3 integrins. J Gen Virol 86:7–10
- 123. Philpott NJ, Nociari M, Elkon KB, Falck-Pedersen E (2004) Adenovirus-induced maturation of dendritic cells through a PI3 kinase-mediated TNF-alpha induction pathway. Proc Natl Acad Sci USA 101:6200–6205
- 124. Plow EF, Haas TA, Zhang L, Loftus J, Smith JW (2000) Ligand binding to integrins. J Biol Chem 275:21785–21788
- 125. Pucci E, Giuliani G, Solari A, Simi S, Minozzi S, Di Pietrantonj C, Galea I (2011) Natalizumab for relapsing remitting multiple sclerosis. Cochrane Database Syst Rev 10:CD007621
- 126. Raghu H, Sharma-Walia N, Veettil MV, Sadagopan S, Caballero A, Sivakumar R, Varga L, Bottero V, Chandran B (2007) Lipid rafts of primary endothelial cells are essential for Kaposi's sarcoma-associated herpesvirus/human herpesvirus 8-induced phosphatidylinositol 3-kinase and RhoA-GTPases critical for microtubule dynamics and nuclear delivery of viral DNA but dispensable for binding and entry. J Virol 81:7941–7959
- 127. Raymond T, Gorbunova E, Gavrilovskaya IN, Mackow ER (2005) Pathogenic hantaviruses bind plexin-semaphorin-integrin domains present at the apex of inactive, bent alphavbeta3 integrin conformers. Proc Natl Acad Sci USA 102:1163–1168
- 128. Ricart AD, Tolcher AW, Liu G, Holen K, Schwartz G, Albertini M, Weiss G, Yazji S, Ng C, Wilding G (2008) Volociximab, a chimeric monoclonal antibody that specifically binds alpha5-beta1 integrin: a phase I, pharmacokinetic, and biological correlative study. Clin Cancer Res 14:7924–7929
- 129. Roivainen M, Hyypia T, Piirainen L, Kalkkinen N, Stanway G, Hovi T (1991) RGD-dependent entry of coxsackievirus A9 into host cells and its bypass after cleavage of VP1 protein by intestinal proteases. J Virol 65:4735–4740
- 130. Roivainen M, Piirainen L, Hovi T, Virtanen I, Riikonen T, Heino J, Hyypia T (1994) Entry of coxsackievirus A9 into host cells: specific interactions with alpha v beta 3 integrin, the vitronectin receptor. Virology 203:357–365
- Roivainen M, Piirainen L, Hovi T (1996) Efficient RGD-independent entry process of coxsackievirus A9. Arch Virol 141:1909–1919
- Ruoslahti E, Pierschbacher MD (1986) Arg-Gly-Asp: a versatile cell recognition signal. Cell 44:517–518
- 133. Sato S, St-Pierre C, Bhaumik P, Nieminen J (2009) Galectins in innate immunity: dual functions of host soluble beta-galactoside-binding lectins as damage-associated molecular patterns (DAMPs) and as receptors for pathogen-associated molecular patterns (PAMPs). Immunol Rev 230:172–187
- 134. Schmidt K, Keller M, Bader BL, Korytar T, Finke S, Ziegler U, Groschup MH (2013) Integrins modulate the infection efficiency of West Nile virus into cells. J Gen Virol 94:1723–1733
- 135. Schornberg KL, Shoemaker CJ, Dube D, Abshire MY, Delos SE, Bouton AH, White JM (2009) Alpha5beta1-integrin controls ebolavirus entry by regulating endosomal cathepsins. Proc Natl Acad Sci USA 106:8003–8008
- 136. Secchiero P, Sun D, De Vico AL, Crowley RW, Reitz MS Jr, Zauli G, Lusso P, Gallo RC (1997) Role of the extracellular domain of human herpesvirus 7 glycoprotein B in virus binding to cell surface heparan sulfate proteoglycans. J Virol 71:4571–4580
- 137. Seo NS, Zeng CQ, Hyser JM, Utama B, Crawford SE, Kim KJ, Hook M, Estes MK (2008) Integrins alpha1beta1 and alpha2beta1 are receptors for the rotavirus enterotoxin. Proc Natl Acad Sci USA 105:8811–8818

- 138. Shakeel S, Seitsonen JJ, Kajander T, Laurinmaki P, Hyypia T, Susi P, Butcher SJ (2013) Structural and functional analysis of coxsackievirus A9 integrin alphavbeta6 binding and uncoating. J Virol 87:3943–3951
- 139. Sharma-Walia N, Naranatt PP, Krishnan HH, Zeng L, Chandran B (2004) Kaposi's sarcoma-associated herpesvirus/human herpesvirus 8 envelope glycoprotein gB induces the integrin-dependent focal adhesion kinase-Src-phosphatidylinositol 3-kinase-rho GTPase signal pathways and cytoskeletal rearrangements. J Virol 78:4207–4223
- 140. Shayakhmetov DM, Eberly AM, Li ZY, Lieber A (2005) Deletion of penton RGD motifs affects the efficiency of both the internalization and the endosome escape of viral particles containing adenovirus serotype 5 or 35 fiber knobs. J Virol 79:1053–1061
- 141. Shukla D, Liu J, Blaiklock P, Shworak NW, Bai X, Esko JD, Cohen GH, Eisenberg RJ, Rosenberg RD, Spear PG (1999) A novel role for 3-O-sulfated heparan sulfate in herpes simplex virus 1 entry. Cell 99:13–22
- 142. Shukla D, Spear PG (2001) Herpesviruses and heparan sulfate: an intimate relationship in aid of viral entry. J Clin Invest 108:503–510
- 143. Silva LA, Khomandiak S, Ashbrook AW, Weller R, Heise MT, Morrison TE, Dermody TS (2014) A single-amino-acid polymorphism in Chikungunya virus E2 glycoprotein influences glycosaminoglycan utilization. J Virol 88:2385–2397
- 144. Skehel JJ, Wiley DC (2000) Receptor binding and membrane fusion in virus entry: the influenza hemagglutinin. Annu Rev Biochem 69:531–569
- 145. Skrincosky D, Hocknell P, Whetter L, Secchiero P, Chandran B, Dewhurst S (2000) Identification and analysis of a novel heparin-binding glycoprotein encoded by human herpesvirus 7. J Virol 74:4530–4540
- 146. Smith JP, Morris-Downes M, Brennan FR, Wallace GJ, Amor S (2000) A role for alpha4-integrin in the pathology following Semliki Forest virus infection. J Neuroimmunol 106:60–68
- 147. Spear PG (2004) Herpes simplex virus: receptors and ligands for cell entry. Cell Microbiol 6:401–410
- 148. Srichai MB, Zent R (2010) Integrin structure and function. In: Zent R, Pozzi A (eds) Cell extracellular matrix interaction in cancer. Sprimger Science + Business Media, New York, pp 19–41
- 149. Stergiou L, Bauer M, Mair W, Bausch-Fluck D, Drayman N, Wollscheid B, Oppenheim A, Pelkmans L (2013) Integrin-mediated signaling induced by simian virus 40 leads to transient uncoupling of cortical actin and the plasma membrane. PLoS One 8:e55799
- Stewart PL, Nemerow GR (2007) Cell integrins: commonly used receptors for diverse viral pathogens. Trends Microbiol 15:500–507
- 151. Summerford C, Bartlett JS, Samulski RJ (1999) AlphaVbeta5 integrin: a co-receptor for adeno-associated virus type 2 infection. Nat Med 5:78–82
- 152. Suzuki S, Sawa H, Komagome R, Orba Y, Yamada M, Okada Y, Ishida Y, Nishihara H, Tanaka S, Nagashima K (2001) Broad distribution of the JC virus receptor contrasts with a marked cellular restriction of virus replication. Virology 286:100–112
- 153. Suzuki T, Suzuki Y (2006) Virus infection and lipid rafts. Biol Pharm Bull 29:1538–1541
- 154. Svennerholm B, Jeansson S, Vahlne A, Lycke E (1991) Involvement of glycoprotein C (gC) in adsorption of herpes simplex virus type 1 (HSV-1) to the cell. Arch Virol 120:273–279
- 155. Takada Y, Ye X, Simon S (2007) The integrins. Genome Biol 8:215

- 156. Tamura M, Natori K, Kobayashi M, Miyamura T, Takeda N (2004) Genogroup II noroviruses efficiently bind to heparan sulfate proteoglycan associated with the cellular membrane. J Virol 78:3817–3826
- 157. Tapon N, Hall A (1997) Rho, Rac and Cdc42 GTPases regulate the organization of the actin cytoskeleton. Curr Opin Cell Biol 9:86–92
- Taube S, Jiang M, Wobus CE (2010) Glycosphingolipids as receptors for non-enveloped viruses. Viruses 2:1011–1049
- 159. Tkac J, Bertok T, Nahalka J, Gemeiner P (2014) Perspectives in glycomics and lectin engineering. Methods Mol Biol 1200:421–445
- 160. Toschi E, Bacigalupo I, Strippoli R, Chiozzini C, Cereseto A, Falchi M, Nappi F, Sgadari C, Barillari G, Mainiero F, Ensoli B (2006) HIV-1 Tat regulates endothelial cell cycle progression via activation of the Ras/ERK MAPK signaling pathway. Mol Biol Cell 17:1985–1994
- 161. Tran T, Barlow B, O'Rear L, Jarvis B, Li Z, Dickeson K, Dupont W, Zutter M (2011) Loss of the alpha2beta1 integrin alters human papilloma virus-induced squamous carcinoma progression in vivo and in vitro. PLoS One 6:e26858
- 162. Triantafilou K, Triantafilou M, Takada Y, Fernandez N (2000) Human parechovirus 1 utilizes integrins alphavbeta3 and alphavbeta1 as receptors. J Virol 74:5856–5862
- 163. Triantafilou K, Takada Y, Triantafilou M (2001) Mechanisms of integrin-mediated virus attachment and internalization process. Crit Rev Immunol 21:311–322
- 164. Triantafilou K, Triantafilou M (2003) Lipid raft microdomains: key sites for Coxsackievirus A9 infectious cycle. Virology 317:128–135
- 165. Trybala E, Liljeqvist JA, Svennerholm B, Bergstrom T (2000) Herpes simplex virus types 1 and 2 differ in their interaction with heparan sulfate. J Virol 74:9106–9114
- 166. Tugizov SM, Berline JW, Palefsky JM (2003) Epstein–Barr virus infection of polarized tongue and nasopharyngeal epithelial cells. Nat Med 9:307–314
- 167. Uyama T, Ishida M, Izumikawa T, Trybala E, Tufaro F, Bergstrom T, Sugahara K, Kitagawa H (2006) Chondroitin 4-Osulfotransferase-1 regulates E disaccharide expression of chondroitin sulfate required for herpes simplex virus infectivity. J Biol Chem 281:38668–38674
- 168. Veesler D, Cupelli K, Burger M, Graber P, Stehle T, Johnson JE (2014) Single-particle EM reveals plasticity of interactions between the adenovirus penton base and integrin alphaVbeta3. Proc Natl Acad Sci USA 111:8815–8819
- 169. Veettil MV, Sadagopan S, Sharma-Walia N, Wang FZ, Raghu H, Varga L, Chandran B (2008) Kaposi's sarcoma-associated herpesvirus forms a multimolecular complex of integrins (al-phaVbeta5, alphaVbeta3, and alpha3beta1) and CD98-xCT during infection of human dermal microvascular endothelial cells, and CD98-xCT is essential for the postentry stage of infection. J Virol 82:12126–12144
- 170. Vellon L, Menendez JA, Lupu R (2005) AlphaVbeta3 integrin regulates heregulin (HRG)-induced cell proliferation and survival in breast cancer. Oncogene 24:3759–3773
- 171. Villard R, Hammache D, Delapierre G, Fotiadu F, Buono G, Fantini J (2002) Asymmetric synthesis of water-soluble analogues of galactosylceramide, an HIV-1 receptor: new tools to study virus-glycolipid interactions. Chembiochem 3:517–525
- 172. Viswanathan K, Chandrasekaran A, Srinivasan A, Raman R, Sasisekharan V, Sasisekharan R (2010) Glycans as receptors for influenza pathogenesis. Glycoconj J 27:561–570
- 173. Vogel BE, Lee SJ, Hildebrand A, Craig W, Pierschbacher MD, Wong-Staal F, Ruoslahti E (1993) A novel integrin specificity exemplified by binding of the alpha v beta 5 integrin to the basic

domain of the HIV Tat protein and vitronectin. J Cell Biol 121:461-468

- 174. Walker LR, Hussein HA, Akula SM (2014) Disintegrin-like domain of glycoprotein B regulates Kaposi's sarcoma-associated herpesvirus infection of cells. J Gen Virol 95:1770–1782
- 175. Weis WI, Drickamer K (1996) Structural basis of lectin-carbohydrate recognition. Annu Rev Biochem 65:441–473
- Whitton JL, Cornell CT, Feuer R (2005) Host and virus determinants of picornavirus pathogenesis and tropism. Nat Rev Microbiol 3:765–776
- 177. Wickham TJ, Mathias P, Cheresh DA, Nemerow GR (1993) Integrins alpha v beta 3 and alpha v beta 5 promote adenovirus internalization but not virus attachment. Cell 73:309–319
- 178. Wickham TJ, Filardo EJ, Cheresh DA, Nemerow GR (1994) Integrin alpha v beta 5 selectively promotes adenovirus mediated cell membrane permeabilization. J Cell Biol 127:257–264
- 179. Wilde A, Beattie EC, Lem L, Riethof DA, Liu SH, Mobley WC, Soriano P, Brodsky FM (1999) EGF receptor signaling stimulates SRC kinase phosphorylation of clathrin, influencing clathrin redistribution and EGF uptake. Cell 96:677–687
- 180. Williams CH, Kajander T, Hypia T, Jackson T, Sheppard D, Stanway G (2004) Integrin alpha v beta 6 is an RGD-dependent receptor for coxsackievirus A9. J Virol 78:6967–6973
- Wu Z, Asokan A, Samulski RJ (2006) Adeno-associated virus serotypes: vector toolkit for human gene therapy. Mol Ther 14:316–327
- 182. Xiao T, Takagi J, Coller BS, Wang JH, Springer TA (2004) Structural basis for allostery in integrins and binding to fibrinogen-mimetic therapeutics. Nature 432:59–67

- 183. Xiong JP, Stehle T, Diefenbach B, Zhang R, Dunker R, Scott DL, Joachimiak A, Goodman SL, Arnaout MA (2001) Crystal structure of the extracellular segment of integrin alpha Vbeta3. Science 294:339–345
- 184. Yoon CS, Kim KD, Park SN, Cheong SW (2001) alpha(6) Integrin is the main receptor of human papillomavirus type 16 VLP. Biochem Biophys Res Commun 283:668–673
- 185. Zaidel-Bar R, Itzkovitz S, Ma'ayan A, Iyengar R, Geiger B (2007) Functional atlas of the integrin adhesome. Nat Cell Biol 9:858–867
- 186. Zarate S, Espinosa R, Romero P, Guerrero CA, Arias CF, Lopez S (2000) Integrin alpha2beta1 mediates the cell attachment of the rotavirus neuraminidase-resistant variant nar3. Virology 278:50–54
- 187. Zhang J, Pekosz A, Lamb RA (2000) Influenza virus assembly and lipid raft microdomains: a role for the cytoplasmic tails of the spike glycoproteins. J Virol 74:4634–4644
- 188. Zhao Y, Bachelier R, Treilleux I, Pujuguet P, Peyruchaud O, Baron R, Clement-Lacroix P, Clezardin P (2007) Tumor alphavbeta3 integrin is a therapeutic target for breast cancer bone metastases. Cancer Res 67:5821–5830
- 189. Zhu J, Motejlek K, Wang D, Zang K, Schmidt A, Reichardt LF (2002) beta8 integrins are required for vascular morphogenesis in mouse embryos. Development 129:2891–2903