

# Bifurcation for a free boundary problem modeling the growth of a tumor with a necrotic core

Wenrui Hao<sup>a,1,\*</sup>, Jonathan D. Hauenstein<sup>b,2</sup>, Bei Hu<sup>a</sup>, Yuan Liu<sup>a</sup>, Andrew J. Somnese<sup>a,1</sup>, Yong-Tao Zhang<sup>a</sup>

<sup>a</sup>*Department of Applied and Computational Mathematics and Statistics, University of Notre Dame, Notre Dame, IN 46556*

<sup>b</sup>*Department of Mathematics, Mailstop 3368, Texas A&M University, College Station, TX 77843*

---

## Abstract

We consider a free boundary problem for a system of partial differential equations, which arises in a model of tumor growth with a necrotic core. For any positive numbers  $\rho < R$ , there exists a radially symmetric stationary solution with tumor boundary  $r = R$  and necrotic core boundary  $r = \rho$ . The system depends on a positive parameter  $\mu$ , which describes the tumor aggressiveness. There also exists a sequence of values  $\mu_2 < \mu_3 < \dots$  for which branches of symmetry-breaking stationary solutions bifurcate from the radially symmetric solution branch.

*Keywords:* Bifurcation, free boundary problem, tumor model, necrotic core

---

## Introduction

Tumor growth models are challenging both theoretically and numerically. They are free boundary problems where the changing shape of the tumor is of prime importance. Spherical solutions are often straightforward to compute, but values of the controlling parameter, called the tumor-aggressiveness factor, where bifurcations occur are nontrivial to compute. Moreover, analytically finding nonspherical solutions on a branch far from a spherical solution is intractable. Another difficult and important question is to determine the linear stability of the solution branches.

---

\*Corresponding author

*Email addresses:* [whao@nd.edu](mailto:whao@nd.edu) (Wenrui Hao), [jhauenst@math.tamu.edu](mailto:jhauenst@math.tamu.edu) (Jonathan D. Hauenstein), [b1hu@nd.edu](mailto:b1hu@nd.edu) (Bei Hu), [yliu7@nd.edu](mailto:yliu7@nd.edu) (Yuan Liu), [sommese@nd.edu](mailto:sommese@nd.edu) (Andrew J. Somnese), [y Zhang@nd.edu](mailto:y Zhang@nd.edu) (Yong-Tao Zhang)

*URL:* [www.math.tamu.edu/~jhauenst](http://www.math.tamu.edu/~jhauenst) (Jonathan D. Hauenstein), [www.nd.edu/~b1hu](http://www.nd.edu/~b1hu) (Bei Hu), [www.nd.edu/~sommese](http://www.nd.edu/~sommese) (Andrew J. Somnese), [www.nd.edu/~y Zhang](http://www.nd.edu/~y Zhang) (Yong-Tao Zhang)

<sup>1</sup>This author was supported by the Duncan Chair of the University of Notre Dame and NSF grant DMS-0712910.

<sup>2</sup>This author was supported by Texas A&M University and NSF grant DMS-0915211.

In this article, we analyze a highly nontrivial tumor growth model. The paper is organized into five sections. In §1, we discuss tumor models and explicitly set up the system of differential equations controlling tumor growth for the necrotic core model. In §2, we derive an analytical formula for the radially symmetric stationary solutions. In §3, we establish the existence of bifurcation branches from radially symmetric stationary, explicitly provide the formula describing where the bifurcation branches occur, and study the bifurcation diagram. In §4, we derive the linearized system and determine linear stability in a small neighborhood of the bifurcation at  $\mu_2$ .

## 1. The model

Mathematical models of solid tumor growth, which consider the tumor tissue as a density of proliferating cells, have been developed and studied in many papers; see [1; 2; 3; 4; 5; 7; 9; 10; 11; 13; 14; 16; 17; 18; 19; 20] and the references provided therein. The case of radially symmetric tumors have been discussed extensively.

Tumors grown in vitro have a nearly spherical shape, but tumors grown in vivo may develop different kinds of protrusions. It is therefore interesting to explore the existence of non-spherical solutions of tumor models.

If dead cells are not removed in an efficient manner from the tumor, they accumulate inside to form a necrotic core [7] and [9]. A necrotic tumor growth model consists of a core of necrotic cells and shell adjacent to this necrotic core of proliferating cells. In particular, let  $\Omega(t)$  denote the tumor domain at time  $t$ , and  $D(t) \subset \Omega(t)$  be the necrotic core within the tumor domain.

Let  $p$  be the pressure within the tumor resulting from proliferation of the tumor cells. The density of the cells,  $c$ , depends on the concentration of nutrients,  $\sigma$ , and, assuming that this dependence is linear, we simply identify  $c$  with  $\sigma$ . We also assume the proliferation rate,  $S$ , depends linearly upon  $\sigma$  in the living tumor region. That is,

$$S = \mu(\sigma - \tilde{\sigma}) \quad \text{in } \Omega(t) \setminus D(t),$$

where  $\tilde{\sigma} > 0$  is a threshold concentration and  $\mu$  is a positive parameter measuring the aggressiveness of the tumor. This equation represents the balance of the growth of the tumor by cell division and the contraction of the tumor by necrosis. The linear approximation  $\mu(\sigma - \tilde{\sigma})$  used here is the result of first order Taylor expansion for the fully nonlinear model.

We assume that there is no proliferation in the necrotic core, i.e,  $S = 0$  in  $D(t)$ . Combining these two equations, we have

$$S = \mu(\sigma - \tilde{\sigma})\chi_{\{\Omega(t) \setminus D(t)\}}(x) \quad \text{in } \Omega(t), \quad (1)$$

where  $\chi_{\{\Omega(t) \setminus D(t)\}}(x)$  is the indicator function of  $\Omega(t) \setminus D(t)$ .

If we assume that necrotic cells do not consume nutrients and the consumption rate of nutrients by the living tumor cells is proportional to the concentration of the nutrients, then after normalization,  $\sigma$  satisfies

$$\sigma_t - \Delta\sigma = -\sigma\chi_{\{\Omega(t) \setminus D(t)\}}(x) \quad \text{in } \Omega(t) \quad \text{and} \quad \sigma = 1 \quad \text{on } \partial\Omega(t). \quad (2)$$

Additionally, if we assume that the density of cells in the necrotic core remains constant, we have  $\sigma = \underline{\sigma}$  in  $D(t)$ .

Most tumor models assume that the tissue has the structure of a porous medium so that Darcy's law holds. In particular,  $\vec{v} = -\nabla p$  where  $\vec{v}$  is the velocity of the cells and  $p$  is the pressure. By conservation of mass,  $\text{div } \vec{v} = S = \mu(\sigma - \tilde{\sigma})\chi_{\{\Omega(t) \setminus D(t)\}}(x)$  and thus  $\Delta p = -\mu(\sigma - \tilde{\sigma})\chi_{\{\Omega(t) \setminus D(t)\}}(x)$  in  $\Omega(t)$ . As in [6], the cell-to-cell adhesiveness condition at the tumor boundary is represented by  $p = \kappa$  on  $\partial\Omega(t)$ . To simplify notation, let  $\chi(x, t) = \chi_{\{\Omega(t) \setminus D(t)\}}(x)$ . The necrotic core system is

$$\begin{cases} \sigma_t - \Delta\sigma = -\sigma\chi(x, t) & \text{and} & -\Delta p = \mu(\sigma - \tilde{\sigma})\chi(x, t) & \text{in } \Omega(t), \\ \sigma = \underline{\sigma} & & & \text{in } D(t), \\ \sigma = 1; \quad p = \kappa; & \text{and} & \frac{\partial p}{\partial n} = -V_n & \text{on } \partial\Omega(t), \end{cases} \quad (3)$$

where  $n$  denotes the exterior normal vector. Additionally, it is reasonable to assume  $\underline{\sigma} < \tilde{\sigma} < 1$ .

An important question is the existence and stability of steady-state solutions. Biologically, a stable steady-state solution means that the tumor can remain inside a patient for a long time without causing much trouble. An unstable tumor is a tumor that is very likely to spread beyond the region it occupies. The steady-state necrotic core tumor model is

$$\Delta\sigma = \sigma\chi(x) \quad \text{in } \Omega, \quad (4)$$

$$-\Delta p = \mu(\sigma - \tilde{\sigma})\chi(x) \quad \text{in } \Omega, \quad (5)$$

$$\sigma = \underline{\sigma} \quad \text{in } D, \quad (6)$$

$$\sigma = 1 \quad \text{on } \partial\Omega, \quad (7)$$

$$p = \kappa \quad \text{on } \partial\Omega, \quad (8)$$

$$\frac{\partial p}{\partial n} = 0 \quad \text{on } \partial\Omega. \quad (9)$$

## 2. Radially symmetric stationary solutions

We recall some basic formulas for the modified bessel functions  $K_n(x)$  and  $I_n(x)$ . These functions form a fundamental solution set of  $x^2 y'' + xy' - (x^2 + n^2)y = 0$ . In particular,

$$I_{n+1}(x) = I_{n-1}(x) - \frac{2n}{x}I_n(x), \quad K_{n+1}(x) = K_{n-1}(x) + \frac{2n}{x}K_n(x), \quad n \geq 1, \quad (1)$$

$$I'_n(x) = \frac{1}{2}[I_{n-1}(x) + I_{n+1}(x)], \quad K'_n(x) = -\frac{1}{2}[K_{n-1}(x) + K_{n+1}(x)], \quad n \geq 1, \quad (2)$$

$$I'_n(x) = I_{n-1}(x) - \frac{n}{x}I_n(x), \quad K'_n(x) = -K_{n-1}(x) - \frac{n}{x}K_n(x), \quad n \geq 1, \quad (3)$$

$$I'_n(x) = \frac{n}{x}I_n(x) + I_{n+1}(x), \quad K'_n(x) = \frac{n}{x}K_n(x) - K_{n+1}(x), \quad n \geq 0. \quad (4)$$

Additionally,

$$I'_n(x) > 0 \quad \text{and} \quad K'_n(x) < 0. \quad (5)$$

In this paper, we will only consider the case of two space dimensions. The 3-dimensional case can be considered in a similar manner, except that the formula are more complicated.

Solving (4) with the boundary conditions (6) and (7), we obtain

$$\sigma_s(r) = \frac{(-\underline{\sigma}K_0(R) + K_0(\rho))I_0(r) - (I_0(\rho) - \underline{\sigma}I_0(R))K_0(r)}{-I_0(\rho)K_0(R) + I_0(R)K_0(\rho)} \quad \rho < r < R; \quad (6)$$

Solving (5), we obtain

$$p_s(r) = \frac{1}{4}\mu\tilde{\sigma}r^2 + c_1(R, \rho) \ln r + c_2(R, \rho) - \mu\sigma_s(r) \quad \rho < r < R; \quad (7)$$

where  $c_1(R, \rho)$  and  $c_2(R, \rho)$  are constants to be determined. The boundary conditions (8) and (9) imply that

$$p_s(R) = \frac{1}{R}, \quad \frac{\partial p_s}{\partial r}(R) = 0, \quad (8)$$

and (5) implies that  $p_s$  is a constant with the necrotic core. Since the first order derivative of  $p_s$  and  $\sigma_s$  are continuous across the boundary of necrotic core,

$$\frac{\partial p_s}{\partial r}(\rho) = 0 \quad \text{and} \quad \frac{\partial \sigma_s}{\partial r}(\rho) = 0. \quad (9)$$

Substituting (7) into (8) and (9), we have

$$\frac{1}{4}\mu\tilde{\sigma}R^2 + c_1(R, \rho) \ln R + c_2(R, \rho) - \mu = \frac{1}{R}, \quad (10)$$

$$\frac{1}{2}\mu\tilde{\sigma}R + c_1(R, \rho)\frac{1}{R} - \mu\frac{\partial \sigma_s}{\partial r}(R) = 0, \quad (11)$$

$$\frac{1}{2}\mu\tilde{\sigma}\rho + c_1(R, \rho)\frac{1}{\rho} - \mu\frac{\partial \sigma_s}{\partial r}(\rho) = 0. \quad (12)$$

From (12) and (9),  $c_1(R, \rho) = -\frac{1}{2}\mu\tilde{\sigma}\rho^2$ .

Substituting  $c_1$  into (10), we obtain  $c_2(R, \rho) = \frac{1}{R} + \mu - \frac{1}{4}\mu\tilde{\sigma}R^2 + \frac{1}{2}\mu\tilde{\sigma}\rho^2 \ln R$ . Thus

$$p_s(r) = \frac{1}{4}\mu\tilde{\sigma}(r^2 - R^2) + \frac{1}{R} + \frac{1}{2}\mu\tilde{\sigma}\rho^2 \ln \frac{R}{r} - \mu(\sigma_s(r) - 1), \quad \rho < r < R; \quad (13)$$

It remains to verify that (9) and (11) hold. We shall use (9) and (11) to determine  $\rho, R, \tilde{\sigma}$ . By (4),  $I_0'(r) = I_1(r)$  and  $K_0'(r) = -K_1(r)$ . Using (6), the equation (9) is equivalent to

$$(-\underline{\sigma}K_0(R) + K_0(\rho))I_1(\rho) + (I_0(\rho) - \underline{\sigma}I_0(R))K_1(\rho) = 0, \quad (14)$$

and (11) can be rewritten as

$$\frac{1}{2}\tilde{\sigma}R - \frac{1}{2}\tilde{\sigma}\frac{\rho^2}{R} - \frac{\partial \sigma_s}{\partial r}(R) = 0, \quad (15)$$

where, by (6),

$$\frac{\partial \sigma_s}{\partial r}(R) = \frac{(-\underline{\sigma}K_0(R) + K_0(\rho))I_1(R) + (I_0(\rho) - \underline{\sigma}I_0(R))K_1(R)}{-I_0(\rho)K_0(R) + I_0(R)K_0(\rho)}. \quad (16)$$

For this system, we need to compute  $\rho$  and  $R$  for given  $\mu, \underline{\sigma}$  and  $\tilde{\sigma}$ . For technical reasons, we will instead solve for  $\tilde{\sigma}$  for a given  $\rho$ . It will produce radially symmetric solution for any given  $\rho$ .

**Lemma 2.1.** *For any given  $\mu > 0$  and  $\rho > 0$ , there exists a unique  $\tilde{\sigma}$  such that a stationary solution  $(\sigma_s(r), p_s(r), \rho, R)$  is given by (6) and (13) where*

- (a)  $R$  is uniquely determined by (14),  $R > \rho$ , and
- (b)  $\tilde{\sigma}$  is uniquely determined over the interval  $(\underline{\sigma}, 1)$  by (15).

*Proof.* To establish (a), we denote

$$f(s) = (-\underline{\sigma}K_0(s) + K_0(\rho))I_1(\rho) + (I_0(\rho) - \underline{\sigma}I_0(s))K_1(\rho),$$

then, recalling that  $\bar{\sigma} < 1$ ,

$$f(\rho) = (1 - \underline{\sigma})(K_0(\rho)I_1(\rho) + I_0(\rho)K_1(\rho)) > 0$$

and  $f(s) \rightarrow -\infty$  as  $s \rightarrow \infty$ . Therefore,  $f(s)$  must have at least one root, say  $s = R$ , in the interval  $(\rho, \infty)$ .

To show that  $f(s)$  is monotone, consider  $g(s) = \frac{K_1(s)}{I_1(s)}$ . We know

$$\frac{dg}{ds}(s) = -\frac{I_1(s)K_0(s) + K_1(s)I_0(s)}{I_1(s)^2} < 0.$$

Therefore,  $g$  is decreasing function. In particular, if  $s > \rho$ , then  $g(s) < g(\rho)$ . This implies  $K_1(s)I_1(\rho) < K_1(\rho)I_1(s)$  yielding

$$\frac{df}{ds}(s) = -\underline{\sigma}(K_1(s)I_1(\rho) - I_1(s)K_1(\rho)) > 0.$$

This implies that  $R$  is the unique solution to (14) on  $(\rho, \infty)$ .

To establish (b), we note that it is easy to solve for  $\tilde{\sigma}$  using (15). Thus, we only need to establish that  $\underline{\sigma} < \tilde{\sigma} < 1$ .

In fact, from equations (14) and (15), we can compute  $\underline{\sigma}$  and  $\tilde{\sigma}$ :

$$\underline{\sigma} = \frac{K_0(\rho)I_1(\rho) + I_0(\rho)K_1(\rho)}{K_0(R)I_1(\rho) + I_0(R)K_1(\rho)}, \quad (17)$$

$$\tilde{\sigma} = \frac{2R \frac{\partial \sigma_s}{\partial r}(R)}{R^2 - \rho^2}. \quad (18)$$

We next establish that  $\tilde{\sigma} > \underline{\sigma}$ . After simplification using (16), recalling also that the denominator in (16) satisfies (by (5))

$$-I_0(\rho)K_0(R) + I_0(R)K_0(\rho) > K_0(R)(-I_0(\rho) + I_0(R)) > 0,$$

we find that this inequality is equivalent to

$$\begin{aligned} & 2R(K_0(\rho)I_1(R) + I_0(\rho)K_1(R))(K_0(R)I_1(\rho) + I_0(R)K_1(\rho)) \\ & > \{(R^2 - \rho^2)[I_0(R)K_0(\rho) - I_0(\rho)K_0(R)] + 2R[K_0(R)I_1(R) + I_0(R)K_1(R)]\} \cdot \\ & \quad (K_0(\rho)I_1(\rho) + I_0(\rho)K_1(\rho)). \end{aligned}$$

We let  $f_1(s)$  be the function consisting of the difference of the left-hand side and the right-hand side of the above inequality after replacing  $\rho$  by  $s$ , i.e., after simplification,

$$\begin{aligned} f_1(s) &= I_0(R)I_0(s)I_1(s) \left( \frac{K_0(s)}{I_0(s)} - \frac{K_0(R)}{I_0(R)} \right) \cdot \\ & \quad \left\{ 2RI_1(R) \left( \frac{K_1(s)}{I_1(s)} - \frac{K_1(R)}{I_1(R)} \right) - (R^2 - s^2)I_0(s) \left( \frac{K_0(s)}{I_0(s)} + \frac{K_1(s)}{I_1(s)} \right) \right\}. \end{aligned} \quad (19)$$

We want to establish  $f_1(s) > 0$  for  $s < R$ . Clearly, for  $s < R$ , we know  $I_0(R)I_0(s)I_1(s) \left( \frac{K_0(s)}{I_0(s)} - \frac{K_0(R)}{I_0(R)} \right) > 0$ . Thus, it suffices to verify that, for  $s < R$ ,

$$h_1(s) := 2RI_1(R) \left( \frac{K_1(s)}{I_1(s)} - \frac{K_1(R)}{I_1(R)} \right) - (R^2 - s^2)I_0(s) \left( \frac{K_0(s)}{I_0(s)} + \frac{K_1(s)}{I_1(s)} \right) > 0. \quad (20)$$

It is clear that  $h_1(R) = 0$  and

$$h_1'(s) = \frac{\left( 2(sI_1(s) - RI_1(R)) + (R^2 - s^2)I_0(s) \right) \left( I_1(s)K_0(s) + I_0(s)K_1(s) \right)}{I_1(s)^2}.$$

Let  $i_1(s) = 2(sI_1(s) - RI_1(R)) + (R^2 - s^2)I_0(s)$ , then

$$i_1'(s) = sI_0(s) + (R^2 - s^2)I_1(s) > 0 \quad \text{and} \quad i_1(R) = 0.$$

Thus,  $i_1(s) > 0$  for any  $s \in (0, R)$ . Utilizing (20), we find  $h_1'(s) < 0$  for  $s < R$ . Combining this with  $h_1(R) = 0$ , we conclude that  $h_1(s) > 0$  for  $s < R$ .

Finally, we will prove that  $\tilde{\sigma} < 1$ , which is equivalent to

$$\begin{aligned} & 2R(K_0(\rho)I_1(R) + I_0(\rho)K_1(R))(K_0(R)I_1(\rho) + I_0(R)K_1(\rho)) \\ & < (R^2 - \rho^2) \left( I_0(R)K_0(\rho) - I_0(\rho)K_0(R) \right) \left( K_0(R)I_1(\rho) + I_0(R)K_1(\rho) \right) \\ & \quad + 2R \left( K_0(R)I_1(R) + I_0(R)K_1(R) \right) \left( K_0(\rho)I_1(\rho) + I_0(\rho)K_1(\rho) \right). \end{aligned} \quad (21)$$

Similarly, we let  $f_2(s)$  be the function consisting of the difference of the left-hand side and the right-hand side of (22) after replacing  $\rho$  by  $s$ . After simplification, we find

$$\begin{aligned} f_2(s) &= I_0(R)I_0(s)I_1(s) \left( \frac{K_0(s)}{I_0(s)} - \frac{K_0(R)}{I_0(R)} \right) \cdot \\ & \quad \left\{ 2RI_1(R) \left( \frac{K_1(s)}{I_1(s)} - \frac{K_1(R)}{I_1(R)} \right) - (R^2 - s^2)I_0(R) \left( \frac{K_0(R)}{I_0(R)} + \frac{K_1(s)}{I_1(s)} \right) \right\}. \end{aligned}$$

We want to show that  $f_2(s) < 0$ , for  $s < R$ . It suffices to establish

$$h_2(s) := 2RI_1(R) \left( \frac{K_1(s)}{I_1(s)} - \frac{K_1(R)}{I_1(R)} \right) - (R^2 - s^2)I_0(R) \left( \frac{K_0(R)}{I_0(R)} + \frac{K_1(s)}{I_1(s)} \right) < 0. \quad (22)$$

It is clear that  $h_2(R) = 0$  and

$$h_2'(s) = \frac{\left( 2(sI_1(s)m(s) - RI_1(R)) + (R^2 - s^2)I_0(R) \right) \left( I_1(s)K_0(s) + I_0(s)K_1(s) \right)}{I_1(s)^2},$$

where  $m(s) = \frac{I_1(s)K_0(R) + K_1(s)I_0(R)}{I_1(s)K_0(s) + K_1(s)I_0(s)}$ . We will show  $m(s) > 1$  for  $s < R$ , so that

$$h_2'(s) > \frac{i_2(s) \left( I_1(s)K_0(s) + I_0(s)K_1(s) \right)}{I_1(s)^2}, \quad (23)$$

where  $i_2(s) = 2(sI_1(s) - RI_1(R)) + (R^2 - s^2)I_0(R)$ . Clearly,  $i_2'(s) = 2sI_0(s) - 2sI_0(R) < 0$  and  $i_2(R) = 0$ . Thus,  $i_2(s) > 0$  for  $s < R$ . Since  $h_2(R) = 0$ , we have established (22).

To finish the proof, we need to show  $m(s) > 1$  for  $s < R$ . Clearly,

$$\begin{aligned} \frac{d}{dR} & \left( (I_1(s)K_0(R) + K_1(s)I_0(R)) - (I_1(s)K_0(s) + K_1(s)I_0(s)) \right) \\ & = I_1(s)I_1(R) \left( -\frac{K_1(R)}{I_1(R)} + \frac{K_1(s)}{I_1(s)} \right) > 0, \end{aligned}$$

Since  $(I_1(s)K_0(R) + K_1(s)I_0(R))|_{R=s} - (I_1(s)K_0(s) + K_1(s)I_0(s))|_{R=s} = 0$ , for  $R > s$ , we know

$$\left( (I_1(s)K_0(R) + K_1(s)I_0(R)) - (I_1(s)K_0(s) + K_1(s)I_0(s)) \right) > 0.$$

Hence  $m(s) > 1$  for  $s < R$ .  $\square$

### 3. Bifurcation from radially symmetric stationary solution

We now turn our attention to compute nonradially symmetric stationary solutions of the form

$$\sigma = \sigma_s + \epsilon\sigma_1 + O(\epsilon^2), \quad (1)$$

$$p = p_s + \epsilon p_1 + O(\epsilon^2), \quad (2)$$

$$\partial\Omega_\epsilon : \quad r = R + \epsilon S(\theta) + O(\epsilon^2), \quad (3)$$

$$\partial D_\epsilon : \quad r = \rho + \epsilon\rho_1 S(\theta) + O(\epsilon^2), \quad (4)$$

where  $\sigma_1$ ,  $p_1$ , and  $\rho_1$  are constants to be determined.

### 3.1. Formal expansion

In order to compute the Frechét derivative in the direction  $S(\theta)$ , we first formally compute  $\sigma_1$ ,  $\rho_1$ , and  $p_1$ .

**Computation of  $\sigma_1$ :** Clearly,

$$\epsilon\sigma_1|_{\partial B_R} = \epsilon\sigma_1|_{\partial\Omega_\epsilon} + O(\epsilon^2) = 1 - \sigma_s(R + \epsilon S) + O(\epsilon^2) = -\epsilon\sigma'_s(R)S(\theta) + O(\epsilon^2).$$

Similarly  $\epsilon\sigma_1|_{\partial D_\rho} = -\epsilon\sigma'_s(\rho)\rho_1 S(\theta) + O(\epsilon^2) = O(\epsilon^2)$ . Thus, after dropping higher terms,

$$\begin{cases} -\Delta\sigma_1 + \sigma_1 = 0 & \text{in } B_R \setminus D_\rho, \\ \sigma_1 = -\sigma'_s(R)S & \text{on } \partial B_R, \\ \sigma_1 = 0 & \text{on } \partial D_\rho. \end{cases}$$

It follows that, for  $S(\theta) = \cos(l\theta)$ ,  $l = 2, 3, 4, 5, \dots$ , by separation of variables

$$\sigma_1(r) = -\sigma'_s(R) \cos(l\theta) Q_l(r; \rho, R), \quad (5)$$

where

$$Q_l(r; \rho, R) = \frac{K_l(\rho)I_l(r) - I_l(\rho)K_l(r)}{K_l(\rho)I_l(R) - K_l(R)I_l(\rho)} \quad (6)$$

satisfies

$$r^2 Q_l'' + rQ_l' - (r^2 + l^2)Q_l = 0, \quad Q_l(\rho; \rho, R) = 0, \quad Q_l(R; \rho, R) = 1, \quad (7)$$

where  $Q_l'$  and  $Q_l''$  are the first and second derivatives of  $Q_l$  with respect to  $r$ , respectively.

**Computation of  $\rho_1$ :** To compute  $\rho_1$ , we note that

$$\frac{\partial(\sigma_s + \epsilon\sigma_1)}{\partial n} \Big|_{\partial D_\epsilon} = O(\epsilon^2), \quad (8)$$

that is,  $\epsilon \left( \frac{\partial^2 \sigma_s(\rho)}{\partial r^2} \rho_1 S + \frac{\partial \sigma_1}{\partial r}(\rho) \right) = O(\epsilon^2)$ . After dropping higher terms, we obtain

$$\frac{\partial \sigma_1}{\partial r}(\rho) = -\frac{\partial^2 \sigma_s(\rho)}{\partial r^2} \rho_1 S(\theta), \quad (9)$$

which implies that

$$\rho_1 = \rho_1(l; \rho, R) = \frac{\sigma'_s(R)}{\sigma''_s(\rho)} \cdot Q'_l(\rho; \rho, R) = \frac{\sigma'_s(R)}{\sigma_0} \cdot Q'_l(\rho; \rho, R). \quad (10)$$

**Computation of  $p_1$ :** Similarly, for  $p_1$  (see [15])

$$\kappa = \frac{1}{R} - \frac{\epsilon}{R^2} (S + S'') + O(\epsilon^2). \quad (11)$$

After dropping the higher order terms, we have

$$\begin{cases} -\Delta p_1 = \mu\sigma_1 & \text{in } B_R \setminus D_\rho, \\ -\Delta p_1 = 0 & \text{in } D_\rho. \end{cases}$$



As for boundary conditions, when  $S(\theta) = \cos(l\theta)$ ,

$$-\frac{1}{R^2}(S + S'') = \left(\frac{1}{R^2} + \frac{l^2}{R^2}\right) \cos(l\theta), \quad (12)$$

so that, after dropping the higher order terms,  $p_1 = \left(-\frac{1}{R^2} + \frac{l^2}{R^2}\right) \cos(l\theta)$  on  $\partial B_R$ .

We next formally derive two boundary conditions on  $\partial D_\rho$ . Clearly,

$$-\Delta p_s - \epsilon \Delta p_1 = \mu(\sigma_s - \tilde{\sigma})\chi_{\Omega_\epsilon \setminus D_\epsilon} + \epsilon \mu \sigma_1 \chi_{\Omega_\epsilon \setminus D_\epsilon} + O(\epsilon^2). \quad (13)$$

Since  $-\Delta p_s = \mu(\sigma_s - \tilde{\sigma})\chi_{B_R \setminus D_\rho}$ , the function  $p_1$  satisfies

$$-\Delta p_1 = \mu \sigma_1 \chi_{\Omega_\epsilon \setminus D_\epsilon} + \frac{\mu}{\epsilon}(\sigma_s - \tilde{\sigma})f_\epsilon, \quad (14)$$

where  $f_\epsilon$  is given by  $f_\epsilon = \chi_{\Omega_\epsilon \setminus D_\epsilon} - \chi_{B_R \setminus D_\rho}$ . Clearly,  $\mu \sigma_1 \chi_{\Omega_\epsilon \setminus D_\epsilon} \rightarrow \mu \sigma_1 \chi_{B_R \setminus D_\rho}$  as  $\epsilon \rightarrow 0$ . It is also clear that  $\frac{1}{\epsilon} f_\epsilon \rho_1 S(\theta)$  converges to a  $\delta$ -function on  $\partial D_\rho$ , thus  $p_1$  is continuous on  $\partial D_\rho$ ,  $p_1(\rho^+) = p_1(\rho^-)$ , where we use  $\rho^+$  to denote the limit from the tumor side and  $\rho^-$  to denote the limit from necrotic core side.

To derive a second condition across the free boundary  $\{r = \rho\}$ , consider

$$\begin{cases} -\Delta q_1^\epsilon &= \mu \sigma_1 \chi_{\Omega_\epsilon \setminus D_\epsilon} & \text{in } \Omega_\epsilon, \\ q_1^\epsilon &= p_1 & \text{on } \partial \Omega_\epsilon, \end{cases}$$

yielding

$$\begin{cases} -\Delta(p_1 - q_1^\epsilon) &= \frac{\mu}{\epsilon}(\sigma_s - \tilde{\sigma})f_\epsilon & \text{in } \Omega_\epsilon, \\ q_1^\epsilon - p_1 &= 0 & \text{on } \partial \Omega_\epsilon. \end{cases}$$

For any test function  $\zeta$  with compact support in  $B_R$ , then

$$-\iint_{\Omega_\epsilon} \zeta \Delta(p_1 - q_1^\epsilon) = \iint_{\Omega_\epsilon} \frac{\mu}{\epsilon}(\sigma_s - \tilde{\sigma})f_\epsilon \zeta,$$

that is,

$$\iint_{\Omega_\epsilon} \nabla \zeta \nabla(p_1 - q_1^\epsilon) = \int_{\partial D_\epsilon} \mu(\sigma_s - \tilde{\sigma})\rho_1 S(\theta) \zeta + O(\epsilon).$$

In the limit as  $\epsilon \rightarrow 0$ ,

$$\iint_{B_R} \nabla \zeta \nabla(p_1 - q_1) = \int_{\partial D_\rho} \mu(\sigma_s - \tilde{\sigma})\rho_1 S(\theta) \zeta.$$

Note that in the limit  $-\Delta(p_1 - q_1) \equiv 0$  in  $D_\rho \cup (B_R \setminus \overline{D_\rho})$  so that integration by parts yields

$$0 = \iint_{B_R} \nabla \zeta \nabla(p_1 - q_1) - \int_{\partial(B_R \setminus D_\rho)} \zeta \frac{\partial(p_1 - q_1)}{\partial n} - \int_{\partial D_\rho} \zeta \frac{\partial(p_1 - q_1)}{\partial n},$$

where  $\vec{n}$  on  $\partial D_\rho$  points into the tumor region while  $\vec{n}$  on  $\partial(B_R \setminus D_\rho)$  points into the necrotic core. Thus, in the limit,

$$\int_{\partial(B_R \setminus D_\rho)} \zeta \frac{\partial(p_1 - q_1)}{\partial n} + \int_{\partial D_\rho} \zeta \frac{\partial(p_1 - q_1)}{\partial n} = \int_{\partial D_\rho} \mu(\sigma_s - \tilde{\sigma})\rho_1 S(\theta) \zeta. \quad (15)$$

Since  $\nabla q_1$  is continuous across the necrotic core,

$$\frac{\partial p_1}{\partial n}(\rho+) + \frac{\partial p_1}{\partial n}(\rho-) = \mu(\underline{\sigma} - \tilde{\sigma})\rho_1 S(\theta). \quad (16)$$

or, recalling the definition of  $\vec{n}$ ,  $-\frac{\partial p_1}{\partial r}(\rho+) + \frac{\partial p_1}{\partial r}(\rho-) = \mu(\underline{\sigma} - \tilde{\sigma})\rho_1 S(\theta)$ .

Solving  $p_1$  in the regions  $\{r < \rho\}$  and  $\{\rho < r < R\}$  respectively, we obtain

$$\begin{cases} p_1 = c_4 r^l \cos(l\theta) & 0 < r < \rho, \\ p_1 + \mu\sigma_1 = \left(-\frac{1}{R^2} - \mu\sigma'_s(R) + \frac{l^2}{R^2}\right) \frac{r^l}{R^l} \cos(l\theta) + c_3 \left(r^{-l} - \frac{r^l}{R^{2l}}\right) \cos(l\theta) & \rho < r < R, \end{cases}$$

where the constants  $c_3$  and  $c_4$  are determined by matching all the boundary conditions

$$\left(-\frac{1}{R^2} - \mu\sigma'_s(R) + \frac{l^2}{R^2}\right) \frac{\rho^l}{R^l} + c_3 \left(\rho^{-l} - \frac{\rho^l}{R^{2l}}\right) = c_4 \rho^l, \quad (17)$$

and

$$\begin{aligned} \left(-\frac{1}{R^2} - \mu\sigma'_s(R) + \frac{l^2}{R^2}\right) \frac{l\rho^{l-1}}{R^l} - c_3 \left(l\rho^{-l-1} + \frac{l\rho^{l-1}}{R^{2l}}\right) + \mu\sigma''_s(\rho)\rho_1 \\ = c_4 l\rho^{l-1} - \rho_1 \mu(\underline{\sigma} - \tilde{\sigma}). \end{aligned} \quad (18)$$

The combination  $l \times (17) - \rho \times (18)$  gives

$$c_3 2l\rho^{-l} - \mu\underline{\sigma}\rho_1\rho = -\rho_1 \mu(\underline{\sigma} - \tilde{\sigma})\rho \quad (19)$$

yielding  $c_3 = \frac{\rho_1 \rho^{l+1} \mu \tilde{\sigma}}{2l}$ . We can now solve for  $c_4$  using (17).

For each  $S(\theta)$  and  $\mu$ , define

$$F(S, \mu) = \frac{\partial p}{\partial n} \Big|_{\partial\Omega_\epsilon}. \quad (20)$$

Then,  $S$  induces a stationary solution if and only if  $F(S, \mu) = 0$ . Clearly

$$\frac{\partial p}{\partial n} \Big|_{r=R+\epsilon S} = \epsilon \left( \frac{\partial^2 p_s(R)}{\partial r^2} S + \frac{\partial p_1}{\partial r} \right) + O(\epsilon^2). \quad (21)$$

Thus, formally, the Frechét derivative in the direction  $\cos(l\theta)$  is given by

$$\left[ \frac{\partial F}{\partial S}(0, \mu) \right] \cos(l\theta) = \frac{\partial^2 p_s(R)}{\partial r^2} \cos(l\theta) + \frac{\partial p_1}{\partial r}. \quad (22)$$

We claim that the condition for bifurcation is

$$\left[ \frac{\partial F}{\partial S}(0, \mu) \right] \cos(l\theta) \equiv 0, \quad (23)$$

and this will determine  $\mu = \mu_l$ . We now proceed to rigorously derive this. From (13), we have

$$\frac{\partial^2 p_s(R)}{\partial r^2} = \frac{1}{2}\mu\tilde{\sigma} + \frac{1}{2}\mu\tilde{\sigma}\frac{\rho^2}{R^2} - \mu\sigma_s''(R). \quad (24)$$

Using the formula for  $p_1$ ,

$$\frac{\partial p_1}{\partial r}(R) + \mu \frac{\partial \sigma_1}{\partial r}(R) = \left( -\frac{1}{R^2} - \mu\sigma_s'(R) + \frac{l^2}{R^2} \right) \frac{l}{R} \cos(l\theta) - c_3 \frac{2l}{R^{l+1}} \cos(l\theta). \quad (25)$$

Using (8), we know  $\frac{\partial \sigma_1}{\partial r}(R) = -\sigma_s'(R) \cos(l\theta) Q_l'(R; \rho, R)$ . Substituting this into (25) and using (22) and (23), we find

$$\begin{aligned} \frac{1}{2}\mu\tilde{\sigma} \left( 1 + \frac{\rho^2}{R^2} \right) - \mu\sigma_s''(R) + \left( -\frac{1}{R^2} - \mu\sigma_s'(R) + \frac{l^2}{R^2} \right) \frac{l}{R} - c_3 \frac{2l}{R^{l+1}} \\ + \mu\sigma_s'(R) Q_l'(R; \rho, R) = 0. \end{aligned}$$

Substituting  $c_3$  into the above equation, we have

$$\begin{aligned} \frac{1}{2}\mu\tilde{\sigma} \left( 1 + \frac{\rho^2}{R^2} \right) - \mu\sigma_s''(R) + \left( -\frac{1}{R^2} - \mu\sigma_s'(R) + \frac{l^2}{R^2} \right) \frac{l}{R} - \frac{\rho_1 \rho^{l+1}}{R^{l+1}} \mu\tilde{\sigma} \\ + \mu\sigma_s'(R) Q_l'(R; \rho, R) = 0. \end{aligned}$$

Thus,  $A_l \cdot \mu_l + B_l = 0$  where

$$A_l = \frac{1}{2}\tilde{\sigma} \left( 1 + \frac{\rho^2}{R^2} \right) - \sigma_s''(R) - \frac{l}{R} \sigma_s'(R) - \frac{\rho_1 \rho^{l+1}}{R^{l+1}} \tilde{\sigma} + \sigma_s'(R) Q_l'(R; \rho, R), \quad (26)$$

$$B_l = \left( -\frac{1}{R^2} + \frac{l^2}{R^2} \right) \frac{l}{R}. \quad (27)$$

Clearly, if  $A_l \neq 0$ , then  $\mu_l = -\frac{B_l}{A_l}$ ,

### 3.2. Bifurcation

We shall apply the following Crandall-Rabinowitz theorem.

**Theorem 3.1** (see [8, Theorem 1.7]). *Let  $X, Y$  be real Banach spaces and  $F(x, \mu)$  a  $C^p$  map,  $p \geq 3$ , of a neighborhood  $(0, \mu_0)$  in  $X \times \mathbb{R}$  into  $Y$ . Suppose*

- (i).  $F(0, \mu) = 0$  for all  $\mu$  in a neighborhood of  $\mu_0$ ,
- (ii).  $\text{Ker } F_x(0, \mu_0)$  is one dimensional space, spanned by  $x_0$ ,
- (iii).  $\text{Im } F_x(0, \mu_0) = Y_1$  has codimension 1,
- (iv).  $F_{\mu x}(0, \mu_0)x_0 \notin Y_1$ .

Then  $(0, \mu_0)$  is a bifurcation point of the equation  $F(x, \mu) = 0$  in the following sense: In a neighborhood of  $(0, \mu_0)$  the set of solutions of  $F(x, \mu) = 0$  consists of two  $C^{p-2}$  smooth curves  $\Gamma_1$  and  $\Gamma_2$  which intersect only at the point  $(0, \mu_0)$ ;  $\Gamma_1$  is the curve  $(0, \mu)$  and  $\Gamma_2$  can be parameterized as follows:

$$\Gamma_2 : (x(\varepsilon), \mu(\varepsilon)), \quad |\varepsilon| \text{ small}, \quad (x(0), \mu(0)) = (0, \mu_0), \quad x'(0) = x_0.$$

As in [12] we introduce the Banach spaces: for  $k \geq 2$ ,

$$X_0^{k+\alpha} = \left\{ S \in C^{k+\alpha}(-\infty, \infty), S \text{ is } 2\pi\text{-periodic} \right\}, \quad (28)$$

$$X_1^{k+\alpha} = \text{closure of the linear space spanned by} \\ \{\cos(j\theta), j = 0, 1, 2, 3, \dots\} \text{ in } X^{k+\alpha}, \quad (29)$$

$$X_2^{k+\alpha} = \text{closure of the linear space spanned by} \\ \{\cos(j\theta), j = 0, 2, 4, 6, \dots\} \text{ in } X^{k+\alpha}. \quad (30)$$

We shall take  $X = X_2^{k+\alpha}$  and  $Y = X_2^{k-1+\alpha}$  and define  $F$  by (20), for any  $S \in X_2^{k+\alpha}$ . Then  $F : X_2^{k+\alpha} \rightarrow X_2^{k-1+\alpha}$ . It is also clear that  $F : X_j^{k+\alpha} \rightarrow X_j^{k-1+\alpha}$ ,  $j = 0$  and  $j = 1$ .

**Lemma 3.2.** *If  $j \in \{0, 1\}$  and  $F : X_j^{k+\alpha} \rightarrow X_j^{k-1+\alpha}$ , then*

$$[F_S(0, \mu)] \cos(l\theta) \equiv 0$$

*if and only if  $l = 1$ , or  $l \geq 2$  and  $\mu = \mu_j$ ,  $j \geq 2$ .*

*Proof.* If  $l = 1$ , clearly  $B_1 = 0$  in (27). To verify  $A_1 = 0$ , we evaluate the various terms in (26). By plugging (19) into (16), and after a lengthy simplification, we find that

$$\sigma'_s(R) = \frac{K_1(\rho)I_1(R) - K_1(R)I_1(\rho)}{K_0(R)I_1(\rho) + I_0(R)K_1(\rho)}.$$

Since by (5),  $K'_l(r) < 0$  and  $I'_l(r) > 0$ , we derive  $\sigma'_s(R) > 0$ . Moreover, using (3), we obtain

$$Q'_1(r; \rho, R) = \frac{K_1(\rho)I_0(r) + I_1(\rho)K_0(r)}{K_1(\rho)I_1(R) - K_1(R)I_1(\rho)} - \frac{1}{r} \frac{K_1(\rho)I_1(r) - I_1(\rho)K_1(r)}{K_1(\rho)I_1(R) - K_1(R)I_1(\rho)},$$

so that  $Q'_1(R; \rho, R) = \frac{K_0(R)I_1(\rho) + I_0(R)K_1(\rho)}{K_1(\rho)I_1(R) - K_1(R)I_1(\rho)} - \frac{1}{R}$  and  $Q'_1(\rho; \rho, R) = \frac{K_1(\rho)I_0(\rho) + I_1(\rho)K_0(\rho)}{K_1(\rho)I_1(R) - K_1(R)I_1(\rho)}$ . Then, substituting  $\sigma_s(R), \underline{\sigma}$  into (10), we derive  $\rho_1(1; \rho, R) = 1$ . Moreover, by the equation for  $\sigma_s$  and the boundary condition,  $\sigma''_s(R) + \frac{1}{R}\sigma'_s(R) = \Delta\sigma_s(R) = \sigma_s(R) = 1$ . It follows that  $A_1 = \sigma'_s(R) \left( \frac{1}{R} + Q'_1(R; \rho, R) \right) - 1 = 0$ .

The case for  $l \geq 2$  is a corollary of Theorem 3.3 and  $A_l \neq 0$  for  $l \geq 2$ .  $\square$

We remark that  $l = 1$  corresponds to the translation of the original radially symmetric solution.

From this lemma it is clear that  $\cos \theta \in \ker[F_S(0, \mu)]$  for all  $\mu$  if we choose our space to be  $X_0^{k+\alpha}$  or  $X_1^{k+\alpha}$  and Crandall-Rabinowitz theorem cannot be used. For this reason we choose our space to be  $X_2^{k+\alpha}$  (so that  $\cos(\theta)$  is excluded). The above lemma then implies that, in  $X_2^{k+\alpha}$ ,

$$\ker[F_S(0, \mu)] = \{0\} \quad \text{if } \mu \neq \mu_2, \mu_4, \mu_6, \dots. \quad (31)$$

It is also clear that, for each fixed  $\mu_l$ ,

$$\ker[F_S(0, \mu_l)] = \text{closure of the linear space spanned by } \{\cos(m\theta); \mu_m = \mu_l\}.$$

Thus, for each fixed  $l \geq 2$ , the dimension of  $\text{Ker } F_S(0, \mu_l)$  is 1 if and only if  $\mu_l \neq \mu_m$  for all  $m \neq l$ .

**Theorem 3.3.** *The points  $(0, \mu_n)$  ( $n \geq 2$  and even) are bifurcation points for the problem (6)-(9), and the corresponding free boundaries are of the form*

$$\partial\Omega_\epsilon: \quad r = R + \epsilon S(\theta) + O(\epsilon^2), \quad \partial D_\epsilon: \quad r = \rho + \epsilon \rho_1 S(\theta) + O(\epsilon^2),$$

where  $\rho_1$  is a constant given by (10).

We note that theorem establishes the bifurcations only for even  $n \geq 2$ . For odd  $n \geq 3$  we may apply the Crandall-Rabinowitz theorem in a more delicate manner.

To use the theorem, we need to show that all  $\mu_l$  are distinct. Now we consider the property of  $Q'_l(r; \rho, R)$ .

**Lemma 3.4.** (i).  $\{Q'_l(\rho; \rho, R)\}$  is a positive decreasing sequence in  $l$ ;  
(ii).  $\{Q'_l(R; \rho, R)\}$  is a positive increasing sequence in  $l$ .

*Proof.* We rewrite (7) as

$$\begin{cases} -\Delta Q_l + (\frac{l^2}{r^2} + 1)Q_l = 0 & \text{in } B_R \setminus \bar{D}_\rho, \\ Q_l = 1 & \text{on } \partial B_R, \\ Q_l = 0 & \text{on } \partial D_\rho. \end{cases}$$

By the maximum principle,  $0 < Q_l < (r/R)^l$  for  $\rho < r < R$ , so that by the strong maximum principle

$$Q'_l(\rho; \rho, R) > 0 \quad \text{and} \quad Q'_l(R; \rho, R) > \frac{l}{R}. \quad (32)$$

Let  $F = \frac{\partial Q_l}{\partial l}$ , then

$$\begin{cases} -\Delta F + (\frac{l^2}{r^2} + 1)F = -\frac{2l}{r^2}Q_l & \text{in } B_R \setminus \bar{D}_\rho, \\ F = 0 & \text{on } \partial B_R, \\ F = 0 & \text{on } \partial D_\rho. \end{cases}$$

Since  $Q_l > 0$  in  $B_R \setminus \bar{D}_\rho$ , we know  $F < 0$  in  $B_R \setminus \bar{D}_\rho$ . Thus,  $\frac{\partial F}{\partial r}|_{r=\rho} < 0$  and  $\frac{\partial F}{\partial r}|_{r=R} > 0$ , i.e.,  $Q'_l(\rho; \rho, R) > 0$  is a decreasing sequence in  $l$ , and  $Q'_l(R; \rho, R)$  is an increasing sequence in  $l$ .  $\square$

- Lemma 3.5.** (i). For any  $\rho < r \leq R$ ,  $\left\{ \frac{Q'_l(r; \rho, R)}{Q_l(r; \rho, R)} - \frac{l}{r} \right\}$  is a positive decreasing sequence in  $l$ ;
- (ii). for  $l \geq 2$ ,  $-\frac{l}{R} + Q'_l(R; \rho, R) < -\frac{l+1}{R} + Q'_{l+1}(R; \rho, R) + \frac{d_l}{l+1}$ , where  $d_l = \max_{\rho \leq r \leq R} \frac{2r^2}{R} \left\{ Q'_l(r; \rho, R) - \frac{l}{r} Q_l(r; \rho, R) \right\}$ .

*Proof.* For any  $\rho < s \leq R$ , it is clear that

$$\psi_1(r) = \frac{r}{s} Q_l(r; \rho, R) - \frac{Q_l(s; \rho, R)}{Q_{l+1}(s; \rho, R)} Q_{l+1}(r; \rho, R)$$

satisfies

$$\begin{cases} -\Delta \psi_1 + \left( \frac{(l+1)^2}{r^2} + 1 \right) \psi_1 &= \frac{2}{s} \left\{ \frac{l}{r} Q_l - Q'_l \right\} & \text{for } \rho < r < s, \\ \psi_1 &= 0 & \text{for } r = \rho, \\ \psi_1 &= 0 & \text{for } r = s. \end{cases}$$

Using (6) and (4), we find that

$$\left\{ \frac{l}{r} Q_l - Q'_l \right\} = \frac{-\left( K_l(\rho) I_{l+1}(r) + I_l(\rho) K_{l+1}(r) \right)}{K_l(\rho) I_l(R) - K_l(R) I_l(\rho)} < 0. \quad (33)$$

Thus we can use the maximum principle to derive  $\psi_1 < 0$  for  $\rho < r < s$ . In particular,  $\left. \frac{d\psi_1}{dr} \right|_{r=s} > 0$ , and this implies that

$$-\frac{l}{s} + \frac{Q'_l(s; \rho, R)}{Q_l(s; \rho, R)} > -\frac{l+1}{s} + \frac{Q'_{l+1}(s; \rho, R)}{Q_{l+1}(s; \rho, R)}. \quad (34)$$

Replacing  $s$  by  $r$  in the above inequality, we obtain (i).

To establish (ii), we let

$$\psi_2(r) = \frac{r}{R} Q_l(r; \rho, R) - Q_{l+1}(r; \rho, R) + \frac{d_l}{(l+1)^2} \left( 1 - \frac{r^{l+1}}{R^{l+1}} \right),$$

A direct computation shows

$$\begin{cases} -\Delta \psi_2 + \left( \frac{(l+1)^2}{r^2} + 1 \right) \psi_2 &= G & \text{for } \rho < r < R, \\ \psi_2 &= \frac{d_l}{(l+1)^2} \left( 1 - \frac{\rho^{l+1}}{R^{l+1}} \right) > 0 & \text{for } r = \rho, \\ \psi_2 &= 0 & \text{for } r = R, \end{cases}$$

where  $G = \frac{2}{R} \left\{ \frac{l}{r} Q_l - Q'_l \right\} + \frac{d_l}{(l+1)^2} \left( 1 - \frac{r^{l+1}}{R^{l+1}} \right) + \frac{d_l}{r^2}$ . By the definition of  $d_l$  it is clear that  $G > 0$ . In this case, we can apply the maximum principle to conclude  $\psi_2 > 0$  for  $\rho < r < R$  and hence  $\left. \frac{d\psi_2}{dr} \right|_{r=s} < 0$ . This implies that, for  $l \geq 2$ ,

$$-\frac{l}{R} + Q'_l(R; \rho, R) < -\frac{l+1}{R} + Q'_{l+1}(R; \rho, R) + \frac{d_l}{R(l+1)}. \quad (35)$$

This establishes the lemma.  $\square$

The following lemma will be established using computations performed in a reasonable range of  $R$  and  $\rho$ , namely,  $0 < R \leq 10$  and  $0 < \rho \leq 10$ .

**Lemma 3.6.**  $\mu_2 < \mu_3 < \mu_4 < \dots$

*Proof.* We can rewrite (26) as  $A_l \triangleq -\alpha - \beta_l - \gamma_l$ , where  $\alpha = -\frac{1}{2}\tilde{\sigma}(1 + \frac{\rho^2}{R^2}) + \sigma_s''(R)$  is independent of  $l$ , and using (10),

$$\beta_l = \frac{\rho_1 \rho^{l+1}}{R^{l+1}} \tilde{\sigma} = \sigma_s'(R) \frac{\tilde{\sigma}}{\underline{\sigma}} \left( \frac{\rho}{R} \right)^{l+1} Q_l'(\rho; \rho, R) \quad \text{and} \quad \gamma_l = -\sigma_s'(R) \left( Q_l'(R; \rho, R) - \frac{l}{R} \right).$$

Since  $\sigma_s'(R) > 0$ , the preceding lemma implies that

$$\gamma_l - \gamma_{l+1} \geq -\sigma_s'(R) \frac{d_l}{R(l+1)} \quad \text{for } l \geq 2. \quad (36)$$

From Lemmas 3.3 and 3.4, and the fact that  $\sigma_s'(R) > 0$ , it is clear that  $\beta_l$  is a decreasing sequence and that  $\gamma_l$  is an increasing sequence. The inequality  $\mu_l < \mu_{l+1}$  is equivalent to:

$$\frac{l^3 - l}{\alpha + \beta_l + \gamma_l} < \frac{(l+1)^3 - (l+1)}{\alpha + \beta_{l+1} + \gamma_{l+1}}. \quad (37)$$

We next show that  $\alpha + \beta_l + \gamma_l > 0$ . Clearly,

$$\alpha = -\frac{1}{2}\tilde{\sigma}(1 + \frac{\rho^2}{R^2}) + \sigma_s''(R) = 1 - \sigma_s'(R) \left( \frac{2R}{R^2 - \rho^2} \right).$$

By (22), we know  $\alpha > 0$ . On the other hand,  $\alpha + \beta_l + \gamma_l \geq \alpha + \gamma_l$ , the right-hand side of the above inequality is an increasing sequence in  $l$ . We will show  $\alpha + \gamma_2 > 0$ , which will then imply  $\alpha + \beta_l + \gamma_l > 0$  for all  $l \geq 2$ .

Clearly, (37) is equivalent to

$$3\alpha + (l+2)\beta_l - (l-1)\beta_{l+1} + (l+2)\gamma_l - (l-1)\gamma_{l+1} > 0. \quad (38)$$

We now proceed to estimate the left-hand side of the above inequality.

$$\begin{aligned} j(l, \rho, R) &:= 3\alpha + (l+2)\beta_l - (l-1)\beta_{l+1} + (l+2)\gamma_l - (l-1)\gamma_{l+1} \\ &= 3(\alpha + \beta_l + \gamma_l) + (l-1)(\beta_l - \beta_{l+1}) + (l-1)(\gamma_l - \gamma_{l+1}) \\ &> 3(\alpha + \gamma_l) - (l-1)\sigma_s'(R) \frac{d_l}{R(l+1)}. \end{aligned} \quad (39)$$

By Lemma 3.4 (i),  $d_l > d_{l+1}$  for all  $l \geq 1$ . Thus

$$j(l, \rho, R) > 3(\alpha + \gamma_4) - \sigma_s'(R) \frac{d_4}{R} \quad \text{for } l \geq 4, \quad (40)$$

$$j(3, \rho, R) > 3(\alpha + \gamma_3) - \frac{1}{2}\sigma_s'(R) \frac{d_3}{R}, \quad (41)$$

$$j(2, \rho, R) > 3(\alpha + \gamma_2) - \frac{1}{3}\sigma_s'(R) \frac{d_2}{R}. \quad (42)$$

The right-hand sides of the above inequalities are functions of  $\rho$  and  $R$ . We explicitly computed these three functions as functions of  $\rho$  and  $R$  and found that they are all positive (see Figures 1, 2, and 3). This also implies that  $\alpha + \gamma_2 > 0$ , which was used in the proof.  $\square$




Figure 1:  $3(\alpha + \gamma_4) - \sigma'_s(R) \frac{d_4}{R} \geq 0$




Figure 2:  $3(\alpha + \gamma_3) - \frac{1}{2} \sigma'_s(R) \frac{d_3}{R} \geq 0$

Consider

$$Y_1 = \text{closure of the linear space spanned by } \{\cos(m\theta); m \neq l\}.$$

We know

$$\text{Ker } [F_S(0, \mu_l)] = \text{span } \{\cos(l\theta)\}, \quad (43)$$

$$\text{Im } [F_S(0, \mu_l)] = Y_1, \quad (44)$$

$$[F_{S\mu}(0, \mu_l)] \cos(l\theta) \notin Y_1. \quad (45)$$






Figure 3:  $3(\alpha + \gamma_2) - \frac{1}{3}\sigma'_s(R) \geq 0$

It is clear that  $Y_1$  has codimension 1 since its direct sum with  $\{\cos(l\theta)\}$  gives the whole space. Thus we can apply Theorem 3.1 to get bifurcation branches for all  $\mu = \mu_l$  ( $l \geq 2$  even).

#### 4. Linear stability

Assume

$$\partial\Omega_0 : \quad r = R_0(\theta). \quad (1)$$

Then the linearization of the two boundaries  $\partial\Omega(t)$  and  $\partial D(t)$  are given by

$$\partial\Omega_\epsilon : \quad r = R_0(\theta) + \epsilon R_1(\theta, t) + O(\epsilon^2), \quad (2)$$

$$\partial D_\epsilon : \quad r = \rho_0(\theta) + \epsilon \rho_1(\theta, t) + O(\epsilon^2), \quad (3)$$

$$\sigma = \sigma_0(r, \theta) + \epsilon \sigma_1(r, \theta, t) + O(\epsilon^2), \quad (4)$$

$$p = p_0(r, \theta) + \epsilon p_1(r, \theta, t) + O(\epsilon^2). \quad (5)$$

The mean curvature is given by

$$\kappa = \frac{r^2 + 2r_\theta^2 - rr_{\theta\theta}}{(r^2 + r_\theta^2)^{\frac{3}{2}}}. \quad (6)$$

So that the linearization of  $\kappa$  is given by

$$\begin{aligned} \kappa|_{\partial\Omega_\epsilon} &= \frac{r^2 + 2r_\theta^2 - rr_{\theta\theta}}{(r^2 + r_\theta^2)^{\frac{3}{2}}} \\ &= \frac{(R_0 + \epsilon R_1)^2 + 2(R_{0\theta} + \epsilon R_{1\theta})^2 - (R_0 + \epsilon R_1)(R_{0\theta\theta} + \epsilon R_{1\theta\theta})}{[(R_0 + \epsilon R_1)^2 + (R_{0\theta} + \epsilon R_{1\theta})^2]^{\frac{3}{2}}} + O(\epsilon^2) \\ &\doteq \kappa_0 + \epsilon \kappa_1 + O(\epsilon^2), \end{aligned}$$

where

$$\kappa_0 = \frac{R_0^2 + 2R_{0\theta}^2 - R_0R_{0\theta\theta}}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}}, \quad (7)$$

and

$$\begin{aligned} \kappa_1 &= \frac{2R_0R_1 + 4R_{0\theta}R_{1\theta} - R_1R_{0\theta\theta} - R_0R_{1\theta\theta}}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}} \\ &\quad - \frac{3}{2} \frac{(R_0^2 + 2R_{0\theta}^2 - R_0R_{0\theta\theta})(2R_0R_1 + 2R_{0\theta}R_{1\theta})}{(R_0^2 + R_{0\theta}^2)^{\frac{5}{2}}}. \end{aligned} \quad (8)$$

Clearly, the linearized equations for  $\sigma$  and  $p$  are:

$$\begin{cases} \sigma_{1t} - \Delta\sigma_1 + \sigma_1 = 0 & \text{in } \Omega_0 \\ -\Delta p_1 = \mu\sigma_1 & \text{in } \Omega_0. \end{cases} \quad (9)$$

The boundary conditions are

$$\begin{aligned} 1 &= \sigma|_{r=R_0+\epsilon R_1+O(\epsilon^2)} \\ &= \sigma_0|_{r=R_0} + \epsilon R_1\sigma_{0r}|_{r=R_0} + \epsilon\sigma_1|_{r=R_0} + O(\epsilon^2) \\ &= 1 + \epsilon R_1\sigma_{0r}|_{r=R_0} + \epsilon\sigma_1|_{r=R_0} + O(\epsilon^2). \end{aligned}$$

It follows that the boundary condition for  $\sigma_1$  is

$$\sigma_1|_{r=R_0} = -R_1 \cdot \sigma_{0r}|_{r=R_0}. \quad (10)$$

Similarly,

$$\begin{aligned} (\kappa_0 + \epsilon\kappa_1)|_{\partial\Omega_\epsilon} &= \kappa|_{r=R_0+\epsilon R_1+O(\epsilon^2)} \\ &= (p_0 + \epsilon p_1)|_{r=R_0+\epsilon R_1} + O(\epsilon^2) \\ &= p_0|_{r=R_0} + p_{0r}|_{r=R_0} \cdot \epsilon R_1 + \epsilon p_1|_{r=R_0} + O(\epsilon^2) \\ &= \kappa_0 + p_{0r}|_{r=R_0} \cdot \epsilon R_1 + \epsilon p_1|_{r=R_0} + O(\epsilon^2) \end{aligned}$$

that is,

$$p_1|_{r=R_0} = \kappa_1 - R_1 \cdot p_{0r}|_{r=R_0}. \quad (11)$$

To derive the equation for  $R_1$ , we note that in the polar coordinate system

$$\begin{cases} \vec{e}_r = \cos\theta \vec{e}_1 + \sin\theta \vec{e}_2; \\ \vec{e}_\theta = -\sin\theta \vec{e}_1 + \cos\theta \vec{e}_2. \end{cases} \quad (12)$$

If we write in vector form

$$\vec{e}_r = (\cos\theta, \sin\theta)^T, \quad \vec{e}_\theta = (-\sin\theta, \cos\theta)^T \quad (13)$$

we then have

$$\nabla_{(x,y)} = \vec{e}_r \partial_r + \vec{e}_\theta \cdot \frac{1}{r} \partial_\theta, \quad (14)$$

that is

$$\begin{pmatrix} \partial_x \\ \partial_y \end{pmatrix} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \partial_r + \frac{1}{r} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \partial_\theta.$$

For the surface  $F(x, y) = 0$ , the normal is given by  $\vec{n} = \frac{\pm \nabla F}{|\nabla F|}$ , thus, we have

$$\vec{n} \Big|_{r=R_0+\epsilon R_1} = \frac{\vec{e}_r - \frac{1}{r}(R_{0\theta} + \epsilon R_{1\theta})\vec{e}_\theta}{\sqrt{1 + \frac{1}{r^2}(R_{0\theta} + \epsilon R_{1\theta})^2}}. \quad (15)$$

In particular,

$$\begin{aligned} \vec{n} \Big|_{r=R_0+\epsilon R_1} &= \frac{r\vec{e}_r - (R_{0\theta} + \epsilon R_{1\theta})\vec{e}_\theta}{\sqrt{r^2 + (R_{0\theta} + \epsilon R_{1\theta})^2}} \\ &= \frac{(R_0 + \epsilon R_1)\vec{e}_r - (R_{0\theta} + \epsilon R_{1\theta})\vec{e}_\theta}{((R_0 + \epsilon R_1)^2 + (R_{0\theta} + \epsilon R_{1\theta})^2)^{\frac{1}{2}}} \\ &\doteq \vec{n}_0 + \epsilon \vec{n}_1 + O(\epsilon^2), \end{aligned}$$

where

$$\begin{aligned} \vec{n}_0 &= \frac{R_0\vec{e}_r - R_{0\theta}\vec{e}_\theta}{(R_0^2 + R_{0\theta}^2)^{\frac{1}{2}}}, \\ \vec{n}_1 &= \frac{R_1\vec{e}_r - R_{1\theta}\vec{e}_\theta}{(R_0^2 + R_{0\theta}^2)^{\frac{1}{2}}} - \frac{(R_0R_1 + R_{0\theta}R_{1\theta})(R_0\vec{e}_r - R_{0\theta}\vec{e}_\theta)}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}} \\ &= \frac{R_1R_{0\theta}^2 - R_0R_{0\theta}R_{1\theta}}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}}\vec{e}_r + \frac{R_0R_1R_{0\theta} - R_0^2R_{1\theta}}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}}\vec{e}_\theta. \end{aligned}$$

Since

$$\begin{cases} p_{0r} \Big|_{r=R_0+\epsilon R_1} = p_{0r} \Big|_{r=R_0} + p_{0rr} \Big|_{r=R_0} \cdot \epsilon R_1 + O(\epsilon^2); \\ p_{0\theta} \Big|_{r=R_0+\epsilon R_1} = p_{0\theta} \Big|_{r=R_0} + p_{0\theta r} \Big|_{r=R_0} \cdot \epsilon R_1 + O(\epsilon^2), \end{cases} \quad (16)$$

we have

$$\begin{aligned} \frac{\partial p}{\partial n} \Big|_{r=R_0+\epsilon R_1} &= \left\{ \vec{e}_r(p_{0r} + \epsilon p_{1r}) + \frac{\vec{e}_\theta}{r}(p_{0\theta} + \epsilon p_{1\theta}) \right\} \Big|_{r=R_0+\epsilon R_1} \cdot (\vec{n}_0 + \epsilon \vec{n}_1) \\ &= \left\{ \vec{e}_r(p_{0r} + \epsilon R_1 p_{0rr} + \epsilon p_{1r}) + \frac{\vec{e}_\theta(p_{0\theta} + \epsilon R_1 p_{0\theta r} + \epsilon p_{1\theta})}{R_0 + \epsilon R_1} \right\} \cdot (\vec{n}_0 + \epsilon \vec{n}_1) \\ &= \left\{ (\vec{e}_r p_{0r} + \vec{e}_\theta \frac{p_{0\theta}}{R_0}) + \epsilon [(R_1 p_{0rr} + p_{1r})\vec{e}_r + \frac{R_0 R_1 p_{0\theta r} + R_0 p_{1\theta} - R_1 p_{0\theta}}{R_0^2} \vec{e}_\theta] \right\} \\ &\quad \cdot (\vec{n}_0 + \epsilon \vec{n}_1). \end{aligned}$$

Notice that  $p_0$  is a stationary solution with

$$0 = \frac{\partial p_0}{\partial n_0} \Big|_{r=R_0} = \frac{p_{0r}}{(1 + \frac{1}{r^2} R_{0\theta}^2)^{\frac{1}{2}}} - \frac{\frac{1}{r^2} p_{0\theta} R_{0\theta}}{(1 + \frac{1}{r^2} R_{0\theta}^2)^{\frac{1}{2}}}. \quad (17)$$

The  $O(\epsilon)$  term is

$$\begin{aligned}
& (\vec{e}_r p_{0r} + \vec{e}_\theta \frac{p_{0\theta}}{R_0}) \cdot \vec{n}_1 + [(R_1 p_{0rr} + p_{1r}) \vec{e}_r + \frac{R_0 R_1 p_{0\theta r} + R_0 p_{1\theta} - R_1 p_{0\theta}}{R_0^2} \vec{e}_\theta] \cdot \vec{n}_0 \\
&= \frac{p_{0r}(R_1 R_{0\theta}^2 - R_0 R_{0\theta} R_{1\theta})}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}} + \frac{p_{0\theta}(R_0 R_1 R_{0\theta} - R_0^2 R_{1\theta})}{R_0 (R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}} + \frac{R_0 (R_1 p_{0rr} + p_{1r})}{(R_0^2 + R_{0\theta}^2)^{\frac{1}{2}}} \\
&\quad - \frac{R_{0\theta}(R_0 R_1 p_{0\theta r} + R_0 p_{1\theta} - R_1 p_{0\theta})}{R_0^2 (R_0^2 + R_{0\theta}^2)^{\frac{1}{2}}}.
\end{aligned}$$

On the other hand, the velocity of the free boundary  $\partial\Omega_\epsilon$  in the direction  $\vec{n}$  would be

$$\begin{aligned}
v_n &= \frac{(R_0 + \epsilon R_1)_t}{\sqrt{1 + \frac{1}{(R_0 + \epsilon R_1)^2} (R_{0\theta} + \epsilon R_{1\theta})^2}} \\
&= \frac{\epsilon R_{1t}}{\sqrt{1 + \frac{1}{(R_0 + \epsilon R_1)^2} (R_{0\theta}^2 + 2\epsilon R_{0\theta} R_{1\theta} + \epsilon^2 R_{1\theta}^2)}} \\
&= \epsilon \frac{R_{1t}}{\sqrt{1 + \frac{1}{R_0^2} R_{0\theta}^2}} + O(\epsilon^2)
\end{aligned}$$

that is,  $-v_n = -\epsilon \frac{R_{1t} R_0}{(R_0^2 + R_{0\theta}^2)^{\frac{1}{2}}} + O(\epsilon^2)$ .

Since  $\frac{\partial p}{\partial n} = -v_n$  on  $\partial\Omega_t$ ,

$$\begin{aligned}
-R_{1t} R_0 &= \frac{p_{0r}(R_1 R_{0\theta}^2 - R_0 R_{0\theta} R_{1\theta})}{R_0^2 + R_{0\theta}^2} + \frac{p_{0\theta}(R_0 R_1 R_{0\theta} - R_0^2 R_{1\theta})}{R_0 (R_0^2 + R_{0\theta}^2)} \\
&\quad + R_0 (R_1 p_{0rr} + p_{1r}) - \frac{R_{0\theta}(R_0 R_1 p_{0\theta r} + R_0 p_{1\theta} - R_1 p_{0\theta})}{R_0^2}.
\end{aligned}$$

We derive

$$\begin{aligned}
-R_{1t} &= \frac{p_{0r} R_1 \rho_{0\theta}^2 - p_{0r} \rho_0 \rho_{0\theta} \rho_{1\theta} + p_{0\theta} \rho_1 \rho_{0\theta} - p_{0\theta} \rho_{1\theta} \rho_0}{\rho_0 (\rho_0^2 + \rho_{0\theta}^2)} + \rho_1 p_{0rr} + p_{1r} \\
&\quad - \rho_{0\theta} \frac{\rho_1 p_{0\theta r} + p_{1\theta}}{\rho_0^2} + \frac{\rho_{0\theta} \rho_1 p_{0\theta}}{\rho_0^3} \\
&= \left[ \frac{p_{0r} \rho_{0\theta}^2}{\rho_0 (\rho_0^2 + \rho_{0\theta}^2)} + \frac{p_{0\theta} \rho_{0\theta}}{\rho_0 (\rho_0^2 + \rho_{0\theta}^2)} + p_{0rr} - \frac{\rho_{0\theta} p_{0\theta r}}{\rho_0^2} + \frac{\rho_{0\theta} p_{0\theta}}{\rho_0^3} \right] \rho_1 \\
&\quad - \left[ \frac{p_{0r} \rho_{0\theta}}{\rho_0^2 + \rho_{0\theta}^2} + \frac{p_{0\theta}}{\rho_0^2 + \rho_{0\theta}^2} \right] \rho_{1\theta} + \left[ p_{1r} - \frac{\rho_{0\theta} p_{1\theta}}{\rho_0^2} \right].
\end{aligned}$$

Thus

$$R_{1t} = \left[ -\frac{p_{0r}\rho_{0\theta}^2}{\rho_0(\rho_0^2 + \rho_{0\theta}^2)} - \frac{p_{0\theta}\rho_{0\theta}}{\rho_0(\rho_0^2 + \rho_{0\theta}^2)} - p_{0rr} + \frac{\rho_{0\theta}p_{0\theta r}}{\rho_0^2} - \frac{\rho_{0\theta}p_{0\theta}}{\rho_0^3} \right] \rho_1 \\ + \left[ \frac{p_{0r}\rho_{0\theta}}{\rho_0^2 + \rho_{0\theta}^2} + \frac{p_{0\theta}}{\rho_0^2 + \rho_{0\theta}^2} \right] \rho_{1\theta} + \left[ \frac{\rho_{0\theta}p_{1\theta}}{\rho_0^2} - p_{1r} \right].$$

From (13) and (20), we can get the linear equations on the boundary  $\partial D_0$ :

$$\sigma_{1r}|_{r=\rho_0} = -\rho_1 \frac{\partial^2 \sigma_0}{\partial n^2} |_{r=\rho_0} \\ \frac{\partial p_1}{\partial n} |_{+} - \frac{\partial p_1}{\partial n} |_{-} = \mu(\sigma - \tilde{\sigma})\rho_1,$$

where


$$\frac{\partial}{\partial n} |_{r=\rho} = \frac{\rho^2 \partial_r - \rho_{\theta} \partial_{\theta}}{\rho \sqrt{\rho^2 + \rho_{\theta}^2}}, \\ \frac{\partial^2}{\partial n^2} |_{r=\rho} = \frac{\rho_{\theta}^2 (\rho + \rho_{\theta\theta}) \partial_r + \rho^2 (\rho^2 + \rho_{\theta}^2) \partial_{rr} - 2\rho_{\theta} (\rho^2 + \rho_{\theta}^2) \partial_{\theta r}}{(\rho^2 + \rho_{\theta}^2)^2} \\ + \frac{\rho \rho_{\theta} (2\rho^2 + \rho_{\theta}^2 + \rho \rho_{\theta\theta}) \partial_{\theta} + \rho_{\theta}^2 (\rho^2 + \rho_{\theta}^2) \partial_{\theta\theta}}{\rho^2 (\rho^2 + \rho_{\theta}^2)^2}.$$

Combing all the above computation, the linearized system is

$$\begin{aligned} \sigma_{1t} &= \Delta \sigma_1 - \sigma_1 \text{ in } \Omega_0, \\ \Delta p_1 &= -\mu \sigma_1 \text{ in } \Omega_0, \\ \sigma_{1r} &= -\rho_1 \frac{\partial^2 \sigma_0}{\partial n^2} \text{ on } \partial D_0, \\ \mu(\sigma - \tilde{\sigma})\rho_1 &= \frac{\partial p_1}{\partial n} |_{+} - \frac{\partial p_1}{\partial n} |_{-} \text{ on } \partial D_0, \\ \sigma_1 &= -R_1 \sigma_{0r} \text{ on } \partial \Omega_0, \\ \frac{\partial p}{\partial n} &= 0 \text{ on } \partial \Omega_0, \\ p_1 &= \frac{2R_0 R_1 + 4R_{0\theta} R_{1\theta} - R_1 R_{0\theta\theta} - R_0 R_{1\theta\theta}}{(R_0^2 + R_{0\theta}^2)^{\frac{3}{2}}} - R_1 p_{0r} \\ &\quad - \frac{3}{2} \frac{(R_0^2 + 2R_{0\theta}^2 - R_0 R_{0\theta\theta})(2R_0 R_1 + 2R_{0\theta} R_{1\theta})}{(R_0^2 + R_{0\theta}^2)^{\frac{5}{2}}} \text{ on } \partial \Omega_0, \\ R_{1t} &= \left( -\frac{p_{0r} R_{0\theta}^2}{R_0 (R_0^2 + R_{0\theta}^2)} - p_{0rr} + \frac{p_{0\theta r} R_{0\theta}}{R_0^2} - \frac{p_{0\theta} R_{0\theta}}{R_0^3} \right. \\ &\quad \left. - \frac{p_{0\theta} R_{0\theta}}{R_0 (R_0^2 + R_{0\theta}^2)} \right) R_1 + \left( \frac{p_{0r} R_{0\theta}}{R_0^2 + R_{0\theta}^2} + \frac{p_{0\theta}}{R_0^2 + R_{0\theta}^2} \right) R_{1\theta} \\ &\quad + \left( \frac{p_{1\theta} R_{0\theta}}{R_0^2} - p_{1r} \right) \text{ on } \partial \Omega_0. \end{aligned} \tag{18}$$

We tested numerically for linear stability. For radial symmetric solutions, we use (6) and (7) to obtain the solutions. For non-radial symmetric solutions, we use (1)-(4) and drop the higher term to get approximations of solutions in a small neighborhood of  $\mu_2$ : we take  $\epsilon$  to be a small value such as  $10^{-2}, 10^{-3}$ . Once the solution is computed, we can recover the parameter  $\mu$  from the linearized equation (18) and test linear stability by checking the spectrum of the linearized system. Doing this, the linear stability in a small neighborhood of  $\mu_2$  has been numerically verified and is shown in Figure 4.

Figure 4: Linear stability of the solution branches



## 5. Conclusion

Although the tumor model with a necrotic core analyzed in this paper is quite simple, we may nevertheless draw some interesting biological conclusions from the mathematical results. Tumors grown in culture are typically sphere. However, tumor grown in vivo may have a variety of shapes. In particular, a tumor with a necrotic core is associated with the growth of protrusions. In our model, these protrusions are expressed by the shape

$$r = R + \epsilon \cos(n\theta) + O(\epsilon^2), r = \rho + \epsilon \rho_1 \cos(n\theta) + O(\epsilon^2)$$

of the free boundaries; the number of protrusions depend on  $n$ . The aggressiveness of a tumor is measured by the parameter  $\mu$ . the larger the  $\mu$  is the more aggressive the tumor is. As this parameter increases, the tumor will lose its spherical shape, develop fingers, and become invasive. The linear stability asserts that radial instability occurs when  $\mu$  reaches the first bifurcation point  $\mu_2$ . Moreover, there is non-radial stability for part of a neighborhood of  $\mu_2$  on a non-radial branch. The numerical algorithms we use to carry out these computations will be presented in another paper, which in particular show that the non-radial solutions have non-spherical shapes.

## References

- [1] J.A.ADAM, General aspect of modeling tumor growth and immune response, A Survey of Models for Tumor-Immune System Dynamics (J.A.Adam and N. Bellomo, eds.) Birkhäuser, Boston, 14–87, (1996).
- [2] J.A.ADAM AND S.A. MAGGELAKIS, Diffusion regulated growth characteristics of a spherical prevascular carcinoma, *Bull. Math. Biol.*, Vol 52, 549–582, (1990).
- [3] B. BAZALLY AND A. FRIEDMAN, Global existence and asymptotic stability for an elliptic-parabolic free boundary problem: an application to a model of tumor growth, *Indiana Univ. Math. J.*, Vol 52, 1265–1304, (2003).
- [4] N. BRITTON AND M.A.J. CHAPLAIN, A qualitative analysis of some models of tissue growth, *Math. Biosci.*, Vol 113, 77–89, (1993).
- [5] H.M. BYRNE AND M.A.J. CHAPLAIN, Growth of nonnecrotic tumors in the presence and absence of inhibitors, *Math. Biosci.*, Vol 130, 151–181, (1995).
- [6] H.M. BYRNE AND M.A.J. CHAPLAIN, Modelling the role of cell-cell adhesion in the growth and development of carcinomas, *Math. Comput. Modelling*, Vol 12, 1–17, (1996).
- [7] M.A.J. CHAPLAIN, The development of a spatial pattern in a model for cancer growth, *Experimental and Theoretical Advances in Biological Pattern Formation* (H.G. Othmer, P.K. Maini, and J.D. Murray, eds), Plenum Press, 45–60, (1993).
- [8] M. G. CRANDALL AND P. H. RABINOWITZ, Bifurcation from simple eigenvalues, *J. Funct. Anal.*, Vol 8, 321 C 340, (1971).
- [9] S. CUI AND A. FRIEDMAN, Analysis of a Mathematical Model of the Growth of Necrotic Tumors, *Journal of Mathematical Analysis and Applications* 255, 636–677, (2001).
- [10] S. CUI AND J. ESCHER ,Bifurcation Analysis of an Elliptic Free Boundary Problem Modelling the Growth of Avascular Tumors, *SIAM Journal on Mathematical Analysis* Vol 39, 210-235 (2007)

- [11] S. CUI AND J. ESCHER ,Asymptotic Behaviour of Solutions of a Multidimensional Moving Boundary Problem Modeling Tumor Growth, *Communications in Partial Differential Equations* Vol 32, 636 - 655 (2008)
- [12] M. FONTELOS AND A. FRIEDMAN, Symmetry-breaking bifurcations of free boundary problems in three dimensions, *Asymptotic Analysis*, Vol 35, 187–206, (2003).
- [13] A. FRIEDMAN AND B. HU, Bifurcation from stability to instability for a free boundary problem arising in a tumor model *Arch. Rat. Mech. Anal.*, Vol. 180, No. 2, (2006).
- [14] A. FRIEDMAN AND F. REITICH, Analysis of a mathematical model for growth of tumor, *J. Math. Biology*, Vol 38, 262–284, (1999).
- [15] A. FRIEDMAN AND F. REITICH, Nonlinear stability of a quasi-static Stefan problem with surface tension: A continuation approach, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, Vol. 30(4), pp. 341–403,(2001).
- [16] H.P. GREENSPAN, Models for the growth of a solid tumor by diffusion, *Studies Appl. Math*, Vol 52, 317–340, (1972).
- [17] H.P. GREENSPAN, On the growth of cell culture and solid tumors, *Theoretical Biology*, Vol 56, 229–242, (1976).
- [18] S.A. MAGGELAKIS AND J.A. ADAM, Mathematical model for prevasculat growth of a spherical carcinoma, *Math. Comp. Modeling*, Vol 13, 23–38, (1990).
- [19] D.L.S. MCEWAIN AND L.E. MORRIS, Apoptosis as a volume loss mechanism in mathematical models of solid tumor growth, *Math. Biosci.*, Vol 39, 147–157, (1978).
- [20] F. ZHOU, J. ESCHER AND S. CUI, Well-posedness and stability of a free boundary problem modeling the growth of multi-layer tumors , *Journal of Differential Equations* Vol 224, 2909-2933 (2008)