

 Open access • Journal Article • DOI:10.1137/0521081

Bifurcations of relative equilibria — [Source link](#)

Martin Krupa

Institutions: University of Minnesota

Published on: 01 Sep 1990 - Siam Journal on Mathematical Analysis (Society for Industrial and Applied Mathematics)

Topics: Equivariant map

Related papers:

- [Singularities and groups in bifurcation theory](#)
- [Equivariant dynamical systems](#)
- [Dynamics and symmetry. Predictions for modulated waves in rotating fluids](#)
- [Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields](#)
- [Elements of applied bifurcation theory](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/bifurcations-of-relative-equilibria-8fqcrr0qs>

BIFURCATIONS OF RELATIVE EQUILIBRIA

By

Martin Krupa

IMA Preprint Series # 615

February 1990

BIFURCATIONS OF RELATIVE EQUILIBRIA

Martin Krupa
Institute for Mathematics and its Applications
University of Minnesota
Minneapolis, MN. 55455

November 12, 1989

Abstract.

In this paper we discuss the dynamics and bifurcation theory of equivariant dynamical systems near *relative equilibria*, that is group orbits invariant under the flow of an equivariant vector field. The theory we develop applies, in particular, to secondary steady-state bifurcations from invariant equilibria. Let Γ be a compact group of symmetries of \mathbb{R}^n and let x_0 be in \mathbb{R}^n . Suppose that f is a smooth Γ -equivariant vector field and Σ the isotropy group of x_0 . We show that there exists a Σ -equivariant vector field f_N , defined on the space normal to X at x_0 , and that the local asymptotic dynamics of f are closely related to the local asymptotic dynamics of f_N . Next we study bifurcations of X occurring when an eigenvalue of $(df_N)_x$ crosses the imaginary axis. Properties of the vector field f_N imply that branches of equilibria and periodic orbits of f_N correspond to trajectories of f which are dense in tori. Field [1980] found bounds on the dimensions of these tori. We extend some of his results. We apply this theory to the following specific problems:

1. Bifurcations of systems with $O(2)$ symmetry.
2. Bifurcations of steady-state solutions of the Kuramoto-Sivashinsky equation.
3. Secondary bifurcations in the planar Bénard problem.

Introduction.

In this work we discuss the dynamics and bifurcation theory of equivariant dynamical systems near group orbits invariant under the action of the flow. Such group orbits are called *relative equilibria*. The simplest example of a relative equilibrium is a group orbit of equilibria. A group orbit of equilibria can be characterized as a relative equilibrium on which the flow is trivial. The symmetry groups we consider are compact and have positive dimension, so, in particular, they must contain a subgroup isomorphic to $SO(2)$. For such groups the flow trajectories on the relative equilibria can be nontrivial. A well known example of such nontrivial trajectories on relative equilibria are rotating waves, that is solutions given by $x(t) = \theta(t)x_0$, where $\theta(t)$ parametrizes $SO(2)$.

A special case of a relative equilibrium is an invariant equilibrium, that is an equilibrium invariant under all the symmetries of the system. Such equilibria often arise in applications and their bifurcations have been extensively studied. Often bifurcations of invariant equilibria are characterized by symmetry breaking, that is, the invariant equilibrium bifurcates to branches of equilibria no longer invariant under the action of the symmetry group. In other words, nontrivial relative equilibria often occur as a result of bifurcations of invariant equilibria. In this context bifurcations of relative equilibria correspond to secondary bifurcations from an invariant equilibrium.

Let X be a group orbit of equilibria of an equivariant vector field f . If X has positive dimensions then the conditions determining when f can undergo a bifurcation near X are quite different than in the case of an invariant equilibrium. In particular, no element of X can be hyperbolic, since the directions along the group orbit must be neutrally stable. More precisely, for any $x \in X$ the tangent space $T_x X$ is contained in the kernel of the derivative $(df)_x$. It follows that X will be normally hyperbolic if $(df)_x$ has no purely imaginary eigenvalues and the algebraic multiplicity of zero as an eigenvalue of $(df)_x$ equals the dimension of $T_x X$. A bifurcation of X will occur if $(df)_x$ has an eigenvalue on the imaginary axis whose (generalized) eigenvector is not contained in the tangent space of X .

An interesting phenomenon has been observed in examples of bifurcations of relative equilibria: an orbit of equilibria can lose stability by having an eigenvalue passing through zero and bifurcate to a group orbit consisting of nontrivial flow trajectories. The flow on the new relative equilibria is a slow drift given by the action of a curve of group elements. Several authors, who studied bifurcations of relative equilibria, found that the resulting dynamics could be described in terms of dynamics related to standard bifurcations modulated by a drift along the group orbit. The following articles focused on bifurcations

of relative equilibria where this feature has been observed. Chossat [1986] showed that the bifurcation of standing waves in the problem of degenerate Hopf bifurcation with $O(2)$ symmetry leads to quasiperiodic motion on a group invariant two dimensional torus. Iooss [1986] showed that a Hopf-Hopf mode interaction in the Taylor-Couette problem ($O(2) \times SO(2)$ symmetry) led to a three frequency flow. Danglemayr [1986] found a rotating wave in the problem of steady state mode interaction with $O(2)$ symmetry. Chossat and Golubitsky [1988] studied a related problem of Hopf bifurcation of a group orbit of standing waves and discovered that this bifurcation led to a three frequency motion, with one of the frequencies given by the drift along the orbit. In their paper Chossat and Golubitsky formulated the following theorem: the flow near a relative equilibrium can be decomposed into the flow in the direction along the orbit and the flow in the direction normal to the orbit. The precise statement and the proof of this theorem is the starting point of this work.

Section 1 of the paper contains some background information on Lie group theory. The remaining part of the paper is divided into two parts. The first part, Sections 2-5, is devoted to the theoretical aspects of the problem. The second part, Sections 6-8, focuses on specific group actions and specific dynamical systems and is designed to show the application of the ideas developed in the first part. The reader more interested in the second part of the paper will only need to know the definitions and the statements of theorems contained in the first part. The following is a brief description of the topics discussed in each of the sections.

In Section 2 we give a precise description of how the previously mentioned decomposition of the vector field can be accomplished. We show that near relative equilibria the vector field can be written as a sum of equivariant components: one tangent to the group orbits and the other normal to the original orbit X (Theorem 2.1). As a consequence of this decomposition each bounded solution near a relative equilibrium is contained in the group orbit of a solution of the normal vector field f_N (Theorem 2.2). In the remainder of Section 2 we show that the asymptotic dynamics of f can be determined by the asymptotic dynamics of the normal vector field modulo drifts along the orbit. Some results of Section 2, including an alternative proof of Theorem 2.1, can be found in Vanderbauwhede, Krupa and Golubitsky [1989].

The results of Section 2 imply that bifurcations of f can be analyzed in two steps. The first step is to describe bifurcations of the normal vector field f_N and the second step is to find the corresponding drifts along group orbits. Let x be in X . In Section 3 we

argue that generic bifurcations of f_N can be described as bifurcations of a generic Σ_x -equivariant vector field, where Σ_x is the isotropy subgroup of x .

Suppose that f describes a family of vector fields, rather than a single vector field. In Section 4 we study bifurcations of relative equilibria occurring when an eigenvalue of $(df_N)_x$ passes through zero. We analyze the case when a relative equilibrium X bifurcates to another relative equilibrium Y . Let $y \in Y$, let Σ be the isotropy subgroup of y and $N(\Sigma)$ the normalizer of Σ . Field [1980] proves a theorem stating that the flow on Y is given by a linear flow on a torus whose dimension is bounded by, and generically equal to $\text{rank}(N(\Sigma)/\Sigma)$ ($\text{rank}(N(\Sigma)/\Sigma)$ equals the dimension of a maximal torus in $N(\Sigma)/\Sigma$). The main theorem of Section 4 (Theorem 4.1') states that there exists a generic set of perturbations of f whose elements have the following property: for all except countably many values of the parameter the dimension of the flow on Y is maximal.

In Section 5 we study Hopf bifurcations of relative equilibria, that is bifurcations occurring when an eigenvalue of $(df_N)_x$ passes through a nonzero point on the imaginary axis. We apply the standard Hopf bifurcation theorems to find periodic solutions of the normal component f_N . Let Y be a periodic orbit of f_N and let Σ be the isotropy subgroup of the elements of Y . Field [1980] shows that the corresponding trajectories of f are dense in tori whose dimension is bounded by $\text{rank}(N(\Sigma)/\Sigma) + 1$. Let $\tilde{\Sigma}$ be the group consisting of all the symmetries that leave Y invariant. In Theorem 5.1 we derive a new bound, given by $\text{rank}(N(\tilde{\Sigma})/\tilde{\Sigma}) + 1$ and show that this bound is attained for a generic vector field. Next, in Theorem 5.2, we consider a family of vector fields f , such that f_N has a Hopf bifurcation and show that there exists a generic set of perturbations of f whose elements are such that for all except countably many values of the parameter the dimension of the flow on the manifolds ΓY is maximal.

In Section 6 we present a classification of generic secondary steady-state and Hopf bifurcations with symmetry group $O(2)$. In this context bifurcations of the normal vector field correspond to steady state and Hopf bifurcations with D_k symmetry. Using the results of Sections 3 and 4 we determine for which bifurcations of the vector field f_N the bifurcating solutions of the full vector field f generically have nontrivial drift along group orbits.

In Section 7 we analyze bifurcations of the zero solution of the Kuramoto-Shivashinsky equation, which has $O(2)$ symmetry. We summarize the results of a computer assisted study done by Kevrekidis, Nicolaenco and Scovel [1988] and compare

their numerical results with the predictions of $O(2)$ bifurcations, as described in Section 6.

In Section 8 we classify the possible generic steady-state bifurcations in the planar Bénard problem. The generic primary bifurcations in the Bénard problem are to two types of equilibria: hexagons (with symmetry D_6) and rolls (with symmetry $O(2) \oplus Z_2$). We consider secondary steady-state bifurcations of hexagons and rolls and show that the resulting trajectories are either equilibria or rotating waves.

The author would like to thank Marty Golubitsky for very helpful suggestions and comments. The author is also grateful to Mike Field and André Vanderbauwhede for helpful discussions. This research was supported in part by a grant from the ACMP program of DARPA.

1. Preliminaries

Let Γ be a compact Lie group. We consider a smooth linear action of Γ on \mathbb{R}^n . With no loss of generality we can assume that this action is orthogonal and hence identify Γ with a subgroup of $O(n)$ (see I, 3.5 in Bredon [1972]). Let X be a compact and Γ -invariant submanifold of \mathbb{R}^n . For $x \in X$ let N_x be the set of vectors normal to X . Note that N_x is a vector subspace of \mathbb{R}^n , since it passes through 0. Let $N(X)$ be the bundle with base space X and fibers N_x ; $N(X)$ is called the *normal bundle* of X . The bundle $N(X)$ is smooth (see Guillemin and Pollack [1974] pg.71). The action of Γ on $N(X)$ is defined by the formula $\gamma(x, v) = (\gamma x, \gamma v)$. To see that this action is well defined observe that the orthogonality of the action of Γ implies that $\gamma N_x = N_{\gamma x}$. Let $\beta : N(X) \rightarrow \mathbb{R}^n$ be defined as $\beta((x, u)) = x + u$. It is easy to see that the map β is Γ -equivariant and a local diffeomorphism. It follows that an invariant neighborhood of X in \mathbb{R}^n can be identified, via the map β , with a neighborhood of the zero section in $N(X)$. For $x \in X$ let $\hat{N}_x = \{ (x, v) : v \in N_x \}$. Note that $\hat{N}_x \subset N(X)$ and the image of \hat{N}_x under β is in N_x . Given a Γ -equivariant vector field $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ let β^*f be defined by $\beta^*f(y) = [d\beta_{\beta(y)}]^{-1}f(\beta(y))$, $y \in N(X)$. The map β^*f is called the *pullback* of f to $N(X)$. Let $x \in \mathbb{R}^n$ and $X = \Gamma x$. It follows that studying local dynamics of f near x is equivalent to studying the dynamics of its pullback to $N(X)$.

For $x \in \mathbb{R}^n$ (or $N(X)$) let

$$\Sigma_x = \{ \sigma \in \Gamma : \sigma x = x \}$$

be the *isotropy subgroup* of x .

The homogeneous space Γ/Σ_x is not in general a group, but it has the structure of a smooth manifold and the quotient map $\pi : \gamma \mapsto \gamma\Sigma_x$ is a surjection (see Bredon [1972], p. 302). The map $\gamma\Sigma_x \mapsto \gamma x$ is a diffeomorphism between Γ/Σ_x and the orbit Γx of x (see Bredon [1972], VI, 1.2). Fix $x \in \mathbb{R}^n$ and let $\Sigma = \Sigma_x$. Suppose that there exists a neighborhood U of $e\Sigma$ in Γ/Σ and a map $\sigma : U \rightarrow \Gamma$ such that $\pi\sigma(u) = u$ for all $u \in U$. The map σ is called a *local cross section* of π (for a more general definition see Bredon [1972], p. 39). We now construct a local cross section of π . Let Δ be a submanifold of Γ transverse to Σ at e with $e \in \Sigma$ and $\dim \Sigma + \dim \Delta = \dim \Gamma$. Note that a neighborhood \hat{U} of e in Δ is diffeomorphic to a neighborhood of $e\Sigma$ in Γ/Σ and this diffeomorphism is given by $\pi|_{\hat{U}}$. Let $\sigma = (\pi|_{\hat{U}})^{-1}$. Clearly the map σ is a local cross section of π .

Let σ be a cross section of π defined on a neighborhood U of $e\Sigma$ in Γ/Σ . A simple argument shows that the map $\phi : \sigma(U) \times N_x \rightarrow \mathbb{R}^n$ defined as $\phi(u,y) = \sigma(u)y$ is a local diffeomorphism. Let N_x^ε be a disc of radius ε around x and let ε be chosen so that $\phi : \sigma(U) \times N_x^\varepsilon$ is a diffeomorphism. Let $V^\varepsilon = \Gamma N_x^\varepsilon$. Orthogonality of the action implies that N_x and N_x^ε are Σ_x -invariant. Let $X = \Gamma x$. We chose ε so that the set V^ε is equivariantly diffeomorphic to a neighborhood of the zero section in $N(X)$. Observe that if $y \in \hat{N}_x$ then it is clear that $\Sigma_y \subset \Sigma_x$. Hence if $y \in N_x^\varepsilon$ then $\Sigma_y \subset \Sigma_x$. We have the following proposition:

Proposition 1.1

Every smooth and Σ_x -equivariant vector field $g : N_x^\varepsilon \rightarrow \mathbb{R}^n$ has a unique smooth and Γ -equivariant extension $f : V^\varepsilon \rightarrow \mathbb{R}^n$.

Proof

We define f by requiring that $f(\gamma y) = \gamma g(y)$ for $\gamma \in \Gamma$, $y \in N_x^\varepsilon$. To see that f is well defined let $\gamma_1 y = \gamma_2 y$, $y \in N_x^\varepsilon$, $\gamma_1, \gamma_2 \in \Gamma$. Then $\gamma_1^{-1} \gamma_2 y = y$, so $\gamma_1^{-1} \gamma_2 \in \Sigma_y \subset \Sigma_x$ and $g(\gamma_1^{-1} \gamma_2 y) = g(y)$, implying $\gamma_1 g(y) = \gamma_2 g(y)$. Hence f is well defined.

Let U , σ and ϕ be as defined prior to the statement of Proposition 1.1. Let $\hat{U} = \phi(U)$. Then $f|_{\hat{U}} = \phi \circ \text{id} \times h \circ \phi^{-1}$. It follows that f is smooth on \hat{U} . Smoothness of f on V^ε follows from equivariance and smoothness of the action.

2. Dynamics near relative equilibria.

Let Γ be a Lie subgroup of $O(n)$ acting orthogonally on \mathbb{R}^n and let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a C^∞ smooth Γ -equivariant vector field. Fix x_0 in \mathbb{R}^n and let X denote the group orbit of x_0 . We say that the set X is a *relative equilibrium* of f if X is invariant under the flow of f . The subject of this work is studying bifurcations of relative equilibria. In this section we develop a systematic way of analyzing dynamics near a relative equilibrium X . We first describe our results, deferring the proofs to the end of the section.

We begin by defining the concepts of a tangent vector field and a normal vector field. Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$. We say that g is a *tangent vector field* if $g(u)$ is tangent to the group orbit of x for all x in \mathbb{R}^n . For x in X let N_x be the space of vectors normal to X at x . We say that g is a *normal vector field* if for every x in X the space N_x is invariant under the flow of g . Note that a normal vector field does not have to be normal to group orbits other than X .

This section contains two main theorems. The first theorem states that near the group orbit X the vector field f can be written as a sum of a smooth Γ -equivariant normal vector field f_N and a smooth Γ -equivariant tangent vector field f_T . We refer to this theorem as the decomposition theorem. The second theorem states that near X the dynamics of f can be described as dynamics of f_N modulated by drifts along group orbits.

The decomposition theorem is a consequence of a technical lemma. The lemma states that near X there exists a smooth, Γ -invariant bundle K whose fibers are tangent to group orbits and whose restriction to X is the tangent bundle of X . Before stating the lemma we discuss the concept of the normal bundle of X . We observe that a Γ -invariant neighborhood of X can be identified with a neighborhood of the zero section in the normal bundle of X . We conclude that the dynamics of f can be understood in terms of the dynamics of the pullback of f to the normal bundle. In the proof of the decomposition theorem we identify f with its pullback to the normal bundle.

Following the proof of the decomposition theorem we discuss the most important implications of the two main theorems. We remark that Γ -equivariance of f_N implies that the dynamics of f_N is completely determined by its dynamics on the invariant set N_{x_0} . We also describe a way of finding a global center manifold near a relative equilibrium X . When X is an orbit of equilibria we show that the global center manifold is the union of local center manifolds constructed for each normal space N_x .

Next we discuss we discuss a method of explicit computation of f_N . We present the general form of "coordinates along group orbits" and "in the direction normal to the

orbit". Such coordinates have been used to study specific examples of bifurcations of relative equilibria.

We begin by assuming that $X = \Gamma x_0$ is the group orbit of x_0 , but not necessarily a relative equilibrium of f . The following theorem is the first of the two main results of this section:

Theorem 2.1

There exists a Γ -invariant neighborhood U of X in \mathbb{R}^n , a smooth and Γ -equivariant normal vector field f_N and a smooth and Γ -equivariant tangent vector field f_T such that

$$f(u) = f_T(u) + f_N(u)$$

for all u in U .

Let g denote the restriction of f_N to the space N_{x_0} . Let U be the neighborhood defined in Theorem 2.1 and suppose that $u(t)$ is a trajectory of f contained in U for all $t \geq 0$. We now state the second of the two main theorems of this section:

Theorem 2.2

There exists a smooth curve of group elements $\gamma(t)$ and a trajectory $y(t)$ of the vector field g such that $\gamma(t)y(t) = u(t)$ for all $t \geq 0$.

Let $\Pi : N(X) \rightarrow X$ be the bundle projection, that is $\Pi((x, v)) = x$. The following lemma is the main technical result necessary to prove Theorem 2.1.

Lemma 2.3

There exists a smooth Γ -invariant subbundle K of $TN(X)$ such that for all $y \in N(X)$

- (i) $K_y \subset T_y\Gamma y$
- (ii) $K_y \oplus N_{\Pi(y)} = \mathbb{R}^n$.

Note that we cannot define K_y as $T_y\Gamma y$ since the dimension of group orbits may increase near x_0 (the fact that it cannot decrease is a consequence of the inclusion $\Sigma_y \subset \Sigma_{\Pi(y)}$ for $y \in \hat{N}_{\Pi(y)}$). In fact proving Lemma 2.3 is the main technical difficulty of this section. We defer the proof to the end of the chapter. The proof of Theorem 2.2 is also deferred, since it relies on the proof of Lemma 2.3. The proof of Theorem 2.1 is a simple

consequence of Lemma 2.3. In the proof we assume that f is a vector field on $N(X)$, that is $f : N(X) \rightarrow TN(X)$.

Proof of Theorem 2.1

Suppose $y \in N(X)$. Let $P : TN(X) \rightarrow TN(X)$ be defined by $P(y,v) = P_y v$, where P_y is a projection with $\ker P_y = K_y$ and $\text{Im } P_y = N_{\Pi(y)}$. The map P is smooth since the spaces $\ker P_y$ and $\text{Im } P_y$ vary smoothly with y . Equivariance of P follows from invariance of K and equivariance of Π . Let $f_N(y) = P(y, f(y))$, $f_T(y) = f(y) - f_N(y)$. As required f_N is a normal vector field, f_T is a tangent vector field and they are both smooth and Γ -equivariant.

We now discuss some implications of the two main theorems. Recall that orthogonality of the action implies that N_{x_0} is Σ_{x_0} -invariant. Let g be the vector field defined following the statement of Theorem 2.1, that is, g is the restriction of f_N to the space N_{x_0} . Since f_N is Γ -equivariant it follows that g is Σ_{x_0} -equivariant. Let $k = \text{codim } X$. Note that $\dim N_{x_0} = k$. It follows that in order to understand the dynamics of f near X one needs to carry out two steps:

- (a) Analyze the dynamics of the k -dimensional Σ_{x_0} -equivariant vector field g .
- (b) Find the drift along group orbits $\gamma(t)$.

Suppose that X is a relative equilibrium; in this case every x in X is an equilibrium of f_N . In particular, the point x_0 is an equilibrium of g . Let m be a positive integer. The equivariant center manifold theorem (cf. Ruelle [1973], Theorem 1.2) implies that near x_0 the vector field g has a C^m smooth Σ_{x_0} -invariant center manifold. Let M_{x_0} denote such a center manifold. Let $M = \Gamma M_{x_0}$. The smoothness of the action and Σ_{x_0} -invariance of M_{x_0} together imply that M is C^m smooth. Theorem 2.2 also implies that all trajectories of f contained in a sufficiently small neighborhood of X approach M as time goes to infinity. We say that M is the *center manifold* of the relative equilibrium X for the vector field f .

When X consists of equilibria it is natural to ask whether the global center manifold M is a local center manifold for every element of X . We answer this question in the affirmative by verifying that for all x in X the tangent space to M at x equals the center subspace of $(df)_x$. Let E^c be the restriction of the tangent bundle of M to the

relative equilibrium X . The bundle E^c is called the *center bundle* of X . We have the following proposition:

Proposition 2.4

Let x be in X . The fiber of the center bundle E^c at x is the center subspace of $(df)_x$.

Proof

We first prove that $T_x X$ is contained in $\ker (df)_x$. Any vector $u \in T_x X$ can be written as $\frac{d}{ds} \gamma(s)x \big|_{s=0}$. We use the chain rule and the fact that $f(\gamma(s)x) = 0$ to obtain

$$(df)_x u = (df)_x \left(\frac{d}{ds} \gamma(s)x \big|_{s=0} \right) = \frac{d}{ds} f(\gamma(s)x) \big|_{s=0} = 0.$$

Hence 0 is an eigenvalue of $(df)_x$ with multiplicity greater than or equal $\dim X$. Let $v \in N_x$. Theorem 2.1 implies that $(df)_x v = (df_N)_x v + (df_T)_x v$. We show that $(df_N)_x v \in N_x$ and $(df_T)_x v \in T_x X$. This implies that $(df)_x$ can be written in the form:

$$(df)_x = \begin{pmatrix} 0 & (df_T)_x \\ 0 & (df_N)_x \end{pmatrix} \quad (2.1)$$

The proposition follows from equation (2.1), since (2.1) implies that all nonzero eigenvalues of $(df)_x$ are also eigenvalues of $(df_N)_x$.

We now prove that (2.1) is valid. The vector field f_N is Γ -equivariant and $f_N(x) = 0$. The argument presented at the beginning of this proof implies that $T_x X \subset \ker (df_N)_x$. Since $f = f_N + f_T$ it follows that $T_x X \subset \ker (df_T)_x$. Recall from the proof of Theorem 2.1 that $f_N(y) = P_y f(y)$, where P_y is a projection with $\ker(P_y) = T_y \Gamma y$ and $\text{Im}(P_y) = N_{\Pi(y)}$. Hence by the chain rule

$$(df_N)_x = (d_y P_y f(x))_x + P_x (df)_x. \quad (2.2)$$

But $f(x) = 0$, so

$$(df_N)_x = P_x (df)_x. \quad (2.3)$$

It follows that $(df_T)_x(N_x) \subset T_x X$ and $(df_N)_x(N_x) \subset N_x$. Equation (2.1) now follows.

In many applications a bifurcation of a group orbit of equilibria X occurs when $(df)_{x_0}$ maps a vector $v \in N_{x_0}$ to $T_{x_0}X$. The vector v then becomes a generalized nullvector of $(df)_{x_0}$. We have the following proposition:

Proposition 2.5

The vector v is a nullvector of $(df_N)_{x_0}$.

Proof

Proposition 2.5 follows from identity (2.3).

In applications one needs to explicitly compute the vector field f_N . This can be done by changing variables to coordinates in the normal space N_{x_0} and a complementary set of coordinates "along group orbits". Such coordinates have been used by Chossat [1986], Iooss [1986], Danglemayr [1986] and others to study bifurcations of relative equilibria. In the form presented here they were suggested by Pascal Chossat and can be found in Moutrane [1988]. In our presentation we assume that f is a vector field on the normal bundle $N(X)$, that is $f : N(X) \rightarrow TN(X)$. For a Lie group Δ let $L(\Delta)$ denote the Lie algebra of Δ . Let $\exp : L(\Gamma) \rightarrow \Gamma$ be the exponential mapping. Let $V \subset L(\Gamma)$ be the orthogonal complement of $L(\Sigma_{x_0})$ in $L(\Gamma)$ (the space V will be defined more precisely in the proof of Lemma 2.3). Let $\theta : V \times \hat{N}_{x_0} \rightarrow N(X)$ be given by $\theta(\xi, y) = (\exp \xi)(y)$. The linear map $(d\theta)_{(0, x_0)}$ is an isomorphism and hence θ is a local diffeomorphism. Let $h = \theta^*f$. Note that for every $y \in N(X)$ the fiber of the tangent bundle $T_y N(X)$ can be written as $T_y N(X) = V \oplus N_{\Pi(y)}$. The vector field h is defined on $V \times \hat{N}_{x_0}$ and has the following property:

Proposition 2.6

If h is written in the form $h = (h_1, h_2)$, with $h_1 \in V$ and $h_2 \in N_{x_0}$ then $h_2(0, y) = f_N(y)$ for all $y \in \hat{N}_{x_0}$.

The proof of Proposition 2.6 relies on the proof of Lemma 2.3 and therefore will be given at the end of the section.

Proof of Lemma 2.3

Let $L(\Gamma)$ denote the Lie algebra of Γ and let $\exp : L(\Gamma) \rightarrow \Gamma$ be the exponential

mapping. We begin by recalling two concepts related to the Lie algebra $L(\Gamma)$. The *action* of $L(\Gamma)$ on $N(X)$ is defined by

$$\xi y = \frac{d}{dt} (\text{expt} \xi)y \Big|_{t=0} \quad \text{for } \xi \in L(\Gamma), y \in N(X).$$

The *adjoint action* of Γ on $L(\Gamma)$ is defined by

$$\text{Ad}_\gamma \xi = \frac{d}{dt} \gamma(\text{expt} \xi)\gamma^{-1} \Big|_{t=0} \quad \text{for } \gamma \in \Gamma, \xi \in L(\Gamma).$$

Note that

$$\gamma \xi y = \text{Ad}_\gamma \xi \gamma y \quad \text{for } \gamma \in \Gamma, \xi \in L(\Gamma), y \in N(X). \quad (2.4)$$

Recall that $k = \text{codim } X$. We prove that finding the bundle K is equivalent to finding a bundle E over $N(X)$ whose fibers are k dimensional subspaces of $L(\Gamma)$ having the following property:

$$E_{\gamma y} = \text{Ad}_\gamma E_y \quad \text{for } \gamma \in \Gamma. \quad (2.5)$$

Suppose that the bundle E has been found. Then we define the fiber K_y of the bundle K as the set of all images of y under the action of elements of E_y , that is

$$K_y = \{ \xi y : \xi \in E_y \}.$$

If $v \in K_y$ then (2.4) implies that $\gamma v \in K_{\gamma y}$. Hence K is Γ -invariant. We now prove that K is a smooth bundle. For $y \in N(X)$ let $\Phi_y : \Gamma \rightarrow N(X)$ be defined by $\Phi_y(\gamma) = \gamma y$, $\gamma \in \Gamma$. For $\xi \in L(\Gamma)$ we have

$$\xi y = d\Phi_y(e)\xi.$$

It follows that $K_y = d\Phi_y(e)E_y$. Smoothness of K now follows from smoothness of E and from smooth dependence of Φ_y on y . The definition of the action of $L(\Gamma)$ implies that the fibers K_y are tangent to group orbits.

It remains to prove the existence of the bundle E . Suppose $\langle \cdot, \cdot \rangle$ is an inner product on $L(\Gamma)$ invariant with respect to the adjoint action. Such an inner product always

exists for a finite dimensional action of a compact Lie group (see for example Golubitsky, Stewart and Shaeffer [1988], Proposition XI, 1.3). Let V be the orthogonal complement of $L(\Sigma_{x_0})$ taken with respect to $\langle\langle \cdot, \cdot \rangle\rangle$, that is $V = L(\Sigma_{x_0})^\perp$. Let E be the bundle over $N(X)$ with $E_y = \text{Ad}_\gamma V$, $y \in \gamma \hat{N}_{x_0}$. To see that E is well defined suppose that $\gamma_1 y = \gamma_2 y$, $y \in \hat{N}_{x_0}$. From the properties of the normal bundle it follows that $\gamma_1 \gamma_2^{-1} \in \Sigma_{x_0}$. It follows that $\text{Ad}_{\gamma_1 \gamma_2^{-1}} V = V$ or $\text{Ad}_{\gamma_1} V = \text{Ad}_{\gamma_2} V$, implying that $E_{\gamma_1 y} = E_{\gamma_2 y}$. Also E_y is defined for all $y \in N(X)$ since $N(X) = \bigcup_{\gamma \in \Gamma} \gamma \hat{N}_{x_0}$. Equation (2.4) is automatically satisfied for the fibers of E .

For smoothness of E let U be a neighborhood of $e\Sigma_{x_0}$ in Γ/Σ_{x_0} and let $\sigma : U \rightarrow \Gamma$ be a local cross section of π . Let $\phi : U \times N_{x_0} \rightarrow N(X)$ be given as $\phi(u, y) = \sigma(u)(x_0, y)$. The map ϕ is a local diffeomorphism near $(e\Sigma_{x_0}, 0)$. It follows that the map $\Psi : U \times N_{x_0} \times V \rightarrow E$ given by $\Psi(u, y, \xi) = (\sigma(u)y, \text{Ad}_{\sigma(u)}\xi)$ is a local bundle diffeomorphism. This shows smoothness of E near $(x_0, 0)$. To show smoothness near $(\gamma x_0, 0)$ we use the map $\gamma\Psi$ and the relation (2.4).

Proof of Theorem 2.2

Suppose $u(0) = u_0$. Let γ_0 be the element of Γ such that $u_0 \in \gamma_0 N_{x_0}$. Let $y_0 = \gamma_0^{-1} u_0$ and let $y(t)$ be the integral curve of f_N with $y(0) = y_0$. Let $\dot{\cdot}$ denote differentiation with respect to t . To prove the theorem we need to find a curve $\gamma(t)$ with $\gamma(0) = \gamma_0$ and such that

$$(\gamma(t)y(t))' = f(\gamma(t)y(t)). \quad (2.6)$$

The idea of the proof is to reduce (2.6) to an initial value problem on Γ . We observe that the left hand side of (2.6) can be written as

$$\frac{d}{ds} \gamma(t+s)y(t) \Big|_{s=0} + \gamma(t)\dot{y}(t).$$

By assumption $\dot{y}(t) = f_N(y(t))$. It follows that (2.6) can be rewritten as

$$\frac{d}{ds} \gamma(t+s)y(t) \Big|_{s=0} = \gamma(t)f_T(y(t)). \quad (2.7)$$

Let V be the subspace of $L(\Gamma)$ defined in the proof of Lemma 2.3. It follows from the proof of Lemma 2.3 and from the construction of the vector field f_T that there exists a curve $\xi(t)$ of elements of V such that $f_T(y(t)) = \xi(t)y(t)$. Equation (2.7) can now be rewritten as

$$\frac{d}{ds} \gamma(t+s)y(t) |_{s=0} = \gamma(t)\xi(t)y(t). \quad (2.8)$$

Consider the initial value problem:

$$\begin{aligned} \dot{\gamma} &= \gamma(t)\xi(t) \\ \gamma(0) &= \gamma_0. \end{aligned} \quad (2.9)$$

By standard theory of ordinary differential equations (2.9) has a unique solution $\gamma(t)$. It is clear that if $\gamma(t)$ is a solution of (2.9) then $\gamma(t)y(t)$ satisfies (2.8). The theorem now follows.

Proof of Proposition 2.6

Let $y \in \hat{N}_{x_0}$. The proof of Lemma 2.3 and the construction of f_T imply that there exists a unique $\xi_0 \in V$ such that $f_T(y) = \frac{d}{dt} (\text{expt}\xi_0)y |_{t=0}$. Note that $\frac{d}{dt} (\text{expt}\xi_0)y |_{t=0} = (d\theta)_{(0,y)}(\xi_0, 0)$ implying

$$(d\theta)_{(0,y)}(\xi_0, 0) = f_T(y). \quad (2.10a)$$

Also $\theta |_{\hat{N}_{x_0}} = \text{id}$. Hence

$$(d\theta)_{(0,y)}(0, f_N(y)) = f_N(y). \quad (2.10b)$$

Combining (2.10a) and (2.10b) we obtain $(d\theta)_{(0,y)}(\xi_0, f_N(y)) = f_N(y) + f_T(y) = f(y)$. Also, since $y \in \hat{N}_{x_0}$ we have $\theta^*f(y) = (d\theta)_{(0,y)}^{-1}f(y)$. Hence $h_2(0, y) = f_N(y)$.

3. Bifurcations of the normal vector field.

Let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ be a family of vector fields and assume that $F(x_0) \in T_{x_0}X$, that is, X is a relative equilibrium. The results of Section 2 imply that the dynamics of F can be described as follows: the trajectory of F with initial condition y_0 is contained in the group orbit of the trajectory of F_N with the same initial condition. We will utilize this property of the dynamics and divide the bifurcation analysis into two steps. The first step will be to analyze bifurcations of the normal vector field. Then, given a bifurcating solution of the normal vector field, say $y(t)$, we will study the dynamics of F on the set $Y = \{ \gamma y(t) : \gamma \in \Gamma, t \in \mathbb{R} \}$. Note that Y is Γ -invariant and, by Theorem 2.2, it is invariant under the flow of F . This program will be carried out for two kinds of trajectories of F_N – equilibria and periodic orbits.

In this section we discuss the first part of the bifurcation analysis, that is bifurcations of the normal vector field. Suppose that $\dim N_{x_0} = k$. We prove that generic bifurcations of F_N can be described in terms of generic bifurcations of Σ_{x_0} -equivariant vector fields on \mathbb{R}^k . More specifically, we show that a property generic in the class of smooth, Σ_{x_0} -equivariant vector fields on \mathbb{R}^k is also generic in the class of normal vector fields on N_{x_0} . Let $G(\cdot, \lambda)$ be the restriction of the vector field $F_N(\cdot, \lambda)$ to N_{x_0} . Let $g = G(\cdot, 0)$. Suppose that $(dg)_{x_0}$ has an eigenvalue on the imaginary axis and let E be the center subspace of $(dg)_{x_0}$. Suppose that G has a steady state bifurcation, that is $(dg)_{x_0}$ has a 0 eigenvalue. Then we have the following proposition:

Proposition 3.1

Generically the space E equals the nullspace of $(dg)_{x_0}$ and the action of Σ_{x_0} on E is irreducible.

Proposition 3.1 follows from Proposition 1.1 and standard results in equivariant bifurcation theory (see for example Golubitsky, Stewart and Schaeffer [1988], Proposition XII, 3.4).

Suppose that W is a subspace of \mathbb{R}^k . We say that the action of Σ_{x_0} on W is *Γ -simple* if it is irreducible but not absolutely irreducible or if there exists a space V such that $W = V \oplus V$ and the action of Σ_{x_0} on V is absolutely irreducible.

Suppose now that G has a Hopf bifurcation, that is $(dg)_{x_0}$ has a purely imaginary eigenvalue $i\omega$. The following proposition gives a characterization of the space E :

Proposition 3.2

Generically the space E is the generalized eigenspace of $i\omega$ for $(dg)_{x_0}$ and the action of Σ_{x_0} on E is Γ -simple.

Proposition 3.2 follows from Proposition 1.1 and standard results in bifurcation theory (see for example Golubitsky, Stewart and Schaeffer [1988], Proposition XVI, 1.4).

A center manifold reduction coupled with a change of coordinates allows us to reduce the original bifurcation problem for G to a bifurcation problem posed on $E \times \mathbb{R}$. We divide the analysis to two cases:

- (i) The action of Σ_{x_0} on E is trivial.
- (ii) The action of Σ_{x_0} on E is nontrivial.

The case (i) is much simpler and can be analyzed simultaneously for all groups Γ . In particular no symmetry breaking takes place. The following proposition summarizes the bifurcation analysis for this case:

Proposition 3.3

Suppose that the action of Σ_{x_0} on E is trivial. If $(dg)_{x_0}$ has a 0 eigenvalue then generically G has a limit point bifurcation. If $(dg)_{x_0}$ has a purely imaginary eigenvalue $i\omega$ then generically $i\omega$ is a simple eigenvalue of $(dg)_{x_0}$ and G has a Hopf bifurcation to a unique periodic solution.

The proof of Proposition 3.3 follows from standard results in bifurcation theory. In the remainder of this work, unless otherwise stated, we will assume that the action of Σ_{x_0} on E is nontrivial.

Let $y = y(\lambda)$ be a branch of equilibria of G and $Y = Y(\lambda)$ a branch of periodic orbits of G . Let Σ be the isotropy subgroup of y and Σ_Y the group of symmetries mapping Y into itself. In Sections 4 and 5 we show that the trajectories of F on Γy are dense in tori whose dimension is bounded by $\text{rank}(N(\Sigma)/\Sigma)$ (the dimension of a maximal torus in $N(\Sigma)/\Sigma$) and the trajectories on ΓY are dense in tori whose dimension is bounded by $\text{rank}(N(\Sigma_Y)/\Sigma_Y) + 1$. Generically these tori are of maximal dimension. In the context of Proposition 3.3 this maximal dimension equals $\text{rank}(N(\Sigma_{x_0})/\Sigma_{x_0})$ for trajectories on Γy and $\text{rank}(N(\Sigma_{x_0})/\Sigma_{x_0}) + 1$ for trajectories on ΓY .

4. Steady state bifurcations.

Let x_0 be in \mathbb{R}^n and let $X = \Gamma x_0$. Suppose that $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ is a smooth family of equivariant vector fields and X is a relative equilibrium of F for all values of λ . Theorem 2.1 guarantees that F can be decomposed as $F = F_N + F_T$, where F_N is a family of normal vector fields and F_T is a family of tangent vector fields. Let $G(\cdot, \lambda)$ denote the restriction of $F_N(\cdot, \lambda)$ to the normal space N_{x_0} . Note that x_0 is an equilibrium of G for all values of λ . We call x_0 the *trivial equilibrium* of G . We say that the family F has a *steady state bifurcation* near X , if there exists a branch of nontrivial equilibria of G emanating from x_0 . Note that such a bifurcation will generically occur if $(dG)_{(x_0, 0)}$ has a zero eigenvalue and the action of the isotropy subgroup Σ_{x_0} on the center subspace of $(dG)_{(x_0, 0)}$ is nontrivial.

Suppose that F has a steady state bifurcation. Let $y(\lambda)$, $0 \leq \lambda < \lambda_0$, be a bifurcating branch of nontrivial equilibria of G . We assume that all the equilibria $y(\lambda)$ have the same isotropy subgroup Σ . We also assume that the map $\lambda \mapsto y(\lambda)$ is smooth on the open interval $(0, \lambda_0)$. Let $Y(\lambda)$ denote the group orbits of the equilibria $y(\lambda)$. Theorem 2.2 guarantees that the sets $Y(\lambda)$ are invariant under the flow of F . The goal of this section is to analyze that flow of F on the sets $Y(\lambda)$.

Let $z(\epsilon, t)$ be the trajectory of F with initial condition $y(\lambda)$. Equivariance of F implies that each trajectory on $Y(\epsilon)$ is given as $\gamma z(\lambda, t)$, for some $\gamma \in \Gamma$. Hence, to understand the dynamics on Y it suffices to analyze the structure of $z(\lambda, t)$. Let $N(\Sigma)$ denote the normalizer of Σ . Our analysis is based on the following observations:

- (a) The trajectory $z(\lambda, t)$ is contained in $N(\Sigma)y(\lambda)$.
- (b) There exists an integer $k \geq 0$ such that $z(\lambda, t)$ can be described as k -frequency drift along the group orbit Y . More precisely there exists a k -torus $\mathbb{T} \subset \Gamma$ such that $z(\epsilon, t)$ is dense in $\mathbb{T}y$.

Field [1980] (Proposition B1) proved that the number of independent frequencies of the drift is bounded by the dimension of a maximal torus in $N(\Sigma)/\Sigma$. The result of Field can be easily deduced from properties (a) and (b).

We now state the main result of this chapter:

Theorem 4.1

For a generic family F the dimension of the drift along the orbit $Y(\lambda)$ equals the dimension of a maximal torus in $N(\Sigma)/\Sigma$ for all except countably many values of λ .

Theorem 4.1 is an extension of Proposition B1 in Field [1980]. Dancer [1980] also obtained results relevant to the problem discussed in this section. Suppose that f is a smooth Γ -equivariant vector field and y is an equilibrium of f . Let Σ be the isotropy subgroup of y . In the Proposition on p. 88 Dancer proved that if $\dim N(\Sigma) > \dim \Sigma$ then generically all equilibria of f which are sufficiently near y lie in the group orbit of y . Property (b) is a generalization of this result.

In the latter part of this section we state a more precise version of Theorem 4.1. In order to do this we need to review some concepts and results from Lie group theory.

Before we can prove Theorem 4.1 we need to analyze the flow on a relative equilibrium of a single vector field f . Let Y be a relative equilibrium of f and suppose that Σ is the isotropy subgroup of some $y \in Y$. We prove that a trajectory on Y is dense in a k -dimensional torus and that generically k equals the dimension of a maximal torus in $N(\Sigma)/\Sigma$.

In Proposition 4.6 we prove an important technical result stating that the set of all images of a point y in R^n under a smooth Γ -equivariant map is $\text{Fix}(\Sigma_y)$. This result is stated without proof in Field [1980], Lemma A.

Proposition 4.10, which is stated following the proof of Theorem 4.1' describes what happens when the drift fails to be of maximal dimension. The proposition asserts that for a generic family the dimension of the drift can only decrease by one. Field [1989] proves that if the dimension of a maximal torus in $N(\Sigma)/\Sigma$ equals one then for a generic family F the set $Y(\lambda)$ contains no equilibria. This result does not follow from Theorem 4.1.

In order to state a more precise version of Theorem 4.1 we need to review the concepts of maximal tori and rank of a Lie group. Let Δ be a Lie group. We say that a Lie subgroup \mathbb{T} of Δ is a *torus* if \mathbb{T} is compact, abelian and connected. A torus is called *maximal* if it is not properly contained in any other torus. The following is the main result on maximal tori:

Theorem 4.2

In a Lie group Δ any two maximal tori are conjugate, and every element of Δ is contained in a maximal torus.

The proof of Theorem 4.2 can be found in Bröcker and tom Dieck [1985], Theorem (1.6), pg. 159.

Theorem 4.2 implies that all maximal tori are of the same dimension. The dimension of maximal tori in Δ is called the *rank* of Δ .

Let $l = \text{rank } \Delta$ and let $\xi \in L(\Delta)$. We say that ξ *generates a maximal torus* in Δ if the set $\{ \exp t\xi : t \in \mathbb{R} \}$ is dense in a torus of dimension l . We have the following proposition:

Proposition 4.3

The set of $\xi \in L(\Delta)$ which generate a maximal torus is residual (an intersection of open and dense sets).

Proof

Let \mathbb{T} be a maximal torus in Δ . We identify \mathbb{T} with $\mathbb{R}^l / \mathbb{Z}^l$ and $L(\mathbb{T})$ with \mathbb{R}^l (see Bröcker and tom Dieck [1985]), Corollary I, (3.7)). Let

$$P^m = \{ \xi \in L(\mathbb{T}) : \xi = (\xi_1, \xi_2, \dots, \xi_l) \text{ and } \sum m_j \xi_j = 0 \}$$

and let

$$E^m = \bigcup_{\sigma \in \Delta} \text{Ad}_\sigma P^m.$$

Since the group Δ is compact it follows that the image of $L(\Delta)$ under the exponential mapping is the connected component of the identity in Δ (see Bröcker and tom Dieck [1985] Theorem IV, (2.2)). Hence, by Theorem 4.2, each $\zeta \in L(\Delta)$ has the form $\text{Ad}_\sigma \xi$, $\xi \in L(\mathbb{T})$. Since $\exp \text{Ad}_\sigma \xi = \sigma \exp \xi \sigma^{-1}$ it follows that ζ generates a maximal torus if and only if ξ does. Also ξ generates a maximal torus if it is in the complement of the sets E^m for all $m \in \mathbb{Z}^l$. The sets E^m are nowhere dense (see proof of Theorem 4.1'), so the complement of their union is a residual set.

Throughout we assume the following conditions on the family of vector fields F :

- (S1) The orbit X is a trivial relative equilibrium of F . In other words $F_N(\lambda, x_0) = 0$ for all values of λ .
- (S2) There exists $\lambda_0 > 0$ and a branch of relative equilibria of F_N , parametrized as $y(\lambda)$, $0 < \lambda < \lambda_0$. The mapping $\lambda \mapsto y(\lambda)$ is smooth on $(0, \lambda_0)$. The points $y(\lambda)$ have isotropy Σ .

Let $C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ denote the space of smooth families of equivariant tangent vector fields on \mathbb{R}^n . For a family F satisfying (S1) and (S2) let $Y(\lambda) = \Gamma y(\lambda)$. We now state Theorem 4.1 more precisely.

Theorem 4.1'

Suppose that a family of vector fields F satisfies (S1) and (S2). Then

- (i) Trajectories on the manifolds $Y(\lambda)$ are dense in tori of dimension bounded by $\text{rank}(N(\Sigma)/\Sigma)$.
- (ii) There exists a residual set $B \subset C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ such that for every $H \in B$ there exists a countable set $I_0 \subset (0, \lambda_0)$ such that for every $\lambda \in (0, \lambda_0) \setminus I_0$ trajectories of $F + H$ on the manifolds $Y(\lambda)$ are dense in tori of dimension equal to $\text{rank}(N(\Sigma)/\Sigma)$.

Note that Theorem 4.1' is more general than Theorem 4.1: we assume that $y(\lambda)$ is a branch of relative equilibria of F_N rather than a branch of equilibria of F_N . This assumption does not increase the complexity of the proof.

Before proving Theorem 4.1' we analyse the following simpler situation. Suppose that $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Γ -equivariant vector field with the following properties:

- (VS1) The orbit X is a relative equilibrium of g .
- (VS2) There exists $y_0 \in \mathbb{R}^n$, $y_0 \notin X$ such that $Y = \Gamma y_0$ is a relative equilibrium of g .

The problem of finding the dynamics on Y has been solved by Field [1980]. Here we briefly present his results. We start with the following proposition:

Proposition 4.4

Suppose that $g(y_0) = v$. Let $\xi \in L(\Gamma)$ be such that $\xi y_0 = v$. Then $y(t) = \exp(t\xi)y_0$ is the integral curve of g with $y(0) = y_0$.

Proof

$$\begin{aligned} \dot{y}(\tau) &= \frac{d}{dt} \exp(t\xi)y_0 \Big|_{t=\tau} \\ &= \exp(\tau\xi) \frac{d}{dt} \exp((t - \tau)\xi)y_0 \Big|_{t=\tau} \\ &= \exp(\tau\xi) \frac{d}{ds} \exp(s\xi)y_0 \Big|_{s=0} \end{aligned}$$

By definition $\frac{d}{ds} \exp(s\xi)y_0 \Big|_{s=0} = \xi y_0$. Hence

$$\begin{aligned} \dot{y}(\tau) &= \exp(\tau\xi) \xi y_0 = \exp(\tau\xi)g(y_0) \\ &= g(\exp(\tau\xi)y_0) = g(y(\tau)). \end{aligned}$$

For $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$ let $\|h\| = \sup_{x \in \mathbb{R}^n} |h(x)|$. The following theorem gives a complete description of dynamics on relative equilibria of a vector field g :

Theorem 4.5 (Field [1980], Proposition B1)

Suppose that $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is an equivariant vector field satisfying (VS1) and (VS2) and let Σ be the isotropy subgroup of y_0 . Then

(i) Every flow trajectory contained in Y has the form $z(t) = \exp(t\xi)y$, for some $\xi \in L(N(\Sigma))$.

(ii) The dimension of the torus $\mathbb{T} = \text{cl} \{ \exp(t\xi)y_0 : t \in \mathbb{R} \}$ is less or equal to $\text{rank}(N(\Sigma)/\Sigma)$.

(iii) For every $\varepsilon > 0$ there exists a vector field h , such that $\|h\| < \varepsilon$, Y is a relative equilibrium of h , and the dimension of the closure of trajectories of $g+h$ on Y equals $\text{rank}(N(\Sigma)/\Sigma)$.

To prove of Theorem 4.5 we need to answer the following question: what are the possible images of the vector y_0 under Γ -equivariant vector fields? Suppose that V is the space of all possible images of y under Γ -equivariant vector fields, that is

$$V = \{h(y), h : \mathbb{R}^n \rightarrow \mathbb{R}^n \text{ is a } \Gamma\text{-equivariant vector field}\}.$$

If h is a Γ -equivariant vector field then it follows that $h(y_0)$ is fixed by all elements in Σ_{y_0} . Hence $V \subset \text{Fix}(\Sigma_{y_0})$. The following proposition shows that the other containment also occurs:

Proposition 4.6

The space V is equal to the fixed point space of Σ_{y_0} , that is

$$V = \text{Fix}(\Sigma_{y_0}).$$

Proof

Let $Y = \Gamma y_0$ and let $\Sigma = \Sigma_{y_0}$. Suppose that $v \in \text{Fix}(\Sigma)$. We first show that there exists a smooth and Γ -equivariant vector field $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $g(y_0) = v$. Let $N(Y)$ be the normal bundle of Y . Recall that $N(Y)$ can be identified with an invariant neighborhood U of Y in \mathbb{R}^n . It follows that $TN(Y)$ can be identified with \mathbb{R}^n . We define a vector field $h : N_{y_0} \rightarrow TN(Y)$ by $h(z) = v$. Clearly h is smooth and Σ -equivariant. By Proposition 1.1 we can extend h to a smooth, Γ -equivariant vector field g_1 on $N(Y)$. The properties of the normal bundle $N(Y)$ imply that the vector field g_1 can be identified with a vector field g_2 defined on U . Let $\alpha : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth, invariant function such that $\alpha(y_0) = 1$ and $\alpha(x) = 0$ for all $x \notin U$. Let g be defined as follows:

$$g(x) = \begin{cases} \alpha(x)g_2(x) & \text{if } x \in U \\ 0 & \text{if } x \notin U \end{cases}$$

Clearly g is smooth, Γ -equivariant and $g(y_0) = v$.

Remark 4.7

The vector field h , described in Theorem 4.5 (iii) can be chosen so that $g+h$ is a polynomial vector field. The existence of such h can be shown using the equivariant version of the Stone-Weierstrass approximation theorem (see Poenaru [1976], proof of Proposition 1, p. 20).

The final ingredient necessary to prove Theorem 4.4 is given by the following elementary lemma:

Lemma 4.8

The following equality holds for any $y \in \mathbb{R}^n$:

$$T_y Y \cap \text{Fix}(\Sigma_y) = \{ \xi y : \xi \in L(N(\Sigma_y)) \}.$$

Proof

Let $\xi \in L(\Gamma)$, $y \in \mathbb{R}^n$. Then $\xi y \in \text{Fix}(\Sigma_y)$ if and only if $\exp \xi y \in \text{Fix}(\Sigma_y)$. This implies that $\sigma \exp \xi y = \exp \xi y$ for all $\sigma \in \Sigma_y$. It follows that there exists $\eta \in L(N(\Sigma))$ such that $\eta y = \xi y$. The lemma now follows.

Proof of Theorem 4.5

The theorem is an easy consequence of Proposition 4.4, Proposition 4.6 and Lemma 4.8.

In the remainder of this section we prove Theorem 4.1'. Let F be a family of vector fields satisfying (S1) and (S2). We begin by defining a map which assigns to each $H \in C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ a curve ξ in $L(N(\Sigma)/\Sigma)$ such that $(F + H)(y(\lambda), \lambda) = \xi(\lambda)y(\lambda)$ for each λ in some interval I . Recall that $T_y Y \cap \text{Fix} \Sigma = \{ \xi y : \xi \in L(N(\Sigma)) \}$. Let V_Σ be the set of $y \in \mathbb{R}^n$ with isotropy group Σ . For $y \in V_\Sigma$ the space $\{ \xi y : \xi \in L(N(\Sigma)) \}$ is isomorphic to $L(N(\Sigma)/\Sigma)$. Let $\Xi : T_y Y \cap \text{Fix} \Sigma \rightarrow L(N(\Sigma)/\Sigma)$ denote this isomorphism. It is clear that Ξ changes smoothly as y is being varied in V_Σ . Let I be a subinterval of $(0, \lambda_0)$. The map $\Theta : C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n) \rightarrow C^\infty(I, L(N(\Sigma)/\Sigma))$ is defined as

$$\Theta(H)(\lambda) = \Xi((F + H)(y(\lambda), \lambda)).$$

We prove Theorem 4.1' by showing that for a residual set of $H \in C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ the curve $\Theta(H)$ is transverse to all the sets E^m (see Proposition 4.3). Before presenting the proof we review some concepts related to the Whitney C^∞ topology. For a more complete treatment of this topic see Golubitsky and Guillemin [1973]. Let Z, W be smooth manifolds. For a positive integer q let $J^q(Z, W)$ denote the space of q -jets of smooth maps from Z to W . We describe a neighborhood basis of a map f in the Whitney C^∞ topology on $C^\infty(Z, W)$. Let q be a positive integer and let d_q be a metric on $J^q(Z, W)$ compatible with its topology (such a metric exists by (I, 5.9) in Golubitsky and Guillemin [1973]). Let $\delta : Z \rightarrow \mathbb{R}^+$ be a continuous function. Let

$$U_{q,\delta} = \{ g \in C^\infty(Z, W) : d_q(j^q f(x), j^q g(x)) < \delta(x) \text{ for all } x \in Z \}.$$

The collection of the sets $U_{q,\delta}$ for all choices of q and δ forms a neighborhood basis of f in the Whitney C^∞ topology.

Suppose that Z is an open subset of \mathbb{R}^p , for some p , and W is a vector space. Then the above mentioned metrics d_q can be chosen as follows. Suppose $s = \dim W$. We identify W with \mathbb{R}^s . For a positive integer q and $g \in J^q(Z, W)$ let

$$\|g\|_q(x) = |x| + |g(x)| + \sum_{1 \leq |\alpha| \leq q} \left| \frac{\partial^{|\alpha|} g(x)}{\partial x^\alpha} \right|.$$

Here α denotes a p -vector of nonnegative integers. We define d_q on $J^q(Z, W)$ as:

$$d_q(\sigma_1, \sigma_2) = \|g_1 - g_2\|_q(x),$$

where $\sigma_1, \sigma_2 \in J^q(Z, W)$ and g_1, g_2 are such that $\sigma_1 = j^q g_1(x)$ and $\sigma_2 = j^q g_2(x)$. It is easy to see that d_q agrees with the topology on $J^q(Z, W)$.

Let I be the interval used in the definition of the map Θ . Then

Lemma 4.9

If $\bar{I} \subset (0, \lambda_0)$ then Θ is continuous in the Whitney C^∞ topology.

Proof

Let $C(I) = \{ (y(\lambda), \lambda) : \lambda \in I \}$. In this proof we use the metrics d_q described prior to the statement of Lemma 4.9. The map Θ can be written as

$$\Theta(H) = \Xi \circ H | C(I).$$

Hence Θ is a composition of two maps: a map Θ_1 given as

$$\Theta(H) = H | C(I)$$

and a map Θ_2 defined as

$$\hat{H} \rightarrow \Xi \circ \hat{H}.$$

The map Θ_1 does not, in general, have to be continuous, but it is continuous if $\bar{I} \subset (0, \lambda_0)$. This follows, since $\bar{I} \subset (0, \lambda_0)$ implies that for any given q all the partial derivatives of the function $\lambda \mapsto y(\lambda)$ are bounded on I . Hence continuity of Θ_1 can be established through repeated application of the chain rule. The map Θ_2 is continuous by (II, 3.5) in Golubitsky and Guillemin [1973].

Proof of Theorem 4.1'

Part (i) of the theorem follows from Theorem 4.5.

We now prove part (ii). Suppose that l is the rank of $N(\Sigma)/\Sigma$. Let \mathbb{T} be a maximal torus in $N(\Sigma)/\Sigma$. As in the proof of Proposition 4.3 we identify \mathbb{T} with R^l/Z^l and $L(\mathbb{T})$ with R^l . Recall the definitions of the sets E^m and P^m (see proof of Proposition 4.3). In the proof of Proposition 4.3 we show that the sets E^m have the following property: a vector $\xi \in L(N(\Sigma)/\Sigma)$ generates a maximal torus if and only if ξ is in the complement of E^m for all m in Z^l .

The sets E^m may not be manifolds, but we show that each E^m is a finite union of manifolds. Let $\xi \in P^m$ and let Δ be the isotropy subgroup of ξ with respect to the adjoint action of $N(\Sigma)/\Sigma$ on $L(N(\Sigma)/\Sigma)$. Let $[\cdot, \cdot]$ denote the bracket in $L(N(\Sigma)/\Sigma)$. It is known (see Bröcker and tom Dieck [1985], I, (2.12)) that

$$[\eta, \zeta] = \frac{d}{dt} \text{Ad}_{\exp t\eta} \zeta \Big|_{t=0}, \text{ for all } \eta, \zeta \in L(N(\Sigma)/\Sigma). \quad (4.1)$$

The containment $P^m \subset L(\mathbb{T})$ implies that $[\xi, \eta] = 0$ for all $\eta \in P^m$. Let $O(\xi) = \bigcup_{\sigma \in N(\Sigma)/\Sigma} \text{Ad}_{\sigma} \xi$ be the orbit of ξ . Let U be a small neighborhood of $e\Delta$ in $(N(\Sigma)/\Sigma)/\Delta$ and let $\sigma : U \rightarrow N(\Sigma)/\Sigma$ be a local cross section. The equation (4.1) implies that $T_{\xi}O(\xi) \cap P^m = \{0\}$. It follows that the map $\Psi : U \times P^m \rightarrow L(N(\Sigma)/\Sigma)$, given by $\Psi(u, \eta) = \text{Ad}_{\sigma(u)}\eta$ is a local diffeomorphism near (e, ξ) .

Let $E^m(\Delta)$ be the set of all elements of E^m whose isotropy subgroup (with respect to the adjoint action) is conjugate to Δ and let $P^m(\Delta) = P^m \cap E^m(\Delta)$. Note that $P^m(\Delta)$ is an open subset of $P^m \cap \text{Fix}(\Delta)$. For every $\xi \in P^m$ the corresponding map Ψ is a local diffeomorphism near (e, ξ) . It follows that $E^m(\Delta)$ is a smooth manifold.

Note that the number of the sets $E^m(\Delta)$ is finite. This follows from the fact that $N(\Sigma)/\Sigma$, being a compact group, has a finite number of conjugacy classes of isotropy subgroups. Clearly $E^m = \bigcup E^m(\Delta)$.

The theorem follows from the following assertion:

(*) For every $m \in \mathbb{Z}^l$, $\mu \in (0, \lambda_0)$ there exists an interval I containing μ and a set $B^m(I) \subset C_T^{\infty}(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ with the following properties:

- (1) $B^m(I)$ is residual in the C^{∞} Whitney topology.
- (2) $\Theta(H)$ is transverse to all the sets $E^m(\Delta)$ at each $\lambda \in I$.

We first show that the theorem follows from (*). To see this let I_1^m, I_2^m, \dots be a sequence of intervals such that $\bigcup_{i=1}^{\infty} I_i^m = (0, \lambda_0)$ and $B^m(I_i^m)$ satisfies the properties (1) and (2). Let $B = \bigcap_{i=1, j=1}^{\infty} B^j(I_i^j)$. It follows that for every $H \in B$ the curve $\Theta(H)$ is transverse to all the sets E^m at every $\lambda \in (0, \lambda_0)$. It is clear that B satisfies the property required in the statement of Theorem 4.1'.

We now prove (*). Fix $m \in \mathbb{Z}^l$, a subgroup $\Delta \subset N(\Sigma)/\Sigma$ and $\mu \in (0, \lambda_0)$. Let $I \subset I_0 \subset (0, \lambda_0)$ be intervals with $\mu \in I$ and $\bar{I}_0 \subset (0, \lambda_0)$. Let A_0 be the set of $\xi \in C^{\infty}(I_0, L(N(\Sigma)/\Sigma))$ such that ξ is transverse to $E^m(\Delta)$ at each $\lambda \in I$. By standard

transversality arguments (see Golubitsky and Guillemin [1973], (II, 4.5)) the set A_0 is open and dense in the Whitney C^∞ topology. We assume that I_0 is the interval used in the definition of the map Θ . Let $A = \Theta^{-1}A_0$. It follows from Lemma 4.9 that A is an intersection of open sets. We now show that A is dense. Fix $H \in C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$. We construct a sequence of families $\{H_i\}$ converging to H and such that each $H_i \in A$. Let $\{\xi_i\}$ be a sequence of elements of $C^\infty(I_0, L(N(\Sigma)/\Sigma))$ such that each curve $\Theta(H) + \xi_i$ is in A_0 and the curves ξ_i converge to the 0 curve as $i \rightarrow \infty$. Such a sequence exists, since A_0 is dense in $C^\infty(I_0, L(N(\Sigma)/\Sigma))$. To show the existence of the sequence $\{H_i\}$ it suffices to prove that for every $\eta \in C^\infty(I_0, L(N(\Sigma)/\Sigma))$ there exists a family H_η such that $\Theta(H_\eta)(\lambda) = \eta(\lambda)$ for all $\lambda \in I$ and such that for every positive integer q the size of partial derivatives of H_η of order less or equal to q can be estimated by the size of partial derivatives of η of order less or equal to q . We now give a more precise description of this estimate. Let $\hat{\eta}$ be a smooth curve of elements of $L(N(\Sigma))$ which projects to η in $L(N(\Sigma)/\Sigma)$. Suppose that α is an n -vector of positive integers and β a positive integer. Then there exists a constant C , depending only on m , and such that

$$\left| \frac{\partial^{|\alpha|+\beta} H_\eta(z, \lambda)}{\partial z^\alpha \partial \lambda^\beta} \right| \leq C \left| \frac{\partial^\beta \hat{\eta}(\lambda)}{\partial \lambda^\beta} \right| \quad (4.2)$$

for all $(z, \lambda) \in V \times I_0$. Moreover $H_\eta(z, \lambda) = 0$ for $(z, \lambda) \notin V \times I_0$.

We now construct H_η . Let $y_0 = y(\mu)$ and $Y = \Gamma y_0$. Recall that $N(Y)$ is equivariantly diffeomorphic to an invariant neighborhood of Y in \mathbb{R}^n . Let V be such a neighborhood. In the sequel we identify V with $N(Y)$. By shrinking the interval I we can assume that $y(\lambda) \in V$ for all $\lambda \in I$. We can assume that $y(\lambda) \in N_{y_0}$ for all $\lambda \in I$. Otherwise we could replace the curve $y(\lambda)$ by a curve $\hat{y}(\lambda) = \gamma(\lambda)y(\lambda)$ with $\gamma(\lambda) \in N(\Sigma)/\Sigma$. We can now define $\hat{H}(z, \lambda)$ as $\hat{\eta}(\lambda)z$ if $z \in N_{y_0}$ and extend this definition by equivariance (see proof of Proposition 4.6). Let U_0 be a small neighborhood of $e\Sigma$ in Γ/Σ and let σ be a local cross section of π (see Section 1). Let $\phi : U_0 \times N_{y_0}^\varepsilon$ be defined as $\phi(u, y) = \sigma(u)y$ (here $N_{y_0}^\varepsilon$ denotes a disc of radius ε around y_0 in N_{y_0}). Recall that for ε small enough ϕ is a diffeomorphism. Let $U = \phi(U_0 \times N_{y_0}^\varepsilon)$. We can express every point $z \in U_0$ in local coordinates as $\sigma(u)y$, $y \in N_{y_0}$, $u \in U$. Then, for every z

$\in U$, $\hat{H}(z, \lambda) = \sigma(u)\hat{\eta}(\lambda)z$. It is clear from this expression and from smoothness of the action of Γ that (4.2) holds for all $z \in U$. From compactness of Y it follows that the bound (4.2) holds on a neighborhood V_1 of Y in \mathbb{R}^n with possibly a different constant C . With no loss of generality we can assume that $V = V_1$. Let W be an invariant neighborhood of y_0 such that $\bar{W} \subset V$ and suppose that I is chosen so that $y(\lambda) \in W$ for all $\lambda \in I$. Let $h : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ be a smooth Γ -invariant cut-off function vanishing on the complement of $V \times I_0$ and equal to 1 on $U \times I$. Let $H_\eta(z, \lambda) = h(z, \lambda)\hat{H}(z, \lambda)$. It is clear that H_η is globally defined and satisfies (4.2).

We complete the proof of (*) by defining $B^m(I)$ as the intersection of the sets A for all choices of Δ .

For $m_1, m_2 \in Z'$ let $E^{m_1, m_2} = E^{m_1} \cap E^{m_2}$. Note that if m_1 and m_2 are not colinear then the sets E^{m_1, m_2} have codimension 2 in $L(N(\Sigma)/\Sigma)$. The union of these sets consists of the elements $\zeta \in L(N(\Sigma)/\Sigma)$ which generate a torus of dimension no less than $\text{rank}(N(\Sigma)/\Sigma) - 1$. In the proof of Theorem 4.1' we could, instead of the sets E^m , use the sets E^{m_1, m_2} . Then, for every $H \in B$, the curve $\Theta(H)$ would be transverse to all E^{m_1, m_2} at each $\lambda \in (0, \lambda_0)$. This would imply that if m_1 and m_2 were not colinear then E^{m_1, m_2} and $\Theta(H)$ would not intersect. This property implies the following proposition:

Proposition 4.10

Suppose that F satisfies (S1) and (S2). Then there exists a residual set $B \subset C_T^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ such that if $H \in B$ then the dimension of the trajectories of $F+H$ on the sets Y is greater or equal to $\text{rank}(N(\Sigma)/\Sigma) - 1$.

5. Hopf bifurcations.

Let x_0 be in \mathbb{R}^n and let $X = \Gamma x_0$. Suppose that $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ is a smooth family of equivariant vector fields and X is a relative equilibrium of F for all values of λ . By Theorem 2.1 $F = F_N + F_T$, where F_N is a family of normal vector fields and F_T is a family of tangent vector fields. Let G be the family defined at the beginning of Section 4, that is $G(\cdot, \lambda)$ is the restriction of $F_N(\cdot, \lambda)$ to the normal space N_{x_0} . Recall that x_0 is the trivial equilibrium of G . We say that the family F has a *Hopf bifurcation* near X , if there exists a branch of nontrivial periodic orbits of G emanating from x_0 . Note that such a bifurcation will generically occur if $(dG)_{(x_0, 0)}$ has a purely imaginary eigenvalue.

Suppose that F has a Hopf bifurcation. Let $Y(\lambda)$, $\lambda_0 > \lambda > 0$, be a branch of periodic orbits of G and let $y(\lambda)$ denote the initial conditions for the trajectories $Y(\lambda)$. We assume that all the points $y(\lambda)$ have the same isotropy subgroup Σ . Let $\Sigma_{Y(\lambda)}$ be the *group of symmetries* of the set $Y(\lambda)$, that is

$$\Sigma_{Y(\lambda)} = \{ \sigma \in \Sigma_{x_0} : \sigma Y(\lambda) = Y(\lambda) \}.$$

We assume that all the sets $Y(\lambda)$ have the same group of symmetries Σ_Y .

Let $Z(\lambda)$ denote the group orbits of the sets $Y(\lambda)$. Theorem 2.2 guarantees that the sets $Z(\lambda)$ are invariant under the flow of F . The goal of this section is to analyze that flow of F on the sets $Z(\lambda)$. Our analysis is based on the following result: every trajectory of F on Z is dense in a $k+1$ -dimensional torus, with k frequencies given by the drift along group orbits and the additional frequency corresponding to the motion along the periodic orbit Y . This result was obtained by Field [1980], Proposition B2. Field also showed that the number of the drift frequencies is bounded by $\text{rank}(N(\Sigma)/\Sigma)$.

This section contains two main results. The first of these results is a modification of the theorem of Field. We assume that f is a smooth Γ -equivariant vector field, X is a relative equilibrium of f and Y is a periodic orbit of f_N . Let $Z = \Gamma Y$. The theorem states that the trajectories on Z are dense in tori of dimension $k+1$, with $k \leq \text{rank } N(\Sigma_Y)/\Sigma_Y$. For some choices of f $\text{rank}(N(\Sigma_Y)/\Sigma_Y) < \text{rank}(N(\Sigma)/\Sigma)$. This is illustrated in Example 5.3.

Our second main result (Theorem 5.2) deals with the dynamics of the family of vector fields F on the sets $Z(\lambda)$. The theorem states that, given a generic family of vector fields F , there exists a countable set $L_0 \subset (0, \lambda_0)$ such that if $\lambda \notin L_0$ then the trajectories on $Z(\lambda)$ are dense in tori of maximal dimension. In Proposition 5.9 we strengthen this result by showing that generically the dimension of trajectories on $Z(\lambda)$ drops only by 1.

We now state the first of the two main theorems. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a smooth, Γ -equivariant vector field with the following properties:

(VH1) The orbit X is a relative equilibrium of f .

(VH2) The vector field f_N has a periodic orbit $Y = \{ y(t) : t \in [0, T] \}$, T is the period of Y .

Let Σ_Y denote the group of symmetries of Y and let $Z = \Gamma Y$. We have:

Theorem 5.1

All trajectories on the set Z are dense in $k+1$ -dimensional tori, $k \leq \text{rank } N(\Sigma_Y)/\Sigma_Y$. For every $\varepsilon > 0$ there exists a smooth and Γ -equivariant vector field h such that $\|h\| \leq \varepsilon$, h satisfies (VH1) and (VH2) and such that the trajectories of $f+h$ on the set Z are dense in tori of dimension equal to $\text{rank } N(\Sigma_Y)/\Sigma_Y + 1$.

We now state the second main theorem. Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a smooth family of Γ -equivariant vector fields with the following properties:

(H1) The orbit X is a relative equilibrium of F for all values of $\lambda \in \mathbb{R}$.

(H2) There exists $\lambda_0 > 0$ and a branch of periodic orbits of F_N , parametrized as $(\lambda, Y(\lambda))$, $0 < \lambda < \lambda_0$, with initial conditions y_λ . All the elements y_λ have the same isotropy subgroups Σ and all the sets $Y(\lambda)$ have the same group of symmetries Σ_Y . The map $\lambda \mapsto y_\lambda$ is smooth on the interval $(0, \lambda_0)$.

For a family of vector fields satisfying (H1) and (H2) let $Z(\lambda) = \Gamma Y(\lambda)$. We have the following theorem:

Theorem 5.2

If F is a family of vector fields satisfying (H1) and (H2) then there exists a set $B \subset C^\infty(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ with the following properties:

- (i) For every $H \subset B$ there exists a countable set $I_0 \subset (0, \lambda_0)$ such that for every $\lambda \in (0, \lambda_0) \setminus I_0$ trajectories of $F+H$ on the manifolds $Z(\lambda)$ are dense in tori of maximal dimension.
- (ii) The set B is residual in the Whitney C^∞ topology.

We now present an example of a vector field f for which $\text{rank } N(\Sigma)/\Sigma > \text{rank } N(\Sigma_Y)/\Sigma_Y$.

Example 5.3

Let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ be a smooth family of vector fields equivariant under the action of $O(2)$. Let $\kappa \in O(2) \setminus SO(2)$. Suppose that the invariant equilibrium $x = 0$ bifurcates to a branch of equilibria $x(\lambda)$ with $\Sigma_{x(\lambda)} = \{e, \kappa\}$. Consider a secondary Hopf bifurcation occurring along the branch $x(\lambda)$. In other words suppose that $(dF_N)_{(x(\lambda_0), \lambda_0)}$ has, for some λ_0 , a pair of purely imaginary eigenvalues $i\omega$. Let V be the real eigenspace of $i\omega$. We assume that V is two dimensional and that the action of κ on V is nontrivial. Then, by standard Hopf bifurcation theorem, F_N has a branch of periodic solutions with period T and such that $y_\lambda(t + \frac{T}{2}) = \kappa y_\lambda(t)$. By Theorem 2.2 the trajectory of F corresponding to $y_\lambda(t)$ is $z_\lambda(t) = y(t)y_\lambda(t)$, where $y(t) \in SO(2)$ and $y(0) = e$. Let Φ_t denote the flow of F . Then $x_\lambda(t) = \Phi_T y_\lambda(0) = \Phi_T y_\lambda(0) = \Phi_{T/2}(\Phi_{T/2} y_\lambda(0)) = \Phi_{T/2}(y(\frac{T}{2})\kappa y_\lambda(0))$. Equivariance of $\Phi_{T/2}$ implies that $\Phi_{T/2}(y(\frac{T}{2})\kappa y_\lambda(0)) = (y(\frac{T}{2})\kappa)^2 y_\lambda(0) = y_\lambda(0) = x_\lambda(0)$. It follows that $x_\lambda(t)$ must be a periodic solution. Note that $\Sigma = \{e\}$, so $N(\Sigma)/\Sigma = O(2)$, but $\Sigma_Y = \{e, \kappa\}$, so $N(\Sigma_Y)/\Sigma_Y$ is discrete. Hence $\text{rank } N(\Sigma)/\Sigma > \text{rank } N(\Sigma_Y)/\Sigma_Y$.

In the remainder of the section we prove Theorems 5.1 and 5.2. In the proof of Theorem 5.1 we will use two lemmas and some background information from Lie group theory. We begin by stating and proving the first of the lemmas. Note that $\Sigma \subset \Sigma_Y \subset N(\Sigma)$. Let $\Delta = \Sigma_Y/\Sigma$. We have

Lemma 5.3

The group Δ is finite and cyclic or Δ is isomorphic to S^1 .

Proof

Let y_0 be the initial condition of the periodic orbit Y . We assume, with no loss of generality, that the period of the solution $y(t)$ is 1. We identify S^1 with \mathbb{R}/\mathbb{Z} . Note that Y is diffeomorphic to S^1 via the map $t \mapsto y(t)$. Let $p : \Delta \rightarrow S^1$ be defined by the identity

$$y(p(\delta)) = \delta y_0, \quad \delta \in \Delta.$$

Clearly p is smooth and well defined. Note that the action of Δ on Y is free, so p must be injective. Equivariance of f implies that p is a Lie group homomorphism. It follows that $p(\Delta)$ (which is isomorphic to Δ) is a Lie subgroup of S^1 and therefore must be isomorphic to either S^1 or \mathbb{Z}_l for some l .

We now review some of the concepts from Lie group theory, which will be used in the proof of Theorem 5.1. For a Lie group H let H_0 denote the connected component of the identity in H . We say that a subgroup K of a compact Lie group H is *topologically cyclic* if there exists $\delta \in K$ such that $K = \text{cl} \{ \delta^n : n \text{ is an integer} \}$. The element δ is called the *generator* of K . We say that K is a *Cartan subgroup* of H if K is topologically cyclic and $N(K)/K$ is discrete. In the proof of Theorem 5.1 we will use the following two propositions.

Proposition 5.4

Each element $h \in H$ is contained in a Cartan subgroup K of H such that K/K_0 is generated by hK_0 .

Proposition 5.5

If K is a Cartan subgroup of H generated by z , then any $h \in H_0z$ is conjugate to an element of K_0z via conjugation by an element of H_0 .

The statements and the proofs of Propositions 5.4 and 5.5 can be found in Bröcker and tom Dieck [1985], IV, (4.2) and (4.3).
 If K is a Cartan subgroup and K_1 a topologically cyclic group then K/K_1 must be discrete. Proposition 5.4 implies that each topologically cyclic group is contained in a Cartan subgroup. It follows that a Cartan subgroup can be defined as a topologically cyclic group of maximal dimension.

Let K be a topologically cyclic subgroup of H . Then K_0 must be a torus in H_0 and K/K_0 must be finite and cyclic. It follows that K is isomorphic to $K_0 \times Z_l$, for some l (see Bröcker and tom Dieck [1985], I, (4.14)).

Let f be a smooth Γ -equivariant vector field satisfying (VH1) and (VH2). Theorem 2.2 guarantees that the trajectory of f with initial condition $y(0)$ has the form $\gamma(t)y(t)$, where $\gamma(t) \in (N(\Sigma)/\Sigma)_0$. We assume that Δ is finite and cyclic with generator δ_0 (see Lemma 5.3) and let T_0 be defined by the identity $y(T_0) = \delta_0 y_0$. In order to prove Theorem 5.1 we need to prove the following lemma:

Lemma 5.6

Suppose that $\gamma(T_0) = \gamma_0$ and that $\gamma_1 \in (N(\Sigma)/\Sigma)_0$. Then there exist a smooth, Γ -equivariant tangent vector field h and a curve $\gamma^1(t)$ such that:

(i) $\gamma^1(0) = e$.

(ii) $\gamma^1(T_0) = \gamma_1$.

(iii) $\gamma^1(t)y(t)$ is the trajectory of $f+h$.

Proof

Let $\xi(t) = \gamma(t)^{-1}\gamma(t)$. Note that $f_T(y(t)) = \xi(t)y(t)$. Let $\sigma = \gamma_1\gamma_0^{-1}$ and let $\zeta = \exp^{-1}\sigma$. Let $\eta(t)$ be a smooth function such that $\eta(0) = 0$, $\eta(T_0) = 1$ and

$$\frac{d^j}{dt^j} \eta(0) = \frac{d^j}{dt^j} \eta(T_0) = 0, \quad j = 1, 2, \dots$$

Let $\gamma^1(t) = \exp(\eta(t)\zeta)\gamma(t)$. We define a curve $\xi(t) = \xi(t) + \eta(t)\gamma(t)^{-1}\zeta\gamma(t)$. Let $\xi^1(t) = \xi(t) - \xi(t)$. Note that

$$\frac{d^j}{dt^j} \xi^1(0) = \frac{d^j}{dt^j} \xi^1(T_0) = 0, \quad j = 1, 2, \dots \quad (5.1)$$

We extend the definition of ξ^1 to all of R by requiring that $\xi^1(t+T_0) = \text{Ad}\xi^1(t)$, where δ is as defined in Lemma 5.3. The equation (5.1) implies that this extension is smooth. Let $h(y(t)) = \xi^1(t)y(t)$. Let $Z = \Gamma Y$. We extend h to Z by equivariance. Let $N(Z)$ be the normal bundle of Z . Recall that $N(Z)$ can be identified with a Γ -invariant neighborhood of Z . We extend h to $N(Z)$ by letting $h(w) = h(z)$ for $w \in N_z$. We use an invariant cut-off function to extend the definition of h to all of R^n . A simple

computation shows that $\gamma^t(t)y(t)$ is a trajectory of $f+h$. It is also clear that $\|h\|$ goes to 0 as σ approaches e .

Proof of Theorem 5.1

Let Δ be as defined prior to the statement of Lemma 5.3, that is $\Delta = \Sigma/\Sigma$. Lemma 5.3 implies that Δ is either finite and cyclic or isomorphic to S^1 . We divide the analysis in two cases:

- (1) Δ is isomorphic to S^1 .
- (2) Δ is finite and cyclic.

Case (1).

It follows from the proof of Lemma 5.3 that Z is a relative equilibrium of f . Therefore the dynamics on Z is described by Theorem 4.5. Hence the trajectories on Z are dense in tori of maximal dimension equal to $\text{rank } N(\Sigma)/\Sigma$. We now show that $\text{rank } N(\Sigma)/\Sigma = \text{rank } N(\Sigma_y)/\Sigma_y + 1$. Note that Δ is a torus of dimension one contained in $N(\Sigma)/\Sigma$. Hence there is a maximal torus \mathbb{T} in $N(\Sigma)/\Sigma$ such that $\Delta \subset \mathbb{T}$. Clearly $\mathbb{T} \subset N(\Sigma_y)/\Sigma_y$. It follows that \mathbb{T}/Δ is a maximal torus in $N(\Sigma_y)/\Sigma_y$, which proves the required equality.

Case (2).

Let $z(t)$ be a trajectory on Z . By Theorem 2.2 $z(t) = \gamma(t)y(t)$, $\gamma(t) \in \Gamma$. Since $\Sigma_y(t) = \Sigma$ we can assume that $\gamma(t) \in N(\Sigma)/\Sigma$. Let T_0 be the number defined prior to the statement of Lemma 5.6 and let $\gamma_0 = \gamma(T_0)\delta_0$. Clearly $z(T_0) = \gamma_0 y_0$. Let $\mathbb{T} = \text{cl } \{z(t) : t \in \mathbb{R}\}$ and let

$$S = \text{cl } \{\gamma_0^k : k \text{ is an integer}\}.$$

Observe that $S\gamma_0 \subset \mathbb{T}$. Let Φ_t be the flow of f . We have

$$\Phi_t \gamma_0 y_0 = \Phi_t \Phi_{T_0} y_0 = \Phi_{t+T_0} y_0 = z(t + T_0).$$

It follows that $\Phi_t S\gamma_0 \subset Z$ for all $t \in \mathbb{R}$. Let us define the action of $\Gamma \times \mathbb{R}$ on \mathbb{R}^n as:

$$(t, \gamma) w = \Phi_t \gamma w, \quad \text{where } (t, \gamma) \in \Gamma \times \mathbb{R}, w \in \mathbb{R}^n. \tag{5.2}$$

It follows that $\mathbb{T} = (S \times R)y_0$. Let Δ_0 be the isotropy subgroup of y_0 with respect to the action defined by (5.2). It is clear that $(S \times R)/\Delta_0$ is compact, connected and abelian. Hence $(S \times R)/\Delta_0$ is isomorphic to a torus. This implies that \mathbb{T} diffeomorphic to a torus. Note that S is topologically cyclic and y_0 is its generator. Hence the dimension of S is less or equal to the dimension of a Cartan subgroup containing y_0 . Note that y_0 and δ_0 lie in the same component of the identity in $N(\Sigma)/\Sigma$, which we denote by $S(\delta_0)$. Let K be a Cartan subgroup generated by an element $\gamma_1 \in S(\delta_0)$. Proposition 5.5 implies that $\dim S \leq \dim K$ and $\dim \mathbb{T} \leq \dim K + 1$.

We now prove that for a small perturbation of f the dimension of the closure of a trajectory on Z equals $\dim K + 1$. Suppose that $\gamma_1 \in S(\delta_0)$ generates a Cartan subgroup. By Lemma 5.6 there exists a vector field h satisfying (VH1) and (VH2) and such that if $\tilde{z}(t) = \tilde{\gamma}(t)y(t)$ is the trajectory of $f+h$ with initial condition y_0 then $\tilde{\gamma}(T^0)\delta_0 = \gamma_1$. Clearly the dimension of the closure of $\tilde{z}(t)$ equals $\dim K + 1$.

To conclude the proof we need to show that $\dim K = \text{rank } N(\Sigma^y)/\Sigma^y$. By Proposition 5.4 we can choose K so that $\Delta \subset K$. The definition of Δ and the fact that Δ is discrete imply that $\text{rank } (N(\Sigma^y)/\Sigma) = \text{rank } (N(\Sigma^y)/\Sigma^y)$. Let $N(\Delta)$ denote the normalizer of Δ in $N(\Sigma)/\Sigma$. Note that $N(\Delta) = N(\Sigma^y)/\Sigma$. Clearly $K \subset N(\Delta)$. We show that $\text{rank } N(\Delta) = \dim K$. Suppose that $\phi \in N(\Delta)_0$ and let \mathbb{T}_0 be the torus generated by ϕ . Let $\phi_0 = \phi\delta_0$ and let K be the topologically cyclic subgroup generated by ϕ_0 . Since $\phi \in N(\Delta)$ we have $\phi\Delta\phi^{-1} = \Delta$, which implies that

$$\phi\delta_0 = \delta_m^0\phi, \text{ for some } m. \tag{5.3}$$

Note that continuity implies that m is independent of ϕ . It follows that for any positive integer j we must have $\phi^j = \delta_s^0\phi^j$, for some s (depending on j but independent of ϕ). Since ϕ_0 is the generator of K it follows that for some l we must have $\phi^l \in K_0 \subset N(\Delta)_0$. By continuity we must have $\phi^l \in N(\Delta)_0$ for all $\phi \in N(\Delta)_0$. Now (5.3) implies that $\delta_s^0\phi^l \in N(\Delta)_0$, for some s (independent of $\phi \in N(\Delta)_0$). It follows that for some ϕ

the torus generated by $\delta_0 \phi'$ is a maximal torus in $N(\Delta)$. It follows that $\dim \mathbb{T}_0 \leq \dim \mathbb{K} \leq \dim K$. The inequality $\dim K \geq \text{rank } N(\Delta)$ follows from the fact that K_0 is a connected abelian subgroup of a compact Lie group, hence it is contained in a maximal torus. It follows that $\text{rank } N(\Delta) \geq \dim K$.

Let F be a family of vector fields satisfying (H1) and (H2). Suppose that Δ is finite and cyclic and let T_0 be as defined prior to the statement of Lemma 5.6. Let $x_\lambda(t)$ be the trajectory of $F(\cdot, \lambda)$ with initial condition y_λ . By Theorem 2.2 $x_\lambda(t) = \gamma_\lambda(t) y_\lambda(t)$ ($y_\lambda(t)$ is the periodic orbit of $F_N(\cdot, \lambda)$). Let $\gamma(\lambda) = \gamma_\lambda(T_0)$. The proof of Theorem 5.2 is based on the following assertion: a generic F_T gives rise to a generic curve γ . Given H satisfying (H1) and (H2) let $\gamma_H^\lambda(t) y_\lambda(t)$ be the trajectory of $H(\cdot, \lambda)$. We now define a map Θ which assigns to every family of vector fields satisfying (H1) and (H2) the corresponding curve $\gamma(\lambda)$. Let $I \subset (0, \lambda_0)$ be an open interval and let

$$\Theta(H)(\lambda) = \gamma_H^\lambda(T_0), \quad \lambda \in I.$$

Let $S(\delta_0)$ be the connected component of $N(\mathbb{Z})/\mathbb{Z}$ containing δ_0 . Note that $\Theta(H)$ is an element of $C^\infty(I, S(\delta_0))$.

Proof of Theorem 5.2

The proof is analogous to the proof of Theorem 4.1' (ii). We will therefore only outline the proof and omit the technical details. If Δ is isomorphic to S^1 then F satisfies the assumptions of Theorem 4.1' with $y(\lambda) = \gamma_\lambda$ being the curve of relative equilibria. Hence the theorem follows from Theorem 4.1' and part (ii) of the proof of Theorem 5.1.

Suppose that Δ is finite and cyclic. Let $Q \subset S(\delta_0)$ be the set of elements which do not generate a Cartan subgroup. We show that Q is a countable union of submanifolds of $S(\delta_0)$, each of codimension greater or equal to 1. Let K be a Cartan subgroup containing δ_0 and generated by an element of $S(\delta_0)$. The existence of K follows from Proposition 5.4. Recall that K is isomorphic to $K_0 \times Z_l$ with $K_0 \times \{1\}$ corresponding to $\delta_0 K_0$ (we identify Z_l with $\{0, 1, 2, \dots, l-1\}$). In K_0 we define the sets P_m (see proof of Proposition 4.3). It follows that $\delta_0 K_0 \cap Q$ is the union of the sets $P_m \times \{1\}$. Let E_m be the union of all conjugacy classes of $P_m \times \{1\}$ by the elements of $N(\mathbb{Z})/\mathbb{Z}$. Proposition 5.5 implies that Q is the union of the sets E_m . Consider the action of $(N(\mathbb{Z})/\mathbb{Z})_0$ on $N(\mathbb{Z})/\mathbb{Z}$ defined as conjugation by a group element. As in the

proof of Theorem 4.1' we can partition $E_m(\Delta)$ into manifolds $E_m(\Delta)$, consisting of all elements of E_m whose isotropy with respect to this action is conjugate to Δ .

The remaining part of the proof is analogous to the proof of Theorem 4.1'. The main objective is to show that there exists a residual subset $B \subset C_\infty^T(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ such that if $H \in B$ then for all choices of m and Δ the curve $\Theta(H)$ is transverse to $E_m(\Delta)$ at every $\lambda \in (0, \lambda_0)$. This is done by showing that for some fixed m and Δ there exists a residual set $B_m(\Delta)$, whose elements are transverse to $E_m(\Delta)$, and then taking the intersection of the sets $B_m(\Delta)$.

We now fix m and Δ and regard Θ as a mapping from $C_\infty^T(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ to $C_\infty(I, S(\delta_0))$. To show existence of $B_m(\Delta)$ we need to prove the following properties of Θ :

(1) If the interval I is such that $I \subset (0, \lambda_0)$ then Θ is continuous in the Whitney C_∞ topology.

(2) For each $\mu \in (0, \lambda_0)$ there exists an interval I such that $\mu \in I$ and $I \subset (0, \lambda_0)$ and a residual set $A \subset C_\infty^T(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ such that if $H \in A$ then $\Theta(H)$ is transverse to $E_m(\Delta)$ at all $\lambda \in I$.

The property (1) follows from standard theorems on smooth dependence of solutions of ordinary differential equations on parameters (see also proof of Theorem 2.2).

We now indicate how to prove the property (2). We choose an interval I_0 such that $\mu \in I_0$ and $I_0 \subset (0, \lambda_0)$. Given some interval $I \subset I_0$ we define A_0 as the set of elements of $C_\infty(I_0, S(\delta_0))$ which are transverse to $E_m(\Delta)$ at all $\lambda \in I$. Standard transversality theory implies that A_0 is residual in the Whitney C_∞ topology on $C_\infty(I_0, S(\delta_0))$. Let $A = \Theta^{-1}(A_0)$. The property (1) implies that A is an intersection of open sets. If I is small enough then A is dense in the Whitney C_∞ topology on $C_\infty^T(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$.

To see this suppose that $\gamma \in C_\infty(I_0, S(\delta_0))$ is a small perturbation of the curve $\Theta(F)$. Then there exists a small perturbation H of the family F such that $\Theta(F+H) = \gamma$. The proof of the existence of H is a straightforward generalization of Lemma 5.6.

The methods used in the proof of Theorem 5.2 can be easily generalized to prove the following proposition.

Proposition 5.9

Suppose that F satisfies (H1) and (H2). Then there exist a residual set $B \subset C_{\infty}^1(\mathbb{R}^n \times \mathbb{R}, \mathbb{R}^n)$ such that if $H \in B$ then the dimension of the trajectories of $F+H$ on the sets Y is greater or equal to $\text{rank}(N(\Sigma_Y)/\Sigma_Y)$.

6. Bifurcations of relative equilibria with $O(2)$ symmetry.

In this section we discuss bifurcation problems with symmetry group $O(2)$. Let τ denote semi direct product. Recall that $O(2) = SO(2) \rtimes Z_2(\kappa)$, where $Z_2(\kappa) = \{1, \kappa\}$, κ is an orientation reversing element of $O(2)$ and $SO(2)$ is the subgroup of $O(2)$ consisting of orientation preserving rotations. We assume that $O(2)$ acts on \mathbb{R}^n and that $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ is a smooth and equivariant family of vector fields. We restrict our attention to bifurcations of group orbits of equilibria whose isotropy subgroups are either $Z_2(\kappa)$ or D_k , $k \geq 2$. Here D_k denotes the group of symmetries of a regular k -gon. The groups $Z_2(\kappa)$ or D_k occur as maximal isotropy subgroups for the various irreducible representations of $O(2)$, hence the bifurcations we study can occur as secondary bifurcations from an invariant equilibrium.

Let R_θ denote the rotation of the plane by the angle θ . The groups D_k are generated by the reflection κ and the rotation $R_{2\pi/k}$. Let Z_k denote the cyclic group generated by $R_{2\pi/k}$. We have $D_k = Z_k \rtimes Z_2(\kappa)$. In the sequel we use D_1 to denote $Z_2(\kappa)$ and Z_1 to denote the trivial subgroup 1 .

Fix $x_0 \in \mathbb{R}^n$, let $X = O(2)x_0$ and let Σ_{x_0} be the isotropy subgroup of x_0 . We assume that $\Sigma_{x_0} = D_k$, $k \geq 1$ and that X is a relative equilibrium of F . Let G be as defined in Section 3, that is G is the restriction of F_N to the normal space N_{x_0} . Let $g = G(\cdot, 0)$ and let E be the center subspace of $(dg)_{x_0}$. In this section we analyze steady state and Hopf bifurcations of F near X . More precisely we consider the following situations:

- (a) $(dg)_{x_0}$ has a zero eigenvalue.
- (b) $(dg)_{x_0}$ has a purely imaginary eigenvalue $i\omega$.

Let $K(\Sigma_{x_0}) = \{ \sigma \in \Sigma_{x_0} : \sigma v = v \text{ for all } v \in E \}$ be the kernel of the action of Σ_{x_0} on E . We assume that the action of Σ_{x_0} on E is nontrivial, that is $K(\Sigma_{x_0})$ is properly contained in Σ_{x_0} . We now state the two main results of this section – a steady state bifurcation theorem and a Hopf bifurcation theorem. We begin with the steady state bifurcation theorem. Suppose that $(dg)_{x_0}$ has a zero eigenvalue. We make a generic assumption that E is the nullspace of $(dg)_{x_0}$ and that the action of Σ_{x_0} on E is absolutely irreducible. The following theorem describes all the generic types of bifurcating solutions and gives the number of distinct nonconjugate branches:

Theorem 6.1

All the generic types of bifurcating solutions of F are listed in Table 6.1.

TABLE 6.1

kernel of isotropy	type of solution	number of branches
Z_k	rotating wave	1
$D_m, k = 2m$	steady state	1
$Z_l, l/k, l < k$	steady state	2

We now state the Hopf bifurcation theorem. Suppose that $(dg)_{x_0}$ has a purely imaginary eigenvalue $i\omega$. We make a generic assumption that E is the eigenspace of $i\omega$ and that the action of Σ_{x_0} on E is T -simple. The following theorem describes all the generic types of bifurcating solutions and gives the number of distinct nonconjugate branches:

Theorem 6.2

All the generic types of bifurcating solutions of F are listed in Table 6.2.

TABLE 6.2

kernel of isotropy	type of solution	number of branches
Z_k	periodic orbit	1
$D_m, k = 2m$	periodic orbit	1
$Z_l, l/k, l < k$	periodic orbit	2
	two-torus	1

In the remainder of this section we prove Theorem 6.1 and Theorem 6.2. We begin by classifying all the possible kernels of the action of Σ_{x_0} on E . We prove the following lemma:

Lemma 6.3

One of the following statements must hold:

- (i) $K(\Sigma_{x_0}) = Z_l$, $l \leq k$, l divides k , $l \neq \frac{k}{2}$.
- (ii) $k = 2m$ and $K(\Sigma_{x_0})$ is isomorphic to D_m .

Proof

Note that $K(\Sigma_{x_0})$ is normal in Σ_{x_0} . Hence finding all the possible groups $K(\Sigma_{x_0})$ is equivalent to classifying the normal subgroups of Σ_{x_0} .

We consider two cases: $K(\Sigma_{x_0}) \subset SO(2)$ or $K(\Sigma_{x_0}) \not\subset SO(2)$. Suppose that $K(\Sigma_{x_0}) \subset SO(2)$. Then $K(\Sigma_{x_0})$ is a subgroup of Z_k . Hence $K(\Sigma_{x_0}) = Z_l$, $l \leq k$, l divides k .

Suppose now that $K(\Sigma_{x_0}) \not\subset SO(2)$. Let $\zeta \in K(\Sigma_{x_0})$, $\zeta \notin SO(2)$. Since $K(\Sigma_{x_0})$ is normal in Σ_{x_0} we have $R_{4\pi/k} \zeta = R_{2\pi/k} \zeta \in K(\Sigma_{x_0})$. Hence $R_{4\pi/k} \in K(\Sigma_{x_0})$. If $k = 2m$ then it follows that $K(\Sigma_{x_0})$ is generated by $R_{4\pi/k}$ and ζ and therefore is isomorphic to D_m . Otherwise $k + 1$ is divisible by 2 and $(R_{4\pi/k})^{(k+1)/2} = R_{2\pi/k}$. This implies that $K(\Sigma_{x_0}) = D_k = \Sigma_{x_0}$, which is a contradiction.

The case $K(\Sigma_{x_0}) = Z_l$, $l \neq \frac{k}{2}$ cannot occur, since then $\Sigma_{x_0}/K(\Sigma_{x_0})$ would be isomorphic to D_2 and every irreducible action of D_2 has a nontrivial kernel.

Proof of Theorem 6.1

We begin by describing the bifurcation problem for the family G . We assume that the center manifold reduction has been carried out, that is G is a family of Σ_{x_0} -equivariant vector fields on E . Recall that $K(\Sigma_{x_0})$ is normal in Σ_{x_0} . Let $\tau : \Sigma_{x_0} \rightarrow \Sigma_{x_0}/K(\Sigma_{x_0})$ be the natural projection. We define the action of $\tau(\Sigma_{x_0})$ on E by $\tau(\sigma)v = \sigma v$, $v \in E$, $\sigma \in \Sigma_{x_0}$. Note that this action is well defined, since $\ker \tau = K(\Sigma_{x_0})$. It follows that G is Σ_{x_0} -equivariant if and only if it is $\tau(\Sigma_{x_0})$ -equivariant. We replace the action of Σ_{x_0} by the action of $\tau(\Sigma_{x_0})$.

As indicated in Table 6.1 we divide the analysis into three cases:

(1) $K(\Sigma^{x_0}) = Z_l, l < k, l$ divides k .

(2) $K(\Sigma^{x_0}) = Z_k$.

(3) $k = 2m$ and $K(\Sigma^{x_0})$ is isomorphic to D_m .

Case (1).

Let $m = \frac{l}{k}$. Since $l < k$ it follows that $m \geq 2$. Clearly $\tau(\Sigma^{x_0})$ is isomorphic to

D_m . Since the action of $\tau(\Sigma^{x_0})$ is faithful it follows that $\dim E > 1$. This implies that

$\dim E = 2$, since all the irreducible representations of D_m are one or two dimensional.

Also $m \geq 3$, since any irreducible representation of D_2 has a nontrivial kernel. It follows

that the action of $\tau(\Sigma^{x_0})$ is isomorphic to the standard action of D_m on C . According to

Table XIII, 5.2 in Golubitsky, Stewart and Schaeffer [1988] a generic family G has two

branches of steady state solutions $y_1(\lambda)$ and $y_2(\lambda)$. Let $Y_1 = O(2)Y_1$ and $Y_2 = O(2)Y_2$.

We now show that the sets Y_1 and Y_2 consist of equilibria of F . The results of

Golubitsky, Stewart and Schaeffer also imply that the isotropy subgroups of Y_1 and Y_2

with respect to the action of D_m are two element groups, each generated by an element not

contained in Z_m . Let Σ_{Y_1} and Σ_{Y_2} denote the isotropy subgroups of Y_1 and Y_2 in Σ^{x_0} .

It follows that $\Sigma_{Y_1} \subset SO(2)$ and $\Sigma_{Y_2} \subset SO(2)$. Hence the normalizers of Σ_{Y_1} and Σ_{Y_2}

are discrete. Theorem 4.1 implies that the group orbits Y_1 and Y_2 consist of equilibria of

F .

Case (2).

Note that $\tau(\Sigma^{x_0})$ is isomorphic to Z_2 . Hence $\dim E = 1$. Since the action of Z_2

is nontrivial it follows that Z_2 acts on E as minus identity. Hence generically G

undergoes a pitchfork bifurcation, that is, G has a unique (up to conjugacy) branch of

equilibria $y(\lambda)$ with trivial isotropy. For more information on Z_2 -equivariant bifurcation

see Golubitsky and Schaeffer [1985], Chapter XVI.

Let $Y = O(2)y$. We show that generically the trajectories of F on Y are rotating

waves. Let Σ_y be the isotropy subgroup of y in Σ^{x_0} . It follows that $\Sigma_y = K(\Sigma^{x_0}) =$

Z_k . We conclude that $N(\Sigma_y) = O(2)$. By Theorem 4.1 generically the trajectories of

$F(\cdot, \lambda)$ on $Y(\lambda)$ are given by drift along circles.

Case (3)

As in Case (2) $\tau(\Sigma_{x_0}^2)$ is isomorphic to Z_2 . Hence G has a branch of equilibria $\Sigma_y(\lambda)$, whose isotropy subgroup in Z_2 is trivial. Let Σ_y be the isotropy subgroup of y in $\Sigma_{x_0}^2$. Since $\Sigma_y = K(\Sigma_{x_0}^2)$ and $K(\Sigma_{x_0}^2)$ is isomorphic to D_m it follows that $\Sigma_y \not\subset SO(2)$ and that $N(\Sigma_y)$ is discrete. By Theorem 4.1 the orbit $Y = O(2)y$ consists of equilibria of F .

Before proving Theorem 6.2 we give some background on Hopf bifurcation from an invariant equilibrium. The results we review will be used in the analysis of bifurcations of the family G . Let $\Gamma \subset O(n)$ be a Lie group acting on \mathbb{R}^n and suppose that this action is nontrivial. Let $H : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n \times \mathbb{R}$ be a family of smooth, Γ -equivariant vector fields. Let $h = H(\cdot, 0)$ and suppose that $(dh)_0$ has a purely imaginary eigenvalue ω . Suppose that the center manifold reduction has been carried out, that is \mathbb{R}^n is the real part of the sum of the eigenspaces of $\pm \omega i$. We make a generic assumption that the action of Γ on \mathbb{R}^n is Γ -simple. Then the group $\{\exp(Lt) : t \in \mathbb{R}\}$ is isomorphic to S^1 . We define the action of S^1 on \mathbb{R}^n as

$$(\gamma, \theta) x = \gamma \exp(L\theta)x, \text{ where } (\gamma, \theta) \in \Gamma \times S^1 \text{ and } x \in \mathbb{R}^n.$$

The following theorem is the Equivariant Hopf Bifurcation Theorem (see Golubitsky and Stewart [1985], Theorem 5.1 or Golubitsky, Stewart and Shaeffer [1988], Theorem XVI, 4.1):

Theorem 6.4

Suppose that Δ is a maximal isotropy subgroup of $\Gamma \times S^1$ and $\dim \text{Fix}(\Delta) = 2$. Then H has a branch of small amplitude periodic solutions $x_\lambda(t)$ such that $\sigma x_\lambda(t + \theta) = x_\lambda(t)$ for every pair $(\sigma, \theta) \in \Delta$.

Suppose that $x_\lambda(t)$ is a branch of periodic solutions described in the statement of Theorem 6.4. Let $X(\lambda) = \{x_\lambda(t) : t \in \mathbb{R}\}$. Recall the definition of the group of symmetries of the set X , denoted by Σ_X , as the set of all $\sigma \in \Gamma$ such that $\sigma x = x$ for all $x \in \mathbb{R}^n$. Clearly Σ_X is obtained by projecting Δ onto the first component of $\Gamma \times S^1$, that is

$$\Sigma_X = \{\sigma \in \Gamma : (\sigma, \theta) \in \Delta \text{ for some } \theta \in S^1\}.$$

We refer to the group Δ as the group of spatial-temporal symmetries of the periodic orbit X.

We will now describe generic Hopf bifurcations in two cases: $\Gamma = Z_2$ and $\Gamma = D_k$, $k \geq 3$. Assume that $\Gamma = Z_2$. We have the following proposition:

Proposition 6.5

Generally the family H has a branch of periodic orbits $Y(\lambda)$ with $Z_{Y(\lambda)} = Z_2$.

Proof

The irreducible representations of Z_2 are absolutely irreducible and one dimensional. Hence a Γ -simple representation of Z_2 will be two dimensional and will have the form $R \oplus R$. The action on each of the copies of R will be given as reflection with respect to the origin. Let ζ be the nontrivial element in Z_2 . Then, for $(x, y) \in R^2$ $\zeta(x, y) = (-x, -y)$. Also

$$L = \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix}$$

and $\exp(Lt)$ is a rotation by angle t . It is easy to see that $Z_2(\zeta, \pi) = \{ (0, 0), (\zeta, \pi) \}$ is a maximal isotropy subgroup of $Z_2 \times S^1$. It follows from Theorem 6.4 that H has a branch of periodic orbits $Y(\lambda)$ with $Z_{Y(\lambda)} = Z_2$. Using normal form theory one can show that generically this branch is unique.

We now discuss that case when $\Gamma = D_k$, $k \geq 3$. We assume that the action of D_k on R^n is faithful. Let $\xi = \frac{k}{2\pi}$. We define the following subgroups of $D_k \times S^1$:

$$\begin{aligned} Z_k &= \left\{ \left(\frac{2\pi m}{k}, \frac{2\pi m}{k} \right) : m = 0, 1, \dots, k \right\} \\ Z_2(k) &= \{ (0, 0), (k, 0) \} \\ Z_2(k, \pi) &= \{ (0, 0), (k, \pi) \} \\ Z_2(k, \xi) &= \{ (0, 0), (k, \xi) \} \end{aligned}$$

and when k is even

$$Z_{2^c} = \{ (0, 0), (\pi, \pi) \}.$$

We have the following theorem:

Theorem 6.6

Generally the family H has three branches of periodic orbits: $Y_1(\lambda)$, $Y_2(\lambda)$ and $Y_3(\lambda)$. The groups of spatial-temporal symmetries of $Y_1(\lambda)$, $Y_2(\lambda)$ and $Y_3(\lambda)$ respectively are given by:

- (a) \tilde{Z}_k , $Z_2(k)$ and $Z_2(\kappa, \pi)$ if k is odd.
- (b) \tilde{Z}_k , $Z_2(k) \oplus Z_2^c$ and $Z_2(\kappa, \pi) \oplus Z_2$ if $k \equiv 2 \pmod{4}$.
- (c) \tilde{Z}_k , $Z_2(k) \oplus Z_2^c$ and $Z_2(\kappa, \xi) \oplus Z_2$ if $k \equiv 0 \pmod{4}$.

Theorem 6.6 is a consequence of Theorem XVIII, 3.1 in Golubitsky, Stewart and Shaeffer [1988].

We now present the proof of Theorem 6.2.

Proof of Theorem 6.2

We begin by describing the bifurcation problem for the family G . We assume that the center manifold reduction has been carried out, that is G is a family of Z_{x_0} -equivariant vector fields on E . Recall that τ is the natural projection from Z_{x_0} onto $Z_{x_0}/K(Z_{x_0})$. As in the proof of Theorem 6.1 we replace the action of Z_{x_0} by the action of $\tau(Z_{x_0})$. We consider three cases:

- (1) $K(Z_{x_0}) = Z_l$, $l > k$, l divides k .
- (2) $K(Z_{x_0}) = Z_k$.
- (3) $k = 2m$ and $K(Z_{x_0})$ is isomorphic to D_m .

Case (1). Let $m = \frac{k}{l}$. We have $\tau(\Sigma_{x_0}) = D_m$. Since the action of $\tau(\Sigma_{x_0})$ on E is faithful it follows that $m \geq 3$. The action of $\tau(\Sigma_{x_0})$ on E is Γ -simple, so we are in position to apply Theorem 6.6. Hence G has three branches of solutions $Y_1(\lambda)$, $Y_2(\lambda)$ and $Y_3(\lambda)$ whose groups of spatial-temporal symmetries in D_m are as indicated in Theorem 6.6. Let Σ_{Y_1} , Σ_{Y_2} and Σ_{Y_3} be the groups of symmetries of these trajectories inside of Σ_{x_0} . Observe that $\Sigma_{Y_1} = Z_k$ and the groups Σ_{Y_2} and Σ_{Y_3} are not contained in $SO(2)$. It follows that $N(\Sigma_{Y_1}) = O(2)$ and the normalizers of the groups Σ_{Y_2} and Σ_{Y_3} are discrete. Let $Z_1(\lambda) = O(2)Y_1(\lambda)$, $Z_2(\lambda) = O(2)Y_2(\lambda)$ and $Z_3(\lambda) = O(2)Y_3(\lambda)$. Theorem 5.2 implies that generically the trajectories of F on Z_1 are dense in two dimensional tori and the trajectories of F on Z_2 and Z_3 are periodic orbits.

Case (2). We have $\tau(\Sigma_{x_0}) = Z_2$. By Proposition 6.5 G has a unique branch of periodic orbits $Y(\lambda)$. Let ξ be the nontrivial element in Z_2 . It follows from the proof of Proposition 6.5 that (ξ, π) is a spatial-temporal symmetry of Y . Let Σ_Y denote the group of symmetries of Y inside of Σ_{x_0} . We have $\tau(\kappa) = \xi$, so $\Sigma_Y \not\subset SO(2)$ and $N(\Sigma_Y)$ is discrete. Let $Z = O(2)Y$. It follows that the flow of F on Z consists of periodic orbits.

Case (3). We have $\tau(\Sigma_{x_0}) = Z_2$. By Proposition 6.5 G has a unique branch of periodic orbits $Y(\lambda)$. Since $K(\Sigma_{x_0}) \not\subset SO(2)$ it follows that $\Sigma_Y \not\subset SO(2)$ and $N(\Sigma_Y)$ is discrete. Let $Z = O(2)Y$. It follows that the flow of F on Z consists of periodic orbits.

7. The Kuramoto-Sivashinsky equation.

The Kuramoto-Sivashinsky equation is used to model several physical and chemical phenomena, for example flame propagation and some aspects of the Belousov-Zhabotynski reaction. The following is the Kuramoto-Sivashinsky equation in one space variable:

$$u_t + 4 u^{xxxx} + \alpha (u_{xx} + \frac{1}{2} u^2) = 0 \tag{7.1}$$

In this section we study a bifurcation problem derived from (7.1). The equation (7.1) is equivariant with respect to translations and reflections in the space variable. An approach often used in such situations is to impose periodic boundary conditions with period $L > 0$. Then L becomes an additional parameter in the problem. The space variable x can be rescaled so that the boundary conditions become 2π periodic. As a result we obtain the following boundary value problem:

$$v_t + 4 v^{xxxx} + \alpha (v_{xx} + \frac{1}{2} v^2) = 0$$

$$v(x + 2\pi, t) = v(x, t) \tag{7.2}$$

where the period L has been absorbed into the parameter α . The boundary value problem (7.2) is $O(2)$ -equivariant. Hence the theory developed in Section 6 will apply to bifurcations of relative equilibria of (7.2).

An interesting aspect of the bifurcation analysis of the Kuramoto-Sivashinsky equation is that one can easily find primary branches of solutions with isotropy D_k , for all $k \geq 2$. This is a consequence of the following observation, which we later prove in Lemma 7.1. Suppose that u is a steady-state solution of (7.2). If we extend u by periodicity to the interval $[0, 2k\pi]$ and rescale the space variable by k , then the so obtained function is an equilibrium solution of (7.2) for a different value of the parameter α . The new equilibrium is called a *replicated solution*. Note that this solution is $2\pi/k$ periodic, which implies that its isotropy subgroup contains Z_k . It is easy to see that $u = 0$ is an equilibrium of (7.2). This equilibrium is stable for α near 0. As α is increased the solution $u = 0$ loses stability and bifurcates to a branch of solutions with isotropy group $Z_2(k)$. Hence for each $k \geq 2$ there exists a branch of replicated equilibria with symmetry D_k .

One might expect that the secondary branches of solutions bifurcating along the replicated branches would be replications of the secondary branches bifurcating from the

primary branch. According to the analysis of Section 6, however, secondary bifurcations from the replicated branches can be different from secondary bifurcations from the branch with symmetry $Z_2(\kappa)$. In particular we expect the branch with isotropy $Z_2(\kappa)$ to bifurcate to a rotating wave, and the branches with isotropy D_k to bifurcate to group orbits of equilibria. Kevrekedis, Nicolaenco and Scovel [1988] carried out a computer assisted study of secondary and tertiary bifurcations from the branches with isotropy groups $Z_2(\kappa)$, D_2 and D_3 . Their results fit the predictions of Theorems 6.1 and 6.2, in particular the first bifurcation along the branch of the equilibria with isotropy group $Z_2(\kappa)$ is to a rotating wave and the first bifurcations along the branches of equilibria with isotropy groups D_2 and D_3 are to orbits of equilibria. In this section we discuss the results of Kevrekedis et al. and compare them with the predictions of Theorem 6.1 and 6.2.

The numerical results of Kevrekedis et al. also indicate existence of quasiperiodic solutions and dynamics related to homoclinic and heteroclinic connections. None of these arise as a result of the bifurcations discussed in Section 6. Armbruster, Guckenheimer and Holmes [1988] analysed the $O(2)$ equivariant problem of interaction of two steady-state modes, one with isotropy group $Z_2(\kappa)$ and the other with isotropy group D_2 . The dynamics they found was much like the dynamics found by Kevrekedis et al. near the $Z_2(\kappa)$ and D_2 branches.

Let us now give a more detailed description of the bifurcation problem derived from the Kuramoto-Sivashinsky equation. We start by modifying the coordinates in (7.2) so that the solutions are bounded (see Kevrekedis et al. [1988]):

$$m(t) = \int_{2\pi}^0 v(x, t) dx.$$

We use (7.2) and the fact that the integrals $\int_{2\pi}^0 v_x dx$ and $\int_{2\pi}^0 v_{xxx} dx$ vanish to show:

$$\dot{m}(t) = -\frac{\alpha}{2} \int_{2\pi}^0 v^2 dx.$$

We now modify the coordinates by letting $u(x, t) = v(x, t) - m(t)$. We obtain:

$$u_t + 4u_{xxxx} + \alpha(u_x)^2 + \frac{1}{2}u_x^2 + \dot{m}(t) = 0 \tag{7.3}$$

$$u(x, t) = u(x + 2\pi, t).$$

Let us now describe the symmetries of (7.2) and (7.3). Let X be a space of four times differentiable functions $u(x, t)$, 2π periodic in the space variable x . The $O(2)$ action on X is generated by:

$$\begin{aligned} \theta u(x, t) &= u(x + \theta, t), \quad \theta \in SO(2) \\ \kappa u(x, t) &= u(-x, t). \end{aligned}$$

It is easy to see that (7.2) and (7.3) are equivariant with respect to this action.

Let us now explain in more detail how one obtains the replicated steady state solutions. The ideas we present can be found in Kevrekidis et al. Consider the steady state problem corresponding to (7.3):

$$4u_{xxxx} + \alpha(u_{xx} + \frac{1}{2}u_x^2) + \int_{2\pi}^0 u_x^2 dx = 0. \tag{7.4}$$

We assert the following: suppose $u(x)$ is a steady state solution of (7.3) with $\alpha = \alpha_0$ and let k be a positive integer. Then $w(x) = u(kx)$ is a solution of (7.3) with $\alpha = 4k^2\alpha_0$. To prove the assertion we apply the left hand side of (7.3) to w and use the fact that u is a solution.

Note that $u = 0$ is a trivial solution of (7.2). In order to determine the stability of 0 we write (7.2) as

$$u_t = F(u),$$

where

$$F(u) = -4u_{xxxx} + \alpha(u_{xx} + \frac{1}{2}u_x^2) + \dot{m}(t).$$

Then

$$dF|_{u=0} h = -4h_{xxxx} + \alpha h_{xx}.$$

It is easy to see that the functions $e^{2\pi i k x}$, where k is an integer, form a complete set of eigenvectors of $(dF)_{u=0}$ and the corresponding eigenvalues are $(2\pi)^2 k^2 (4k^2 - \alpha)$. The first instability occurs at $k = 1$ and $\alpha = 4$. As α crosses 4 a branch of equilibria with isotropy group $Z_2(k)$ bifurcates from the trivial solution. We will refer to it as the unimodal branch. It follows from Lemma 7.1 that this branch is replicated for $\alpha = 4k^2$.

These replicated branches will be referred to as the k -modal branches. Kevrekedis et al. describe some secondary and tertiary bifurcations discovered in their numerical studies of the Kuramoto-Sivashinsky equation. If we believe that those bifurcations are generic in the sense discussed in Section 6, then each one of them must match one of the cases described in Theorems 6.1 and 6.2. In what follows we summarize the findings of Kevrekedis et al. and relate them to the results of Theorems 6.1 and 6.2. The bifurcation diagram based on the results of Kevrekedis et al. is given in Fig. 7.1. The solid lines represent branches of asymptotically stable solutions and the dotted ones represent branches of unstable solutions.

Let U be the branch of equilibria with isotropy $Z_2(k)$, B the branch of equilibria with isotropy D_2 and T the branch of equilibria with isotropy D_3 . We discuss the secondary bifurcations found by Kevrekedis et al. along each of these branches. We first consider steady state bifurcations.

1. Steady state bifurcations from the branch U .

At $\alpha = 13.005$ the computations of Kevrekedis et al. reveal that a real eigenvalue passes through 0. In this case Theorem 6.1 predicts a bifurcation to a rotating wave. The numerical experiments confirm the existence of a rotating wave. Moreover Kevrekedis et al. give an analytical proof of the existence of the rotating wave, based on the ideas of Iooss [1986].

2. Steady state bifurcations from the branch B .

a) The first bifurcation on the branch B occurs at $\alpha = 16.1399$. Theorem 6.1 predicts a bifurcation to a unique branch of equilibria with isotropy group isomorphic to Z_2 (provided that the kernel of the action of D_2 on the nullspace is not contained in $SO(2)$). This is in agreement with the computations, which show that the branch U merges with the branch B .

b) An analogous bifurcation is observed for $\alpha = 22.559$. The isotropy group bifurcating branch is $Z_2(k)$. We label this branch BT since it later joins with the trimodal branch.

3. Steady state bifurcations from the branch T .

At $\alpha = 36.235$ two real eigenvalues of the branch T pass through 0. According to Theorem 6.1 there are two branches of equilibria bifurcating from the branch T . The isotropy of these equilibria is $Z_2(k)$. The numerical results are in complete agreement with this prediction. One of the bifurcating branches is the branch BT . Kevrekedis et

al. refer to the other branch as continuation of BT and also label it BT. For this reason we label this branch as BT.

4. Steady state bifurcations from the branch BT.

Kevekedis et al. also find a bifurcation point related to a zero eigenvalue on the branch BT. They conjecture that the corresponding bifurcation is to a rotating wave. Theorem 6.1 also predicts a bifurcation to a rotating wave and hence supports the conjecture.

Kevekedis et al. discuss three Hopf bifurcation points, marked in Fig. 7.1 as H1, H2, H3. The following are the predictions of the nature of these bifurcations derived from Theorem 6.2.

The Hopf bifurcation point 1

Point H1 corresponds to a Hopf bifurcation from the branch of rotating waves R. The group of symmetries of the branch of rotating waves is $SO(2)$. Theorem 6.2 implies that generically the bifurcating trajectories are dense in two-tori. This agrees with the predictions of Kevekedis, Nicolaenco and Scovel (closing remarks of § 5a), who conclude from the structure of the dynamics that a doubly periodic solution is likely to exist.


The Hopf bifurcation point 2

Point H2 corresponds to a Hopf bifurcation occurring along the branch B. The isotropy group of the equilibria on the branch B is D_2 . Theorem 6.2 implies that this Hopf bifurcation leads to a periodic flow. The numerical results of Kevekedis, Nicolaenco and Scovel indicate that the bifurcating solutions are periodic.

The Hopf bifurcation point 3

Point H3 corresponds to Hopf bifurcation occurring along the branch BT. Let Z be the isotropy group of the equilibria on that branch. We have previously remarked that $Z = Z_2(\kappa)$ or $Z = Z_2(\kappa, \pi)$. It follows from Theorem 6.2 that the bifurcating solutions must be periodic orbits. Kevekedis, Nicolaenco and Scovel do not comment on the dynamics related to this bifurcation.

Fig. 7.1 Secondary and tertiary bifurcation of the Kuramoto-Shivashinsky equation.



8. The Bénard problem.

In this section we analyze secondary steady state bifurcations of a dynamical system equivariant with respect to the group $\Gamma = D_6 + \mathbb{T}^2$, where \mathbb{T}^2 is a two dimensional torus and D_6 is the group of symmetries of a regular hexagon. A bifurcation problem with this symmetry arises in the analysis of the mathematical model of convection between two infinite planes. This problem is called the planar Bénard problem. We now briefly describe the symmetries of the model and the derivation of the bifurcation problem with symmetry group Γ . Detailed information on this topic and the analysis of primary bifurcations can be found in Buzano and Golubitsky [1983] or in Golubitsky, Stewart and Schaeffer [1988], Case Study 4.

Let x, y be the coordinates in a horizontal plane and z the coordinate in the vertical direction. The model of convection is equivariant with respect to translations, reflections and rotations in the xy plane. The group generated by these transformations is called the group of Euclidian motions in the plane and is denoted by E_2 . Let w be in \mathbb{R}^2 and let T_w denote translation by w . The the group of translations in the plane is isomorphic to \mathbb{R}^2 with the isomorphism defined by the assignment $w \mapsto T_w$. The group E_2 can be thought of as the semidirect product $O(2) \rtimes \mathbb{R}^2$ with multiplication defined by:

$$(\sigma_1, T_{w_1})(\sigma_2, T_{w_2}) = (\sigma_1\sigma_2, T_{w_1 + \sigma_1 w_2}) \quad \sigma_1, \sigma_2 \in O(2), w_1, w_2 \in \mathbb{R}^2. \quad (8.1)$$

Let $e \in \mathbb{R}^2$ be an arbitrary vector and let f be obtained by rotating e by $\frac{\pi}{3}$. The hexagonal lattice H_6 is given as:

$$H_6 = \{ ne + mf : \text{for all pairs of integers } n \text{ and } m \}$$

Note that H_6 is a subgroup of \mathbb{R}^2 . The solutions of the convection problem have the form $u(t, x, y, z)$. We restrict our attention to those which are periodic in both directions of the lattice. Let

$$X = \{ u(t, x, y, z) : u(t, (x, y, z) + (e, 0)) = u(t, (x, y, z) + (f, 0)) = u(t, x, y, z) \}$$

Clearly the only elements of $O(2)$ that leave X invariant are the elements of D_6 . Also the action of H_6 on X is trivial. Hence the group of symmetries of the convection problem restricted to X is given by Γ with $\mathbb{T}^2 = \mathbb{R}^2/H_6$.

Multiplication in Γ is induced by multiplication in E_2 . Let D_6 act on \mathbb{R}^2 by the standard action. Let 1 denote the identity in D_6 , 0 the identity in \mathbb{T}^2 and e the identity in Γ . For $p \in \mathbb{R}^2$ let p' denote the image of p under the natural projection $\mathbb{R}^2 \rightarrow \mathbb{T}^2$. For $\sigma \in D_6$ we define $\sigma \bullet p' = (\sigma p)'$. Multiplication in Γ is given as follows:

$$(\sigma_1, p_1') (\sigma_2, p_2') = (\sigma_1 \sigma_2, p_1' + \sigma_1 \bullet p_2'). \tag{8.2}$$

Here $\sigma_1 \sigma_2$ is the product of σ_1 and σ_2 in D_6 and $p_1' + \sigma_1 \bullet p_2'$ is the sum of vectors in \mathbb{T}^2 .

The Bénard problem has an invariant equilibrium (the pure conduction state). There exists a region in the parameter space where this equilibrium is stable. The known primary bifurcations are to two types of equilibria with maximal isotropy subgroups. These bifurcations are D_6 and $D_2 + S^1 (= Z_2 \oplus O(2))$. The equilibria with isotropy D_6 are called hexagons, and the equilibria with isotropy $Z_2 \oplus O(2)$ are called rolls. In what follows we describe the kinds of steady state bifurcations each one of these solutions can undergo.

From now on we consider an abstract Γ -equivariant bifurcation problem. We assume that $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n \times \mathbb{R}$ (for some n) is a smooth Γ -equivariant family of vector fields and that F has a branch of equilibria with isotropy group D_6 (which we refer to as hexagons) and a branch of solutions with isotropy group $Z_2 \oplus O(2)$ (which we refer to as rolls). We analyze the generic bifurcations of these solutions.

a. Bifurcations of hexagons.

Suppose $X = \Gamma x_0$ is a group orbit of hexagons. Let G be the restriction of F_N to the normal space $x_0 + N_{x_0}$ and let $g = G(\cdot, 0)$. We assume that $(dg)_{x_0}$ has a zero eigenvalue. Let E be the center subspace of $(dg)_{x_0}$. We make a generic assumption that E is the nullspace of $(dg)_{x_0}$ and that the action of D_6 on E is absolutely irreducible. Our bifurcation analysis will depend on the action of D_6 on E . Let $K(D_6)$ be the kernel of the action of D_6 on E . We assume that the bifurcation is symmetry breaking, that is, $K(D_6)$ is a proper subgroup of D_6 . According to Lemma 6.3 either $K(D_6) = Z_m$, $m = 1, 2, 6$, or $K(D_6)$ is isomorphic to D_3 . The following proposition gives a classification of generic bifurcations of hexagons.

Proposition 8.1

All generic types of bifurcating solutions of F are listed in Table 8.1.

TABLE 8.1

kernel of isotropy	type of solution	number of half branches
Z_6 or D_3	steady state	1
Z_2	steady state	2
trivial	periodic orbit	2

We now state and prove a lemma necessary to prove Proposition 8.1. Suppose that Σ is a subgroup of D_6 and let $N(\Sigma)$ denote the normalizer of Σ in Γ . Let $\text{Fix}(\Sigma)$ denote the fixed point space of Σ taken with respect to the standard action of D_6 on \mathbb{R}^2 . We have:

Lemma 8.2

$$\dim N(\Sigma) = \dim \text{Fix}(\Sigma).$$

Proof

Let Γ_0 denote the connected component of the identity in Γ . Since Γ is a compact group it suffices to show that the normalizer of Σ in Γ_0 has the same dimension as $\text{Fix}(\Sigma)$. The group Γ_0 consists of elements of form $(1, p)$, $p' \in \mathbb{T}^2$. Suppose that $(\sigma, 0)$ is in Σ . The element $(1, p)$ is in $N(\Sigma)$ if $(1, -p)(\sigma, 0)(1, p)$ is in Σ . We have

$$(1, -p)(\sigma, 0)(1, p) = (\sigma, \sigma \bullet p' - p'). \tag{8.3}$$

If (8.1) holds then we must have $\sigma \bullet p' - p' = 0$ which is equivalent to $(\sigma p - p)' = 0$. The lemma now follows.

Proof of Proposition 8.1

Let $\tau : D_6 \rightarrow D_6/K(D_6)$ be the natural projection. In the analysis of bifurcations of the family G we replace the action of D_6 be the action of $\tau(D_6)$. As indicated in Table 8.1 we divide the analysis into three cases:

- (1) $K(D_6) = Z_6$ or $K(D_6)$ is isomorphic to D_3 .
- (2) $K(D_6) = Z_2$.
- (3) $K(D_6)$ is trivial.

Case (1).

Observe that $\tau(D_6)$ is isomorphic to Z_2 . In this case a generic family G has a unique branch of equilibria $y(\lambda)$, whose isotropy subgroup in D_6 is trivial (see proof of Theorem 6.1). The isotropy group of $y(\lambda)$ in $\tau(D_6)$ is trivial. Let Z_y be the isotropy group of $y(\lambda)$ in D_6 . It follows that Z_y is $K(D_6)$. The group $K(D_6)$ contains a nontrivial rotation. It follows that the fixed point space of $K(D_6)$ with respect to the standard action of D_6 on R^2 is trivial. By Lemma 8.2 $\dim N(Z) = 0$. Let $Y(\lambda) = \tau y(\lambda)$. By Theorem 4.1 the set $Y(\lambda)$ consists of equilibria of F .

Case (2).

Observe that $\tau(D_6)$ is isomorphic to D_3 . It follows from Table XIII, 5.1 in Golubitsky, Stewart and Schaeffer [1988] that a generic family G has two half branches of equilibria $y_1(\lambda)$ and $y_2(\lambda)$. Let Z_{y_1} and Z_{y_2} denote the isotropy subgroups of y_1 and y_2 in D_6 . Both of these groups must contain $K(D_6)$. Since $K(D_6)$ contains the rotation by π it follows that the fixed point spaces of Z_{y_1} and Z_{y_2} are trivial. Lemma 8.2 implies that the normalizers of these groups are discrete. Let $Y_1(\lambda) = \tau y_1(\lambda)$, $Y_2(\lambda) = \tau y_2(\lambda)$. Theorem 4.1 implies that the sets $Y_1(\lambda)$ and $Y_2(\lambda)$ consist of equilibria of F .

Case (3).

According to Table XIII, 5.2 in Golubitsky, Stewart and Schaeffer [1988] a generic family G has two half branches of equilibria $y_1(\lambda)$, $y_2(\lambda)$, with $Z_{y_1} = \{1, \kappa\}$ and $Z_{y_2} = \{1, \kappa\pi\}$. Here κ denotes the reflection with through the x-axis and $\kappa\pi$ denotes the reflection through the y-axis. Clearly these groups have one dimensional fixed point spaces, hence, by Lemma 8.2, their normalizers are one dimensional. Let $Y_1(\lambda) =$

$\Gamma_{Y_1}(\lambda)$, $Y_2(\lambda) = \Gamma_{Y_2}(\lambda)$. Theorem 4.1 implies that generically the trajectories of the flow of F on the sets $Y_1(\lambda)$ and $Y_2(\lambda)$ are rotating waves.

b. Bifurcations of rolls.

Suppose $X = \Gamma x_0$ is a group orbit of rolls. Let G be the restriction of F_N to the normal space N_{x_0} and let $g = G(\cdot, 0)$. We assume that $(dg)_{x_0}$ has an eigenvalue on the imaginary axis. Let Z_r denote the isotropy subgroup of rolls. Let E be the center subspace of $(dg)_{x_0}$. We make a generic assumption that the action of Z_r on E is irreducible. We will show that this implies that the action of Z_r on E is absolutely irreducible. This implies that $(dg)_{x_0}$ must have a zero eigenvalue and hence the bifurcation we consider is a steady state bifurcation.

The group Z_r is generated by translations along the y -axis, reflection through the y -axis and rotation by π (see Golubitsky, Stewart and Schaeffer [1988] p. 154). The projection of the y -axis into the torus \mathbb{T}^2 is a circle which we denote by S^1 . Let ξ correspond to the element of D_6 which acts on \mathbb{R}^2 as rotation by π and let ζ denote the element of D_6 which acts as reflection through the y -axis. The element ζ commutes with the elements of S^1 and the element ξ anticommutes with the elements of S^1 . It follows that Z_r is isomorphic to $Z_2 \oplus O(2)$, with S^1 corresponding to $SO(2)$, ξ corresponding to a reflection in $O(2)$ and ζ corresponding to the nontrivial element of Z_2 . The irreducible representations of $Z_2 \oplus O(2)$ are given by the irreducible representations of $O(2)$, with ζ acting as identity or as minus identity. Since the irreducible representations of $O(2)$ are absolutely irreducible it follows that $(dg)_{x_0}|_E \equiv 0$.

We now divide that analysis into two cases:

- (1) S^1 acts trivially on E .
- (2) S^1 acts nontrivially on E .

Case (1).

In the analysis of bifurcations of G we can replace the action of Z_r on E by the action of $D_2 = \{1, \xi, \zeta, \xi\zeta\}$. The space E must be one dimensional and the kernel of the action of D_2 must be on of the groups:

$$\begin{aligned}
 Z_2(\xi) &= \{1, \xi\} \\
 Z_2(\zeta) &= \{1, \xi\} \\
 Z_2(\xi\zeta) &= \{1, \xi\zeta\}.
 \end{aligned}
 \tag{8.4}$$

It follows that in the analysis of bifurcations of G the action of D_2 can be replaced by the action of a group isomorphic to Z_2 . Hence a generic family G has a unique branch of equilibria $y(\lambda)$ and the isotropy group of $y(\lambda)$ in D_2 is equal to one of the groups listed in (8.4). Let Z_y be the isotropy subgroup of $y(\lambda)$ in Z_T . We have $Z_y = \langle S^1, \xi \rangle$, or $Z_y = \langle S^1, \xi \rangle$ or $Z_y = \langle S^1, \xi \rangle$. In other words Z_y is generated by the elements of S^1 and ξ , ξ or $\xi\xi$. Let $Y(\lambda) = \Gamma y(\lambda)$. We have the following proposition:

Proposition 8.3

If $Z_y = \langle S^1, \xi \rangle$ then the trajectories of flow of F on $Y(\lambda)$ are rotating waves. Otherwise $Y(\lambda)$ consists of equilibria.

Proof

We prove that if $Z_y = \langle S^1, \xi \rangle$ then $\dim N(Z_y) = 2$ and otherwise $\dim N(Z_y) = 1$. The proposition will then follow from Theorem 4.1.

By compactness of Γ it suffices to show that the normalizer of Z_y in Γ_0 has the indicated dimension. Recall that Γ_0 consists of the elements of Γ of the form $(1, p)$, $p \in \mathbb{T}^2$. Suppose that $(\sigma, 0) \in Z_y$, $\sigma \in D_2$. The identity (8.3) implies that

$$\sigma \cdot p' - p' \in S^1. \tag{8.4}$$

Recall that S^1 is the image of the y -axis under the natural projection $\mathbb{R}^2 \rightarrow \mathbb{T}^2$. Hence (8.4) implies that $\sigma p - p = (0, q)$, $q \in \mathbb{R}$.

Suppose that $p = (p_1, p_2)$. The element $\xi\xi$ acts on \mathbb{R}^2 as reflection through the x -axis, that is, $\xi\xi(p_1, p_2) = (p_1, -p_2)$. It follows that $\xi\xi p - p = (0, -2p_2)$. Hence (8.4) holds for all p' in S^1 . It follows that if $Z_y = \langle S^1, \xi\xi \rangle$ then $\dim N(Z_y) = 2$.

The element ξ acts on \mathbb{R}^2 as rotation by π , that is, $\xi(p_1, p_2) = (-p_1, -p_2)$. It follows that $\xi p - p = (-2p_1, -2p_2)$. Hence (8.4) holds if $p' \in S^1$. It follows that if $Z_y = \langle S^1, \xi \rangle$ then $\dim N(Z_y) = 1$.

The element ξ acts on \mathbb{R}^2 as reflection through the x -axis, that is, $\xi(p_1, p_2) = (-p_1, p_2)$. It follows that $\xi p - p = (-2p_1, 0)$. Hence (8.4) holds if $p' \in S^1$. It follows that if $Z_y = \langle S^1, \xi \rangle$ then $\dim N(Z_y) = 1$.

Case (2)

We show that if S^1 acts nontrivially then generically the flow on the bifurcating relative equilibria is trivial. Namely we prove the following proposition:

Proposition 8.4

A generic family F has a unique branch of bifurcating relative equilibria $Y(\lambda)$.

The flow on the sets $Y(\lambda)$ is trivial.

Proof

We first consider bifurcations of the family G . Let $K(\Sigma_T)$ be the kernel of the action of Σ_T on E . We show that $K(\Sigma_T)$ must be nontrivial. If ξ acts on E as identity then $(\xi, 0) \in K(\Sigma_T)$, which implies the assertion. Suppose that ξ acts on E as minus identity. Let p_0 be the element of S^1 acting on E as rotation by π . Then $(\xi, p_0) \in K(\Sigma_T)$, so $K(\Sigma_T)$ is nontrivial.

Since the action of S^1 is nontrivial it follows that the action of ξ on E is a reflection. Hence $K(\Sigma_T)$ is generated by a cyclic subgroup of S^1 and either $(\xi, 0)$ or (ξ, p_0) . It follows that $\Sigma_T/K(\Sigma_T)$ is isomorphic to $O(2)$. In the analysis of the bifurcations of G we replace the action of Σ_T on E by the action of $\Sigma_T/K(\Sigma_T)$ on E . By standard results on $O(2)$ -equivariant bifurcation G has a unique branch of equilibria $y(\lambda)$ and Σ_y contains a reflection. Since ξ acts as reflection we can assume (by possibly replacing $y(\lambda)$ by a conjugate branch) that $(\xi, 0) \in \Sigma_y$.

We now prove that the normalizer of Σ_y in Γ_0 is discrete. As we argued before this implies that normalizer of Σ_y in Γ is discrete. The group Σ_y is generated by $(\xi, 0)$ and the elements of $K(\Sigma_T)$. Suppose that $(1, p) \in N(\Sigma_y)$, $p = (p_1, p_2)$. Recall that ξ acts on \mathbb{R}^2 as rotation by π , that is, $\xi(p_1, p_2) = (-p_1, -p_2)$. The identity (8.3) implies that $\xi \bullet p' - p' \in \Sigma_y$. Also $\xi p - p = (-2p_1, -2p_2) = -2p'$. It follows that $2p'$ must be in $K(\Sigma_T) \cap S^1$, which, by assumption is a discrete group. It follows that $N(\Sigma_y)$ is discrete and by Theorem 4.1 the relative equilibria $Y(\lambda) = \Gamma y(\lambda)$ consist of equilibria of F .

REFERENCES

- D. Armbruster, J. Guckenheimer and P.J. Holmes [1987] Heteroclinic cycles and modulated traveling waves in a system with $O(2)$ symmetry. *Physica, 29D*, pp. 257-282.
- G. Bredon [1972] *Introduction to Compact Transformation Groups*. Pure and Applied Mathematics, v. 46, Academic Press, New York and London.
- T. Bröcker and T. tom Dieck [1985] *Representations of Compact Lie Groups*. Graduate Texts in Mathematics, Springer, New York.
- E. Buzano and M. Golubitsky [1983] Bifurcations on the hexagonal lattice and the Bénard problem. *Phil. Trans. R. Soc. Lond. A* 308, 617 - 667.
- P. Chossat [1986] Bifurcation secondaire de solution quasi périodiques dans un problème de bifurcation de Hopf par symétrie $O(2)$. *C.R. Acad. Sc. Paris*, v. 302, Série I, No. 15, 539 - 451.
- P. Chossat and M. Golubitsky [1988] Iterates of maps with symmetry. *Siam J. of Math. Anal.* Vol. 19, No. 6, November 1988.
- E. N. Dancer. [1980] An implicit function theorem with symmetries and its application to nonlinear eigenvalue problems. *Bull. Austral. Math. Soc.* 21, 404 - 437.
- G. Dangelmayr [1986] Steady state mode interactions in the presence of $O(2)$ symmetry. *Dynamics and Stability of Systems I*, 159 - 185.
- M. J. Field [1980] Equivariant dynamical systems. *Transactions of the AMS* 259, No. 1, May 1980, p. 185 - 205.
- M. J. Field [1988] *Equivariant Bifurcation Theory and Symmetry Breaking*. Preprint. University of Sydney.
- M. Golubitsky and V. W. Guillemin [1974] *Stable Mappings and Their Singularities* Springer, New York.
- M. Golubitsky and D. Schaeffer [1985] *Singularities and Groups in Bifurcation Theory*. Vol. I, Appl. Math Sci. 51, Springer, New York.
- M. Golubitsky and I. Stewart [1985] Hopf bifurcation in the presence of symmetry. *Arch. Rat. Mech. Anal.* Vol. 87, No. 2, 107 - 165.
- M. Golubitsky and I. Stewart, D. Schaeffer [1988] *Singularities and Groups in Bifurcation Theory*. Vol. II, Appl. Math Sci. 69, Springer, New York.
- V. Guillemin and A. Pollack [1974] *Differential Topology*. Prentice Hall 1974.
- G. Iooss [1986] Secondary bifurcations of the Taylor vortices into wavy inflow and outflow boundaries. *J. Fluid. Mechanics*, 173, 273 - 288.

- I. G. Kevrekedis, B. Nicolaenco and J. C. Scovel [1988] Back in the saddle again: a computer assisted study of the Kuramoto - Sivashinsky equation. *SIAM J. of Appl Math.*, to appear.
- E. Moutrane [1988] Interaction de modes spheriques dans le probleme de Bénard entre deux spheres. *Ph.D. thesis, Université Nice.*
- V. Poenaru [1976] *Singularités C^∞ en Présence de Symétrie*. Springer Lecture Notes, v. 510.
- D. Ruelle [1973] Bifurcations in presence of a symmetry group. *Arch. Rat. Mech. Anal.* 51, 131-172.
- A. Vanderbauwhede, M. Krupa and M. Golubitsky [1989] Secondary bifurcations in symmetric systems. *Proceedings of the Equadiff Conference 1987*, Lecture Notes in Pure and Applied Mathematics, v.118, Marcel Dekker, Inc., New York and Basel.

Recent IMA Preprints

#	Author/s	Title
547	M. Slemrod	Dynamics of Measured Valued Solutions to a Backward-Forward Heat Equation
548	Avner Friedman and Jürgen Sprekels	Steady States of Austenitic-Martensitic-Domains in the Ginzburg-Landau Theory of Shape Memory Alloys
549	Avner Friedman and Bei Hu	Degenerate Hamilton-Jacobi-Bellman Equations in a Bounded Domain
550	E.G. Kalnins, Willard Miller, Jr., and M.V. Tratnik	Families of Orthogonal and Biorthogonal Polynomials on the N-Sphere
551	Heinrich Freistühler	On Compact Linear Degeneracy
552	Matthew Witten	Quantifying the Concepts of Rate and Acceleration/Deceleration of Aging
553	J.P. Albert and J.L. Bona	Total Positivity and the Stability of Internal Waves in Stratified Fluids of Finite Depth
554	Brian Coomes and Victor Zurlkowski	Linearization of Polynomial Flows and Spectra of Derivations
555	Yuriko Renardy	A Couette-Poiseuille Flow of Two Fluids in a Channel
556	Michael Renardy	Short wave instabilities resulting from memory slip
557	Daniel D. Joseph and Michael Renardy	Stokes' first problem for linear viscoelastic fluids with finite memory
558	Xiaxi Ding	Superlinear Conservation Law with Viscosity
559	J.L. Ericksen	Liquid Crystals with Variable Degree of Orientation
560	F. Robert Ore, Jr. and Xinfu Chen	Electro-Optic Modulation in an Arbitrary Cross-Section Waveguide
561	M.V. Tratnik	Multivariable biorthogonal continuous-discrete Wilson and Racah polynomials
562	Yisong Yang	Existence of Solutions for a Generalized Yang-Mills Theory
563	Peter Grtitzmann, Laurent Habstieger and Victor Klee	Good and Bad Radii of Convex Polygons
564	Martin Golubitsky, Martin Krupa and Chjan. C. Lim	Time-Reversibility and Particle Sedimentation
565	G. Yin	Recent Progress in Parallel Stochastic Approximations
566	G. Yin	On H-Valued SA: Finite Dimensional Approximations
567	Chien-Cheng Chang	Accurate Evaluation of the Effect of Diffusion and Conductivity in Certain Equations
568	Chien-Cheng Chang and Ruey-Ling Chern	The Effect of Viscous Diffusion in Discrete Vortex Dynamics for Slightly Viscous Flows
569	Li Ta-Tsien (Li Da-qian) and Zhao Yan-Chun	Global Existence of Classical Solutions to the Typical Free Boundary Problem for General Quasilinear Hyperbolic Systems and its Applications
570	Thierry Cazenave and Fred B. Weissler	The Structure of Solutions to the Pseudo-Conformally Invariant Nonlinear Schrödinger Equation
571	Marshall Slemrod and Athanasios E. Tzavaras	A Limiting Viscosity Approach for the Riemann Problem in Isentropic Gas Dynamics
572	Richard D. James and Scott J. Spector	The Formation of Filamentary Voids in Solids in P.J. Vassiliou, On the Geometry of Semi-Linear Hyperbolic Partial Differential Equations in the Plane Integrable by the Method of Darboux
573	Jerome V. Moloney and Alan C. Newell	Nonlinear Optics
574	Keti Tenenblat	A Note on Solutions for the Intrinsic Generalized Wave and Sine-Gordon Equations
575	P. Szmolany	Heteroclinic Orbits in Singularly Perturbed Differential Equations
576	Wenxiong Liu	A Parabolic System Arising in Film Development
577	Daniel B. Dix	Temporal Asymptotic Behavior of Solutions of the Benjamin-Ono-Burgers Equation
578	Michael Renardy and Yuriko Renardy	On the nature of boundary conditions for flows with moving free surfaces
580	Werner A. Stahel	Robust Statistics: From an Intellectual Game to a Consumer Product
581	Avner Friedman and Fernando Reich	The Stefan Problem with Small Surface Tension
582	E.G. Kalnins and W. Miller, Jr.	Separation of Variables Methods for Systems of Differential Equations in Mathematical Physics
583	Mitchell Luskin and George R. Sell	The Construction of Inertial Manifolds for Reaction-Diffusion Equations by Elliptic Regularization
584	Konstantin Mischaikow	Dynamic Phase Transitions: A Connection Matrix Approach
585	Philippe Le Floch and Li Ta-Tsien	A Global Asymptotic Expansion for the Solution to the Generalized Riemann Problem
586	Matthew Witten, Ph.D.	Computational Biology: An Overview
587	Matthew Witten, Ph.D.	Peering Inside Living Systems: Physiology in a Supercomputer

#	Author/s	Title
588	Michael Renardy,	An existence theorem for model equations resulting from kinetic theories of polymer solutions
589	Daniel D. Joseph and Luigi Preziosi,	Reviews of Modern Physics: Addendum to the Paper "Heat Waves"
590	Luigi Preziosi,	An Invariance Property for the Propagation of Heat and Shear Waves
591	Gregory M. Constantine and John Bryant,	Sequencing of Experiments for Linear and Quadratic Time Effects
592	Prabir Daripa,	On the Computation of the Beltrami Equation in the Complex Plane
593	Philippe Le Floch,	Shock Waves for Nonlinear Hyperbolic Systems in Nonconservative Form
594	A.L. Gorin, D.B. Roe and A.G. Greenberg,	On the Complexity of Pattern Recognition Algorithms On a Tree-Structured Parallel Computer
595	Mark J. Friedman and Eusebius J. Doedel,	Numerical computation and continuation of invariant manifolds connecting fixed points
596	Scott J. Spector,	Linear Deformations as Global Minimizers in Nonlinear Elasticity
597	Denis Serre,	Richness and the classification of quasilinear hyperbolic systems
598	L. Preziosi and F. Rosso,	On the stability of the shearing flow between pipes
599	Avner Friedman and Wenxiong Liu,	A system of partial differential equations arising in electrophotography
600	Jonathan Bell, Avner Friedman, and Andrew A. Lacey,	On solutions to a quasilinear diffusion problem from the study of soft tissue
601	David G. Schaeffer and Michael Shearer,	Loss of hyperbolicity in yield vertex plasticity models under nonproportional loading
602	Herbert C. Kranzer and Barbara Lee Keyfitz,	A strictly hyperbolic system of conservation laws admitting singular shocks
603	S. Laedrich and M. Levi,	Qualitative dynamics of planar chains
604	Milan Miklavcic,	A sharp condition for existence of an inertial manifold
605	Charles Collins, David Kinderlehrer, and Mitchell Luskin,	Numerical approximation of the solution of a variational problem with a double well potential
606	Todd Arbogast,	Two-phase incompressible flow in a porous medium with various nonhomogeneous boundary conditions
607	Peter Polacik,	Complicated dynamics in scalar semilinear parabolic equations in higher space dimension
608	Bei Hu,	Diffusion of penetrant in a polymer: a free boundary problem
609	Mohamed Sami ElBialy,	On the smoothness of the linearization of vector fields near resonant hyperbolic rest points
610	Max Jodet, Jr. and Peter J. Olver,	On the equation $\text{grad } f = M \text{ grad } g$
611	Shui-Nee Chow, Kenning Lu, and Yun-Qiu Shen,	Normal form and linearization for quasiperiodic systems
612	Prabir Daripa,	Theory of one dimensional adaptive grid generation
613	Michael C. Mackey and John G. Milton,	Feedback, delays and the origin of blood cell dynamics
614	D.G. Aronson and S. Kamih,	Disappearance of phase in the Stefan problem: one space dimension
615	Martin Krupa,	Bifurcations of relative equilibria
616	D.D. Joseph, F. Singh, and K. Chen,	Couette flows, rollers, emulsions, tall Taylor cells, phase separation and inversion, and a chaotic bubble in Taylor-Couette flow of two immiscible liquids
617	Artemio Gonzalez-Lopez, Niky Karaman, and Peter J. Olver,	Lie algebras of differential operators in two complex variables
618	L.E. Fraenkel,	On a linear, partly hyperbolic model of viscoelastic flow past a plate
619	Stephen Schecter and Michael Shearer,	Undercompressive shocks for nonstrictly hyperbolic conservation laws
620	Xinfu Chen,	Axially symmetric jets of compressible fluid
621	J. David Logan,	Wave propagation in a qualitative model of combustion under equilibrium conditions
622	M.T. Zeman,	Hopf bifurcations in competitive three-dimensional Lotka-Volterra Systems
623	Allan P. Fordy,	Isospectral flows: their Hamiltonian structures, Miura maps and master symmetries
624	Daniel D. Joseph, John Nelson, Michael Renardy, and Yuriko Renardy,	Two-Dimensional cusped interfaces