

## Binge Drinking in Adolescents: A Review of Neurophysiological and Neuroimaging Research

Géraldine Petit<sup>1</sup>, Pierre Maurage<sup>2</sup>, Charles Kornreich<sup>1</sup>, Paul Verbanck<sup>1</sup> and Salvatore Campanella<sup>1,\*</sup>

<sup>1</sup>Laboratory of Psychological Medicine and Addictology, ULB Neuroscience Institute (UNI), Université Libre de Bruxelles (ULB), Brussels, Belgium and

<sup>2</sup>Laboratory for Experimental Psychopathology, Institute of Psychology, Catholic University of Louvain, Louvain-la-Neuve, Belgium

\*Corresponding author: The Belgian Fund for Scientific Research (FNRS), CHU Brugmann, Department of Psychiatry, 4 Place Vangehuchten, 1020 Brussels, Belgium. Tel.: +32-24773465; E-mail: salvatore.campanella@chu-brugmann.be

**Abstract** — **Aims:** While the relationship between chronic exposure to alcohol and neurobiological damage is well established, deleterious brain effects of binge drinking in youths have only recently been studied. **Methods:** Narrative review of studies of brain disturbances associated with binge drinking as assessed by neuroimaging (EEG and IRMf techniques in particular) in adolescent drinkers. **Results:** Some major points still deserved to be investigated; directions for future research are suggested. **Conclusions:** Information and prevention programs should emphasize that binge drinking is not just inoffensive social fun, but if carried on, may contribute to the onset of cerebral disturbances possibly leading to alcohol dependence later in life.

### INTRODUCTION

The practice of drinking to intoxication (i.e. binge drinking) has become the peer norm among some groups of young people, and alcohol stands as the central favorite in a repertoire of psychoactive substances employed to facilitate pleasure and the enjoyment of time out with friends (Johnston *et al.*, 2009). It has been known for a long time that, because of alcohol neurotoxicity, alcohol abuse leads to deleterious effects on the central nervous system, such as brain atrophy and/or dysfunction (e.g. Nicolas *et al.*, 1997). Until recently, most research on brain damage from alcohol drinking has concentrated on heavy long-term use in adults (usually around age 45). A few studies focused on the effects of alcohol use disorders (AUDs) in adolescents and young adults (of 13–24 years). AUDs in 13- to 21-year age group have been linked to abnormal thalamic and putamen volumes and decreased gray matter density (Fein *et al.*, 2013), reductions in bilateral hippocampal (De Bellis *et al.*, 2000) and in prefrontal lobe volumes (De Bellis *et al.*, 2005), to increased functional anisotropy (FA), decreased mean diffusivity in the corpus callosum (De Bellis *et al.*, 2008), and to increased parietal activity during a spatial memory task (Tapert *et al.*, 2004). Researchers in recent years have focused on the effects of binge drinking patterns in younger individuals. It is known from animal studies that (a) during maturation (which is typically not completed until up to 25 years in humans), the brain is particularly sensitive to the effects of alcohol (e.g. Baron *et al.*, 2005) and (b) the repeated withdrawals due to alternation between periods of alcohol intoxication and abstinence may be deleterious for brain function due to excitotoxic cell death (Obernier *et al.*, 2002; Bijl *et al.*, 2007). Thus the binge drinking population seems to be at risk of brain damage.

At the *structural level*, binge drinking has been linked to compromised white matter (WM) in frontal, cerebellar, temporal and parietal regions (McQueeney *et al.*, 2009), reduced WM integrity in both association fiber pathways as well as in the corona radiata (Jacobus *et al.*, 2009), smaller cerebellar volumes (Lisdahl *et al.*, 2013) and changes in cortical thickness (Squeglia *et al.*, 2012a). These structural changes may account for the discovery by Courtney and Polich (2010) of alterations of delta and fast-beta activity among binge drinkers during electroencephalographic passive recording (EEG).

On the other hand, at the functional level, event-related potentials (ERPs) and functional magnetic resonance imaging (fMRI) recorded during diverse cognitive tasks showed that binge drinking is associated with impaired neural processes. Using ERPs, Ehlers *et al.* (2007) showed a decrease in the P3a component latency during a facial discrimination task, a component linked to automatic attention processing of stimulus deviance (see Polich and Criado, 2006). Crego *et al.* (2009, 2010) reported anomalies of the N2, the P3 and the late positive (LP) components combined with hypoactivation of the right anterior prefrontal cortex in binge drinkers during a visual working memory task, possibly reflecting some impairment of working memory processes. Similar to López-Caneda *et al.* (2013), they also demonstrated increased P3b amplitudes in a simple visual oddball task (Crego *et al.*, 2012), suggesting anomalies in neural processes mediating attention processing. Maurage *et al.* (2009) showed diminished activity during an emotional auditory task, indexed by delayed latencies for the P1, the N2 and the P3b components and delayed P3 latencies and impairments related to earlier processes (P100, N100, N170, P2, N2b) during a facial detection task (Maurage *et al.*, 2012). Finally, López-Caneda *et al.* (2012) reported altered Go- and NoGo-P3 components during response execution and inhibition which may represent a neural antecedent of difficulties in cognitive control.

fMRI demonstrated in binge drinkers increased frontal and parietal activations and decreased occipito-hippocampal activation during verbal encoding (Schweinsburg *et al.*, 2010, 2011), thereby indicating greater engagement of working memory systems during encoding and disadvantaged processing of novel verbal information. Binge drinkers also display abnormal patterns of activation during working memory tasks (Squeglia *et al.*, 2011; Campanella *et al.*, 2013). Finally, Xiao *et al.* (2013) showed higher activity in the left amygdala and insula during a decision-making task in binge drinkers, suggesting alterations of the neural circuitry implicated in the execution of emotional and incentive-related behaviors.

### PRE-EXISTING OR ALCOHOL-INDUCED DEFICITS?

While all of these studies undeniably describe cerebral aberrations associated with binge drinking, the majority of them

either carried out only one testing session or followed binge drinkers without controlling for past alcohol consumption, which makes it difficult to be certain of the cause of the deficits. To our knowledge, only three studies explored this question by performing a test–retest study on groups of young individuals with no past alcohol consumption and with comparable baseline results on the tasks proposed. *Squeglia et al.* (2009, 2012b) showed that after 3 years of heavy alcohol use, adolescents exhibited lower visuospatial memory and sustained attention performances as well as abnormal frontal, parietal and occipital activation during a visual working memory task, compared with controls. Using ERPs, *Maurage et al.* (2009) showed delayed latencies of P100, N200 and P300 components only after 9 months in students who had initiated binge drinking habits when compared with students who had persisted in very low alcohol consumption. These few studies indicated that heavy and binge drinking can lead to marked cerebral dysfunction, in the absence of any pre-existing cerebral impairment.

#### EFFECT OF DRINKING PATTERN

One could wonder whether the deficits identified are due to the particular binge drinking pattern (i.e. intense but episodic alcohol consumption episodes) or to the global heavy alcohol intake, independent of the rate and frequency of alcohol absorption. As almost all of the studies cited above only compared binge drinkers with control non-drinkers or very low drinkers, they did not differentiate impairments due to cumulative total alcohol intake versus a specific binge pattern. *Maurage et al.* (2012) explored this hypothesis by comparing the results in groups with different drinking pattern in quantity and frequency, in a simple ERP cognitive task. Where there was a similar cumulative consumption (15–29 doses per week), binge drinkers (i.e. concentrating their weekly 15–59 doses to 2–3 drinking occasions) presented stronger cerebral impairments than daily drinkers (i.e. spreading the weekly 15–59 doses on 5–7 occasions). Similarly, in their recent work that explored alcohol-related bias in binge drinkers, *Petit et al.* (submitted) showed that among the different drinking characteristics of the binge drinkers studied (i.e. the number of alcohol doses per week, the number of alcohol doses per occasion, the number of occasion of drinking per week), the best predictor of the alcohol-related processing bias was the number of alcohol doses consumed on each occasion. Finally, in their study, *Campanella et al.* (2013) identified a specific link between the effects they observed in binge drinkers (i.e. higher activity in the dorsomedial prefrontal cortex) and the number of alcohol doses consumed per occasion. These latter observations indicate that, apart from the traditional AUDs, and the different patterns of misuse of alcohol (as risky, harmful, hazardous alcohol use), the repeated alternation between intoxication and withdrawal appears to have the more deleterious consequences.

#### BINGE DRINKING COMPARED CHRONIC ALCOHOL MISUSE

There are two important observations from these studies on binge drinking. Firstly, despite the broad variety of neuronal mechanisms and structures that they explored, it is evident that

binge drinking is not only associated with damages or deficits, but that these anomalies mirror those observed in alcoholism. Reduced cerebellar volumes, WM integrity and abnormal cortical thickness observed in binge drinkers (*Jacobus et al.*, 2009; *McQueeny et al.*, 2009; *Squeglia et al.*, 2012a; *Lisdahl et al.*, 2013) align volume deficits, FA diminutions and aberrant thickness observed in adults with AUDs (e.g. *Sullivan et al.*, 2000; *Momenan et al.*, 2012). Abnormal central nervous system neuroelectric activity (*Courtney and Polich*, 2010) seen in binge drinkers refers to abnormalities in EEG profile shown alcoholics (e.g. *Bauer*, 2001). Defects in the LPC and the P3 components of the ERP (*Ehlers et al.*, 2007; *Crego et al.*, 2009, 2010, 2012; *Maurage et al.*, 2009, 2012, 2013; *López-Caneda et al.*, 2012) and in components associated with earlier cognitive processes in bingers (*Crego et al.*, 2009; *Maurage et al.*, 2009, 2012; *Petit et al.*, 2012a) are in line with changes in both the amplitude and latency of the LPC and the P300 component (e.g. *Brecher et al.*, 1987; *Porjesz and Begleiter*, 2003; *George et al.*, 2004; *Easdon et al.*, 2005) and with low-level cognitive function impairments observed in chronic alcoholism (e.g. *Nicolas et al.*, 1997; *Verma et al.*, 2006; *Maurage et al.*, 2007; *Fein et al.*, 2009). Aberrant patterns of brain activations during verbal encoding (*Schweinsburg et al.*, 2010, 2011), working memory (*Crego et al.*, 2010; *Squeglia et al.*, 2011; *Campanella et al.*, 2013) and decision-making tasks (*Xiao et al.*, 2013) observed in binge drinkers align poor verbal learning capacities evidenced in chronic heavy drinkers (for a review, see *Grant*, 1987), abnormal brain activation during working memory task observed in adolescents with AUDs (*Tapert et al.*, 2004) and decision-making deficits linked to dysfunctional brain activity reported in alcoholics (*Bechara et al.*, 2001), respectively. Secondly, these studies also show that the similarities invariably observed in binge drinkers are either less serious than those observed in chronic alcohol misuse or are detected by neuroimaging tools but remain unexpressed at a behavioral level. Indeed, if a few studies (*Squeglia et al.*, 2011, 2012a; *Xiao et al.*, 2013) linked binge drinking with decreased performance, most of the others (*Ehlers et al.*, 2007; *Crego et al.*, 2009, 2010, 2012; *Maurage et al.*, 2009, 2012; *Schweinsburg et al.*, 2010, 2011; *López-Caneda et al.*, 2012, 2013; *Petit et al.*, 2012a, 2013, submitted) found neurophysiological differences without any significant behavioral modification. These results emphasize the necessity of using neuroimaging techniques to correctly estimate the actual level of impairment that could go unnoticed at the behavioral level, in a population of binge drinkers, in which abnormalities are not as marked as in pathological populations (*Maurage et al.*, 2009; *Campanella et al.*, 2013) (Fig. 1).

#### IS BINGE DRINKING A PATHWAY TO CHRONIC ALCOHOLISM?

Although less obvious, binge drinkers appear to present with the same pattern of impairments as alcohol-dependent individuals, since the deficits concern the same cognitive functions. The similarities in brain alterations between adolescent binge drinkers and adult alcohol dependents have led some authors (e.g. *Wagner and Anthony*, 2002; *McCarthy et al.*, 2004; *Enoch*, 2006) to suggest the ‘continuum hypothesis’, in which binge drinking and chronic alcohol dependence should be regarded as two stages of the same phenomenon, inducing




Fig. 1. The use of imaging techniques as event-related potentials (ERPs) and functional magnetic resonance imaging (fMRI) allowed measuring the brain's response during each processing steps involved in specific cognitive tasks. fMRI and ERPs data allowed to highlight brain areas that are activated by the presentation of stimuli in binge drinkers when compared with matched controls, and to stress the presence of abnormal electrophysiological processes when participants are for instance confronted with alcohol vs. non-alcohol cues. Such highly sensitive procedures allowed visualizing even minor cognitive restrictions, which may be observable despite normal behavioral performance.

analogous deficits, and not as independent pathologies. This further encouraged the strong suggestion that binge drinking during adolescence could constitute a first step towards the development of alcohol dependence during adulthood (Schulenberg *et al.*, 1996; Tucker *et al.*, 2003). Others have presented contrary data which show that binge drinking is an adolescence's related normative feature (e.g. Gotham *et al.*, 1997) which declines with the increased responsibility due to life transitions such as employment, marriage or parenthood (e.g. Muthén and Muthén, 2000; Wood *et al.*, 2000). Should one therefore be concerned, 'Youth will have its fling'. But what if, for some individuals, this transitional period leaves damage influencing their future? Indeed, knowing that current epidemiological studies have suggested that binge drinking in youths is associated with an increased risk of alcohol abuse/dependence in adulthood (Chassin *et al.*, 2002; Bonomo *et al.*, 2004; McCarty *et al.*, 2004; Viner and Taylor, 2007), it is reasonable to believe that some deficits and/or neurobiological changes that occur during the binge drinking period could play a role in the maintenance of alcohol use and abuse, and could cause difficulties in curtailing consumption leading to long-term alcohol problems (e.g. Hiller-Sturmhofel and Swartzwelder, 2004; King *et al.*, 2006; Haller *et al.*, 2010). However, while this idea sounds relevant, as it has not been explored in depth, the causal relationship between binge drinking deficits and the development of alcohol dependence remains unclear. To elucidate the question of the possible role of such emerging brain abnormalities in binge drinking as risk factors to chronic

alcoholism, future research should focus on two objectives. The first must be to investigate the similarities that exist between the neurocognitive deficits and/or abnormalities detectable in binge drinkers and those observed in adult chronic alcoholics *by focusing specifically* on those that have shown to play a role in the emergence and/or the maintenance of persistent drinking habits in chronic alcoholics. In this regard, as impaired inhibitory control has been identified as a risk factor for substance abuse (Porjesz *et al.*, 2005), López-Caneda *et al.* (2012) focused on analyzing the inhibitory capacity of young binge drinkers. Neurophysiological aberrations indexing impaired inhibition capacities were evident in young binge drinkers which suggested that these alterations could constitute a risk factor for developing alcohol dependence in these individuals. Similarly Petit *et al.* (2012a,b, 2013, submitted) identified a differential electrophysiological processing of alcohol-related stimuli in binge drinkers, a feature demonstrated in alcoholism (e.g. Herrmann, 2000; Namkoong, 2004) and overall, proposed to play a role in the maintenance of drug dependence (Field and Cox, 2008) and considered to be a risk marker for alcohol dependence (Bartholow *et al.*, 2007, 2010). The second important point should be to study the evolution of these deficits and/or abnormalities over time in parallel with the evolution of the consumption of the study population. Although their studies have not been conducted over a lengthy period (bingers were followed for almost 2 years), this allowed López-Caneda *et al.* (2012) and Petit *et al.* (submitted) to demonstrate that the emergence of the electrophysiological

Table 1. Neurophysiological and neuroimaging studies of binge-drinking

Authors (year)	Types of study	Sample characteristics	For longitudinal studies: verification of the absence of difference <i>before starting</i> BD habits	Age (years)	Cognitive task	Neuroimaging tools	Cognitive outcome	Neurophysiological and neuroimaging outcome	Examples of studies and reviews depicting similar defects in alcoholic population
Ehlers <i>et al.</i> (2007)	Cross-sectional	BD ( $n = 30$ ): >5 drinks in 1 occasion during adolescence (before age 18) C ( $n = 36$ ): no BD history during adolescence	NA	18–25	Facial discrimination task	ERP	BD = C	P3 latency: BD < C	Porjesz and Begleiter (2003)
Crego <i>et al.</i> (2009)	Cross-sectional	BD ( $n = 42$ ): ≥6 drinks on 1 occasion at a speed of ≥3 drinks/h C ( $n = 53$ ): ≤6 drinks on 1 occasion at a speed of ≤3 drinks/h	NA	18–20	Visual working memory task	ERP	BD = C	N2 amplitudes: BD > C P3 amplitudes: BD ≠ C	Porjesz and Begleiter (2003), Fein <i>et al.</i> (2009) and Verma <i>et al.</i> (2006)
Jacobus <i>et al.</i> (2009)	Cross-sectional	BD ( $n = 28$ ): history of at least 1 episode of ≥5 (M) or 4 (F) drinks on one occasion C ( $n = 14$ ): very limited if any substance use history	NA	16–19	None	DTI	NA	WM indexed by FA: BD < C	Pfefferbaum <i>et al.</i> (2006)
McQueeney <i>et al.</i> (2009)	Cross-sectional	BD ( $n = 14$ ): ≥5 (M) or 4 (F) drinks in 1 sitting during the last 3 months C ( $n = 14$ ): no history of BD episodes	NA	16–19	None	DTI	NA	WM integrity (indexed by FA): BD < C	Pfefferbaum <i>et al.</i> (2006)
Maurage <i>et al.</i> (2009)	Cohort (prospective) – over 9 months	BD ( $n = 18$ ): 12.52 DPO; 2.33 NOW; 35 DPW C ( $n = 18$ ): no BD	Yes	T1: 18–19	Emotional auditory discrimination task	ERP	T1 and T2: BD = C	T1: BD = C T2: P1, N2, P3b latencies: BD > C	Maurage <i>et al.</i> (2008) and Kathmann <i>et al.</i> (1996)
Courtney and Polich (2010)	Cross-sectional	Low-BD ( $n = 32$ ): 5/4–7/6 drinks in within 2 h on >1 occasion within the past 6 months High-BD ( $n = 32$ ): ≥10 drinks within 2 h on >1 occasion within the past 6 months Controls ( $n = 32$ ): >1 to 5/4 drinks within 2 h on >1 occasion within the past 6 months	NA	18–22	None	EEG	NA	Spectral power in the delta (0–4 Hz) and fast-beta (20–35 Hz) bands : C = low-BD < high-BD	Bauer (2001)
Crego <i>et al.</i> (2010)	Cross-sectional	BD ( $n = 42$ ): ≥6 drinks on 1 occasion at a speed of ≥3 drinks/h 1 or more times per month. C ( $n = 53$ ): ≤6 drinks on 1 occasion at a speed of ≤3 drinks/h	NA	18–20	Visual working memory task	ERP and eLORETA	BD = C	LPC amplitude and aPFC: BD < C	Tapert <i>et al.</i> (2004) and Brecher <i>et al.</i> (1987)

Continued

Table 1. Continued

Authors (year)	Types of study	Sample characteristics	For longitudinal studies: verification of the absence of difference <i>before starting</i> BD habits	Age (years)	Cognitive task	Neuroimaging tools	Cognitive outcome	Neurophysiological and neuroimaging outcome	Examples of studies and reviews depicting similar defects in alcoholic population
Schweinsburg <i>et al.</i> (2010)	Cross-sectional	BD ( $n = 12$ ): $\geq 5$ (M) or 4 (F) drinks in 1 occasion C ( $n = 15$ ): $\leq 5$ (M) or 4 (F) drinks in 1 occasion	NA	16–18	Verbal encoding task	fMRI	BD = C	BOLD response during novel verbal encoding: BD > C	Tapert <i>et al.</i> (2004)
Schweinsburg <i>et al.</i> (2011)	Cross-sectional	BD ( $n = 16$ ): $\geq 5$ (M) or 4 (F) drinks in 1 occasion during the last 3 months C ( $n = 22$ ): very limited alcohol use experience	NA	16–18	Verbal encoding task	fMRI	BD = C	Frontal BOLD response during novel verbal encoding: BD $\neq$ C	Tapert <i>et al.</i> (2004)
Squeglia <i>et al.</i> (2011)	Cross-sectional	BD ( $n = 40$ ) C ( $n = 55$ )	NA	16–19	Neuropsychological battery + spatial working memory task	fMRI	Working memory performance: BD $\neq$ C	Neural activation: BD $\neq$ C	Tapert <i>et al.</i> (2004) and Ambrose <i>et al.</i> (2001)
Crego <i>et al.</i> (2012)	Cross-sectional	BD ( $n = 32$ ): $\geq 6$ drinks on 1 occasion at a speed of $\geq 3$ drinks/h $\geq 1$ times per month C ( $n = 53$ ): $\leq 6$ drinks on 1 occasion at a speed of $\leq 3$ drinks/h	NA	18–20	Visual oddball task	ERP	BD = C	P3b amplitudes: BD > C	Porjesz and Begleiter (2003)
Lopez-Caneda <i>et al.</i> (2012)	Cohort (prospective) – over 2 years	BD ( $n = 23$ ): $\geq 6$ drinks on 1 occasion at a speed of $\geq 3$ drinks/h $\geq 1$ times per month or $\geq 6$ drinks on 1 occasion $\geq 1$ times per week C ( $n = 25$ ): $\leq 6$ drinks on 1 occasion	No	T1: 18–19	Go/NoGo task	ERP and eLORETA	T1 and T2: BD = C	NoGo-P3 amplitude: T1: BD = C; T2: BD > C Go-P3 amplitude: T1 and T2: BD > C Activation in rIFC: T1: BD = C; T2: BD > C	Cohen <i>et al.</i> (1997) and Kamarajan <i>et al.</i> (2004)
Maurage <i>et al.</i> (2012)	Cross-sectional	BD1 ( $n = 20$ ): 5–12 ADO; 2–3 DOW; ADH >3; 15–29 ADW BD2 ( $n = 20$ ): ADO >10; 3–4 DOW; ADH >3; ADW >30 C ( $n = 20$ ): ADO <2; DOW <0.5; ADH <1; ADW <2	NA	19–24	Visual oddball task (face detection)	ERP	BD1 = BD2 = C	P100, N100, N170/P2, N2b/P3a and P3b amplitudes: BD1 < C; BD2 < C P100, N100 and N2b/P3a latencies: BD2 > C P3b latency: BD1 > C; BD2 > C	Porjesz and Begleiter (2003), Maurage <i>et al.</i> (2007) and Nicolas <i>et al.</i> (1997)
Petit <i>et al.</i> (2012a)	Cross-sectional	BD ( $n = 18$ ): $\geq 6$ drinks on 1 occasion at a speed of $\geq 3$ drinks/h C ( $n = 18$ ): $\leq 6$ drinks on 1 occasion at a speed of $\leq 3$ drinks/h	NA	19–25	Alcohol-modified oddball task	ERP	BD = C	P100 amplitude in response to alcohol cues: BD > C	Herrman <i>et al.</i> (2000)
Squeglia <i>et al.</i> (2012a)	Cross-sectional	BD ( $n = 29$ ) C ( $n = 30$ )	NA	16–19	Neuropsychological battery	MRI	Visuospatial, inhibition and attention performances: BD $\neq$ C	Cortical thickness: BD $\neq$ C	Momenan <i>et al.</i> (2012)

Campanella <i>et al.</i> (2013)	Cross-sectional	BD: 9.2 ADO, 3.4 ADH, 2.7 DOW C: 3.7 ADO, 1.5 ADH, 0.5 DOW	NA	18–25	Working memory N-back task	fMRI	BD = C	Activity in the pre-supplementary motor area: BD > C	Moselhy <i>et al.</i> (2001)
Lisdahl <i>et al.</i> (2013)	Cross-sectional	BD ( <i>n</i> = 46) C ( <i>n</i> = 60)	NA	16–19	None	MRI	NA	Cerebellar volumes: BD < C	Sullivan <i>et al.</i> (2000)
Lopez-Caneda <i>et al.</i> (2013)	Cohort (prospective) – over 2 years	BD ( <i>n</i> = 26): ≥6 drinks on 1 occasion at a speed of ≥3 drinks/h ≥1 times per month or ≥6 drinks on 1 occasion ≥1 times per week C ( <i>n</i> = 31): ≤6 drinks on 1 occasion	No	T1: 18–19	Visual oddball task	ERP	T1 and T2: BD = C	P3b amplitudes: T1: BD > C T2: BD > + C	Porjesz and Begleiter (2003)
Petit <i>et al.</i> (2013)	Cross-sectional	BD ( <i>n</i> = 29): ≥6 drinks on 1 occasion at a speed of ≥3 drinks/h C ( <i>n</i> = 27): ≤6 drinks on 1 occasion at a speed of ≤3 drinks/h	NA	18–27	Alcohol-modified oddball task	ERP	BD = C	P3 amplitudes to alcohol cues: BD > C	Herrmann <i>et al.</i> (2000)
Petit <i>et al.</i> (submitted)	Cohort (prospective) – over 1 year	BD ( <i>n</i> = 15): ≥6 drinks on 1 occasion at a speed of ≥3 drinks/h C ( <i>n</i> = 15): ≤6 drinks on 1 occasion at a speed of ≤3 drinks/h	No	19–25	Alcohol-modified oddball task	ERP	T1 and T2: BD = C	T1: P3 amplitudes to alcohol vs. non-alcohol cues: BD = C T2: P3 amplitudes to alcohol vs. non-alcohol cues: BD ≠ C	Herrmann <i>et al.</i> (2000)
Xiao <i>et al.</i> (2013)	Cross-sectional	BD ( <i>n</i> = 14) C ( <i>n</i> = 14)	NA	16–18	Iowa Gambling Task (IGT)	fMRI	Performance on IGT: BD ≠ C	Neural activity: BD > C	Bechara <i>et al.</i> (2001)

C, controls; BD, binge drinkers; C = B, no significant difference between control and binge group; BD ≠ C, binge drinkers significantly differed from controls; BD > + C, more pronounced difference in BD vs. controls (here, compared with T1); EEG, electroencephalogram; ERP, event-related potentials; DTI, diffusion tensor imaging; WM, white matter; FA, fractional anisotropy; eLORETA, exact low-resolution brain electromagnetic tomography; LPC, late positive component; A PFC, activation of the right anterior prefrontal cortex; MRI, magnetic resonance imaging; fMRI, functional magnetic resonance imaging; ADO, mean number of alcohol doses per drinking occasion; DOW, mean number of drinking occasions per week; ADH, mean number of alcohol doses per hour (consumption speed); ADW, mean number of alcohol doses per week; rIFC, right Inferior Frontal Cortex.

abnormalities observed was associated with the continuation of binge drinking habits. These preliminary results suggest that a vicious circle could start with the habit of binge drinking. In other words, alterations in cue reactivity or reduced inhibition, two characteristics that the contemporary dual process model theories associate with the development of alcohol abuse (e.g. Stacy and Wiers, 2010), could arise with the emergence of binge drinking habits and then be amplified due to alcohol neurotoxicity. The stronger neuropsychological impairments and their link to neurobiological changes would then reinforce the alcohol consumption, preventing the normal decrease of heavy drinking with maturation and leading to the perpetuation of risky drinking habits and eventually to alcoholism.

### CONCLUSION: STILL A LONG WAY TO GO

The neuroscience approach of binge drinking has already drawn attention to the structural and functional brain damage associated with this practice (Hermens *et al.*, 2012). Research on binge drinking in young people is however still in its infancy and many questions remain unresolved. A first important question is whether the brain alterations observed among binge drinkers are the result of alcohol misuse or whether these changes may be pre-existing. This issue has only been properly assessed in a few studies and still needs to be confirmed and extended in more longitudinal studies, notably in regards of the control of other variables that precede and could also predispose to the development of binge drinking habits, as for example brain modifications related to family history of alcoholism. In their neuroimaging longitudinal study, Squeglia *et al.* (2012b) suggested that some cerebral deficits observed in heavy drinkers could already be present before alcohol misuse and be involved in the onset of alcohol consumption. Also, the earlier Ehlers *et al.*'s study (2007) already mentioned that a positive family history for alcohol dependence acted as a significant covariate in some of their findings on binge drinking while Norman *et al.* (2011) and Wetherill *et al.* (2013) showed in their longitudinal studies that pre-existing abnormalities in neural activity during response inhibition observed in early adolescence were predictors of future alcohol heavy use. Controlled longitudinal studies will be crucial. Finally, longitudinal studies on binge drinkers may reveal some who start to abstain from binge drinking and reversibility of abnormalities should be looked for. Meanwhile, given the data available, there is a case to develop adapt information and prevention programs to emphasize the message that binge drinking is not just inoffensive social fun, but if carried on, may contribute to the onset of cerebral disturbances leading to alcohol dependence later in life, even at a stage at which behavioral manifestation is not yet evident (Campanella *et al.*, 2013). In Table 1, we summarize neurophysiological and neuroimaging studies of binge-drinking. We especially focused on (1) whether these studies reported cognitive impairments in addition to neural mechanisms impairments (2) the similarities of the neurobiological alterations underlined with defects known in chronic alcoholism and (3) whether the alterations are proven to have occurred after binge drinking started rather than possibly be present before.

*Funding* — G.P. (research fellow), P.M. (research associate) and S.C. (research associate) are funded by the Belgian Fund for Scientific Research (F.R.S.-F.N.R.S., Belgium).

*Conflict of interest statement.* None declared.

### REFERENCES

- Ambrose ML, Bowden SC, Whelan G. (2001) Working memory impairments in alcohol-dependent participants without clinical amnesia. *Alcohol Clin Exp Res* **25**:185–91.
- Barron S, White A, Swartzwelder HS *et al.* (2005) Adolescent vulnerabilities to chronic alcohol or nicotine exposure: findings from rodent models. *Alcohol Clin Exp Res* **29**:1720–5.
- Bartholow BD, Henry EA, Lust SA. (2007) Effects of alcohol sensitivity on P3 event-related potential reactivity to alcohol cues. *Psychol Addict Behav* **21**:555–63.
- Bartholow BD, Lust SA, Traggesser SL. (2010) Specificity of P3 event-related potential reactivity to alcohol cues in individuals low in alcohol sensitivity. *Psychol Addict Behav* **24**:220–8.
- Bauer LO. (2001) CNS recovery from cocaine, cocaine and alcohol, or opioid dependence: a P300 study. *Clin Neurophysiol* **112**:1508–15.
- Bechara A, Dolan S, Denburg N *et al.* (2001) Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. *Neuropsychologia* **39**:376–89.
- Bijl S, de Bruin EA, Böcker KE *et al.* (2007) Effects of chronic drinking on verb generation: an event related potential study. *Hum Psychopharmacol* **22**:157–66.
- Bonomo YA, Boves G, Coffey C *et al.* (2004) Teenage drinking and the onset of alcohol dependence: a cohort study over seven years. *Addiction* **99**:1520–8.
- Brecher M, Porjesz B, Begleiter H. (1987) Late positive component amplitude in schizophrenics and alcoholics in two different paradigms. *Biol Psychiatry* **22**:848–56.
- Campanella S, Peigneux P, Petit G *et al.* (2013) Increased cortical activity in binge drinkers during working memory task: a preliminary assessment through a functional magnetic resonance imaging study. *PLoS One* **8**:e62260.
- Chassin L, Pitts SC, Prost J. (2002) Binge drinking trajectories from adolescence to emerging adulthood in a high-risk sample: predictors and substance abuse outcomes. *J Consult Clin Psychol* **70**:67–78.
- Cohen HL, Porjesz B, Begleiter H *et al.* (1997) Neurophysiological correlates of response production and inhibition in alcoholics. *Alcohol Clin Exp Res* **21**:1398–406.
- Courtney KE, Polich J. (2010) Binge drinking effects on EEG in young adult humans. *Int J Environ Res Public Health* **7**:2325–36.
- Crego A, Holguín SR, Parada M *et al.* (2009) Binge drinking affects attentional and visual working memory processing in young university students. *Alcohol Clin Exp Res* **33**:1870–9.
- Crego A, Rodríguez-Holguín S, Parada M *et al.* (2010) Reduced anterior prefrontal cortex activation in young binge drinkers during a visual working memory task. *Drug Alcohol Depend* **109**:45–56.
- Crego A, Cadaveira F, Parada M *et al.* (2012) Increased amplitude of P3 event-related potential in young binge drinkers. *Alcohol* **46**:415–25.
- De Bellis MD, Clark DB, Beers SR *et al.* (2000) Hippocampal volume in adolescent-onset alcohol use disorders. *Am J Psychiatry* **157**:737–44.
- De Bellis MD, Narasimhan A, Thatcher DL *et al.* (2005) Prefrontal cortex thalamus and cerebellar volumes in adolescents and young adults with adolescent-onset alcohol use disorders and comorbid mental disorders. *Alcohol Clin Exp Res* **29**:1590–600.
- De Bellis MD, Van Voorhees E, Hooper SR *et al.* (2008) Diffusion tensor measures of the corpus callosum in adolescents with adolescent onset alcohol use disorders. *Alcohol Clin Exp Res* **32**:395–404.
- Easdon C, Izenberg A, Armilio ML *et al.* (2005) Alcohol consumption impairs stimulus- and error-related processing during a Go/No-Go Task. *Brain Res Cogn Brain Res* **25**:873–83.
- Ehlers CL, Phillips E, Finnerman G *et al.* (2007) P3 components and adolescent binge drinking in Southwest California Indians. *Neurotoxicol Teratol* **29**:153–63.
- Enoch MA. (2006) Genetic and environmental influences on the development of alcoholism: resilience vs risk. *Ann N Y Acad Sci* **1094**:193–201.
- Fein G, Shimotsu R, Chu R *et al.* (2009) Parietal gray matter volume loss is related to spatial processing deficits in long-term abstinent alcoholic men. *Alcohol Clin Exp Res* **33**:1806–14.

- Fein G, Greenstein D, Cardenas VA *et al.* (2013) Cortical and subcortical volumes in adolescents with alcohol dependence but without substance or psychiatric comorbidities. *Psychiatry Res* **214**:1–8.
- Field M, Cox WM. (2008) Attentional bias in addictive behaviors: a review of its development, causes, and consequences. *Drug Alcohol Depend* **97**:1–20.
- George MR, Potts G, Kothman D *et al.* (2004) Frontal deficits in alcoholism: an ERP study. *Brain Cogn* **54**:245–7.
- Gotham HJ, Sher KJ, Wood PK. (1997) Predicting stability and change in frequency of intoxication from the college years to beyond: individual-difference and role transition variables. *J Abnorm Psychol* **106**:619–29.
- Grant I. (1987) Alcohol and the brain: neuropsychological correlates. *J Consult Clin Psychol* **55**:310–24.
- Haller M, Handley E, Chassin L *et al.* (2010) Developmental cascades: linking adolescent substance use, affiliation with substance use promoting peers, and academic achievement to adult substance use disorders. *Dev Psychopathol* **22**:899.
- Hermens DF, Lagopoulos J, Tobias-Webb J *et al.* (2012) Pathways to alcohol-induced brain impairment in young people: a review. *Cortex* **49**:3–17.
- Herrmann MJ, Weijers HG, Wiesbeck GA *et al.* (2000) Event-related potentials and cue-reactivity in alcoholism. *Alcohol Clin Exp Res* **24**:1724–9.
- Hiller-Sturmhofel S, Swartzwelder HS. (2004) Alcohol's effects on the adolescent brain: what can be learned from animal models. *Alcohol Res Health* **28**:213.
- Jacobus J, McQueeney T, Bava S *et al.* (2009) White matter integrity in adolescents with histories of marijuana use and binge drinking. *Neurotoxicol Teratol* **31**:349–55.
- Johnston LD, O'Malley PM, Bachman JG *et al.* (2009) Monitoring the future national results on adolescent drug use: overview of key findings. In *Abuse*. Bethesda, MD: National Institute on Drug Abuse. NIH Publication No. 09–7401.
- Kamarajan C, Porjesz B, Jones KA *et al.* (2004) The role of brain oscillations as functional correlates of cognitive systems: a study of frontal inhibitory control in alcoholism. *Int J Psychophysiol* **51**:155–80.
- Kathmann N, Soyka M, Bickel R *et al.* (1996) ERP changes in alcoholics with and without alcohol psychosis. *Biol Psychiatry* **39**:873–81.
- King A, Munisamy G, de Wit H *et al.* (2006) Attenuated cortisol response to alcohol in heavy social drinkers. *Int J Psychophysiol* **59**:203–9.
- Lisdahl KM, Thayer R, Squeglia LM *et al.* (2013) Recent binge drinking predicts smaller cerebellar volumes in adolescents. *Psychiatry Res* **211**:17–23.
- López-Caneda E, Cadaveira F, Crego A *et al.* (2012) Hyperactivation of right inferior frontal cortex in young binge drinkers during response inhibition: a follow-up study. *Addiction* **107**:1796–808.
- López-Caneda E, Cadaveira F, Crego A *et al.* (2013) Effects of a persistent binge drinking pattern of alcohol consumption in young people: a follow-up study using event-related potentials. *Alcohol Alcohol* **48**:464–71.
- Maurage P, Philippot P, Verbanck P *et al.* (2007) Is the P300 deficit in alcoholism associated with early visual impairments (P100 N170)? An oddball paradigm. *Clin Neurophysiol* **118**:633–44.
- Maurage P, Philippot P, Joassin F *et al.* (2008) The auditory-visual integration of anger stimuli is impaired in alcoholism: an ERP study. *J Psychiatry Neurosci* **33**:111–22.
- Maurage P, Pesenti M, Philippot P *et al.* (2009) Latent deleterious effects of binge drinking over a short period of time revealed only by electrophysiological measures. *J Psychiatry Neurosci* **34**:111–8.
- Maurage P, Joassin F, Speth A *et al.* (2012) Cerebral effects of binge drinking: respective influences of global alcohol intake and consumption pattern. *Clin Neurophysiol* **123**:892–901.
- McCarty CA, Ebel BE, Garrison MM *et al.* (2004) Continuity of binge and harmful drinking from late adolescence to early adulthood. *Pediatrics* **114**:714–9.
- McQueeney T, Schweinsburg BC, Schweinsburg AD *et al.* (2009) Altered white matter integrity in adolescent binge drinkers. *Alcohol Clin Exp Res* **33**:1278–85.
- Momenan R, Steckler LE, Saad ZS *et al.* (2012) Effects of alcohol dependence on cortical thickness as determined by magnetic resonance imaging. *Psychiatry Res* **204**:101–11.
- Moselhy HF, Georgiou G, Kahn A. (2001) Frontal lobe changes in alcoholism: a review of the literature. *Alcohol Alcohol* **36**:357–68.
- Muthén BO, Muthén LK. (2000) The development of heavy drinking and alcohol-related problems from ages 18 to 37 in a US national sample. *J Stud Alcohol Drugs* **61**:290.
- Namkoong K, Lee E, Lee CH *et al.* (2004) Increased P3 amplitudes induced by alcohol-related pictures in patients with alcohol dependence. *Alcohol Clin Exp Res* **28**:1317–23.
- Nicolas JM, Estruch R, Salamero M *et al.* (1997) Brain impairment in well-nourished chronic alcoholics is related to ethanol intake. *Ann Neurol* **41**:590–8.
- Norman AL, Pulido C, Squeglia LM *et al.* (2011) Neural activation during inhibition predicts initiation of substance use in adolescence. *Drug Alcohol Depend* **119**:216–23.
- Obernier JA, White AM, Swartzwelder HS *et al.* (2002) Cognitive deficits and CNS damage after a 4-day binge ethanol exposure in rats. *Pharmacol Biochem Behav* **72**:521–32.
- Petit G, Kornreich C, Maurage P *et al.* (2012a) Early attentional modulation by alcohol-related cues in young binge drinkers: an event-related potentials study. *Clin Neurophysiol* **123**:925–36.
- Petit G, Kornreich C, Noël X *et al.* (2012b) Alcohol-related context modulates performance of social drinkers in a visual Go/No-Go task: a preliminary assessment of event-related potentials. *PLoS One* **7**:e37466.
- Petit G, Kornreich C, Verbanck P *et al.* (2013) Gender differences in reactivity to alcohol cues in binge drinkers: a preliminary assessment of event-related potentials. *Psychiatry Res* **209**:494–503.
- Petit G, Kornreich C, Noël X *et al.* (Submitted) Continuity of binge drinking leads to differential electrophysiological processing of alcohol-related stimuli: a follow up study.
- Pfefferbaum A, Adalsteinsson E, Sullivan EV. (2006) Supratentorial profile of white matter microstructural integrity in recovering alcoholic men and women. *Biol Psychiatry* **59**:364–72.
- Polich J, Criado JR. (2006) Neuropsychology and neuropharmacology of P3a and P3b. *Int J Psychophysiol* **60**:172–85.
- Porjesz B, Begleiter H. (2003) Alcoholism and human electrophysiology. *Alcohol Res Health* **27**:153–60.
- Porjesz B, Rangaswamy M, Kamarajan C *et al.* (2005) The utility of neurophysiological markers in the study of alcoholism. *Clin Neurophysiol* **116**:993–1018.
- Schulenberg J, O'Malley PM, Bachman JG *et al.* (1996) Getting drunk and growing up: trajectories of frequent binge drinking during the transition to young adulthood. *J Stud Alcohol* **57**:289–304.
- Schweinsburg AD, McQueeney T, Nagel BJ *et al.* (2010) A preliminary study of functional magnetic resonance imaging response during verbal encoding among adolescent binge drinkers. *Alcohol* **44**:111–7.
- Schweinsburg AD, Schweinsburg BC, Nagel BJ *et al.* (2011) Neural correlates of verbal learning in adolescent alcohol and marijuana users. *Addiction* **106**:564–73.
- Squeglia LM, Spadoni AD, Infante MA *et al.* (2009) Initiating moderate to heavy alcohol use predicts changes in neuropsychological functioning for adolescent girls and boys. *Psychol Addict Behav* **23**:715–22.
- Squeglia LM, Schweinsburg AD, Pulido C *et al.* (2011) Adolescent binge drinking linked to abnormal spatial working memory brain activation: differential gender effects. *Alcohol Clin Exp Res* **35**:1831–41.
- Squeglia LM, Sorg SF, Schweinsburg AD *et al.* (2012a) Binge drinking differentially affects adolescent male and female brain morphology. *Psychopharmacology* **220**:529–39.
- Squeglia LM, Pulido C, Wetherill RR *et al.* (2012b) Brain response to working memory over three years of adolescence: influence of initiating heavy drinking. *J Stud Alcohol Drugs* **73**:749–60.
- Stacy AW, Wiers RW. (2010) Implicit cognition and addiction: a tool for explaining paradoxical behavior. *Ann Rev Clin Psychol* **6**:551–75.
- Sullivan EV, Deshmukh A, Desmond JE *et al.* (2000) Cerebellar volume decline in normal aging, alcoholism, and Korsakoff's syndrome: relation to ataxia. *Neuropsychology* **14**:341–52.
- Tapert SF, Schweinsburg AD, Barlett VC *et al.* (2004) Blood oxygen level dependent response and spatial working memory in adolescents with alcohol use disorders. *Alcohol Clin Exp Res* **28**:1577–86.
- Tucker JS, Orlando M, Ellickson PL. (2003) Patterns and correlates of binge drinking trajectories from early adolescence to young adulthood. *Health Psychol* **22**:79–87.



- Verma RK, Panda NK, Basu D *et al.* (2006) Audiovestibular dysfunction in alcohol dependence. Are we worried? *Am J Otolaryngol* **27**:225–8.
- Viner RM, Taylor B. (2007) Adult outcomes of binge drinking in adolescence: findings from a UK national birth cohort. *J Epidemiol Community Health* **61**:902–7.
- Wagner FA, Anthony JC. (2002) From first drug use to drug dependence. Developmental periods of risk for dependence upon marijuana cocaine and alcohol. *Neuropsychopharmacology* **26**:479–88.
- Wetherill RR, Squeglia LM, Yang TT *et al.* (2013) A longitudinal examination of adolescent response inhibition: neural differences before and after the initiation of heavy drinking. *Psychopharmacology* (in press).
- Wood MD, Sher KJ, McGowan AK. (2000) Collegiate alcohol involvement and role attainment in early adulthood: findings from a prospective high-risk study. *J Stud Alcohol* **61**: 278–89.
- Xiao L, Bechara A, Gong Q *et al.* (2013) Abnormal affective decision making revealed in adolescent binge drinkers using a functional magnetic resonance imaging study. *Psychol Addict Behav* **27**:443–54.