

## Bioactive Phenolic Constituents from the Culms of *Phyllostachys bambusoides*

Hyun Jung Kim<sup>1</sup>, Min-Ah Kang<sup>2</sup>, So-Hyun Kim<sup>3</sup>, Soon-Ho Yim<sup>4</sup>, and Ik-Soo Lee\*

College of Pharmacy and Research Institute of Drug Development, Chonnam National University,  
Gwangju 500-757, Republic of Korea

<sup>1</sup>Research Center for Resistant Cells, School of Medicine, Chosun University, Gwangju 501-759, Republic of Korea

<sup>2</sup>Kolmar Korea Co., Ltd., 618-3, Sinjeong-ri, Jeonui-myeon, Yeongi-gun, Chungcheongnam-do 339-853, Republic of Korea

<sup>3</sup>Biobank of Chonnam National University Hwasun Hospital, Ilsim-ri, Hwasun-eup, Hwasun-gun,  
Jeollanam-do 519-763, Republic of Korea

<sup>4</sup>Gist Technology Institute, Gwangju Institute of Science and Technology, Gwangju 500-712,  
Republic of Korea

**Abstract** – In our search for bioactive phenolics from plants, the culms of *Phyllostachys bambusoides* has been selected for investigation of anti-cariogenic and 1,1-diphenylpicrylhydrazyl (DPPH) radical scavenging agents based on the initial screening results. Fractionation process of *n*-hexane and CHCl<sub>3</sub> extracts afforded four phenolic constituents, ferulic acid (**1**), vanillin (**2**), coniferaldehyde (**3**), and coniferyl alcohol (**4**) as guided by their DPPH free radical scavenging activities. Additionally, activity-guided fractionation of EtOAc extract with anti-cariogenic activity has resulted in the isolation of coniferaldehyde (**3**), 2,6-dimethoxy-*p*-benzoquinone (**5**), *p*-methoxy-cinnamic acid (**6**), (±)-balanophonin (**7**), and 6-methoxychromanone (**8**). The structures of **1** - **8** were determined by spectroscopic data interpretation, and also by comparison of their data with the published values. Phenolic compounds **1** - **4** exhibited similar DPPH radical scavenging activities compared with the synthetic antioxidant, butylated hydroxytoluene (BHT), and compounds **3** and **5** - **8** showed significant antibacterial activity against cariogenic oral streptococci, *Streptococcus mutans* and *S. sobrinus*.

**Keywords** – *Phyllostachys bambusoides*, Gramineae, phenolics, anti-cariogenic, DPPH radical scavenger

### Introduction

The *Phyllostachys* species including *P. bambusoides*, *P. nigra* var. *henonis* and *P. pubescens* are woody perennial evergreen plants belonging to the family Gramineae, which are very popular plants in Asia. Their culms were traditionally used as medicinal materials including *Bambusae Caulis* in Liguamen and *Bambusae Caulis* in Taenis and *Bambusae Concretio Salicea*, which have been used as antipyretic, antitussive and antidiuretic agents (Bae, 2000). The dried mass of a secretion from the culms of *P. bambusoides* S. et Z. has been used in traditional Korean medicine, and it was reported to be useful for the clinical treatment of degenerative neuronal disorders (Lee, 1986; Ko *et al.*, 1994). In addition, the culms of *Phyllostachys* were widely used as grain storage and a wrapping material for foods. Several biological activities (Nikaido *et al.*, 1984; Nishina *et al.*, 1991; Cowan, 1999) and phytochemical constituents (Kweon *et al.*, 2001;

Tanaka *et al.*, 2003; Suga *et al.*, 2003) were revealed, however, little is known about its biological evaluation and chemical composition in spite of the fact that the culms of bamboos have been used extensively as a foodstuff.

In the course of screening for bioactive compounds from MeOH extract of the culms of *P. bambusoides*, phenolic compounds **1** - **8** were found to have free radical scavenging effect on DPPH radical or anti-bacterial activities against cariogenic oral streptococci, *Streptococcus mutans* and *S. sobrinus*. This paper deals with the structure elucidation and biological evaluation of the isolated phenolic compounds.

### Experimental

*General Experimental Procedures.* <sup>1</sup>H and <sup>13</sup>C NMR spectra, with tetramethylsilane (TMS) as the internal standard were recorded using a Varian Unity INOVA 500 spectrometer (Varian, Inc., U.S.A.) for 1D and 2D NMR experiments in MeOH-*d*<sub>4</sub> and CDCl<sub>3</sub>. The chemical shifts

\*Author for correspondence  
Tel: +82-62-530-2932; E-mail: islee@chonnam.ac.kr

( $\delta$ ) were expressed in parts per million (ppm) and coupling constants ( $J$ ) were in Hz. Mass spectra were measured on a JMS-700 (Jeol, Japan) and Varian 1200, Platform II (Varian, U.S.A.) spectrometers. IR spectra were obtained on a JASCO FT/IR-300E spectrometer (Jasco Corp., Japan) and UV spectra were recorded on a JASCO V-530 UV/Vis spectrophotometer (Jasco Corp., Japan). Optical rotation was measured on a JASCO DIP-1000 digital polarimeter.

TLC chromatographic analyses were carried out on precoated Silica gel 60 F<sub>254</sub> plates, and RP-18 F<sub>254</sub> plates (Merck). Visualization of the silica gel TLC was performed using 10% (v/v) H<sub>2</sub>SO<sub>4</sub> followed by charring at 110 °C for 10 min. The adsorbent used for column chromatography was silica gel 60/70-230 mesh. The flash chromatography (MPLC) was performed with a Lobar<sup>TM</sup> glass prepacked column (11 mm × 300 mm), and Sephadex LH-20 (Pharmacia Biotech Co., Ltd.) was used for size exclusion chromatography. The gentamicin, gentamicin disc (10 µg), Vitamin C, BHA, and BHT were purchased from Sigma Chemical and BD Biosciences, U.S.A.

**Plant material.** The culms of *Phyllostachys bambusoides* were collected in Damyang, Jeonnam Province, Korea, in 2005 and were pulverized by using a grinder. A voucher specimen has been deposited in the College of Pharmacy, Chonnam National University, Korea.

**Extraction and isolation.** The dried bamboo powder (5 kg) was extracted with 5 L of methanol-water mixture (8 : 2) three times. After filtration, the 80% MeOH extract was combined and concentrated *in vacuo* using rotary evaporator to give a dark green residue (210 g). This 80% MeOH extract was suspended in water (1 L) and partitioned with *n*-hexane, CHCl<sub>3</sub>, EtOAc, *n*-BuOH, and water, successively.

The *n*-hexane extract (5.2 g) was subjected to silica gel column chromatography with a gradient of CHCl<sub>3</sub>-MeOH (400 : 1, 200 : 1, 50 : 1, 20 : 1, 10 : 1) mixture to give six fractions. Fraction 3 (841.1 mg) was column chromatographed on a silica gel (70 - 230 mesh, Merck), eluting with CHCl<sub>3</sub>-MeOH gradient mixture to give further fractions. Subfraction 3 (202.2 mg) was subjected to MPLC (CHCl<sub>3</sub>-MeOH = 200 : 1) to give three subfractions and to yield compound **1** (1.9 mg, yield 0.0009%)

The CHCl<sub>3</sub> soluble extract (7.5 g) was subjected to silica gel column chromatography with a gradient of CH<sub>2</sub>Cl<sub>2</sub>-MeOH (50 : 1, 20 : 1, 10 : 1) mixture to give five fractions. Fraction 1 (95 mg) was column chromatographed on Sephadex LH-20 using 100% MeOH as eluent to provide 5 fractions. Subfraction 2 was subjected to MPLC (CH<sub>2</sub>Cl<sub>2</sub>-MeOH = 100 : 1) to give three subfrac-

tions and compound **2** (2.3 mg, yield 0.0011%). Fraction 3 (207.2 mg) was column chromatographed on silica gel (70 - 230 mesh, Merck), eluting with CH<sub>2</sub>Cl<sub>2</sub>-MeOH gradient mixture to give further four fractions. Among them, subfraction 3 was separated by MPLC using CH<sub>2</sub>Cl<sub>2</sub>-MeOH (80 : 1) to give seven fractions and compound **3** (1.2 mg, yield 0.00057%), and subfraction 5 was subjected to MPLC (CH<sub>2</sub>Cl<sub>2</sub>-MeOH = 60 : 1) to yield compound **4** (2.8 mg, yield 0.0013%).

The EtOAc-soluble extract (15 g) was subjected to silica gel column chromatography with a gradient of CHCl<sub>3</sub>-MeOH (100 : 1, 50 : 1, 20 : 1, 10 : 1) mixture to give nine fractions. Fraction 6 (320 mg) was chromatographed on a silica gel eluted with CHCl<sub>3</sub> to give seven fractions (a-g). Subfraction d was subjected to MPLC (*n*-hexane-CH<sub>2</sub>Cl<sub>2</sub> = 2 : 8) to yield compound **3** (3.5 mg, yield 0.0016%). Fraction 7 (82 mg) was dissolved in CHCl<sub>3</sub>, and 90% MeOH was added to solution and filtered. And then filtered solution was concentrated under reduced pressure using a rotary evaporator and dissolved in CHCl<sub>3</sub> to give compound **5** (20 mg, yield 0.0095%). Fraction 8 (1.34 g) was chromatographed on a silica gel, eluting with CH<sub>2</sub>Cl<sub>2</sub>-MeOH gradient mixture to give further fractions (A-G). Fraction B was purified further on Sephadex LH-20 using MeOH-CHCl<sub>3</sub> (3 : 2) mixture to give compound **6** (5.8 mg, yield 0.0027%), and fraction C was purified by preparative TLC using CH<sub>2</sub>Cl<sub>2</sub>-MeOH (20 : 1) mixture to give compound **7** (9.2 mg, yield 0.0043%) and compound **8** (1.3 mg, yield 0.0006%).

**Ferulic acid (1):** White powder; UV  $\lambda_{\max}$  (MeOH): 256, 309 nm; EI-MS ( $m/z$ ): 194; Molecular formula: C<sub>10</sub>H<sub>10</sub>O<sub>4</sub>; IR (KBr) cm<sup>-1</sup>: 1207, 1730 and 3345; <sup>1</sup>H-NMR (500 MHz in MeOH-*d*<sub>4</sub>)  $\delta$  3.89 (3H, s, OCH<sub>3</sub>), 6.30 (1H, d,  $J$  = 16 Hz, H-8), 6.80 (1H, d,  $J$  = 8 Hz, H-5), 7.06 (1H, dd,  $J$  = 8.0, 2.0 Hz, H-6), 7.17 (1H, d,  $J$  = 2.0 Hz, H-2), 7.60 (1H, d,  $J$  = 16 Hz, H-7). <sup>13</sup>C-NMR (125 MHz in MeOH-*d*<sub>4</sub>)  $\delta$  56.6 (OCH<sub>3</sub>), 111.9 (C-2), 116.1 (C-8), 116.6 (C-5), 124.1 (C-6), 127.9 (C-1), 147.1 (C-7), 149.5 (C-3), 150.6 (C-4), 171.1 (C-9).

**Vanillin (2):** Yellowish powder; UV  $\lambda_{\max}$  (MeOH): 229, 278 and 304 nm; EI-MS ( $m/z$ ): 152; Molecular formula: C<sub>8</sub>H<sub>8</sub>O<sub>3</sub>; IR (KBr) cm<sup>-1</sup>: 1676 and 3205; <sup>1</sup>H-NMR (500 MHz in CDCl<sub>3</sub>)  $\delta$  3.96 (3H, s, OCH<sub>3</sub>), 6.22 (1H, brs, H-2), 7.03 (1H, d,  $J$  = 8.4 Hz, H-5), 7.43 (1H, dd,  $J$  = 8.4, 1.5 Hz, H-6), 9.83 (1H, s, H-7). <sup>13</sup>C-NMR (125 MHz in CDCl<sub>3</sub>)  $\delta$  56.1 (OCH<sub>3</sub>), 108.8 (C-2), 114.4 (C-5), 127.5 (C-6), 129.8 (C-1), 147.1 (C-3), 151.7 (C-4), 190.8 (C-7).

**Coniferaldehyde (3):** White amorphous solid; UV  $\lambda_{\max}$  (MeOH): 223, 333 nm; EI-MS ( $m/z$ ): 178, 161, 147, 107; Molecular formula: C<sub>10</sub>H<sub>10</sub>O<sub>3</sub>; IR (KBr) cm<sup>-1</sup>: 833, 1167,

1601 and 3380; <sup>1</sup>H-NMR (500 MHz in CDCl<sub>3</sub>) δ 3.95 (3H, s, OCH<sub>3</sub>), 6.60 (1H, dd, *J* = 15.7, 8.0 Hz, H-8), 6.97 (1H, d, *J* = 8.5 Hz, H-5), 7.07 (1H, dd, *J* = 8.5, 1.5 Hz, H-6), 7.12 (1H, d, *J* = 1.5 Hz, H-2), 7.38 (1H, d, *J* = 15.5 Hz, H-7), 9.65 (1H, d, *J* = 8 Hz, H-9). <sup>13</sup>C-NMR (125 MHz in CDCl<sub>3</sub>) δ 55.9 (OCH<sub>3</sub>), 109.3 (C-2), 114.9 (C-5), 124.0 (C-6), 126.4 (C-8), 126.6 (C-1), 146.9 (C-3), 148.9 (C-4), 153.0 (C-7), 193.5 (C-9).

**Coniferyl alcohol (4):** White-pale yellow solid: UV λ<sub>max</sub> (MeOH): 230, 333 nm; EI-MS (*m/z*): 180, 137, 124, 119, 91 and 77; Molecular formula: C<sub>10</sub>H<sub>12</sub>O<sub>3</sub>; IR (KBr) cm<sup>-1</sup>: 1510, 1600, 3000, 3525 and 3595; <sup>1</sup>H-NMR (500 MHz in MeOH-*d*<sub>4</sub>) δ 3.85 (3H, s, OCH<sub>3</sub>), 4.19 (2H, dd, *J* = 6.0, 1.1 Hz, H-9), 6.19 (1H, dt, *J* = 15.6, 6.0 Hz, H-8), 6.51 (1H, brd, *J* = 15.9 Hz, H-7), 6.74 (1H, d, *J* = 8.0 Hz, H-5), 6.84 (1H, dd, *J* = 8.0, 1.8 Hz, H-6), 6.93 (1H, d, *J* = 1.8 Hz, H-2). <sup>13</sup>C-NMR (125 MHz in MeOH-*d*<sub>4</sub>) δ 56.1 (OCH<sub>3</sub>), 63.4 (C-9), 109.9 (C-2), 115.1 (C-5), 120.6 (C-6), 128.0 (C-8), 130.2 (C-1), 130.5 (C-7), 147.2 (C-4), 148.5 (C-3).

**2,6-Dimethoxy-*p*-benzoquinone (5):** Yellow needle: EI-MS (*m/z*): 168; Molecular formula C<sub>8</sub>H<sub>8</sub>O<sub>4</sub>; IR (KBr) cm<sup>-1</sup>: 1593, and 1727; UV λ<sub>max</sub> (CHCl<sub>3</sub>): 286 nm; <sup>1</sup>H-NMR (500 MHz in CDCl<sub>3</sub>) δ 3.82 (6H, s, OCH<sub>3</sub>) and 5.85 (2H, s, H-3,5); <sup>13</sup>C-NMR (125 MHz in CDCl<sub>3</sub>) δ 56.3 (OCH<sub>3</sub>), 107.3 (C-3,5), 157.1 (C-2,6), 176.3 (C-4), 186.2 (C-1).

***p*-Methoxycinnamic acid (6):** White powder: EI-MS (*m/z*): 178; Molecular formula C<sub>10</sub>H<sub>10</sub>O<sub>3</sub>; IR (KBr) cm<sup>-1</sup>: 1456, 1732, 2853 and 2924; UV λ<sub>max</sub> (MeOH): 217, 323 nm; <sup>1</sup>H-NMR (500 MHz in CDCl<sub>3</sub>) δ 3.69 (3H, s, OCH<sub>3</sub>), 6.38 (1H, d, *J* = 16 Hz, H-8), 6.81 (2H, d, *J* = 8.5 Hz, H-3, H-5), 7.62 (1H, d, *J* = 16 Hz, H-7), 7.45 (2H, d, *J* = 8.5 Hz, H-2, 6); <sup>13</sup>C-NMR (125 MHz in CDCl<sub>3</sub>) δ 52.1 (OCH<sub>3</sub>), 115.0 (C-8), 116.9 (C-3, 5), 127.2 (C-1), 131.2 (C-2, 6), 146.7 (C-7), 161.4 (C-4), 169.8 (COOH).

**(±)-Balanophonin (7):** Pale yellow oil; [α]<sub>D</sub> -0.39° (*c* = 0.34, CHCl<sub>3</sub>); EI-MS (*m/z*): 356; Molecular formula C<sub>20</sub>H<sub>20</sub>O<sub>6</sub>; IR (KBr) cm<sup>-1</sup>: 1595, 2923, 3411; UV λ<sub>max</sub> (MeOH): 216, 339 nm; <sup>1</sup>H-NMR (500 MHz in MeOH-*d*<sub>4</sub>) δ 3.56 (1H, brs, H-8), 3.81 (3H, s, OCH<sub>3</sub>), 3.83 (2H, brs, H-9), 3.90 (3H, s, OCH<sub>3</sub>), 5.60 (1H, d, *J* = 6.6 Hz, H-7), 6.63 (1H, dd, *J* = 16.5, 7.8 Hz, H-8'), 6.77 (1H, d, *J* = 8.1 Hz, H-5), 6.84 (1H, dd, *J* = 8.1, 1.8 Hz, H-6), 6.94 (1H, d, *J* = 1.8 Hz, H-2), 7.22 (1H, s, H-2'), 7.27 (1H, s, H-6'), 7.60 (1H, d, *J* = 16.5 Hz, H-7'), 9.57 (1H, d, *J* = 7.8 Hz, H-9'). <sup>13</sup>C-NMR (125 MHz in MeOH-*d*<sub>4</sub>) δ 54.7 (C-8), 56.5 (OCH<sub>3</sub>), 56.9 (OCH<sub>3</sub>), 64.7 (C-9), 90.2 (C-7), 110.7 (C-5), 114.4 (C-2'), 116.4 (C-2), 119.9 (C-6), 120.1 (C-6'), 127.2 (C-8'), 129.7 (C-1'), 131.4 (C-5') 134.0 (C-1), 146.1

(C-3'), 147.9 (C-3), 149.3 (C-4), 153.1 (C-4'), 156.2 (C-7'), 196.3 (C-9').

**6-Methoxychromanone (8):** Pale yellow powder; EI-MS (*m/z*): 178; Molecular formula: C<sub>10</sub>H<sub>10</sub>O<sub>3</sub>. IR (KBr): 1277, 1599 and 3424 cm<sup>-1</sup>; UV λ<sub>max</sub> (CHCl<sub>3</sub>): 257, 218, 208 nm; <sup>1</sup>H-NMR (500 MHz in CDCl<sub>3</sub>) δ 3.90 (3H, s, OCH<sub>3</sub>), 3.16 (2H, t, *J* = 6.5 Hz, H-3), 3.96 (2H, t, *J* = 6.5 Hz, H-2), 6.85 (1H, d, *J* = 8.5 Hz, H-8), 7.54 (1H, d, *J* = 2.0 Hz, H-5), 7.57 (1H, dd, *J* = 8.5, 2.0 Hz, H-7). <sup>13</sup>C-NMR (125 MHz in CDCl<sub>3</sub>) δ 41.0 (C-3), 56.3 (OCH<sub>3</sub>), 59.0 (C-2), 112.0 (C-5), 115.9 (C-8), 124.9 (C-7), 130.7 (C-10), 153.5 (C-9), 199.8 (C-4).

*Scavenging effect on DPPH radical.* The method of Blois with modification was used to measure scavenging capacity of 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical (Blois, 1958; Kitts *et al.*, 2000). Reaction mixtures containing test samples dissolved in EtOH and 300 μM DPPH in ethanolic solution in a 96-well microtiter plate were incubated at 37 °C for 30 min. Vitamin C and BHT were used as positive control. After the reaction, absorbance was read at 517 nm by a microplate reader and mean value was obtained from triplicate, and percent inhibition was calculated. IC<sub>50</sub> (inhibitory concentration) values denote the concentration of sample required to scavenge 50% of the DPPH free radical.

*Antibacterial activity and MIC determination.* Two bacterial strains, *S. mutans* KTCT 3065, *S. sobrinus* KTCT 3288, were used for the determination of antibacterial activity and were cultured in Brain Heart Infusion (BHI, Difco, Detroit, U.S.A.) medium under aerobic conditions at 37 °C. The screening of antibacterial activity for fractions was conducted using a disk diffusion method (Shapiro *et al.*, 1994). Briefly, test samples were dissolved in DMSO to a final concentration of 50 mg/mL. 100 μL of prepared culture containing 108 CFU/ml of bacteria was spread on BHI agar. The discs of 8 mm in diameter placed on the inoculated agar were impregnated with 50 μL (2.5 mg/mL) of extract. 10 μL of DMSO was used as vehicle control and gentamicin disc was also used as positive control. The inoculated plates were incubated for 24 h. Antibacterial activity was evaluated by measuring the zone of inhibition against the test bacterial strains.

MIC values were determined for single compounds from active fractions with the highest antibacterial activity in the disc diffusion assay, using a micro-well dilution method (Shapiro *et al.*, 1994). Growth inhibitory concentration of isolated compounds was tested against *Streptococcus mutans* and *S. sobrinus*, which were incubated for 24 h at 37 °C. At the end of the incubation period, optical density of the cultures was adjusted to

0.500 ± 0.050 with sterile BHI. Sterile 96-well microtiter plates were used by the respective growth medium. Four samples were used for each test concentration, and the experiments were performed in triplicate. Plates with wells containing 300 µL of BHI (100% growth controls) and 170 µL of BHI with 100 µL of substance to be tested were covered with plastic lids, then inoculated with 30 µL of bacterial culture adjusted to an optical density at 550 nm (OD<sub>550</sub>). All plates were incubated at 37 °C under appropriate atmospheric conditions and growth was estimated spectrophotometrically (630 nm) after 24 h using a microtiter plate reader. The controls included the inoculated growth medium without test compounds. Sample blanks contained uninoculated medium only.

### Results and discussion

The dried powder of the culms of *P. bambusoides* was extracted with 80% aqueous MeOH, and concentrated to yield greenish brown extract. Concentrated aqueous MeOH extract was partitioned with *n*-hexane, CHCl<sub>3</sub>, EtOAc, *n*-BuOH, and water. Of them, *n*-hexane and CHCl<sub>3</sub> extracts showed mild antioxidant activities against DPPH radical (IC<sub>50</sub>: 273 µg/mL for *n*-hexane and 244 µg/mL for CHCl<sub>3</sub>), and its EtOAc extract exhibited significant antibacterial effects against oral cariogenic *Streptococcus mutans* (10 mm of zone inhibition). Repeated column chromatography on Si gel, Sephadex

LH-20, MPLC and HPLC of the *n*-hexane, CHCl<sub>3</sub>, and EtOAc soluble fractions led to the isolation of eight bioactive simple phenolic compounds (**1 - 8**) (Fig. 1). Phenolics **1 - 4** were isolated as radical scavengers on DPPH radical from its *n*-hexane and CHCl<sub>3</sub> extracts, and phenolic compounds **3** and **5 - 8** were identified as anti-cariogenic agents from EtOAc layer. To determine the structures of compounds, the combined analyses with a series of 1D and 2D NMR, infrared, and mass spectra were accomplished. In addition, all physical and spectroscopic data obtained in this present study were compared with those of previously published manuscripts.

Phenolics **1**, **3** and **4** showed characteristic aromatic proton signals corresponding to a 1,3,4-trisubstituted aromatic ring which is composed of two doublets ( $J = \sim 8$  Hz and  $\sim 2$  Hz, respectively), doublet of doublets ( $J = \sim 2$  and 8 Hz), as well as two doublets with a large  $J$  value of coupling constant ( $J = \sim 16$  Hz) assignable to a *trans*-double bond. On the basis of 2D NMR spectra, the position of a methoxyl and a hydroxyl in an aromatic ring and a substituted moiety in a side chain was unambiguously elucidated. Thus, structures of these compounds were assigned as ferulic acid (**1**) (Young *et al.*, 1992), coniferaldehyde (**3**) (Carpinella *et al.*, 2005), and coniferyl alcohol (**4**) (Heravi *et al.*, 2004). Phenolic compound **6** displayed two doublets with a large coupling constant ( $J = \sim 16$  Hz) assignable to a *trans*-double bond, however, two *ortho*-coupled aromatic proton doublets




Fig. 1. Structures of phenolic compounds (**1 - 8**) from the culms of *P. bambusoides*.

were observed at  $\delta$  6.81 (2H, d,  $J = 8.5$  Hz) and 7.45 (2H, d,  $J = 8.5$  Hz). Therefore, compound **6** was identified as *p*-methoxycinnamic acid (Haruna *et al.*, 1982).

The  $^1\text{H-NMR}$  spectrum of **2** showed a singlet at  $\delta$  9.83 indicating an aldehyde group, in addition to two aromatic doublets ( $J = 8.4$  and 1.5 Hz, respectively), showing a splitting pattern of a doublet of doublets, which indicated that it is a trisubstituted benzene. The  $^{13}\text{C}$  NMR spectral data revealed the presence of a ketone carbonyl at  $\delta$  190.8, six aromatic ring carbons at  $\delta$  114.4, 127.5, 129.8, 147.1, 151.7, and a methoxyl carbon signal at  $\delta$  56.1. The structure of this compound was determined as vanillin (Harish *et al.*, 2005).

Compound **5** showed a molecular ion peak at  $m/z$  168, and its  $^1\text{H-NMR}$  spectrum exhibited only two singlet proton signals, a methoxyl signal at  $\delta$  3.82 and an olefinic proton signal at  $\delta$  5.85. The  $^{13}\text{C-NMR}$  spectrum showed five resonance signals, consisting of two carbonyl carbons at  $\delta$  176.3 and 186.2, and two carbons for an oxygenated olefinic bond at  $\delta$  157.1 and 107.3, and a methoxyl at  $\delta$  56.3. On the basis of this evidence and of a comparison of the published data (Nishina *et al.*, 1991), the structure of **5** was determined to be 2,6-dimethoxy-*p*-benzoquinone.

Phenolic compound **7** exhibited two singlet methoxyl signals at  $\delta$  3.81 and 3.90, a trisubstituted aromatic ring at  $\delta$  6.77, 6.84, and 6.94, one pair of meta-coupled benzene signals at  $\delta$  7.22 and 7.27. In addition, dihydrobenzofuran type signals at  $\delta$  5.60 and 3.83, a hydroxymethylene at  $\delta$  64.7, two *trans*-olefinic proton signals at  $\delta$  6.63 and 7.60 with  $J$  value of 16.5 Hz, and an aldehyde group at  $\delta$  9.57 were also observed in the  $^1\text{H-NMR}$  spectrum. Based on these spectral data, compound **7** was assumed to be a hgnan, and its unambiguous structure was established to be a neolignan, ( $\pm$ )-balanophonin by 2D NMR experiments and comparison with structural information of previously published data (Haruna *et al.*, 1982; Yeun *et al.*, 1998).

Compound **8** exhibited signals for three aromatic protons at  $\delta$  6.86, 7.57, and 7.54 corresponding to a typical trisubstituted aromatic ring. Two methylenes connected to ether oxygen atom showed as two triplets at  $\delta$  3.16 (2H, t,  $J = 6.5$  Hz) and 3.93 (2H, t,  $J = 6.5$  Hz). The  $^{13}\text{C-NMR}$  spectral data indicated the presence of a ketone carbonyl at  $\delta$  199.8, six aromatic ring carbons at  $\delta$  112.0, 115.9, 124.9, 130.7, 149.2, 153.5, and two methylene carbons connected to an oxygen atom at  $\delta$  41.8 and 59.0. The position of a methoxyl and a ketone were confirmed by long-range correlations with the ring protons in the HMBC spectrum. By the comparison of its spectral data with those of literature values (DeWald *et al.*, 1990), the structure of **8** was identified as 6-methoxy-

**Table 1.** DPPH radical scavenging activity of compounds **1 - 4**

Compounds	IC <sub>50</sub> ( $\mu\text{g/mL}$ )
ferulic acid ( <b>1</b> )	149
vanillin ( <b>2</b> )	135
coniferaldehyde ( <b>3</b> )	120
coniferyl alcohol ( <b>4</b> )	146
vitamin C	90
BHT	126

**Table 2.** Antibacterial activity of compounds **3, 5 - 8** isolated from the culms of *P. bambusoides*<sup>a,b</sup>

Compounds	Antibacterial activity MIC value ( $\mu\text{g/mL}$ )	
	<i>S. mutans</i>	<i>S. sobrinus</i>
gentamicin	< 50	< 50
coniferaldehyde ( <b>3</b> )	300	567
2,6-dimethoxy- <i>p</i> -benzoquinone ( <b>5</b> )	110	166
<i>p</i> -methoxycinnamic acid ( <b>6</b> )	352	470
( $\pm$ )-balanophonin ( <b>7</b> )	182	490
6-methoxychromanone ( <b>8</b> )	127	58

<sup>a</sup>MIC was determined by twofold serial broth dilution method.

<sup>b</sup>Vehicle control (0.1% DMSO) showed no MIC.

chromanone.

Biological activities of the isolated compounds **1 - 8** were summarized in the Tables **1** and **2**. As shown in Table **1**, phenolic compounds **1 - 4** isolated from *n*-hexane and  $\text{CHCl}_3$  extracts of *P. bambusoides* culms showed weaker radical scavenging activities on DPPH than vitamin C, however, similar antioxidant activities to that of BHT, a commercially available synthetic antioxidant. Coniferaldehyde (**3**) showed the strongest radical scavenging activity with an IC<sub>50</sub> value of 120  $\mu\text{g/mL}$ . The five antibacterial compounds, **3** and **5 - 8** identified from EtOAc extract exhibited inhibitory activity against the gram-positive cariogenic oral streptococci, *S. mutans* and *S. sobrinus*, frequently associated with human periodontitis, with MIC values ranging from 58 to 567  $\mu\text{g/mL}$ . Antibacterial activities of all the compounds were weaker than the positive control, gentamicin. Of the isolated compounds, 2,6-dimethoxy-*p*-benzoquinone (**5**) demonstrated the strongest activity against the bacterial strain, *S. mutans* with an MIC value of 110  $\mu\text{g/mL}$ . In contrast, 6-methoxychromanone (**8**) exhibited the strongest activity against *S. sobrinus* with an MIC value of 58  $\mu\text{g/mL}$ .

### Acknowledgments

We thank Gwangju Center of the Korea Basic Science

Institute (KBSI) for running NMR experiments. This work was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-220-E00042).

## References

- Bae, K.H., "The medicinal plants of Korea", Kyo-Hak Publishing Co., Ltd., Seoul, Korea, pp. 565-566 (2000).
- Blois, M.S., Antioxidant determination by the use of a stable free radical. *Nature*, **181**, 1199-1200 (1958).
- Carpinella, M.C., Ferrayoli, C.G., and Palacios, S.M., Antifungal synergistic effect of scopoletin, a hydroxycoumarin isolated from *Melia azedarach* L. fruits. *J. Agric. Food Chem.*, **53**(8), 2922-2927 (2005).
- Cowan, M.M., Plant products as antimicrobial agents. *Clin. Microbiol. Rev.*, **12**(4), 564-582 (1999).
- DeWald, H.A., Heffner, T.G., Jaen, J.C., Lustgarten, D.M., McPhail, A.T., Meltzer, L.T., Pugsley, T.A., and Wise, L.D., Synthesis and dopamine agonist properties of ( $\pm$ )-*trans*-3,4,4a,10b-tetrahydro-4-propyl-2H,5H-[1]benzopyrano[4,3-*b*]-1,4-oxazin-9-ol and its enantiomers. *J. Med. Chem.*, **33**, 445-450 (1990).
- Harada, H., Sakagami, H., Nagata, K., Ohhara, T., Kawazoe, Y., Ishihama, A., Hata, N., Misawa, Y., Terada, H., and Konno, K., Possible involvement of lignin structure in anti-influenza virus activity. *Antivir. Res.*, **15**(1), 41-49 (1991).
- Harish, R., Divakar, S., Srivastava, A., and Shivanandappa, T., Isolation of antioxidant compounds from the methanolic extract of the roots of *Decalepis hamiltonii* (Wight and Arn.). *J. Agric. Food Chem.*, **53**, 7709-7714 (2005).
- Haruna, M., Koube, T., Ito, K., and Murata, H., Balanophonin, A new neo-lignan from *Balanophora japonica* Makino. *Chem. Pharm. Bull.*, **30**(4), 1525-1527 (1982).
- Heravi, M., Masoum, S., and Shahbazikhah, P., Simulation of  $^{13}\text{C}$  nuclear magnetic resonance spectra of lignin compounds using principal component analysis and artificial neural networks. *J. Magn. Reson.*, **171**, 176-185 (2004).
- Kitts, D.D., Wijewickreme, A.N., and Hu, C., Antioxidant properties of a North American ginseng extract. *Mol. Cell. Biol.*, **20**(3), 1-10 (2000).
- Ko, S.W., Laupattarakasem, W., Saengnipanthkul, S., Mahaisavariya, B., and Theapongpackdee, S., Charcoal bamboo as a bone substitute: an animal study. *J. Med. Assoc. Thai.*, **77**, 496-500 (1994).
- Kweon, M.H., Hwang, H.J., and Sung, H.C., Identification and antioxidant activity of novel chlorogenic acid derivatives from bamboo (*Phyllostachys edulis*). *J. Agr. Food Chem.*, **49**(10), 4646-4655 (2001).
- Lee, S.I., "Clinical application of oriental medicinal herbs", Sung Bo Publishing Co., Seoul, Korea (1986).
- Nikaido, T., Sung, Y.I., Ohmoto, T., and Sankawa, U., Inhibitors of cyclic adenosine 3,5-monophosphate phosphodiesterase in *Phyllostachys nigra* Munro var. *henonis* Stapf. and *Phragmites communis* Trin., and inhibition by related compounds. *Chem. Pharm. Bull.*, **32**(2), 578-584 (1984).
- Nishina, A., Hasegawa, K., Uchibori, T., Seino, H., and Osawa, T.J., 2,6-Dimethoxy-*p*-benzoquinone as an antibacterial substance in the bark of *Phyllostachys heterocyclus* var. *pubescens*, a species of thick-stemmed bamboo. *J. Agric. Food Chem.*, **39**(2), 266-269 (1991).
- Shapiro, S., Meier, A., and Guggenheim, B., The antimicrobial activity of essential oils and essential oil components towards oral bacteria. *Oral Microbiol. Immunol.*, **9**(4), 202-208 (1994).
- Suga, A., Takaishi, Y., Goto, S., Munakata, T., Yamauchi, I., and Kogure, K., Two lignan dimers from bamboo stems (*Phyllostachys edulis*). *Phytochemistry*, **64**, 991-996 (2003).
- Tanaka, E., Tanaka, C., Mori, N., Kuwahara, Y., and Tsuda, M., Phenylpropanoid amides of serotonin accumulate in witches' broom diseased bamboo. *Phytochemistry*, **64**(5), 965-969 (2003).
- Young, H.S., Kim, M.S., Park, H.J., Chung, H.Y., and Choi, J.S., Phytochemical study on *Catalpa ovate*. *Arch. Pharm. Res.*, **15**, 322-327 (1992).
- Yuen, M.S.M., Xue, F., Mak, T.C.W., and Wong, H.N.C., On the absolute structure of optically active neolignans containing a dihydrobenzo [*b*]furan skeleton. *Tetrahedron*, **54**(41), 12429-12444 (1998).

Received September 17, 2011

Revised October 25, 2011

Accepted October 30, 2011