



Published in final edited form as:

Angew Chem Int Ed Engl. 2019 June 11; 58(24): 8029–8033. doi:10.1002/anie.201902489.

Biocompatible SuFEx Click Chemistry: Thionyl Tetrafluoride (SOF₄) Derived Connective Hubs For Bioconjugation to DNA and Proteins

Feng Liu^{§,a,b,g}, Hua Wang^{§,b}, Suhua Li^c, Grant A. L. Bare^b, Xuemin Chen^d, Chu Wang^d, John E. Moses^e, Peng Wu^f, and K. Barry Sharpless^b

^[a]School of Perfume and Aroma Technology, Shanghai Institute of Technology, Shanghai 201418, P.R. China.

^[b]Department of Chemistry, The Scripps Research Institute. La Jolla, CA 92037, USA.

^[c]School of Chemistry, Sun Yat-sen University, Guangzhou, 510275, P. R.China.

^[d]College of Chemistry and Molecular Engineering, Peking University, Beijing, 100871, P.R. China.

^[e]La Trobe Institute For Molecular Science, La Trobe University, Melbourne, VIC 3086, Australia.

^[f]Department of Molecular Medicine, The Scripps Research Institute, La Jolla, CA, 92037, USA.


^[g]Department of Chemistry, Fudan University, Shanghai, 200438, P.R. China.

Abstract

We report here the development of a suite of biocompatible SuFEx transformations from the SOF₄-derived iminosulfur oxydifluoride-hub in aqueous buffer conditions. These biocompatible SuFEx reactions of iminosulfur oxydifluorides (R-N=SOF₂) with primary amines give sulfamides (8 examples, up to 98%), while the reaction with secondary amines furnish sulfuramidimidoyl fluoride products (8 examples, up to 97%). Likewise, under mild buffered conditions, phenols react with the iminosulfur oxydifluorides (Ar-N=SOF₂) to produce sulfurofluoridoimidates (13 examples, up to 99%), which can themselves be further modified by nucleophiles. These transformations open the potential for asymmetric and trisubstituted linkages projecting from the sulfur(VI) center, including versatile S-N and S-O connectivity (9 examples, up to 94%). Finally, the SuFEx bioconjugation of iminosulfur oxydifluorides to amine-tagged single-stranded DNA and to BSA protein demonstrate the potential of SOF₄ derived SuFEx click chemistry in biological applications.

Graphical Abstract

[§]These two authors contributed equally to this work.



We describe the first high yielding and biocompatible SOF_4 based SuFEx reaction methodology that operates at low concentration in aqueous solution under mild pH and temperature conditions. The biocompatible SuFEx reactions open the potential for asymmetric and trisubstituted products at the sulfur(VI) center with versatile S-N and S-O connectivity. The SuFEx bioconjugation of iminosulfur oxydifluorides to either BSA protein or to single-stranded DNA support the development of these useful reactions in bioconjugation applications.

Keywords

SuFEx; Click Chemistry; Bioconjugation; Protein modification; DNA Encoded Libraries

Synthetic methods that allow the chemoselective derivatization of biomolecules, such as proteins, DNA, RNA, and carbohydrates etc., have enabled many important discovery-based investigations, and are essential tools for interrogating biomolecules in their native environment.^[1]

The development of bio-orthogonal click chemistry over the past decade^[2], and in particular CuAAC and the strain promoted copper-free Huisgen cycloaddition reactions have had enormous impact as tools for enabling chemical conjugations. Despite these major advances, there remains a growing demand for engineering fast, selective, and high yielding transformations that can be conducted in a complex biological milieu under physiological conditions. This is particularly pertinent in the burgeoning field of DNA-Encoded Chemical Libraries (DELs). Pioneered by Lerner and Brenner in 1992, DELs comprise individual organic molecules with distinctive DNA tags that serve as amplifiable identification bar codes to allow the construction and screening of combinatorial libraries of unprecedented size.^[3]

To avoid damaging the DNA during the course of synthesizing the library, there are a number of requirements that a reaction must meet. For example: the chemistry must be compatible with aqueous media and proceed at moderate temperatures (< 90 °C) within pH 4–14; the reaction rate constant should be high enough to accommodate low concentrations of reactants (< 1 mM); and finally, the reactants should be inert to phosphates and nucleotides. Unsurprisingly, only a handful of reactions are available (Figure 1),^[4] and there is an urgent need to expand the repertoire of DNA compatible transformations.

Sulfur-fluoride exchange (SuFEx) is a new generation of click chemistry that enables new connections to be formed through S-F exchange with a range of nucleophiles, to give stable S-N, S-O, and S-C covalent linkages. Much like nature's own phosphate hubs, SuFEx connectors enable the modular construction of functional molecules through SuFExable hubs including: sulfonyl fluoride (SO_2F_2),^[5] ethenesulfonyl fluoride (ESF),^[6] and thionyl tetrafluoride (SOF_4).^[7]

A growing number of applications of SuFEx click chemistry have been reported since it was first introduced in 2014,^[5] including new synthetic methodologies,^[8] polymer preparation,^[5,9] and bioconjugation.^[10] However, it wasn't until the evolution of Thionyl tetrafluoride (SOF_4) as the first multidimensional SuFEx hub that the true power of polyvalent click chemistry was realized—opening the door to the 3-dimensional world and the domain of biological space.^[7]

SOF_4 is a reactive gas that under SuFEx catalysis undergoes S-F exchange with primary amines, including anilines and aliphatic primary amines, to afford tetrahedral iminosulfur oxydifluoride products **I** (Figure 2).^[7a-b] The remaining two SuFExable S-F handles of **I** can themselves undergo sequential SuFEx reactions with N-, O-, and C- nucleophiles, to form a diverse array of sulfur fluoride compounds and downstream multi-substituted derivatives **II–VII**. Collectively, these transformations form the framework for creating SOF_4 derived SuFEx platforms and libraries of extreme, hub-based special variety.^[7]

SuFEx enabled “Schotten-Baumann-like”, linkage reactions with amines and phenols are nearly perfect at aqueous buffer interfaces, but better in that unwanted hydrolysis is even more suppressed. This interfacial enhanced mode of S-F reactivity is deemed perfect for bioconjugation endeavors where SuFEx activation can be achieved with nucleophilic catalysts (e.g. DBU (1,8-diazabicyclo[5.4.0]undec-7-ene)), and also by vigorous stirring with an immiscible buffered aqueous phase. The blending of water with miscible solvents such as THF or acetonitrile also aid the process, although we posit that most SuFEx reactions will benefit from two-phase conditions in which the aqueous phase contains bifluoride (KFHF). We anticipated that such conditions would be amenable for the development of biocompatible SuFEx chemistry, and report here the first example of SOF_4 -derived SuFEx reactions that proceed in dilute aqueous solutions under mild conditions of pH and temperature. We demonstrate the power of the new bioconjugation methodology by the joining of small molecules to amine-tagged single-stranded DNA and also to BSA-protein.^[11]

The *p*-ethynyl-aniline iminosulfur oxydifluoride **1a** and 2-phenylethan-1-amine **2a** were selected as model connective partners for the development of the SuFEx reactions in aqueous conditions (Table S1, see supporting information for condition optimization). Rigorous screening of conditions revealed the reaction of **2a** and iminosulfur oxydifluoride **1a** proceeded rapidly at low concentration (1 mM in pH 7.3 aq. PBS buffer/MeCN 1:1 mixture) at room temperature. Notably, the sulfuramidimidoyl fluoride intermediate **3'** was not detected in the reaction mixture—itself spontaneously hydrolyzing to furnish the sulfamide product **3** (Table 1).

Under the optimized reaction conditions, the iminosulfur oxydifluorides **1** underwent the substitution-hydrolysis sequence with **2a** to give the corresponding sulfamide products in excellent yield (**3a-3h**, 80–98%). The aliphatic amine iminosulfur oxydifluorides **2d** and **2e** were found to be slightly less reactive than the aniline counterparts, although the reactions were generally complete within 1 hour to giving products **3d** (91%) and **3e** (80%). It is noteworthy that the reaction of the amino acid L-alanine **2f** and **1a** gave product **3f** in 95% yield, demonstrating that the SuFEx is compatible with free carboxylic acid functionality. Similarly, the long chain aliphatic primary amine *N*-Boc-lysine methyl ester **2g** coupled successfully with both **1a** and **1h** to form the products **3g** and **3h** respectively.

Next, the substitution by secondary amines **4** under mild aqueous conditions were examined (Table 2). A slight increase of the reaction temperature (37 °C) and prolonged reaction time (12 h) were required to yield the corresponding sulfuramidimidoyl fluorides **5** in excellent yield (**5a-h**, 88–97%).

In the context of bioconjugation, the potential SuFEx chemistry of tyrosine was considered particularly important, since it is found abundantly in naturally occurring proteins. We then examined the reactivity of a number of phenols **6** with iminosulfur oxydifluorides **1** under buffered aqueous conditions (pH 8.0). As illustrated in table 3, the monosubstituted sulfurofluoridoimidates **7a-7l** (95–99%) were readily obtained with stirring at 37 °C for 16 h in slightly basic pH 8.0 aq. PBS buffer/MeCN 1:1 mixtures. A wide selection of aromatic substituents on both the iminosulfur oxydifluorides **1** and the phenols **6** were well tolerated, although the reaction of the aliphatic iminosulfur oxydifluoride **1m** and phenol produced **7m** in only moderate yield (19%), even when a large excess of difluoride **1m** (10 equiv.) was used. In this particular case, most of the starting material **1m** was hydrolyzed to the (2-phenoxyethyl)sulfamic acid. Significantly, the products **7j-k**, and **7l**, derived from the reaction with *N*-Boc-tyrosine methyl ester and the *N*-Boc-tyrosine respectively, were isolated in excellent yield. The results support the notion that bioconjugation to protein tyrosine residues through the SOF₄-derived iminosulfur oxydifluoride-hub is viable.

The stability of the monofluoride species **5** and **7** were next tested in 1mM aqueous buffered solutions as a function of pH (Figure S1 and S2). The compound **7a** was found to be stable after agitation at 37 °C for 48 h from pH 6 to pH 8, but hydrolyzed to the sulfonate at more acidic or basic pH. On the other hand, compound **5a** was more stable even after subjection to agitation at 37 °C for 96 h in pH 2 to pH 14 solutions. Based upon these results, we studied the reaction of monofluoride **7a** and pyrrolidine **4j** (Table S2), to implement a second SuFEx reaction.

Screening different reaction conditions revealed that a large excess of nucleophile (50 equiv.) under more extreme conditions (80 °C, pH 10) were required to achieve satisfactory conversion. The reaction with a range of secondary amines **4a-e** gave the corresponding sulfonimidates **8a-e** in good yields, whereas reaction with phenols **6a-d** gave excellent conversion to the sulfurimidates **9a-d** (86–94%) (Table 4).

In sharp contrast, efforts to substitute the S-F bond of the sulfuramidimidoyl fluoride **5** using ArOH or secondary amine as nucleophile were unsuccessful, even under forcing conditions

(100 °C, pH 10, 48 h)—this is consistent with the increased stability imparted by the greater electron density on sulfur associated with two nitrogen-sulfur bonds.

With optimized conditions in hand, we next investigated the suitability of the new biocompatible SOF_4 derived SuFEx conditions in the labelling of DNA, and hence as new technology DEL (Figure 3). The single-stranded DNA **11**, comprising a long chain primary amine tail overhang and hairpin structure was selected as a suitable model substrate for conjugation reaction. Agitation of a reaction mixture comprising difluoride **1a** and **11** at room temperature for 6 h resulted in successful conjugation and formation of the product **12** in 70% yield (determined by HPLC). The phenol-tailed single-stranded DNA **14** was then synthesized by DNA-amide condensation and desilylation with linker molecule **13**. Coupling of **14** with the difluoride **1a** gave the corresponding small molecule-DNA bioconjugate with the CuAAC-ready alkyne **15**.^[11g]

To explore the new SOF_4 -derived reagents for protein modification, bovine serum albumin (BSA) was chosen as a model protein to react with the alkyne-bearing difluoride **1a** (1–100 equivalents). The resultant BSA-SuFEx bioconjugates were purified and analyzed using liquid chromatography-mass spectrometry (LC-MS) (Figure 4a). Using 1 molar equivalent of **1a** to react with BSA (15 μM in pH 7.3 PBS) for 1 hour at 25 °C, a single modification of the target protein was observed. With 10 equivalents of **1a** under the same condition, a triple modification was detected in the MS spectrum, and one modification site was confirmed as Lys548 by LC-MS/MS (Figure S11). When 100 equivalents of difluoride **1a** were used, BSA was found to be modified by 8 molecules of **1a** (Scheme S7–S9). Remarkably, in all cases, only lysine residues were found to be modified. The alkyne-modified BSA conjugates enabled further reaction with biotin-PEG₃-azide via ligand-assisted CuAAC^[12]. Likewise, the biotin-PEG₃-amine difluoride **16** could be directly connected to BSA via an exposed amine such as a lysine residue (Figure 4b). However, due to lower reactivity of this aliphatic amine derived difluoride, only moderate SuFEx conversion of ~30% was observed (Scheme S12).

In conclusion, we have developed a series of high yielding and biocompatible SOF_4 based SuFEx reactions that operate at low concentration, in aqueous solution under mild pH and temperature conditions. A diverse selection of chemical space was sampled through the polyvalent and multidimensional *N*- and *O*-nucleophilic substitutions from the SOF_4 hub. The efficient labelling of BSA protein and amine-tagged single-stranded DNA was also accomplished, demonstrating the potential of the biocompatible SuFEx ligation sequence as a useful tool for bioconjugation applications. Collectively, the experimental results validate the suitability for aqueous phase combinatorial chemistry and we posit that SOF_4 mediated SuFEx will prove to be a generally useful addition to the click chemistry toolbox with wide application in chemical biology, biophysics, bioengineering, and materials science.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors gratefully acknowledge financial support from the NIH (R01 GM117145 to K.B.S. R01GM093282 to P.W.), the ARC for supporting a Future Fellowship FT170100156 (J.E.M), and NSFC 21302126(F. L).

References

- [1]. Sletten EM, Bertozzi CR, *Angew. Chem. Int. Ed.* 2009, 48, 6974.
- [2]. a) Kolb HC, Finn MG, Sharpless KB, *Angew. Chem. Int. Ed.* 2001, 40, 2004. b) Baskin JM, Bertozzi CR, *Qsar & Comb. Sci.* 2007, 26, 1211. c) Tornøe CW, Christensen C, Meldal M, *J. Org. Chem.* 2002, 67, 3057. [PubMed: 11975567] d) Rostovtsev VV, Green LG, Fokin VV, Sharpless KB, *Angew. Chem. Int. Ed.* 2002, 41, 2596. e) Baskin JM, Prescher JA, Laughlin ST, Agard NJ, Chang PV, Miller IA, Lo A, Codelli JA, Bertozzi CR, *Proc. Natl. Acad. Sci.* 2007, 104, 16793. [PubMed: 17942682]
- [3]. a) Brenner S, Lerner RA, *Proc. Natl. Acad. Sci.* 1992, 89, 5381 [PubMed: 1608946] b) Brenner S, Lerner RA, *Angew. Chem. Int. Ed.* 2017, 56, 1164. c) Neri D, Lerner RA, *Annu. Rev. Biochem.* 2018, 87, 479. [PubMed: 29328784]
- [4]. a) Malone ML, Paegel BM, *ACS Comb. Sci.* 2016, 18, 182. [PubMed: 26971959] b) Nicholas N, Bassi G, Scheuermann J, Neri D, *FEBS Lett.* 2018, 592, 2168. [PubMed: 29683493]
- [5]. Dong J, Krasnova L, Finn MG, Sharpless KB, *Angew. Chem. Int. Ed.* 2014, 53, 9430.
- [6]. a) Krutak JJ, Burpitt RD, Moore WH, Hyatt JA, *J. Org. Chem.* 1979, 44, 3847; b) Zheng Q, Dong J, Sharpless KB, *J. Org. Chem.* 2016, 81, 11360. [PubMed: 27764941] c) Qin H, Zheng Q, Bare GAL, Wu P, Sharpless KB, *Angew. Chem. Int. Ed.* 2016, 55, 14155. d) Chinthakindi PK, Govender KB, Kurnar AS, Kruger HG, Govender T, Naicker T, Arvidsson PI, *Org. Lett.* 2017, 19, 480. [PubMed: 28075600] e) Zha G, Zheng Q, Leng J, Wu P, Qin H, Sharpless KB, *Angew. Chem. Int. Ed.* 2017, 56, 4849.
- [7]. a) Li S, Wu P, Moses JE, Sharpless KB, *Angew. Chem. Int. Ed.* 2017, 56, 2903. b) Gao B, Li S, Wu P, Moses JE, Sharpless KB, *Angew. Chem. Int. Ed.* 2018, 56, 2903.
- [8]. a) Kende AS, Mendoza JS, *J. Org. Chem.* 1990, 55, 1125; b) Chen W, Dong J, Li S, Liu Y, Wang Y, Yoon L, Wu P, Sharpless KB, Kelly JW, *Angew. Chem. Int. Ed.* 2016, 55, 1835. c) Zhang E, Tang J, Li S, Wu P, Moses JE, Sharpless KB, *Chem. Eur. J.* 2016, 22, 5692. [PubMed: 26990693] d) Ren G, Zheng Q, Wang H, *Org. Lett.* 2017, 19, 1582. [PubMed: 28332844] e) Dondoni A, Marra A, *Org. Biomol. Chem.* 2017, 15, 1549. [PubMed: 28116403]
- [9]. a) Dong J, Sharpless KB, Kwisnek L, Oakdale JS, Fokin VV, *Angew. Chem. Int. Ed.* 2014, 53, 9466. b) Li S, Beringer LT, Chen S, Averick S, *Polymer* 2015, 78, 37. c) Yatvin J, Brooks K, Locklin J, *Chem. Eur. J.* 2016, 22, 16348. [PubMed: 27557871] d) Oakdale JS, Kwisnek L, Fokin VV, *Macromolecules* 2016, 49, 4473. e) Wang H, Zhou F, Ren G, Zheng Q, Chen H, Gao B, Klivansky L, Liu Y, Wu B, Xu Q, Lu J, Sharpless KB, Wu P, *Angew. Chem. Int. Ed.* 2017, 56, 11203. f) Brendel JC, Martin L, Zhang J, Perrier S, *Polym. Chem.* 2017, 8, 7475. g) Wang H, Ren F, Zhou G, Zheng Q, Chen H, Gao B, Klivansky L, Liu Y, Wu B, Xu Q, *Angew. Chem. Int. Ed.* 2017, 56, 11203. i) Zhu H, Chen D, Li N, Xu Q, Li H, He J, Wang H, Wu P, Lu J, *Chemistry* 2017, 23, 14712. [PubMed: 28881405] j) Gao B, Zhang L, Zheng Q, Zhou F, Klivansky LM, Lu J, Liu Y, Dong J, Wu P, Sharpless KB, *Nat. Chem.* 2017, 9, 1083. [PubMed: 29064495]
- [10]. a) Grimster NP, Connelly S, Baranczak A, Dong J, Krasnova LB, Sharpless KB, Powers ET, Wilson IA, Kelly JW, *J. Am. Chem. Soc.* 2013, 135, 5656. [PubMed: 23350654] b) Baranczak A, Liu Y, Connelly S, Du W, Greiner ER, Genereux JC, Wiseman RL, Eisele YS, Bradbury NC, Dong J, Noodleman L, Sharpless KB, Wilson IA, Encalada SE, Kelly JW, *J. Am. Chem. Soc.* 2015, 137, 7404. [PubMed: 26051248] c) Zelli R, Tommasone S, Dumy P, Marra A, Dondoni A, *Eur. J. Org. Chem.* 2016, 30, 5102. d) Li S, Cohen-Karni D, Beringer LT, Wu C, Kallick E, Edington H, Passineau MJ, Averick S, *Polymer* 2016, 99, 7. e) Hoppmann C, Wang L, *Chem. Commun.* 2016, 52, 5140. f) Chen W, Dong J, Plate L, Mortenson DE, Brighty GJ, Li S, Liu Y, Galmozzi A, Lee PS, Hulce JJ, Cravatt BF, Saez E, Powers ET, Wilson IA, Sharpless KB, Kelly JW, *J. Am. Chem. Soc.* 2016, 138, 7353. [PubMed: 27191344] g) Álvarez NH, van de Langemheen H, Brouwer AJ, Liskamp RM, *Bioorganic & Medicinal Chemistry* 2017, 25, 5055. [PubMed: 28734665] i) Mortenson DE, Brighty GJ, Plate L, Bare G, Chen W, Li S, Wang H,

Cravatt BF, Forli S, Powers ET, Sharpless KB, Wilson IA, Kelly JW, J. Am. Chem. Soc. 2018, 140, 200. [PubMed: 29265822]

- [11]. a) Hermanson GT, Bioconjugate Techniques (Academic Press, San Diego, CA, 1996). b) Griffin BA, Adams SR, Tsien RY, Science, 1998, 281, 269. [PubMed: 9657724] c) Wu P, Shui W, Carlson BL, Hu N, Rabuka D, Lee J, Bertozzi CR Proc. Natl. Acad. Sci. USA, 2009, 106, 3000. [PubMed: 19202059] d) Casi G, Huguenin-Dezot N, Zuberbuhler K, Scheuermann J, Neri D. J. Am. Chem. Soc. 2012, 134, 5887. [PubMed: 22394212] e) MacDonald JI, Munch HK, Moore T, Francis MB. Nat. Chem. Biol. 2015, 11, 326. [PubMed: 25822913] f) Zhang C, Welborn M, Zhu T, Yang NJ, Santos MS, Van Voorhis T, Pentelute BL, Nat. Chem. 2016, 8, 120. [PubMed: 26791894] g) Choi EJ, Jung D, Kim J, Lee Y, Kim BM, Chem. Eur. J. 2018, 24, 10948. [PubMed: 29935027],
- [12]. Soriano del Amo D, Wang W, Jiang H, Besanceney C, Yan AC, Levy M, Liu Y, Marlow FL, Wu P, J. Am. Chem. Soc. 2010, 132, 16893. [PubMed: 21062072]




Figure 1. Reactions that are commonly used and compatible for DELs chemistry. ^[4f]




Figure 2. SuFEx connections from the polyvalent and multidimensional hub of SOF_4 .






Figure 3.
Bioconjugation of difluoride **1a** to amine-tagged ss-DNA.

**Figure 4.**

- a) Sequential SuFEx bioconjugation and biotin CuAAC labeling of BSA protein.
 b) Modification of BSA protein with biotin-PEG3-amine difluoride 16.

Table 1.


Connection of iminosulfur oxydifluorides **1** and 1° amines **2**.^[a, b]



^[a]Reactions were performed with **1** (0.10 mmol), **2** (0.10 mmol), MeCN 50 mL, pH 7.3 aq. PBS buffer 50 mL, at $25\text{ }^\circ\text{C}$ for 5 minutes unless otherwise stated.

^[b]Isolated yield.


^[c]For **3 d** and **3 e**, the reaction was carried out for 1 hour.

Table 2.Connection of iminosulfur oxydifluorides **1** and 2° amines **4**.^[a, b]

^[a] Reactions were performed with **1** (0.10 mmol), **4** (0.10 mmol), MeCN 50 mL, pH 7.3 aq. PBS buffer 50 mL, at 37 °C for 12 hours.

^[b] Isolated yield.

Table 3.


Connection of iminosulfur oxydifluorides **1** and phenols **6**.^[a, b]

[a] Reactions were performed with **1** (0.10 mmol), **6** (0.10 mmol), MeCN 50 mL, pH 8.0 aq. PBS buffer 50 mL, at 37 °C for 16 hours unless otherwise stated.

[b] Isolated yield.

[c] For **7 m**, 1 mmol (10 eq.) of **1m** was used.

Table 4.

3-dimension SuFEx reactions of monofluoride 7.^[a, b]

^[a] Reactions were performed with 7 (0.10 mmol), 4 or 6 (5 mmol, 50 eq), MeCN 50 mL, pH 10 aq. PB buffer 50 mL, at 80°C for 16 hours.

^[b] Isolated yield.