

Research Article

Biodegradation and utilization of waste cooking oil by *Yarrowia lipolytica* CECT 1240

Alberto Domínguez, Francisco J. Deive, M. Angeles Sanromán and María A. Longo

Department of Chemical Engineering. University of Vigo, Campus Universitario Lagoas–Marcosende, Vigo, Spain

1 Biodegradation of waste cooking oil and its application as lipase production inducer in cultures
2 of *Yarrowia lipolytica* CECT 1240 have been investigated, both in shake flasks and a bench-scale
3 bioreactor. The ability of this strain to degrade the spent oil was evaluated by monitoring
4 COD throughout the cultures, and a remarkable decrease was recorded (almost 90% decrease
5 in oil COD after 3 days in bioreactor). Moreover, the addition of waste cooking oil to the
6 medium led to a significant augmentation in extracellular lipase production by the yeast,
7 compared to oil-free cultures. This confirms the suitability of the studied residue as an inducer
8 of lipase biosynthesis, which is a very interesting fact, from an economic standpoint. These results
9 were confirmed when a fed-batch strategy was proposed. Finally, some properties of the crude
10 enzyme were studied, and compared to the enzymes obtained when non-used oil was added to
11 the medium.

12
13 **Practical application:** Nowadays the search of new strategies to valorize wastes from the food and agro
14 industries is attracting a great scientific interest due to the important advantages offered from an
15 economic and environmental point of view. For this reason, the yeast *Yarrowia lipolytica* CECT 1240
16 is proposed for degrading waste cooking oils. This approach entails also another benefit in terms of
17 lipolytic enzyme synthesis, since the addition of used up oils has a lipase inducer effect. The enormous
18 interest of lipases is reflected in the number of applications that they present, which justify the research
19 work proposed here. The process was successfully carried out both in shake flasks and a bench-scale
20 bioreactor, allowing producing high levels of lipolytic activity at the same time that the COD was
21 diminished up to nearly 90%.

22
23 **Keywords:** biodegradation / cooking oil / lipase / waste / *Yarrowia lipolytica*

24 Received: February 11, 2010 / Revised: August 5, 2010 / Accepted: August 11, 2010

25 DOI: 10.1002/ejlt.201000049

1 Introduction

26 Large amounts of cooking oils, subjected to high tempera-
27 tures (160–200 °C) for relatively long periods of time, are
28 daily produced throughout the world. As a result of the harsh
29 conditions, oxidation of fatty acids and their subsequent

transformations may cause unpleasant flavors and odors, 1
which severely limits long-term repeated use [1, 2]. Some 2
procedures have been proposed to eliminate free fatty acids 3
and extend oil usability [3]; however, they are seldom uti- 4
lized. Disposal of this kind of polluting wastes has become an 5
increasingly distressing matter. The immiscibility of oil in 6
water, albeit facilitating separation by flotation, hinders trans- 7
portation through pipelines, disposal, and biodegradation. 8
When poured down drains, they result in coating and event- 9
ual occlusion of drainage and sewage pipes [4]. 10

From an economic standpoint, the development of fea- 11
sible ways for reutilization of waste edible oils and fats would 12
be highly desirable. Traditionally, they have been used for 13
soap fabrication, as additives in domestic animals feedstock, 14
or for other creative applications such as the fabrication of an 15

Correspondence: Dr. María A. Longo, Department of Chemical
Engineering. University of Vigo, Campus Lagoas–Marcosende, E-36310
Vigo, Spain

E-mail: mlongo@uvigo.es

Fax: +34 986 812380

Abbreviation: COD, chemical oxygen demand; OMW, olive mill
wastewater

1 inexpensive cleaning agent for soot and carbon residues on
2 machine parts and firearms [5]. However, most recent
3 research contributions on this topic focus on the utilization
4 of waste vegetable oils as raw material for the production of
5 biodiesel [6], an alternative environmentally friendly fuel for
6 Diesel engines composed of monoalkyl esters of long chain
7 fatty acids. Utilization of waste oil would help to reduce
8 production costs for this combustible, therefore contributing
9 to resolve one of the main obstacles to its extensive utilization
10 [1, 7–9]. Additionally, the use of spent vegetable oil as diesel
11 engine fuel, following thermal cracking to reduce its viscosity,
12 has been proposed [10].

13 Biodegradation of vegetable oils has been studied mainly
14 in relation to their incorporation (after chemical modifi-
15 cation) to the formulae of lubricants and greases [11], or
16 the treatment of accidental oil spills [12–14]. A wide number
17 of microorganisms, both aerobic and anaerobic, are able to
18 break up vegetable oils, although chemical characteristics of
19 the oil could determine its biodegradability [12]. Vegetable
20 oil biodegradation occurs in a similar way both in the pres-
21 ence or absence of oxygen, through hydrolysis of triglycerides
22 and beta oxidation of released fatty acids. High oil concen-
23 trations may inhibit anaerobic degradation under methano-
24 genic conditions, although this may be solved if iron-reducing
25 conditions are induced [14]. The design of strategies involv-
26 ing microorganisms, more specifically yeasts to simul-
27 taneously degrade waste oils and obtain high added-value
28 products would be an interesting endeavor, both from an
29 economic and environmental angle [15, 16].

30 Microbial lipases (triacylglycerol acylhydrolases, EC
31 3.1.1.3) are considered as extremely useful biocatalysts
32 due to their numerous potential applications, such as food
33 and detergent manufacture and specific synthesis of fine
34 chemicals or chiral drugs [17]. They catalyze both hydrolysis
35 and synthesis reactions (*i.e.*, esterification, transesterifica-
36 tion), and are usually stable in organic solvents and at rela-
37 tively high temperatures [18–21]. Many microorganisms are
38 able to produce extracellular lipases, although nutritional and
39 physico-chemical factors (*i.e.*, temperature, pH, nitrogen,
40 and carbon sources, presence of lipids, inorganic salts, agi-
41 tation, oxygen concentration) strongly influence the levels of
42 enzyme expression [17, 22]. Although the addition of veg-
43 etable oil to the culture medium has been repeatedly used to
44 trigger microbial lipase production, the possibility of using
45 waste cooking oil has not been evaluated so far. If successful,
46 this approach could help to diminish lipase production costs,
47 and also contribute to lessen the environmental problem
48 posed by used frying oils.

49 The lipase production ability of several strains of *Yarrowia*
50 *lipolytica* has been investigated [23–35]. Some researchers
51 have proposed the utilization of this yeast to reduce contami-
52 nation in olive mill wastewater (OMW). They obtained
53 promising results in terms of COD reduction, and a signifi-
54 cant decrease in polyphenol content [36]. In previous works,
our group indicated rather high levels of extracellular lipase in

1 submerged and solid-state cultures of *Y. lipolytica* CECT
2 1240 [27, 28]. The influence of several lipids and surfactants
3 on biomass and lipolytic enzyme biosynthesis was examined,
4 and the best results were obtained when vegetable oils were
5 added to the culture medium.

6 In this work, the ability of *Yarrowia lipolytica* CECT 1240
7 (ATCC 18942) to grow on a liquid medium supplemented
8 with waste cooking oil has been investigated. The variation in
9 chemical oxygen demand (COD) has been monitored
10 throughout the cultures, in order to assess the viability of
11 the process as a decontamination method. Also, the levels of
12 lipase production have been determined to evaluate the effi-
13 ciency of the spent oil as lipolytic enzyme production inducer.

2 Materials And Methods 14

2.1 Microorganism and media 15

16 *Yarrowia lipolytica* CECT 1240 (ATCC 18942) was grown
17 on peptone–malt extract agar plates. The medium consisted
18 of (g dm⁻³, in distilled water): glucose 10, peptone 5, yeast
19 extract 3, malt extract 3, and agar 20. Plates were incubated
20 at 30 °C for 7 days and stored at 4 °C between transfers.

21 Submerged cultures were carried out using a basal
22 medium [23] prepared in distilled water, containing (per
23 dm³): 20 g glucose, 2 g urea, 1 g KH₂PO₄, 0.5 g
24 MgSO₄·7H₂O, 0.1 g CaCl₂, 0.1 g NaCl, 0.5 mg H₃BO₃,
25 0.2 mg FeCl₃·4H₂O, 0.4 mg ZnSO₄·7H₂O, 0.4 mg
26 MnSO₄·H₂O, 0.1 mg KI, 0.04 mg CuSO₄·5H₂O, 200 µg
27 thiamin, 8 µg D-biotin. The medium without urea and vita-
28 mins was sterilized at 121 °C for 20 min. After cooling, the
29 vitamins and urea previously sterilized by microfiltration
30 (0.22 µm) were added to the basal medium.

2.2 Culture conditions 31

32 First, the microorganism was grown in 0.250 dm³
33 Erlenmeyer flasks containing 0.15 dm³ of basal medium,
34 in order to obtain the inoculum for oil degradation and lipase
35 production. A volume of 0.01 dm³ of 0.9% NaCl solution was
36 added to a previously grown agar plate, and 10⁻³ dm³ of the
37 resulting suspension was added to the liquid medium as
38 inoculum. The flasks were incubated in an orbital shaker
39 (New Brunswick, Innova 4400) at 30 °C, 150 rpm, for 48 h.

40 Flask cultures for the investigation of waste oil biodegra-
41 dation and lipase production were carried out in 0.25 dm³
42 Erlenmeyer flasks containing 0.15 dm³ of basal medium, to
43 which a fixed concentration (1–30 g dm⁻³) of waste cooking
44 oil was added. This oil was collected in local restaurants and
45 was mainly composed by oleic acid (31%) and linoleic acid
46 (55%), with a free fatty acid content of 5.2%. The culture
47 medium was emulsified for 7 min at 995 rad·s⁻¹
48 (Homogeniser IKA T25) before autoclaving. The flasks were
49 inoculated 2% v/v with the freshly prepared cultures of
Yarrowia lipolytica, and incubated in an orbital shaker at

1 150 rpm and 30 °C. They were capped with cellulose stop-
2 pers, which permit passive aeration. Some experiments were
3 also performed on the same medium with only 5 g dm⁻³
4 glucose and 20 g dm⁻³ oil. Control cultures were carried out
5 without addition of waste oil.

6 In order to evaluate and compare the characteristics of the
7 produced enzymes, additional flask cultures were performed,
8 using the basal medium with 5 g dm⁻³ glucose and 10 g
9 dm⁻³ olive oil, sunflower oil and waste-cooking oil,
10 respectively.

11 Bioreactor cultures were carried out in a 5-L stirred tank
12 bioreactor (Biostat B, Braun, Germany), containing 3 L of
13 basal medium with 5 g dm⁻³ glucose and 30 g dm⁻³ waste
14 cooking oil. Airflow and agitation speed were 1 L min⁻¹ and
15 400 rpm, respectively.

16 Experiments were done in duplicate and samples were
17 analyzed in triplicate. The values in the figures correspond to
18 mean values with a deviation less than 15%.

19 2.3 Sample preparation

20 In all cases, samples (whole flask content or 10 mL sample in
21 flask and bioreactor cultures, respectively) were centrifuged
22 for 10 min at 5000 g and 5 °C. The supernatant was recov-
23 ered, homogenized at 9500 rpm for 1 min (Homogenizer
24 IKA T25), and utilized for COD and lipolytic activity deter-
25 mination. The solid residue was thoroughly washed, and
26 resuspended in distilled water for biomass determination.

27 2.4 Analytical methods

28 Lipolytic activity was determined spectrophotometrically
29 using *p*-nitrophenyl myristate (pNPM) as substrate.
30 The reaction medium consisted of 10⁻⁴ dm³ sample,
31 8 × 10⁻⁴ dm³ 50 mmol dm⁻³ Tris-HCl buffer (pH 7.2)
32 containing 40 mM CaCl₂, and 10⁻⁴ dm³ pNPM 25 mmol
33 dm⁻³ dissolved in ethanol. The hydrolysis reaction was
34 allowed to proceed for 20 min at 30 °C and afterwards
35 stopped by adding 2.5 × 10⁻⁴ dm³ of 2 mmol dm⁻³
36 Na₂CO₃. The mixture was kept on ice for at least 15 min
37 before being centrifuged (5000 g, 10 min). The absorbance
38 of the supernatant was monitored at 400 nM in a Unicam
39 Helios β (Thermo Electron Corp.) spectrophotometer.
40 Under alkaline conditions, the molar extinction coefficient
41 was 17.215 ± 0.002 mmol⁻¹ cm⁻¹. One activity unit was
42 defined as the amount of enzyme that produced 1 μmol of
43 *p*-nitrophenol per min under standard assay conditions. The
44 activities were expressed in U dm⁻³.

45 Biomass concentration was measured via turbidimetry at
46 600 nm, and the obtained values were converted to g cell dry
47 wt dm⁻³ using a previously determined calibration curve.

48 The COD was determined following standard method
49 UNE 77004 [37]. Oil-associated COD in the cultures was
50 calculated by subtracting the COD obtained in control
cultures, to which oil had not been added.

2.5 Enzyme characterization

2 Enzyme activity-pH profiles were determined by measuring
3 lipolytic activity on *p*-nitrophenyl myristate, as indicated
4 above, using different buffer solutions in the reaction mix-
5 ture: citrate-phosphate for pH 4, phosphate for pH 6-7, Tris-
6 HCl for pH 8-9. Enzyme activity-temperature profiles were
7 obtained by determination of lipolytic activity by the
8 described method, at pH 5.0 and different incubation tem-
9 peratures (30–60 °C). The results were expressed as percent-
10 ages, referred to the maximum activity value.

11 Enzyme stability toward pH was determined by incubat-
12 ing samples in suitable buffer solutions at 30 °C, and measur-
13 ing residual lipolytic activity after 24 h. Enzyme
14 thermostability was assessed by incubation of samples in
15 a pH 5.0 buffer at several temperatures, and measuring
16 residual lipolytic activity after 1 h.

3 Results And Discussion

17 There are several end-uses for waste frying oils, such as the
18 production of soaps or of energy by anaerobic digestion,
19 thermal cracking, and more recently, the production of bio-
20 diesel. The development of a procedure allowing simul-
21 taneous biodegradation and reutilization of waste cooking
22 oils, while producing a high added-value product would be
23 of great interest, both from an economic and environmental
24 point of view. In this work, used up vegetable frying oil has
25 been added to the culture medium of *Yarrowia lipolytica*
26 CECT 1240, a lipase producer. The level of oil biodegrada-
27 tion has been assessed, as well as the ability of this residual
28 material to act as lipase production inducer by the
29 microorganism.
30

3.1 Shake-flask cultures

31 Biodegradation of waste cooking oil in submerged cultures of
32 *Yarrowia lipolytica* CECT 1240 has been investigated.
33 Initially, the yeast was grown in Erlenmeyer flasks on a liquid
34 medium to which a certain amount (20 g dm⁻³) of the spent
35 oil was added, and the variation in COD after 6-days culture
36 time was compared to that observed in control cultures (no
37 added oil). Two different initial glucose concentrations
38 (5 and 20 g dm⁻³) in the culture medium were considered,
39 as model concentrations providing different levels of cell
40 growth [38]. The aim of this experiment was to find out if
41 carbon source depletion could further oil consumption by the
42 microorganism as reported Carvalho *et al.* [39]. A glucose
43 consumption rate of 1.8 g dm⁻³ day⁻¹ was observed through-
44 out the cultivation period (6-days). In the culture with 5 g
45 dm⁻³ of glucose, total glucose depletion was found after 3
46 days. Nevertheless, 8 g dm⁻³ of glucose remains in the media
47 carried out with 20 g dm⁻³ after the culture time. The results
48 confirmed the ability of *Y. lipolytica* CECT 1240 to metab-
49 olize waste cooking oil, since around 80 and 50% decrease in

oil-associated COD was observed at the end of the cultivation time with 5 and 20 g dm⁻³ of glucose, respectively. These data suggest that glucose depletion furthers consumption of the waste cooking oil by the microorganism.

The highest biomass levels yielded (9 g dm⁻³) were obtained for the culture containing 20 g dm⁻³ of glucose. The values are lower than those reported by Corzo and Revah and Deive *et al.* [23, 38], which reported maximum values of biomass of 30 and 17 g dm⁻³, respectively, when they supplemented the basal medium with olive oil. This fact indicates that the waste cooking oil composition might exert a considerable influence at cell metabolism level. On account of the higher degradation levels attained with low glucose concentration, 5 g dm⁻³ initial glucose was added to the culture medium in all subsequent experiments.

Afterwards, the microorganism was grown in shake flasks to which different concentrations of waste oil were added (1–30 g dm⁻³). COD evolution during the cultures is shown in Fig. 1. The importance of evaluating initial lipidic concentration has already been stressed by Papanikolaou *et al.* [40], who reported that stearin (trioctadecanoin, or a glyceryl ester of stearic acid) concentrations higher than 10 g dm⁻³ were adequate for lipid accumulation in *Y. lipolytica*. Deive *et al.* [38] determined that 3 days was the minimum time required to reach the stationary phase by *Y. lipolytica* CECT 1240 in analogous environmental conditions. Therefore, samples were taken out at 3 and 6 days, to have representative values of degradation. Final degradation close to 100% was attained for the lowest oil concentrations (1 and 5 g dm⁻³), while around 80% of the waste oil was metabolized after 6 days culture in the other cases. As it has been previously reported in several research works [40, 41, 15] fatty acid composition of the fatty substrate can be playing a major role in the lipid

uptake by *Y. lipolytica*, explaining thus the lower percentages of degradation at high lipidic concentrations. Cell growth could also be affected by this variable. Additionally, the values are comparable to those obtained by Lanciotti *et al.* [36]. They analyzed the COD reduction after 3 days of cultivation by 62 strains of *Y. lipolytica*, and obtained that just three strains were able to provide about 30–40% of COD reduction after 3 days of cultivation. Hence, the results obtained here support the suitability of the studied strain to treat waste cooking oil.

Additionally, extracellular lipolytic activity was measured in the culture medium (Fig. 2) and significant enzyme concentrations were found in experiments supplemented with waste cooking oil. Domínguez *et al.* and Deive *et al.* [27, 28, 38] pointed out the necessity of the addition of vegetable oils in order to trigger the lipase synthesis by this specific strain, and the possibility of using waste oil instead of non-used oil is confirmed in this work. This fact is of interest, because it entails the valorization of a residue as enzyme inducer. Maximum enzyme activity increased with oil concentration up to 5 g dm⁻³, then decreased, but in all cases enzyme production was higher when waste oil had been added to the culture medium. It should be emphasized that a limitation of the lipase production has been reported in other investigations when increased concentrations of oil were used in the medium, due to oxygen transfer limitations [42]. On the other hand, some authors have described decreased lipase concentrations as oil (*i.e.*, carbon source) was depleted in the culture medium [35, 38].

Also, a linear relationship could be established between the decrease in COD rate (mg O₂...dm⁻³ day⁻¹) and the oil concentration, as it can be seen in Fig. 3. Higher contents of waste cooking oil led to higher decrease in COD, which seems




Figure 1. Evolution of oil-associated COD in shake-flask cultures of *Yarrowia lipolytica* CECT 1240 on basal medium containing 5 g dm⁻³ glucose and different concentrations of waste cooking oil, at 30 °C and 150 rpm: (●) 1 g dm⁻³, (○) 5 g dm⁻³, (▲) 10 g dm⁻³, (△) 20 g dm⁻³ and (▽) 30 g dm⁻³.




Figure 2. Lipolytic activity in shake-flask cultures of *Yarrowia lipolytica* CECT 1240 on basal medium containing 5 g dm⁻³ glucose and different concentrations of waste cooking oil, at 30 °C and 150 rpm: (◆) No added oil, (●) 1 g dm⁻³, (○) 5 g dm⁻³, (▲) 10 g dm⁻³, (△) 20 g dm⁻³ and (▽) 30 g dm⁻³.




Figure 3. Linear relationship between decrease in COD and waste cooking oil concentration in shake-flask cultures of *Yarrowia lipolytica* CECT 1240 on basal medium containing 5 g dm⁻³ glucose, at 30 °C and 150 rpm (COD = 143.18 · [Oil concentration]).

to indicate that COD reduction is not solely associated with lipase production, but also with the synthesis of other metabolites during the biological process. Following this behavior, Lanciotti *et al.* [36] also described different trends for COD reduction and lipolytic enzyme activity in cultures of several yeasts.

An additional promising application of this strain could be the treatment of other oil-based wastes, such as OMW. The residual oil that OMW contains in relatively large quantities (from 0.3 to 3.5 g dm⁻³ depending on the extraction process efficiency) makes this waste a potentially suitable medium for lipase production, as it has been recently reported [43–45].

3.2 Bioreactor cultures

Taking into account the results achieved in flask cultures, the efficiency of the studied strain for waste cooking oil degradation and lipase production was assessed in a bench scale bioreactor. A medium containing the highest oil concentration (30 g dm⁻³) and lowest amount of glucose (5 g dm⁻³) previously assayed was employed, and COD and lipolytic activity were monitored during the cultivation time (6 days). The results are shown in Fig. 4. More rapid oil consumption was recorded in the bioreactor, compared to flask cultures. Almost 90% of initial COD was eliminated in just 3 days, taking into account that glucose consumption rate is 10 g dm⁻³ day⁻¹. This degradation level is similar to that reported by Scioli and Vollarò [46], which indicated about 80% COD reduction after carrying out the biological process in a 3.5 L fermenter. The same value was also obtained by De Felice *et al.* [47] using *Y. lipolytica* 20255. On the other hand, lipolytic enzyme levels progressively increase from day 2 onwards, attaining values close to 800 U dm⁻³ after 6-days culture time.




Figure 4. COD and lipolytic activity in *Yarrowia lipolytica* CECT 1240 cultures carried out in a stirred tank bioreactor (400 rpm, 1 dm³ min⁻¹ airflow), on basal culture medium containing 5 g dm⁻³ glucose and 30 g dm⁻³ waste cooking oil: (•) COD, (○) Lipolytic activity.

In view of the rapid COD reduction and high lipase production levels observed, the feasibility of a fed-batch strategy was assessed, as a means to further the usefulness of the culture for COD reduction and waste cooking oil valorization. Deive *et al.* [38] had previously assayed the culture of *Y. lipolytica* CECT 1240 in fed-batch and continuous mode, and obtained very promising results. Thus, 30 g of waste cooking olive oil were added after 6 days of cultivation, and COD and enzyme activity levels were monitored afterwards. The main results are presented in Table 1. The studied strain was capable of reducing COD values nearly by 50% within 24 h of oil addition. Also, further increase in enzyme activity was observed. Thus, the microorganism was able to degrade additional waste cooking oil while producing lipolytic enzymes after a new addition of the residue, although at a lower rate than before. These results support the interest of the studied strain, and its potential for the treatment of oily wastes.

Moreover, it is noteworthy that the maximum lipolytic enzyme production attained in the bioreactor culture (927 U dm⁻³) was more than 5-fold higher than that detected in flask cultures when an analogous culture medium was used. This fact might be related to the specific hydrodynamic conditions inside the bioreactor, in terms of agitation and aeration. On the one hand, an increased transfer area is furthered by forming small sized bubbles which are dispersed in the culture medium, increasing their residence time on the liquid. On the other hand, a good agitation favors turbulence which leads to a decrease on the thickness of the liquid layer on the gas/liquid interface, reducing the resistance to oxygen mass transfer. This statement is in agreement with the conclusions reported by Lopes *et al.* [48], who recorded an increase of 96% in lipase activity values when used a 5-bar air pressure instead of 1-bar pressure during the culture of *Yarrowia*

Table 1. COD reduction and lipolytic activity production during fed batch operation in *Yarrowia lipolytica* CECT 1240 cultures carried out in a stirred tank bioreactor (400 rpm, 1 dm³ min⁻¹ airflow). Oil addition: 30 g waste cooking oil added at day 6.

	Before oil addition	After oil addition
Maximum lipolytic activity (U dm ⁻³)	778	927
Lipolytic enzyme production rate (U dm ⁻³ day ⁻¹)	129.7	21.3
Maximum COD reduction (%)	89.4	51.6
Maximum COD reduction rate (mg O ₂ dm ⁻³ day ⁻¹)	10 182	5454

1 *lipolytica*. Also, an efficient agitation could improve the contact between the microorganism and the oil, thus facilitating
2 its utilization. However, the relevance of the strain selected
3 must be taken into account since, contrarily to the results
4 presented here, Alonso *et al.* [49] concluded that increased
5 agitation and aeration rates negatively affected lipase production by another strain of *Y. lipolytica*.

6
7
8 In summary, the studied biological system appears as a
9 promising tool for the treatment and valorization of oily
10 residues from the food industry, such as waste cooking oil.

11 3.3 Enzyme characteristics


12 Some characteristics of the crude extracts (*i.e.*, cell-free concentrated extracellular liquid) were assayed in order to evaluate if the use of waste cooking oil as carbon source and lipase inducer, instead of non-used oil, could alter the main catalytic properties of the synthesized enzymes. Thus, shake flask cultures of *Y. lipolytica* CECT 1240 were carried out in culture media supplemented with waste cooking oil, non-used olive oil and non-used sunflower oil. Identical environmental conditions were employed in all cases, and the enzyme crude solutions were recovered at the end of the cultures.

22

1 First, the influence of temperature and pH on lipase activity was assessed in the enzyme solutions obtained from the abovementioned cultures. The selected operating conditions were a temperature range of 30–60 °C and a pH varying between 4 and 9, respectively. As it can be seen in Fig. 5, the enzyme extracts showed maximum activity at pH values around 7.0, as a number of microbial lipases described in previous reports, such as those from *Bacillus* sp., *Acinetobacter* sp., and a mutant *Yarrowia lipolytica* strain [50, 51, 26]. The activity-pH profiles were rather close for the three enzyme solutions assayed, and more so for those obtained in waste cooking oil and sunflower oil-supplemented cultures. Since sunflower oil is commonly used as commercial frying oil in our geographical area, it would be reasonable to expect a certain degree of similarity between the effects of both oils, related to their basic composition.

17 As for enzyme thermoactivity, a crucial feature in industrial biocatalysis, all the crude enzymes showed maximum activities at temperature values between 30 and 40 °C, and the activity-temperature profiles were also quite similar. These values of thermoactivity are typical for lipases obtained from mesophilic microorganisms, and have been found for other strains of this yeast, such as *Y. lipolytica* 681 [23].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

**Figure 5.** Influence of pH and temperature on lipolytic enzyme activity for different *Yarrowia lipolytica* CECT 1240 enzyme extracts: (●) Cultures with sunflower oil, (○) Cultures with olive oil, (▲) Cultures with waste cooking oil.




Figure 6. Influence of pH and temperature on lipolytic enzyme stability for different *Yarrowia lipolytica* CECT 1240 enzyme extracts: (●) Cultures with sunflower oil, (○) Cultures with olive oil, (▲) Cultures with waste cooking oil.

Thermal stability is another important property, often investigated during the evaluation of enzyme characteristics and applicability. The deactivation of the three enzyme extracts was assessed after 24 h incubation at 30 °C and different pH values ranging from 3.0 to 9.0. The results are shown in Fig. 6. The observed profiles follow a similar trend. All the enzyme solutions show the highest stability in acidic environments, while a dramatic fall in stability is registered at pH values over 6.5.

Regarding thermal stability (at pH 5), the enzymes produced in cultures supplemented with sunflower oil and waste-cooking oil appeared to be more resistant to thermal deactivation than that obtained from olive oil-supplemented cultures (Fig. 6). This effect is noticeable at temperatures above 40 °C. Some authors have reported the production of different lipolytic isoenzymes by strains of *Y. lipolytica*, and they identified the genes associated with their production [52, 34]. Those isoenzymes showed different biochemical properties. In the present case, the culture conditions (*i.e.*, type of added oil) could lead to variation in the isoenzymes ratio in the enzyme extracts, and this could possibly explain the differences observed between their properties.

4 Conclusions

The necessity of new strategies to valorize wastes from the food and agro industries supports the interest of the present work. The potential application of *Yarrowia lipolytica* CECT 1240 to degrade waste cooking oils is confirmed, as well as the efficiency of this residue to promote the production of lipolytic enzymes by the strain. The process was successfully carried out both in shake flasks and a bench-scale bioreactor, allowing producing high levels of lipolytic activity at the same time that the COD was diminished up to nearly 90%. Besides, the efficient degradation of repeated waste oil

charges was ascertained. Also, the catalytic potential of the crude enzymes obtained in cultures supplemented with waste cooking oil was assessed, and they compared favorably with those obtained when non-used oils were added to the culture medium.

This work was financed by Xunta de Galicia (Project PGIDIT06PXIB314376PR). Francisco Deive thanks Fundación Juana de Vega for a postdoctoral grant.

The authors have declared no conflict of interest.

References

- [1] Cvengroš, J., Cvengrošová, Z., Used frying oils and fats and their utilization in the production of methyl esters of higher fatty acids. *Biomass Bioenerg.* 2004, 27, 173–181.
- [2] Isidori, M., Parrella, A., Genotoxicity of aqueous extract from heated cooking oils and its suppression by *Lactobacilli*. *Food Sci. Technol. Int.* 2009, 15, 267–273.
- [3] Mulflur, M. W. Munson: J. R. US Patent 4681768 (1987).
- [4] Takeno, K., Yamaoka, Y., Sasaki, K., Treatment of oil-containing sewage wastewater using immobilized photosynthetic bacteria. *World J. Microbiol. Biotechnol.* 2005, 21, 1385–1391.
- [5] Chua: A. S. PCT Patent WO9830668 (1998).
- [6] Adamczak, M., Bornscheuer, W. T., Bednarski, W., The application of biotechnological methods for the synthesis of biodiesel. *Eur. J. Lipid Sci. Technol.* 2009, 111, 800–813.
- [7] Zhang, Y., Dubé, M. A., Malean, D. D., Kates, M., Biodiesel production from waste cooking oil: 1. Process design and technological assessment. *Bioresour. Technol.* 2003a, 89, 1–16.
- [8] Zhang, Y., Dubé, M. A., Malean, D. D., Kates, M., Biodiesel production from waste cooking oil: 2. Economic assessment and sensitivity analysis. *Bioresour. Technol.* 2003b, 90, 229–240.

- [9] Çetinkaya, M., Ulusoy, Y., Tekin, Y., Karaosmanoğlu, F., Engine and winter road test performances of used cooking oil originated biodiesel. *Energy Conv. Manage.* 2005, 46, 1279–1291.
- [10] Zaher, F. A., Megahed, O. A., El Kinawy, O. S., Utilization of used frying oil as diesel engine fuel. *Energy Sources* 2003, 25, 819–826.
- [11] Battersby, N. S., The biodegradability and microbial toxicity testing of lubricants – some recommendations. *Chemosphere* 2000, 41, 1011–1027.
- [12] Pereira, M. G., Mudge, S. M., Latchford, J., Vegetable oil spills on salt marsh sediments; comparison between sunflower and linseed oils. *Mar. Environ. Res.* 2003, 56, 367–385.
- [13] Li, Z., Wrenn, B. A., Venosa, A. D., Anaerobic biodegradation of vegetable oil and its metabolic intermediates in oil-enriched freshwater sediments. *Biodegradation.* 2005, 16, 341–352.
- [14] Li, Z., Wrenn, B. A., Effects of ferric hydroxide on the anaerobic biodegradation kinetics and toxicity of vegetable oil in freshwater sediments. *Water Res.* 2004, 38, 3859–3868.
- [15] Papanikolaou, S., Aggelis, G., Selective uptake of fatty acids by the yeast *Yarrowia lipolytica*. *Eur. J. Lipid Sci. Technol.* 2003, 105, 651–655.
- [16] Bednarski, W., Adamczak, M., Tomasik, J., Paszczyk, M., Application of oil refinery waste in the biosynthesis of glycolipids by yeast. *Bioresour. Technol.* 2004, 95, 15–18.
- [17] Sharma, R., Chisti, Y., Banerjee, U., Production, purification, characterization and applications of lipases. *Biotechnol. Adv.* 2001, 19, 627–662.
- [18] Gotor, V., Biocatalysis applied to the preparation of pharmaceuticals. *Org. Process Res. Dev.* 2002, 6, 420–426.
- [19] Hills, G., Industrial use of lipases to produce fatty acid esters. *Eur. J. Lipid Sci. Technol.* 2003, 105, 601–607.
- [20] Abate, A., Brenna, E., Fuganti, C., Gatti, F. G., Serra, S., Lipase-catalysed preparation of enantiomerically enriched odorants. *J. Mol. Catal. B-Enzym.* 2004, 32, 33–51.
- [21] Reetz, M. T., Lipases as practical biocatalysts. *Curr. Opin. Chem. Biol.* 2002, 6, 145–150.
- [22] Gupta, R., Gupta, N., Rathi, P., Bacterial lipases: an overview of production, purification and biochemical properties. *Appl. Microbiol. Biotechnol.* 2004, 64, 763–781.
- [23] Corzo, G., Revah, S., Production and characteristics of the lipase from *Yarrowia lipolytica* 681. *Bioresour. Technol.* 1999, 70, 173–180.
- [24] Ota, Y., Gomi, K., Kato, S., Sugiura, T., Minoda, Y., Purification and some properties of cell-bound lipase from *Saccharomycopsis lipolytica*. *Agric. Biol. Chem.* 1982, 46, 2885–2893.
- [25] Pereira-Meirelles, F. V., Rocha-Leao, M. H. M., Sant’Ana, G. L. J., Lipase location in *Yarrowia lipolytica* cells. *Biotechnol. Lett.* 2000, 22, 71–75.
- [26] Destain, J., Roblain, D., Thonart, P., Improvement of lipase production from *Yarrowia lipolytica*. *Biotechnol. Lett.* 1997, 19, 105–107.
- [27] Domínguez, A., Deive, F. J., Sanroman, M. A., Longo, M. A., Effect of lipids and surfactants on extracellular lipase production by *Yarrowia lipolytica*. *J. Chem. Technol. Biotechnol.* 2003a, 78, 1166–1170.
- [28] Domínguez, A., Costas, M., Longo, M. A., Sanromán, M. A., A novel application of solid state culture: production of lipases by *Yarrowia lipolytica*. *Biotechnol. Lett.* 2003b, 25, 1225–1229.
- [29] Ionita, A., Moscovici, M., Dragolici, A., Eremia, M. *et al.*, Lipase production in discontinuous operation system using a *Candida lipolytica* strain. *Roumanian Biotechnol. Lett.* 2002, 7, 547–552.
- [30] Nicaud, J. M., Madzak, C., van den Broek, P., Gysler, C. *et al.*, Protein expression and secretion in the yeast *Yarrowia lipolytica*. *FEMS Yeast Res.* 2002, 2, 371–379.
- [31] Fickers, P., Nicaud, J. M., Destain, J., Thonart, P., Overproduction of lipase by *Yarrowia lipolytica* mutants. *Appl. Microbiol. Biotechnol.* 2003, 63, 136–142.
- [32] Fickers, P., Nicaud, J. M., Gaillardin, C., Destain, J., Thonart, P., Carbon and nitrogen sources modulate lipase production in the yeast *Yarrowia lipolytica*. *J. Appl. Microbiol.* 2004, 96, 742–749.
- [33] Fickers, P., Fudalej, F., Nicaud, J. M., Destain, J., Thonart, P., Selection of new over-producing derivatives for the improvement of extracellular lipase production by the non-conventional yeast *Yarrowia lipolytica*. *J. Biotechnol.* 2005a, 115, 379–386.
- [34] Fickers, P., Fudalej, F., Le Dall, M. T., Casaregola, S. *et al.*, Identification and characterisation of LIP7 and LIP8 genes encoding two extracellular triacylglycerol lipases in the yeast *Yarrowia lipolytica*. *Fungal Genet. Biol.* 2005b, 42, 264–274 (2005b).
- [35] Papanikolaou, S., Chevalot, I., Galiotou-Panayotou, M., Komaitis, M. *et al.*, Industrial derivative of tallow: a promising renewable substrate for microbial lipid, single-cell protein and lipase production by *Yarrowia lipolytica*. *Electron. J. Biotechnol.* 2007, 10, 425–435.
- [36] Lanciotti, R., Gianotti, A., Baldi, D., Angrisani, R. *et al.*, Use of *Yarrowia lipolytica* strains for the treatment of olive mill wastewater. *Bioresour. Technol.* 2005, 96, 317–322.
- [37] AENOR. 2002, UNE 77004: Calidad del agua. Determinación de la demanda química de oxígeno (DQO). Método del dicromato. (Water Quality – Determination of Chemical Oxygen Demand (COD)).
- [38] Deive, F. J., Sanroman, M. A., Longo, M. A., A comprehensive study of lipase production by *Yarrowia lipolytica* CECT 1240 (ATCC 18942): from shake flask to continuous bioreactor. *J. Chem. Technol. Biotechnol.* 2010, 85, 258–266.
- [39] Carvalho, M. B., Martins, I., Leitão, M. C., Garcia, H. *et al.*, Screening pentachlorophenol degradation ability by environmental fungal strains belonging to the phyla *Ascomycota* and *Zygomycota*. *J. Ind. Microbiol. Biotechnol.* 2009, 36, 1249–1256.
- [40] Papanikolaou, S., Chevalot, I., Komaitis, M., Marc, I., Aggelis, G., Single cell oil production by *Yarrowia lipolytica* growing on an industrial derivative of animal fat in batch cultures. *Appl. Microbiol. Biotechnol.* 2002, 58, 308–312.
- [41] Papanikolaou, S., Chevalot, I., Komaitis, M., Aggelis, G., Marc, I., Kinetic profile of the cellular lipid composition in an oleaginous *Yarrowia lipolytica* capable of producing a cocoa-butter substitute from industrial fats. *Antonie van Leeuwenhoek* 2001, 80, 215–224.
- [42] Lima, V. G. M., Krieger, N., Sarquis, M. I. M., Mitchell, D. A. *et al.*, Effect of nitrogen and carbon sources on lipase production by *Penicillium aurantiogriseum*. *Food Technol. Biotechnol.* 2003, 41, 105–110.

- 1 [43] D'Annibale, A., Sermanni, G., Federici, F., Petruccioli, M.,
2 Olive-mill wastewaters: a promising substrate for microbial
3 lipase production. *Bioresour. Technol.* 2006, 97, 1828–1833.
- 4 [44] Federici, F., Fava, F., Kalogerakis, N., Mantzavinos, D.,
5 Valorisation of agroindustrial by products effluents and
6 waste: concept, opportunities and the case of olive mill
7 wastewaters. *J. Chem. Technol. Biotechnol.* 2009, 84, 895–
8 900.
- 9 [45] Gonçalves, C., Lopes, M., Ferreira, J. P., Belo, I., Biological
10 treatment of olive mill wastewater by non-conventional
11 yeasts. *Bioresour. Technol.* 2009, 100, 3759–3763.
- 12 [46] Scioli, C., Vollaro, L., The use of *Yarrowia lipolytica* to reduce
13 pollution in olive mill wastewaters. *Water Resour.* 1997, 31,
14 2520–2524.
- 15 [47] De Felice, B., Pontecorvo, G., Carfagna, M., Degradation of
16 waste water from olive oil mills by *Yarrowia lipolytica* ATCC
17 20255 and *Pseudomonas putida*. *Acta Biotechnol.* 1997, 17,
18 231–239.
- [48] Lopes, M., Gomes, N., Mota, M., Belo, I., *Yarrowia lipolytica*
1 growth under increased air pressure: influence on enzyme pro-
2 duction. *Appl. Biochem. Biotechnol.* 2009, 159, 46–53. 3
- [49] Alonso, F. O. M., Oliveira, E. B. L., Dellamora-Ortiz, G. M.,
4 Pereira-Meirelles, F. V., Improvement of lipase production at
5 different stirring speeds and oxygen levels. *Braz. J. Chem.*
6 *Eng.* 2005, 22, 9–18. 7
- [50] Sugihara, A., Tadaaki, T., Tominaga, Y., Purification and
8 characterization of a novel thermostable lipase from *Bacillus*
9 sp. *J. Biochem.* 1991, 109, 211–216. 10
- [51] Barbaro, S. E., Trevors, J. T., Inniss, W. E., Effect of low
11 temperature, cold shock, and various carbon sources on
12 esterase and lipase activities and expolysaccharide pro-
13 duction by a psychotrophic *Acinetobacter* sp. *Can. J.*
14 *Microbiol.* 2001, 47, 194–205. 15
- [52] Pignede, G., Wang, H., Fudalej, F., Gaillardin, C. *et al.*,
16 Characterization of an extracellular lipase encoded by LIP2
17 in *Yarrowia lipolytica*. *J. Bacteriol.* 2000, 182, 2802–2810. 18
19
20