

Biofouling in marine aquaculture: a review of recent research and developments

Jana Bannister^a , Michael Sievers^b , Flora Bush^a  and Nina Bloecher^c 

^aInstitute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia; ^bAustralian Rivers Institute - Coast and Estuaries, Griffith University, Gold Coast, Queensland, Australia; ^cSINTEF Ocean, Trondheim, Norway

ABSTRACT

Biofouling in marine aquaculture is one of the main barriers to efficient and sustainable production. Owing to the growth of aquaculture globally, it is pertinent to update previous reviews to inform management and guide future research. Here, the authors highlight recent research and developments on the impacts, prevention and control of biofouling in shellfish, finfish and seaweed aquaculture, and the significant gaps that still exist in aquaculturalists' capacity to manage it. Antifouling methods are being explored and developed; these are centred on harnessing naturally occurring antifouling properties, culturing fouling-resistant genotypes, and improving farming strategies by adopting more sensitive and informative monitoring and modelling capabilities together with novel cleaning equipment. While no simple, quick-fix solutions to biofouling management in existing aquaculture industry situations have been developed, the expectation is that effective methods are likely to evolve as aquaculture develops into emerging culture scenarios, which will undoubtedly influence the path for future solutions.

ARTICLE HISTORY

Received 24 May 2019
Accepted 2 July 2019

KEYWORDS

Antifouling; fish; bivalve; seaweed; epiphyte; net


Introduction

In 2016, global aquaculture production was 110.2 million tonnes (US\$243.5 billion), including 54.1 million tonnes of finfish (US\$138.5 billion), 17.1 million tonnes of molluscs (US\$29.2 billion) and 30.1 million tonnes of aquatic plants (US\$11.7 billion) (FAO 2018). Despite annual growth rates of global aquaculture slowing to an average of 5.8% during the period 2000–2016, aquaculture continues to grow faster than other major food production sectors (FAO 2018). More specifically, marine shellfish aquaculture has expanded substantially over the last few decades, with Asia dominating production and accounting for >90% of global tonnage (FAO 2018). Supped oysters (*Crassostrea* spp.) and the Japanese carpet shell (*Ruditapes philippinarum*; a clam) account for more than half of global production (FAO 2018). Finfish aquaculture has increased its volume by >70% over the last decade and freshwater fish, most of them cyprinids, make up 86%. Salmonids, including Atlantic salmon (*Salmo salar*), which are predominantly featured in the scientific literature on biofouling in finfish aquaculture and its management, account for 43% of non-freshwater aquaculture (FAO 2018). Finally,

aquatic plants are harvested globally for direct human consumption, and for the production of thickening agents for pharmaceutical and cosmetics, and dried ingredients for animal feed, fertilisers and other products (FAO 2018). Global aquatic plant production has tripled from 10.1 million tonnes in 2000 to 30.1 million tonnes in 2016 (FAO 2002, 2018) with the increased production of the tropical seaweeds *Kappaphycus alvarezii* and *Euचेuma* spp. in Indonesia (as raw material for carrageenan extraction) being the major contributor to this growth (FAO 2018).

Biofouling in marine aquaculture is one of the main barriers to efficient and sustainable production (Dürr and Watson 2010). It is the settlement and development of unwanted aquatic species on natural and artificial surfaces, and plagues shellfish, finfish and seaweed culture globally. The direct economic costs of managing biofouling in the aquaculture industry are estimated to be 5–10% of production costs (Lane and Willemsen 2004). However, the cost of biofouling often varies considerably between aquaculture locations, species and companies, as farmers use differing management approaches and cost

CONTACT Jana Bannister  jana.bannister@utas.edu.au

 Supplemental data for this article can be accessed at <https://doi.org/10.1080/08927014.2019.1640214>.

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

accounting (Iversen et al. 2016). Furthermore, many of the indirect impacts remain largely unassessed, so the overall economic cost of biofouling in marine aquaculture is still uncertain but is likely to be significantly underestimated (Fitridge et al. 2012).

The aim of this review article is to present and discuss recent research and developments on the impacts, prevention and control of biofouling in marine aquaculture, and thereby update the seminal review article by Fitridge et al. (2012). In addition to marine shellfish and finfish aquaculture, the authors also include a comprehensive evaluation of epiphytism/biofouling in seaweed aquaculture, as this industry has grown significantly over recent years.

The impact of biofouling on shellfish aquaculture

A wide variety of biofouling organisms are associated with shellfish aquaculture globally (Table S1 Supplemental material). The impacts of these species growing on shellfish and farm infrastructure can be partitioned into three broad categories: (1) increased weight, (2) physical damage, and (3) reductions to shellfish fitness. These impacts, along with the cost of biofouling management, can culminate in substantially reduced farm productivity.

Increased weight

Biofouling adds considerable weight to both stock and culture equipment, causing stock detachment in long-line culture and increasing the costs associated with buoyancy and anchoring systems (see Fitridge et al. 2012 and references within). Relative to finfish and seaweed culture, the impacts of increases to weight to shellfish culture are less frequently quantified, so impacts to productivity are comparatively unknown.

Physical damage

Boring organisms, such as clionid sponges and polychaete worms from the genus *Polydora*, can penetrate shellfish, causing blisters, cavities and tunnels (Sato-Okoshi and Abe 2013; Carroll et al. 2015). This causes disrupted shell and hinge formation and results in brittle, thin shells that are susceptible to parasites, disease and predation (see Fitridge et al. 2012 and references within). Several other calcareous fouling species settle on shellfish and do not cause direct damage to the shellfish but do affect aesthetics and often lead to the product being devalued or discarded, with

considerable financial losses to affected farms (Campbell and Kelly 2002).

Reductions to shellfish fitness

One of the most important impacts from biofouling in shellfish aquaculture is the reduction in shellfish fitness (ie survival, growth, condition and weight). These reductions are typically attributed to direct competition for food, oxygen and other resources, or indirectly, *via* smothering or interfering with proper valve functioning (Lodeiros and Himmelman 1996; Pit and Southgate 2003; Woods et al. 2012). In an empirical study, Sievers et al. (2013) added biofouling to experimental mussel lines and found clear evidence of commercially relevant reductions in growth and flesh weights in mussels (*Mytilus galloprovincialis*) fouled by the widespread tunicates *Ciona intestinalis* and *Styela clava* and the hydroid *Ectopleura crocea*. However, in general, scant evidence exists in the literature to confirm fitness reductions (Sievers et al. 2017). Much of the evidence comes from observational studies, where initial differences between fouled and unfouled stock such as health, size and antifouling (AF) capacity may confound observed patterns. For example, slower growing stock may be more susceptible to the settlement of fouling on shells, potentially leading to erroneous conclusions that fouling reduces growth where fouled and unfouled stock are compared. Despite a general lack of empirical evidence, considerable observational and anecdotal evidence (eg Daigle and Herbingier 2009; Fitridge and Keough 2013) strongly suggests that biofouling can impact cultured shellfish fitness and thus warrants considerable attention from the industry.

Farm productivity

The aforementioned impacts coupled with the substantial cost of removal has spurred considerable interest in quantifying the total impact of biofouling on farm productivity. For example, based on impacts to yield and seed-stock supply, fouling within green-lipped mussel farms by *M. galloprovincialis* was estimated to cost the New Zealand industry US\$16.4 million per year, representing ~10% of the industry's value (Forrest and Atalah 2017). In the same region, the direct and combined economic impacts of two key biofoulers (the tunicate *Styela clava* and the polychaete *Sabella spallanzanii*) were estimated at NZ\$26.4 million over a 24-year period (the timeframe at which models predict total farm infestation would

occur [Soliman and Inglis 2018]). Similar estimates from around the world (eg Adams et al. 2011) mean that biofouling prevention and treatment are critical areas of research and that suitable AF methods are pertinent to efficient production within the shellfish industry.

The control of biofouling in shellfish aquaculture

The diverse impacts from biofouling drive persistent, resource intensive control efforts across the shellfish industry. Husbandry strategies to reduce the impact of biofouling involve preventing the development of fouling and treating fouled stock and infrastructure (Fitridge et al. 2012; Sievers et al. 2017) with the most effective biofouling management strategies incorporating aspects of both.

Prevention

Preventing harmful organisms from affecting aquaculture offers considerable advantages over treatments (Bui et al. 2019). A greater focus on avoiding biofouling will yield production benefits by reducing both the direct impacts of fouling organisms, and the frequency and intensity of treatments. In locations with predictable seasonal fouling patterns, spatial and temporal avoidance of biofouling may be feasible (Bullard et al. 2013; Sievers et al. 2014; Holthuis et al. 2015), and considerable recent effort has gone into monitoring biofouling development at farms around the world (eg Carraro et al. 2012; Antoniadou et al. 2013; Watts et al. 2015; Casso et al. 2018; das Chagas et al. 2018). Modelling biofouling settlement and development is also paving the way for sophisticated avoidance strategies. For example, computer modelling of mussel spat-fall in New Zealand, together with the creation of a readily accessible interactive application, has allowed farmers to actively avoid mussel fouling by informing when and where to deploy lines (Atalah, Rabel, et al. 2016; Atalah et al. 2017). In addition to using monitoring and modelling data to avoid culturing stock in specific locations during times of heavy fouling by particularly harmful species, these data can lead to more effective decisions on the timing of spat collection, and also inform when AF treatments and husbandry strategies that incidentally remove fouling, such as re-socking and grading, should occur (Cyr et al. 2007; Sievers et al. 2014).

Alternative preventative measures include strategic selection of rope types or culture methods, applying

AF shell coatings, and culturing fouling resistant genotypes. For example, Sievers et al. (2019) examined both spat collection and biofouling accumulation rates on seven different rope types, noting substantial and consistent variability amongst ropes, with clear advantages of selecting particular types in long-line mussel culture. Modifying stocking densities can also influence fouling rates (Dunham and Marshall 2012), and the surface wettability and microtopography of spat collectors has been altered to both enhance spat settlement and reduce fouling loads (Carl et al. 2012; Vucko et al. 2013). Finally, selective breeding of fouling resistant stock holds potential as a future preventative measure. In New Zealand, the fouling resistant properties of genetic variants of GreenshellTM mussels (*Perna calaniticulus*) are being investigated, with efforts extending beyond pilot-scale trials to large-scale intensive efforts (Camara and Symonds 2014).

Treatment

Despite the clear advantages of prevention, some biofouling will inevitably develop. On-site treatment methods which remove biofouling effectively, cheaply, easily and with minimal environmental impact are needed. Since Fitridge et al. (2012), several treatment methods have been tested experimentally and used commercially with varying levels of success, including manual removal (Li et al. 2018), exposure to air (Hillock and Costello 2013; Hopkins et al. 2016), freshwater (Fletcher et al. 2013), heat (Sievers et al. 2019), organic acids and bases (Rolheiser et al. 2012), pressure washing (Davidson et al. 2012), applying silicone release coatings (Tettelbach et al. 2014), adding a culture medium (a substratum within suspended bag culture that physically dislodges biofouling) (Dunham and Marshall 2012; Marshall and Dunham 2013), and employing biocontrol (Atalah et al. 2014; Sterling et al. 2016).

An interesting avenue of continued research which may prove useful for many cultured species involves combining multiple treatments. This approach has considerable appeal as (1) it may be more effective against a broader range of fouling species; (2) treatments will be effective using lower chemical concentrations or temperatures when applied simultaneously (safer for farmers and the environment); and (3) the effective exposure times are likely to be shorter. For example, recent evidence suggests that combining heat and acid treatments is more effective against numerous fouling species at lower intensities than

either in isolation (Sievers et al. 2019). Strategic timing of treatment application also offers considerable benefits over a random or intermittent approach. For instance, cost–benefit analysis of high pressure washing to remove the tunicate *Ciona intestinalis* found that beginning treatment when tunicates were small was most effective, and that three to four repeat treatments produced the greatest increase to overall farm profitability (Davidson et al. 2012).

In general, specific treatment methods are often tailored for the species cultured (eg oysters, mussels, scallops), the culture method (eg longline, trays, lanterns) and the composition of the fouling community. It is thus difficult to provide general guidelines for biofouling management within shellfish culture. Furthermore, although often effective at removing fouling, many current and previous treatment methods have little benefit for shellfish fitness or farm productivity, and at times can be more detrimental to cultured shellfish than biofouling (Sievers et al. 2017). For example, although the addition of culture media reduced biofouling by 70%, it also significantly reduced shellfish fitness due to the media limiting access to fresh seawater and food (Marshall and Dunham 2013). These issues have spurred considerable research into novel AF methods since Fitridge et al. (2012).

Novel antifouling control technologies

Natural compounds that inhibit larval metamorphosis may be useful antifoulants in shellfish aquaculture (Cahill et al. 2012; Cahill, Burritt, et al. 2013; Moodie et al. 2017). These products typically have a contact active mode of action, whereby they are effective while remaining bound within a stable matrix, so effects are limited to coated surfaces (Cahill and Kuhajek 2014). Although these compounds are suggested to have little environmental impact, be applicable to both farm infrastructure and shellfish, and reduce biofouling (Cahill, Heasman, et al. 2013), the present authors are not aware of any commercial-scale trials to test the effectiveness or feasibility of this method. Other natural compounds, such as extracts from shellfish periostracum, exhibit strong AF properties and can be designed for commercial use. For example, periostracum dichloromethane extracts containing oleamide reduce algal spore settlement (Kang et al. 2016), and crude periostracum extracts inhibit the attachment of barnacles, diatoms and marine bacteria (Bers et al. 2006). Shellfish periostraca and biogenically derived microtopographies present a non-toxic, environmentally friendly substratum to prevent

the settlement and attachment of a range of fouling organisms. However, many of the tested topographies that reduce biofouling settlement are species-specific and thus not particularly useful as a broad AF treatment (Bers and Wahl 2004; Magin et al. 2010; Scardino and de Nys 2011; Vucko et al. 2013). Furthermore, developing suitable methods to apply these types of compounds to shellfish is a major challenge. Coating shellfish with an AF formulation may be unviable from a logistical, fiscal, or consumer perspective. However, with additional research these products may become – at a minimum – a useful future tool for the shellfish aquaculture industry to protect infrastructure from biofouling.

An emerging strategy currently being tested to combat non-indigenous biofouling species *via* biosecurity incursion responses is encapsulation, whereby fouled structures are wrapped in material (eg PVC pallet wrapping), denying the organisms oxygen, nutrients and light (Roche et al. 2015; Atalah, Brook, et al. 2016). Toxic compounds also build up and contribute to high mortality rates of biofoulers (Coutts and Forrest 2007; Vaquer-Sunyer and Duarte 2010). Empirical evidence also suggests that combining encapsulation with chemical dosing using acetic acid may greatly reduce effective treatment times (Forrest et al. 2007; Denny 2008). Encapsulation was largely developed for use on boat hulls, pontoons and piles (see Atalah, Brook, et al. 2016 and references within), and is unlikely to be appropriate for use directly on shellfish (eg wrapping mussel lines). As with coatings, encapsulation may, however, be a viable option for removing biofouling from infrastructure such as mooring lines, buoys and trays.

The long-term genetic improvement of broodstock through selective breeding offers several advantages for the shellfish industry, including a potential means to combat biofouling (Camara and Symonds 2014; Hollenbeck and Johnston 2018). Selective breeding of Greenshell™ mussel and Pacific Oyster genetic variants have begun to improve farm productivity and overall profit yields, with indications that these variants accumulate hard-bodied biofoulers at lower rates (Camara and Symonds 2014). Ultimately, the practicality and efficacy of genetic improvements, and indeed all novel AF methods, need to be rigorously evaluated, with particular attention given to how such treatments affect stock fitness and farm productivity.

The impact of biofouling on finfish aquaculture

There are four main concerns regarding the growth of biofouling organisms on fish cages and

infrastructure in finfish aquaculture: (1) modified hydrodynamics in and around the cage affecting water quality and the cage's volume and stability; (2) increased disease risk due to biofoulers and associated pathogens; (3) behavioural impacts to cleaner fish used as biological control against sea lice; and (4) reservoirs for non-indigenous species.

Modified hydrodynamics

The occlusion of nets by biofouling restricts water exchange, which reduces oxygen levels, waste flushing and cage volumes, and increases the effective stocking densities, impacting fish health and welfare (reviewed in Fitridge et al. 2012). In addition, biofouling can increase the hydrodynamic load on nets up to 10-fold (Bi et al. 2018), deforming cages and adding strain to moorings. Impacts differ with species composition (Gansel et al. 2015, 2017) and organism size (Lader et al. 2015), and percentage net-aperture occlusion is a good predictor of flow reduction and drag increase associated with biofouling (Gansel et al. 2015; Bi and Xu 2018).

Biofouling and associated pathogens

Direct contact with cnidarian biofouling can be harmful to the fish, as organisms bearing nematocysts (stinging cells), such as the hydroid *Ectopleura larynx* and the anemone *Anthothoe albocincta*, have the potential to cause gill and skin damage (Baxter et al. 2012; Wybourne 2013; Bloecher, Powell, et al. 2018). In addition, biofouling poses a health risk to cultured fish as it can facilitate and amplify the presence of pathogens by harbouring viral, bacterial, and parasitic organisms that cause various diseases (reviewed in Fitridge et al. 2012). Resuspended faeces of the mussel *Mytilus edulis* containing the bacterium *Vibrio anguillarum* can infect cod with vibriosis in nearby cages and cause mortalities (Pietrak et al. 2012). This issue is also concerning for integrated multi trophic aquaculture (IMTA) sites where fish and shellfish are intentionally cultured in close vicinity. Vibriosis bacteria have also been found in biofilms on cage nets in Malaysia, where their abundance correlated with outbreaks of the disease (Albert and Ransangan 2013). The parasitic amoeba responsible for amoebic gill disease (AGD) in Atlantic salmon, *Paramoeba perurans*, is associated with several key biofouling organisms during acute AGD outbreaks, including hydroids, bryozoans, tunicates, and molluscs (Hellebø et al. 2017). Although the amoeba's reservoir between

outbreaks is still unknown, biofouling organisms could act as reinfection agents for recently treated or uninfected fish in nearby cages. Finally, other parasites, such as blood flukes (*Cardicola* spp.) that infect bluefin tuna in Japan, can be found growing on ropes, floats, and frames around cages where their intermediate hosts, terebellid polychaetes, live in balanid shells (Shirakashi and Hirano 2015; Sugihara et al. 2015).

Effects on cleaner fish

Cleaner fish, such as lumpfish (*Cyclopterus lumpus*) and ballan wrasse (*Labrus bergylta*), are increasingly used for biological control of sea lice in salmon farming (Powell et al. 2018). They are opportunistic, omnivorous feeders, targeting multiple food sources and switching their choice of prey organisms to what becomes available in their environment (Deady et al. 1995; Kvenseth 1996; Imsland et al. 2015; Eliassen et al. 2018). As their prey includes biofoulers, such as seaweeds, crustaceans, hydrozoans and mussels (Deady et al. 1995; Imsland et al. 2015; Eliassen et al. 2018), they can effectively reduce biofouling on cage nets (Kvenseth 1996). Consequently, to encourage lice feeding behaviour, farmers were recommended to keep nets free of biofouling (Deady et al. 1995; Kvenseth 1996; Powell et al. 2018). However, recent research has found the presence of biofouling to have a positive effect on the prevalence of sea lice in lumpfish stomachs, possibly because of a more active foraging behaviour of the lumpfish, or the provision of a sheltered environment resulting in better lumpfish welfare (Eliassen et al. 2018). Furthermore, cleaning cage nets to remove biofouling does not affect the behaviour (depth distribution, activity and habitat use) of lumpfish and ballan wrasse (Leclercq et al. 2018). Based on these findings, the salmon producer HiddenFjord on the Faroe Islands reduced the frequency of net cleaning to allow biofouling to accumulate for the benefit of cleaner fish, reporting positive results (E. Patursson, HiddenFjord, pers. comm.). In Norway and Scotland, however, regular net cleaning remains the norm.

Non-indigenous species reservoir

Fish farm biofouling communities can act as reservoirs for non-indigenous species (NIS) that can affect fish health, while farming practices can result in their range expansion, with potential ecological impacts (Table S2; Mineur et al. 2012; Simkanin et al. 2012).

For example, the movement of infrastructure into new areas increases the risk of spreading harmful algal blooms (HABs) when biofouling includes certain tunicates (eg *Didemnum vexillum* and *Molgula manhattensis*) (Rosa et al. 2013). HABs reduce fish welfare and can even cause mass mortality in caged fish with considerable economic consequences (Rensel and Whyte 2003; Cook et al. 2012). In some jurisdictions where the presence of NIS is of significant environmental concern, more stringent biofouling management and waste disposal practices are required (Rosa et al. 2013), with the presence of NIS resulting in stricter controls and potential removal of the site. However, from an operational farm perspective, biofouling by NIS is no more harmful to finfish culture than fouling by native species unless they have a faster growth rate, greater biomass or introduce additional health risks to cultured species than compared to their native counterparts, and thus the presence of NIS does not necessarily increase the need for additional biofouling management practices. To the best of the authors' knowledge, no studies exploring the economic implications of NIS vs native biofouling communities on fish aquaculture have been published.

The control of biofouling in finfish aquaculture

The management of biofouling in finfish aquaculture is an increasingly important issue for regulators as the impacts of different control methods become realised (Sim-Smith and Forsythe 2013; Floerl et al. 2016; Scianni et al. 2017). While the use of antifoulants is generally declining, the onset of other biofouling management techniques (eg *in situ* cleaning) with different environmental implications has spurred the development of aquaculture biofouling best management practices. This development has been largely industry-led and, in some jurisdictions, statutory aquaculture biofouling regulation has not yet been established. Table S3 summarises the current statutory and recommended best management practices for select countries.

Net cleaning and exchange

In situ net cleaning is one of the most common methods to manage biofouling on fish cage nets (Floerl et al. 2016). The technology, based on a rig equipped with rotating discs that expel high pressure water through nozzles, is moving towards remote operation

and autonomy by attaching rigs to remotely operated vehicles or equipping them with crawl belts (eg 'RONC' by MPI, 'NCL-LX' by Yanmar) or propulsion units (eg 'FNC8' by AKVAgrou, 'Manta' by Stranda Prolog, 'Stealth Cleaner' by Ocein). Alternative systems apply suction parallel to (and potentially independent of) high-pressure cleaning (eg 'MIC2.0' by PFG group), while others rely on cavitation-based systems (eg Cavitator Underwater Surface Cleaners).

Net cleaning or net exchange is often conducted *ad hoc*, determined by biofouling accumulation rates and requirements of the cultured species and/or statutory requirements. While this can be as seldom as once a year, in some cultures more frequent biofouling control is required. Further, in many salmon farms in Norway and Scotland concern over cleaner fish performance has led to considerable increases in cleaning frequency within the last decade (Guenther et al. 2010; Bloecher et al. 2015), now commonly fixed at fortnightly intervals (Bloecher et al. 2015). During the main biofouling season, intervals may even be reduced to as little as five days, especially for nets without antifouling coatings (SINTEF Ocean, unpublished data). For coated nets, increased net cleaning frequency leads to abrasion of the AF coating, considerably reducing the life-time of the coating (SINTEF Ocean, unpublished data) and contributing to the release of harmful AF products into the environment (Skarbøvik et al. 2017).

Other farm infrastructure such as mooring lines and chains, walkways and buoys are cleaned at much lower frequencies, occasionally only following the grow-out phase. Therefore, biofouling accumulation on these structures can be considerable (Bloecher et al. 2015) and may act as a reservoir for pathogens (Sugihara et al. 2014; Shirakashi and Hirano 2015), likely exacerbating biofouling recruitment to cage nets (Bloecher et al. 2015).

Net cleaning can also facilitate the spread of NIS by fracturing colonial species (Hopkins et al. 2011; Aldred and Clare 2014; Floerl et al. 2016) and triggering the simultaneous release of gametes, causing rapid recolonization (Carl et al. 2011; Floerl et al. 2016). Furthermore, the release of cleaning waste containing fragments of biofouling organisms and, potentially, particles of abraded copper coating, can severely impact fish health. Salmon farmers report agitated behaviour and reduced appetite during net cleaning and have observed gill and skin disorders afterwards. Upon contact, cnidarian biofoulers expel nematocysts that can penetrate fish skin and deliver poison (Helmholz et al. 2010; Cegolon et al. 2013), even after

fragmentation following pressure-washing (Bloecher, Floerl, et al. 2018). For example, the hydroid *Ectopleura larynx* causes gill injuries in Atlantic salmon (Baxter et al. 2012; Bloecher, Powell, et al. 2018), and white-striped anemones *Anthothoe albocincta* are suspected to cause skin damage in Chinook salmon (Wybourne 2013). Management methods for other issues (eg parasitic sea lice) can further exacerbate these impacts. For example, lice skirts (Stien et al. 2018) can trap cleaning waste within cages, leading to greater interaction and risk of stinging by nematocysts.

Finally, *in situ* cleaning without waste retention can release large volumes of organic material into the surrounding environment, exacerbating existing issues around the build of organic matter below salmon farms (Sim-Smith and Forsythe 2013; Floerl et al. 2016). For this reason, *in situ* cleaning has been discontinued in New Brunswick and Nova Scotia in areas where the benthic environment is anoxic or hypoxic. While *in situ* cleaning is still conducted in New Zealand, Sim-Smith and Forsythe (2013) recommend that it should not be used at sites with low flow due to concerns for the benthic environment (see Table S1).

Net materials and treatments

While salmon farmers in Australia and New Zealand have recently abandoned biocidal antifoulants due to environmental concerns (Floerl et al. 2016, Table S3), their use is common elsewhere. Copper continues to be the main active biocide (Guardiola et al. 2012; Makridis et al. 2018), with Norway's salmon industry using ~1,250 t of copper for AF coatings annually (Skarbøvik et al. 2017). Copper coatings protect nylon nets from fouling for up to seven months at sea (Edwards et al. 2015) and can result in better growth and feed conversion rates in seabass compared to those in unprotected nylon nets (Yigit et al. 2018). However, if biofouling pressure is high, coatings may fail after only eight weeks at sea (Bloecher, Floerl, et al. 2018). Furthermore, ~85% of the copper on Norwegian farms is released into the sea because of leaching and abrasive net cleaning (Skarbøvik et al. 2017). While copper accumulation in marketable tissue of cultured fish does not exceed food safety standards (Cotou et al. 2012; Nikolaou et al. 2014; Kalantzi et al. 2016), impacts on the fish (Azizishirazi et al. 2015) and non-target organisms (Fitridge et al. 2012; Guardiola et al. 2012) have been reported.

Alternative biocides termed 'booster biocides' can be added to enhance the efficacy of copper coatings

(Guardiola et al. 2012; Amara et al. 2018), or can be used as the main active ingredient, offering alternatives to traditional coatings based on cuprous oxide. The most commonly used 'booster biocide' compounds in net coatings include copper pyrithione, zinc pyrithione, and tralopyril ('Econea') (Bloecher, Floerl, et al. 2018). In a comparative study, only copper pyrithione-based coatings performed similar to conventional copper coating (Bloecher, Floerl, et al. 2018). While booster biocides are often marketed as environmentally friendly alternatives to conventional copper coatings, there are still environmental concerns regarding their toxicity. Copper pyrithione, for example, reduced salmonid gill health in laboratory assays (Borg and Trombetta 2010) and impacted non-target organisms (Bao et al. 2011; Oliveira et al. 2014; Oliveira et al. 2016; Amara et al. 2018).

Another alternative to traditional copper coatings is copper alloy metal (CAM) nets. CAM nets are able to prevent most biofouling (Chambers et al. 2012) and offer higher form stability in strong currents due to reduced drag (Tsukrov et al. 2011). Growth and feed conversion rates in CAM nets are equal to or better than in nylon net pens treated with conventional copper coating (Chambers et al. 2012; Yigit et al. 2018). In comparison to conventional copper coatings, leaching rates are higher during the first six months at sea before they reach similar levels (Kalantzi et al. 2016). It is assumed, however, that the reduced frequency of cleaning required for CAM nets will result in an overall lower leaching rate for a growing season (Kalantzi et al. 2016).

Most non-biocidal alternatives seal net surfaces under wax- or resin-based coatings. The aim is to reduce surface structure and texture to create a less favourable settlement surface in addition to increasing resilience to net cleaning (Swain and Shinjo 2014; Edwards et al. 2015; Baum et al. 2017). Alternative net materials such as HDPE ('Dyneema') or PET monofilaments ('Kikko net') are also used with similar intention. While this technology can delay the onset and reduce the accumulation of biofouling compared to regular, uncoated raschel-knitted nylon nets (Edwards et al. 2015; Baum et al. 2017), they do not outperform conventional copper coated nets (Edwards et al. 2015; Bloecher, Floerl, et al. 2018).

Biological control

While there are a range of invertebrate and fish species that feed on specific biofouling organisms and can theoretically be co-cultured in cages or on nets,

not all biofoulers have natural predators suitable as biofouling control agents (reviewed in Fitridge et al. 2012; Madin and Ching 2015). Since the publication of Fitridge et al. (2012), no significant advances in the use of biological controls have been made and therefore their use in finfish aquaculture remains in the experimental stage.

Novel biofouling control technologies

Given the current reliance on copper for biofouling control, efforts have been put into developing methods to minimise the release of copper into the environment whilst maintaining the use of its AF properties (Liu et al. 2017). The combination of polymer coatings with embedded copper biocide has shown potential in deterring biofouling while limiting the release of copper on hard surfaces (Vucko, King, et al. 2014) and netting (Sato et al. 2012; Ashraf and Edwin 2016; Ashraf et al. 2017). Although many studies have explored the use of natural compounds (eg from plants, bacteria, fungi, algae and sponges) as antifoulants with varying success (reviewed in Almeida and Vasconcelos 2015), to date none are available for commercial application in finfish aquaculture.

As a biocide-free approach to biofouling prevention, manipulation of surface texture and wettability to create a less favourable settlement and adhesion surface has been explored (Scardino and de Nys 2011; Bloecher et al. 2013; Nir and Reches 2016). However, results highlight the limited use of this approach in finfish farming as different species are impacted by different textures (Vucko, Poole, et al. 2014) and deterrence of key pest species, such as hydroids, is ineffective (Bloecher et al. 2013).

Silicone-based fouling-release (FR) technologies have been trialled to develop environmentally benign coatings that reduce adhesion and improve the ability to clean nets (see references in Nurioglu et al. 2015, Pradhan et al. 2018, and Gevaux et al. 2019 for examples). Although these coatings provide little preventative benefit when applied to fish cage nets, silicon-based FR coatings offer the potential for improved biofouling control in conjunction with mechanical cleaning methods (Swain and Shinjo 2014; Edwards et al. 2015).

With regard to mechanical cleaning, recent trends have shifted towards preventing the development of mature biofouling communities by frequent disturbance using autonomous brush systems which continuously clean the net (eg 'HALO Net Maintenance

System' by AquaRobotics, 'Netrobot' by Mørenot). However, as these systems are just entering the market, their efficacy has not yet been assessed independently.

While ambitions to develop environmentally benign AF materials and cleaning strategies remain, further developments are needed before existing alternatives compete with the biofouling management methods currently employed by the finfish aquaculture industry.

The impacts of biofouling in seaweed aquaculture

Biofouling exerts a range of negative impacts on the commercial production of seaweeds (Table S4). These can be categorised into three groups: (1) competition for light, space and dissolved nutrients; (2) physical damage; and (3) interference with seaweed culture infrastructure. The culmination of these can reduce the productivity of seaweed farms.

Competition for light, space and dissolved nutrients

Fouling species compete with cultured seaweed species for light, space and dissolved nutrients. Studies on the cultured red algae *Gracilaria chilensis* (Buschmann and Gómez 1993) and *Kappaphycus alvarezii* (Marroig and Reis 2016) show that biofouling significantly reduces levels of solar irradiance reaching cultured stock, leading to lower photosynthetic rates and photosynthetic efficiency than unfouled stock (Borlongan et al. 2016). Interestingly, higher chlorophyll-*a* and phycobilin content in heavily fouled stock suggests that some seaweeds can acclimatise to low-irradiance conditions (Borlongan et al. 2016). Furthermore, opportunistic benthic algae growing on cultured seaweed and farm infrastructure, such as cultivation ropes, raceways and rafts, directly compete with cultured seaweed for substratum, space, and dissolved nutrients, such as ammonia, nitrogen and inorganic carbon (Buschmann and Gómez 1993; Fletcher 1995; Veeragurunathan et al. 2015).

Physical damage

Biofouling adds considerable weight to cultured seaweeds, making them prone to breakage and dislodgement. Some fouling species, such as the encrusting bryozoans *Membranipora membranacea* and *Electra pilosa*, make the lamina of cultivated kelps brittle, which increases susceptibility to breakage (Førde et al.

2016). In addition, the presence of tunicates, predominantly *Ciona intestinalis*, and the hydroid *Obelia geniculata* can deteriorate the appearance and quality of cultured kelp blades (Park and Hwang 2012; Rolin et al. 2017). Furthermore, some epiphytic fouling species penetrate deep into the host cell tissues, causing disorganisation or even destruction of the host's cells close to the infection (Leonardi et al. 2006). Epiphytic red algae from the genus *Neosiphonia* (syn. *Polysiphonia*) can cause pit-like structures to form on cultured seaweeds, leading to future infections (Hurtado et al. 2006; Vairappan et al. 2008).

Interference with seaweed culture infrastructure

Fouling organisms on ropes and rafts used to culture seaweeds can cause infrastructure to sink below the surface, requiring labour-intensive cleaning and leading to a loss of productivity (Marroig and Reis 2011, 2016). Biofouling on infrastructure can also cause environmental effects beyond geographic farm boundaries. For example, the recurring large-scale green tides in the Yellow Sea of China originated from the green alga *Ulva prolifera* that was scraped off rafts at *Pyropia* (syn. *Porphyra*) *yezoensis* farms in the Yellow Sea (Fan et al. 2015; Song et al. 2018). The decomposition of green tides can result in hypoxia and acidification, induce red tides and have long lasting effects on the coastal carbon cycle and ecosystem health (reviewed in Zhang et al. 2019).

Reduction in productivity, quality and commercial value

Similar to shellfish and finfish culture, the culmination of these negative impacts can lead to a loss in productivity of seaweed culture by decreasing growth rates and biomass and reducing the quality and commercial value of the end product (Kuschel and Buschmann 1991; Park and Hwang 2012; Bruhn et al. 2016; Førde et al. 2016; Marroig and Reis 2016). Although the impact of biofouling is consistently cited as a leading issue in seaweed aquaculture (Lüning and Pang 2003; Kim et al. 2017), the authors are unaware of any attempt to quantify the financial implications of productivity reductions or increases to time and labour costs.

The control of biofouling in seaweed aquaculture

The economic feasibility of seaweed aquaculture requires the control of various factors, including

biofouling (Zuniga-Jara and Marin-Riffo 2016; Kim et al. 2017). Prevention, inhibition and treatment of biofouling on commercially cultivated seaweed species includes understanding and harnessing the natural antifouling defences of seaweeds, and strategic farm management and husbandry practices.

Prevention and inhibition

Strategic farm management and husbandry strategies can minimise the presence and impact of biofouling and optimise biomass yields. Like shellfish and finfish culture, farmers can choose to culture seaweed in areas less likely to develop harmful biofouling by considering levels of exposure and water movement, water temperature, cultivation period, timing of harvest, and through the choice of infrastructure materials, which all influence biofouling rates. Farming seaweed at exposed locations is one promising strategy, but it is highly species-specific. For example, exposed sites are linked with reduced levels of biofouling for the cultured kelps *Undaria pinnatifida*, *Saccharina latissima* and *Laminaria digitata* (Andersen et al. 2011; Peteiro and Freire 2013; Rolin et al. 2017) and increased water movement with less siltation is recommended for the cultivated red alga *Kappaphycus alvarezii* (Hurtado et al. 2006). In contrast, softer-bodied seaweeds such as the red alga *Gracilaria chilensis* may lose biomass when cultured at exposed locations (reviewed in Buck et al. 2018). However, cultivating seaweeds at exposed sites may also present other environmental challenges as severe storms can damage seaweeds and displace aquaculture structures, leading to reductions in biomass and farm productivity (Rolin et al. 2017).

Water temperature also influences the accumulation of fouling species. In general, it has been recommended that cultured seaweeds are harvested before sea temperatures rise in spring and summer to avoid the negative impacts of seasonal biofouling (Park and Hwang 2012; Ateweberhan et al. 2015; Marinho et al. 2015; Førde et al. 2016; Keesing et al. 2016).

In terms of infrastructure material, the efficacy of microtextured surfaces in deterring the settlement of specific fouling organisms has been increasingly studied, with mixed findings (reviewed in Scardino and de Nys 2011). For example, polyethylene tubes used in the culture of *Gracilaria* promoted the unwanted accumulation of *Giffordia* (Kuschel and Buschmann 1991), but treated bamboo poles used in the culture of *Pyropia yezoensis* inhibited the attachment and germination of green alga *Ulva prolifera*

micro-propagules (Song et al. 2018). The use of such bamboo poles might therefore play a role in mitigating recurring large-scale green tides in the Yellow Sea caused by *U. prolifera* (Song et al. 2018).

Biological aspects of farm management and husbandry strategies to prevent and inhibit biofouling include the selection of seed stock and stocking density. The selection of clean and healthy seedlings that are free of epiphytes to initiate cultivation is strongly recommended to curb biofouling problems later in the cultivation cycle (Hurtado et al. 2006; Hayashi et al. 2010). Epiphytic biofouling can be controlled by growing seaweeds at high densities in rope cultures in the sea or in tank cultures on land (reviewed in Lüning and Pang 2003), although the efficacy of this method appears species-specific, with the culture density for some species such as *Gracilaria* sp. and the kelp *Alaria esculenta* not affecting the abundance, species richness or composition of fouling species (Kuschel and Buschmann 1991; Walls et al. 2017).

In addition, the role of surface-associated secondary metabolites as a chemical AF defence mechanism has been demonstrated for several seaweed species (reviewed in Jormalainen and Honkanen 2008; da Gama et al. 2014; Othmani et al. 2016; Pereira et al. 2017), including the cultivated red alga *Gracilaria vermiculophylla* (Wang et al. 2018) and the brown alga *Fucus vesiculosus* (Lachnit et al. 2013). These mechanisms could be enhanced by farmers by using Acadian Marine Plant Extract Powder (AMPEP), a commercial product derived from the brown alga *Ascophyllum nodosum* and used extensively to increase the productivity of agricultural and horticultural crops for over 30 years (Hurtado and Critchley 2013). More recently, AMPEP has been tested and used as a culture medium to propagate and cultivate the red seaweed *Kappaphycus alvarezii* (Hurtado et al. 2009; Hurtado et al. 2012; Hurtado and Critchley 2013; Marroig and Reis 2016) and mitigate various biotic and abiotic stressors including biofouling (reviewed in Hurtado and Critchley 2018). For example, dipping *K. alvarezii* seedlings in AMPEP solutions helps prevent the occurrence of epiphytic infections and epibiont biomass during their culture period (Borlongan et al. 2011; Marroig and Reis 2016). The use of AMPEP and alternative extracts to prevent and inhibit epiphytes is a promising method, which may lead to further studies on the mode of action and metabolic pathways (Hurtado and Critchley 2018) and to the broader adoption of this method in seaweed aquaculture.

Treatment and mitigation

Seaweeds, particularly long-lived macroalgae species, have numerous natural defences that effectively remove biofouling, including blade abandonment and simultaneous rapid proliferation (Littler and Littler 1999), and periodic epidermal shedding (Filion-Myklebust and Norton 1981; Nylund and Pavia 2005; Yamamoto et al. 2013; Halat et al. 2015). For example, the commercially harvested brown alga *Ascophyllum nodosum* from the intertidal zone of Nova Scotia, Canada, sheds ~25% of its frond epidermis per week for at least nine months of the year to remove epiphytes, such as the host-specific red alga *Vertebrate lanosa* (syn. *Polysiphonia lanosa*) and the facultative brown algae *Elachista fucicola* and *Pylaiella littoralis* (Longtin et al. 2009; Halat et al. 2015). Combining knowledge of these natural defence mechanisms together with knowledge of the seasonal occurrence of biofouling could provide seaweed farmers with an increased capacity to limit biomass losses by harvesting at strategic times.

Strategic farm management and husbandry practices to control epiphytes include the exposure of desiccation-tolerant intertidal seaweed species, such as *Porphyra*, to air or the application of organic acids onto the cultivation nets to control the pH (Harrison and Hurd 2001; Kim et al. 2017). For seaweed species cultured in tanks, such as *Gracilaria*, epiphytes can also be controlled by nutrient pulsing, thereby starving the epiphytes of nitrogen between pulses, while not significantly affecting the growth of *Gracilaria* (reviewed in Harrison and Hurd 2001).

Biological control methods include encouraging grazing by herbivores, such as amphipods, isopods, and some fish species. The isopod *Paridotea reticulata* grazes on the epiphytic red alga *Ceramium diaphanum* growing on commercially farmed red alga *Gracilaria gracilis* (Anderson et al. 1998; Smit et al. 2003). However, the relationships between cultured seaweed, epiphytes and herbivores are dynamic, and the benefits of herbivores preferentially consuming epiphytes can be quickly overshadowed by negative effects of herbivores grazing on cultured seaweed when herbivore densities become too high and alternative food sources scarce (Shacklock and Doyle 1983; Smit et al. 2003; Cruz-Rivera and Friedlander 2011). In *Gracilaria* culture, isopod densities can be controlled by short-term freshwater exposure (Smit et al. 2003). Furthermore, in a food preference study with 11 fish species, the fish *Aphanius dispar* and *Tilapia zillii*, acclimatised to seawater, displayed selective feeding on epiphytes and were identified as

good candidates to control epiphytes in the mass culture of *Gracilaria conferta* (Friedlander et al. 1996). Future efforts should focus on harnessing the utility of biological control agents within culture settings.

Finally, since biofouling on seaweed culture infrastructure can be a reservoir for epibionts, periodic manual removal of fouling organisms on infrastructure and/or cultured species occurs in some farming regions (Kuschel and Buschmann 1991; Hurtado et al. 2006; Marroig and Reis 2011; Liu et al. 2013). However, manual removal is labour-intensive, and the gains in productivity must be evaluated against the increases in labour costs (Kuschel and Buschmann 1991). Furthermore, for several cultured seaweeds, the logistics of removing epiphytes make this method almost impossible. For example, epiphytic filamentous red algae from the genus *Polysiphonia* penetrate the cells of commercially cultured *Kappaphycus* (Hurtado et al. 2006; Leonardi et al. 2006), making manual removal difficult without harming the stock.

Although seaweed aquaculture technologies have developed significantly over the last decades, simple, economically viable biofouling management solutions have not yet been realised. Further research in this area is required (Kim et al. 2017), including trials of promising methods at commercial scales to facilitate future implementation by the industry.

Conclusion and future directions

Although there have been considerable recent advances in knowledge of biofouling in aquaculture, there are still significant gaps in aquaculturalists' capacity to manage biofouling. Issues surrounding biofouling differ among shellfish, finfish and seaweed aquaculture industries, yet the direction of emerging biofouling management strategies are similar. These strategies aim to take advantage of improved scientific understanding and technological advances to develop robust and proactive preventative approaches. The use of monitoring and modelling to inform preventative management decisions (eg farm locations, timing of treatment, harvesting cycles) can reduce the cost of biofouling management and maximise the benefit of any subsequent control treatments. The most effective strategies are likely to include a combination of both prevention and treatment. Likewise, though successes have been identified at varying scales by applying methods in isolation, the combination of multiple treatments may enable more success in combating biofouling in aquaculture.

Future research will need to address the unresolved issues in existing aquaculture practice, in addition to broadening the scope to encompass a wider variety of culture species and habitats as the aquaculture industry moves into new culture scenarios (eg offshore aquaculture and closed culture systems). By understanding the settlement and development of biofouling in these environments, novel materials and/or AF coatings for infrastructure, and technological advances, such as equipment for more efficient monitoring, modelling and autonomous cleaning can be developed. Aquaculture may help feed the world into the future, but biofouling stands out as a clear barrier that needs to be overcome first.

Acknowledgements

The authors would like to thank Isla Fitridge for her comments on the manuscript.

Disclosure statement

The authors declare no conflict of interest, financial interest or benefit from the direct applications of this research. Any use of trade, firm and product names is for descriptive purposes only and does not imply endorsement.

Funding

This work was supported by internal funds from the Institute of Marine and Antarctic Studies, The Global Wetlands Project [private charitable trust], and the Research Council of Norway [postdoctoral grant number 244444/E40].

ORCID

Jana Bannister  <http://orcid.org/0000-0002-5281-7800>

Michael Sievers  <http://orcid.org/0000-0001-7162-1830>

Flora Bush  <http://orcid.org/0000-0003-0596-0601>

Nina Bloecher  <http://orcid.org/0000-0002-8009-331X>

References

- Adams CM, Shumway SE, Whitlatch RB, Getchis T. 2011. Biofouling in marine molluscan shellfish aquaculture: a survey assessing the business and economic implications of mitigation. *J World Aquacult Soc.* 42:242–252. doi:10.1111/j.1749-7345.2011.00460.x
- Albert V, Ransangan J. 2013. Effect of water temperature on susceptibility of culture marine fish species to vibriosis. *Int J Res Pure Appl Microbiol.* 3:48–52.
- Aldred N, Clare AS. 2014. Mini-review: Impact and dynamics of surface fouling by solitary and compound ascidians. *Biofouling.* 30:259–270. doi:10.1080/08927014.2013.866653

- Almeida JR, Vasconcelos V. 2015. Natural antifouling compounds: effectiveness in preventing invertebrate settlement and adhesion. *Biotechnol Adv.* 33:343–357. doi:10.1016/j.biotechadv.2015.01.013
- Amara I, Miled W, Slama RB, Ladhari N. 2018. Antifouling processes and toxicity effects of antifouling paints on marine environment. A review. *Environ Toxicol Pharmacol.* 57:115–130. doi:10.1016/j.etap.2017.12.001
- Andersen GS, Steen H, Christie H, Fredriksen S, Moy FE. 2011. Seasonal patterns of sporophyte growth, fertility, fouling, and mortality of *Saccharina latissima* in Skagerrak, Norway: implications for forest recovery. *J Mar Biol.* 2011:1–8. doi:10.1155/2011/690375
- Anderson BC, Smit AJ, Bolton JJ. 1998. Differential grazing effects by isopods on *Gracilaria gracilis* and epiphytic *Ceramium diaphanum* in suspended raft culture. *Aquaculture.* 169:99–109. doi:10.1016/S0044-8486(98)00322-6
- Antoniadou C, Voultziadou E, Rayann A, Chintiroglou C. 2013. Sessile biota fouling farmed mussels: diversity, spatio-temporal patterns, and implications for the basibiont. *J Mar Biol Ass.* 93:1593–1607. doi:10.1017/S0025315412001932
- Ashraf PM, Edwin L. 2016. Nano copper oxide incorporated polyethylene glycol hydrogel: an efficient antifouling coating for cage fishing net. *Int Biodeter Biodegr.* 115:39–48. doi:10.1016/j.ibiod.2016.07.015
- Ashraf PM, Sasikala KG, Thomas SN, Edwin L. 2017. Biofouling resistant polyethylene cage aquaculture nettings: a new approach using polyaniline and nano copper oxide. *Arab J Chem.* <http://dx.doi.org/10.1016/j.arabjc.2017.1008.1006>.
- Atalah J, Brook R, Cahill PL, Fletcher LM, Hopkins GA. 2016. It's a wrap: encapsulation as a management tool for marine biofouling. *Biofouling.* 32:277–286. doi:10.1080/08927014.2015.1137288
- Atalah J, Newcombe EM, Hopkins GA, Forrest BM. 2014. Potential biocontrol agents for biofouling on artificial structures. *Biofouling.* 30:999–1010. doi:10.1080/08927014.2014.956734
- Atalah J, Rabel H, Forrest B. 2016. Blue mussel over-settlement predictive model and web application. Nelson (New Zealand): Cawthron Institute.
- Atalah J, Rabel H, Forrest BM. 2017. Modelling long-term recruitment patterns of blue mussels *Mytilus galloprovincialis*: a biofouling pest of green-lipped mussel aquaculture in New Zealand. *Aquacult Environ Interact.* 9:103–114. doi:10.3354/aei00216
- Atalah J, Smith KF. 2015. Assessment of stinging cells and identity of anemones at Ruakaka Bay salmon farm. Nelson (New Zealand): Cawthron Institute.
- Ateweberhan M, Rougier A, Rakotomahazo C. 2015. Influence of environmental factors and farming technique on growth and health of farmed *Kappaphycus alvarezii* (cottonii) in south-west Madagascar. *J Appl Phycol.* 27:923–934. doi:10.1007/s10811-014-0378-3
- Azizishirazi A, Dew WA, Bougas B, Bernatchez L, Pyle GG. 2015. Dietary sodium protects fish against copper-induced olfactory impairment. *Aquat Toxicol.* 161:1–9. doi:10.1016/j.aquatox.2015.01.017
- Bao VW, Leung KM, Qiu JW, Lam MH. 2011. Acute toxicities of five commonly used antifouling booster biocides to selected subtropical and cosmopolitan marine species. *Mar Pollut Bull.* 62:1147–1151. doi:10.1016/j.marpolbul.2011.02.041
- Baum C, Rohde P, Friedrichs L, Carl S, Kalmer L, Bendig T. 2017. Simulated hydrodynamic qualities of a virtually coated Raschel net fragment related to biofouling. *Aquac Res.* 48:5159–5165. doi:10.1111/are.13214
- Baxter E, Sturt M, Ruane N, Doyle T, McAllen R, Rodger H. 2012. Biofouling of the hydroid *Ectopleura larynx* on aquaculture nets in Ireland: implications for finfish health. *Fish Vet J.* 13:18–30.
- Bers AV, D'Souza F, Klijnstra JW, Willemsen PR, Wahl M. 2006. Chemical defence in mussels: antifouling effect of crude extracts of the periostracum of the blue mussel *Mytilus edulis*. *Biofouling.* 22:251–259. doi:10.1080/08927010600901112
- Bers AV, Wahl M. 2004. The influence of natural surface microtopographies on fouling. *Biofouling.* 20:43–51. doi:10.1080/08927010410001655533
- Bi C-W, Xu T-J. 2018. Numerical study on the flow field around a fish farm in tidal current. *Turk J Fish Aquat Sci.* 18.
- Bi C-W, Zhao Y-P, Dong G-H, Wu Z-M, Zhang Y, Xu T-J. 2018. Drag on and flow through the hydroid-fouled nets in currents. *Ocean Eng.* 161:195–204. doi:10.1016/j.oceaneng.2018.05.005
- Bloecher N, de Nys R, Poole AJ, Guenther J. 2013. The fouling hydroid *Ectopleura larynx*: a lack of effect of next generation antifouling technologies. *Biofouling.* 29:237–246. doi:10.1080/08927014.2013.763228
- Bloecher N, Floerl O, Frank K, Fore M. 2018. Evaluating the efficacy of alternative methods for biofouling control in aquaculture. Trondheim (Norway): SINTEF Ocean; Report No. 2018:00733.
- Bloecher N, Floerl O, Sunde LM. 2015. Amplified recruitment pressure of biofouling organisms in commercial salmon farms: potential causes and implications for farm management. *Biofouling.* 31:163–172. doi:10.1080/08927014.2015.1012713
- Bloecher N, Powell M, Hytterod S, Gjessing M, Wiik-Nielsen J, Mohammad SN, Johansen J, Hansen H, Floerl O, Gjevne AG. 2018. Effects of cnidarian biofouling on salmon gill health and development of amoebic gill disease. *PLoS One.* 13:e0199842. doi:10.1371/journal.pone.0199842
- Borg DA, Trombetta LD. 2010. Toxicity and bioaccumulation of the booster biocide copper pyrithione, copper 2-pyridinethiol-1-oxide, in gill tissues of *Salvelinus fontinalis* (brook trout). *Toxicol Ind Health.* 26:139–150. doi:10.1177/0748233710362381
- Borlongan IAG, Luhan MRJ, Padilla PIP, Hurtado AQ. 2016. Photosynthetic responses of '*Neosiphonia* sp. epiphyte-infected' and healthy *Kappaphycus alvarezii* (Rhodophyta) to irradiance, salinity and pH variations. *J Appl Phycol.* 28:2891–2902. doi:10.1007/s10811-016-0833-4
- Borlongan IAG, Tibubos KR, Yunque DAT, Hurtado AQ, Critchley AT. 2011. Impact of AMPEP on the growth and occurrence of epiphytic *Neosiphonia* infestation on two varieties of commercially cultivated *Kappaphycus alvarezii* grown at different depths in the Philippines. *J Appl Phycol.* 23:615–621. doi:10.1007/s10811-010-9649-9

- Bruhn A, Tørring DB, Thomsen M, Canal-Vergés P, Nielsen MM, Rasmussen MB, Eybye KL, Larsen MM, Balsby TJS, Petersen JK. 2016. Impact of environmental conditions on biomass yield, quality, and bio-mitigation capacity of *Saccharina latissima*. *Aquacult Environ Interact*. 8:619–636. doi:10.3354/aei00200
- Buck BH, Troell MF, Krause G, Angel DL, Grote B, Chopin T. 2018. State of the art and challenges for off-shore Integrated Multi-Trophic Aquaculture (IMTA). *Front Mar Sci*. 5:165. doi:10.3389/fmars.2018.00165
- Bui S, Oppedal F, Sievers M, Dempster T. 2019. Behaviour in the toolbox to outsmart parasites and improve fish welfare in aquaculture. *Rev Aquacult*. 11:168–186. doi:10.1111/raq.12232
- Bullard SG, Davis CV, Shumway SE. 2013. Seasonal patterns of ascidian settlement at an aquaculture facility in the Damariscotta River, Maine. *J Shellfish Res*. 32: 255–264. doi:10.2983/035.032.0202
- Buschmann AH, Gómez P. 1993. Interaction mechanisms between *Gracilaria chilensis* (Rhodophyta) and epiphytes. *Hydrobiologia*. 260-261:345–351. doi:10.1007/BF00049039
- Cahill PL, Burritt D, Heasman K, Jeffs AG, Kuhajek J. 2013. Screening for antioxidant and detoxification responses in *Perna canaliculus* Gmelin exposed to an antifouling bioactive intended for use in aquaculture. *Chemosphere*. 93:931–938. doi:10.1016/j.chemosphere.2013.05.058
- Cahill PL, Heasman K, Jeffs A, Kuhajek J, Mountfort D. 2012. Preventing ascidian fouling in aquaculture: screening selected allelochemicals for anti-metamorphic properties in ascidian larvae. *Biofouling*. 28:39–49. doi:10.1080/08927014.2011.648624
- Cahill PL, Heasman K, Jeffs AG, Kuhajek J. 2013. Laboratory assessment of the antifouling potential of a soluble-matrix paint laced with the natural compound polygodial. *Biofouling*. 29:967–975. doi:10.1080/08927014.2013.822488
- Cahill PL, Kuhajek JM. 2014. Polygodial: a contact active antifouling biocide. *Biofouling*. 30:1035–1043. doi:10.1080/08927014.2014.966305
- Camara M, Symonds J. 2014. Genetic improvement of New Zealand aquaculture species: programmes, progress and prospects. *New Zeal J Mar Freshw*. 48:466–491. doi:10.1080/00288330.2014.932291
- Campbell DA, Kelly MS. 2002. Settlement of *Pomatoceros triqueter* (L.) in two Scottish Lochs, and factors determining its abundance on mussels grown in suspended culture. *J Shellfish Res*. 21:519–527.
- Carl C, Guenther J, Sunde LM. 2011. Larval release and attachment modes of the hydroid *Ectopleura larynx* on aquaculture nets in Norway. *Aquac Res*. 42:1056–1060. doi:10.1111/j.1365-2109.2010.02659.x
- Carl C, Poole AJ, Sexton BA, Glenn FL, Vucko MJ, Williams MR, Whalan S, de Nys R. 2012. Enhancing the settlement and attachment strength of pediveligers of *Mytilus galloprovincialis* by changing surface wettability and microtopography. *Biofouling*. 28:175–186. doi:10.1080/08927014.2012.662676
- Carraro JL, Rupp GS, Mothes B, Lerner C, Wuerdig NL. 2012. Characterization of the fouling community of macroinvertebrates on the scallop *Nodipecten nodosus* (Mollusca, Pectinidae) farmed in Santa Catarina, Brazil. *Cienc Mar*. 38:577–588. doi:10.7773/cm.v38i3.1982
- Carroll JM, O'Shaughnessy KA, Diedrich GA, Finelli CM. 2015. Are oysters being bored to death? Influence of *Cliona celata* on *Crassostrea virginica* condition, growth and survival. *Dis Aquat Org*. 117:31–44. doi:10.3354/dao02928
- Casso M, Navarro M, Ordóñez V, Fernández-Tejedor M, Pascual M, Turon X. 2018. Seasonal patterns of settlement and growth of introduced and native ascidians in bivalve cultures in the Ebro Delta (NE Iberian Peninsula). *Reg Stud Mar Sci*. 23:12–22. doi:10.1016/j.rsm.2017.11.002
- Cegolon L, Heymann WC, Lange JH, Mastrangelo G. 2013. Jellyfish stings and their management: a review. *Mar Drugs*. 11:523–550. doi:10.3390/md11020523
- Chambers M, Bunker J, Watson W, Howell W. 2012. Comparative growth and survival of juvenile Atlantic cod (*Gadus morhua*) cultured in copper and nylon net pens. *J Aquacult Res Dev*. 03:137.
- Cook EJ, Davidson K, Fox C, Black K. 2012. Monitoring and eradication of invasive and non-native species in aquaculture units. Oban (Scotland): Scottish Association for Marine Science; Report No. SARF087.
- Cotou E, Henry M, Zeri C, Rigos G, Torrealba A, Catsiki VA. 2012. Short-term exposure of the European sea bass *Dicentrarchus labrax* to copper-based antifouling treated nets: copper bioavailability and biomarkers responses. *Chemosphere*. 89:1091–1097. doi:10.1016/j.chemosphere.2012.05.075
- Coutts AD, Forrest BM. 2007. Development and application of tools for incursion response: lessons learned from the management of the fouling pest *Didemnum vexillum*. *J Exp Mar Biol Ecol*. 342:154–162. doi:10.1016/j.jembe.2006.10.042
- Cruz-Rivera E, Friedlander M. 2011. Feeding preferences of mesograzers on aquacultured *Gracilaria* and sympatric algae. *Aquaculture*. 322:218–222.
- Cyr C, Myrand B, Cliche G, Desrosiers G. 2007. Weekly spat collection of sea scallop, *Placopecten magellanicus*, and undesirable species as a potential tool to predict an optimal deployment period of collectors. *J Shellfish Res*. 26:1045–1055.2.0.CO;2] doi:10.2983/0730-8000(2007)26 [1045:WSCOSS]2.0.CO;2
- da Gama BAP, Plouguerne E, Pereira RC. 2014. The antifouling defence mechanisms of marine macroalgae. In: Bourgougnon N, editor. *Advances in botanical research: Sea Plants*. Vol. 71. Academic Press, Elsevier Ltd; p. 413–440.
- Daigle RM, Herbinger CM. 2009. Ecological interactions between the vase tunicate (*Ciona intestinalis*) and the farmed blue mussel (*Mytilus edulis*) in Nova Scotia, Canada. *Aquat Invasions*. 4:177–187. doi:10.3391/ai.2009.4.1.18
- das Chagas RA, Barros MRF, dos Santos WCR, Herrmann M. 2018. Composition of the biofouling community associated with oyster culture in an Amazon estuary, Para State, North Brazil. *Rev Biol Mar Oceanogr*. 53:9–17. doi:10.4067/S0718-19572018000100009
- Davidson JD, Landry T, Johnson G, Duncan A. 2012. Evaluating the productivity and cost benefit of high pressure water treatment regimes on *Ciona intestinalis*

- infested mussel stocks on Prince Edward Island. *J Shellfish Res.* 31:274–274.
- Deady S, Varian SJA, Fives JM. 1995. The use of cleanerfish to control sea lice on two Irish salmon (*Salmo salar*) farms with particular reference to wrasse behaviour in salmon cages. *Aquaculture.* 131:73–90. doi:10.1016/0044-8486(94)00331-H
- Denny CM. 2008. Development of a method to reduce the spread of the ascidian *Didemnum vexillum* with aquaculture transfers. *ICES J Mar Sci.* 65:805–810. doi:10.1093/icesjms/fsn039
- Dunham A, Marshall RD. 2012. Using stocking density modifications and novel growth medium to control shell deformities and biofouling in suspended culture of bivalves. *Aquaculture.* 324:234–241.
- Dürr S, Watson DI. 2010. Biofouling and antifouling in aquaculture. In: Dürr S, Thomason JC, editors. *Biofouling.* West Sussex (UK): Wiley-Blackwell; p. 267–287.
- Edwards CD, Pawluk KA, Cross SF. 2015. The effectiveness of several commercial antifouling treatments at reducing biofouling on finfish aquaculture cages in British Columbia. *Aquac Res.* 46:2225–2235. doi:10.1111/are.12380
- Eliassen K, Danielsen E, Johannesen Á, Joensen LL, Patursson EJ. 2018. The cleaning efficacy of lumpfish (*Cyclopterus lumpus* L.) in Faroese salmon (*Salmo salar* L.) farming pens in relation to lumpfish size and seasonality. *Aquaculture.* 488:61–65. doi:10.1016/j.aquaculture.2018.01.026
- Fan S, Fu M, Wang Z, Zhang X, Song W, Li Y, Liu G, Shi X, Wang X, Zhu M. 2015. Temporal variation of green macroalgal assemblage on *Porphyra* aquaculture rafts in the Subei Shoal, China. *Estuar Coast Shelf Sci.* 163: 23–28. doi:10.1016/j.ecss.2015.03.016
- FAO. 2002. *The State of World Fisheries and Aquaculture 2002.* Rome (Italy).
- FAO. 2018. *The State of World Fisheries and Aquaculture 2018—meeting the sustainable development goals.* Rome (Italy).
- Filion-Myklebust C, Norton TA. 1981. Epidermis shedding in the brown seaweed *Ascophyllum nodosum* (L.) Le Jolis, and its ecological significance. *Mar Biol Lett.* 2:45–51.
- Fitridge I, Dempster T, Guenther J, de Nys R. 2012. The impact and control of biofouling in marine aquaculture: a review. *Biofouling.* 28:649–669. doi:10.1080/08927014.2012.700478
- Fitridge I, Keough MJ. 2013. Ruinous resident: the hydroid *Ectopleura crocea* negatively affects suspended culture of the mussel *Mytilus galloprovincialis*. *Biofouling.* 29: 119–131. doi:10.1080/08927014.2012.752465
- Fletcher LM, Forrest BM, Bell JJ. 2013. Impacts of the invasive ascidian *Didemnum vexillum* on green-lipped mussel *Perna canaliculus* aquaculture in New Zealand. *Aquacult Environ Interact.* 4:17–30. doi:10.3354/aei00069
- Fletcher RL. 1995. Epiphytism and fouling in *Gracilaria* cultivation: an overview. *J Appl Phycol.* 7:325–333. doi: 10.1007/BF00004006
- Floerl O, Sunde LM, Bloecher N. 2016. Potential environmental risks associated with biofouling management in salmon aquaculture. *Aquacult Environ Interact.* 8: 407–417. doi:10.3354/aei00187
- Førde H, Forbord S, Handå A, Fossberg J, Arff J, Johnsen G, Reitan KI. 2016. Development of bryozoan fouling on cultivated kelp (*Saccharina latissima*) in Norway. *J Appl Phycol.* 28:1225–1234. doi:10.1007/s10811-015-0606-5
- Forrest BM, Atalah J. 2017. Significant impact from blue mussel *Mytilus galloprovincialis* biofouling on aquaculture production of green-lipped mussels in New Zealand. *Aquacult Environ Interact.* 9:115–126. doi:10.3354/aei00220
- Forrest BM, Hopkins GA, Dodgshun TJ, Gardner J. 2007. Efficacy of acetic acid treatments in the management of marine biofouling. *Aquaculture.* 262:319–332. doi:10.1016/j.aquaculture.2006.11.006
- Friedlander M, Weintraub N, Freedman A, Sheer J, Snovsky Z, Shapiro J, Kissil GW. 1996. Fish as potential biocontrollers of *Gracilaria* (Rhodophyta) culture. *Aquaculture.* 145: 113–118. doi:10.1016/S0044-8486(96)01350-6
- Gansel LC, Endresen PC, Steinhovden KB, Dahle SW, Svendsen E, Forbord S, Jensen Ø. 2017. Drag on nets fouled with blue mussel (*Mytilus edulis*) and sugar kelp (*Saccharina latissima*) and parameterization of fouling. *ASME Proceedings of the 36th International Conference on Ocean, Offshore and Arctic Engineering, Volume 6, Ocean Space Utilization: V006T005A008.*
- Gansel LC, Plew DR, Endresen PC, Olsen AI, Misimi E, Guenther J, Jensen O. 2015. Drag of clean and fouled net panels—measurements and parameterization of fouling. *PLoS One.* 10:e0131051. doi:10.1371/journal.pone.0131051
- Gevaux L, Lejars M, Margailan A, Briand J-F, Bunet R, Bressy C. 2019. Hydrolyzable additive-based silicone elastomers: a new approach for antifouling coatings. *Polymers.* 11:305. doi:10.3390/polym11020305
- Guardiola FA, Cuesta A, Meseguer J, Esteban MA. 2012. Risks of using antifouling biocides in aquaculture. *Int J Mol Sci.* 13:1541–1560. doi:10.3390/ijms13021541
- Guenther J, Misimi E, Sunde LM. 2010. The development of biofouling, particularly the hydroid *Ectopleura larynx*, on commercial salmon cage nets in Mid-Norway. *Aquaculture.* 300:120–127. doi:10.1016/j.aquaculture.2010.01.005
- Halat L, Galway ME, Gitto S, Garbary DV. 2015. Epidermal shedding in *Ascophyllum nodosum* (Phaeophyceae) seasonality, productivity and relationship to harvesting. *Phycologia.* 54:599–608. doi:10.2216/15-32.1
- Harrison PJ, Hurd CL. 2001. Nutrient physiology of seaweeds: application of concepts to aquaculture. *Cah Biol Mar.* 42:71–82.
- Hayashi L, Hurtado AQ, Msuya FE, Bleicher-Lhonneur G, Critchley AT. 2010. A review of *Kappaphycus* farming: prospects and constraints. In: Israel A, Einav R, Seckbach J, editors. *Seaweeds and their role in globally changing environments.* New York (NY): Springer; p. 251–283.
- Hellebø A, Stene A, Aspehaug V. 2017. PCR survey for *Paramoeba perurans* in fauna, environmental samples and fish associated with marine farming sites for Atlantic salmon (*Salmo salar* L.). *J Fish Dis.* 40:661–670. doi:10.1111/jfd.12546
- Helmholz H, Johnston BD, Ruhnau C, Prange A. 2010. Gill cell toxicity of northern boreal scyphomedusae *Cyanea capillata* and *Aurelia aurita* measured by an in vitro cell

- assay. *Hydrobiologia*. 645:223–234. doi:10.1007/s10750-010-0216-9
- Hillock KA, Costello MJ. 2013. Tolerance of the invasive tunicate *Styela clava* to air exposure. *Biofouling*. 29: 1181–1187. doi:10.1080/08927014.2013.832221
- Hollenbeck CM, Johnston IA. 2018. Genomic tools and selective breeding in molluscs. *Front Genet*. 9:253. doi: 10.3389/fgene.2018.00253
- Holthuis TD, Bergström P, Lindegarth M, Lindegarth S. 2015. Monitoring recruitment patterns of mussels and fouling tunicates in mariculture. *J Shellfish Res*. 34: 1007–1018. doi:10.2983/035.034.0327
- Hopkins GA, Forrest BM, Piola RF, Gardner J. 2011. Factors affecting survivorship of defouled communities and the effect of fragmentation on establishment success. *J Exp Mar Biol Ecol*. 396:233–243. doi:10.1016/j.jembe.2010.10.027
- Hopkins GA, Prince M, Cahill PL, Fletcher LM, Atalah J. 2016. Desiccation as a mitigation tool to manage biofouling risks: trials on temperate taxa to elucidate factors influencing mortality rates. *Biofouling*. 32:1–11. doi:10.1080/08927014.2015.1115484
- Hurtado AQ, Critchley AT. 2013. Impact of Acadian Marine Plant Extract Powder (AMPEP) in *Kappaphycus* production. *Malay J Sci*. 32:239–252.
- Hurtado AQ, Critchley AT. 2018. A review of multiple bio-stimulant and bioeffector benefits of AMPEP, an extract of the brown alga *Ascophyllum nodosum*, as applied to the enhanced cultivation and micropropagation of the commercially important red algal carrageenophyte *Kappaphycus alvarezii* and its selected cultivars. *J Appl Phycol*. 30:2859–2873. doi:10.1007/s10811-018-1407-4
- Hurtado AQ, Critchley AT, Trespoey A, Lhonneur GB. 2006. Occurrence of *Polysiphonia* epiphytes in *Kappaphycus* farms at Calaguas Is., Camarines Norte, Philippines. *J Appl Phycol*. 18:301–306. doi:10.1007/s10811-006-9032-z
- Hurtado AQ, Joe M, Sanares RC, Fan D, Prithiviraj B, Critchley AT. 2012. Investigation of the application of Acadian Marine Plant Extract Powder (AMPEP) to enhance the growth, phenolic content, free radical scavenging, and iron chelating activities of *Kappaphycus* Doty (Solieriaceae, Gigartinales, Rhodophyta). *J Appl Phycol*. 24:601–611. doi:10.1007/s10811-011-9785-x
- Hurtado AQ, Yunque DA, Tibubos K, Critchley AT. 2009. Use of Acadian marine plant extract powder from *Ascophyllum nodosum* in tissue culture of *Kappaphycus* varieties. *J Appl Phycol*. 21:633–639. doi:10.1007/s10811-008-9395-4
- Imsland AK, Reynolds P, Eliassen G, Hangstad TA, Nytrø AV, Foss A, Vikingstad E, Elvegård TA. 2015. Feeding preferences of lumpfish (*Cyclopterus lumpus* L.) maintained in open net-pens with Atlantic salmon (*Salmo salar* L.). *Aquaculture*. 436:47–51. doi:10.1016/j.aquaculture.2014.10.048
- Jormalainen V, Honkanen T. 2008. Macroalgal chemical defenses and their roles in structuring temperate marine communities. In: Amsler CD, editor. *Algal chemical ecology*. Berlin (Germany): Springer; p. 57–89.
- Kalantzi I, Zeri C, Catsiki V-A, Tsangaris C, Stroglyoudi E, Kaberi H, Vergopoulos N, Tsapakis M. 2016. Assessment of the use of copper alloy aquaculture nets: potential impacts on the marine environment and on the farmed fish. *Aquaculture*. 465:209–222. doi:10.1016/j.aquaculture.2016.09.016
- Kang J-Y, Bangoura I, Cho J-Y, Joo J, Choi YS, Hwang DS, Hong Y-K. 2016. Antifouling effects of the periostracum on algal spore settlement in the mussel *Mytilus edulis*. *Fish Aquat Sci*. 19:1–6.
- Keesing JK, Liu D, Shi Y, Wang Y. 2016. Abiotic factors influencing biomass accumulation of green tide causing *Ulva* spp. on *Pyropia* culture rafts in the Yellow Sea, China. *Mar Pollut Bull*. 105:88–97. doi:10.1016/j.marpollbul.2016.02.051
- Kim JK, Yarish C, Hwang EK, Park M, Kim Y. 2017. Seaweed aquaculture: cultivation technologies, challenges and its ecosystem services. *Algae*. 32:1–13. doi:10.4490/algae.2017.32.3.3
- Kuschel FA, Buschmann AH. 1991. Abundance, effects and management of epiphytism in intertidal cultures of *Gracilaria* (Rhodophyta) in southern Chile. *Aquaculture*. 92:7–19. doi:10.1016/0044-8486(91)90004-Q
- Kvenseth PG. 1996. Large-scale use of wrasse to control sea lice and net fouling in salmon farms in Norway. In: Sayer MDJ, Treasurer JW, Costello MJ, editors. *Wrasse: biology and use in aquaculture*. West Sussex (UK): Wiley-Blackwell; p. 196–203.
- Lachnit T, Fischer M, Kunzel S, Baines JF, Harder T. 2013. Compounds associated with algal surfaces mediate epiphytic colonization of the marine macroalga *Fucus vesiculosus*. *FEMS Microbiol Ecol*. 84:411–420. doi:10.1111/1574-6941.12071
- Lader P, Fredriksson DW, Guenther J, Volent Z, Blocher N, Kristiansen D, Gansel L, Decew J. 2015. Drag on hydroid-fouled nets—an experimental approach. *China Ocean Eng*. 29:369–389. doi:10.1007/s13344-015-0026-y
- Lane A, Willemsen PR. 2004. Collaborative effort looks into biofouling. *Fish Farming Int*. 2004:34–35.
- Leclercq E, Zerafa B, Brooker AJ, Davie A, Migaud H. 2018. Application of passive-acoustic telemetry to explore the behaviour of ballan wrasse (*Labrus bergylta*) and lumpfish (*Cyclopterus lumpus*) in commercial Scottish salmon sea-pens. *Aquaculture*. 495:1–12. doi:10.1016/j.aquaculture.2018.05.024
- Leonardi PI, Miravalles AB, Faugeton S, Flores V, Beltrán J, Correa JA. 2006. Diversity, phenomenology and epidemiology of epiphytism in farmed *Gracilaria chilensis* (Rhodophyta) in northern Chile. *Eur J Phycol*. 41: 247–257. doi:10.1080/09670260600645659
- Li J, Yang C, Wang Q, Du X, Deng Y. 2018. Growth and survival of host pearl oyster *Pinctada fucata martensii* (Dunker, 1880) treated by different biofouling-clean methods in China. *Estuar Coast Shelf Sci*. 207:104–108. doi:10.1016/j.ecss.2018.04.009
- Littler MM, Littler DS. 1999. Blade abandonment/proliferation: a novel mechanism for rapid epiphyte control in marine macrophytes. *Ecology*. 80:1736–1746. doi:10.1890/0012-9658(1999)080[1736:BAPANM]2.0.CO;2
- Liu D, Keesing JK, He P, Wang Z, Shi Y, Wang Y. 2013. The world's largest macroalgal bloom in the Yellow Sea, China: formation and implications. *Estuar Coast Shelf Sci*. 129:2–10. doi:10.1016/j.ecss.2013.05.021
- Liu Y, Suo XK, Wang Z, Gong YF, Wang X, Li H. 2017. Developing polyimide-copper antifouling coatings with

- capsule structures for sustainable release of copper. *Mater Design*. 130:285–293. doi:10.1016/j.matdes.2017.05.075
- Lodeiros CJM, Himmelman JH. 1996. Influence of fouling on the growth and survival of the tropical scallop, *Euvola (Pecten) ziczac* (L. 1758) in suspended culture. *Aquaculture Res*. 27:749–756. doi:10.1111/j.1365-2109.1996.tb01233.x
- Longtin CM, Scrosati RA, Whalen GB, Garbary DJ. 2009. Distribution of algal epiphytes across environmental gradients at different scales: intertidal elevation, host canopies, and host fronds(1). *J Phycol*. 45:820–827. doi:10.1111/j.1529-8817.2009.00710.x
- Lüning K, Pang S. 2003. Mass cultivation of seaweeds: current aspects and approaches. *J Appl Phycol*. 15:115–119. doi:10.1023/A:1023807503255
- Madin J, Ching CV. 2015. Biofouling challenge and management methods in marine aquaculture. In: Mustafa S, Shapawi R, editors. *Aquaculture ecosystems: adaptability and sustainability*. West Sussex (UK): John Wiley & Sons, Ltd; p. 107–138.
- Magin CM, Cooper SP, Brennan AB. 2010. Non-toxic antifouling strategies. *Mater Today*. 13:36–44. doi:10.1016/S1369-7021(10)70058-4
- Makridis P, Mente E, Grundvig H, Gausen M, Koutsikopoulos C, Bergheim A. 2018. Monitoring of oxygen fluctuations in seabass cages (*Dicentrarchus labrax* L.) in a commercial fish farm in Greece. *Aquac Res*. 49:684–691. doi:10.1111/are.13498
- Marinho GS, Holdt SL, Birkeland MJ, Angelidaki I. 2015. Commercial cultivation and bioremediation potential of sugar kelp, *Saccharina latissima*, in Danish waters. *J Appl Phycol*. 27:1963–1973. doi:10.1007/s10811-014-0519-8
- Marroig RG, Reis RP. 2011. Does biofouling influence *Kappaphycus alvarezii* (Doty) Doty ex Silva farming production in Brazil? *J Appl Phycol*. 23:925–931. doi:10.1007/s10811-010-9602-y
- Marroig RG, Reis RP. 2016. Biofouling in Brazilian commercial cultivation of *Kappaphycus alvarezii* (Doty) Doty ex P. C. Silva. *J Appl Phycol*. 28:1803–1813. doi:10.1007/s10811-015-0713-3
- Marshall RD, Dunham A. 2013. Effects of culture media and stocking density on biofouling, shell shape, growth, and survival of the Pacific oyster (*Crassostrea gigas*) and the Manila clam (*Venerupis philippinarum*) in suspended culture. *Aquaculture*. 406:68–78.
- Mineur F, Cook EJ, Minchin D, Bohn K, MacLeod A, Maggs CA. 2012. Changing coasts: marine aliens and artificial structures. *Oceanogr Mar Biol*. 50:189–233.
- Moodie LWK, Trepos R, Cervin G, Larsen L, Larsen DS, Pavia H, Hellio C, Cahill P, Svenson J. 2017. Probing the structure–activity relationship of the natural antifouling agent polygodial against both micro-and macrofoulers by semisynthetic modification. *J Nat Prod*. 80:515–525. doi:10.1021/acs.jnatprod.6b01056
- Nikolaou M, Neofitou N, Skordas K, Castritsi-Catharios I, Tziantziou L. 2014. Fish farming and anti-fouling paints: a potential source of Cu and Zn in farmed fish. *Aquacult Environ Interact*. 5:163–171. doi:10.3354/aei00101
- Nir S, Reches M. 2016. Bio-inspired antifouling approaches: the quest towards non-toxic and non-biocidal materials. *Curr Opin Biotechnol*. 39:48–55. doi:10.1016/j.copbio.2015.12.012
- Nurioglu AG, Esteves ACC, de With G. 2015. Non-toxic, non-biocide-release antifouling coatings based on molecular structure design for marine applications. *J Mater Chem B*. 3:6547–6570. doi:10.1039/C5TB00232J
- Nylund GM, Pavia H. 2005. Chemical versus mechanical inhibition of fouling in the red alga *Dilsea carnosa*. *Mar Ecol Prog Ser*. 299:111–121. doi:10.3354/meps299111
- Oliveira IB, Beiras R, Thomas KV, Suter MJ, Barroso CM. 2014. Acute toxicity of tralopyryl, capsaicin and triphenylborane pyridine to marine invertebrates. *Ecotoxicology*. 23:1336–1344. doi:10.1007/s10646-014-1276-9
- Oliveira IB, Groh KJ, Stadnicka-Michalak J, Schonenberger R, Beiras R, Barroso CM, Langford KH, Thomas KV, Suter MJ. 2016. Tralopyryl bioconcentration and effects on the gill proteome of the Mediterranean mussel *Mytilus galloprovincialis*. *Aquat Toxicol*. 177:198–210. doi:10.1016/j.aquatox.2016.05.026
- Othmani A, Briand JF, Aye M, Molmeret M, Culioli G. 2016. Surface metabolites of the brown alga *Taonia atomaria* have the ability to regulate epibiosis. *Biofouling*. 32:801–813. doi:10.1080/08927014.2016.1198954
- Park CS, Hwang EK. 2012. Seasonality of epiphytic development of the hydroid *Obelia geniculata* on cultivated *Saccharina japonica* (Laminariaceae, Phaeophyta) in Korea. *J Appl Phycol*. 24:433–439. doi:10.1007/s10811-011-9755-3
- Pereira RC, da Silva Costa E, Sudatti DB, da Gama B. 2017. Inducible defenses against herbivory and fouling in seaweeds. *J Sea Res*. 122:25–33. doi:10.1016/j.seares.2017.03.002
- Peteiro C, Freire O. 2013. Epiphytism on blades of the edible kelps *Undaria pinnatifida* and *Saccharina latissima* farmed under different abiotic conditions. *J World Aquacult Soc*. 44:706–715. doi:10.1111/jwas.12065
- Pietrak MR, Molloy SD, Bouchard DA, Singer JT, Bricknell I. 2012. Potential role of *Mytilus edulis* in modulating the infectious pressure of *Vibrio anguillarum* 02β on an integrated multi-trophic aquaculture farm. *Aquaculture*. 326-329:36–39. doi:10.1016/j.aquaculture.2011.11.024
- Pit JH, Southgate PC. 2003. Fouling and predation; how do they affect growth and survival of the blacklip pearl oyster, *Pinctada margaritifera*, during nursery culture? *Aquacult Int*. 11:545–555. doi:10.1023/B:AQUI.0000013310.17400.97
- Powell A, Treasurer JW, Pooley CL, Keay AJ, Lloyd R, Imsland AK, Garcia de Leaniz C. 2018. Use of lumpfish for sea-lice control in salmon farming: challenges and opportunities. *Rev Aquacult*. 10:683–702. doi:10.1111/raq.12194
- Pradhan S, Kumar S, Mohanty S, Nayak SK. 2018. Environmentally benign fouling-resistant marine coatings: a review. *Polym Plast Technol Mater*. 58:498–518.
- Rensel JE, Whyte JNC. 2003. Finfish mariculture and harmful algal blooms. In: Hallegraeff GM, Anderson DM, Cembella AD, editors. *Manual on harmful marine microalgae*. Paris (France): UNESCO Publishing; p. 693–722.
- Roche RC, Monnington JM, Newstead RG, Sambrook K, Griffith K, Holt RHF, Jenkins SR. 2015. Recreational

- vessels as a vector for marine non-natives: developing biosecurity measures and managing risk through an in-water encapsulation system. *Hydrobiologia*. 750:187–199. doi:10.1007/s10750-014-2131-y
- Rolheiser KC, Dunham A, Switzer SE, Pearce CM, Therriault TW. 2012. Assessment of chemical treatments for controlling *Didemnum vexillum*, other biofouling, and predatory sea stars in Pacific oyster aquaculture. *Aquaculture*. 364:53–60.
- Rolin C, Inkster R, Laing J, McEvoy L. 2017. Regrowth and biofouling in two species of cultivated kelp in the Shetland Islands. *J Appl Phycol*. 29:2351–2361. doi:10.1007/s10811-017-1092-8
- Rosa M, Holohan BA, Shumway SE, Bullard SG, Wikfors GH, Morton S, Getchis T. 2013. Biofouling ascidians on aquaculture gear as potential vectors of harmful algal introductions. *Harmful Algae*. 23:1–7. doi:10.1016/j.hal.2012.11.008
- Sato T, Fujimori Y, Nakayama T, Gotoh Y, Sunaga Y, Nemoto M, Matsunaga T, Tanaka T. 2012. Assessment of the anti-biofouling potentials of a copper iodide-doped nylon mesh. *Appl Microbiol Biotechnol*. 95:1043–1050. doi:10.1007/s00253-011-3720-6
- Sato-Okoshi W, Abe H. 2013. Morphology and molecular analysis of the 18S rRNA gene of oyster shell borers, *Polydora* species (Polychaeta: Spionidae), from Japan and Australia. *J Mar Biol Assoc UK*. 93:1279–1286. doi:10.1017/S002531541200152X
- Scardino AJ, de Nys R. 2011. Mini review: biomimetic models and bioinspired surfaces for fouling control. *Biofouling*. 27:73–86. doi:10.1080/08927014.2010.536837
- Scianni C, Falkner M, DeBruyckere L. 2017. Biofouling in the U.S. Pacific States and British Columbia. White Paper prepared for the Coastal Committee of the Western Regional Panel on Aquatic Nuisance Species.
- Shacklock PF, Doyle RW. 1983. Control of epiphytes in seaweed cultures using grazers. *Aquaculture*. 31:141–151. doi:10.1016/0044-8486(83)90308-3
- Shirakashi S, Hirano C. 2015. Accumulation and distribution of skin fluke *Neobenedenia girellae* eggs on a culture cage. *Aquaculture*. 443:1–4. doi:10.1016/j.aquaculture.2015.02.036
- Sievers M, Dempster T, Fitridge I, Keough MJ. 2014. Monitoring biofouling communities could reduce impacts to mussel aquaculture by allowing synchronisation of husbandry techniques with peaks in settlement. *Biofouling*. 30:203–212. doi:10.1080/08927014.2013.856888
- Sievers M, Dempster T, Keough M, Fitridge I. 2019. Methods to prevent and treat biofouling in shellfish aquaculture. *Aquaculture*. 505:263–270. doi:10.1016/j.aquaculture.2019.02.071
- Sievers M, Fitridge I, Bui S, Dempster T. 2017. To treat or not to treat: a quantitative review of the effect of biofouling and control methods in shellfish aquaculture to evaluate the necessity of removal. *Biofouling*. 33:755–767. doi:10.1080/08927014.2017.1361937
- Sievers M, Fitridge I, Dempster T, Keough MJ. 2013. Biofouling leads to reduced shell growth and flesh weight in the cultured mussel *Mytilus galloprovincialis*. *Biofouling*. 29:97–107. doi:10.1080/08927014.2012.749869
- Simkanin C, Davidson IC, Dower JF, Jamieson G, Therriault TW. 2012. Anthropogenic structures and the infiltration of natural benthos by invasive ascidians. *Mar Ecol-Evol Persp*. 33:499–511. doi:10.1111/j.1439-0485.2012.00516.x
- Sim-Smith C, Forsythe A. 2013. Comparison of the international regulations and best management practices for marine finfish farming. Ministry for Primary Industries, New Zealand Government.
- Skarbøvik E, Allan I, Sample JE, Greipsland I, Selvik JR, Schanke LB, Beldring S, Stålnacke P, Kaste Ø. 2017. Riverine inputs and direct discharges to Norwegian coastal waters - 2016. Norwegian Institute for Water Research; NIVA Report No. 7217-2017.
- Smit AJ, Fourie AM, Robertson BL, Du Preez DR. 2003. Control of the herbivorous isopod, *Paridotea reticulata*, in *Gracilaria gracilis* tank cultures. *Aquaculture*. 217:385–393. doi:10.1016/S0044-8486(02)00412-X
- Soliman T, Inglis GJ. 2018. Forecasting the economic impacts of two biofouling invaders on aquaculture production of green-lipped mussels *Perna canaliculus* in New Zealand. *Aquacult Environ Interact*. 10:1–12. doi:10.3354/aei00249
- Song W, Jiang M, Wang Z, Wang H, Zhang X, Fu M. 2018. Source of propagules of the fouling green macroalgae in the Subei Shoal. *Acta Oceanol Sin*. 37:102–108. doi:10.1007/s13131-018-1169-5
- Sterling AM, Cross SF, Pearce CM. 2016. Co-culturing green sea urchins (*Strongylocentrotus droebachiensis*) with mussels (*Mytilus* spp.) to control biofouling at an integrated multi-trophic aquaculture site. *Aquaculture*. 464:253–261. doi:10.1016/j.aquaculture.2016.06.010
- Stien LH, Lind MB, Oppedal F, Wright DW, Seternes T. 2018. Skirts on salmon production cages reduced salmon lice infestations without affecting fish welfare. *Aquaculture*. 490:281–287. doi:10.1016/j.aquaculture.2018.02.045
- Sugihara Y, Yamada T, Ogawa K, Yokoyama F, Matsukura K, Kanai K. 2015. Occurrence of the bluefin tuna blood fluke *Cardicola opisthorchis* in the intermediate host *Terebella* sp. *Fish Pathol*. 50:105–111. doi:10.3147/jsfp.50.105
- Sugihara Y, Yamada T, Tamaki A, Yamanishi R, Kanai K. 2014. Larval stages of the bluefin tuna blood fluke *Cardicola opisthorchis* (Trematoda: Aporocotylidae) found from *Terebella* sp. (Polychaeta: Terebellidae). *Parasitol Int*. 63:295–299. doi:10.1016/j.parint.2013.11.010
- Swain G, Shinjo N. 2014. Comparing biofouling control treatments for use on aquaculture nets. *Int J Mol Sci*. 15:22142–22154. doi:10.3390/ijms151222142
- Tettelbach ST, Tetrault K, Carroll J. 2014. Efficacy of Netminder®silicone release coating for biofouling reduction in bay scallop grow-out and comparative effects on scallop survival, growth and reproduction. *Aquac Res*. 45:234–242. doi:10.1111/j.1365-2109.2012.03220.x
- Tsukrov I, Drach A, DeCew J, Robinson Swift M, Celikkol B. 2011. Characterization of geometry and normal drag coefficients of copper nets. *Ocean Eng*. 38:1979–1988. doi:10.1016/j.oceaneng.2011.09.019
- Vairappan CS, Chung CS, Hurtado AQ, Soya FE, Lhonneur GB, Critchley A. 2008. Distribution and symptoms of epiphyte infection in major carrageenophyte-producing

- farms. *J Appl Phycol.* 20:477–483. doi:10.1007/s10811-007-9299-8
- Vaquer-Sunyer R, Duarte CM. 2010. Sulfide exposure accelerates hypoxia-driven mortality. *Limnol Oceanogr.* 55:1075. doi:10.4319/lo.2010.55.3.1075
- Veeragurunathan V, Eswaran K, Malarvizhi J, Gobalakrishnan M. 2015. Cultivation of *Gracilaria dura* in the open sea along the southeast coast of India. *J Appl Phycol.* 27:2353–2365. doi:10.1007/s10811-014-0514-0
- Vucko MJ, King PC, Poole AJ, Hu Y, Jahedi MZ, de Nys R. 2014. Assessing the antifouling properties of cold-spray metal embedment using loading density gradients of metal particles. *Biofouling.* 30:651–666. doi:10.1080/08927014.2014.906584
- Vucko MJ, Poole AJ, Carl C, Sexton BA, Glenn FL, Whalan S, de Nys R. 2014. Using textured PDMS to prevent settlement and enhance release of marine fouling organisms. *Biofouling.* 30:1–16. doi:10.1080/08927014.2013.836507
- Vucko MJ, Poole AJ, Sexton BA, Glenn FL, Carl C, Whalan S, de Nys R. 2013. Combining a photocatalyst with microtopography to develop effective antifouling materials. *Biofouling.* 29:751–762. doi:10.1080/08927014.2013.803193
- Walls AM, Edwards MD, Firth LB, Johnson MP. 2017. Successional changes of epibiont fouling communities of the cultivated kelp *Alaria esculenta*: predictability and influences. *Aquacult Environ Interact.* 9:57–71. doi:10.3354/aei00215
- Wang S, Weinberger F, Lenz M. 2018. Fluctuations in the strength of chemical antifouling defenses in a red macroalga in response to variations in epibiont colonization pressure. *Mar Biol.* 165:107. doi:10.1007/s00227-018-3365-4
- Watts AM, Hopkins GA, Goldstien SJ. 2015. Characterising biofouling communities on mussel farms along an environmental gradient: a step towards improved risk management. *Aquacult Environ Interact.* 8:15–30. doi:10.3354/aei00159
- Woods CMC, Floerl O, Hayden BJ. 2012. Biofouling on Greenshell™ mussel (*Perna canaliculus*) farms: a preliminary assessment and potential implications for sustainable aquaculture practices. *Aquacult Int.* 20:537–557. doi:10.1007/s10499-011-9484-2
- Wybourne B. 2013. Defensive acontia of the white-striped anemone (*Actinotoe albocincta*) from New Zealand King Salmon Clay Point salmon farm. Report prepared for NZ King Salmon Company Ltd. Skretting.
- Yamamoto K, Endo H, Yoshikawa S, Ohki K, Kamiya M. 2013. Various defense ability of four sargassacean algae against the red algal epiphyte *Neosiphonia harveyi* in Wakasa Bay, Japan. *Aquat Bot.* 105:11–17. doi:10.1016/j.aquabot.2012.10.008
- Yigit M, Celikkol B, Ozalp B, Bulut M, Dwyer RL, Yilmaz S, Maita M, Buyukates Y. 2018. Comparison of copper alloy mesh with conventional nylon nets in offshore cage farming of gilthead seabream (*Sparus aurata*). *Aquacult Stud.* 18:57–65.
- Zhang Y, He P, Li H, Li G, Liu J, Jiao F, Zhang J, Huo Y, Shi X, Su R. 2019. *Ulva prolifera* green tide outbreaks and their environmental impact in the Yellow Sea, China. *Natl Sci Rev.* nzw026. doi:10.1093/nsr/nzw026