

Biological evaluation of non-steroidal anti-inflammatory drugs-cobalt(II) complexes

**Filitsa Dimiza¹, Athanasios N. Papadopoulos², Vassilis Tangoulis¹, Vassilis Psycharis³,
Catherine P. Raptopoulou³, Dimitris P. Kessissoglou¹, George Psomas^{1,*}**

¹ *Department of General and Inorganic Chemistry, Faculty of Chemistry, Aristotle University of Thessaloniki, P.O. Box 135, GR-54124 Thessaloniki, GREECE*

² *Department of Nutrition and Dietetics, Faculty of Food Technology and Nutrition, Alexandrion Technological Educational Institution, Sindos, Thessaloniki, GREECE.*

³ *Institute of Materials Science, NCSR "Demokritos", GR-15310 Aghia Paraskevi Attikis, GREECE*

Supplementary material

* Corresponding author: Dr. George Psomas, Tel.:+30+2310997790; Fax:+30+2310997738; E-mail: gepsomas@chem.auth.gr

Figure S1. X-Band EPR spectra of (A) **1** and (B) **2** in powder samples (up) and in frozen dmsol solution (down) at 4K.




Figure S2. Stern-Volmer quenching plot of EB bound to CT DNA for Hmef and complexes **1-4**.




Figure S3. Stern-Volmer quenching plot of BSA for Hmef and complexes **1-4**.




Figure S4. Scatchard plot of BSA for Hmef and complexes **1-4**.




Figure S5. Stern-Volmer quenching plot of HSA for Hmef and complexes **1-4**.




Figure S6. Scatchard plot of HSA for Hmef and complexes **1-4**.

