

1 Biomass-modulated fire dynamics during the last glacial-interglacial
2 transition at the Central Pyrenees (Spain)

3 Graciela Gil-Romera^{1*}, Penélope González-Sampériz¹, Laura Lasheras-Álvarez¹, Miguel Sevilla-Callejo¹,
4 Ana Moreno¹, Blas Valero-Garcés¹, Lourdes López-Merino², José S. Carrión³, Ana Pérez Sanz¹,
5 Josu Aranbarri¹, Eduardo García-Prieto Fonce¹

6 1 Quaternary Palaeoenvironments and Global Change Dept., Instituto Pirenaico de Ecología-CSIC,
7 Campus Aula Dei, Avda. Montañana 1005, 50059 Zaragoza (Spain) | 2 Institute for the Environment,
8 Brunel University, Uxbridge, London, Middlesex UB8 3PH (UK) | Department of Plant Biology, Facultad de
9 Biología, Universidad de Murcia. Campus del Espinardo, Murcia (Spain)

10 * Corresponding author - graciela.gil@ipe.csic.es

11

12

13 **Abstract**

14 Understanding long-term fire ecology is essential for current day interpretation of ecosystem fire
15 responses. However paleoecology of fire is still poorly understood, especially at high-altitude mountain
16 environments, despite the fact that these are fire-sensitive ecosystems and their resilience might be
17 affected by changing fire regimes. We reconstruct wildfire occurrence since the Lateglacial (14.7 cal ka
18 BP) to the Mid-Holocene (6 cal ka BP) and investigate the climate-fuel-fire relationships in a sedimentary
19 sequence located at the treeline in the Central Spanish Pyrenees. Pollen, macro- and micro-charcoal were
20 analysed for the identification of fire events (FE) in order to detect vegetation post-fire response and to
21 define biomass-fire interactions. Mean fire intervals (MFI) reduced since the Lateglacial, peaking at 9-7.7
22 cal ka BP while from 7.7 to 6 cal ka BP no fire is recorded. We hypothesize that Early Holocene maximum
23 summer insolation, as climate forcing, and mesophyte forest expansion, as a fuel-creating factor, were
24 responsible for accelerating fire occurrence in the Central Pyrenees treeline. We also found that fire had
25 long-lasting negative effects on most of the treeline plant communities and that forest contraction from 7.7
26 cal ka BP is likely linked to the ecosystem's threshold response to high fire frequencies.

27 **Keywords**

28 Fire history; Quaternary; Lateglacial; palaeoecology; historical biogeography; Iberian Peninsula.

29

30 Taxa nomenclature is based on Castroviejo et al. (1986).

32 Fire is a well-known transforming agent of the Earth system that, while well studied across different spatial
33 scales, it is best understood at short-term temporal scales. Fire occurrence is determined by a range of
34 factors that interact dynamically, such as climate, fuel availability and flammability, landscape structure,
35 amongst others, defining fire regimes. Sensitive ecosystems to the current global environmental change
36 experience fire regime changes which are often responsible for deep modifications of their biodiversity and
37 resilience patterns (Bond and Keeley, 2005). This could be the case for mountain ranges, which have
38 been spotlighted as one of the most endangered continental ecosystems because of their high levels of
39 plant endemism (Médail and Quézel, 1999) and their particular sensitivity to global change (Engler et al.,
40 2011; Gottfried et al., 2012). Assessing the potential ecosystem transformations caused by future changes
41 in fire regime at mountain ranges requires a comprehensive knowledge of their long-term fire ecology
42 (Gobet et al., 2003; Stähli et al., 2006; Tinner et al., 1999, 2005a). Thus, changes in fire regimes at
43 centennial to millennial time-scales have been thoroughly documented in the Alps (Blarquez and
44 Carcaillet, 2010; Colombaroli et al., 2009, 2010; Stähli et al., 2006; Tinner and Lotter, 2001) and at lower
45 altitudes of the Northern Pyrenees (Rius et al., 2009, 2011a, 2011b). In the Iberian Peninsula, past fire
46 histories and vegetation responses have been explored in the Mediterranean area (Carrión, 2002; Carrión
47 et al., 2004, 2007, 2010a, 2010b; Carrión-Marco, 2005; Franco Múgica et al., 2005; Gil-Romera et al.,
48 2010; Jiménez-Moreno et al., 2013; López-Merino et al., 2009), and in the Eurosiberian region of the
49 Iberian Peninsula (Carrión-Marco et al., 2010; Kaal et al., 2011; López-Merino et al., 2012; Morales-Molino
50 et al., 2013). However, the Iberian investigations are focused on regional fire activity, through examination
51 of micro-charcoal records so they do not tackle locally-occurred fire events due to the absence of
52 contiguous macro-charcoal datasets, thus precluding the estimation of fire frequencies or return intervals
53 (Power et al., 2008; Whitlock et al., 2010a, 2010b). Studies addressing long-term local fire events have
54 been barely carried out in the Iberian Peninsula, and they do not investigate the climate-fire-biomass
55 relationships or the ecosystem response to fire, as the record of Estany de Burg (Fig. 1) (Bal et al., 2011) .

56 The study presented here is focused on the long-term fire ecology of a fuel-limited alpine site located at
57 the Central Spanish Pyrenees, El Portalet, where a previous palynological analysis described a sequence
58 of environmental changes with high sensitivity to abrupt climate fluctuations since the Lateglacial
59 (González-Sampériz et al., 2006). Our objectives are to 1) analyse the Lateglacial to Mid-Holocene (ca.
60 14.7-6 cal ka BP) interactions amongst the different agents determining fire occurrence, i.e. climate and
61 fuel availability, in a fuel-limited environment, and 2) determine long-term plant responses and ecosystem
62 resilience to fire disturbance.

63 **2. Material and methods**

64 *2.1 Study area*

65 El Portalet site (1802 m asl) is located in the Central-western Pyrenees (Fig. 1), lying on Devonian and
66 Carboniferous shales and limestones. The site is currently a peatland situated within a small glacial cirque.
67 Climate is alpine, with high rainfall (mean annual precipitation ca. 1800 mm) mainly linked to the activity of
68 Atlantic fronts. Annual mean air temperature is 5°C and daily temperatures are below 0° C for more than
69 200 days along the year while mean summer temperature is ca. 12° C (AEMET-IM, 2011).

70 The area is currently devoid of woody vegetation, presenting some bushed spots formed by *Vaccinium*
71 *myrtillus* and mixed pasture-alpine grasslands (Fig. 2) probably established during the Middle Ages, when
72 intense deforestation was regularly practiced due to pastoral and slash-and-burn activities (Montserrat,
73 1992). However, the closest west-to-east valleys (Fig. 2) present *Pinus sylvestris* and *P. uncinata*
74 communities reaching the treeline (between 1500 and 1800 m asl), with a mixed scrubland formed by
75 *Juniperus* (*J. communis* ssp. *alpina*, *J. communis* ssp. *hemisphaerica*) and by *Buxus sempervirens* at
76 lower altitudes. *Abies alba* and *Fagus sylvatica* mixed forest appear in shady slopes between 1500 and
77 1700 m asl. Mixed temperate forests formed by a variety of mesophytes (*Corylus*, *Acer*, *Salix*, *Fraxinus*,
78 etc.) can be found at lower altitudes (1500 to 1600 m asl), often confined to riparian environments. *Betula*
79 communities (*B. pendula* and some remnants of *B. nana*) are present at different altitudes, forming
80 disperse forests, sheltered in areas controlled by moisture availability and reduced sun exposure.

81 Evergreen and semi-deciduous *Quercus* (*Q. ilex* ssp. *ballota* and *Q. faginea*) become abundant below
82 1500 m asl, where the latter prefers cooler and wetter locations.

83 2.2 Chronological framework and sampling resolution

84 El Portalet age-depth model was constructed from 13 radiocarbon dates obtained from pollen
85 concentrates and a wood fragment, along a 6.08 m long core, recording a time-span between ca. 32 and
86 5.5 cal ka BP (González-Sampérez et al., 2006). The analyses in this new study have been performed
87 between 14.7 to 12.5 cal ka BP (470-382 cm) and 10 to 6 cal ka BP (364-160 cm), the sections with higher
88 and relatively constant sedimentation rates (Table 1) and that provide examples of different climates and
89 biomass availabilities. The criterion to select these intervals is based on sedimentation issues, as the 382-
90 364 cm interval (ca. 12.5 and 10 cal ka BP) corresponds to a period of abrupt changes in the
91 sedimentation rate, with some missing cm of sediments, including some possible hiatuses during the
92 Younger Dryas stadial (Table 1) (González-Sampérez et al., 2006)

93 Different sampling resolution was applied for the pollen analysis of the two selected sections. The
94 Holocene section (10-6 cal ka BP), with sedimentation rate ca. 15 yr cm⁻¹, (Table 1), enabled a high-
95 resolution approach. Pollen, micro-charcoal (between 10µm and 150µm) and macro-charcoal
96 (particles > 150µm) particles were studied from contiguously taken samples (1 cm), permitting time-series
97 analyses and cross-correlation amongst the three proxies. The Lateglacial section (14.7-12.5 cal ka BP),
98 with a sedimentation rate ca. 24 yr cm⁻¹, was contiguously sampled for macro-charcoal analyses (1 cm),
99 and every 2 to 5 cm for pollen and micro-charcoal content (Table 1).

100 In this new study, a total of 292 samples were analysed for macro-charcoal, from which 235 were used for
101 micro-charcoal and pollen analyses. The 2006 study was limited to 100 pollen samples for this period, and
102 both micro- and macro-charcoal were not count (14.7-12.5 and 10-6 cal ka BP).

103 2.3 *Laboratory methods*

104 2.3.1 Charcoal analyses

105 Sedimentary macro-charcoal particles (>150 μm) have been proved to be a good indicator of past local
106 fire events (FE) at different time scales (Clark and Royall, 1996; Clark et al., 1996; Higuera et al., 2010;
107 Whitlock and Larsen, 2002). Estimating mean fire intervals (MFI) implies detecting local FE from macro-
108 charcoal particles (>150 μm) (see discussion in Whitlock and Larsen, 2002). Thus, macro-charcoal
109 particles were chemically processed and sieved at 150 μm mesh and retrieved particles were counted on
110 a binocular microscope (x40). Charcoal identification was accomplished according to existing literature,
111 counting opaque, angular particles (Clark et al., 1996; Finsinger and Tinner, 2005; Tinner and Hu, 2003;
112 Turner et al., 2004). Micro-charcoal particles (>10 μm) were counted on pollen slides under an optical
113 microscope (x250) being a proxy for regional fire activity. More details on laboratory methods can be
114 followed in Supplementary data.

115 Micro- and macro-charcoal concentrations (charcoal particles cm^{-3}) were transformed into accumulation
116 rates (particles $\text{cm}^{-3} \text{yr}^{-1}$). The macro-charcoal accumulation rate is referred as CHAR hereafter.

117 2.3.2 Pollen analysis

118 Palynological samples were chemically treated and a minimum of 300 terrestrial pollen grains were
119 counted, despite some of the samples displayed low counts (minimum=215, maximum=560,
120 mean=402.84, STD=106.7), excluding from the pollen sum spores, hydro- and hygrophyte pollen types,
121 and expressed in abundance (%). Mesophytes, other trees, shrubs and herbs pollen types were grouped
122 for representation and data analyses purposes. See Supplementary data for more details.

123 No plant macro-remains have been found in El Portalet sequence, and therefore fossil pollen is our only
124 proxy for vegetation change.

125

126 2.4 Data analyses

127 2.4.1 Fire events (FE) and Mean Fire Intervals (MFI)

128 Identifying FE – one or several fires occurring within the time resolution of a sample in a core (Higuera et
129 al., 2010; Whitlock and Larsen, 2002)– implies isolating charcoal peaks equivalent to the signal related to
130 local fire occurrence (Carcaillet et al., 2001; Gavin et al., 2006; Lynch et al., 2002). Peak identification was
131 performed using CharAnalysis (Higuera et al., 2009). MFI for particular periods are calculated as the age
132 between the first and the last fire event divided by the number of fire intervals in that period. Fire
133 frequencies (FF, number of fires/time unit) have been estimated every 500 years. See more details on
134 methods in Supplementary data.

135 2.4.2 Vegetation dynamics: numerical approach

136 Pollen assemblage zones and diagram plotting were performed with software Psimpoll 4.27 (Bennett,
137 2009). Palynological richness was estimated by rarefaction analysis, implemented in the open software
138 Analytic Rarefaction 2.0, and a Spearman's R correlation with different taxa and group of taxa was
139 performed to check whether richness is led by any of them. These were done in R v. 2.0.5 (R Core Team,
140 2012).

141 Cross-correlation analyses between CHAR and those pollen taxa with expected fire interactions following
142 the ecological literature (Paula et al., 2009) were performed to explore the effect of biomass accumulation
143 on fire occurrence and the potential feedbacks between local wildfire and vegetation response. Thus, 235
144 samples (corresponding to the 10-6 cal ka BP section) analysed at the same time resolution for pollen and
145 macro-charcoal were used after potential trends were removed by linearly detrending both data series
146 (Blarquez and Carcaillet, 2010; Colombaroli et al., 2007, 2008). CHAR was cross-correlated to pollen
147 percentages as a way to control potential common trends in sedimentation rates, avoiding potential
148 spurious correlations (Tinner et al., 1999). Cross-correlation coefficients were calculated at ± 10 lags
149 corresponding to ± 150 years (each lag corresponds to the mean time difference between two adjacent

150 samples, i.e. 15 years for the Holocene section of the record). Negative lags represent the effect of the
151 second variable (taxa abundance) over the first (CHAR) while positive lags are a measure of the influence
152 of the first variable (CHAR) over the second (taxa abundance). Cross-correlation analyses were completed
153 with the package TSA of R v. 2.0.5 (R Core Team, 2012). There is more related information in
154 Supplementary data.

155 **3. Results and discussion**

156 We interpret the Lateglacial and Holocene fire dynamics in El Portalet (Fig 3) as naturally produced, with
157 lightning as the main ignition source as it happens currently in high altitude areas of the Pyrenees
158 (Amatulli et al., 2007). Neither evidence of human indicators appeared in the pollen assemblages at the
159 studied period, nor archaeological sites are present in the area. Magdalenian, Azilian or Mesolithic
160 settlements at the subalpine belt have not been found. Some trans-Pyrenean migratory routes existed in
161 the area, but they were located in lowland natural passes (Utrilla and Rodanés, 1997; Utrilla and Montes,
162 2007).

163 *3.1 Climate-biomass-fire interactions at the Central Pyrenees: wetter is better*

164 Macro-charcoal analyses resulted in the identification of eighteen FE, most of them (seventeen) occurred
165 between 13.7-12.5 and 9.8-7.7 cal ka BP (Fig. 3B), whereas a single FE appears at ca. 14.6 cal ka BP.
166 Between 7.7 and 6 cal ka BP there was very low to nil macro-charcoal counts and no fire episodes were
167 detected (Fig. 3A). MFI decreased from ca. 314 years during the Lateglacial (14.7-12.5 cal ka BP) to 190
168 years during the Early-Mid Holocene (9.8-7.7 cal ka BP; (Fig. 3A).

169 Unlike the Mediterranean ecosystems, where fire regimes have played an important role shaping the plant
170 landscape before and after human presence (Keeley et al., 2012 and quotes herein), the high altitude
171 plant communities of the Pyrenees are not usually exposed to high wildfire frequencies (Amatulli et al.,
172 2007). Hence, the MFIs found at El Portalet (between ca. 190 to 314 years) (Figs. 3 and 4) points to the
173 low frequency of natural wildfire occurrence, which agrees with present-day fire frequency in the high
174 altitude areas of the Pyrenees, where virtually no natural fire has been recorded from 1974 and 2000

175 (Vázquez de la Cueva et al., 2006). Despite general low frequencies for the period analysed, a clear
176 pattern on wildfire behaviour in El Portalet is detected as MFIs reduced since the Lateglacial to the Early-
177 Mid Holocene. Similarly, lower fire activities during the Lateglacial than during the Early Holocene have
178 been reported in several Northern Hemisphere sites (Carcaillet et al., 2012; Higuera et al., 2011, 2009;
179 Power et al., 2008) and particularly in high-altitude European mountain sites (Feurdean et al., 2012; Tinner
180 and Lotter, 2001; Tinner et al., 2005; Vescovi et al., 2007), where climate was colder and drier during this
181 time compared to the Holocene. Changes in temperature during the Lateglacial did not play a significant
182 role in MFIs variability. The nearest chironomid-derived palaeotemperature curve in the Col d'Ech
183 Northern Pyrenees site (700 m asl, Fig. 1, Lourdes Basin, France) (Millet et al., 2012), shows an important
184 increase in temperature of ca. 2°C at ca. 14.7 cal ka BP, coeval with the Bølling onset. Likewise, the also
185 chironomid-inferred paleotemperature from Laguna de la Roya (NW Spain) (Muñoz Sobrino et al., 2013)
186 indicates an increase of 2.5°C during the Bølling. This thermic rise did not apparently promote large
187 wildfire ignition in El Portalet, with a single FE recorded (Fig. 4). On the one hand, it is possible that the
188 temperature rise during this period would have been gradual rather than abrupt as it has been found for
189 the Mediterranean Sea (Cacho et al., 2001) and continental Iberia (Moreno et al., 2012), where maximum
190 temperature values were identified during the Allerød (Fig. 4). Considering the complex Lateglacial
191 Interstadial palaeoclimate scenario and El Portalet's altitude (1800 m asl), we argue that the gradual
192 increase in temperature during the Bølling, with both cool summers – with still reduced summer solar
193 insolation (Fig. 4) – and extreme winter temperatures, could explain the low fire frequency between 14.6
194 and 13.9 cal ka BP. Afterwards, during the warmer intervals GI-1c (13.9-13.3 cal ka BP) and GI-1a (13.1-
195 12.9 cal ka BP) (Lowe et al., 2008), and as summer temperature progressively increased during the
196 Allerød, fire frequency increased (Fig. 4), supporting the climatically driven character of these fires.
197 Similarly, and somehow coeval with the more frequent warm episodes within the Allerød (Fig. 4), FE in El
198 Portalet area modestly increased ca. 13.5 cal ka BP from 2 to 3 fires per 500 years, and similarly regional
199 fire activity moderately enhanced as indicated by micro-charcoal influx (Fig. 5).

200 Regardless of the climate setting during this period, vegetation in El Portalet between the Bølling (GI-1e)
201 and part of the GI-1c events (ca. 14.6-13.6 cal kyr BP) (Fig 4) was regionally dominated by pines, most
202 likely Scots and mountain pines (*Pinus sylvestris* and *P. uncinata* respectively), while local vegetation was
203 probably composed of less pollen productive types as junipers, most likely *J. communis* ssp. *alpina* and *J.*
204 *communis* ssp. *hemisphaerica*, and patchy grasslands of Poaceae, *Artemisia* and Chenopodiaceae (Fig.
205 5), pine forests would have probably provided enough fuel for regional fires to happen while the local
206 sparse vegetation would have not represented an important and well-connected biomass to feed local
207 fires, despite relatively warmer conditions.

208 The mesophyte forest developed to unprecedented extensions from 13.5 cal ka BP, during the latest
209 Allerød warm phase (González-Sampéris et al., 2006) (Fig. 5). The rising temperatures and still likely high
210 amounts of winter precipitation at a regional scale (Genty et al., 2006; Morellón et al., 2009; Moreno et al.,
211 2010) probably favoured temperate, drought-intolerant communities during this time in El Portalet and this
212 is a well-documented pattern across European sites and in some other Iberian records (Carrión et al.,
213 2004; Fletcher et al., 2010; López-Merino et al., 2012). The location of our site in a south-facing, less
214 steep slopes of the Central Spanish Pyrenees, would have promoted the faster upwards migration of birch
215 forests from its lowland locations when conditions improved. We suggest that since ca. 13.5 cal ka BP, El
216 Portalet held locally dense mesophyte communities formed by dwarf birches (*Betula alba* ssp. *nana*)
217 similar to those currently present in the nearby valleys at the same altitude (Fig. 2). Birch at El Portalet
218 Lateglacial landscapes would have played a pioneer role as it is a well-dispersing tree in poor, oligotrophic
219 soils and its heliophytic character favours its development at early successional stages. Interestingly,
220 intensified fire frequencies from 13.5 cal ka BP concurs with expanding moisture-demanding communities,
221 which may seem a counterintuitive idea as wetter environments would prevent large wildfires. However,
222 these conditions induce a major biomass production (Clark et al., 1989; Daniau et al., 2007; Power et al.,
223 2012) facilitating the expansion of mesophytic communities that grow faster than conifers (Blarquez and
224 Carcaillet, 2010), and creating a remarkable cover of flammable fuel necessary for intense wildfires

225 (Krawchuk and Moritz, 2010). This controlling role of a particular forest composition on local fire could be
226 even more patent during the Early Holocene in El Portalet.

227 Fire frequencies varied from 1 fire per 500 years at 9.8-9 cal ka BP, to 3 fires per 500 years at 9-7.7 cal ka
228 BP (Figs. 3C and 4) and the *Pinus* : mesophytes ratio varying drastically (Fig. 4). Climatically, the Early
229 Holocene in the Mediterranean region was marked by wetter and warmer conditions than in Lateglacial
230 times (Cacho et al., 2002; Roberts et al., 2004), which in El Portalet stimulated the continuous
231 development of mesic forests (González-Sampériz et al., 2006) (Fig. 5). Accordingly, vegetation
232 landscapes in El Portalet became then even more biomass-rich than during the Allerød period, as a
233 denser *Corylus*-dominated shrubland spread synchronous with the rise in temperature, followed by a new
234 expansion of *Betula* ca. 9.3 cal ka BP. Thus, increasing fire frequencies between 9-7.7 cal ka BP would
235 have most likely been driven by the mesophytes expansion (Figs. 4 and 5).

236 Mesophytic biomass control of fire frequency during the Holocene is supported by cross-correlation results
237 (Fig. 6), where birch and other mesophytes, as well as total tree percentages, have a positive effect on
238 local wildfire at different lags since 150 years before fire occurrence. Bushes and herbs are,
239 unsurprisingly, negatively affecting fire occurrence as they prevent large wildfires in a less connected
240 landscape. Regarding trees, the major effect of mesic communities on local fire responds to the biomass
241 accumulated from 120 to 30 years prior to a fire event (mesophyte cross-correlation, Fig. 6). Our results
242 are coherent with a recent study by Pausas and Paula (2012) on the present role of plant productivity,
243 community structure and climate-fire relationships on fire dynamics in the Iberian Peninsula. These
244 authors showed that differences in fire activity across an aridity gradient are not linked to the frequency of
245 fire-prone climate conditions but by the vegetation sensitivity to fire under those conditions, which resulted
246 higher in wetter and more productive regions. Conversely, herbs and shrubs are negatively affecting fire
247 ignition in El Portalet (Fig. 6), reinforcing the hypothesis of more likely wildfire incidence at dense,
248 encroached temperate forests. Indeed, pine presence does not seem to positively influence fire
249 occurrence, reflecting either the less flammable character of *Pinus uncinata*/*P. sylvestris* stands than
250 *Betula* (Fernandes et al., 2008; Tapias et al., 2004), or a forest structure with relatively thin pine

251 woodlands and young stands with no significant ageing processes, both preventing large, more frequent
252 fires.

253 The Early to Mid-Holocene transition reflects a sharp decline in local fire activity and after 7.7 cal ka BP no
254 fire is recorded. We postulate that relatively humid conditions could have partially prevented fire
255 occurrence, while the gradual contraction of tree cover probably yielded less effective fuel to be burnt
256 precluding large local wildfires. Summer insolation decreased during the Early to Mid-Holocene transition,
257 favouring relatively cooler summers and milder winters (Webb III et al., 1993). Regional indicators of
258 moisture availability are less clear, and while some Northern Mediterranean and Northern European
259 records showed drier conditions (Magny et al., 2003; Roberts et al., 2004), nearer sequences indicate a
260 regional increase in water availability (Morellón et al., 2008; Pérez-Sanz et al., 2013; Aranbarri et al.,
261 2014).

262 The Holocene decrease of local FE does not find a counterpart in El Portalet regional fire activity, as
263 micro-charcoal is still recorded along the sequence (Fig. 5), proving that charcoal sources might not be
264 associated. The vegetation-regional fire relationships should be cautiously taken though, as pollen and
265 micro-charcoal may have different source areas difficult to identify. Some experiments trying to overcome
266 this issue proved that charcoal particles <50 µm are likely deposited from 10 to 15 km away from a 200 m
267 diameter lake (Duffin et al., 2008). The original surface area of El Portalet lake is unknown but the current
268 peatland diameter is over 200 m, and the whole basin is not much larger (15 ha), and micro-charcoal
269 source area most likely exceeds 10 km, implying fire happening in the lowlands, where active burning
270 might still have occurred. Relevant source area for pollen, as well as its productivity, are unidentified;
271 however, the immediate response of some taxa (*Pinus*, *Juniperus*, and *Betula*) to local fire occurrence
272 indicates a local to sub-regional pollen input (<10 km) with some regional contribution, particularly from
273 those highly productive and well-dispersed taxa as *Pinus*.

274 Considering that *Betula* and *Corylus* pollen proportions decreased to less than 15% after 7.7 cal ka BP,
275 we suggest that both taxa locally disappeared and only remained in patched formations at closer valleys
276 and at lower elevations, as they occur nowadays (Fig. 2). These small patches would be located farther

277 away than the 10 km charcoal source area limit and, consequently, they would not contribute to the
278 macro-charcoal input to El Portalet site. Most importantly, this proves that the local sensitivity to fire-
279 climate relationship is mostly mediated by flammable biomass and not only by climate (Pausas and Paula,
280 2012), as this site would have received a relatively high rainfall amount during the Holocene.

281 These findings confirm the trends found in other Holocene European records, where fire intensity and
282 frequency are strengthened since the beginning of the Holocene until the Early-Mid Holocene transition
283 (Colombaroli et al., 2007; Feurdean et al., 2012; Feurdean et al., 2013; Tinner et al., 1999; Vescovi et al.,
284 2007). However, in the closest Holocene Col d'Ech fire record (Lourdes basin, France, ca. 700 m asl; Rius
285 et al., 2011a, 2011b), a different fire regime pattern is found (Fig 3F). Despite little temporal overlap with El
286 Portalet, fire activity intensification in the Lourdes basin from 7.8 to 4 cal ka BP is discussed by the authors
287 considering three features: a change in the sedimentation rate (from silt to peat ca. 8.2 cal ka BP); an
288 increasing effect of human activity; and progressively warmer summers and flammable-prone conditions
289 due to a lagged effect on temperature of Laurentide ice sheet deglaciation (Renssen et al., 2009). El
290 Portalet presents a change in sedimentation rate much later than Col d'Ech (ca. 6.5 cal ka BP), thus, in
291 our study the lack of fire activity from 7.7 cal ka BP would not be associated with depositional factors. As
292 discussed before, anthropogenically-induced environmental changes are less likely at higher altitudes and,
293 unlike at the Col d'Ech, no evidence of human activity has been documented in the area during the Early
294 Holocene. Mid-Holocene warmer summers (Holocene Thermal Maximum) have been modelled at the
295 Northern hemisphere spatial scale, suggesting that cooler summers than expected due to summer
296 insolation before 7 cal ka BP arose from a combination of the inhibition of Labrador Sea deep convection
297 by the flux of melt water from the ice sheet, which weakened northward heat transport by the ocean, and
298 the high surface albedo of the ice sheet (Renssen et al., 2009). However, our findings in El Portalet may
299 not fit in this model as the occurrence of a high local fire activity from 9 to 7 cal ka BP in a relatively fuel-
300 limited environment as the Central Spanish Pyrenees treeline, supports the importance of orbital forcing
301 on the mesophyte development (Fig. 5) and the role of the latter in modulating fire activity (Daniau et al.,
302 2007; Pausas and Paula, 2012; Power et al., 2011).

303 3.2 Vegetation–fire interactions, plant responses to disturbance and ecosystem resilience

304 The current plant communities at the mid to high altitudes of the Pyrenees do not show any fire trait
305 implying a clear adaptation to fire (Fulé et al., 2008; Montané et al., 2009; Paula et al., 2009). Modern fire
306 frequencies are relatively higher in the Central Pyrenees compared to the MFI found in El Portalet (1 to 5
307 FE, both wildfire and attempted arsons, for the last 10 years, Spanish Forestry Database (MAGRAMA,
308 2012)). Hence, it should be considered that FE identified in El Portalet sequence responds to large
309 wildfires, probably implying both crown and surface fires, while current statistics do not distinguish these
310 criteria (size or fire type). It is not surprising then that local wildfires have a long-term negative effect at
311 different time-lags on most studied tree taxa in El Portalet and that taxa favoured by fire are those forming
312 open-landscape communities, as herbs and shrubs, which are also those preventing fire occurrence (Fig.
313 6). An exception to this pattern is *Pinus*, most likely represented by *Pinus sylvestris* / *P. uncinata* in El
314 Portalet, which are not fire-adapted pines, unlike the Mediterranean pines (*Pinus pinaster*, *P. pinea*, *P.*
315 *nigra* and *P. halepensis*) that are showing serotiny, bark thickness and bud resistance to fire (Fernandes
316 et al., 2008; Paula et al., 2009). Despite some previous studies in the Alps having proved a positive
317 relationship between fire and *Pinus sylvestris* (Stähli et al., 2006), we suggest that in El Portalet the *Pinus*-
318 fire correlation might be associated with temporal exclusion of *Pinus* competitors, rather than to a direct
319 positive effect of fire over pine occurrence, enabling pine recolonization from the edge of burnt areas or
320 from sheltered refugia in rocky outcrops (Keeley et al., 2012).

321 The long-term role of birch in the sub-alpine/alpine belt of the Central Spanish Pyrenees is partially in
322 agreement with its documented early-successional character after disturbances, as revealed by other
323 Holocene sequences (Feurdean and Astalos, 2005; Morales-Molino et al., 2011; Pérez-Sanz et al., 2013)
324 and modern ecology studies on their resprouting ability (Reyes and Casal, 1998). Interestingly, *Betula* in
325 El Portalet is a main factor building-up fuel but its development is negatively affected by fire in the first 150
326 years after the fire event. This is common with most of the mountain trees, supporting the idea of the
327 alpine ecosystem as a long-term fire-sensitive environment. Indeed it is worth considering that more
328 recurrent fires during the Early-Mid Holocene transition would have had a long-term impact on the birch

329 development as it is the only mesophyte that clearly decreases from 7.5 cal kyr BP. However, this does
330 not necessarily preclude birch communities pioneering the treeline recuperation as one amongst the first
331 forestalls types able to spread after fires, especially considering that most trees respond negatively to fire
332 (Fig. 6).

333 Palynological richness (rarefaction index) gradually increased as large wildfires reduced from 7.7 cal ka
334 BP (Fig. 5), and despite its long-term variation, richness does not show any clear pattern linked to local fire
335 occurrence (Fig. 6). A long-term positive fire effect on plant richness has often been argued, especially in
336 those ecosystems where plants present fire-resistant seedbanks (Beckage and Stout, 2000; Johst and
337 Huth, 2005; Montané et al., 2009). However, we argue that El Portalet increasing palynological richness
338 during the Holocene is not led by fire disturbance, but it seems to be a consequence of the tree cover
339 decline and the spread of non-arboreal communities, as semi-open landscapes represent a main driver of
340 palynological richness and show larger diversity than closed forests (Colombaroli and Tinner, 2013;
341 Feurdean and Astalos, 2005). We actually found that the non-arboreal pollen abundance presents a
342 Spearman's $R = 0.75$ ($p < 0.05$) (Table 2) with the rarefaction values for the 10-6 cal ka BP period (Fig. 5),
343 supporting the idea of richer ecosystem in more open landscapes but not necessarily linked to fire
344 disturbance.

345 **4. Conclusions**

346 The high resolution pollen and charcoal records at El Portalet enabled long-term analyses on fire regime,
347 biomass-mediated fire occurrence and vegetation response to disturbance at the Central Spanish
348 Pyrenees treeline. The main findings of our study are:

- 349 1) Local wildfire frequency varied since the Lateglacial (14.7 cal ka BP) to the Mid-Holocene (6 cal ka
350 BP) at El Portalet, where the mean fire intervals progressively decreased during the fire-active
351 periods, between 13.6-12.5 and 9.8-7.7 cal ka BP.
- 352 2) Variation in fire activity was not linked to human presence in El Portalet but to climate and biomass
353 fluctuations. Mesophyte forest expansion (led by *Betula* and *Corylus* communities), as a fuel-creating

354 factor, triggered increasing fire frequencies in the Central Spanish Pyrenees treeline. Hence, fire
355 regimes in this region proved to be long-term sensitive to insolation changes and biomass
356 abundance.

357 3) Fire has long-lasting negative effects on most of the treeline plant communities. These are not
358 currently fire-prone forests, exhibiting no clear fire-trait, and El Portalet sequence shows that this is a
359 long-term ecological feature of the subalpine/alpine Pyrenees communities.

360 4) Forest contraction from 7.7 cal ka BP could be linked to a threshold response of the system to
361 recurrent fire frequencies as climate was not unfavourable for forest development. Palynological
362 richness increased from that time coeval with the openness of the landscape.

363 Long-term fire ecology is proved to be a useful tool to explore Quaternary fire regime changes, especially
364 in those fire-sensitive ecosystems where changing fire regimes may determine vegetation composition
365 and resilience.

366 Besides, our findings are relevant under the current and predicted scenarios of global change, as warming
367 would stimulate more frequent fire events in biomass-rich environments and therefore influencing
368 environmental resilience vulnerability levels. Interestingly, these results from Quaternary scenarios concur
369 with the local knowledge on current forest management, as both would stress the importance of keeping
370 low levels of woody encroachment and promoting traditional forest use in order to prevent fuel
371 accumulation and avoid virulent fires, with catastrophic effects both in this fire-sensitive ecosystem and in
372 the local communities.

373 **Acknowledgements**

374 This research has been funded by the projects DINAMO (CGL2009-07992) (funding EGPF – grant ref.
375 BES-2010-038593 and MSC), DINAMO2 (CGL2012-33063), ARAFIRE (2012 GA LC 064), GRACCIE-
376 CONSOLIDER (CSD2007-00067). GGR was funded by the Juan de la Cierva Program (grant ref.
377 JCI2009-04345) and JAE-Doc CSIC Program, LLM was supported by a postdoctoral MINT fellowship

378 funded by the Institute for the Environment (Brunel University), AMC is a Ramón y Cajal fellow (ref: RYC-
379 2008-02431), APS holds a grant funded by the Aragon Government (ref. 17030 G/5423/480072/14003)
380 and JAE holds a grant funded by the Basque Country Government (BFI-2010-5). We are indebted to
381 Beatriz Bueno and Aida Adsuar for sampling processing and laboratory analyses and to César Morales
382 del Molino for productive discussions on the ideas of this paper. We are grateful to Steven M. Holland for
383 providing the free software to calculate rarefaction (Analytic Rarefaction), which is available at
384 <http://strata.uga.edu/software/>.

385 **References**

- 386 AEMET-IM (2011). Atlas Climático Ibérico (Agencia Estatal de Meteorología Ministerio de Medio Ambiente y Medio
387 Rural y Marino).
- 388 Amatulli, G., Pérez-Cabello, F., and de la Riva, J. (2007). Mapping lightning/human-caused wildfires occurrence
389 under ignition point location uncertainty. *Ecological Modelling* 200, 321–333.
- 390 Aranbarri, J., González-Sampériz, P., Valero-Garcés, B., Moreno, A., Gil-Romera, G., Sevilla-Callejo, M., García-
391 Prieto, E., Di Rita, F., Mata, M.P., Morellón, M., Magri, D., Rodríguez-Lázaro, J., Carrión, J.S., 2014. Rapid climatic
392 changes and resilient vegetation during the Lateglacial and Holocene in a continental region of south-western
393 Europe. *Global and Planetary Change* 114, 50–65.
- 394 Bal, M.-C., Pelachs, A., Perez-Obiol, R., Julia, R., and Cunill, R. (2011). Fire history and human activities during the
395 last 3300 cal yr BP in Spain's Central Pyrenees: The case of the Estany de Burg. *Palaeogeography,*
396 *Palaeoclimatology, Palaeoecology* 300, 179–190.
- 397 Beckage, B., and Stout, I.J. (2000). Effects of repeated burning on species richness in a Florida pine savanna: a test
398 of the intermediate disturbance hypothesis. *Journal of Vegetation Science* 11, 113–122.
- 399 Bennett, K. (2009). Documentation for Psimpoll 4.27 and Pscomb 1.03: C Programs for Plotting Pollen Diagrams and
400 Analysing Pollen Data (Queen's University of Belfast, Department of Archaeology and Palaeoecology).
- 401 Blarquez, O., and Carcaillet, C. (2010). Fire, Fuel Composition and Resilience Threshold in Subalpine Ecosystem.
402 *PLoS ONE* 5, e12480.
- 403 Bond, W.J., and Keeley, J.E. (2005). Fire as a global “herbivore”: the ecology and evolution of flammable
404 ecosystems. *Trends in Ecology and Evolution* 20, 387–394.
- 405 Cacho, I., Grimalt, J.O., Canals, M., Saffi, L., Shackleton, N.J., Schönfeld, J., and Zahn, R. (2001). Variability of the
406 western Mediterranean Sea surface temperature during the last 25,000 years and its connection with the Northern
407 Hemisphere climatic changes. *Paleoceanography* 16, 40–52.
- 408 Cacho, I., Grimalt, J., and Canals, M. (2002). Response of the Western Mediterranean Sea to rapid climatic
409 variability during the last 50,000 years: a molecular biomarker approach. *Journal of Marine Systems* 34, 253–272.

- 410 Carcaillet, C., Bergeron, Y., Richard, P.J.H., Frechette, B., Gauthier, S., and Prairie, Y.T. (2001). Change of fire
411 frequency in the eastern Canadian boreal forests during the Holocene: does vegetation composition or climate trigger
412 the fire regime? *Journal of Ecology* 89, 930–946.
- 413 Carcaillet, C., Hörnberg, G., and Zackrisson, O. (2012). Woody vegetation, fuel and fire track the melting of the
414 Scandinavian ice-sheet before 9500 cal yr BP. *Quaternary Research* 540–548.
- 415 Carrión, J.S. (2002). Patterns and processes of Late Quaternary environmental change in a montane region of
416 southwestern Europe. *Quaternary Science Reviews* 21, 2047–2066.
- 417 Carrión, J.S., Yll, E.I., Willis, K., and Sanchez, P. (2004). Holocene forest history of the eastern plateaux in the
418 Segura Mountains (Murcia, Southern Spain). *Review of Palaeobotany and Palynology* 132, 219–236.
- 419 Carrión, J.S., Fuentes, N., González-Sampériz, P., Sanchez Quirante, L., Finlayson, J.C., Fernández, S., and
420 Andrade, A. (2007). Holocene environmental change in a montane region of southern Europe with a long history of
421 human settlement. *Quaternary Science Reviews* 26, 1455–1475.
- 422 Carrión, J.S., Fernández, S., Jiménez-Moreno, G., Fauquette, S., Gil-Romera, G., González-Sampériz, P., and
423 Finlayson, C. (2010a). The historical origins of aridity and vegetation degradation in southeastern Spain. *Journal of*
424 *Arid Environments* 74, 731–736.
- 425 Carrión, J.S., Fernández, S., González-Sampériz, P., Gil-Romera, G., Badal, E., Carrión-Marco, Y., López-Merino,
426 L., López-Sáez, J.A., Fierro, E., and Burjachs, F. (2010b). Expected trends and surprises in the Lateglacial and
427 Holocene vegetation history of the Iberian Peninsula and Balearic Islands. *Review of Palaeobotany and Palynology*
428 162, 458–475.
- 429 Carrión-Marco, Y. (2005). *La vegetación mediterránea y atlántica de la península Ibérica. Nuevas secuencias*
430 *antracológicas*. (Valencia: Servicio de Investigación prehistórica. Diputación provincial de Valencia).
- 431 Carrión-Marco, Y., Kaal, J., López-Sáez, J.A., López-Merino, L., and Cortizas, A.M. (2010). Holocene vegetation
432 changes in NW Iberia revealed by anthracological and palynological records from a colluvial soil. *The Holocene* 20,
433 53–66.
- 434 Castroviejo, S. (Coord) (1986). *Flora Iberica* (Madrid: Real Jardín Botánico. CSIC).
- 435 Clark, J.S., and Royall, P.D. (1996). Local and regional sediment charcoal evidence for fire regimes in presettlement
436 north-eastern North America. *Journal of Ecology* 84, 365–382.
- 437 Clark, J.S., Merkt, J., and Muller, H. (1989). Post-Glacial Fire, Vegetation, and Human History on the Northern Alpine
438 Forelands, South-Western Germany. *Journal of Ecology* 77, 897–925.
- 439 Clark, J.S., Royall, P.D., and Chumbley, C. (1996). The Role of Fire During Climate Change in an Eastern Deciduous
440 Forest at Devil's Bathtub, New York. *Ecology* 77, 2148–2166.
- 441 Colombaroli, D., and Tinner, W. (2013). Determining the long-term changes in biodiversity and provisioning services
442 along a transect from Central Europe to the Mediterranean. *The Holocene* 23, 1625–1634.
- 443 Colombaroli, D., Marchetto, A., and Tinner, W. (2007). Long-term interactions between Mediterranean climate,
444 vegetation and fire regime at Lago di Massaciuccoli (Tuscany, Italy). *Journal of Ecology* 95, 755–770.
- 445 Colombaroli, D., Vanniere, B., Emmanuel, C., Magny, M., and Tinner, W. (2008). Fire-vegetation interactions during
446 the Mesolithic–Neolithic transition at Lago dell'Accesa, Tuscany, Italy. *The Holocene* 18, 679–692.

- 447 Colombaroli, D., Tinner, W., van Leeuwen, J., Noti, R., Vescovi, E., Vanni re, B., Magny, M., Schmidt, R., and
448 Bugmann, H. (2009). Response of broadleaved evergreen Mediterranean forest vegetation to fire disturbance during
449 the Holocene: insights from the peri-Adriatic region. *Journal of Biogeography* 36, 314–326.
- 450 Colombaroli, D., Henne, P.D., Kaltenrieder, P., Gobet, E., and Tinner, W. (2010). Species responses to fire, climate
451 and human impact at tree line in the Alps as evidenced by palaeo-environmental records and a dynamic simulation
452 model. *Journal of Ecology* 98, 1346–1357.
- 453 Daniau, A.-L., S nchez-Go ni, M.F., Beaufort, L., Laggoun-D farge, F., Loutre, M.-F., and Duprat, J. (2007).
454 Dansgaard-Oeschger climatic variability revealed by fire emissions in southwestern Iberia. *Quaternary Science*
455 *Reviews* 26, 1369–1383.
- 456 Dearing, J.A. (2008). Landscape change and resilience theory: a palaeoenvironmental assessment from Yunnan,
457 SW China. *The Holocene* 18, 117–127.
- 458 Duffin, K.I., Gillson, L., and Willis, K.J. (2008). Testing the Sensitivity of Charcoal as an Indicator of Fire Events in
459 Savanna Environments: Quantitative Predictions of Fire Proximity, Area and Intensity. *The Holocene* 18, 279–291.
- 460 Engler, R., Randin, C.F., Thuiller, W., Dullinger, S., Zimmermann, N.E., Ara jo, M.B., Pearman, P.B., Le Lay, G.,
461 Piedallu, C., Albert, C.H., et al. (2011). 21st century climate change threatens mountain flora unequally across
462 Europe. *Global Change Biology* 17, 2330–2341.
- 463 Fernandes, P.M., Vega, J.A., Jimenez, E., and Rigolot, E. (2008). Fire resistance of European pines. *Forest Ecology*
464 *and Management* 256, 246–255.
- 465 Feurdean, A., and Astalos, C. (2005). The impact of human activities in the Gutaiului Mountains, Romania. *Studia*
466 *Universitatis Babe -Bolyai, Geologia* 50, 63–72.
- 467 Feurdean, A., Spessa, A., Magyari, E.K., Willis, K.J., Veres, D., and Hickler, T. (2012). Trends in biomass burning in
468 the Carpathian region over the last 15,000 years. *Quaternary Science Reviews* 45, 111–125.
- 469 Feurdean, A., Liakka, J., Vanni re, B., Marinova, E., Hutchinson, S.M., Mosburgger, V., Hickler, T., 2013. 12,000-
470 Years of fire regime drivers in the lowlands of Transylvania (Central-Eastern Europe): a data-model approach.
471 *Quaternary Science Reviews* 81, 48–61.
- 472 Fletcher, W.J., Sanchez Go ni, M.F., Peyron, O., and Dormoy, I. (2010). Abrupt climate changes of the last
473 deglaciation detected in a Western Mediterranean forest record. *Clim. Past* 6, 245–264.
- 474 Franco M gica, F., Garc a Ant n, M., Maldonado Ruiz, J., Morla Juaristi, C., and Sainz Ollero, H. (2005). Ancient
475 pine forest on inland dunes in the Spanish northern meseta. *Quaternary Research* 63, 1–14.
- 476 Ful , P.Z., Ribas, M., Guti rrez, E., Vallejo, R., and Kaye, M.W. (2008). Forest structure and fire history in an old
477 *Pinus nigra* forest, eastern Spain. *Forest Ecology and Management* 255, 1234–1242.
- 478 Garc a-Ruiz, J.M., Valero-Garc s, B.L., Mart -Bono, C., and Gonz lez-Samp riz, P. (2003). Asynchronicity of
479 maximum glacier advances in the central Spanish Pyrenees. *Journal of Quaternary Science* 18, 61–72.
- 480 Gavin, D.G., Hu, F.S., Lertzman, K., and Corbett, P. (2006). Weak climatic control of stand-scale fire history during
481 the Late Holocene. *Ecology* 87, 1722–1732.
- 482 Genty, D., Blamart, D., Ghaleb, B., Plagnes, V., Causse, C., Bakalowicz, M., Zouari, K., Chkir, N., Hellstrom, J.,
483 Wainer, K., et al. (2006). Timing and dynamics of the last deglaciation from European and North African $\delta^{13}C$
484 stalagmite profiles—comparison with Chinese and South Hemisphere stalagmites. *Quaternary Science Reviews* 25,
485 2118–2142.

- 486 Gil-Romera, G., Carrión, J.S., Pausas, J.G., Sevilla-Callejo, M., Lamb, H.F., Fernández, S., and Burjachs, F. (2010).
487 Holocene fire activity and vegetation response in South-Eastern Iberia. *Quaternary Science Reviews* 29, 1082–1092.
- 488 Gobet, E., Tinner, W., Hochuli, P.A., van Leeuwen, J.F.N., and Ammann, B. (2003). Middle to Late Holocene
489 vegetation history of the Upper Engadine (Swiss Alps): the role of man and fire. *Vegetation History and*
490 *Archaeobotany* 12, 143–163.
- 491 González-Sampériz, P., Valero-Garcés, B.L., Moreno, A., Jalut, G., García-Ruiz, J.M., Martí-Bono, C., Delgado-
492 Huertas, A., Navas, A., Otto, T., and Dedoubat, J.J. (2006). Climate variability in the Spanish Pyrenees during the
493 last 30,000 yr revealed by the El Portalet sequence. *Quaternary Research* 66, 38–52.
- 494 Gottfried, M., Pauli, H., Futschik, A., Akhalkatsi, M., Barančok, P., Alonso, J.L.B., Coldea, G., Dick, J., Erschbamer,
495 B., Calzado, M.R.F., et al. (2012). Continent-wide response of mountain vegetation to climate change. *Nature*
496 *Climate Change* 2, 111–115.
- 497 Higuera, P., Chipman, M., Barnes, J., Urban, M., and Hu, F.S. (2011). Variability of tundra fire regimes in Arctic
498 Alaska: millennial-scale patterns and ecological implications. *Ecological Applications*.
- 499 Higuera, P.E., Brubaker, L.B., Anderson, P.M., Hu, F.S., and Brown, T.A. (2009). Vegetation mediated the impacts of
500 postglacial climate change on fire regimes in the south-central Brooks Range, Alaska. *Ecological Monographs* 79,
501 201–219.
- 502 Higuera, P.E., Gavin, D.G., Bartlein, P.J., and Hallett, D.J. (2010). Peak detection in sediment–charcoal records:
503 impacts of alternative data analysis methods on fire-history interpretations. *Int. J. Wildland Fire* 19, 996–1014.
- 504 Jiménez-Moreno, G., García-Alix, A., Hernández-Corbalán, M.D., Anderson, R.S., and Delgado-Huertas, A. (2013).
505 Vegetation, fire, climate and human disturbance history in the southwestern Mediterranean area during the late
506 Holocene. *Quaternary Research*.
- 507 Johst, K., and Huth, A. (2005). Testing the intermediate disturbance hypothesis: when will there be two peaks of
508 diversity? *Diversity and Distributions* 11, 111–120.
- 509 Kaal, J., Carrión Marco, Y., Asouti, E., Martín Seijo, M., Martínez Cortizas, A., Costa Casáis, M., and Criado Boado,
510 F. (2011). Long-term deforestation in NW Spain: linking the Holocene fire history to vegetation change and human
511 activities. *Quaternary Science Reviews* 30, 161–175.
- 512 Keeley, J.E., Bond, W.J., Bradstock, R.A., Pausas, J.G., and Rundel, P. (2012). *Fire in Mediterranean Ecosystems:*
513 *Ecology, Evolution and Management* (Cambridge University Press).
- 514 Krawchuk, M.A., Moritz, M.A., 2010. Constraints on global fire activity vary across a resource gradient. *Ecology* 92,
515 121–132.
- 516 Laskar, J. R. (2004). A long-term numerical solution for the insolation quantities of the Earth. *Astronomy and*
517 *Astrophysics* 428, 261–285.
- 518 López-Merino, L., López-Sáez, J.A., Alba-Sánchez, F., Pérez-Díaz, S., and Carrión, J.S. (2009). 2000 years of
519 pastoralism and fire shaping high-altitude vegetation of Sierra de Gredos in central Spain. *Review of Palaeobotany*
520 *and Palynology* 158, 42–51.
- 521 López-Merino, L., Silva Sánchez, N., Kaal, J., López-Sáez, J.A., and Martínez Cortizas, A. (2012). Post-disturbance
522 vegetation dynamics during the Late Pleistocene and the Holocene: An example from NW Iberia. *Global and*
523 *Planetary Change* 92–93, 58–70.

- 524 Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J.C., and Yu, Z.C. (2008).
525 Synchronisation of palaeoenvironmental events in the North Atlantic region during the Last Termination: a revised
526 protocol recommended by the INTIMATE group. *Quaternary Science Reviews* 27, 6–17.
- 527 Lynch, J.A., Clark, J.S., Bigelow, N.H., Edwards, M.E., and Finney, B.P. (2002). Geographic and temporal variations
528 in fire history in boreal ecosystems of Alaska. *Journal of Geophysical Research-Atmospheres* 108.
- 529 Magny, M., Bégeot, C., Guiot, J., and Peyron, O. (2003). Contrasting patterns of hydrological changes in Europe in
530 response to Holocene climate cooling phases. *Quaternary Science Reviews* 22, 1589–1596.
- 531 MAGRAMA (2012). Estadísticas de Incendios forestales (Ministerio de Agricultura, Alimentación y Medio Ambiente).
- 532 Médail, F., and Quézel, P. (1999). Biodiversity Hotspots in the Mediterranean Basin: Setting Global Conservation
533 Priorities. *Conservation Biology* 13, 1510–1513.
- 534 Millet, L., Rius, D., Galop, D., Heiri, O., and Brooks, S.J. (2012). Chironomid-based reconstruction of Lateglacial
535 summer temperatures from the Ech palaeolake record (French western Pyrenees). *Palaeogeography,*
536 *Palaeoclimatology, Palaeoecology* 315-316, 86–99.
- 537 Montané, F., Casals, P., Taull, M., Lambert, B., and Dale, M.R.T. (2009). Spatial patterns of shrub cover after
538 different fire disturbances in the Pyrenees. *Annals of Forest Science* 66, 612–612.
- 539 Morales-Molino, C., Postigo-Mijarra, J.M., García-Antón, M., and Zazo, C. (2011). Vegetation and environmental
540 conditions in the Doñana Natural Park coastal area (SW Iberia) at the beginning of the last glacial cycle. *Quaternary*
541 *Research* 75, 205–212.
- 542 Morales-Molino, C., García-Antón, M., Postigo-Mijarra, J.M., and Morla, C. (2013). Holocene vegetation, fire and
543 climate interactions on the westernmost fringe of the Mediterranean Basin. *Quaternary Science Reviews* 59, 5–17.
- 544 Morellón, M., Valero-Garcés, B., Moreno, A., González-Sampériz, P., Mata, P., Romero, O., Maestro, M., and Navas,
545 A. (2008). Holocene palaeohydrology and climate variability in northeastern Spain: The sedimentary record of Lake
546 Estanya (Pre-Pyrenean range). *Quaternary International* 181, 15–31.
- 547 Morellón, M., Valero-Garcés, B., Vegas-Vilarrúbia, T., González-Sampériz, P., Romero, Ó., Delgado-Huertas, A.,
548 Mata, P., Moreno, A., Rico, M., and Corella, J.P. (2009). Lateglacial and Holocene palaeohydrology in the western
549 Mediterranean region: The Lake Estanya record (NE Spain). *Quaternary Science Reviews* 28, 2582–2599.
- 550 Moreno, A., Stoll, H.M., Jiménez-Sánchez, M., Cacho, I., Valero-Garcés, B., Ito, E., and Edwards, L.R. (2010). A
551 speleothem record of rapid climatic shifts during last glacial period from Northern Iberian Peninsula. *Global and*
552 *Planetary Change* 71, 218–231; doi:10.1016/j.gloplacha.2009.10.002.
- 553 Moreno, A., González-Sampériz, P., Morellón, M., Valero-Garcés, B.L., and Fletcher, W.J. (2012). Northern Iberian
554 abrupt climate change dynamics during the last glacial cycle: A view from lacustrine sediments. *Quaternary Science*
555 *Reviews*.
- 556 Muñoz Sobrino, C., Heiri, O., Hazekamp, M., van der Velden, D., Kirilova, E.P., García-Moreiras, I., and Lotter, A.F.
557 (2013). New data on the Lateglacial period of SW Europe: a high resolution multiproxy record from Laguna de la
558 Roya (NW Iberia). *Quaternary Science Reviews* 80, 58–77.
- 559 Paula, S., Arianoutsou, M., Kazanis, D., Tavsanoğlu, Ç., Lloret, F., Buhk, C., Ojeda, F., Luna, B., Moreno, J.M.,
560 Rodrigo, A., et al. (2009). Fire-related traits for plant species of the Mediterranean Basin. *Ecology* 90, 1420.
- 561 Pausas, J.G., and Paula, S. (2012). Fuel shapes the fire–climate relationship: evidence from Mediterranean
562 ecosystems. *Global Ecology and Biogeography* 21, 1074–1082.

- 563 Pérez-Sanz, A., González-Sampériz, P., Moreno, A., Valero-Garcés, B., Gil-Romera, G., Rieradevall, M., Tarrats, P.,
564 Lasheras-Álvarez, L., Morellón, M., Belmonte, A., et al. (2013). Holocene climate variability, vegetation dynamics and
565 fire regime in the central Pyrenees: the Basa de la Mora sequence (NE Spain). *Quaternary Science Reviews* 73,
566 149–169.
- 567 Power, M.J., Marlon, J., Ortiz, N., Bartlein, P.J., Harrison, S.P., Mayle, F.E., Ballouche, A., Bradshaw, R.H.W.,
568 Carcaillet, C., Cordova, C., et al. (2008). Changes in fire regimes since the last glacial maximum: An assessment
569 based on a global synthesis and analysis of charcoal data. *Climate Dynamics* 30, 887–907.
- 570 Power M.J., Whitlock C., Bartlein P.J. (2011) Postglacial fire, vegetation, and climate history across an elevational
571 gradient in the Northern Rocky Mountains, USA and Canada. *Quaternary Science Reviews* 30, 2520–2533.
- 572 Power M.J., Mayle F.E., Bartlein P.J., Marlon J.R., Anderson R.S., Behling H., Brown K.J., Carcaillet C., Colombaroli
573 D., Gavin D.G., Hallett D.J., Horn S.P., Kennedy L.M., Lane C.S., Long C.J., Moreno P.I., Paitre C., Robinson G.,
574 Taylor Z., Walsh M.K. (2013) Climatic control of the biomass-burning decline in the Americas after ad 1500.
575 *Holocene* 23, 3–13.
- 576 R Core Team (2012). *R: A Language and Environment for Statistical Computing* (Vienna, Austria: R Foundation for
577 Statistical Computing).
- 578 Renssen, H., Seppä, H., Heiri, O., Roche, D.M., Goosse, H., and Fichet, T. (2009). The spatial and temporal
579 complexity of the Holocene thermal maximum. *Nature Geoscience* 2, 411–414.
- 580 Reyes, O., and Casal, M. (1998). Are the dominant species in N.W. Spain fire-prone?. In *Fire Management and*
581 *Landscape Ecology*, (Fairfield, Washington), pp. 177–188.
- 582 Rius, D., Vannière, B., and Galop, D. (2009). Fire frequency and landscape management in the northwestern
583 Pyrenean piedmont, France, since the early Neolithic (8000 cal. BP). *The Holocene* 19, 847 –859.
- 584 Rius, D., Vannière, B., and Galop, D. (2011a). Holocene history of fire, vegetation and land use from the central
585 Pyrenees (France). *Quaternary Research* 77, 54–64.
- 586 Rius, D., Vannière, B., Galop, D., and Richard, H. (2011b). Holocene fire regime changes from multiple-site
587 sedimentary charcoal analyses in the Lourdes basin (Pyrenees, France). *Quaternary Science Reviews* 30, 1696–
588 1709.
- 589 Roberts, N., Stevenson, T., Davis, B., Cheddadi, R., Brewster, S., and Rosen, A. (2004). Holocene climate,
590 environment and cultural change in the circum-Mediterranean region. In *Past Climate Variability through Europe and*
591 *Africa*, R.W. Battarbee, F. Gasse, and C.E. Stickley, eds. (Springer Netherlands), pp. 343–362.
- 592 Stähli, M., Finsinger, W., Tinner, W., and Allgöwer, B. (2006). Wildfire history and fire ecology of the Swiss National
593 Park (Central Alps): new evidence from charcoal, pollen and plant macrofossils. *The Holocene* 16, 805–817.
- 594 Tapias, R., Climent, J., Pardos, J.A., and Gil, L. (2004). Life histories of Mediterranean pines. *Plant Ecology* 171, 53–
595 68.
- 596 Tinner, W., and Lotter, A.F. (2001). Central European vegetation response to abrupt climate change at 8.2 ka.
597 *Geology* 29, 551 –554.
- 598 Tinner, W., Hubschmid, P., Wehrli, M., Ammann, B., and Conedera, M. (1999). Long-Term Forest Fire Ecology and
599 Dynamics in Southern Switzerland. *Journal of Ecology* 87, 273–289.
- 600 Tinner, W., Conedera, M., Ammann, B., and Lotter, A. (2005). Fire ecology north and south of the Alps since the last
601 ice age. *The Holocene* 15, 1214–1226.

- 602 Utrilla, P., and Montes, L. (2007). El Paleolítico superior al sur de los Pirineos. Contactos entre fronteras. In
603 *Frontières Naturelles et Frontières Culturelles Dans Les Pyrénées Préhistoriques*, N. Cazals, J.E. González Urquijo,
604 and X. Terradas, eds. (Tarascon sur Ariège), pp. 205–223.
- 605 Utrilla, P., and Rodanés, J.M. (1997). La actuación del hombre sobre el paisaje durante la Prehistoria en el Valle
606 Medio del Ebro. In *Acción Humana Y Desertificación En Ambientes Mediterráneos.*, García-Ruiz, J.M., and López
607 García, P., eds. (Zaragoza: Instituto Pirenaico de Ecología-CSIC), pp. 61–98.
- 608 Vázquez de la Cueva, A., García del Barrio, J.M., Quero, M.O., and Palomares, O.S. (2006). Recent fire regime in
609 peninsular Spain in relation to forest potential productivity and population density. *International Journal of Wildland*
610 *Fire* 15, 397.
- 611 Vescovi, E., Ravazzi, C., Arpentí, E., Finsinger, W., Pini, R., Valsecchi, V., Wick, L., Ammann, B., and Tinner, W.
612 (2007). Interactions between climate and vegetation during the Lateglacial period as recorded by lake and mire
613 sediment archives in Northern Italy and Southern Switzerland. *Quaternary Science Reviews* 26, 1650–1669.
- 614 Webb III, T., Ruddiman, W.F., Street-Perrot, F.A., Markgraf, V., Kutzbach, J.E., Bartlein, P.J., Wright Jr, H.E., and
615 Prell, W.L. (1993). Climatic changes during the Past 18000 years: regional syntheses, mechanisms and causes. In
616 *Global Climates since the Last Glacial Maximum*, (University of Minnesota Press), pp. 515–535.
- 617 Whitlock, C., and Larsen, C. (2002). *Tracking Environmental Change Using Lake Sediments*. (Springer Netherlands),
618 pp. 75–97.
- 619 Whitlock, C., Higuera, P.E., McWethy, D.B., and Briles, C.E. (2010a). Paleocological Perspectives on Fire Ecology:
620 Revisiting the Fire-Regime Concept. *TOECOLJ* 3, 6–23.
- 621 Whitlock, C., Tinner, W., Newman, Louise, and Kiefer, Thorsten (2010b). Fire in the Earth System: a
622 palaeoperspective. *Pages News* 18, 55–56.

623 **Tables**

624 Table 1: Criteria used for the selection of different core sections and analyses carried out in each of them.

625 Table 2: Spearman's R of some taxa and group of taxa and palynological richness. Asterisks are
626 significant values at $p < 0.05$.

627 **Figures**

628 Figure 1: Digital elevation model of the Western-central Southern Pyrenees. Sites mentioned in the text: 1)
629 El Portalet, 2) Estany de Burg, 3) Col d'Ech.

630 Figure 2: Vegetation map of the principal plant communities of the Western-central Spanish Pyrenees The
631 upper panel displays in red the mesophyte forest, including *Betula* and *Corylus* distributions in the area.
632 See Supplementary data in Supporting Information for taxa included in the different groups. Colour figure
633 available online.

634 Figure 3: A) Raw charcoal accumulation rate values (CHAR) (grey bars); transformed Log C interpolated
635 (variance stabilisation is achieved by log-transforming C interpolated series into Log C interpolated) (solid
636 black line); Log C background (Detrending Log C interpolated resulted in background charcoal (Log C
637 background) (red line). B) Fire events (FE) (red crosses) and detected fire peaks that did not pass the
638 threshold for FE (red dots). C) Fire return intervals (FRI) for the two studied periods, expressed as years
639 between fire events. D) Fire frequency (FF) every 500 years. E) Microcharcoal- accumulation rate. No
640 samples were analysed between 364 and 382 cm - more details are given in the text -. Colour figure
641 available online. F) Fire frequency (FF) every 500 years and fire events (FE) (black crosses) at Col d'Ech
642 (site 3 in Fig 1) during the Holocene period concurrent with our site. See more details on the methods in
643 the Supporting Information.

644 Figure 4: *Pinus*:mesophyte ratio plotted against local fire occurrence (fire frequency and peak magnitude)
645 and July summer insolation (42°N) (Laskar, 2004) in an age scale. Chronology from Lateglacial climate
646 events are defined after Lowe et al. (2008) except for the Allerød warm event as El Portalet pollen and

647 sedimentary signals found in González-Sampériz et al. (2006), marked as *Allerød/GI-1a** with a dotted-line
648 box in this figure, do not chronologically agree with the recent chronology presented in Lowe et al. (2008).
649 Colour figure available online.

650 Figure 5: Synthetic pollen diagram with relevant selected taxa. Pollen values are abundances (%)
651 represented as filled black areas. Fire activity is characterized by micro-charcoal area concentration, and
652 local fire events are represented by red crosses. Palynological richness is represented with confidence
653 intervals and measured in number of taxa. See Supplementary data for taxa included in the different
654 groups. Colour figure available online.

655 Figure 6: Cross-correlograms of detrended macro-charcoal influx versus detrended pollen percentages for
656 the 6-10 cal kyr BP period. Every lag corresponds to the average time difference between two adjacent
657 samples, both containing pollen and charcoal (15 yr cm⁻¹). Lag 0 is the ordinary linear correlation
658 coefficient between two variables at a particular time. Positive lags measure the influence of the first
659 variable on the second, ie., how charcoal affects pollen, after a particular time lag; negative lags measure
660 the influence of the second variable (pollen) on the first (charcoal) with reference to the lag. Dotted lines
661 correspond to the significance level $p < 0.05$.



662


Table 1


Depth (cm)	Age (cal kyr BP)	Analysis	Criteria
60-160	5 to 6	No analyses	Peatland to lake transition; different environmental settings for charcoal deposition
160-364	6 to 10	High resolution pollen, micro- and macro-charcoal (1 cm ³ every 1 cm)	Sedimentation rate meets the demands for decomposition of time series and peak-fire detection (15 yr/cm)
364-381	10 to 11.5	No analyses	Sedimentation rate too low in the very few available samples
381-382	11.5 to 12.5	No analyses	Possible hiatus corresponding to the Younger Dryas
382-471	12.5-14.7	Pollen and micro-charcoal every 2-5 cm, macro-charcoal every 1 cm	Sedimentation rate is high enough (24 yr/cm) fitting the required for peak-fire detection while pollen analysis was performed in the available samples from the original core
471-608	14.7-32	No analyses	Glacial influence and low presence of potentially local taxa were not suitable for the proposed objectives.


Table 2

Taxa or group of taxa	Spearman's R
Trees	-0.42 *
Mesophytes	0.27
Shrubs	0.65 *
Poaceae	0.34
Herbs	0.83 *
Non arboreal (shrub and all herbs)	0.75 *
<i>Pinus</i>	-0.20
<i>Betula</i>	-0.32
<i>Corylus</i>	-0.29












El Portalet
1802 masl




Detrended macrocharcoal influx vs




Pinus




Betula




Corylus




Deciduous *Quercus*




Mesophytes




Evergreen *Quercus*




Poaceae




Palynological richness




Microcharcoal > 10



Trees
Lag



Shrubs
Lag



Herbs
Lag