



# Bioreactor type affects the accumulation of phenolic acids and flavonoids in microshoot cultures of *Schisandra chinensis* (Turcz.) Baill.

Agnieszka Szopa<sup>1</sup> · Adam Kokotkiewicz<sup>2</sup> · Marcelina Bednarz<sup>1</sup> · Karolina Jafernik<sup>1</sup> · Maria Luczkiewicz<sup>2</sup> · Halina Ekiert<sup>1</sup>

Received: 14 March 2019 / Accepted: 20 August 2019 / Published online: 28 August 2019  
© The Author(s) 2019

## Abstract

Microshoots of the East Asian medicinal plant species *Schisandra chinensis* (Chinese magnolia vine) were grown in bioreactors characterized by different construction and cultivation mode. The tested systems included two continuous immersion systems—a cone-type bioreactor (CNB) and a cylindrical tube bioreactor (CTB), a nutrient sprinkle bioreactor (NSB), and two temporary immersion systems (TIS)—RITA® and Plantform. Microshoots were grown for 30 and 60 days in the MS medium enriched with 1 mg l<sup>-1</sup> NAA and 3 mg l<sup>-1</sup> BA. The accumulation of two groups of phenolic compounds: phenolic acids and flavonoids in the bioreactor-grown *S. chinensis* biomass, was evaluated for the first time. In the microshoot extracts, seven phenolic acids: chlorogenic, gallic, p-hydroxybenzoic, protocatechuic, syringic, salicylic and vanillic, and three flavonoids: kaempferol, quercitrin and rutoside, were identified. The highest total amount of phenolic acids (46.68 mg 100 g<sup>-1</sup> DW) was recorded in the biomass maintained in the CNB for 30 days. The highest total content of flavonoids (29.02 mg 100 g<sup>-1</sup> DW) was found in the microshoots maintained in the NSB for 30 days. The predominant metabolites in all the tested systems were: gallic acid (up to 10.01 mg 100 g<sup>-1</sup> DW), protocatechuic acid (maximal concentration 16.30 mg 100 g<sup>-1</sup> DW), and quercitrin (highest content 21.00 mg 100 g<sup>-1</sup> DW).

## Key message

The influence of bioreactor type on the accumulation of phenolic acids and flavonoids in microshoot cultures of *Schisandra chinensis* was proven and optimized.

**Keywords** Plant-dedicated bioreactors · Chinese magnolia vine · Schizandra · Phenolic secondary metabolites · Protocatechuic acid · Quercitrin

## Abbreviations

ADB Accumulated dry biomass  
BA 6-Benzyladenine  
CNB Cone-type bioreactor  
CTB Cylindrical tube bioreactor

DW Dry weight  
FW Fresh weight  
Gi Growth index  
HPLC–DAD High-performance liquid chromatography coupled with diode-array detection  
NAA 1-Naphthaleneacetic acid  
 $\mu$  Specific growth rate  
MS Murashige and Skoog  
NSB Nutrient sprinkle bioreactor  
TCM Traditional Chinese medicine  
TIS Temporary immersion systems

Communicated by Danny Geelen.

✉ Agnieszka Szopa  
a.szopa@uj.edu.pl

<sup>1</sup> Chair and Department of Pharmaceutical Botany, Jagiellonian University, Collegium Medicum, ul. Medyczna 9, 30-688 Kraków, Poland

<sup>2</sup> Chair and Department of Pharmacognosy, Faculty of Pharmacy, Medical University of Gdansk, al. gen. J. Hallera 107, 80-416 Gdańsk, Poland

Research on secondary metabolites production in bioreactor-grown biomasses, constitute an important topic in plant


biotechnology. The results of such studies are important from a practical point of view and the prospects of producing valuable, biologically active metabolites in such systems continue to draw scientists' attention (Verpoorte et al. 2002; Karuppusamy 2009). The latest studies have demonstrated the possibility of large-scale production of metabolites like: artemisinin (antimalarial drug) in cultures of *Artemisia annua* grown in a specially-designed stirred tank bioreactor, plumbagin (anticancer compound) in *Plumbago rosea* cultures maintained in a customized reaction kettle (Jose et al. 2016), and steviol glycosides (low-calorie glucoside sweeteners) in *Stevia rebaudiana* cultures grown in a temporary immersion bioreactor (Vives et al. 2017). In all studies involving large-scale in vitro systems, the selection of proper bioreactor type and optimization of process parameters is crucial for maximizing secondary metabolite production.

The role of plant-derived phenolic substances in diverse areas such as medicine, cosmetology and food industry, is invaluable. Secondary metabolites like flavonoids and phenolic acids exhibit many important, scientifically proven, biological activities, e.g. antioxidant (Young and Woodside 2001; Krishnaiah et al. 2011), anti-inflammatory (Zhang and Tsao 2016), or anticancer (Roleira et al. 2015). These compounds are used either in pure form or as constituents of herbs, plant extracts and functional foods.

In the current study, highly differentiated microshoot cultures of rare, East Asian plant species—*Schisandra chinensis* (Turcz.) Baill. (Chinese magnolia vine) were

investigated as a source of flavonoids and phenolic acids. *S. chinensis* fruits constitute raw pharmacopoeial material and are especially well-known in the traditional medicine of East-Asian countries. However, they have also been included in current phytotherapy all over the world (Barnes and Anderson 2007; World Health Organization 2007; European Directorate for the Quality of Medicines 2017; Szopa et al. 2017a). Their medicinal properties, e.g. hepatoprotective, immunostimulant, adaptogenic and anti-cancer, are determined mainly by the presence of dibenzocyclooctadiene type lignans (aka “schisandra lignans”), a *Schisandra* genus-specific group of secondary metabolites (Opletal et al. 2004; Szopa et al. 2017a). However, recent studies have demonstrated that phenolics present in *S. chinensis* act synergistically with lignans, thus serving as co-adjuvants and increasing biological activities of the plant (Szopa and Ekiert 2012; Cheng et al. 2013; Mocan et al. 2014). The compounds of particular interest are phenolic acids and flavonoids which themselves exhibit beneficial biological effects.

In this study, in vitro microshoot cultures of *S. chinensis* were evaluated for the production of phenolic compounds in five bioreactors designed for differentiated plant in vitro cultures (systems nomenclature after Kim et al. 2004; Steingroewer et al. 2013). Of these, three were custom-made systems: a cone-type balloon bioreactor (CNB) a cylindrical tube bioreactor (CTB) with internal rack for biomass immobilization and a nutrient sprinkle bioreactor (NSB), and two



**Fig. 1** Microshoot cultures of *Schisandra chinensis* grown in different bioreactor systems: **a** cone-type bioreactor (CNB); **b** cylindrical tube bioreactor (CTB); **c** nutrient sprinkle bioreactor (NSB); **d** RITA® bioreactor (open lid); **e** Platform bioreactor (open lid)

were commercially available, temporary immersion bioreactors—RITA® (Vitropic, France) and Plantform (PlantForm, Sweden) (Fig. 1). The construction details and setup of the bioreactors had been described by us previously (Szopa et al. 2017c). To our knowledge, this is the first work aimed at optimizing conditions for favourable biomass growth and phenolic acids and flavonoids production in bioreactor-grown *S. chinensis* microshoots.

As described previously, the microshoots were cultivated in the Murashige & Skoog (MS) medium (Murashige and Skoog 1962) supplemented with 1.0 mg l<sup>-1</sup> 1-naphthalene acetic acid (NAA) and 3.0 mg l<sup>-1</sup> 6-benzyladenine (BA), which had previously been demonstrated to provide the best biomass growth and secondary metabolites production in *S. chinensis* in vitro cultures (Szopa and Ekiert 2013, 2015; Szopa et al. 2016). All bioreactor systems were inoculated at 3/100 w/v ratio which corresponds to 15 g of microshoots per 500 ml of medium in CNB, CTB, NSB and Plantform bioreactors, and 6 g of microshoots per 200 ml of medium for RITA® container. The cultures were grown in constant artificial light (Philips fluorescent white lamps, 90 ± 2 μmol m<sup>-2</sup> s<sup>-1</sup>), at 24 ± 2 °C. The bioreactor cultures were maintained for 30 and 60 days and the collected biomasses were evaluated for growth parameters including fresh (FW) and dry (DW) weight, growth index (Gi), accumulated dry biomass (ADB) and specific growth rate (μ). The recorder values and formulae used are presented in Table 1. Afterwards, the microshoots and corresponding media samples (40 ml) were freeze dried (Lyovac GT2, Finland) and subjected to

phytochemical analysis. The bioreactor experiments were done in four repetitions.

The lyophilized and pulverized biomass samples (0.5 g from each bioreactor, four samples for each bioreactor type) were extracted with methanol (50 ml) for 2 h under reflux condenser (Harborne 1984). The extracts were transferred to crystallizers through a filter paper (Whatman paper) and left at room temperature to evaporate the solvent. The dry residue was dissolved in 2 ml of methanol (HPLC grade purity, Merck) and subjected to HPLC analysis.

Qualitative and quantitative HPLC–DAD analyses were performed according to the previously published protocols (Ellnain-Wojtaszek and Zgorzka 1999; Sułkowska-Ziaja et al. 2017; Szopa et al. 2017b). The equipment used and method parameters were as follows: HPLC–DAD system (Merck-Hitachi), Purospher Merck analytical RP-18 column (4 × 250 mm, 5 μm), mobile phase (gradient program): A—methanol with 0.5% acetic acid (1:4 v/v); B—methanol, flow—1 ml min<sup>-1</sup>, injection—10 μl, λ—254 nm, temp. 25 °C. The compounds were identified by co-chromatography with reference substances and comparison of their retention times and quantified using calibration curves. The phenolic acid standards (including benzoic and cinnamic acid and their derivatives) were as follows (t<sub>r</sub> values in minutes): gallic (3.86), neochlorogenic (4.98), protocatechuic (5.91), 3,4-dihydroxyphenylacetic (6.93), gentisic (10.00), chlorogenic (11.20), p-hydroxybenzoic (12.53), hydrocaffeic (15.37), vanillic (16.44), caffeic

**Table 1** Growth parameters of *Schisandra chinensis* microshoots maintained for 30 and 60 days in different bioreactor systems

Bioreactor type	Growth period (days)	FW (g l <sup>-1</sup> )	DW (g l <sup>-1</sup> )	Gi <sup>a</sup>	ADB (g l <sup>-1</sup> ) <sup>b</sup>	μ (d <sup>-1</sup> ) <sup>c</sup>	Biomass productivity (g l <sup>-1</sup> d <sup>-1</sup> )
CNB	30	120.29 ± 17.15	6.88 ± 0.93	3.01 ± 0.57	4.68 ± 0.93	0.0377 ± 0.0048	0.229 ± 0.031
	60	266.99 ± 25.55	15.41 ± 0.88	7.90 ± 0.85	13.21 ± 0.88	0.0324 ± 0.0010	0.257 ± 0.015
CTB	30	128.09 ± 22.80	8.08 ± 0.94	3.22 ± 0.75	5.87 ± 0.94	0.0431 ± 0.0040	0.269 ± 0.031
	60	233.91 ± 13.30	14.63 ± 0.71	6.73 ± 0.47	12.43 ± 0.71	0.0315 ± 0.0008	0.244 ± 0.012
NSB	30	126.34 ± 20.30	7.76 ± 1.00	3.21 ± 0.68	5.56 ± 1.00	0.0417 ± 0.0043	0.259 ± 0.033
	60	274.01 ± 21.13	15.50 ± 0.43	8.13 ± 0.70	13.29 ± 0.43	0.0325 ± 0.0005	0.258 ± 0.007
RITA®	30	134.62 ± 5.84	7.49 ± 0.21	3.49 ± 0.19	5.28 ± 0.21	0.0408 ± 0.0009	0.250 ± 0.007
	60	353.37 ± 20.04	17.86 ± 0.72	10.78 ± 0.67	15.65 ± 0.72	0.0349 ± 0.0007	0.298 ± 0.012
Plantform	30	105.78 ± 3.68	6.23 ± 0.18	2.53 ± 0.12	4.02 ± 0.18	0.0346 ± 0.0010	0.208 ± 0.006
	60	294.19 ± 19.30	17.07 ± 1.14	8.81 ± 0.64	14.86 ± 1.14	0.0341 ± 0.0011	0.284 ± 0.019

Growth parameters of *Schisandra chinensis* microshoots maintained for 30 and 60 days in different bioreactor systems: FW fresh weight; DW dry weight; Gi growth index; ADB accumulated dry biomass; μ specific growth rate and biomass productivity

<sup>a</sup>According to (Grzegorzczuk and Wysokińska 2008):  $Gi = [(Fw_1 - Fw_0) / Fw_0]$ , where Gi is the growth index, Fw<sub>1</sub> is the weight of microshoots at the end of experiment and Fw<sub>0</sub> is the fresh weight of the inoculum

<sup>b</sup>According to (Pavlov et al. 2007):  $ADB = FDB - IDB$  where ADB is the accumulated dry biomass (in g/L), FDB is the final dry biomass (g/L) and IDB is the initial dry biomass (g/L)

<sup>c</sup>According to (Homova et al. 2010):  $\mu = \ln(X/X_0) / \Delta t$  where μ is the specific growth rate (1/d), X<sub>0</sub> and X are the initial and final biomasses (g/l) and Δt is the culture time interval (d)

(17.83), syringic (20.14), ferulic (30.28), salicylic (31.76), p-coumaric (34.31), sinapic (35.41), o-coumaric (36.62), m-coumaric (37.81), isoferulic (39.18) and rosmarinic (44.32) acids (Sigma-Aldrich, USA). Standards used to estimate flavonoids included ( $t_r$  values in minutes): vitexin (42.07), cynaroside (43.57), hyperoside (46.04), myricetin (48.01), rutoside (49.26), quercitrin (50.29), apigetrin (53.27), trifolin (54.47), quercetin (58.32), luteolin (60.97) and kaempferol (64.42) (Sigma-Aldrich, USA). The results were statistically analyzed with one-way ANOVA and the Tukey's range test (Statistica 12, Poland).

In all the types of bioreactors employed, *S. chinensis* microshoots showed vigorous growth, and no necrosis or medium browning was observed (Fig. 1). After the biomasses were collected, their growth parameters were determined (Table 1). The Gi parameter ranged from 2.53 to 3.49 and from 6.73 to 10.78 for microshoots cultivated for 30 and 60 days, respectively. Of the custom made systems, the NSB yielded the highest ADB after 60 days of cultivation. In terms of Gi, ADB and biomass productivity, the TIS bioreactors performed substantially better than other systems employed. The highest biomass increments were obtained with RITA<sup>®</sup> bioreactors which was reflected by high values of all growth parameters, most notably ADB and biomass productivity (Table 1).

Based on HPLC-DAD quantitative analyses, the total contents of phenolic acids varied from 19.16 mg 100 g<sup>-1</sup> DW (CTB, 60-day growth period) to 46.68 mg 100 g<sup>-1</sup> DW (CNB, 30-day growth period). Comparing the amounts of phenolic acids, we found that the highest total quantities were obtained in biomass of the cultures grown over 30-day growth periods in all the tested types of bioreactor. The maximal total amount of phenolic acids achieved in biomass extracts from the CNB bioreactor was, respectively, 1.50, 1.18, 1.43 and 1.35 times higher than in the CTB, NSB, RITA<sup>®</sup> and Plantform bioreactors (Table 2).

Out of the nineteen phenolic acids, the analyses confirmed the presence of seven compounds: chlorogenic, gallic, p-hydroxybenzoic, protocatechuic, syringic, salicylic and vanillic acids (Table 2). The quantities of individual phenolic acids ranged from 0.72 mg 100 g<sup>-1</sup> DW for syringic acid (CTB, 60-day growth period) to 16.30 mg 100 g<sup>-1</sup> DW for protocatechuic acid (CNB, 30-day growth period). Besides protocatechuic acid, a considerable amounts of gallic acid (maximal 10.01 mg 100 g<sup>-1</sup> DW, CNB, 30-day growth period) and salicylic acid (maximal 6.74 mg 100 g<sup>-1</sup> DW, SB, 30-day growth period) were also confirmed (Table 2).

The total amounts of estimated flavonoids ranged from 12.90 mg 100 g<sup>-1</sup> DW (CTB, 60-day growth period) to 29.02 mg 100 g<sup>-1</sup> DW (NSB, 30-day growth period). Comparing the amounts of flavonoids in the respective systems, we found that the highest total contents were obtained in microshoots grown over 30-day growth periods in all the

tested types of bioreactor. The maximal total amount of flavonoids achieved in biomass extracts from the NSB bioreactor was, respectively, 1.40, 1.32, 1.36 and 1.42 times higher than in the CNB, CTB, RITA<sup>®</sup> and Plantform bioreactors (Table 3).

From the eleven flavonoids estimated, three compounds were found: kaempferol, quercitrin and rutoside (Table 3). The quantities of kaempferol ranged from 0.18 mg 100 g<sup>-1</sup> DW (CTB, 60-day growth period) to 6.49 mg 100 g<sup>-1</sup> DW (SB, 30-day growth period), and rutoside from 0.88 mg 100 g<sup>-1</sup> DW (CTB, 60-day growth period) to 1.87 mg 100 g<sup>-1</sup> DW (Plantform, 30-day growth period). Quercitrin was the main metabolite; its amounts ranged from 11.85 mg 100 g<sup>-1</sup> DW (CTB, 60-day growth period) up to 21.00 mg 100 g<sup>-1</sup> DW (NSB, 30-day growth period) (Table 3).

In all experiments, none of the estimated phenols were found in the media samples, indicating that the investigated compounds were stored solely intracellularly.

Our study proved that biomass growth and secondary metabolite contents in *S. chinensis* microshoots depended on the type of bioreactor used and its mode of operation, including medium circulation and aeration, which is in agreement with previous reports (Liu et al. 2003; Zobayed et al. 2004; Paek et al. 2005), including our work on the production of dibenzocyclooctadiene type lignans in bioreactor-grown *S. chinensis* microshoots (Szopa et al. 2017c). In that study, the best conditions were provided by the Plantform bioreactor, in which, during a 30-day growth period, microshoots had accumulated the highest amounts of lignans (546.98 mg 100 g<sup>-1</sup> DW), with gomisins A, deoxyschisandrin and schisandrin as the major constituents (67.86, 77.66 and 118.59 mg 100 g<sup>-1</sup> DW, respectively) (Szopa et al. 2017c). As mentioned before, the pharmacological action of *S. chinensis* is the result of synergistic action of all the plant's components; in this case, phenolic compounds seem to play a major role besides lignans (Mocan et al. 2014; Szopa et al. 2017b). We had proved before that different types of in vitro cultures of *S. chinensis* were able to produce phenolic acids and flavonoids, and that their yield depended on the in vitro conditions used, such as the basal media formulation, concentration of phytohormones, duration of the growth period, lighting conditions, as well as the mode of cultivation (agar, agitated, stationary liquid) (Szopa and Ekiert 2012, 2016; Szopa et al. 2017b).

Under the current study, we made the first step towards scaling-up the production of phenolic compounds using *S. chinensis* microshoot cultures. Of the tested systems, the RITA<sup>®</sup> TIS bioreactor provided the highest biomass yields (Table 1). However, the total phenolic acids content was considerably greater in the CNB (Table 2), while that of flavonoids in the NSB (Table 3). In the CNB system, aeration-induced mechanical stress caused by tissue-medium contact resulted in lower biomass growth (Gi equal to 3.01

**Table 2** Phenolic acid contents (mg 100 g<sup>-1</sup> DW ± SD) in extracts from biomasses of *Schisandra chinensis* microshoots cultivated in different types of bioreactors (n = 4, p < 0.05)

Phenolic acids	Bioreactor type									
	CNB		CTB		NSB		RITA®		Plantform	
	Growth period (days)									
	30	60	30	60	30	60	30	60	30	60
Chlorogenic acid	4.65 ± 0.3 <sup>cederghij</sup>	4.23 ± 0.14 <sup>cdelghij</sup>	2.46 ± 0.11 <sup>abefi</sup>	2.40 ± 0.17 <sup>abefi</sup>	3.75 ± 0.29 <sup>abcdeghij</sup>	3.56 ± 0.88 <sup>abcdeghij</sup>	2.32 ± 0.3 <sup>labefi</sup>	2.14 ± 0.98 <sup>abefi</sup>	2.98 ± 0.27 <sup>abef</sup>	2.81 ± 0.45 <sup>gher</sup>
Gallie acid	10.01 ± 1.04 <sup>bdhij</sup>	6.67 ± 0.30 <sup>acdeghij</sup>	8.50 ± 0.52 <sup>bcghj</sup>	2.75 ± 0.13 <sup>abcefergi</sup>	8.87 ± 0.35 <sup>bdhij</sup>	3.64 ± 0.13 <sup>abcdeghij</sup>	8.77 ± 0.62 <sup>bdhij</sup>	3.25 ± 0.14 <sup>abcdeghj</sup>	7.45 ± 0.32 <sup>acdfhij</sup>	2.44 ± 0.27 <sup>abcdehij</sup>
p-Hydroxybenzoic acid	3.58 ± 0.46 <sup>bcdeghij</sup>	2.56 ± 0.08 <sup>acdeh</sup>	1.82 ± 0.09 <sup>abef</sup>	1.74 ± 0.08 <sup>abefgi</sup>	3.12 ± 0.28 <sup>cdghij</sup>	3.11 ± 0.37 <sup>bedghij</sup>	2.23 ± 0.54 <sup>abefh</sup>	1.79 ± 0.41 <sup>abefgi</sup>	2.18 ± 0.22 <sup>acdeh</sup>	2.08 ± 0.38 <sup>acdeh</sup>
Protocatechuic acid	16.30 ± 1.15 <sup>cederghij</sup>	15.97 ± 0.66 <sup>cdelghij</sup>	8.38 ± 0.23 <sup>abdefi</sup>	5.07 ± 0.12 <sup>abceferghij</sup>	12.20 ± 0.47 <sup>abcdeghij</sup>	9.33 ± 0.63 <sup>abdefi</sup>	9.87 ± 0.84 <sup>abde</sup>	8.52 ± 0.68 <sup>abdefij</sup>	10.85 ± 0.58 <sup>abceh</sup>	9.01 ± 0.42 <sup>abdefi</sup>
Salicylic acid	5.66 ± 0.06 <sup>deij</sup>	5.91 ± 0.19 <sup>deij</sup>	5.09 ± 0.11 <sup>efi</sup>	5.00 ± 0.09 <sup>abefj</sup>	6.74 ± 0.18 <sup>abcdeghj</sup>	6.05 ± 0.86 <sup>abdefhij</sup>	5.27 ± 0.25 <sup>cedefj</sup>	4.87 ± 0.36 <sup>abefi</sup>	5.84 ± 0.36 <sup>cdelghj</sup>	4.29 ± 0.40 <sup>bcdehij</sup>
Syringic acid	0.77 ± 0.03 <sup>efgi</sup>	0.74 ± 0.02 <sup>efgi</sup>	1.92 ± 0.06 <sup>abdefghij</sup>	0.72 ± 0.02 <sup>efgi</sup>	0.84 ± 0.11 <sup>efgi</sup>	0.83 ± 0.17 <sup>efgi</sup>	0.98 ± 0.22 <sup>abcdehij</sup>	0.74 ± 0.11 <sup>efgi</sup>	1.25 ± 0.11 <sup>abcehijghj</sup>	0.87 ± 0.09 <sup>bcgij</sup>
Vanillic acid	5.71 ± 0.50 <sup>bcdehij</sup>	3.85 ± 0.28 <sup>acdeghij</sup>	2.92 ± 0.05 <sup>abdefhi</sup>	1.50 ± 0.04 <sup>abceferghij</sup>	4.03 ± 0.27 <sup>acdeghj</sup>	3.91 ± 0.13 <sup>acdeghj</sup>	3.21 ± 0.21 <sup>abdefhi</sup>	1.86 ± 0.54 <sup>abcdfghij</sup>	4.01 ± 0.35 <sup>acdeghj</sup>	2.88 ± 0.32 <sup>abdefhi</sup>
Total content	46.68 ± 3.61 <sup>bcdehij</sup>	39.92 ± 1.66 <sup>acdfghij</sup>	31.10 ± 1.18 <sup>abdefhij</sup>	19.16 ± 0.65 <sup>abceferghij</sup>	39.56 ± 1.95 <sup>acdfghij</sup>	30.44 ± 3.17 <sup>abdefhij</sup>	32.65 ± 2.99 <sup>abcdehij</sup>	23.17 ± 3.22 <sup>abcdehij</sup>	34.56 ± 2.21 <sup>abdefhij</sup>	24.38 ± 2.33 <sup>abcdehij</sup>

<sup>a</sup>p < 0.05 versus CNB cultures 30 days growth period

<sup>b</sup>p < 0.05 versus CNB cultures 60 days growth period

<sup>c</sup>p < 0.05 versus CTB cultures 30 days growth period

<sup>d</sup>p < 0.05 versus CTB cultures 60 days growth period

<sup>e</sup>p < 0.05 versus NSB cultures 30 days growth period

<sup>f</sup>p < 0.05 versus NSB cultures 60 days growth period

<sup>g</sup>p < 0.05 versus RITA® bioreactor cultures 30 days growth period

<sup>h</sup>p < 0.05 versus RITA® bioreactor cultures 60 days growth period

<sup>i</sup>p < 0.05 versus Plantform bioreactor cultures 30 days growth period

<sup>j</sup>p < 0.05 versus Plantform bioreactor cultures 60 days growth period



**Table 3** Flavonoid contents (mg 100 g<sup>-1</sup> DW ± SD) in extracts from biomasses of *Schisandra chinensis* microshoots cultivated in in different types of bioreactors (n = 4, p < 0.05)

Flavonoids	Bioreactor type																		
	CNB			CTB			NSB			RITA®			Plantform						
	Growth period (days)																		
	30		60		30		60		30		60		30		60				
Kaempferol	1.14 ± 0.07 <sup>def</sup>	1.05 ± 0.06 <sup>cdef</sup>	2.07 ± 0.06 <sup>abdehij</sup>	0.18 ± 0.02 <sup>abcdehij</sup>	6.49 ± 0.62 <sup>abcdehij</sup>	3.16 ± 0.08 <sup>abcdehij</sup>	1.11 ± 0.17 <sup>cdef</sup>	1.08 ± 0.28 <sup>cdef</sup>	0.98 ± 0.12 <sup>cdefg</sup>	18.39 ± 0.43 <sup>hij</sup>	17.71 ± 0.36 <sup>degh</sup>	18.34 ± 0.34 <sup>fhj</sup>	11.85 ± 0.09 <sup>abcdegi</sup>	21.00 ± 0.22 <sup>abcdegi</sup>	13.27 ± 0.69 <sup>klhij</sup>	12.55 ± 0.96 <sup>abcdegi</sup>	17.53 ± 0.85 <sup>dehij</sup>	13.25 ± 0.54 <sup>abcdegi</sup>	
Quercitrin	1.26 ± 0.04 <sup>cdegi</sup>	1.25 ± 0.06 <sup>cdegi</sup>	1.62 ± 0.02 <sup>abdfj</sup>	0.88 ± 0.01 <sup>abcdehij</sup>	1.53 ± 0.07 <sup>abdf</sup>	1.01 ± 0.03 <sup>seg</sup>	1.27 ± 0.30 <sup>cde</sup>	1.87 ± 0.21 <sup>abdfhij</sup>	1.22 ± 0.73 <sup>cdegi</sup>	20.78 ± 0.54 <sup>cdehij</sup>	20.00 ± 0.48 <sup>cdehij</sup>	22.03 ± 0.42 <sup>abdehij</sup>	12.90 ± 0.12 <sup>abcdegi</sup>	29.02 ± 1.38 <sup>abcdehij</sup>	17.44 ± 1.38 <sup>abcdehij</sup>	13.93 ± 1.43 <sup>abcdehij</sup>	20.48 ± 1.34 <sup>cdehij</sup>	15.45 ± 1.39 <sup>abcdegi</sup>	
Total content																			

<sup>a</sup>p < 0.05 versus CNB cultures 30 days growth period<sup>b</sup>p < 0.05 versus CNB cultures 60 days growth period<sup>c</sup>p < 0.05 versus CTB cultures 30 days growth period<sup>d</sup>p < 0.05 versus CTB cultures 60 days growth period<sup>e</sup>p < 0.05 versus NSB cultures 30 days growth period<sup>f</sup>p < 0.05 versus NSB cultures 60 days growth period<sup>g</sup>p < 0.05 versus RITA® bioreactor cultures 30 days growth period<sup>h</sup>p < 0.05 versus RITA® bioreactor cultures 60 days growth period<sup>i</sup>p < 0.05 versus Plantform bioreactor cultures 30 days growth period<sup>j</sup>p < 0.05 versus Plantform bioreactor cultures 60 days growth period

and 7.90, after 30- and 60-day growth cycles, respectively) (Table 1), but also in higher production of phenolic acids (Table 2). Phenolic compounds are the plant's "stress metabolites", hence the high production (Halliwell 2003, 2007; Akula and Ravishankar 2011). Moreover, it is worth noting that the 'bubble (air-lift) bioreactors' had been successfully applied before for the production of caffeic acid derivatives in *Echinacea purpurea* adventitious root cultures (Jeong et al. 2009), ginsenosides in adventitious root cultures of *Panax ginseng* (Kim et al. 2004) as well as phenolic acids and flavonoids in *Eleutherococcus senticosus* somatic embryos (Shohael et al. 2006).

In this study, the gas-phase nutrient sprinkle bioreactor (NSB) proved to be the best for flavonoid production (Table 3). The characteristic feature of this bioreactor design is the lack of mechanical stress exerted on growing biomass, as well as good gas circulation and nutrient accessibility (Steingroewer et al. 2013; Georgiev et al. 2014). Similar to our study, the correlative experiment on *Artemisia annua* in vitro shoots had showed an improvement in artemisinin output in a gas-phase bioreactor (Liu et al. 2006). Moreover, shoots grown in a nutrient sprinkle bioreactor have been employed for effective biomass growth and production of phenolic compounds, expressed as total phenolic acids and flavonoids, in medicinal plants such as *Rehmannia glutinosa* (Piątczak et al. 2014) and *Scutellaria alpina* (Grzegorzczak-Karolak et al. 2017).

To conclude, our work showed that *S. chinensis* in vitro microshoots grown in different types of bioreactors (Fig. 1, Table 1), could provide biologically active phenolic compounds. Moreover, we proved, for the first time, the influence of bioreactor type on the phenolic acid and flavonoid production in *S. chinensis* biomass (Tables 2 and 3). The obtained results are thus a good starting point for further studies, involving scale-up experiments.

**Acknowledgements** This study was funded by National Science Centre, Poland—No. 2016/23/D/NZ7/01316, and project supported by the Polish Ministry of Science and Higher Education—No. K/DSC/004297.

**Author contributions** AS, AK, MB and KJ performed the research and elaborated the data. AK performed the bioreactor experiments. AS and MB performed the chemical analysis. AS performed the statistical analysis and prepared the manuscript. AK, ML and HE checked and corrected the manuscript.

## Compliance with ethical standards

**Conflict of interest** All authors approved the manuscript and declare that there are no conflicts of interests.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

[mmons.org/licenses/by/4.0/](http://creativecommons.org/licenses/by/4.0/)), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Akula R, Ravishankar GA (2011) Influence of abiotic stress signals on secondary metabolites in plants. *Plant Signal Behav* 6:1720–1731. <https://doi.org/10.4161/psb.6.11.17613>
- Barnes J, Anderson LA (2007) *Herbal medicines*, 2nd edn. Pharmaceutical press, London
- Cheng N, Ren N, Gao H et al (2013) Antioxidant and hepatoprotective effects of *Schisandra chinensis* pollen extract on CCl<sub>4</sub>-induced acute liver damage in mice. *Food Chem Toxicol* 55:234–240. <https://doi.org/10.1016/j.fct.2012.11.022>
- Ellnain-Wojtaszek M, Zgorzka G (1999) High-performance liquid chromatography and thin-layer chromatography of phenolic acids from *Ginkgo biloba* L. leaves collected within vegetative period. *J Liq Chromatogr Relat Technol* 22:1457–1471. <https://doi.org/10.1081/JLC-100101744>
- European Directorate for the Quality of Medicines (2017) *Schisandra* fruit. *European Pharmacopoeia* 9.0. Strasburg, p 1514
- Georgiev V, Schumann A, Pavlov A, Bley T (2014) Temporary immersion systems in plant biotechnology. *Eng Life Sci* 14:607–621. <https://doi.org/10.1002/elsc.201300166>
- Grzegorzczak I, Wysokińska H (2008) Liquid shoot culture of *Salvia officinalis* L. for micropropagation and production of antioxidant compounds; effects of triacontanol. *Acta Soc Bot Pol* 77:99–104. <https://doi.org/10.5586/asbp.2008.013>
- Grzegorzczak-Karolak I, Rytczak P, Bielecki S, Wysokińska H (2017) The influence of liquid systems for shoot multiplication, secondary metabolite production and plant regeneration of *Scutellaria alpina*. *Plant Cell, Tissue Organ Cult* 128:479–486. <https://doi.org/10.1007/s11240-016-1126-y>
- Halliwell B (2003) Oxidative stress in cell culture: an under-appreciated problem? *FEBS Lett* 540:3–6. [https://doi.org/10.1016/S0014-5793\(03\)00235-7](https://doi.org/10.1016/S0014-5793(03)00235-7)
- Halliwell B (2007) Biochemistry of oxidative stress. *Biochem Soc Trans* 35:1147–1150. <https://doi.org/10.1042/BST0351147>
- Harborne JB (1984) *Phytochemical Methods*. Springer, Dordrecht
- Homova V, Weber J, Schulze J et al (2010) Devil's claw hairy root culture in flasks and in a 3-L bioreactor: bioactive metabolite accumulation and flow cytometry. *Z Naturforsch, C: J Biosci* 65:439–446. <https://doi.org/10.1515/znc-2010-7-809>
- Jeong J-A, Wu C-H, Murthy HN et al (2009) Application of an air-lift bioreactor system for the production of adventitious root biomass and caffeic acid derivatives of *Echinacea purpurea*. *Biotechnol Bioprocess Eng* 14:91–98. <https://doi.org/10.1007/s12257-007-0142-5>
- Jose B, Pillai DB, Satheeshkumar K (2016) In vitro cultivation of hairy roots of *Plumbago rosea* L. in a customized reaction kettle for the production of plumbagin—an anticancer compound. *Ind Crops Prod* 87:89–95. <https://doi.org/10.1016/j.indcrop.2016.04.023>
- Karuppusamy S (2009) A review on trends in production of secondary metabolites from higher plants by in vitro tissue, organ and cell cultures. *J Med Plants Res* 3:1222–1239
- Kim Y-S, Hahn E-J, Peak KY (2004) Effects of various bioreactors on growth and ginsenoside accumulation in ginseng adventitious root cultures (*Panax ginseng* C.A. Meyer). *J Plant Biotechnol* 31(3):249–253. <https://doi.org/10.5010/JPB.2004.31.3.249>

- Krishnaiah D, Sarbatly R, Nithyanandam R (2011) A review of the antioxidant potential of medicinal plant species. *Food Bioprod Process* 89:217–233
- Liu CZ, Guo C, Wang YC, Ouyang F (2003) Comparison of various bioreactors on growth and artemisinin biosynthesis of *Artemisia annua* L. shoot cultures. *Process Biochem* 39:45–49. [https://doi.org/10.1016/S0032-9592\(02\)00294-7](https://doi.org/10.1016/S0032-9592(02)00294-7)
- Liu C, Zhao Y, Wang Y (2006) Artemisinin: current state and perspectives for biotechnological production of an antimalarial drug. *Appl Microbiol Biotechnol* 72:11–20. <https://doi.org/10.1007/s00253-006-0452-0>
- Mocan A, Crişan G, Vlase L et al (2014) Comparative studies on polyphenolic composition, antioxidant and antimicrobial activities of *Schisandra chinensis* leaves and fruits. *Molecules* 19:15162–15179. <https://doi.org/10.3390/molecules190915162>
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 15:473–479
- Opletal L, Sovová H, Bártlová M (2004) Dibenzo[a, c]cyclooctadiene lignans of the genus *Schisandra*: importance, isolation and determination. *J Chromatogr B* 812:357–371. <https://doi.org/10.1016/j.jchromb.2004.07.040>
- Paek KY, Chakrabarty D, Hahn EJ (2005) Application of bioreactor systems for large scale production of horticultural and medicinal plants. *Plant Cell, Tissue Organ Cult* 81:287–300. <https://doi.org/10.1007/s11240-004-6648-z>
- Pavlov A, Georgiev M, Bley T (2007) Batch and fed-batch production of betalains by red beet (*Beta vulgaris*) hairy roots in a bubble column reactor. *Z Naturforsch, C: J Biosci* 62:439–446
- Piątczak E, Grzegorzczak-Karolak I, Wysokińska H (2014) Micropropagation of *Rehmannia glutinosa* Libosch.: production of phenolics and flavonoids and evaluation of antioxidant activity. *Acta Physiol Plant* 36:1693–1702. <https://doi.org/10.1007/s11738-014-1544-6>
- Roleira FMF, Tavares-da-Silva EJ, Varela CL et al (2015) Plant derived and dietary phenolic antioxidants: anticancer properties. *Food Chem* 183:235–258. <https://doi.org/10.1016/j.foodchem.2015.03.039>
- Shohael AM, Ali MB, Yu KW et al (2006) Effect of light on oxidative stress, secondary metabolites and induction of antioxidant enzymes in *Eleutherococcus senticosus* somatic embryos in bioreactor. *Process Biochem* 41:1179–1185. <https://doi.org/10.1016/j.procbio.2005.12.015>
- Steingroewer J, Bley T, Georgiev V et al (2013) Bioprocessing of differentiated plant in vitro systems. *Eng Life Sci* 13:26–38. <https://doi.org/10.1002/elsc.201100226>
- Sułkowska-Ziaja K, Maślanka A, Szewczyk A, Muszyńska B (2017) Physiologically active compounds in four species of *Phellinus*. *Nat Prod Commun* 12:363
- Szopa A, Ekiert H (2012) In vitro cultures of *Schisandra chinensis* (Turcz.) Baill. (Chinese magnolia vine)—a potential biotechnological rich source of therapeutically important phenolic acids. *Appl Biochem Biotechnol* 166:1941–1948. <https://doi.org/10.1007/s12010-012-9622-y>
- Szopa A, Ekiert H (2013) Production of deoxyschizandrin and gamma-schizandrin in shoot-differentiating and undifferentiating callus cultures of *Schisandra chinensis* (Turcz.) Baill. (Chinese magnolia vine). *J Biotechnol* 165:209–213
- Szopa A, Ekiert H (2015) Production of schisantherin A and gomisin G in in vitro cultures of *Schisandra chinensis*. *Phytochem Lett* 11:440–444. <https://doi.org/10.1016/J.PHYTOL.2014.12.022>
- Szopa A, Ekiert H (2016) The importance of applied light quality on the production of lignans and phenolic acids in *Schisandra chinensis* (Turcz.) Baill. cultures in vitro. *Plant Cell, Tissue Organ Cult* 127:115–121. <https://doi.org/10.1007/s11240-016-1034-1>
- Szopa A, Kokotkiewicz A, Marzec-Wróblewska U et al (2016) Accumulation of dibenzocyclooctadiene lignans in agar cultures and in stationary and agitated liquid cultures of *Schisandra chinensis* (Turcz.) Baill. *Appl Microbiol Biotechnol* 100:3965–3977. <https://doi.org/10.1007/s00253-015-7230-9>
- Szopa A, Ekiert R, Ekiert H (2017a) Current knowledge of *Schisandra chinensis* (Turcz.) Baill. (Chinese magnolia vine) as a medicinal plant species: a review on the bioactive components, pharmacological properties, analytical and biotechnological studies. *Phytochem Rev* 16:195–218. <https://doi.org/10.1007/s11101-016-9470-4>
- Szopa A, Kokotkiewicz A, Bednarz M et al (2017b) Studies on the accumulation of phenolic acids and flavonoids in different in vitro culture systems of *Schisandra chinensis* (Turcz.) Baill. using a DAD-HPLC method. *Phytochem Lett* 20:462–469. <https://doi.org/10.1016/j.phytol.2016.10.016>
- Szopa A, Kokotkiewicz A, Luczkiewicz M, Ekiert H (2017c) *Schisandra* lignans production regulated by different bioreactor type. *J Biotechnol* 247:11–17. <https://doi.org/10.1016/j.jbiotec.2017.02.007>
- Verpoorte R, Contín A, Memelink J (2002) Biotechnology for the production of plant secondary metabolites. *Phytochem Rev* 1:13–25. <https://doi.org/10.1023/A:1015871916833>
- Vives K, Andújar I, Lorenzo JC et al (2017) Comparison of different in vitro micropropagation methods of *Stevia rebaudiana* B. including temporary immersion bioreactor (BIT®). *Plant Cell, Tissue Organ Cult* 131:195–199. <https://doi.org/10.1007/s11240-017-1258-8>
- World Health Organization (2007) WHO monographs on selected medicinal plants, vol 3. *Fructus Schisandrae*. Geneva
- Young IS, Woodside JV (2001) Antioxidants in health and disease. *J Clin Pathol* 54:176–186
- Zhang H, Tsao R (2016) Dietary polyphenols, oxidative stress and antioxidant and anti-inflammatory effects. *Curr Opin Food Sci* 8:33–42. <https://doi.org/10.1016/j.cofs.2016.02.002>
- Zobayed SMA, Murch SJ, Rupasinghe HPV et al (2004) Optimized system for biomass production, chemical characterization and evaluation of chemo-preventive properties of *Scutellaria baicalensis* Georgi. *Plant Sci* 167:439–446. <https://doi.org/10.1016/j.plantsci.2004.04.022>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.