

**Bis(2,3-diaminopyridinium) bis( $\mu$ -pyridine-2,6-dicarboxylato)- $\kappa^4 O^2, N, O^6; \kappa^4 O^2: O^2, N, O^6$ -bis[aqua-(pyridine-2,6-dicarboxylato- $\kappa^3 O^2, N, O^6$ )-bismuthate(III)] tetrahydrate**

Hossein Aghabozorg,<sup>a\*</sup> Shokoofeh Kazemi,<sup>a</sup> Ali Akbar Agah,<sup>b</sup> Masoud Mirzaei<sup>c</sup> and Behrouz Notash<sup>d</sup>

<sup>a</sup>Faculty of Chemistry, Islamic Azad University, North Tehran Branch, Tehran, Iran, <sup>b</sup>Faculty of Chemistry, Tarbiat Moallem University, 15614 Tehran, Iran, <sup>c</sup>Department of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Mashhad 917791436, Iran, and <sup>d</sup>Department of Chemistry, Shahid Beheshti University, G. C., Evin, Tehran 1983963113, Iran

Correspondence e-mail: haghabozorg@yahoo.com


Received 9 February 2011; accepted 15 February 2011

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(C-C) = 0.012$  Å;  $R$  factor = 0.057;  $wR$  factor = 0.154; data-to-parameter ratio = 17.1.

In the centrosymmetric dinuclear complex anion of the title compound,  $(C_5H_8N_3)_2[Bi_2(C_7H_3NO_4)_4(H_2O)_2] \cdot 4H_2O$ , the Bi<sup>III</sup> atom is eight-coordinated in an  $N_2O_6$  environment and has a distorted bicapped trigonal-prismatic coordination environment. Extensive intermolecular  $O-H \cdots O$ ,  $N-H \cdots O$  and weak  $C-H \cdots O$  hydrogen bonds lead to the stability of the crystal structure. Interactions between one C-H group of the 2,3-diaminopyridinium [(2,3-dapyH)<sup>+</sup>] cation and the aromatic ring of the pyridine-2,6-dicarboxylate (pydc) ligand ( $C-H \cdots$ centroid distance = 2.78 Å) and  $\pi-\pi$  interactions between the (2,3-dapyH)<sup>+</sup> cations and between the (2,3-dapyH)<sup>+</sup> cation and the pydc ligand [centroid-centroid distances = 3.489 (5) and 3.694 (5) Å] are observed.

**Related literature**

For related structures, see: Aghabozorg *et al.* (2008, 2010); Sheshmani *et al.* (2005).



**Experimental**

*Crystal data*

$(C_5H_8N_3)_2[Bi_2(C_7H_3NO_4)_4 \cdot (H_2O)_2] \cdot 4H_2O$   
 $M_r = 1406.76$   
 Triclinic,  $P\bar{1}$   
 $a = 9.3462$  (19) Å  
 $b = 10.726$  (2) Å  
 $c = 11.098$  (2) Å  
 $\alpha = 95.13$  (3)°

$\beta = 91.38$  (3)°  
 $\gamma = 90.47$  (3)°  
 $V = 1107.7$  (4) Å<sup>3</sup>  
 $Z = 1$   
 Mo  $K\alpha$  radiation  
 $\mu = 8.03$  mm<sup>-1</sup>  
 $T = 298$  K  
 $0.33 \times 0.27 \times 0.23$  mm

*Data collection*

Stoe IPDS-2 diffractometer  
 Absorption correction: numerical (*X-SHAPE* and *X-RED32*; Stoe & Cie, 2005)  
 $T_{min} = 0.083$ ,  $T_{max} = 0.156$

12406 measured reflections  
 5940 independent reflections  
 5539 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.113$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.057$   
 $wR(F^2) = 0.154$   
 $S = 1.05$   
 5940 reflections  
 347 parameters  
 9 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{max} = 2.98$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -2.93$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3A...O11 <sup>i</sup>	0.87 (14)	2.02 (14)	2.797 (10)	148 (12)
N4—H4A...O2 <sup>ii</sup>	0.86	2.11	2.924 (9)	158
N4—H4B...O6 <sup>i</sup>	0.86	1.97	2.830 (10)	178
N5—H5A...O3 <sup>iii</sup>	0.86	2.53	3.216 (10)	138
N5—H5B...O6 <sup>i</sup>	0.86	2.11	2.972 (10)	176
O9—H9A...O8 <sup>iv</sup>	0.81 (7)	2.07 (12)	2.746 (10)	141 (15)
O9—H9B...O11 <sup>v</sup>	0.84 (8)	1.99 (10)	2.771 (11)	155 (15)
O10—H10A...O5	0.94 (8)	1.96 (8)	2.862 (9)	160 (11)
O10—H10B...O7 <sup>vi</sup>	0.80 (8)	2.21 (8)	2.972 (10)	160 (13)
O11—H11A...O4 <sup>iii</sup>	0.89 (8)	2.01 (13)	2.717 (11)	136 (14)
O11—H11B...O10 <sup>vii</sup>	0.92 (9)	1.96 (9)	2.836 (12)	158 (13)
C11—H11...O8 <sup>viii</sup>	0.93	2.26	3.050 (10)	142

Symmetry codes: (i)  $-x + 2, -y + 1, -z + 1$ ; (ii)  $-x + 2, -y, -z + 1$ ; (iii)  $-x + 1, -y + 1, -z + 1$ ; (iv)  $-x, -y, -z + 2$ ; (v)  $x - 1, y - 1, z$ ; (vi)  $x + 1, y, z$ ; (vii)  $x, y + 1, z$ ; (viii)  $-x, -y + 1, -z + 2$ .

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to the Islamic Azad University, North Tehran Branch, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2406).

## References

- Aghabozorg, H., Manteghi, F. & Sheshmani, S. (2008). *J. Iran. Chem. Soc.* **5**, 184–227.
- Aghabozorg, H., Moteieyan, E., Salimi, A. R., Mirzaei, M., Manteghi, F., Shokrollahi, A., Derki, S., Ghadermazi, M., Sheshmani, S. & Eshtiagh-Hosseini, H. (2010). *Polyhedron*, **29**, 1453–1464.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheshmani, S., Kheirollahi, P. D., Aghabozorg, H., Shokrollahi, A., Kickelbick, G., Shamsipur, M., Ramezanipour, F. & Moghimi, A. (2005). *Z. Anorg. Allg. Chem.* **631**, 3058–3065.
- Stoe & Cie (2005). *X-AREA*, *X-SHAPE* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.

**supplementary materials**

*Acta Cryst.* (2011). E67, m360-m361 [ doi:10.1107/S1600536811005629 ]

**Bis(2,3-diaminopyridinium) bis( $\mu$ -pyridine-2,6-dicarboxylato)- $\kappa^4 O^2, N, O^6: O^6; \kappa^4 O^2: O^2, N, O^6$ -bis[aqua(pyridine-2,6-dicarboxylato)- $\kappa^3 O^2, N, O^6$ ] bismuthate(III)] tetrahydrate**

**H. Aghabozorg, S. Kazemi, A. A. Agah, M. Mirzaei and B. Notash**

**Comment**

Pyridine-2,6-dicarboxylic acid (pydcH<sub>2</sub>) can form various complexes containing transition and main metals (Aghabozorg *et al.*, 2008). There are complexes in which pydc acts as a bridging ligand between two metal atoms (Aghabozorg *et al.*, 2010; Sheshmani *et al.*, 2005).

Herein, we report the crystal structure of the title compound as another example of bismuth(III) coordination compound, which bears heterocyclic 2,3-diaminopyridine (2,3-dapy) and pydcH<sub>2</sub> ligands. The molecular structure of the title compound is shown in Fig. 1. The centrosymmetric binuclear unit consists of two Bi<sup>III</sup> atoms, four (pydc)<sup>2-</sup> ligands, two coordinated water molecules. Two pydc ligands act as tridentate ligands with an N atom of the pyridine ring and two O atoms of the dicarboxylate groups acting as donors. One of the dicarboxylate O atoms for the other two pydc ligands plays a bridging role between two Bi atoms. The structure also contains two (2,3-dapyH)<sup>+</sup> cations and four uncoordinated water molecules. The Bi<sup>III</sup> atom is eight-coordinated in an N<sub>2</sub>O<sub>6</sub> environment and has a distorted bicapped trigonal-prismatic geometry, as it is shown in Fig. 2. There are extensive intermolecular O—H $\cdots$ O, N—H $\cdots$ O and weak C—H $\cdots$ O hydrogen bonds, which cause the stability of the crystal structure (Fig. 3, Table 1). There are also  $\pi$ – $\pi$  interactions between the (2,3-dapyH)<sup>+</sup> rings and between the (2,3-dapyH)<sup>+</sup> and pydc rings (Fig. 4), with centroid–centroid distances of 3.489 (5) and 3.694 (5) Å, respectively. Furthermore, there is C—H $\cdots$  $\pi$  interaction between C—H group of the (2,3-dapyH)<sup>+</sup> cation and pydc ligand, with an C—H $\cdots$ centroid distance of 2.78 Å (Fig. 5).

**Experimental**

An aqueous solution of Bi(NO<sub>3</sub>)<sub>3</sub> (1 mmol), pydcH<sub>2</sub> (3 mmol) and 2,3-dapy (1 mmol) was refluxed for about 30 min in a 1:3:1 molar ratio. Brown crystals of the title compound were obtained from the solution by slow evaporation of the solvent within two weeks at room temperature.

**Refinement**

H atoms attached to pyridine N and water O atoms were found in a difference Fourier map and refined with  $U_{iso}(H) = 1.0-1.5U_{eq}(N,O)$ . H atoms of the water molecules, H9A, H9B, H10A, H10B, H11A, H11B were refined with distance restraints of O—H = 0.81 (7), 0.84 (8), 0.94 (8), 0.80 (8), 0.89 (8) and 0.92 (9) Å and H $\cdots$ H distance restraints of 1.45 (4) Å for H10A $\cdots$ H10B and 1.40 (4) Å for H11A $\cdots$ H11B. H atoms on C atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 Å and  $U_{iso}(H) = 1.2U_{eq}(C)$ . The highest residual electron density was found at 0.83 Å from Bi1 atom and the deepest hole at 0.74 Å from Bi1 atom.

Figures




Fig. 1. The molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level. [Symmetry code: (i)  $-x+1, -y, -z+2$ .]




Fig. 2. The coordination environment of the  $\text{Bi}^{\text{III}}$  atom, showing a distorted bicapped trigonal-prismatic geometry. [Symmetry code: (i)  $-x+1, -y, -z+2$ .]




Fig. 3. A view of the crystal packing, showing  $\text{O—H}\cdots\text{O}$ ,  $\text{N—H}\cdots\text{O}$  and weak  $\text{C—H}\cdots\text{O}$  hydrogen bonds (dashed lines) in the title compound.




Fig. 4.  $\pi$ - $\pi$  stacking interactions between the aromatic rings of the  $(2,3\text{-dapyH})^+$  cation and pydc ligand.




Fig. 5.  $\text{C—H}\cdots\pi$  interaction between  $\text{C—H}$  group of the  $(2,3\text{-dapyH})^+$  cation and the aromatic ring of the pydc ligand.

**Bis(2,3-diaminopyridinium) bis( $\mu$ -pyridine-2,6-dicarboxylato)- $\kappa^4 O^2, N, O^6: O^6; \kappa^4 O^2: O^2, N, O^6$ -bis[aqua(pyridine-2,6-dicarboxylato- $\kappa^3 O^2, N, O^6$ )bismuthate(III)] tetrahydrate**

*Crystal data*

$(C_5H_8N_3)_2[Bi_2(C_7H_3NO_4)_4(H_2O)_2] \cdot 4H_2O$	$Z = 1$
$M_r = 1406.76$	$F(000) = 680$
Triclinic, $P\bar{1}$	$D_x = 2.109 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.3462 (19) \text{ \AA}$	Cell parameters from 5940 reflections
$b = 10.726 (2) \text{ \AA}$	$\theta = 2.2\text{--}29.2^\circ$
$c = 11.098 (2) \text{ \AA}$	$\mu = 8.03 \text{ mm}^{-1}$
$\alpha = 95.13 (3)^\circ$	$T = 298 \text{ K}$
$\beta = 91.38 (3)^\circ$	Prism, brown
$\gamma = 90.47 (3)^\circ$	$0.33 \times 0.27 \times 0.23 \text{ mm}$
$V = 1107.7 (4) \text{ \AA}^3$	

*Data collection*

Stoe IPDS-2 diffractometer	5940 independent reflections
Radiation source: fine-focus sealed tube graphite	5539 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.113$
Absorption correction: numerical ( <i>X-SHAPE</i> and <i>X-RED32</i> ; Stoe & Cie, 2005)	$\theta_{\text{max}} = 29.2^\circ$ , $\theta_{\text{min}} = 2.2^\circ$
$T_{\text{min}} = 0.083$ , $T_{\text{max}} = 0.156$	$h = -12 \rightarrow 12$
12406 measured reflections	$k = -14 \rightarrow 14$
	$l = -15 \rightarrow 15$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.154$	$w = 1/[\sigma^2(F_o^2) + (0.1042P)^2 + 3.0215P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
5940 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
347 parameters	$\Delta\rho_{\text{max}} = 2.98 \text{ e \AA}^{-3}$
9 restraints	$\Delta\rho_{\text{min}} = -2.93 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.0047 (13)

## supplementary materials

---

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O10	0.8002 (8)	0.0220 (8)	0.8803 (9)	0.0534 (19)
O8	-0.0239 (8)	0.3014 (6)	1.0095 (8)	0.056 (2)
O11	0.9124 (8)	0.8479 (8)	0.7003 (7)	0.0502 (17)
O6	0.6729 (7)	0.3001 (6)	0.7953 (8)	0.0494 (17)
Bi1	0.32379 (2)	0.045644 (19)	0.88303 (2)	0.02387 (13)
N1	0.4091 (7)	-0.0927 (6)	0.7035 (5)	0.0265 (11)
C5	0.3481 (8)	-0.0744 (7)	0.5968 (7)	0.0297 (14)
C1	0.5057 (7)	-0.1819 (6)	0.7123 (6)	0.0247 (12)
C3	0.4867 (12)	-0.2374 (10)	0.4998 (8)	0.047 (2)
H3	0.5131	-0.2871	0.4312	0.056*
C7	0.2415 (9)	0.0299 (7)	0.5988 (8)	0.0359 (16)
C2	0.5469 (10)	-0.2566 (9)	0.6098 (8)	0.0426 (19)
H2	0.6147	-0.3188	0.6163	0.051*
C4	0.3855 (10)	-0.1434 (9)	0.4899 (8)	0.0388 (17)
H4	0.3450	-0.1277	0.4155	0.047*
N4	1.1655 (8)	0.4867 (7)	0.2534 (8)	0.0436 (18)
H4A	1.1986	0.4137	0.2319	0.052*
H4B	1.2133	0.5526	0.2398	0.052*
C15	1.0421 (8)	0.4974 (7)	0.3068 (7)	0.0300 (14)
C16	0.9763 (9)	0.6147 (8)	0.3425 (8)	0.0333 (15)
N3	0.9707 (9)	0.3919 (7)	0.3263 (7)	0.0383 (15)
N5	1.0463 (9)	0.7230 (6)	0.3244 (9)	0.0462 (19)
H5A	1.0086	0.7940	0.3469	0.055*
H5B	1.1281	0.7205	0.2904	0.055*
C17	0.8460 (10)	0.6133 (9)	0.3960 (10)	0.043 (2)
H17	0.8021	0.6885	0.4204	0.051*
C19	0.8402 (11)	0.3898 (9)	0.3806 (9)	0.046 (2)
H19	0.7959	0.3144	0.3938	0.055*
C18	0.7782 (10)	0.5004 (11)	0.4144 (10)	0.048 (2)
H18	0.6892	0.5012	0.4502	0.057*
O3	0.2133 (7)	0.0852 (6)	0.7009 (6)	0.0381 (13)
O7	0.0896 (6)	0.1326 (5)	0.9346 (6)	0.0340 (12)
O5	0.5370 (6)	0.1332 (5)	0.8106 (6)	0.0324 (11)
N2	0.3143 (7)	0.2700 (6)	0.8827 (6)	0.0262 (11)
C8	0.4311 (8)	0.3335 (7)	0.8530 (7)	0.0283 (13)
O4	0.1860 (11)	0.0529 (9)	0.5024 (7)	0.066 (2)
C14	0.0788 (9)	0.2486 (7)	0.9591 (8)	0.0331 (15)
C13	0.5562 (8)	0.2518 (7)	0.8158 (7)	0.0290 (14)
C12	0.1996 (8)	0.3318 (7)	0.9215 (7)	0.0294 (14)
C9	0.4360 (8)	0.4623 (7)	0.8582 (8)	0.0321 (15)
H9	0.5173	0.5045	0.8362	0.039*
C11	0.1952 (8)	0.4614 (7)	0.9293 (8)	0.0324 (15)
H11	0.1135	0.5036	0.9554	0.039*
C10	0.3145 (9)	0.5268 (7)	0.8975 (9)	0.0358 (17)
H10	0.3137	0.6138	0.9023	0.043*

O1	0.5240 (6)	-0.1236 (5)	0.9223 (5)	0.0290 (10)
O2	0.6605 (9)	-0.2791 (7)	0.8471 (6)	0.0489 (17)
C6	0.5695 (8)	-0.1989 (7)	0.8366 (7)	0.0276 (13)
O9	0.1606 (8)	-0.1496 (7)	0.8422 (7)	0.0455 (15)
H10B	0.867 (9)	0.064 (12)	0.906 (13)	0.055*
H11A	0.851 (13)	0.845 (14)	0.638 (9)	0.068*
H11B	0.860 (15)	0.908 (13)	0.742 (11)	0.068*
H9A	0.143 (14)	-0.171 (15)	0.909 (8)	0.068*
H9B	0.090 (13)	-0.126 (14)	0.802 (10)	0.068*
H10A	0.717 (9)	0.070 (11)	0.875 (12)	0.055*
H3A	1.007 (14)	0.325 (13)	0.289 (12)	0.055*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O10	0.036 (3)	0.043 (3)	0.081 (6)	0.007 (3)	-0.001 (3)	0.005 (4)
O8	0.052 (4)	0.029 (3)	0.090 (6)	0.012 (3)	0.045 (4)	0.005 (3)
O11	0.049 (4)	0.054 (4)	0.049 (4)	0.022 (3)	-0.006 (3)	0.011 (3)
O6	0.034 (3)	0.034 (3)	0.080 (5)	-0.004 (2)	0.025 (3)	0.001 (3)
Bi1	0.02420 (17)	0.01863 (16)	0.02853 (18)	0.00371 (9)	0.00376 (9)	-0.00045 (9)
N1	0.030 (3)	0.026 (3)	0.024 (3)	0.004 (2)	0.004 (2)	0.000 (2)
C5	0.032 (3)	0.025 (3)	0.032 (4)	0.000 (3)	0.003 (3)	0.000 (3)
C1	0.025 (3)	0.022 (3)	0.026 (3)	0.002 (2)	0.003 (2)	-0.004 (2)
C3	0.060 (6)	0.049 (5)	0.030 (4)	0.014 (4)	0.012 (4)	-0.010 (4)
C7	0.039 (4)	0.028 (3)	0.042 (4)	0.006 (3)	0.004 (3)	0.006 (3)
C2	0.041 (4)	0.046 (5)	0.037 (4)	0.011 (4)	0.006 (3)	-0.015 (4)
C4	0.045 (4)	0.043 (4)	0.027 (4)	0.004 (3)	0.006 (3)	-0.003 (3)
N4	0.037 (4)	0.030 (3)	0.063 (5)	0.005 (3)	0.019 (3)	-0.004 (3)
C15	0.035 (4)	0.030 (3)	0.025 (3)	0.000 (3)	-0.001 (3)	0.002 (3)
C16	0.032 (4)	0.029 (3)	0.039 (4)	0.000 (3)	0.005 (3)	0.003 (3)
N3	0.045 (4)	0.027 (3)	0.043 (4)	-0.001 (3)	0.010 (3)	-0.002 (3)
N5	0.051 (4)	0.019 (3)	0.069 (5)	0.007 (3)	0.024 (4)	0.002 (3)
C17	0.037 (4)	0.036 (4)	0.055 (6)	0.012 (3)	0.013 (4)	0.002 (4)
C19	0.053 (5)	0.041 (4)	0.044 (5)	-0.009 (4)	0.013 (4)	0.003 (4)
C18	0.033 (4)	0.057 (6)	0.052 (5)	-0.011 (4)	0.015 (4)	-0.008 (4)
O3	0.044 (3)	0.034 (3)	0.036 (3)	0.018 (2)	-0.007 (2)	0.001 (2)
O7	0.029 (2)	0.024 (2)	0.048 (3)	-0.0006 (19)	0.007 (2)	0.004 (2)
O5	0.031 (3)	0.024 (2)	0.042 (3)	0.0096 (19)	0.014 (2)	0.002 (2)
N2	0.027 (3)	0.025 (3)	0.026 (3)	0.005 (2)	0.006 (2)	-0.002 (2)
C8	0.033 (3)	0.022 (3)	0.029 (3)	0.002 (2)	0.006 (3)	0.001 (2)
O4	0.082 (6)	0.073 (5)	0.042 (4)	0.035 (5)	-0.015 (4)	0.003 (4)
C14	0.033 (4)	0.024 (3)	0.043 (4)	0.005 (3)	0.008 (3)	0.008 (3)
C13	0.027 (3)	0.027 (3)	0.032 (3)	0.004 (2)	0.008 (3)	-0.001 (3)
C12	0.028 (3)	0.025 (3)	0.035 (4)	0.002 (2)	0.005 (3)	-0.003 (3)
C9	0.026 (3)	0.025 (3)	0.046 (4)	0.003 (2)	0.008 (3)	0.003 (3)
C11	0.028 (3)	0.026 (3)	0.043 (4)	0.006 (3)	0.007 (3)	-0.001 (3)
C10	0.036 (4)	0.023 (3)	0.048 (5)	0.007 (3)	0.006 (3)	0.000 (3)
O1	0.030 (2)	0.032 (2)	0.023 (2)	0.007 (2)	-0.0015 (19)	-0.0062 (19)



## supplementary materials

---

O2	0.067 (4)	0.040 (3)	0.039 (3)	0.029 (3)	-0.002 (3)	-0.002 (3)
C6	0.027 (3)	0.026 (3)	0.029 (3)	0.004 (2)	0.006 (3)	-0.002 (3)
O9	0.045 (3)	0.041 (3)	0.049 (4)	-0.012 (3)	0.010 (3)	-0.005 (3)

### *Geometric parameters (Å, °)*

O10—H10B	0.80 (8)	C15—C16	1.433 (11)
O10—H10A	0.94 (8)	C16—N5	1.362 (11)
O8—C14	1.239 (10)	C16—C17	1.368 (12)
O11—H11A	0.89 (8)	N3—C19	1.374 (12)
O11—H11B	0.92 (9)	N3—H3A	0.87 (14)
O6—C13	1.238 (9)	N5—H5A	0.8600
Bi1—O3	2.323 (6)	N5—H5B	0.8600
Bi1—O5	2.384 (6)	C17—C18	1.396 (15)
Bi1—N2	2.409 (6)	C17—H17	0.9300
Bi1—O7	2.447 (6)	C19—C18	1.351 (15)
Bi1—N1	2.525 (6)	C19—H19	0.9300
Bi1—O9	2.581 (7)	C18—H18	0.9300
Bi1—O1 <sup>i</sup>	2.627 (5)	O7—C14	1.255 (9)
Bi1—O1	2.673 (5)	O5—C13	1.279 (9)
N1—C1	1.329 (9)	N2—C12	1.325 (9)
N1—C5	1.333 (10)	N2—C8	1.346 (9)
C5—C4	1.394 (11)	C8—C9	1.377 (10)
C5—C7	1.504 (11)	C8—C13	1.510 (10)
C1—C2	1.395 (10)	C14—C12	1.522 (11)
C1—C6	1.516 (10)	C12—C11	1.386 (10)
C3—C2	1.364 (14)	C9—C10	1.394 (10)
C3—C4	1.398 (14)	C9—H9	0.9300
C3—H3	0.9300	C11—C10	1.381 (12)
C7—O4	1.224 (12)	C11—H11	0.9300
C7—O3	1.266 (11)	C10—H10	0.9300
C2—H2	0.9300	O1—C6	1.275 (9)
C4—H4	0.9300	O1—Bi1 <sup>i</sup>	2.627 (5)
N4—C15	1.310 (11)	O2—C6	1.225 (10)
N4—H4A	0.8600	O9—H9A	0.81 (7)
N4—H4B	0.8600	O9—H9B	0.84 (8)
C15—N3	1.346 (10)		
H10B—O10—H10A	112 (9)	N3—C15—C16	117.8 (7)
H11A—O11—H11B	91 (8)	N5—C16—C17	122.4 (8)
O3—Bi1—O5	87.5 (2)	N5—C16—C15	119.2 (7)
O3—Bi1—N2	73.7 (2)	C17—C16—C15	118.4 (8)
O5—Bi1—N2	67.26 (19)	C15—N3—C19	124.1 (8)
O3—Bi1—O7	74.0 (2)	C15—N3—H3A	113 (9)
O5—Bi1—O7	133.22 (18)	C19—N3—H3A	122 (9)
N2—Bi1—O7	66.40 (19)	C16—N5—H5A	120.0
O3—Bi1—N1	66.3 (2)	C16—N5—H5B	120.0
O5—Bi1—N1	70.7 (2)	H5A—N5—H5B	120.0
N2—Bi1—N1	122.1 (2)	C16—C17—C18	120.9 (8)

O7—Bi1—N1	132.5 (2)	C16—C17—H17	119.6
O3—Bi1—O9	79.0 (3)	C18—C17—H17	119.6
O5—Bi1—O9	140.1 (2)	C18—C19—N3	118.1 (9)
N2—Bi1—O9	140.2 (2)	C18—C19—H19	121.0
O7—Bi1—O9	78.7 (2)	N3—C19—H19	121.0
N1—Bi1—O9	69.4 (2)	C19—C18—C17	120.7 (8)
O3—Bi1—O1 <sup>i</sup>	150.1 (2)	C19—C18—H18	119.6
O5—Bi1—O1 <sup>i</sup>	74.6 (2)	C17—C18—H18	119.6
N2—Bi1—O1 <sup>i</sup>	77.3 (2)	C7—O3—Bi1	124.9 (5)
O7—Bi1—O1 <sup>i</sup>	100.97 (19)	C14—O7—Bi1	119.0 (5)
N1—Bi1—O1 <sup>i</sup>	126.40 (18)	C13—O5—Bi1	121.0 (4)
O9—Bi1—O1 <sup>i</sup>	129.7 (2)	C12—N2—C8	119.8 (6)
O3—Bi1—O1	128.33 (18)	C12—N2—Bi1	120.5 (5)
O5—Bi1—O1	76.04 (18)	C8—N2—Bi1	119.4 (5)
N2—Bi1—O1	136.45 (19)	N2—C8—C9	122.5 (7)
O7—Bi1—O1	147.79 (19)	N2—C8—C13	114.3 (6)
N1—Bi1—O1	61.99 (18)	C9—C8—C13	123.2 (7)
O9—Bi1—O1	83.3 (2)	O8—C14—O7	125.2 (8)
O1 <sup>i</sup> —Bi1—O1	70.87 (18)	O8—C14—C12	117.1 (7)
C1—N1—C5	120.5 (6)	O7—C14—C12	117.7 (7)
C1—N1—Bi1	123.4 (5)	O6—C13—O5	122.5 (7)
C5—N1—Bi1	116.0 (5)	O6—C13—C8	120.1 (7)
N1—C5—C4	122.3 (7)	O5—C13—C8	117.4 (6)
N1—C5—C7	115.4 (7)	N2—C12—C11	121.4 (7)
C4—C5—C7	122.3 (8)	N2—C12—C14	114.3 (6)
N1—C1—C2	120.7 (7)	C11—C12—C14	124.3 (7)
N1—C1—C6	117.8 (6)	C8—C9—C10	117.5 (7)
C2—C1—C6	121.5 (7)	C8—C9—H9	121.3
C2—C3—C4	120.3 (8)	C10—C9—H9	121.3
C2—C3—H3	119.8	C10—C11—C12	118.8 (7)
C4—C3—H3	119.8	C10—C11—H11	120.6
O4—C7—O3	124.9 (8)	C12—C11—H11	120.6
O4—C7—C5	117.8 (8)	C11—C10—C9	119.9 (7)
O3—C7—C5	117.3 (7)	C11—C10—H10	120.0
C3—C2—C1	119.3 (8)	C9—C10—H10	120.0
C3—C2—H2	120.3	C6—O1—Bi1 <sup>i</sup>	124.2 (5)
C1—C2—H2	120.3	C6—O1—Bi1	121.5 (5)
C5—C4—C3	116.9 (8)	Bi1 <sup>i</sup> —O1—Bi1	109.13 (18)
C5—C4—H4	121.6	O2—C6—O1	125.5 (7)
C3—C4—H4	121.6	O2—C6—C1	119.2 (7)
C15—N4—H4A	120.0	O1—C6—C1	115.2 (6)
C15—N4—H4B	120.0	Bi1—O9—H9A	105 (10)
H4A—N4—H4B	120.0	Bi1—O9—H9B	106 (10)
N4—C15—N3	118.1 (8)	H9A—O9—H9B	116 (10)
N4—C15—C16	124.0 (7)		
O3—Bi1—N1—C1	-178.5 (6)	O1 <sup>i</sup> —Bi1—O5—C13	-75.3 (6)
O5—Bi1—N1—C1	85.6 (6)	O1—Bi1—O5—C13	-148.9 (6)

## supplementary materials

---

N2—Bi1—N1—C1	131.0 (5)	O3—Bi1—N2—C12	86.6 (6)
O7—Bi1—N1—C1	-142.8 (5)	O5—Bi1—N2—C12	-179.2 (6)
O9—Bi1—N1—C1	-91.8 (6)	O7—Bi1—N2—C12	7.3 (6)
O1 <sup>i</sup> —Bi1—N1—C1	33.0 (6)	N1—Bi1—N2—C12	134.0 (6)
O1—Bi1—N1—C1	1.6 (5)	O9—Bi1—N2—C12	38.0 (7)
O3—Bi1—N1—C5	-0.5 (5)	O1 <sup>i</sup> —Bi1—N2—C12	-100.8 (6)
O5—Bi1—N1—C5	-96.4 (5)	O1—Bi1—N2—C12	-144.4 (5)
N2—Bi1—N1—C5	-51.0 (6)	O3—Bi1—N2—C8	-99.3 (6)
O7—Bi1—N1—C5	35.2 (6)	O5—Bi1—N2—C8	-5.1 (5)
O9—Bi1—N1—C5	86.2 (5)	O7—Bi1—N2—C8	-178.5 (6)
O1 <sup>i</sup> —Bi1—N1—C5	-149.0 (5)	N1—Bi1—N2—C8	-51.9 (6)
O1—Bi1—N1—C5	179.6 (6)	O9—Bi1—N2—C8	-147.8 (5)
C1—N1—C5—C4	-2.4 (11)	O1 <sup>i</sup> —Bi1—N2—C8	73.3 (5)
Bi1—N1—C5—C4	179.5 (6)	O1—Bi1—N2—C8	29.8 (7)
C1—N1—C5—C7	-179.6 (7)	C12—N2—C8—C9	-1.7 (12)
Bi1—N1—C5—C7	2.4 (8)	Bi1—N2—C8—C9	-175.9 (6)
C5—N1—C1—C2	1.1 (11)	C12—N2—C8—C13	177.3 (7)
Bi1—N1—C1—C2	179.0 (6)	Bi1—N2—C8—C13	3.1 (9)
C5—N1—C1—C6	-179.2 (6)	Bi1—O7—C14—O8	-164.6 (8)
Bi1—N1—C1—C6	-1.3 (9)	Bi1—O7—C14—C12	16.9 (10)
N1—C5—C7—O4	177.7 (9)	Bi1—O5—C13—O6	169.5 (7)
C4—C5—C7—O4	0.5 (13)	Bi1—O5—C13—C8	-8.4 (10)
N1—C5—C7—O3	-4.0 (11)	N2—C8—C13—O6	-174.6 (8)
C4—C5—C7—O3	178.8 (8)	C9—C8—C13—O6	4.4 (13)
C4—C3—C2—C1	0.4 (16)	N2—C8—C13—O5	3.3 (11)
N1—C1—C2—C3	-0.2 (14)	C9—C8—C13—O5	-177.7 (8)
C6—C1—C2—C3	-179.8 (9)	C8—N2—C12—C11	1.8 (12)
N1—C5—C4—C3	2.6 (13)	Bi1—N2—C12—C11	175.9 (6)
C7—C5—C4—C3	179.5 (9)	C8—N2—C12—C14	-176.5 (7)
C2—C3—C4—C5	-1.5 (15)	Bi1—N2—C12—C14	-2.4 (9)
N4—C15—C16—N5	-2.3 (13)	O8—C14—C12—N2	171.6 (9)
N3—C15—C16—N5	179.7 (9)	O7—C14—C12—N2	-9.8 (11)
N4—C15—C16—C17	179.0 (9)	O8—C14—C12—C11	-6.6 (14)
N3—C15—C16—C17	1.0 (12)	O7—C14—C12—C11	172.0 (8)
N4—C15—N3—C19	-179.6 (9)	N2—C8—C9—C10	0.8 (13)
C16—C15—N3—C19	-1.5 (13)	C13—C8—C9—C10	-178.1 (8)
N5—C16—C17—C18	-179.2 (11)	N2—C12—C11—C10	-1.0 (13)
C15—C16—C17—C18	-0.6 (15)	C14—C12—C11—C10	177.1 (8)
C15—N3—C19—C18	1.4 (15)	C12—C11—C10—C9	0.2 (13)
N3—C19—C18—C17	-0.9 (17)	C8—C9—C10—C11	-0.1 (13)
C16—C17—C18—C19	0.6 (17)	O3—Bi1—O1—C6	-2.0 (7)
O4—C7—O3—Bi1	-178.0 (9)	O5—Bi1—O1—C6	-77.3 (6)
C5—C7—O3—Bi1	3.8 (11)	N2—Bi1—O1—C6	-110.2 (6)
O5—Bi1—O3—C7	68.1 (7)	O7—Bi1—O1—C6	124.4 (6)
N2—Bi1—O3—C7	135.2 (8)	N1—Bi1—O1—C6	-1.9 (5)
O7—Bi1—O3—C7	-155.3 (8)	O9—Bi1—O1—C6	68.3 (6)
N1—Bi1—O3—C7	-1.9 (7)	O1 <sup>i</sup> —Bi1—O1—C6	-155.5 (7)
O9—Bi1—O3—C7	-74.1 (7)	O3—Bi1—O1—Bi1 <sup>i</sup>	153.5 (2)

O1 <sup>i</sup> —Bi1—O3—C7	120.6 (7)	O5—Bi1—O1—Bi1 <sup>i</sup>	78.3 (2)
O1—Bi1—O3—C7	-1.8 (8)	N2—Bi1—O1—Bi1 <sup>i</sup>	45.4 (4)
O3—Bi1—O7—C14	-91.9 (7)	O7—Bi1—O1—Bi1 <sup>i</sup>	-80.0 (4)
O5—Bi1—O7—C14	-21.3 (8)	N1—Bi1—O1—Bi1 <sup>i</sup>	153.6 (3)
N2—Bi1—O7—C14	-13.0 (6)	O9—Bi1—O1—Bi1 <sup>i</sup>	-136.2 (3)
N1—Bi1—O7—C14	-125.7 (6)	O1 <sup>i</sup> —Bi1—O1—Bi1 <sup>i</sup>	0.0
O9—Bi1—O7—C14	-173.6 (7)	Bi1 <sup>i</sup> —O1—C6—O2	28.4 (11)
O1 <sup>i</sup> —Bi1—O7—C14	57.8 (7)	Bi1—O1—C6—O2	-179.9 (7)
O1—Bi1—O7—C14	129.2 (6)	Bi1 <sup>i</sup> —O1—C6—C1	-149.8 (5)
O3—Bi1—O5—C13	80.5 (6)	Bi1—O1—C6—C1	2.0 (8)
N2—Bi1—O5—C13	7.2 (6)	N1—C1—C6—O2	-178.8 (8)
O7—Bi1—O5—C13	15.4 (7)	C2—C1—C6—O2	0.9 (12)
N1—Bi1—O5—C13	146.3 (7)	N1—C1—C6—O1	-0.6 (10)
O9—Bi1—O5—C13	150.0 (6)	C2—C1—C6—O1	179.1 (8)

Symmetry codes: (i)  $-x+1, -y, -z+2$ .

*Hydrogen-bond geometry* ( $\text{\AA}, ^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N3—H3A $\cdots$ O11 <sup>ii</sup>	0.87 (14)	2.02 (14)	2.797 (10)	148 (12)
N4—H4A $\cdots$ O2 <sup>iii</sup>	0.86	2.11	2.924 (9)	158
N4—H4B $\cdots$ O6 <sup>ii</sup>	0.86	1.97	2.830 (10)	178
N5—H5A $\cdots$ O3 <sup>iv</sup>	0.86	2.53	3.216 (10)	138
N5—H5B $\cdots$ O6 <sup>ii</sup>	0.86	2.11	2.972 (10)	176
O9—H9A $\cdots$ O8 <sup>v</sup>	0.81 (7)	2.07 (12)	2.746 (10)	141 (15)
O9—H9B $\cdots$ O11 <sup>vi</sup>	0.84 (8)	1.99 (10)	2.771 (11)	155 (15)
O10—H10A $\cdots$ O5	0.94 (8)	1.96 (8)	2.862 (9)	160 (11)
O10—H10B $\cdots$ O7 <sup>vii</sup>	0.80 (8)	2.21 (8)	2.972 (10)	160 (13)
O11—H11A $\cdots$ O4 <sup>iv</sup>	0.89 (8)	2.01 (13)	2.717 (11)	136 (14)
O11—H11B $\cdots$ O10 <sup>viii</sup>	0.92 (9)	1.96 (9)	2.836 (12)	158 (13)
C11—H11 $\cdots$ O8 <sup>ix</sup>	0.93	2.26	3.050 (10)	142

Symmetry codes: (ii)  $-x+2, -y+1, -z+1$ ; (iii)  $-x+2, -y, -z+1$ ; (iv)  $-x+1, -y+1, -z+1$ ; (v)  $-x, -y, -z+2$ ; (vi)  $x-1, y-1, z$ ; (vii)  $x+1, y, z$ ; (viii)  $x, y+1, z$ ; (ix)  $-x, -y+1, -z+2$ .

Fig. 1




Fig. 2




Fig. 3




Fig. 4






Fig. 5

