

Bis[4-amino-N-(4-methylpyrimidin-2-yl- κN^3)benzenesulfonamido- κN](2,2'-bi-pyridine- $\kappa^2 N,N'$)mercury(II)

G. M. Golzar Hossain^{a*} and A. J. Amoroso^b

^aDepartment of Chemistry, University of Dhaka, Dhaka 1000, Bangladesh, and

^bSchool of Chemistry, Cardiff University, Cardiff CF10 3AT, Wales

Correspondence e-mail: acsbd@yahoo.com


Received 24 February 2014; accepted 2 March 2014

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.039; wR factor = 0.074; data-to-parameter ratio = 16.8.

The complete molecule of the title complex, $[\text{Hg}(\text{C}_{11}\text{H}_{11}\text{N}_4\text{O}_2\text{S})_2(\text{C}_{10}\text{H}_8\text{N}_2)]$, is generated by crystallographic twofold symmetry, with the mercury cation lying on the rotation axis. The mercury coordination polyhedron can be described as tetrahedral (from the N,N' -bidentate bi-pyridine molecule and the sulfonamide N atoms of the sulfamerazine anions) or as squashed trigonal-prismatic, if two long ($> 2.80\text{ \AA}$) $\text{Hg}-\text{N}$ bonds to pyrimidine N atoms are included. The dihedral angle between the aromatic rings in the anion is $73.3(2)^\circ$. In the crystal, $\text{N}-\text{H}\cdots(\text{N},\text{O})$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds link the molecules into a three-dimensional network.

Related literature

For related structures, see: Garcia-Raso *et al.* (1997, 2000); Saladini *et al.* (2001); Zamora *et al.* (1982); Hergold-Brundić *et al.* (1989). For ligand conformations, see: Hossain & Amoroso (2007, 2012); Hossain *et al.* (2007).



Experimental

Crystal data

$[\text{Hg}(\text{C}_{11}\text{H}_{11}\text{N}_4\text{O}_2\text{S})_2(\text{C}_{10}\text{H}_8\text{N}_2)]$	$V = 3371.4(3)\text{ \AA}^3$
$M_r = 883.37$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 18.7483(8)\text{ \AA}$	$\mu = 4.74\text{ mm}^{-1}$
$b = 15.0824(7)\text{ \AA}$	$T = 150\text{ K}$
$c = 12.1143(6)\text{ \AA}$	$0.12 \times 0.12 \times 0.10\text{ mm}$
$\beta = 100.202(2)^\circ$	

Data collection

Nonius KappaCCD diffractometer	29910 measured reflections
Absorption correction: multi-scan (<i>SORTAV</i> ; Blessing, 1995)]	3874 independent reflections
$T_{\min} = 0.600$, $T_{\max} = 0.648$	3408 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.072$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.074$	$\Delta\rho_{\max} = 1.13\text{ e \AA}^{-3}$
$S = 1.10$	$\Delta\rho_{\min} = -0.75\text{ e \AA}^{-3}$
3874 reflections	
231 parameters	
3 restraints	

Table 1
Selected bond lengths (\AA).

$\text{Hg1}-\text{N11}$	2.214 (4)	$\text{Hg1}-\text{N12}$	2.883 (3)
$\text{Hg1}-\text{N1}$	2.322 (3)		

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N14}-\text{H14A}\cdots\text{O11}^i$	0.90 (1)	2.41 (3)	3.173 (5)	143 (4)
$\text{N14}-\text{H14A}\cdots\text{N13}^i$	0.90 (1)	2.42 (3)	3.125 (6)	135 (4)
$\text{N14}-\text{H14B}\cdots\text{O11}^{ii}$	0.90 (1)	2.13 (1)	2.997 (5)	163 (4)

Symmetry codes: (i) $x, -y + 1, z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT* (Hooft, 1998); cell refinement: *DENZO* and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012).

GMGH is grateful to the Ministry of Science and Technology, Bangladesh, for financial support and to School of Chemistry, Cardiff University, Wales, for the crystallographic services.

Supporting information for this paper is available from the IUCr electronic archives (Reference: HB7201).

References

- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Garcia-Raso, A., Fiol, J. J., Martorell, G., Lopez-Zafra, A. & Quiros, M. (1997). *Polyhedron*, **16**, 613–621.
- Garcia-Raso, A., Fiol, J. J., Rigo, S., Lopez-Lopez, A., Molins, E., Espinosa, E., Borras, E., Alzuet, G., Borras, J. & Castineiras, A. (2000). *Polyhedron*, **19**, 991–1004.
- Hergold-Brundić, A., Kamenar, B. & Jovanovski, G. (1989). *Acta Cryst. C* **45**, 556–558.
- Hooft, R. W. W. (1998). COLLECT. Nonius BV, Delft, The Netherlands.
- Hossain, G. M. G. & Amoroso, A. J. (2007). *Acta Cryst. E* **63**, m759–m760.
- Hossain, G. M. G. & Amoroso, A. J. (2012). *Acta Cryst. E* **68**, m276–m277.
- Hossain, G. M. G., Amoroso, A. J., Banu, A. & Malik, K. M. A. (2007). *Polyhedron*, **23**, 967–974.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Saladini, M., Menabue, L., Ferrari, E. & Iacopino, D. (2001). *J. Chem. Soc. Dalton Trans.* pp. 1513–1519.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zamora, F., Sabat, M. & Lippert, B. (1982). *Inorg. Chim. Acta*, **282**, 237–242.

supporting information

Acta Cryst. (2014). E70, m127–m128 [doi:10.1107/S1600536814004760]

Bis[4-amino-N-(4-methylpyrimidin-2-yl- κ N³)benzenesulfonamido- κ N](2,2'-bipyridine- κ^2 N,N')mercury(II)

G. M. Golzar Hossain and A. J. Amoroso

S1. Comment

The mercury complexes of some related ligands have been reported (Garcia-Raso *et al.*, 1997; Hossain *et al.*, 2007). In the title complex, sulfamerazine behaves as a bidentate anionic ligand and Hg(II) ion is coordinated *via* sulfonamidic N(11) and pyrimido N(12) atoms, the fifth and sixth coordination sites are occupied by bipyridine molecule. The geometry of the complex is trigonal prismatic with the Hg atom lying on a twofold axis as the bpy molecule lies on the one part and the two larger sulfamerazine molecules lie on the other part.

The Hg—N(11) bond distance of 2.216 (3) Å is slightly longer than the values of 2.087 (4) Å (Garcia-Raso *et al.*, 1997), 2.071 (4) Å (Garcia-Raso *et al.*, 2000) and 2.14 (2) Å (Saladini *et al.*, 2001). The pyrimido nitrogen [N(12)] atom makes bond with longer interaction with the distance of 2.881 (3) Å. The C(18)—N(14) bond distance of 1.373 (5) Å is in good agreement with the corresponding bond in free sulfamerazine suggesting the terminal amino group is not coordinated with the Hg atoms. The Hg—N(1) bond distance of 2.323 (3) Å is longer than the corresponding bond lengths of 2.140 (3) and 2.124 (3) Å (Zamora *et al.*, 1982) and 2.130 Å (Hergold-Brundić *et al.*, 1989).

The bond angles around the S atom correspond to a distorted tetrahedral geometry. The bond distance C(18)—N(14) (Terminal amino group) of 1.371 (5) Å and the torsion angle C(15)—S(11)—N(11)—C(11) of 74.4 (4)° are larger than those observed in the related structures (Hossain & Amoroso, 2007; Hossain & Amoroso, 2012). The dihedral angle between the aromatic rings of the sulfamerazinate anion of 73.27 (13)° is larger than the value of 71.10 (14)° (Hossain & Amoroso, 2007) and smaller than the value of 76.60 (8)° (Hossain & Amoroso, 2012) in the sulfadiazinate anion.


The packing of the complex unit (Fig. 2) is governed by weak (intermolecular N—H···N, and N—H···O) hydrogen bonds (Table 2) between the amino (—NH₂) group of one complex unit and the sulfonyl oxygen and the pyrimido nitrogen atoms of the next units.

S2. Experimental


The complex was obtained by dissolving (2.723 g, 2 mmol) sulfamerazine in 50 ml of methanol followed by drop-wise addition of Hg(CH₃COO)₂·4H₂O (1.245 g, 1 mmol) in water with constant stirring on hot plate for 1 h. A white precipitate was formed, filtered and washed with methanol and dried in a desiccator over silica gel. The precipitate was dissolved in DMF, bipyri dine (0.235 g, 1 mmol) was added to this solution and stirred for 30 minutes. The solution was filtered and left for crystallization. A week later, colourless blocks were obtained.

S3. Refinement

The H atoms were positioned geometrically and refined using a riding model except that for terminal amino group N(14) which were located from the difference map and fixed to 0.900 (3) Å, with C—H = 0.95–0.98 Å and with $U_{\text{iso}}(\text{H}) = 1.2$ (1.5 for methyl groups) times $U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of (I), with 50% probability displacement ellipsoids for non-H atoms.

**Figure 2**

The packing of (I), viewed down the a -axis, showing one layer of molecules connected by $\text{N}—\text{H}\cdots\text{N}$ and $\text{N}—\text{H}\cdots\text{O}$ hydrogen bonds (dashed lines).

Bis[4-amino-N-(4-methylpyrimidin-2-yl- κN^3)benzenesulfonamido- κN](2,2'-bipyridine- $\kappa^2\text{N},\text{N}'$)mercury(II)

Crystal data

$[\text{Hg}(\text{C}_{11}\text{H}_{11}\text{N}_4\text{O}_2\text{S})_2(\text{C}_{10}\text{H}_8\text{N}_2)]$

$M_r = 883.37$

Monoclinic, $C2/c$

$a = 18.7483$ (8) Å

$b = 15.0824$ (7) Å

$c = 12.1143$ (6) Å

$\beta = 100.202$ (2)°

$V = 3371.4$ (3) Å³

$Z = 4$

$F(000) = 1744$

$D_x = 1.740 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3874 reflections

$\theta = 2.9\text{--}27.5$ °

$\mu = 4.74 \text{ mm}^{-1}$

$T = 150$ K

Block, colorless

0.12 × 0.12 × 0.10 mm

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube

ω scans

Absorption correction: multi-scan
(*SORTAV*; Blessing, 1995)]

$T_{\min} = 0.600$, $T_{\max} = 0.648$

29910 measured reflections

3874 independent reflections

3408 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.072$

$\theta_{\max} = 27.5$ °, $\theta_{\min} = 3.2$ °

$h = -24\text{--}24$

$k = -19\text{--}19$

$l = -15\text{--}15$

Refinement

Refinement on F^2

$wR(F^2) = 0.074$

Least-squares matrix: full

$S = 1.10$

$R[F^2 > 2\sigma(F^2)] = 0.039$

3874 reflections

231 parameters

3 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0108P)^2 + 13.7149P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 1.13 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.75 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.0000	0.15586 (2)	0.2500	0.02712 (9)
S11	0.14988 (6)	0.26641 (7)	0.20322 (9)	0.0241 (2)
O11	0.19225 (16)	0.26047 (19)	0.1139 (2)	0.0300 (7)
O12	0.17100 (16)	0.20881 (18)	0.2977 (2)	0.0298 (7)
N11	0.0655 (2)	0.2427 (2)	0.1595 (3)	0.0250 (8)
N12	-0.0384 (2)	0.2333 (2)	0.0300 (3)	0.0272 (8)
N13	0.0520 (2)	0.3400 (2)	0.0030 (3)	0.0321 (9)
N14	0.1869 (2)	0.6308 (2)	0.3883 (3)	0.0297 (9)
C11	0.0260 (2)	0.2745 (3)	0.0597 (3)	0.0236 (9)
C12	-0.0808 (2)	0.2607 (3)	-0.0646 (4)	0.0316 (10)
C13	-0.0589 (3)	0.3277 (3)	-0.1283 (4)	0.0382 (12)
H13	-0.0889	0.3474	-0.1954	0.046*
C14	0.0076 (3)	0.3648 (3)	-0.0915 (4)	0.0390 (12)
H14	0.0233	0.4107	-0.1353	0.047*
C15	0.1581 (2)	0.3761 (3)	0.2533 (3)	0.0228 (9)
C16	0.1290 (3)	0.3989 (3)	0.3461 (4)	0.0304 (10)
H16	0.1012	0.3566	0.3783	0.036*
C17	0.1398 (3)	0.4825 (3)	0.3930 (4)	0.0292 (10)
H17	0.1204	0.4965	0.4581	0.035*
C18	0.1790 (2)	0.5466 (3)	0.3453 (3)	0.0254 (9)
C19	0.2096 (2)	0.5225 (3)	0.2518 (4)	0.0290 (10)
H19	0.2379	0.5644	0.2197	0.035*
C20	0.1989 (2)	0.4381 (3)	0.2058 (4)	0.0289 (10)
H20	0.2193	0.4226	0.1421	0.035*
C21	-0.1540 (3)	0.2183 (4)	-0.0972 (5)	0.0514 (15)
H21A	-0.1481	0.1554	-0.1135	0.077*
H21B	-0.1807	0.2478	-0.1641	0.077*
H21C	-0.1811	0.2241	-0.0354	0.077*
N1	-0.0594 (2)	0.0298 (2)	0.1713 (3)	0.0302 (8)
C1	-0.0330 (2)	-0.0493 (3)	0.2060 (3)	0.0275 (10)
C2	-0.0665 (3)	-0.1265 (3)	0.1612 (4)	0.0381 (12)
H2	-0.0474	-0.1826	0.1872	0.046*
C3	-0.1274 (3)	-0.1219 (3)	0.0792 (4)	0.0435 (13)
H3	-0.1510	-0.1744	0.0485	0.052*

C4	-0.1535 (3)	-0.0397 (3)	0.0424 (4)	0.0423 (12)
H4	-0.1944	-0.0344	-0.0160	0.051*
C5	-0.1189 (3)	0.0351 (3)	0.0922 (4)	0.0400 (12)
H5	-0.1379	0.0919	0.0696	0.048*
H14A	0.171 (2)	0.644 (3)	0.452 (2)	0.038 (14)*
H14B	0.2224 (18)	0.666 (2)	0.373 (4)	0.040 (14)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hg1	0.02493 (15)	0.02381 (12)	0.02946 (13)	0.000	-0.00387 (9)	0.000
S11	0.0190 (6)	0.0238 (5)	0.0276 (5)	0.0009 (4)	-0.0009 (4)	-0.0021 (4)
O11	0.0231 (18)	0.0329 (16)	0.0338 (16)	-0.0007 (13)	0.0046 (14)	-0.0080 (14)
O12	0.0218 (17)	0.0271 (15)	0.0366 (17)	0.0026 (13)	-0.0053 (14)	0.0021 (13)
N11	0.025 (2)	0.0279 (18)	0.0192 (16)	0.0003 (15)	-0.0048 (15)	-0.0001 (15)
N12	0.017 (2)	0.0334 (19)	0.0283 (19)	0.0025 (15)	-0.0028 (15)	-0.0003 (16)
N13	0.025 (2)	0.039 (2)	0.0312 (19)	-0.0006 (18)	0.0009 (16)	0.0055 (18)
N14	0.030 (2)	0.0259 (19)	0.033 (2)	-0.0040 (16)	0.0056 (18)	-0.0031 (16)
C11	0.017 (2)	0.028 (2)	0.025 (2)	0.0039 (17)	0.0028 (17)	-0.0006 (18)
C12	0.018 (2)	0.041 (3)	0.033 (2)	0.0021 (19)	-0.0034 (19)	-0.002 (2)
C13	0.027 (3)	0.053 (3)	0.032 (2)	0.004 (2)	-0.004 (2)	0.006 (2)
C14	0.031 (3)	0.049 (3)	0.034 (3)	0.002 (2)	0.002 (2)	0.014 (2)
C15	0.018 (2)	0.0217 (19)	0.026 (2)	-0.0015 (16)	-0.0024 (17)	-0.0016 (17)
C16	0.028 (3)	0.031 (2)	0.034 (2)	-0.0052 (19)	0.010 (2)	0.002 (2)
C17	0.033 (3)	0.031 (2)	0.026 (2)	-0.0009 (19)	0.011 (2)	-0.0022 (19)
C18	0.017 (2)	0.030 (2)	0.026 (2)	-0.0009 (17)	-0.0057 (17)	0.0007 (18)
C19	0.026 (3)	0.028 (2)	0.033 (2)	-0.0038 (19)	0.006 (2)	0.0015 (19)
C20	0.026 (3)	0.032 (2)	0.029 (2)	-0.0007 (19)	0.0073 (19)	-0.0029 (19)
C21	0.023 (3)	0.076 (4)	0.047 (3)	-0.005 (3)	-0.014 (2)	0.020 (3)
N1	0.027 (2)	0.0289 (18)	0.0315 (19)	-0.0013 (16)	-0.0045 (17)	-0.0004 (16)
C1	0.025 (3)	0.030 (2)	0.028 (2)	0.0036 (18)	0.0033 (19)	0.0007 (19)
C2	0.042 (3)	0.025 (2)	0.046 (3)	-0.001 (2)	0.001 (2)	-0.007 (2)
C3	0.043 (3)	0.036 (3)	0.048 (3)	-0.009 (2)	-0.002 (3)	-0.015 (2)
C4	0.035 (3)	0.046 (3)	0.039 (3)	-0.008 (2)	-0.011 (2)	-0.010 (2)
C5	0.034 (3)	0.037 (3)	0.044 (3)	0.002 (2)	-0.010 (2)	0.001 (2)

Geometric parameters (\AA , $^\circ$)

Hg1—N11	2.214 (4)	C15—C20	1.394 (6)
Hg1—N11 ⁱ	2.214 (4)	C16—C17	1.383 (6)
Hg1—N1	2.322 (3)	C16—H16	0.9500
Hg1—N1 ⁱ	2.322 (3)	C17—C18	1.400 (6)
Hg1—N12 ⁱ	2.883 (3)	C17—H17	0.9500
Hg1—N12	2.883 (3)	C18—C19	1.405 (6)
S11—O12	1.436 (3)	C19—C20	1.390 (6)
S11—O11	1.454 (3)	C19—H19	0.9500
S11—N11	1.616 (4)	C20—H20	0.9500
S11—C15	1.760 (4)	C21—H21A	0.9800

N11—C11	1.387 (5)	C21—H21B	0.9800
N12—C12	1.339 (5)	C21—H21C	0.9800
N12—C11	1.349 (5)	N1—C1	1.331 (5)
N13—C11	1.343 (6)	N1—C5	1.337 (6)
N13—C14	1.344 (6)	C1—C2	1.387 (6)
N14—C18	1.371 (5)	C1—C1 ⁱ	1.483 (9)
N14—H14A	0.900 (3)	C2—C3	1.376 (7)
N14—H14B	0.900 (3)	C2—H2	0.9500
C12—C13	1.378 (7)	C3—C4	1.378 (7)
C12—C21	1.503 (7)	C3—H3	0.9500
C13—C14	1.367 (7)	C4—C5	1.386 (6)
C13—H13	0.9500	C4—H4	0.9500
C14—H14	0.9500	C5—H5	0.9500
C15—C16	1.378 (6)		
N11—Hg1—N11 ⁱ	107.50 (18)	C16—C15—C20	119.7 (4)
N11—Hg1—N1	123.31 (12)	C16—C15—S11	119.5 (3)
N11 ⁱ —Hg1—N1	114.81 (13)	C20—C15—S11	120.6 (3)
N11—Hg1—N1 ⁱ	114.81 (13)	C15—C16—C17	120.8 (4)
N11 ⁱ —Hg1—N1 ⁱ	123.31 (12)	C15—C16—H16	119.6
N1—Hg1—N1 ⁱ	70.00 (17)	C17—C16—H16	119.6
N11—Hg1—N12 ⁱ	98.53 (11)	C16—C17—C18	120.7 (4)
N11 ⁱ —Hg1—N12 ⁱ	51.11 (11)	C16—C17—H17	119.7
N1—Hg1—N12 ⁱ	137.29 (11)	C18—C17—H17	119.7
N1 ⁱ —Hg1—N12 ⁱ	85.95 (11)	N14—C18—C17	120.8 (4)
N11—Hg1—N12	51.11 (11)	N14—C18—C19	120.9 (4)
N11 ⁱ —Hg1—N12	98.53 (11)	C17—C18—C19	118.3 (4)
N1—Hg1—N12	85.95 (11)	C20—C19—C18	120.6 (4)
N1 ⁱ —Hg1—N12	137.29 (11)	C20—C19—H19	119.7
N12 ⁱ —Hg1—N12	132.17 (14)	C18—C19—H19	119.7
O12—S11—O11	116.49 (18)	C19—C20—C15	120.0 (4)
O12—S11—N11	104.05 (18)	C19—C20—H20	120.0
O11—S11—N11	111.99 (18)	C15—C20—H20	120.0
O12—S11—C15	107.41 (18)	C12—C21—H21A	109.5
O11—S11—C15	106.76 (19)	C12—C21—H21B	109.5
N11—S11—C15	110.01 (19)	H21A—C21—H21B	109.5
C11—N11—S11	123.3 (3)	C12—C21—H21C	109.5
C11—N11—Hg1	112.3 (3)	H21A—C21—H21C	109.5
S11—N11—Hg1	124.42 (18)	H21B—C21—H21C	109.5
C12—N12—C11	117.0 (4)	C1—N1—C5	119.8 (4)
C12—N12—Hg1	158.4 (3)	C1—N1—Hg1	118.7 (3)
C11—N12—Hg1	82.8 (2)	C5—N1—Hg1	121.6 (3)
C11—N13—C14	114.5 (4)	N1—C1—C2	120.7 (4)
C18—N14—H14A	120 (3)	N1—C1—C1 ⁱ	116.3 (2)
C18—N14—H14B	120 (3)	C2—C1—C1 ⁱ	122.9 (3)
H14A—N14—H14B	115 (3)	C3—C2—C1	120.0 (4)
N13—C11—N12	126.2 (4)	C3—C2—H2	120.0
N13—C11—N11	121.0 (4)	C1—C2—H2	120.0

N12—C11—N11	112.8 (4)	C2—C3—C4	118.8 (4)
N12—C12—C13	121.0 (4)	C2—C3—H3	120.6
N12—C12—C21	118.1 (4)	C4—C3—H3	120.6
C13—C12—C21	120.9 (4)	C3—C4—C5	118.6 (4)
C14—C13—C12	117.6 (4)	C3—C4—H4	120.7
C14—C13—H13	121.2	C5—C4—H4	120.7
C12—C13—H13	121.2	N1—C5—C4	122.0 (4)
N13—C14—C13	123.8 (5)	N1—C5—H5	119.0
N13—C14—H14	118.1	C4—C5—H5	119.0
C13—C14—H14	118.1		

Symmetry code: (i) $-x, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N14—H14A \cdots O11 ⁱⁱ	0.90 (1)	2.41 (3)	3.173 (5)	143 (4)
N14—H14A \cdots N13 ⁱⁱ	0.90 (1)	2.42 (3)	3.125 (6)	135 (4)
N14—H14B \cdots O11 ⁱⁱⁱ	0.90 (1)	2.13 (1)	2.997 (5)	163 (4)

Symmetry codes: (ii) $x, -y+1, z+1/2$; (iii) $-x+1/2, y+1/2, -z+1/2$.