

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

# Bis(guanidinium) bis(4-hydroxypyridine-2,6-dicarboxylato- $\kappa^3 O^2, N, O^6$ )nickel-ate(II) dihydrate

Hossein Aghabozorg,<sup>a\*</sup> Elham Motyeian,<sup>a</sup> Jafar Attar Gharamaleki,<sup>a</sup> Janet Soleimannejad,<sup>b</sup> Mohammad Ghadermazi<sup>c</sup> and Elizabeth Spey Sharon<sup>d</sup>

<sup>a</sup>Faculty of Chemistry, Teacher Training University, Tehran, Iran, <sup>b</sup>Faculty of Science, Department of Chemistry, Ilam University, Iran, <sup>c</sup>Department of Chemistry, Faculty of Science, University of Kurdistan, Sanandaj, Iran, and <sup>d</sup>Department of Chemistry, Sheffield University, Sheffield S3 7HF, England

Correspondence e-mail: haghabozorg@yahoo.com


Received 5 November 2007; accepted 2 December 2007

Key indicators: single-crystal X-ray study;  $T = 150$  K; mean  $\sigma(C-C) = 0.001$  Å;  $R$  factor = 0.026;  $wR$  factor = 0.076; data-to-parameter ratio = 31.4.

The reaction of nickel(II) nitrate hexahydrate, guanidine (G) and 4-hydroxypyridine-2,6-dicarboxylic acid (hypydcH<sub>2</sub>) in a 1:2:2 molar ratio in aqueous solution resulted in the formation of the title compound, (CH<sub>5</sub>N<sub>3</sub>)<sub>2</sub>[Ni(C<sub>7</sub>H<sub>3</sub>NO<sub>5</sub>)<sub>2</sub>].2H<sub>2</sub>O or (GH)<sub>2</sub>[Ni(hypydc)<sub>2</sub>].2H<sub>2</sub>O. The six donor atoms of the two 4-hydroxypyridine-2,6-dicarboxylate or (hypydc)<sup>2-</sup> ligands form a distorted octahedral arrangement around the Ni<sup>II</sup> centre. Considerable C—O... $\pi$  stacking interactions between the CO groups of carboxylate fragments and the pyridine rings of (hypydc)<sup>2-</sup> with a distance of 3.3212 (8) Å are observed. In the crystal structure, a wide range of noncovalent interactions consisting of hydrogen bonding (of the types O—H...O and N—H...O), ion pairing, and  $\pi$ - $\pi$  [centroid-centroid distance 3.8037 (5) Å], N—H... $\pi$  and C—O... $\pi$  stacking interactions connect the various components into a supramolecular structure.

## Related literature

For related literature, see: Aghabozorg, Attar Gharamaleki *et al.* (2007a, 2007b); Aghabozorg, Daneshvar *et al.* (2007).



## Experimental

### Crystal data

(CH<sub>5</sub>N<sub>3</sub>)<sub>2</sub>[Ni(C<sub>7</sub>H<sub>3</sub>NO<sub>5</sub>)<sub>2</sub>].2H<sub>2</sub>O  
 $M_r = 577.13$   
 Triclinic,  $P\bar{1}$   
 $a = 8.9253$  (6) Å  
 $b = 9.9060$  (6) Å  
 $c = 13.2186$  (9) Å  
 $\alpha = 101.415$  (3)°  
 $\beta = 103.099$  (3)°

$\gamma = 91.938$  (3)°  
 $V = 1111.99$  (13) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.96$  mm<sup>-1</sup>  
 $T = 150$  (2) K  
 $0.50 \times 0.32 \times 0.15$  mm

### Data collection

Bruker SMART 1000 diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 1998)  
 $T_{\min} = 0.647$ ,  $T_{\max} = 0.870$

46516 measured reflections  
 10497 independent reflections  
 9650 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.019$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.076$   
 $S = 1.06$   
 10497 reflections

334 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.56$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.61$  e Å<sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

|            |            |            |            |
|------------|------------|------------|------------|
| Ni1—N2     | 1.9665 (6) | Ni1—O10    | 2.1387 (6) |
| Ni1—N1     | 1.9700 (6) | Ni1—O6     | 2.1477 (6) |
| Ni1—O1     | 2.1088 (6) | Ni1—O5     | 2.2149 (6) |
|            |            |            |            |
| N2—Ni1—N1  | 171.30 (3) | O1—Ni1—O10 | 93.41 (3)  |
| O1—Ni1—O5  | 154.97 (2) | O6—Ni1—O5  | 91.30 (2)  |
| O10—Ni1—O6 | 154.87 (2) |            |            |

**Table 2**

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the N1/C2—C6 ring.

| $D-H\cdots A$                | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|------------------------------|-------|-------------|-------------|---------------|
| O3—H3C...O6 <sup>i</sup>     | 0.90  | 1.71        | 2.5901 (8)  | 166           |
| O8—H8...O5 <sup>ii</sup>     | 0.90  | 1.84        | 2.7057 (9)  | 160           |
| O11—H11A...O9 <sup>iii</sup> | 0.90  | 1.86        | 2.7106 (9)  | 156           |
| O11—H11B...O2 <sup>iv</sup>  | 0.90  | 1.88        | 2.7464 (9)  | 161           |
| O12—H12B...O11 <sup>ii</sup> | 0.90  | 2.15        | 3.0425 (12) | 172           |
| O12—H12A...O5                | 0.90  | 2.15        | 3.0244 (13) | 164           |

| $D-H\cdots A$                      | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|------------------------------------|-------|-------------|-------------|---------------|
| O12–H12A $\cdots$ O4               | 0.90  | 2.64        | 3.3823 (12) | 141           |
| N3–H3A $\cdots$ O4 <sup>v</sup>    | 0.90  | 2.15        | 2.9307 (10) | 144           |
| N3–H3B $\cdots$ O10                | 0.90  | 2.07        | 2.8817 (10) | 149           |
| N4–H4A $\cdots$ O4 <sup>v</sup>    | 0.90  | 1.97        | 2.8072 (11) | 154           |
| N4–H4B $\cdots$ O11 <sup>vi</sup>  | 0.90  | 2.06        | 2.9161 (10) | 159           |
| N5–H5A $\cdots$ O10                | 0.90  | 2.19        | 2.9887 (10) | 148           |
| N5–H5B $\cdots$ O2 <sup>ii</sup>   | 0.90  | 1.98        | 2.8656 (10) | 169           |
| N6–H6A $\cdots$ O7 <sup>viii</sup> | 0.90  | 2.28        | 3.0178 (11) | 139           |
| N6–H6A $\cdots$ O1                 | 0.90  | 2.49        | 3.0038 (9)  | 117           |
| N6–H6B $\cdots$ O3 <sup>vii</sup>  | 0.90  | 2.55        | 3.2320 (11) | 133           |
| N7–H7A $\cdots$ O9 <sup>iii</sup>  | 0.90  | 2.01        | 2.8999 (10) | 172           |
| N7–H7B $\cdots$ O12 <sup>ix</sup>  | 0.90  | 1.93        | 2.8282 (13) | 176           |
| N8–H8A $\cdots$ O7 <sup>viii</sup> | 0.90  | 1.94        | 2.7993 (10) | 158           |
| N8–H8B $\cdots$ O11                | 0.90  | 1.99        | 2.8847 (10) | 174           |
| N5–H5A $\cdots$ Cg1                | 0.90  | 3.47        | 3.362 (7)   | 75            |

Symmetry codes: (i)  $-x, -y + 1, -z$ ; (ii)  $-x + 1, -y + 1, -z + 1$ ; (iii)  $-x + 1, -y, -z + 1$ ; (iv)  $-x, -y, -z + 1$ ; (v)  $-x + 1, -y + 1, -z$ ; (vi)  $x, y, z - 1$ ; (vii)  $-x, -y, -z$ ; (viii)  $-x, -y + 1, -z + 1$ ; (ix)  $x, y - 1, z$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINTE* (Bruker, 1998); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Bruker, 2005) and *Mercury* (Version 1.4.2; Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXL97*.

Financial support from Ilam University and the Teacher Training University, Tehran, is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2189).

## References

- Aghabozorg, H., Attar Gharamaleki, J., Ghadermazi, M., Ghasemikhah, P. & Soleimannejad, J. (2007a). *Acta Cryst.* **E63**, m1803–m1804.
- Aghabozorg, H., Attar Gharamaleki, J., Ghasemikhah, P., Ghadermazi, M. & Soleimannejad, J. (2007b). *Acta Cryst.* **E63**, m1710–m1711.
- Aghabozorg, H., Daneshvar, S., Moteyean, E., Ghadermazi, M. & Attar Gharamaleki, J. (2007). *Acta Cryst.* **E63**, m2468–m2469.
- Bruker (1998). *SADABS* (Version 2004/1), *SAINTE* (Version 6.01) and *SMART* (Version 5.059). Bruker AXS Inc., Madison, Wisconsin.
- Bruker (2005). *SHELXTL*. Version 6.10. Bruker AXS Inc., Madison, Wisconsin, USA.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.

## supporting information

*Acta Cryst.* (2008). E64, m144–m145 [https://doi.org/10.1107/S1600536807065026]

**Bis(guanidinium) bis(4-hydroxypyridine-2,6-dicarboxylato- $\kappa^3O^2,N,O^6$ )nickelate(II) dihydrate**

**Hossein Aghabozorg, Elham Motyeian, Jafar Attar Gharamaleki, Janet Soleimannejad, Mohammad Ghadermazi and Elizabeth Spey Sharon**

**S1. Comment**

The non-covalent interactions such as ion pairing, hydrogen bonding and  $\pi$ - $\pi$  stacking are observed in these ionic compounds. The importance of weak hydrogen bonds in the context of crystal engineering, molecular recognition and supramolecular chemistry has been well recognized in recent years. Recently, we have defined a plan to prepare water soluble proton transfer compounds as novel self assembled systems that can function as suitable ligands in the synthesis of metal complexes. In this regard, we have reported cases in which proton transfers from pyridine-2,6-dicarboxylic acid, pydcH<sub>2</sub>, and benzene-1,2,4,5-tetracarboxylic acid, btcH<sub>4</sub>, to propane-1,3-diamine (pn) and 1,10-phenanthroline, (phen), resulted in the formation of novel self-assembled (pnH<sub>2</sub>)(pydc)(pydcH<sub>2</sub>) $\cdot$ 2.5H<sub>2</sub>O, (pnH<sub>2</sub>)<sub>2</sub>(btc)2H<sub>2</sub>O and (phenH)<sub>4</sub>(btcH<sub>3</sub>)<sub>2</sub>(btcH<sub>2</sub>) systems, respectively. The resulting compounds with some remaining sites as electron donors can coordinate to metallic ions (Aghabozorg, Attar Gharamaleki *et al.*, 2007a,b; Aghabozorg, Daneshvar *et al.*, 2007).

The molecular structure of the title compound is presented in Fig. 1. The Ni<sup>II</sup> atom is six-coordinated by two 4-hydroxypyridine-2,6-dicarboxylate, or (hypydc)<sup>2-</sup>, groups, *i.e.* each (hypydc)<sup>2-</sup> is coordinated through one pyridine N atom and two carboxylate O atoms (Table 1). N1 and N2 atoms of the two (hypydc)<sup>2-</sup> fragments occupy the axial positions, while atoms O1, O5, O6 and O10 form the equatorial plane [with Ni—O distances ranging from 2.1088 (6) to 2.2150 (6) Å]. The N1—Ni1—N2 angle [171.30 (3)°] deviates from linearity. Therefore, the coordination around Ni<sup>II</sup> is distorted octahedral. The O5—Ni1—O6 and O1—Ni1—O10 angles are equal to 91.30 (2) and 93.41 (3)°, respectively. On the other hand, O1—Ni1—O6—C8 and O10—Ni1—O5—C7 torsion angles are 93.20 (5) and 92.27 (5) Å, respectively indicating that two (hypydc)<sup>2-</sup> units are almost perpendicular to each other.

A noticeable feature of the title compound is the presence of N—H $\cdots$  $\pi$  stacking interactions between N—H group of guanidinium ions with aromatic rings of (hypydc)<sup>2-</sup> units. The H $\cdots$  $\pi$  distance (measured to the centre of phenyl ring) is 3.475 Å for N5—H5A $\cdots$ Cg1 [Cg1 is the centroid of N1/C2—C6 ring]. There is also  $\pi$  $\cdots$  $\pi$  stacking interaction between two aromatic rings of (hypydc)<sup>2-</sup> units, with distance of 3.8037 (5) Å [-x, 1 - y, -z] (Fig. 2).


Also a considerable C—O $\cdots$  $\pi$  stacking interactions between CO groups of carboxylate fragments with aromatic rings of 4-hydroxypyridine-2,6-dicarboxylate with distances of 3.321 (8) Å for C8—O7 $\cdots$ Cg2 (-x, 1 - y, 1 - z) [Cg2 is the centroid for N2/C9—C13 ring] are observed in the prepared compound (Fig. 3). In the crystal structure, a wide range of non-covalent interactions consisting of hydrogen bonding (of the type of O—H $\cdots$ O and N—H $\cdots$ O with D $\cdots$ A ranging from 2.5901 (8) Å to 3.3823 (12) Å), ion pairing,  $\pi$  $\cdots$  $\pi$ , N—H $\cdots$  $\pi$  and C—O $\cdots$  $\pi$  stacking connect the various components into a supramolecular structure (Table 2, Figs. 4 and 5).

## S2. Experimental

An aqueous solution of sodium hydroxide (80 mg, 2 mmol) was added to guanidine hydrochloride (200 mg, 2 mmol) in a 1:1 molar ratio. The resulting suspension was stirred for 1 h and filtered. An aqueous solution of  $\text{Ni}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  (290 mg, 1 mmol) and 4-hydroxypyridine-2,6-dicarboxylic (360 mg, 2 mmol) acid was added to the filtered solution in a 1:2 molar ratio, and the reaction mixture was heated to boiling point for 2 h. Green crystals were obtained from the solution after two days at room temperature.

## S3. Refinement

Hydrogen atoms were positioned geometrically and refined with a riding model with  $\text{O}-\text{H} = 0.95 \text{ \AA}$ , and with  $U(\text{H})$  constrained to be 1.2 times  $U_{\text{eq}}$  of the carrier atom.



**Figure 1**

The structure of the title compound showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

$\pi\cdots\pi$  Stacking interaction between two aromatic rings of  $(\text{hypydc})^{2-}$  units, with distance of 3.8037 (5) Å  $[-x, 1 - y, -z]$ ; N—H $\cdots\pi$  stacking interaction between N—H groups of guanidinium ions with aromatic rings of  $(\text{hypydc})^{2-}$  units. The H $\cdots\pi$  distance (measured to the centre of phenyl ring) is 2.96 Å for N5—H5A $\cdots$ Cg1 [Cg1 is the centroid of N1/C2—C6 ring].






Figure 3

C—O $\cdots\pi$  stacking interactions between CO groups of carboxylate fragments with aromatic rings of 4-hydroxypyridine-2,6-dicarboxylate with distances of 3.321 (8) Å for C8—O7 $\cdots$ Cg2  $(-x, 1 - y, 1 - z)$  [Cg2 is the centroid for N2/C9—C13 ring].



**Figure 4**

A layered packing diagram. The space between the two layers of  $[\text{Ni}(\text{hypydc})_2]^{2-}$  fragments is filled with a layer of  $(\text{GH})^+$  cations and water molecules.



**Figure 5**

The crystal packing of the title compound view down the  $a$  axis; hydrogen bonds are shown as dashed lines.

**Bis(guanidinium) bis(4-hydroxypyridine-2,6-dicarboxylato- $\kappa^3\text{O}^2, \text{N}, \text{O}^6$ )nickelate(II) dihydrate**

*Crystal data*

$(\text{CH}_6\text{N}_3)_2[\text{Ni}(\text{C}_7\text{H}_3\text{NO}_5)_2] \cdot 2\text{H}_2\text{O}$

$M_r = 577.13$

Triclinic,  $P\bar{1}$

$a = 8.9253(6) \text{ \AA}$

$b = 9.9060(6) \text{ \AA}$

$c = 13.2186(9) \text{ \AA}$

$\alpha = 101.415 (3)^\circ$   
 $\beta = 103.099 (3)^\circ$   
 $\gamma = 91.938 (3)^\circ$   
 $V = 1111.99 (13) \text{ \AA}^3$   
 $Z = 2$   
 $F(000) = 596$   
 $D_x = 1.724 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 Cell parameters from 9699 reflections  
 $\theta = 2.4\text{--}36.3^\circ$   
 $\mu = 0.96 \text{ mm}^{-1}$   
 $T = 150 \text{ K}$   
 Block, green  
 $0.50 \times 0.32 \times 0.15 \text{ mm}$

*Data collection*

Bruker SMART 1000  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 Detector resolution: 8.3 pixels  $\text{mm}^{-1}$   
 $\omega$  scans  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 1998)  
 $T_{\min} = 0.647$ ,  $T_{\max} = 0.870$

46516 measured reflections  
 10497 independent reflections  
 9650 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.019$   
 $\theta_{\max} = 36.4^\circ$ ,  $\theta_{\min} = 1.6^\circ$   
 $h = -14 \rightarrow 11$   
 $k = -16 \rightarrow 16$   
 $l = -21 \rightarrow 21$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.076$   
 $S = 1.06$   
 10497 reflections  
 334 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.043P)^2 + 0.2702P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.002$   
 $\Delta\rho_{\max} = 0.56 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.61 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | <i>x</i>      | <i>y</i>      | <i>z</i>     | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|---------------|---------------|--------------|----------------------------------|
| Ni1 | 0.173117 (11) | 0.423258 (10) | 0.259400 (7) | 0.01287 (3)                      |
| O1  | -0.00784 (7)  | 0.27245 (6)   | 0.24384 (5)  | 0.01774 (10)                     |
| O2  | -0.16765 (9)  | 0.11004 (7)   | 0.12138 (5)  | 0.02439 (13)                     |
| O3  | -0.02586 (9)  | 0.20971 (7)   | -0.21698 (5) | 0.02167 (12)                     |
| H3C | -0.0206       | 0.2745        | -0.2554      | 0.026*                           |
| O4  | 0.35691 (8)   | 0.60277 (7)   | 0.04763 (5)  | 0.02381 (12)                     |
| O5  | 0.31707 (7)   | 0.56247 (6)   | 0.20082 (5)  | 0.01687 (10)                     |
| O6  | 0.05225 (7)   | 0.59263 (6)   | 0.32052 (4)  | 0.01635 (10)                     |
| O7  | 0.02582 (8)   | 0.72029 (7)   | 0.47531 (5)  | 0.02065 (11)                     |

|      |              |              |              |              |
|------|--------------|--------------|--------------|--------------|
| O8   | 0.37838 (8)  | 0.46857 (7)  | 0.73456 (5)  | 0.02054 (11) |
| H8   | 0.4719       | 0.4367       | 0.7525       | 0.025*       |
| O9   | 0.51375 (8)  | 0.19318 (7)  | 0.39007 (5)  | 0.02143 (11) |
| O10  | 0.34554 (7)  | 0.27853 (6)  | 0.26981 (5)  | 0.01768 (10) |
| O11  | 0.32536 (8)  | -0.04310 (7) | 0.73709 (5)  | 0.02032 (11) |
| H11A | 0.3814       | -0.1076      | 0.7089       | 0.024*       |
| H11B | 0.2653       | -0.0814      | 0.7719       | 0.024*       |
| O12  | 0.37899 (13) | 0.87354 (12) | 0.25296 (8)  | 0.0491 (3)   |
| H12B | 0.4704       | 0.9157       | 0.2539       | 0.059*       |
| H12A | 0.3664       | 0.7820       | 0.2257       | 0.059*       |
| N1   | 0.09970 (7)  | 0.36856 (6)  | 0.10372 (5)  | 0.01309 (10) |
| N2   | 0.25270 (7)  | 0.44832 (6)  | 0.41390 (5)  | 0.01274 (10) |
| N3   | 0.50285 (9)  | 0.28895 (8)  | 0.10335 (6)  | 0.02072 (13) |
| H3A  | 0.5659       | 0.3475       | 0.0847       | 0.025*       |
| H3B  | 0.4860       | 0.3064       | 0.1691       | 0.025*       |
| N4   | 0.43549 (11) | 0.16824 (9)  | -0.06920 (7) | 0.02808 (17) |
| H4A  | 0.5014       | 0.2288       | -0.0833      | 0.034*       |
| H4B  | 0.3907       | 0.0927       | -0.1182      | 0.034*       |
| N5   | 0.32885 (10) | 0.09767 (8)  | 0.05716 (6)  | 0.02313 (14) |
| H5A  | 0.3226       | 0.1189       | 0.1254       | 0.028*       |
| H5B  | 0.2719       | 0.0285       | 0.0071       | 0.028*       |
| N6   | 0.12908 (10) | 0.08178 (8)  | 0.38427 (6)  | 0.02258 (13) |
| H6A  | 0.0636       | 0.1481       | 0.3920       | 0.027*       |
| H6B  | 0.1274       | 0.0351       | 0.3182       | 0.027*       |
| N7   | 0.29526 (10) | -0.06930 (9) | 0.45012 (7)  | 0.02577 (15) |
| H7A  | 0.3483       | -0.1044      | 0.5041       | 0.031*       |
| H7B  | 0.3174       | -0.0873      | 0.3859       | 0.031*       |
| N8   | 0.18342 (10) | 0.08088 (8)  | 0.56283 (6)  | 0.02222 (13) |
| H8A  | 0.1141       | 0.1442       | 0.5680       | 0.027*       |
| H8B  | 0.2214       | 0.0388       | 0.6166       | 0.027*       |
| C1   | -0.06666 (9) | 0.20926 (8)  | 0.14912 (6)  | 0.01563 (12) |
| C2   | -0.00601 (8) | 0.26094 (7)  | 0.06391 (6)  | 0.01391 (11) |
| C3   | -0.04990 (9) | 0.20506 (8)  | -0.04397 (6) | 0.01656 (12) |
| H3   | -0.1226      | 0.1295       | -0.0705      | 0.020*       |
| C4   | 0.01756 (9)  | 0.26469 (8)  | -0.11267 (6) | 0.01546 (12) |
| C5   | 0.12882 (9)  | 0.37751 (8)  | -0.06979 (6) | 0.01465 (11) |
| H5   | 0.1754       | 0.4190       | -0.1134      | 0.018*       |
| C6   | 0.16685 (8)  | 0.42490 (7)  | 0.03947 (5)  | 0.01292 (11) |
| C7   | 0.28988 (9)  | 0.53963 (8)  | 0.09859 (6)  | 0.01487 (11) |
| C8   | 0.08025 (9)  | 0.62575 (7)  | 0.42266 (6)  | 0.01421 (11) |
| C9   | 0.19209 (8)  | 0.53834 (7)  | 0.47982 (5)  | 0.01287 (11) |
| C10  | 0.23152 (9)  | 0.54827 (8)  | 0.58878 (6)  | 0.01447 (11) |
| H10  | 0.1879       | 0.6108       | 0.6338       | 0.017*       |
| C11  | 0.33960 (9)  | 0.46069 (8)  | 0.62872 (6)  | 0.01443 (11) |
| C12  | 0.40380 (9)  | 0.36772 (8)  | 0.55896 (6)  | 0.01472 (11) |
| H12  | 0.4767       | 0.3100       | 0.5843       | 0.018*       |
| C13  | 0.35558 (8)  | 0.36443 (7)  | 0.45134 (6)  | 0.01300 (11) |
| C14  | 0.41057 (9)  | 0.26960 (7)  | 0.36476 (6)  | 0.01460 (11) |



|     |             |             |             |              |
|-----|-------------|-------------|-------------|--------------|
| C15 | 0.42237 (9) | 0.18398 (8) | 0.03028 (6) | 0.01729 (12) |
| C16 | 0.20223 (9) | 0.02991 (8) | 0.46651 (7) | 0.01741 (12) |

*Atomic displacement parameters (Å<sup>2</sup>)*

|     | $U^{11}$    | $U^{22}$    | $U^{33}$    | $U^{12}$      | $U^{13}$      | $U^{23}$     |
|-----|-------------|-------------|-------------|---------------|---------------|--------------|
| Ni1 | 0.01507 (5) | 0.01471 (4) | 0.00908 (4) | 0.00167 (3)   | 0.00287 (3)   | 0.00302 (3)  |
| O1  | 0.0216 (3)  | 0.0205 (2)  | 0.0116 (2)  | -0.0011 (2)   | 0.00473 (19)  | 0.00417 (18) |
| O2  | 0.0293 (3)  | 0.0251 (3)  | 0.0181 (3)  | -0.0102 (2)   | 0.0079 (2)    | 0.0024 (2)   |
| O3  | 0.0347 (3)  | 0.0192 (2)  | 0.0091 (2)  | -0.0019 (2)   | 0.0017 (2)    | 0.00284 (18) |
| O4  | 0.0263 (3)  | 0.0261 (3)  | 0.0201 (3)  | -0.0075 (2)   | 0.0076 (2)    | 0.0064 (2)   |
| O5  | 0.0173 (2)  | 0.0194 (2)  | 0.0125 (2)  | -0.00028 (18) | 0.00219 (18)  | 0.00193 (18) |
| O6  | 0.0193 (2)  | 0.0184 (2)  | 0.0119 (2)  | 0.00462 (19)  | 0.00323 (18)  | 0.00461 (18) |
| O7  | 0.0252 (3)  | 0.0201 (3)  | 0.0179 (2)  | 0.0094 (2)    | 0.0069 (2)    | 0.0034 (2)   |
| O8  | 0.0218 (3)  | 0.0283 (3)  | 0.0092 (2)  | 0.0049 (2)    | -0.00023 (19) | 0.0026 (2)   |
| O9  | 0.0243 (3)  | 0.0229 (3)  | 0.0208 (3)  | 0.0117 (2)    | 0.0086 (2)    | 0.0079 (2)   |
| O10 | 0.0230 (3)  | 0.0196 (2)  | 0.0121 (2)  | 0.0061 (2)    | 0.00636 (19)  | 0.00398 (18) |
| O11 | 0.0262 (3)  | 0.0188 (2)  | 0.0177 (2)  | 0.0029 (2)    | 0.0082 (2)    | 0.0044 (2)   |
| O12 | 0.0487 (5)  | 0.0521 (6)  | 0.0417 (5)  | -0.0196 (5)   | 0.0255 (4)    | -0.0149 (4)  |
| N1  | 0.0141 (2)  | 0.0150 (2)  | 0.0107 (2)  | 0.00126 (19)  | 0.00302 (18)  | 0.00398 (18) |
| N2  | 0.0140 (2)  | 0.0139 (2)  | 0.0106 (2)  | 0.00212 (18)  | 0.00304 (18)  | 0.00285 (18) |
| N3  | 0.0222 (3)  | 0.0198 (3)  | 0.0177 (3)  | -0.0040 (2)   | 0.0059 (2)    | -0.0021 (2)  |
| N4  | 0.0392 (4)  | 0.0245 (3)  | 0.0186 (3)  | -0.0094 (3)   | 0.0127 (3)    | -0.0046 (3)  |
| N5  | 0.0268 (3)  | 0.0208 (3)  | 0.0212 (3)  | -0.0058 (3)   | 0.0068 (3)    | 0.0031 (2)   |
| N6  | 0.0266 (3)  | 0.0242 (3)  | 0.0192 (3)  | 0.0068 (3)    | 0.0045 (3)    | 0.0103 (3)   |
| N7  | 0.0300 (4)  | 0.0283 (4)  | 0.0263 (4)  | 0.0142 (3)    | 0.0130 (3)    | 0.0135 (3)   |
| N8  | 0.0262 (3)  | 0.0229 (3)  | 0.0187 (3)  | 0.0076 (3)    | 0.0047 (3)    | 0.0068 (2)   |
| C1  | 0.0180 (3)  | 0.0167 (3)  | 0.0136 (3)  | 0.0009 (2)    | 0.0053 (2)    | 0.0047 (2)   |
| C2  | 0.0154 (3)  | 0.0152 (3)  | 0.0115 (3)  | 0.0008 (2)    | 0.0029 (2)    | 0.0041 (2)   |
| C3  | 0.0202 (3)  | 0.0167 (3)  | 0.0118 (3)  | -0.0015 (2)   | 0.0020 (2)    | 0.0035 (2)   |
| C4  | 0.0205 (3)  | 0.0156 (3)  | 0.0098 (2)  | 0.0018 (2)    | 0.0020 (2)    | 0.0034 (2)   |
| C5  | 0.0179 (3)  | 0.0161 (3)  | 0.0109 (2)  | 0.0018 (2)    | 0.0038 (2)    | 0.0046 (2)   |
| C6  | 0.0137 (3)  | 0.0148 (3)  | 0.0109 (2)  | 0.0017 (2)    | 0.0030 (2)    | 0.0041 (2)   |
| C7  | 0.0148 (3)  | 0.0157 (3)  | 0.0144 (3)  | 0.0010 (2)    | 0.0036 (2)    | 0.0036 (2)   |
| C8  | 0.0150 (3)  | 0.0149 (3)  | 0.0135 (3)  | 0.0021 (2)    | 0.0036 (2)    | 0.0043 (2)   |
| C9  | 0.0140 (3)  | 0.0136 (2)  | 0.0109 (2)  | 0.0013 (2)    | 0.0028 (2)    | 0.0023 (2)   |
| C10 | 0.0159 (3)  | 0.0160 (3)  | 0.0106 (2)  | 0.0012 (2)    | 0.0024 (2)    | 0.0016 (2)   |
| C11 | 0.0151 (3)  | 0.0171 (3)  | 0.0098 (2)  | -0.0002 (2)   | 0.0010 (2)    | 0.0023 (2)   |
| C12 | 0.0154 (3)  | 0.0163 (3)  | 0.0122 (3)  | 0.0022 (2)    | 0.0016 (2)    | 0.0040 (2)   |
| C13 | 0.0140 (3)  | 0.0137 (3)  | 0.0117 (2)  | 0.0017 (2)    | 0.0033 (2)    | 0.0032 (2)   |
| C14 | 0.0168 (3)  | 0.0146 (3)  | 0.0140 (3)  | 0.0027 (2)    | 0.0059 (2)    | 0.0039 (2)   |
| C15 | 0.0180 (3)  | 0.0152 (3)  | 0.0180 (3)  | 0.0007 (2)    | 0.0051 (2)    | 0.0011 (2)   |
| C16 | 0.0171 (3)  | 0.0176 (3)  | 0.0197 (3)  | 0.0023 (2)    | 0.0051 (2)    | 0.0082 (2)   |

*Geometric parameters (Å, °)*

|        |            |        |        |
|--------|------------|--------|--------|
| Ni1—N2 | 1.9665 (6) | N4—H4A | 0.9000 |
| Ni1—N1 | 1.9700 (6) | N4—H4B | 0.9000 |

|            |             |          |             |
|------------|-------------|----------|-------------|
| Ni1—O1     | 2.1088 (6)  | N5—C15   | 1.3267 (11) |
| Ni1—O10    | 2.1387 (6)  | N5—H5A   | 0.9002      |
| Ni1—O6     | 2.1477 (6)  | N5—H5B   | 0.9000      |
| Ni1—O5     | 2.2149 (6)  | N6—C16   | 1.3394 (10) |
| O1—C1      | 1.2668 (9)  | N6—H6A   | 0.9000      |
| O2—C1      | 1.2492 (10) | N6—H6B   | 0.9001      |
| O3—C4      | 1.3394 (9)  | N7—C16   | 1.3247 (11) |
| O3—H3C     | 0.9000      | N7—H7A   | 0.9000      |
| O4—C7      | 1.2357 (9)  | N7—H7B   | 0.8999      |
| O5—C7      | 1.2895 (9)  | N8—C16   | 1.3238 (11) |
| O6—C8      | 1.2880 (9)  | N8—H8A   | 0.9000      |
| O7—C8      | 1.2343 (10) | N8—H8B   | 0.9002      |
| O8—C11     | 1.3485 (9)  | C1—C2    | 1.5198 (10) |
| O8—H8      | 0.9000      | C2—C3    | 1.3832 (10) |
| O9—C14     | 1.2413 (9)  | C3—C4    | 1.4044 (11) |
| O10—C14    | 1.2796 (9)  | C3—H3    | 0.9300      |
| O11—H11A   | 0.9000      | C4—C5    | 1.4055 (11) |
| O11—H11B   | 0.9001      | C5—C6    | 1.3857 (10) |
| O12—H12B   | 0.9000      | C5—H5    | 0.9300      |
| O12—H12A   | 0.9000      | C6—C7    | 1.5111 (10) |
| N1—C6      | 1.3371 (9)  | C8—C9    | 1.5141 (10) |
| N1—C2      | 1.3374 (9)  | C9—C10   | 1.3856 (10) |
| N2—C9      | 1.3363 (9)  | C10—C11  | 1.4032 (11) |
| N2—C13     | 1.3368 (9)  | C10—H10  | 0.9300      |
| N3—C15     | 1.3296 (10) | C11—C12  | 1.4021 (11) |
| N3—H3A     | 0.9001      | C12—C13  | 1.3833 (10) |
| N3—H3B     | 0.9001      | C12—H12  | 0.9300      |
| N4—C15     | 1.3256 (11) | C13—C14  | 1.5143 (10) |
|            |             |          |             |
| N2—Ni1—N1  | 171.30 (3)  | O2—C1—C2 | 118.69 (7)  |
| O1—Ni1—O5  | 154.97 (2)  | O1—C1—C2 | 115.64 (6)  |
| O10—Ni1—O6 | 154.87 (2)  | N1—C2—C3 | 121.56 (7)  |
| N2—Ni1—O1  | 96.84 (2)   | N1—C2—C1 | 112.83 (6)  |
| N1—Ni1—O1  | 78.70 (2)   | C3—C2—C1 | 125.60 (7)  |
| N2—Ni1—O10 | 77.95 (2)   | C2—C3—C4 | 118.69 (7)  |
| N1—Ni1—O10 | 94.76 (2)   | C2—C3—H3 | 120.7       |
| O1—Ni1—O10 | 93.41 (3)   | C4—C3—H3 | 120.7       |
| N2—Ni1—O6  | 77.41 (2)   | O3—C4—C3 | 118.15 (7)  |
| N1—Ni1—O6  | 110.20 (2)  | O3—C4—C5 | 122.66 (7)  |
| O1—Ni1—O6  | 94.31 (2)   | C3—C4—C5 | 119.18 (7)  |
| N2—Ni1—O5  | 108.19 (2)  | C6—C5—C4 | 117.89 (6)  |
| N1—Ni1—O5  | 76.46 (2)   | C6—C5—H5 | 121.1       |
| O10—Ni1—O5 | 91.73 (2)   | C4—C5—H5 | 121.1       |
| O6—Ni1—O5  | 91.30 (2)   | N1—C6—C5 | 122.25 (7)  |
| C1—O1—Ni1  | 114.35 (5)  | N1—C6—C7 | 113.16 (6)  |
| C4—O3—H3C  | 111.0       | C5—C6—C7 | 124.55 (6)  |
| C7—O5—Ni1  | 113.28 (5)  | O4—C7—O5 | 125.12 (7)  |
| C8—O6—Ni1  | 114.63 (5)  | O4—C7—C6 | 119.22 (7)  |

|                |             |              |             |
|----------------|-------------|--------------|-------------|
| C11—O8—H8      | 109.2       | O5—C7—C6     | 115.66 (6)  |
| C14—O10—Ni1    | 114.35 (5)  | O7—C8—O6     | 126.10 (7)  |
| H11A—O11—H11B  | 109.4       | O7—C8—C9     | 119.17 (7)  |
| H12B—O12—H12A  | 115.5       | O6—C8—C9     | 114.73 (6)  |
| C6—N1—C2       | 120.41 (6)  | N2—C9—C10    | 121.93 (7)  |
| C6—N1—Ni1      | 121.23 (5)  | N2—C9—C8     | 112.93 (6)  |
| C2—N1—Ni1      | 117.81 (5)  | C10—C9—C8    | 125.14 (7)  |
| C9—N2—C13      | 120.78 (6)  | C9—C10—C11   | 117.64 (7)  |
| C9—N2—Ni1      | 119.82 (5)  | C9—C10—H10   | 121.2       |
| C13—N2—Ni1     | 119.03 (5)  | C11—C10—H10  | 121.2       |
| C15—N3—H3A     | 119.6       | O8—C11—C12   | 121.73 (7)  |
| C15—N3—H3B     | 119.6       | O8—C11—C10   | 118.30 (7)  |
| H3A—N3—H3B     | 120.4       | C12—C11—C10  | 119.96 (6)  |
| C15—N4—H4A     | 117.6       | C13—C12—C11  | 118.05 (7)  |
| C15—N4—H4B     | 120.5       | C13—C12—H12  | 121.0       |
| H4A—N4—H4B     | 121.3       | C11—C12—H12  | 121.0       |
| C15—N5—H5A     | 115.0       | N2—C13—C12   | 121.62 (7)  |
| C15—N5—H5B     | 119.8       | N2—C13—C14   | 113.12 (6)  |
| H5A—N5—H5B     | 125.0       | C12—C13—C14  | 125.25 (7)  |
| C16—N6—H6A     | 121.8       | O9—C14—O10   | 125.71 (7)  |
| C16—N6—H6B     | 118.1       | O9—C14—C13   | 119.11 (7)  |
| H6A—N6—H6B     | 118.4       | O10—C14—C13  | 115.17 (6)  |
| C16—N7—H7A     | 121.3       | N4—C15—N5    | 120.87 (8)  |
| C16—N7—H7B     | 116.5       | N4—C15—N3    | 119.19 (8)  |
| H7A—N7—H7B     | 121.1       | N5—C15—N3    | 119.93 (8)  |
| C16—N8—H8A     | 116.9       | N8—C16—N7    | 121.25 (8)  |
| C16—N8—H8B     | 119.8       | N8—C16—N6    | 119.57 (8)  |
| H8A—N8—H8B     | 121.8       | N7—C16—N6    | 119.16 (8)  |
| O2—C1—O1       | 125.67 (7)  |              |             |
|                |             |              |             |
| N2—Ni1—O1—C1   | -165.80 (6) | O2—C1—C2—C3  | -1.87 (12)  |
| N1—Ni1—O1—C1   | 6.63 (6)    | O1—C1—C2—C3  | 177.90 (8)  |
| O10—Ni1—O1—C1  | -87.54 (6)  | N1—C2—C3—C4  | -0.79 (12)  |
| O6—Ni1—O1—C1   | 116.39 (6)  | C1—C2—C3—C4  | -179.78 (7) |
| O5—Ni1—O1—C1   | 13.95 (10)  | C2—C3—C4—O3  | -179.99 (7) |
| N2—Ni1—O5—C7   | 170.16 (5)  | C2—C3—C4—C5  | 0.95 (12)   |
| N1—Ni1—O5—C7   | -2.20 (5)   | O3—C4—C5—C6  | -179.04 (7) |
| O1—Ni1—O5—C7   | -9.58 (9)   | C3—C4—C5—C6  | -0.02 (11)  |
| O10—Ni1—O5—C7  | 92.27 (5)   | C2—N1—C6—C5  | 1.36 (11)   |
| O6—Ni1—O5—C7   | -112.67 (5) | Ni1—N1—C6—C5 | 172.68 (5)  |
| N2—Ni1—O6—C8   | -2.86 (5)   | C2—N1—C6—C7  | -176.55 (6) |
| N1—Ni1—O6—C8   | 172.73 (5)  | Ni1—N1—C6—C7 | -5.23 (8)   |
| O1—Ni1—O6—C8   | 93.20 (5)   | C4—C5—C6—N1  | -1.14 (11)  |
| O10—Ni1—O6—C8  | -14.34 (9)  | C4—C5—C6—C7  | 176.52 (7)  |
| O5—Ni1—O6—C8   | -111.22 (5) | Ni1—O5—C7—O4 | -179.06 (7) |
| N2—Ni1—O10—C14 | 3.50 (5)    | Ni1—O5—C7—C6 | 0.27 (8)    |
| N1—Ni1—O10—C14 | -171.70 (6) | N1—C6—C7—O4  | -177.72 (7) |
| O1—Ni1—O10—C14 | -92.77 (6)  | C5—C6—C7—O4  | 4.43 (12)   |

|                |              |                 |             |
|----------------|--------------|-----------------|-------------|
| O6—Ni1—O10—C14 | 14.95 (9)    | N1—C6—C7—O5     | 2.91 (10)   |
| O5—Ni1—O10—C14 | 111.74 (6)   | C5—C6—C7—O5     | -174.94 (7) |
| N2—Ni1—N1—C6   | -119.19 (16) | Ni1—O6—C8—O7    | 179.06 (7)  |
| O1—Ni1—N1—C6   | -179.02 (6)  | Ni1—O6—C8—C9    | -0.37 (8)   |
| O10—Ni1—N1—C6  | -86.48 (6)   | C13—N2—C9—C10   | -0.51 (11)  |
| O6—Ni1—N1—C6   | 90.51 (6)    | Ni1—N2—C9—C10   | 172.42 (5)  |
| O5—Ni1—N1—C6   | 4.16 (6)     | C13—N2—C9—C8    | 178.98 (6)  |
| N2—Ni1—N1—C2   | 52.35 (19)   | Ni1—N2—C9—C8    | -8.09 (8)   |
| O1—Ni1—N1—C2   | -7.48 (6)    | O7—C8—C9—N2     | -174.29 (7) |
| O10—Ni1—N1—C2  | 85.06 (6)    | O6—C8—C9—N2     | 5.18 (9)    |
| O6—Ni1—N1—C2   | -97.95 (6)   | O7—C8—C9—C10    | 5.17 (11)   |
| O5—Ni1—N1—C2   | 175.70 (6)   | O6—C8—C9—C10    | -175.36 (7) |
| N1—Ni1—N2—C9   | -145.37 (15) | N2—C9—C10—C11   | 0.58 (11)   |
| O1—Ni1—N2—C9   | -86.74 (6)   | C8—C9—C10—C11   | -178.84 (7) |
| O10—Ni1—N2—C9  | -178.78 (6)  | C9—C10—C11—O8   | -179.59 (7) |
| O6—Ni1—N2—C9   | 6.18 (5)     | C9—C10—C11—C12  | 0.16 (11)   |
| O5—Ni1—N2—C9   | 93.38 (6)    | O8—C11—C12—C13  | 178.81 (7)  |
| N1—Ni1—N2—C13  | 27.7 (2)     | C10—C11—C12—C13 | -0.94 (11)  |
| O1—Ni1—N2—C13  | 86.32 (6)    | C9—N2—C13—C12   | -0.33 (11)  |
| O10—Ni1—N2—C13 | -5.72 (5)    | Ni1—N2—C13—C12  | -173.32 (5) |
| O6—Ni1—N2—C13  | 179.23 (6)   | C9—N2—C13—C14   | 179.79 (6)  |
| O5—Ni1—N2—C13  | -93.57 (6)   | Ni1—N2—C13—C14  | 6.80 (8)    |
| Ni1—O1—C1—O2   | 174.96 (7)   | C11—C12—C13—N2  | 1.04 (11)   |
| Ni1—O1—C1—C2   | -4.79 (9)    | C11—C12—C13—C14 | -179.09 (7) |
| C6—N1—C2—C3    | -0.36 (11)   | Ni1—O10—C14—O9  | -179.76 (7) |
| Ni1—N1—C2—C3   | -171.97 (6)  | Ni1—O10—C14—C13 | -1.05 (8)   |
| C6—N1—C2—C1    | 178.76 (6)   | N2—C13—C14—O9   | 175.34 (7)  |
| Ni1—N1—C2—C1   | 7.15 (8)     | C12—C13—C14—O9  | -4.54 (11)  |
| O2—C1—C2—N1    | 179.06 (7)   | N2—C13—C14—O10  | -3.46 (9)   |
| O1—C1—C2—N1    | -1.17 (10)   | C12—C13—C14—O10 | 176.66 (7)  |

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

| <i>D</i> —H $\cdots$ <i>A</i>       | <i>D</i> —H | H $\cdots$ <i>A</i> | <i>D</i> $\cdots$ <i>A</i> | <i>D</i> —H $\cdots$ <i>A</i> |
|-------------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| O3—H3C $\cdots$ O6 <sup>i</sup>     | 0.90        | 1.71                | 2.5901 (8)                 | 166                           |
| O8—H8 $\cdots$ O5 <sup>ii</sup>     | 0.90        | 1.84                | 2.7057 (9)                 | 160                           |
| O11—H11A $\cdots$ O9 <sup>iii</sup> | 0.90        | 1.86                | 2.7106 (9)                 | 156                           |
| O11—H11B $\cdots$ O2 <sup>iv</sup>  | 0.90        | 1.88                | 2.7464 (9)                 | 161                           |
| O12—H12B $\cdots$ O11 <sup>ii</sup> | 0.90        | 2.15                | 3.0425 (12)                | 172                           |
| O12—H12A $\cdots$ O5                | 0.90        | 2.15                | 3.0244 (13)                | 164                           |
| O12—H12A $\cdots$ O4                | 0.90        | 2.64                | 3.3823 (12)                | 141                           |
| N3—H3A $\cdots$ O4 <sup>v</sup>     | 0.90        | 2.15                | 2.9307 (10)                | 144                           |
| N3—H3B $\cdots$ O10                 | 0.90        | 2.07                | 2.8817 (10)                | 149                           |
| N4—H4A $\cdots$ O4 <sup>v</sup>     | 0.90        | 1.97                | 2.8072 (11)                | 154                           |
| N4—H4B $\cdots$ O11 <sup>vi</sup>   | 0.90        | 2.06                | 2.9161 (10)                | 159                           |
| N5—H5A $\cdots$ O10                 | 0.90        | 2.19                | 2.9887 (10)                | 148                           |
| N5—H5B $\cdots$ O2 <sup>vii</sup>   | 0.90        | 1.98                | 2.8656 (10)                | 169                           |
| N6—H6A $\cdots$ O7 <sup>viii</sup>  | 0.90        | 2.28                | 3.0178 (11)                | 139                           |

---

|                             |      |      |             |     |
|-----------------------------|------|------|-------------|-----|
| N6—H6A···O1                 | 0.90 | 2.49 | 3.0038 (9)  | 117 |
| N6—H6B···O3 <sup>vii</sup>  | 0.90 | 2.55 | 3.2320 (11) | 133 |
| N7—H7A···O9 <sup>iii</sup>  | 0.90 | 2.01 | 2.8999 (10) | 172 |
| N7—H7B···O12 <sup>ix</sup>  | 0.90 | 1.93 | 2.8282 (13) | 176 |
| N8—H8A···O7 <sup>viii</sup> | 0.90 | 1.94 | 2.7993 (10) | 158 |
| N8—H8B···O11                | 0.90 | 1.99 | 2.8847 (10) | 174 |
| N5—H5A···Cg1(N1/C2–C6)      | 0.90 | 3.47 | 3.362 (7)   | 75  |

---

Symmetry codes: (i)  $-x, -y+1, -z$ ; (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $-x+1, -y, -z+1$ ; (iv)  $-x, -y, -z+1$ ; (v)  $-x+1, -y+1, -z$ ; (vi)  $x, y, z-1$ ; (vii)  $-x, -y, -z$ ; (viii)  $-x, -y+1, -z+1$ ; (ix)  $x, y-1, z$ .