

Boson Stars as Solitary Waves

Journal Article**Author(s):**

Fröhlich, Jürg; Jonsson, B. Lars G.; Lenzmann, Enno

Publication date:

2007

Permanent link:

<https://doi.org/10.3929/ethz-b-000007429>

Rights / license:

In Copyright - Non-Commercial Use Permitted

Originally published in:

Communications in Mathematical Physics 274(1), <https://doi.org/10.1007/s00220-007-0272-9>

Boson Stars as Solitary Waves

Jürg Fröhlich¹, B. Lars G. Jonsson^{1,2}, Enno Lenzmann^{3,4}

¹ Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland.

E-mail: juerg@phys.ethz.ch; ljonsson@kth.se

² Division of Electromagnetic Engineering, School of Electrical Engineering, Royal Institute of Technology, SE-100 44 Stockholm, Sweden

³ Department of Mathematics, ETH Zurich, 8092 Zurich, Switzerland

⁴ Department of Mathematics, MIT, Cambridge, MA 02139, USA.

E-mail: lenzmann@mit.edu

Received: 14 December 2005 / Accepted: 11 December 2006

Published online: 19 June 2007 – © Springer-Verlag 2007

Abstract: We study the nonlinear equation

$$i \partial_t \psi = (\sqrt{-\Delta + m^2} - m) \psi - (|x|^{-1} * |\psi|^2) \psi \quad \text{on } \mathbb{R}^3,$$

which is known to describe the dynamics of pseudo-relativistic boson stars in the mean-field limit. For positive mass parameters, $m > 0$, we prove existence of travelling solitary waves, $\psi(t, x) = e^{it\mu} \varphi_v(x - vt)$, for some $\mu \in \mathbb{R}$ and with speed $|v| < 1$, where $c = 1$ corresponds to the speed of light in our units. Due to the lack of Lorentz covariance, such travelling solitary waves cannot be obtained by applying a Lorentz boost to a solitary wave at rest (with $v = 0$). To overcome this difficulty, we introduce and study an appropriate variational problem that yields the functions $\varphi_v \in H^{1/2}(\mathbb{R}^3)$ as minimizers, which we call boosted ground states. Our existence proof makes extensive use of concentration-compactness-type arguments.

In addition to their existence, we prove orbital stability of travelling solitary waves $\psi(t, x) = e^{it\mu} \varphi_v(x - vt)$ and pointwise exponential decay of $\varphi_v(x)$ in x .

1. Introduction

In this paper and its companion [4], we study solitary wave solutions — and solutions close to such — of the pseudo-relativistic Hartree equation

$$i \partial_t \psi = (\sqrt{-\Delta + m^2} - m) \psi - \left(\frac{1}{|x|} * |\psi|^2 \right) \psi \quad \text{on } \mathbb{R}^3. \quad (1.1)$$

Here $\psi(t, x)$ is a complex-valued wave field, and the symbol $*$ stands for convolution on \mathbb{R}^3 . The operator $\sqrt{-\Delta + m^2} - m$, which is defined via its symbol $\sqrt{k^2 + m^2} - m$ in Fourier space, is the kinetic energy operator of a relativistic particle of mass, $m \geq 0$, and the convolution kernel, $|x|^{-1}$, represents the Newtonian gravitational potential in appropriate physical units.

As recently shown by Elgart and Schlein in [3], Eq. (1.1) arises as an effective dynamical description for an N -body quantum system of relativistic bosons with two-body interaction given by Newtonian gravity. Such a system is a model system for a pseudo-relativistic *boson star*. That is, we consider a regime, where effects of special relativity (accounted for by the operator $\sqrt{-\Delta + m^2} - m$) become important, but general relativistic effects can be neglected. The idea of a mathematical model of pseudo-relativistic boson stars dates back to the works of Lieb and Thirring [10] and of Lieb and Yau [11], where the corresponding N -body Hamiltonian and its relation to the Hartree energy functional $\mathcal{H}(\psi) = 2\mathcal{E}(\psi)$ are discussed, with $\mathcal{E}(\psi)$ defined in (1.3), below.

Let us briefly recap the state of affairs concerning Eq. (1.1) itself. With help of the conserved quantities of charge, $\mathcal{N}(\psi)$, and energy, $\mathcal{E}(\psi)$, given by

$$\mathcal{N}(\psi) = \int_{\mathbb{R}^3} |\psi|^2 dx, \quad (1.2)$$

$$\mathcal{E}(\psi) = \frac{1}{2} \int_{\mathbb{R}^3} \overline{\psi} (\sqrt{-\Delta + m^2} - m) \psi dx - \frac{1}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx, \quad (1.3)$$

results derived so far can be summarized as follows (see also Fig. 1 below).

– **Well-Posedness:** For any initial datum $\psi_0 \in H^{1/2}(\mathbb{R}^3)$, there exists a unique solution

$$\psi \in C^0([0, T]; H^{1/2}(\mathbb{R}^3)) \cap C^1([0, T]; H^{-1/2}(\mathbb{R}^3)), \quad (1.4)$$

for some $T > 0$, where $H^s(\mathbb{R}^3)$ denotes the inhomogeneous Sobolev space of order s . Moreover, we have global-in-time existences (i. e., $T = \infty$) whenever the initial datum satisfies the condition

$$\mathcal{N}(\psi_0) < N_c, \quad (1.5)$$




Fig. 1. Qualitative diagram for the boson star Eq. (1.1) with positive mass parameter $m > 0$. Here $N = \mathcal{N}(\psi_0)$ and $E = \mathcal{E}(\psi_0)$ denote charge and energy for the initial condition $\psi_0 \in H^{1/2}(\mathbb{R}^3)$. In region **I**, all solutions are global in time and the (unboosted) ground states are minimizers of $\mathcal{E}(\psi)$ subject to fixed $\mathcal{N}(\psi_0) = N$ with $0 < N < N_c$. If N exceeds N_c , the energy E can attain values below $-\frac{1}{2}mN$. As shown in [5] for spherically symmetric $\psi_0 \in C_c^\infty(\mathbb{R}^3)$ that belong to region **III**, we have in fact blow-up of $\psi(t)$ within a finite time. Finally, the qualitative behavior of solutions with initial conditions in region **II** seems to be of indefinite nature

where $N_c > 4/\pi$ is some universal constant; see [7] for a detailed study of the Cauchy problem for (1.1) with initial data in $H^s(\mathbb{R}^3)$, $s \geq 1/2$.

- **Solitary Waves:** Due to the focusing nature of the nonlinearity in (1.1), there exist solitary wave solutions, which we refer to as *solitary waves*, given by

$$\psi(t, x) = e^{it\mu} \varphi(x), \tag{1.6}$$

where $\varphi \in H^{1/2}(\mathbb{R}^3)$ is defined as a minimizer of $\mathcal{E}(\psi)$ subject to $\mathcal{N}(\psi) = N$ fixed. Any such minimizer, $\varphi(x)$, is called a *ground state* and it has to satisfy the corresponding Euler-Lagrange equation

$$(\sqrt{-\Delta + m^2} - m)\varphi - \left(\frac{1}{|x|} * |\varphi|^2\right)\varphi = -\mu\varphi, \tag{1.7}$$

for some $\mu \in \mathbb{R}$. An existence proof of ground states, for $0 < \mathcal{N}(\varphi) < N_c$ and $m > 0$, can be found in [11]. The method used there is based on rearrangement inequalities that allow one to restrict one’s attention to radial functions, which simplifies the variational calculus. But in order to extend this existence result to so-called *boosted ground states*, i. e., x in (1.6) is replaced by $x - vt$ and Eq. (1.7) acquires the additional term, $i(v \cdot \nabla)\varphi$, we have to employ concentration-compactness-type methods; see Theorem 2.1 and its proof, below.

- **Blow-Up:** Any spherically symmetric initial datum, $\psi_0 \in C_c^\infty(\mathbb{R}^3)$, with

$$\mathcal{E}(\psi_0) < -\frac{1}{2}m\mathcal{N}(\psi_0) \tag{1.8}$$

leads to blow-up of $\psi(t)$ in a finite time, i. e., we have that $\lim_{t \nearrow T} \|\psi(t)\|_{H^{1/2}} = \infty$ holds, for some $T < \infty$. We remark that (1.8) implies that the smallness condition (1.5) cannot hold. See [5] for a proof of this blow-up result.¹ In physical terms, finite-time blow-up of $\psi(t)$ is indicative of “*gravitational collapse*” of a boson star modelled by (1.1); the constant N_c appearing in (1.5) may then be regarded as a “*Chandrasekhar limit mass*.”

We now come to the main issues of the present paper which focuses on existence and properties of travelling solitary waves for (1.1). More precisely, we consider solutions of the form

$$\psi(t, x) = e^{it\mu} \varphi_v(x - vt) \tag{1.9}$$

with some $\mu \in \mathbb{R}$ and travelling velocity, $v \in \mathbb{R}^3$, such that $|v| < 1$ holds (i. e., below the speed of light in our units). We point out that, since Eq. (1.1) is not Lorentz covariant, solutions such as (1.9) cannot be directly obtained from solitary waves at rest (i. e., we set $v = 0$) and then applying a Lorentz boost. To circumvent this difficulty, we plug the ansatz (1.9) into (1.1). This yields

$$(\sqrt{-\Delta + m^2} - m)\varphi_v + i(v \cdot \nabla)\varphi_v - \left(\frac{1}{|x|} * |\varphi_v|^2\right)\varphi_v = -\mu\varphi_v, \tag{1.10}$$

which is an Euler-Lagrange equation for the following minimization problem

$$\mathcal{E}_v(\psi) := \mathcal{E}(\psi) + \frac{i}{2} \int_{\mathbb{R}^3} \overline{\psi} (v \cdot \nabla) \psi \, dx = \min! \quad \text{subject to } \mathcal{N}(\psi) = N. \tag{1.11}$$

¹ In [5] the energy functional, $\mathcal{E}(\psi)$, is shifted by $+\frac{1}{2}m\mathcal{N}(\psi)$. Thus, condition (1.8) reads $\mathcal{E}(\psi_0) < 0$ in [5].

We refer to such minimizers, $\varphi_v \in \mathbf{H}^{1/2}(\mathbb{R}^3)$, as *boosted ground states* throughout this paper. Indeed, we will prove existence of boosted ground states when $|v| < 1$ and $0 < N < N_c(v)$ holds, as well as non-existence when $N \geq N_c(v)$; see Theorem 2.1, below. Our existence proof rests on concentration-compactness arguments which for our problem need some technical modifications, due to the pseudo-differential operator $\sqrt{-\Delta + m^2}$.

Apart from existence of boosted ground states, we are also concerned with properties such as “*orbital stability*” and exponential decay of $\varphi_v(x)$ in x ; see Theorems 3.1 and 4.1, below. We remark that both properties rely crucially on the positivity of the mass parameter, i. e., we have $m > 0$ in (1.1). By contrast, it is shown, for instance, in [5] that (resting) solitary waves become unstable when $m = 0$, due to nearby initial data leading to blow-up solutions.

In a companion paper [4], we will explore the effective dynamics of (slowly) travelling solitary waves in an external potential; see also Sect. 5 for a short summary of these result.

The plan of this paper is as follows:

- In Sect. 2, we set-up the variational calculus for problem (1.11) and we prove existence of boosted ground states, $\varphi_v \in \mathbf{H}^{1/2}(\mathbb{R}^3)$, for $0 < \mathcal{N}(\varphi_v) < N_c(v)$ and $|v| < 1$, as well as their nonexistence if $\mathcal{N}(\varphi_v) \geq N_c(v)$; see Theorem 2.1, below.
- Section. 3 addresses “orbital stability” of travelling solitary waves $\psi(t, x) = e^{it\mu} \varphi_v(x - vt)$; see Theorem 3.1, below.
- In Sect. 4, we derive pointwise exponential decay and regularity of boosted ground states; see Theorem 4.1, below.
- In Sect. 5, we sketch the main result of [4] describing the effective dynamics of travelling solitary waves in an external potential.
- In App. A–C, we collect and prove several technical statements which we refer to throughout this text.

Notation. Lebesgue spaces of complex-valued functions on \mathbb{R}^3 will be denoted by $L^p(\mathbb{R}^3)$, with norm $\|\cdot\|_p$ and $1 \leq p \leq \infty$. We define the Fourier transform for $f \in \mathcal{S}(\mathbb{R}^3)$ (i. e., Schwartz functions) according to

$$(\mathcal{F}f)(k) = \widehat{f}(k) = \frac{1}{(2\pi)^{3/2}} \int_{\mathbb{R}^3} f(x) e^{-ik \cdot x} dx, \quad (1.12)$$

where \mathcal{F} extends to $\mathcal{S}'(\mathbb{R}^3)$ (i. e., the space of tempered distributions) by duality.

For $s \in \mathbb{R}$, we introduce the operator $(1 - \Delta)^s$ via its multiplier $(1 + |k|^2)^s$ in Fourier space, i. e., we set $(1 - \Delta)^s f = \mathcal{F}^{-1}[(1 + |k|^2)^s \mathcal{F}f]$. Likewise, we define the operator $\sqrt{-\Delta + m^2}$ through its multiplier $\sqrt{|k|^2 + m^2}$ in Fourier space.

We employ Sobolev spaces, $\mathbf{H}^s(\mathbb{R}^3)$, of fractional order $s \in \mathbb{R}$ defined by

$$\mathbf{H}^s(\mathbb{R}^3) := \{f \in \mathcal{S}'(\mathbb{R}^3) : (1 - \Delta)^{s/2} f \in L^2(\mathbb{R}^3)\} \quad (1.13)$$

and equipped with the norm $\|f\|_{\mathbf{H}^s} := \|(1 - \Delta)^{s/2} f\|_2$.

Since we exclusively deal with \mathbb{R}^3 , we often write L^p and \mathbf{H}^s instead of $L^p(\mathbb{R}^3)$ and $\mathbf{H}^s(\mathbb{R}^3)$ in what follows. A further abbreviation we use is given by

$$\int_{\mathbb{R}^3} f dx := \int_{\mathbb{R}^3} f(x) dx. \quad (1.14)$$

We equip $L^2(\mathbb{R}^3)$ with a complex inner product, $\langle \cdot, \cdot \rangle$, defined as

$$\langle f, g \rangle := \int_{\mathbb{R}^3} \bar{f} g \, dx. \quad (1.15)$$

Operator inequalities (in the sense of quadratic forms) are denoted by $A \leq B$, which means that $\langle \psi, A\psi \rangle \leq \langle \psi, B\psi \rangle$ holds for all $\psi \in D(|A|^{1/2}) \subseteq D(|B|^{1/2})$, where A and B are self-adjoint operators on $L^2(\mathbb{R}^3)$ with domains $D(A)$ and $D(B)$, respectively. Finally, we remark that we employ the notation $v \cdot \nabla = \sum_{k=1}^3 v_k \partial_{x_k}$, where $v \in \mathbb{R}^3$ is some fixed vector.

2. Existence of Boosted Ground States

We consider the following minimization problem:

$$E_v(N) := \inf \{ \mathcal{E}_v(\psi) : \psi \in H^{1/2}(\mathbb{R}^3), \mathcal{N}(\psi) = N \}, \quad (2.1)$$

where $\mathcal{N}(\psi)$ is defined in (1.2), and $N > 0$, $v \in \mathbb{R}^3$, with $|v| < 1$, denote given parameters. Furthermore, we set

$$\begin{aligned} \mathcal{E}_v(\psi) := & \frac{1}{2} \langle \psi, (\sqrt{-\Delta + m^2} - m)\psi \rangle + \frac{i}{2} \langle \psi, (v \cdot \nabla)\psi \rangle \\ & - \frac{1}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 \, dx. \end{aligned} \quad (2.2)$$

Any minimizer, $\varphi_v \in H^{1/2}(\mathbb{R}^3)$, for (2.1) has to satisfy the corresponding Euler-Lagrange equation given by

$$(\sqrt{-\Delta + m^2} - m)\varphi_v + i(v \cdot \nabla)\varphi_v - \left(\frac{1}{|x|} * |\varphi_v|^2 \right)\varphi_v = -\mu\varphi_v, \quad (2.3)$$

with some Lagrange multiplier, $-\mu \in \mathbb{R}$, where this sign convention turns out to be convenient for our analysis. In what follows, we refer to such minimizers, φ_v , for (2.1) as *boosted ground states*, since they give rise to moving solitary waves

$$\psi(t, x) = e^{it\mu} \varphi_v(x - vt), \quad (2.4)$$

for (1.1) with travelling speed $v \in \mathbb{R}^3$ with $|v| < 1$.

Concerning existence of boosted ground states, we have the following theorem which generalizes a result derived in [11] for minimizers of (2.1) with $v = 0$.

Theorem 2.1. *Suppose that $m > 0$, $v \in \mathbb{R}^3$, and $|v| < 1$. Then there exists a positive constant $N_c(v)$ depending only on v such that the following holds.*

- i) *For $0 < N < N_c(v)$, problem (2.1) has a minimizer, $\varphi_v \in H^{1/2}(\mathbb{R}^3)$, and it satisfies (2.3), for some $\mu \in \mathbb{R}$. Moreover, every minimizing sequence, (ψ_n) , for (2.1) with $0 < N < N_c(v)$ is relatively compact in $H^{1/2}(\mathbb{R}^3)$ up to translations, i.e., there exists a sequence, (y_k) , in \mathbb{R}^3 and a subsequence, (ψ_{n_k}) , such that $\psi_{n_k}(\cdot + y_k) \rightarrow \varphi_v$ strongly in $H^{1/2}(\mathbb{R}^3)$ as $k \rightarrow \infty$, where φ_v is some minimizer for (2.1).*
- ii) *For $N \geq N_c(v)$, no minimizer exists for problem (2.1), even though $E_v(N)$ is finite for $N = N_c(v)$.*

Remarks. 1) It has been proved in [11] that (2.1) with $v = 0$ has a spherically symmetric minimizer, which can be chosen to be real-valued and nonnegative. But the proof given in [11] crucially relies on symmetric rearrangement arguments that allow to restrict to radial functions in this special case. For $v \neq 0$, such methods cannot be used and a general discussion of (2.1) needs a fundamental change of methods. Fortunately, it turns out that the concentration-compactness method introduced by P.-L. Lions in [12] is tailor-made for studying (2.1). To prove Theorem 2.1, we shall therefore proceed along the lines of [12]. But — due to the presence of the pseudo-differential operator $\sqrt{-\Delta + m^2}$ in (2.2) — some technical modifications have to be taken into account and they are worked out in detail in App. A.

- 2) A corresponding existence result for boosted ground states can also be derived when $-1/|x|$ in (2.2) is replaced by some other attractive two-body potential, e. g., a Yukawa type potential $\Phi(x) = -e^{-\mu|x|}/|x|$ with $\mu > 0$. But then minimal L^2 -norm of minimizers may also arise, i. e., the condition $N > N_*(v; \Phi)$ enters for some $N_*(v; \Phi) > 0$.
- 3) We do not know whether we have *uniqueness* of minimizers (up to phase and translation) for problem (2.1). Even the simpler case, where one assumes $v = 0$, has not been settled yet.

2.1. Setting up the Variational Calculus. Before we turn to the proof of Theorem 2.1, we collect and prove some preliminary results.

First one easily verifies that $\mathcal{E}_v(\psi)$ is real-valued (by using, for instance, Plancherel's theorem for the first two terms in (2.2)). Moreover, the inequality

$$\int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx \leq S_v \langle \psi, (\sqrt{-\Delta} + i v \cdot \nabla) \psi \rangle \langle \psi, \psi \rangle, \quad (2.5)$$

which is proven in App. B, ensures that $\mathcal{E}_v(\psi)$ is well-defined on $H^{1/2}(\mathbb{R}^3)$. As stated in Lemma B.1, inequality (2.5), with $|v| < 1$, has an optimizer, $Q_v \neq 0$, which satisfies

$$\sqrt{-\Delta} Q_v + i(v \cdot \nabla) Q_v - \left(\frac{1}{|x|} * |Q_v|^2 \right) Q_v = -Q_v \quad (2.6)$$

and yields the best constant, S_v , in terms of

$$S_v = \frac{2}{\langle Q_v, Q_v \rangle}. \quad (2.7)$$

Correspondingly, we introduce the constant, $N_c(v)$, by

$$N_c(v) := \frac{2}{S_v}. \quad (2.8)$$

By Lemma B.1, we also have the bounds

$$N_c \geq N_c(v) \geq (1 - |v|) N_c, \quad (2.9)$$

where $N_c(v = 0) = N_c > 4/\pi$ is, of course, the same constant that appeared in Sect. 1.

We now state our first auxiliary result for (2.1).

Lemma 2.1. *Suppose that $m \geq 0$, $v \in \mathbb{R}^3$, and $|v| < 1$. Then the following inequality holds:*

$$2\mathcal{E}_v(\psi) \geq \left(1 - \frac{N}{N_c(v)}\right) \left(\psi, (\sqrt{-\Delta} + iv \cdot \nabla)\psi\right) - mN \quad (2.10)$$

for all $\psi \in \mathbf{H}^{1/2}(\mathbb{R}^3)$ with $\mathcal{N}(\psi) = N$. Here $N_c(v)$ is the constant introduced in (2.8) above.

Moreover, we have that $E_v(N) \geq -\frac{1}{2}mN$ for $0 < N \leq N_c(v)$ and $E_v(N) = -\infty$ for $N > N_c(v)$. Finally, any minimizing sequence for problem (2.1) is bounded in $\mathbf{H}^{1/2}(\mathbb{R}^3)$ whenever $0 < N < N_c(v)$.

Proof (of Lemma 2.1). Let the assumption on m and v stated above be satisfied. Estimate (2.10) is derived by noting that $\sqrt{-\Delta + m^2} \geq \sqrt{-\Delta}$ and using inequality (2.5) together with the definition of $N_c(v)$ in (2.8). Furthermore, that $E_v(N) \geq -\frac{1}{2}mN$ for $N \leq N_c(v)$ is a consequence of (2.10) itself. To see that $E_v(N) = -\infty$ when $N > N_c(v)$, we recall from Lemma B.1 that there exists an optimizer, $Q_v \in \mathbf{H}^{1/2}(\mathbb{R}^3)$, with $\mathcal{N}(Q_v) = N_c(v)$, for inequality (2.5). Using that Q_v turns (2.5) into an equality and noticing that $\sqrt{-\Delta + m^2} - m \leq \sqrt{-\Delta}$ holds, a short calculation yields

$$E_v(N) \leq \mathcal{E}_v(\lambda Q_v) \Big|_{m=0} = -\frac{\lambda^2(\lambda^2 - 1)}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |Q_v|^2\right) |Q_v|^2 dx. \quad (2.11)$$

For $N > N_c(v)$, we can choose $\lambda > 1$ which implies that the right-hand side is strictly negative and, in addition, by L^2 -norm preserving rescalings, $Q_v(x) \mapsto a^{3/2} Q_v(ax)$ with $a > 0$, we find that

$$E_v(N) \leq \mathcal{E}_v(\lambda a^{3/2} Q_v(a \cdot)) \Big|_{m=0} = a \mathcal{E}_v(\lambda Q_v) \Big|_{m=0} \rightarrow -\infty, \quad (2.12)$$

with $\lambda > 1$ fixed and as $a \rightarrow \infty$. Thus, we deduce that $E_v(N) = -\infty$ when $N > N_c(v)$.

To see the $\mathbf{H}^{1/2}(\mathbb{R}^3)$ -boundedness of any minimizing sequence, (ψ_n) , with $0 < N < N_c(v)$, we note that $\sqrt{-\Delta} + iv \cdot \nabla \geq (1 - |v|)\sqrt{-\Delta}$ holds. Hence we see that $\sup_n \langle \psi_n, \sqrt{-\Delta} \psi_n \rangle \leq C < \infty$, thanks to (2.10). This completes the proof of Lemma 2.1. \square

As a next step, we derive an upper bound for $E_v(N)$, which is given by the nonrelativistic ground state energy, $E_v^{\text{nr}}(N)$, defined below. Here the positivity of the mass parameter, $m > 0$, is essential for deriving the following estimate.

Lemma 2.2. *Suppose that $m > 0$, $v \in \mathbb{R}^3$, and $|v| < 1$. Then we have that*

$$E_v(N) \leq -\frac{1}{2}(1 - \sqrt{1 - v^2})mN + E_v^{\text{nr}}(N), \quad (2.13)$$

where $E_v^{\text{nr}}(N)$ is given by

$$E_v^{\text{nr}}(N) := \inf \left\{ \mathcal{E}_v^{\text{nr}}(\psi) : \psi \in \mathbf{H}^1(\mathbb{R}^3), \mathcal{N}(\psi) = N \right\}, \quad (2.14)$$

$$\mathcal{E}_v^{\text{nr}}(\psi) := \frac{\sqrt{1 - v^2}}{4m} \int_{\mathbb{R}^3} |\nabla \psi|^2 dx - \frac{1}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2\right) |\psi|^2 dx. \quad (2.15)$$

Proof (of Lemma 2.2). To prove (2.13), we pick a spherically symmetric function, $\phi \in \mathbf{H}^1(\mathbb{R}^3)$ with $\mathcal{N}(\phi) = N$, and we introduce the one-parameter family

$$\phi_\lambda(x) := e^{i\lambda v \cdot x} \phi(x) = e^{i\lambda |v|z} \phi(x), \quad \text{with } \lambda > 0. \quad (2.16)$$

Here and in what follows, we assume (without loss of generality) that v is parallel to the z -axis, i. e., $v = |v|e_z$. One checks that

$$\frac{i}{2} \langle \phi_\lambda, (v \cdot \nabla) \phi_\lambda \rangle = -\frac{\lambda v^2}{2} N, \quad (2.17)$$

using the fact that $\langle \phi, \nabla \phi \rangle = 0$ holds, by spherical symmetry of $\phi(x)$. Hence, we find that

$$\begin{aligned} \mathcal{E}_v(\phi_\lambda) &= \frac{1}{2} \langle \phi_\lambda, (\sqrt{-\Delta + m^2} - m) \phi_\lambda \rangle + \frac{i|v|}{2} \langle \phi_\lambda, \partial_z \phi_\lambda \rangle \\ &\quad - \frac{1}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\phi_\lambda|^2 \right) |\phi_\lambda|^2 dx \\ &= \frac{1}{2} \left(\langle \phi_\lambda, (\sqrt{-\Delta + m^2} - m) \phi_\lambda \rangle - v^2 \lambda N \right) - \frac{1}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\phi|^2 \right) |\phi|^2 dx \\ &=: A + B. \end{aligned} \quad (2.18)$$

To estimate A in (2.18), we recall the operator inequality

$$\sqrt{-\Delta + m^2} \leq \frac{1}{2\lambda} (-\Delta + m^2 + \lambda^2), \quad (2.19)$$

which follows from the elementary inequality $2ab \leq a^2 + b^2$. Thus, we are led to

$$\begin{aligned} A &\leq \frac{1}{4\lambda} \langle \phi_\lambda, (-\Delta + m^2 + \lambda^2) \phi_\lambda \rangle - \frac{1}{2} m N - \frac{1}{2} v^2 \lambda N \\ &= \frac{1}{4\lambda} (\lambda^2 v^2 N + \langle \phi, -\Delta \phi \rangle + (m^2 + \lambda^2) N) - \frac{1}{2} m N - \frac{1}{2} v^2 \lambda N. \end{aligned} \quad (2.20)$$

By minimizing the upper bound (2.20) with respect to $\lambda > 0$, which is a matter of elementary calculations, we obtain with $\lambda_* = m/\sqrt{1-v^2}$ the estimate

$$\begin{aligned} \mathcal{E}_v(\phi_{\lambda_*}) &\leq -\frac{1}{2} (1 - \sqrt{1-v^2}) m N \\ &\quad + \frac{\sqrt{1-v^2}}{4m} \langle \phi, -\Delta \phi \rangle - \frac{1}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\phi|^2 \right) |\phi|^2 dx \\ &= -\frac{1}{2} (1 - \sqrt{1-v^2}) m N + \mathcal{E}_v^{\text{nr}}(\phi). \end{aligned} \quad (2.21)$$

Next, we remark that $\mathcal{E}_v^{\text{nr}}(\psi)$ is the energy functional for the non-relativistic boson star problem with mass parameter $m_v = m/\sqrt{1-v^2}$. Indeed, it is known from [8] that $\mathcal{E}_v^{\text{nr}}(\psi)$ subject to $\mathcal{N}(\psi) = N$ has a spherically symmetric minimizer, $\phi_* \in \mathbf{H}^1(\mathbb{R}^3)$, with

$$\mathcal{E}_v^{\text{nr}}(N) = \mathcal{E}_v^{\text{nr}}(\phi_*) < 0, \quad (2.22)$$

which completes the proof of Lemma 2.2. \square

By making use of Lemma 2.2, we show that the function $E_v(N)$ satisfies a strict sub-additivity condition. This is essential to the discussion of (2.1) when using concentration-compactness-type methods.

Lemma 2.3. *Suppose that $m > 0$, $v \in \mathbb{R}^3$, and $|v| < 1$. Then $E_v(N)$ satisfies the strict subadditivity condition*

$$E_v(N) < E_v(\alpha) + E_v(N - \alpha) \quad (2.23)$$

whenever $0 < N < N_c(v)$ and $0 < \alpha < N$. Here $N_c(v)$ is the constant of Lemma 2.1.

Moreover, the function $E_v(N)$ is strictly decreasing and strictly concave in N , where $0 < N < N_c(v)$.

Remarks. 1) Condition $m > 0$ is necessary for (2.23) to hold. To see this, note that if $m = 0$ then $\mathcal{E}_v(\psi_\lambda) = \lambda \mathcal{E}_v(\psi)$ holds, where $\psi_\lambda = \lambda^{3/2} \psi(\lambda x)$ and $\lambda > 0$. This leads to the conclusion that $E_v(N)$ is either 0 or $-\infty$ when $m = 0$.

2) The fact that $E_v(N)$ is strictly concave will be needed in our companion paper [4] when making use of the symplectic structure associated with the Hamiltonian PDE (1.1). More precisely, the strict concavity of $E_v(N)$ will enable us to prove the nondegeneracy of the symplectic form restricted to the manifold of solitary waves.

Proof (of Lemma 2.3). By Lemma 2.2 and the fact that $E_v^{\text{nr}}(N) \leq E_{v=0}^{\text{nr}}(N) < 0$ holds, by (2.22), we deduce that

$$E_v(N) < -\frac{1}{2}(1 - \sqrt{1 - v^2})mN. \quad (2.24)$$

Next, we notice the following scaling behavior:

$$E_v(N) = N e_v(N), \quad (2.25)$$

where

$$e_v(N) := \inf_{\psi \in H^{1/2}, \|\psi\|_2^2 = 1} \left\{ \frac{1}{2} \langle \psi, (\sqrt{-\Delta + m^2} - m) \psi \rangle + \frac{i}{2} \langle \psi, (v \cdot \nabla) \psi \rangle - \frac{N}{4} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx \right\}. \quad (2.26)$$

This shows that $e_v(N)$ is strictly decreasing, provided that we know that we may restrict the infimum to elements such that

$$\int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx \geq c > 0 \quad (2.27)$$

holds for some c . Suppose now that (2.27) were not true. Then there exists a minimizing sequence, (ψ_n) , such that

$$\int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_n|^2 \right) |\psi_n|^2 dx \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (2.28)$$

But on account of the fact that (cf. App. C)

$$\langle \psi, (\sqrt{-\Delta + m^2} - m) \psi \rangle + i \langle \psi, (v \cdot \nabla) \psi \rangle \geq -(1 - \sqrt{1 - v^2})mN, \quad (2.29)$$

we conclude that

$$E_v(N) = Ne_v(N) \geq -\frac{1}{2}(1 - \sqrt{1 - v^2})mN, \quad (2.30)$$

which contradicts (2.24). Thus $e_v(N)$ is strictly decreasing. Returning to (2.25) and noting that $e_v(N) < 0$ holds, by (2.24), we deduce that

$$E_v(\vartheta N) < \vartheta E_v(N), \quad \text{for } \vartheta > 1 \text{ and } 0 < N < N_c(v). \quad (2.31)$$

By an argument presented in [12], this inequality leads to the strict subadditivity condition (2.23).

Finally, we show that $E_v(N)$ is strictly decreasing and strictly concave on the interval $(0, N_c(v))$. To see that $E_v(N) = Ne_v(N)$ is strictly decreasing, we notice that $e_v(N)$ is strictly decreasing and negative. Furthermore, we remark that $e_v(N) = \inf\{\text{linear functions in } N\}$ has to be a concave function. Therefore it follows that $E_v(N) = Ne_v(N)$ is strictly concave, since the left- and right-derivatives, $D^\pm E_v(N)$, exist and are found to be strictly decreasing, by using that $e_v(N)$ is concave and strictly decreasing. \square

2.2. Proof of Theorem 2.1. We now come to the proof of Theorem 2.1 and we suppose that $m > 0$, $v \in \mathbb{R}^3$, and $|v| < 1$ holds.

Proof of Part i) Let us assume that

$$0 < N < N_c(v), \quad (2.32)$$

where $N_c(v)$ is the constant defined in (2.8). Furthermore, let (ψ_n) be a minimizing sequence for (2.1), i. e.,

$$\lim_{n \rightarrow \infty} \mathcal{E}_v(\psi_n) = E_v(N), \quad \text{with } \psi_n \in H^{1/2}(\mathbb{R}^3) \text{ and } \mathcal{N}(\psi_n) = N \text{ for all } n \geq 0. \quad (2.33)$$

By Lemma 2.1, we have that $E_v(N) > -\infty$ and that (ψ_n) is a bounded sequence in $H^{1/2}(\mathbb{R}^3)$. We now apply the following concentration-compactness lemma.

Lemma 2.4. *Let (ψ_n) be a bounded sequence in $H^{1/2}(\mathbb{R}^3)$ such that $\mathcal{N}(\psi_n) = \int_{\mathbb{R}^3} |\psi_n|^2 dx = N$ for all $n \geq 0$. Then there exists a subsequence, (ψ_{n_k}) , satisfying one of the three following properties:*

i) *Compactness: There exists a sequence, (y_k) , in \mathbb{R}^3 such that, for every $\epsilon > 0$, there exists $0 < R < \infty$ with*

$$\int_{|x-y_k| < R} |\psi_{n_k}|^2 dx \geq N - \epsilon. \quad (2.34)$$

ii) *Vanishing:*

$$\lim_{k \rightarrow \infty} \sup_{y \in \mathbb{R}^3} \int_{|x-y| < R} |\psi_{n_k}|^2 dx = 0, \quad \text{for all } R > 0.$$

iii) *Dichotomy*: There exists $\alpha \in (0, N)$ such that, for every $\epsilon > 0$, there exist two bounded sequences, (ψ_k^1) and (ψ_k^2) , in $H^{1/2}(\mathbb{R}^3)$ and $k_0 \geq 0$ such that, for all $k \geq k_0$, the following properties hold:

$$\|\psi_{n_k} - (\psi_k^1 + \psi_k^2)\|_p \leq \delta_p(\epsilon), \quad \text{for } 2 \leq p < 3, \quad (2.35)$$

with $\delta_p(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$, and

$$\left| \int_{\mathbb{R}^3} |\psi_k^1|^2 dx - \alpha \right| \leq \epsilon \quad \text{and} \quad \left| \int_{\mathbb{R}^3} |\psi_k^2|^2 dx - (N - \alpha) \right| \leq \epsilon, \quad (2.36)$$

$$\text{dist}(\text{supp } \psi_k^1, \text{supp } \psi_k^2) \rightarrow \infty, \quad \text{as } k \rightarrow \infty. \quad (2.37)$$

Moreover, we have that

$$\liminf_{k \rightarrow \infty} \left(\langle \psi_{n_k}, T \psi_{n_k} \rangle - \langle \psi_k^1, T \psi_k^1 \rangle - \langle \psi_k^2, T \psi_k^2 \rangle \right) \geq -C(\epsilon), \quad (2.38)$$

where $C(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$ and $T := (\sqrt{-\Delta + m^2} - m) + i(v \cdot \nabla)$ with $m \geq 0$ and $v \in \mathbb{R}^3$, $|v| < 1$.

Remark. We refer to App. A for the proof of Lemma 2.4. Part i) and ii) are standard, but part iii) requires some technical arguments, due to the presence of the pseudo-differential operator T .

Invoking Lemma 2.4, we conclude that a suitable subsequence, (ψ_{n_k}) , satisfies either i), ii), or iii). We rule out ii) and iii) as follows.

Suppose that (ψ_{n_k}) exhibits property ii). Then we conclude that

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_{n_k}|^2 \right) |\psi_{n_k}|^2 dx = 0, \quad (2.39)$$

by Lemma A.1. But as shown in the proof of Lemma 2.3, this implies

$$E_v(N) \geq -\frac{1}{2}(1 - \sqrt{1 - v^2})mN, \quad (2.40)$$

which contradicts (2.24). Hence ii) cannot occur.

Let us suppose that iii) is true for (ψ_{n_k}) . Then there exists $\alpha \in (0, N)$ such that, for every $\epsilon > 0$, there are two bounded sequences, (ψ_k^1) and (ψ_k^2) , with

$$\alpha - \epsilon \leq \mathcal{N}(\psi_k^1) \leq \alpha + \epsilon, \quad (N - \alpha) - \epsilon \leq \mathcal{N}(\psi_k^2) \leq (N - \alpha) + \epsilon, \quad (2.41)$$

for k sufficiently large. Moreover, inequality (2.38) and Lemma A.2 allow us to deduce that

$$E_v(N) = \lim_{k \rightarrow \infty} \mathcal{E}_v(\psi_{n_k}) \geq \liminf_{k \rightarrow \infty} \mathcal{E}_v(\psi_k^1) + \liminf_{k \rightarrow \infty} \mathcal{E}_v(\psi_k^2) - r(\epsilon), \quad (2.42)$$

where $r(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$. Since (ψ_k^1) and (ψ_k^2) satisfy (2.41), we infer

$$E_v(N) \geq E_v(\alpha + \epsilon) + E_v(N - \alpha + \epsilon) - r(\epsilon), \quad (2.43)$$

using that $E_v(N)$ is decreasing in N . Passing to the limit $\epsilon \rightarrow 0$ and by continuity of $E_v(N)$ in N (recall that $E_v(N)$ is a concave function on an open set), we deduce that

$$E_v(N) \geq E_v(\alpha) + E_v(N - \alpha) \quad (2.44)$$

holds for some $0 < \alpha < N$. This contradicts the strict subadditivity condition (2.23) stated in Lemma 2.3. Therefore iii) is ruled out.

By the discussion so far, we conclude that there exists a subsequence, (ψ_{n_k}) , such that i) of Lemma 2.4 is true for some sequence (y_k) in \mathbb{R}^3 . Let us now define the sequence

$$\tilde{\psi}_k := \psi_{n_k}(\cdot + y_k). \quad (2.45)$$

Since $(\tilde{\psi}_k)$ is a bounded sequence in $H^{1/2}(\mathbb{R}^3)$, we can pass to a subsequence, still denoted by $(\tilde{\psi}_k)$, such that $(\tilde{\psi}_k)$ converges weakly in $H^{1/2}(\mathbb{R}^3)$ to some $\varphi_v \in H^{1/2}(\mathbb{R}^3)$ as $k \rightarrow \infty$. Moreover, we have that $\tilde{\psi}_k \rightarrow \varphi_v$ strongly in $L^p_{\text{loc}}(\mathbb{R}^3)$ as $k \rightarrow \infty$, for $2 \leq p < 3$, thanks to a Rellich-type theorem for $H^{1/2}(\mathbb{R}^3)$ (see, e. g., [9, Theorem 8.6] for this). But on account of the fact

$$\int_{|x| < R} |\tilde{\psi}_k|^2 dx \geq N - \epsilon, \quad (2.46)$$

for every $\epsilon > 0$ and suitable $R = R(\epsilon) < \infty$, we conclude that

$$\tilde{\psi}_k \rightarrow \varphi_v \text{ strongly in } L^p(\mathbb{R}^3) \text{ as } k \rightarrow \infty, \text{ for } 2 \leq p < 3. \quad (2.47)$$

Next, by the Hardy–Littlewood–Sobolev and Hölder’s inequality, we deduce that

$$\begin{aligned} & \left| \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\tilde{\psi}_k|^2 \right) |\tilde{\psi}_k|^2 dx - \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\varphi_v|^2 \right) |\varphi_v|^2 dx \right| \\ & \leq C (\|\tilde{\psi}_k\|_{12/5}^3 + \|\varphi_v\|_{12/5}^3) \|\tilde{\psi}_k - \varphi_v\|_{12/5}. \end{aligned} \quad (2.48)$$

From (2.47), we have that $\tilde{\psi}_k$ converges strongly to φ_v in $L^{12/5}(\mathbb{R}^3)$, as $k \rightarrow \infty$, and therefore

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\tilde{\psi}_k|^2 \right) |\tilde{\psi}_k|^2 dx = \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\varphi_v|^2 \right) |\varphi_v|^2 dx. \quad (2.49)$$

Moreover, we have that

$$E_v(N) = \lim_{k \rightarrow \infty} \mathcal{E}_v(\tilde{\psi}_k) \geq \mathcal{E}_v(\varphi_v) \geq E_v(N), \quad (2.50)$$

since the functional

$$\mathcal{T}(\psi) := \langle \psi, \sqrt{-\Delta + m^2} \psi \rangle + i \langle \psi, (v \cdot \nabla) \psi \rangle, \quad (2.51)$$

is weakly lower semicontinuous on $H^{1/2}(\mathbb{R}^3)$, see Lemma A.4 in App. A. Thus, we have proved that $\varphi_v \in H^{1/2}(\mathbb{R}^3)$ is a minimizer for (2.1), i. e., we have $E_v(N) = \mathcal{E}_v(\varphi_v)$ and $\mathcal{N}(\varphi_v) = N$.

To prove the relative compactness of minimizing sequences in $H^{1/2}(\mathbb{R}^3)$ (up to translations), we notice that there has to be equality in (2.50), which leads to $\lim_{k \rightarrow \infty} \mathcal{T}(\tilde{\psi}_k) = \mathcal{T}(\varphi_v)$. By Lemma A.4, this fact implies a posteriori that

$$\tilde{\psi}_k \rightarrow \varphi_v \text{ strongly in } H^{1/2}(\mathbb{R}^3) \text{ as } k \rightarrow \infty, \quad (2.52)$$

which completes the proof of part i) of Theorem 2.1.

Proof of Part ii) To complete the proof of Theorem 2.1, we address its part ii). Clearly, no minimizer exists if $N > N_c(v)$, since in this case we have that $E_v(N) = -\infty$, by Lemma 2.1.

Next, we show that $E_v(N) = -\frac{1}{2}mN$ holds if $N = N_c(v)$, which can be seen as follows. We take an optimizer, $Q_v \in H^{1/2}(\mathbb{R}^3)$, for inequality (2.5); see Lemma B.1 and recall that $\mathcal{N}(Q_v) = N_c(v)$. Then

$$E_v(N) \leq \mathcal{E}_v(Q_v^{(\lambda)}) = \frac{1}{2} \langle Q_v^{(\lambda)}, (\sqrt{-\Delta + m^2} - \sqrt{-\Delta}) Q_v^{(\lambda)} \rangle - \frac{1}{2}mN, \quad (2.53)$$

for $N = N_c(v)$, where $Q_v^{(\lambda)}(x) := \lambda^{3/2} Q_v(\lambda x)$ with $\lambda > 0$, so that $\mathcal{N}(Q_v^{(\lambda)}) = \mathcal{N}(Q_v) = N_c(v)$. Using Plancherel's theorem and by dominated convergence, we deduce that

$$\begin{aligned} \langle Q_v^{(\lambda)}, (\sqrt{-\Delta + m^2} - \sqrt{-\Delta}) Q_v^{(\lambda)} \rangle &= \int_{\mathbb{R}^3} |\widehat{Q}_v(k)|^2 (\sqrt{\lambda^2 k^2 + m^2} - \lambda|k|) dk \\ &\rightarrow 0 \quad \text{as } \lambda \rightarrow \infty. \end{aligned} \quad (2.54)$$

Thus, we conclude that $E_v(N) \leq -\frac{1}{2}mN$ for $N = N_c(v)$. In combination with the estimate $E_v(N) \geq -\frac{1}{2}mN$ for $N \leq N_c(v)$ taken from Lemma 2.1, this shows that

$$E_v(N) = -\frac{1}{2}mN \quad \text{for } N = N_c(v). \quad (2.55)$$

Finally, we prove that there does not exist a minimizer for (2.1) with $N = N_c(v)$. We argue by contradiction as follows. Suppose that $\varphi_v \in H^{1/2}(\mathbb{R}^3)$ is a minimizer for (2.1) with $N = N_c(v)$. For $\varphi_v \in H^{1/2}(\mathbb{R}^3)$, $\varphi_v \neq 0$, and $m > 0$, we use $\langle \varphi_v, \sqrt{-\Delta + m^2} \varphi_v \rangle > \langle \varphi_v, \sqrt{-\Delta} \varphi_v \rangle$ to obtain

$$-\frac{1}{2}mN = \mathcal{E}_v(\varphi_v)|_{m>0} > \mathcal{E}_v(\varphi_v)|_{m=0} - \frac{1}{2}mN \geq -\frac{1}{2}mN, \quad (2.56)$$

which is a contradiction. Here we use Lemma 2.1 to estimate $\mathcal{E}_v(\varphi_v)|_{m=0} \geq 0$ for $\mathcal{N}(\varphi_v) = N_c(v)$. Hence no minimizer exists for (2.1) if $N \geq N_c(v)$. This completes the proof of Theorem 2.1. \square

3. Orbital Stability

The purpose of this section is to address “orbital stability” of travelling solitary waves

$$\psi(t, x) = e^{it\mu} \varphi_v(x - vt), \quad (3.1)$$

where $\varphi_v \in H^{1/2}(\mathbb{R}^3)$ is a boosted ground state. By the relative compactness of minimizing sequences (see Theorem 2.1) and by using a general idea presented in [2], we are able to prove the following abstract stability result.

Theorem 3.1. *Suppose that $m > 0$, $v \in \mathbb{R}^3$, $|v| < 1$, and $0 < N < N_c(v)$. Let $S_{v,N}$ denote the corresponding set of boosted ground states, i. e.,*

$$S_{v,N} := \{ \varphi_v \in H^{1/2}(\mathbb{R}^3) : \mathcal{E}_v(\varphi_v) = E_v(N), \mathcal{N}(\varphi_v) = N \},$$

which is non-empty by Theorem 2.1.

Then the solitary waves given in (3.1), with $\varphi_v \in S_{v,N}$, are stable in the following sense. For every $\epsilon > 0$, there exists $\delta > 0$ such that

$$\inf_{\varphi_v \in S_{v,N}} \|\psi_0 - \varphi_v\|_{H^{1/2}} \leq \delta \text{ implies that } \sup_{t \geq 0} \inf_{\varphi_v \in S_{v,N}} \|\psi(t) - \varphi_v\|_{H^{1/2}} \leq \epsilon.$$

Here $\psi(t)$ denotes the solution of (1.1) with initial condition $\psi_0 \in H^{1/2}(\mathbb{R}^3)$.

Remark. It is an interesting and open question whether uniqueness (modulo phase and translation) of boosted ground states holds, i.e., we have that $S_{v,N}$ is of the form $\{e^{i\gamma} \varphi_v(\cdot - y) : \gamma \in \mathbb{R}, y \in \mathbb{R}^3\}$, for some fixed $\varphi_v \in S_{v,N}$.

Proof (of Theorem 3.1). Let m and v satisfy the given assumptions. Since we have $N < N_c(v) \leq N_c$, we can choose $\delta > 0$ sufficiently small such that $\inf_{\phi \in S_{v,N}} \|\psi_0 - \phi\|_{H^{1/2}} \leq \delta$ guarantees that $\mathcal{N}(\psi_0) < N_c$. By the global well-posedness result for (1.1) derived in [7], we have that the corresponding solution, $\psi(t)$, exists for all times $t \geq 0$. Thus, taking $\sup_{t \geq 0}$ is well-defined.

Let us now assume that orbital stability (in the sense defined above) does not hold. Then there exists a sequence on initial data, $(\psi_n(0))$, in $H^{1/2}(\mathbb{R}^3)$ with

$$\inf_{\varphi \in S_{v,N}} \|\psi_n(0) - \varphi\|_{H^{1/2}} \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (3.2)$$

and some $\epsilon > 0$ such that

$$\inf_{\varphi \in S_{v,N}} \|\psi_n(t_n) - \varphi\|_{H^{1/2}} > \epsilon, \quad \text{for all } n \geq 0, \quad (3.3)$$

for a suitable sequence of times (t_n) . Note that (3.2) implies that $\mathcal{N}(\psi_n(0)) \rightarrow N$ as $n \rightarrow \infty$. Since $N < N_c$ by assumption, we can assume — without loss of generality — that $\mathcal{N}(\psi_n(0)) < N_c$ holds for all $n \geq 0$, which guarantees (see above) that the corresponding solution, $\psi_n(t)$, exists globally in time.

Next, we consider the sequence, (β_n) , in $H^{1/2}(\mathbb{R}^3)$ that is given by

$$\beta_n := \psi_n(t_n). \quad (3.4)$$

By conservation of $\mathcal{N}(\psi(t))$ and of $\mathcal{E}_v(\psi(t))$, whose proof can be done along the lines of [7] for the conservation of $\mathcal{E}(\psi(t))$, we have that $\mathcal{N}(\beta_n) = \mathcal{N}(\psi_n(0))$ and $\mathcal{E}_v(\beta_n) = \mathcal{E}_v(\psi_n(0))$, which, by (3.2), implies

$$\lim_{n \rightarrow \infty} \mathcal{E}_v(\beta_n) = E_v(N) \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathcal{N}(\beta_n) = N. \quad (3.5)$$

Defining the rescaled sequence

$$\tilde{\beta}_n := a_n \beta_n, \quad \text{where } a_n := \sqrt{N/\mathcal{N}(\beta_n)}, \quad (3.6)$$

and using the fact (β_n) has to be bounded in $H^{1/2}(\mathbb{R}^3)$, by virtue of Lemma 2.1, we infer that

$$\|\beta_n - \tilde{\beta}_n\|_{H^{1/2}} \leq C|1 - a_n| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.7)$$

By continuity of $\mathcal{E}_v : H^{1/2}(\mathbb{R}^3) \rightarrow \mathbb{R}$, we deduce that

$$\lim_{n \rightarrow \infty} \mathcal{E}_v(\tilde{\beta}_n) = E_v(N) \quad \text{and} \quad \mathcal{N}(\tilde{\beta}_n) = N, \quad \text{for all } n \geq 0. \quad (3.8)$$

Therefore $(\tilde{\beta}_n)$ is a minimizing sequence for (2.1) which, by Theorem 2.1 part i), has to contain a subsequence, $(\tilde{\beta}_{n_k})$, that strongly converges in $H^{1/2}(\mathbb{R}^3)$ (up to translations) to some minimizer $\varphi \in S_{v,N}$. In particular, inequality (3.3) cannot hold when $\beta_n = \psi_n(t_n)$ is replaced by $\tilde{\beta}_n$. But in view of (3.7), this conclusion is easily extended to the sequence (β_n) itself. Thus, we are led to a contradiction and the proof of Theorem 3.1 is complete.

□

4. Properties of Boosted Ground States

Concerning fundamental properties of boosted ground states given by Theorem 2.1, we have the following result.

Theorem 4.1. *Let $m > 0$, $v \in \mathbb{R}^3$, $|v| < 1$, and $0 < N < N_c(v)$. Then every boosted ground state, $\varphi_v \in H^{1/2}(\mathbb{R}^3)$, of problem (2.1) satisfies the following properties.*

- i) $\varphi_v \in H^s(\mathbb{R}^3)$ for all $s \geq 1/2$.
- ii) *The corresponding Lagrange multiplier satisfies $\mu > (1 - \sqrt{1 - v^2})m$. Moreover, we have pointwise exponential decay, i. e.,*

$$|\varphi_v(x)| \leq C e^{-\delta|x|} \quad (4.1)$$

holds for all $x \in \mathbb{R}^3$, where $\delta > 0$ and $C > 0$ are suitable constants.

- iii) *For $v = 0$, the function $\varphi_v(x)$ can be chosen to be radial, real-valued, and strictly positive.*

Remarks. 1) By part i) and Sobolev embeddings, any boosted ground state is smooth: $\varphi_v \in C^\infty(\mathbb{R}^3)$. Moreover, we have that $\varphi_v \in L^1 \cap L^\infty$, due to part ii).

2) Part iii) follows from the discussion presented in [11], except for the strict positivity which we will show below.

3) For a more precise exponential decay estimate for $\varphi_v(x)$, see Lemma C.1 in App. C.

Proof (of Theorem 4.1). Part i): We rewrite the Euler-Lagrange equation (3.1) for φ_v as

$$(H_0 + \lambda)\varphi_v = F(\varphi_v) + (\lambda - \mu)\varphi_v, \quad (4.2)$$

for any $\lambda \in \mathbb{R}$, where

$$H_0 := (\sqrt{-\Delta + m^2} - m) + i(v \cdot \nabla), \quad F(\varphi_v) := \left(\frac{1}{|x|} * |\varphi_v|^2\right)\varphi_v. \quad (4.3)$$

By [7, Lemma 3], we have that $F : H^s(\mathbb{R}^3) \rightarrow H^s(\mathbb{R}^3)$ for all $s \geq 1/2$ (F is indeed locally Lipschitz). Thus, the right-hand side in (4.2) belongs to $H^{1/2}(\mathbb{R}^3)$. Since H_0 is bounded from below, we can choose $\lambda > 0$ sufficiently large such that $(H_0 + \lambda)^{-1}$ exists. This leads to

$$\varphi_v = (H_0 + \lambda)^{-1} [F(\varphi_v) + (\lambda - \mu)\varphi_v]. \quad (4.4)$$

Noting that $(H_0 + \lambda)^{-1} : H^s(\mathbb{R}^3) \rightarrow H^{s+1}(\mathbb{R}^3)$, we see that $\varphi_v \in H^{3/2}(\mathbb{R}^3)$. By repeating the argument, we conclude that $\varphi_v \in H^s(\mathbb{R}^3)$ for all $s \geq 1/2$. This proves part i).

Part ii): The exponential decay follows from Lemma C.1, provided that the Lagrange multiplier, $-\mu$, satisfies

$$-\mu < -(1 - \sqrt{1 - v^2})m, \quad (4.5)$$

which means that $-\mu$ lies strictly below the essential spectrum of H_0 ; see App. C. To prove (4.5), we multiply the Euler-Lagrange equation by $\bar{\varphi}_v$ and integrate to obtain

$$2E_v(N) - \frac{1}{2} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\varphi_v|^2\right) |\varphi_v|^2 dx = -\mu N. \quad (4.6)$$

Using the upper bound (2.24) for $E_v(N)$, we conclude that

$$-\mu N < -(1 - \sqrt{1 - v^2})mN, \quad (4.7)$$

which proves (4.5).

Part iii): For the sake of brevity, we write $\varphi(x) := \varphi_{v=0}(x)$. By [11] problem (2.1), with $v = 0$, has a minimizer that equals its symmetric-decreasing rearrangement, i. e., $\varphi(x) = \varphi^*(x)$. In particular, $\varphi(x)$ is a spherically symmetric, real-valued, nonincreasing function with $\varphi(x) \geq 0$. It remains to show that $\varphi(x) > 0$ holds. To see this, we put $\lambda = \mu$ in (4.4), which is possible by the proof of ii), and we obtain

$$\varphi = ((\sqrt{-\Delta + m^2} - m) + \mu)^{-1} F(\varphi). \quad (4.8)$$

By using functional calculus for the self-adjoint operator $\sqrt{-\Delta + m^2}$ on $L^2(\mathbb{R}^3)$ with domain $H^1(\mathbb{R}^3)$, we find that

$$((\sqrt{-\Delta + m^2} - m) + \mu)^{-1} = \int_0^\infty e^{-t\mu} e^{-t(\sqrt{-\Delta + m^2} - m)} dt. \quad (4.9)$$

Referring to the explicit formula (C.10) for $v = 0$, we see that the integral kernel, $e^{-t(\sqrt{-\Delta + m^2} - m)}(x, y)$, is strictly positive. In view of (4.8), (4.9), and the fact that $F(\varphi) \geq 0$, we conclude that $\varphi(x) > 0$ holds for almost every $x \in \mathbb{R}^3$. But since $\varphi(x)$ is a nonincreasing and continuous function, we deduce that $\varphi(x) > 0$ has to be true for all $x \in \mathbb{R}^3$. This completes the proof of Theorem 4.1. \square

5. Outlook

Our analysis presented so far serves as a basis for the upcoming work in [4] which explores the effective motion of travelling solitary waves in an external potential. More precisely, we consider

$$i \partial_t \psi = (\sqrt{-\Delta + m^2} - m) \psi + V \psi - \left(\frac{1}{|x|} * |\psi|^2 \right) \psi \quad \text{on } \mathbb{R}^3. \quad (5.1)$$

Here the external potential $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ is assumed to be a smooth, bounded function with bounded derivatives. Note that its spatial variation introduces the length scale

$$\ell_{\text{ext}} = \|\nabla V\|_\infty^{-1}. \quad (5.2)$$

In addition, another length scale, ℓ_{sol} , enters through the exponential decay of $\varphi_v(x)$, i. e., we have that

$$\ell_{\text{sol}} = \delta^{-1}, \quad (5.3)$$

where $\delta > 0$ is the constant taken from Theorem 4.1. On intuitive grounds, one expects that if we have that

$$\ell_{\text{sol}} \ll \ell_{\text{ext}} \quad (5.4)$$

holds, then solutions, $\psi(t, x)$, of (5.1) that are initially close to some $\varphi_v(x)$ should approximately behave like point-particles, at least on a large (but possibly finite) interval of time.

We now briefly sketch how this heuristic picture of point-particle behavior of solitary waves is addressed by rigorous analysis in [4]. There we introduce a nondegeneracy assumption on the linearized operator

$$L := \begin{pmatrix} L_1 & 0 \\ 0 & L_2 \end{pmatrix} \quad (5.5)$$

acting on $L^2(\mathbb{R}^3; \mathbb{R}^2)$ with domain $H^1(\mathbb{R}^3; \mathbb{R}^2)$, where

$$L_1 \xi := (\sqrt{-\Delta + m^2} - m + \mu_0) \xi - \left(\frac{1}{|x|} * \varphi^2 \right) \xi - 2 \left(\frac{1}{|x|} * (\varphi \xi) \right) \varphi, \quad (5.6)$$

$$L_2 \xi := (\sqrt{-\Delta + m^2} - m + \mu_0) \xi - \left(\frac{1}{|x|} * \varphi^2 \right) \xi. \quad (5.7)$$

Here $\varphi(x) = \varphi_{v=0}(x)$ is some ground state at rest, whose corresponding Lagrange multiplier we denote by $\mu_0 \in \mathbb{R}$. Note that we can assume that $\varphi_{v=0}(x)$ is spherically symmetric and real-valued, see Theorem 4.1.

The nondegeneracy condition introduced in [4] now reads

$$\ker(L) = \text{span} \left\{ \begin{pmatrix} 0 \\ \varphi \end{pmatrix}, \begin{pmatrix} \partial_{x_1} \varphi \\ 0 \end{pmatrix}, \begin{pmatrix} \partial_{x_2} \varphi \\ 0 \end{pmatrix}, \begin{pmatrix} \partial_{x_3} \varphi \\ 0 \end{pmatrix} \right\}. \quad (5.8)$$

Under this kernel assumption, we then construct in [4] (by an implicit-function-type argument) a map $(v, \mu) \mapsto \varphi_{v, \mu}$, so that $\varphi_{v, \mu} \in H^{1/2}$ solves Eq. (1.7) and (v, μ) belongs to the small neighborhood around $(0, \mu_0)$.

The main result proven in [4] can now be sketched as follows. We consider suitable external potentials of the form

$$V(x) := W(\epsilon x). \quad (5.9)$$

Furthermore, let φ_{v_0, μ_0} with $|v_0| \ll 1$ be given and choose $\epsilon \ll 1$ so that (5.4) holds. Then for any initial datum, $\psi_0(x)$, such that

$$\|\|\psi_0 - e^{i\vartheta_0} \varphi_{v_0, \mu_0}(\cdot - a_0)\|\| \leq \epsilon, \quad \text{for some } \vartheta_0 \in \mathbb{R} \text{ and } a_0 \in \mathbb{R}^3, \quad (5.10)$$

where $\|\|\cdot\|\|$ is some weighted Sobolev norm, the corresponding solution, $\psi(t, x)$, of (5.1) can be uniquely written as

$$\psi(t, x) = e^{i\vartheta} [\varphi_{v, \mu}(x - a) + \xi(t, x - a)], \quad \text{for } 0 \leq t < C\epsilon^{-1}. \quad (5.11)$$

Here $\|\|\xi\|\| = \mathcal{O}(\epsilon)$ holds and the time-dependent functions $\{\vartheta, a, v, \mu\}$ satisfy equations of the following form:

$$\begin{cases} \dot{N} = \mathcal{O}(\epsilon^2), & \dot{\vartheta} = \mu - V(a) + \mathcal{O}(\epsilon^2), \\ \dot{a} = v + \mathcal{O}(\epsilon^2), & \gamma(\mu, v)\dot{v} = -\nabla V(a) + \mathcal{O}(\epsilon^2), \end{cases} \quad (5.12)$$

where $N = \mathcal{N}(\varphi_{v, \mu})$. The term $\gamma(\mu, v)$ can be viewed as an “effective mass” which takes relativistic effects into account.

Finally, we remark that the proof of (5.11) and (5.12) makes extensive use of the Hamiltonian formulation of (5.1) and its associated symplectic structure restricted to the manifold of solitary waves. Moreover, assumption (5.8) enables us to derive additional properties of $\varphi_v(x)$, for $|v| \ll 1$, such as cylindrical symmetry with respect to the v -axis, which is of crucial importance in the analysis presented in [4].

A. Variational and Pseudo-Differential Calculus

In this section of the appendix, we collect and prove results needed for our variational and pseudo-differential calculus.

A.1. Proof of Lemma 2.4. Let (ψ_n) be a bounded sequence in $H^{1/2}(\mathbb{R}^3)$ with $\|\psi_n\|_2^2 = N$ for all n . Along the lines of [12], we define the sequence, (Q_n) , of Lévy concentration functions by

$$Q_n(R) := \sup_{y \in \mathbb{R}^3} \int_{|x-y| < R} |\psi_n|^2 dx, \quad \text{for } R \geq 0. \quad (\text{A.1})$$

As stated in [12], there exists a subsequence, (Q_{n_k}) , such that

$$Q_{n_k}(R) \rightarrow Q(R) \text{ as } k \rightarrow \infty \text{ for all } R \geq 0, \quad (\text{A.2})$$

where $Q(R)$ is a nonnegative, nondecreasing function. Clearly, we have that

$$\alpha := \lim_{R \rightarrow \infty} Q(R) \in [0, N]. \quad (\text{A.3})$$

If $\alpha = 0$, then situation ii) of Lemma 2.4 arises as a direct consequence of definition (A.1). If $\alpha = N$, then i) follows, see [12] for details.

Assume that $\alpha \in (0, N)$ holds, and let $\epsilon > 0$ be given. Suppose that $\xi, \phi \in C^\infty(\mathbb{R}^3)$ with $0 \leq \phi, \xi \leq 1$ such that

$$\xi(x) \equiv 1 \quad \text{for } 0 \leq |x| \leq 1, \quad \xi(x) \equiv 0 \quad \text{for } |x| \geq 2, \quad (\text{A.4})$$

$$\phi(x) \equiv 0 \quad \text{for } 0 \leq |x| \leq 1, \quad \phi(x) \equiv 1 \quad \text{for } |x| \geq 2. \quad (\text{A.5})$$

Furthermore, we put $\xi_R(x) := \xi(x/R)$ and $\phi_R(x) := \phi(x/R)$, for $R > 0$, and we introduce

$$\psi_k^1 := \xi_{R_1}(\cdot - y_k) \psi_{n_k} \quad \text{and} \quad \psi_k^2 := \phi_{R_k}(\cdot - y_k) \psi_{n_k}. \quad (\text{A.6})$$

As shown in [12, Proof of Lemma III.1], there exists

$$R_1(\epsilon) \rightarrow \infty, \quad \text{as } \epsilon \rightarrow 0, \quad (\text{A.7})$$

and a sequence, (R_k) , with

$$R_k \rightarrow \infty, \quad \text{as } k \rightarrow \infty, \quad (\text{A.8})$$

such that (ψ_k^1) and (ψ_k^2) satisfy (2.36) and (2.37) in Lemma 2.4. Moreover, we have that

$$\int_{\mathbb{R}^3} |\psi_{n_k} - (\psi_k^1 + \psi_k^2)|^2 dx \leq 4\epsilon, \quad (\text{A.9})$$

for k sufficiently large.

By [9, Theorem 7.16], we see that ψ_k^1 and ψ_k^2 defined in (A.6) are bounded in $H^{1/2}(\mathbb{R}^3)$. More precisely, using the technique of the proof given there and the explicit formula

$$\langle f, \sqrt{-\Delta} f \rangle = (\text{const.}) \int_{\mathbb{R}^3 \times \mathbb{R}^3} \frac{|f(x) - f(y)|^2}{|x - y|^4} dx dy, \quad \text{for } f \in H^{1/2}(\mathbb{R}^3), \quad (\text{A.10})$$

we deduce that

$$\|gf\|_{H^{1/2}} \leq C(\|g\|_\infty + \|\nabla g\|_\infty) \|f\|_{H^{1/2}}. \quad (\text{A.11})$$

Thus, we find that

$$\|\psi_k^1\|_{H^{1/2}} \leq C\left(1 + \frac{1}{R_1}\right) \quad \text{and} \quad \|\psi_k^2\|_{H^{1/2}} \leq C\left(1 + \frac{1}{R_k}\right), \quad (\text{A.12})$$

for some constant $C = C(M)$, where $M = \sup_{k \geq 0} \|\psi_{n_k}\|_{H^{1/2}} < \infty$. Thus, (ψ_k^1) and (ψ_k^2) are bounded sequences in $H^{1/2}(\mathbb{R}^3)$. This fact together with Hölder's and Sobolev's inequalities leads to

$$\|\psi_{n_k} - (\psi_k^1 + \psi_k^2)\|_p \leq \delta_p(\epsilon), \quad \text{for } 2 \leq p < 3, \quad (\text{A.13})$$

where $\delta_p(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$. This proves (2.35) in Lemma 2.4.

It remains to show property (2.38) in Lemma 2.4. Since

$$\liminf_{k \rightarrow \infty} \left(\langle \psi_{n_k}, (-m)\psi_{n_k} \rangle - \langle \psi_k^1, (-m)\psi_k^1 \rangle - \langle \psi_k^2, (-m)\psi_k^2 \rangle \right) \quad (\text{A.14})$$

$$\geq -mN + m(\alpha - \epsilon) + m(N - \alpha - \epsilon) \geq -2m\epsilon \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0, \quad (\text{A.15})$$

we observe that it suffices to prove the claim

$$\liminf_{k \rightarrow \infty} \left(\langle \psi_{n_k}, A\psi_{n_k} \rangle - \langle \psi_k^1, A\psi_k^1 \rangle - \langle \psi_k^2, A\psi_k^2 \rangle \right) \geq -C(\epsilon), \quad (\text{A.16})$$

for some constant $C(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$, where

$$A := \sqrt{-\Delta + m^2} + i(v \cdot \nabla) + \lambda, \quad (\text{A.17})$$

with $m \geq 0$, $v \in \mathbb{R}^3$, $|v| < 1$, and $\lambda > 0$ is some constant so that

$$A \geq (1 - |v|)\sqrt{-\Delta} + \lambda \geq \lambda > 0. \quad (\text{A.18})$$

In view of (A.14), adding any fixed λ can be done without loss of generality.

Next, we recall definition (A.6) and rewrite the left-hand side in (A.16) as follows

$$\liminf_{k \rightarrow \infty} \langle \psi_{n_k}, (A - \xi_k A \xi_k - \phi_k A \phi_k) \psi_{n_k} \rangle, \quad (\text{A.19})$$

where

$$\xi_k(x) := \xi_{R_1}(x - y_k) \quad \text{and} \quad \phi_k(x) := \phi_{R_k}(x - y_k). \quad (\text{A.20})$$

Using commutators $[X, Y] := XY - YX$, we find that

$$\begin{aligned} A - \xi_k A \xi_k - \phi_k A \phi_k &= A(1 - \xi_k^2 - \phi_k^2) - [\xi_k, A]\xi_k - [\phi_k, A]\phi_k \\ &= \sqrt{A}(1 - \xi_k^2 - \phi_k^2)\sqrt{A} - \sqrt{A}[\sqrt{A}, (\xi_k^2 + \phi_k^2)] \\ &\quad - [\xi_k, A]\xi_k - [\phi_k, A]\phi_k. \end{aligned} \quad (\text{A.21})$$

Note that $\sqrt{A} > 0$ holds, due to $A > 0$. By applying Lemma A.3, we obtain

$$\|[\xi_k, A]\|_{L^2 \rightarrow L^2} \leq C \|\nabla \xi_k\|_\infty \leq \frac{C}{R_1}, \quad (\text{A.22})$$

$$\|[\phi_k, A]\|_{L^2 \rightarrow L^2} \leq C \|\nabla \phi_k\|_\infty \leq \frac{C}{R_k}. \quad (\text{A.23})$$

To estimate the remaining commutator in (A.21), we use (A.58) in the proof of Lemma A.3 to find that

$$\|[\sqrt{A}, (\xi_k^2 + \phi_k^2)]\|_{L^2 \rightarrow L^2} \leq C \left(\frac{1}{R_1} + \frac{1}{R_k} \right) \int_0^\infty \sqrt{s} \left\| \frac{1}{(s+A)} \right\|_{L^2 \rightarrow L^2}^2 ds \quad (\text{A.24})$$

$$\leq C \left(\frac{1}{R_1} + \frac{1}{R_k} \right) \int_0^\infty \frac{\sqrt{s}}{(s+\lambda)^2} ds \quad (\text{A.25})$$

$$\leq C \left(\frac{1}{R_1} + \frac{1}{R_k} \right). \quad (\text{A.26})$$

Returning to (A.19) and using that $\|\psi_{n_k}\|_{H^{1/2}} \leq C$, we conclude, for k large, that

$$\begin{aligned} \langle \psi_{n_k}, (A - \xi_k A \xi_k - \phi_k A \phi_k) \psi_{n_k} \rangle &\geq \langle \sqrt{A} \psi_{n_k}, (1 - \xi_k^2 - \phi_k^2) \sqrt{A} \psi_{n_k} \rangle \\ &\quad - C \left(\frac{1}{R_1} + \frac{1}{R_k} \right) \end{aligned} \quad (\text{A.27})$$

$$\geq -C \left(\frac{1}{R_1} + \frac{1}{R_k} \right), \quad (\text{A.28})$$

since $(1 - \xi_k^2 - \phi_k^2)(x) \geq 0$ when k is sufficiently large. Finally, we note that $R_k \rightarrow \infty$ as $k \rightarrow \infty$ as well as $R_1(\epsilon) \rightarrow \infty$ as $\epsilon \rightarrow 0$ holds, which leads to

$$\liminf_{k \rightarrow \infty} \langle \psi_{n_k}, (A - \xi_k A \xi_k - \phi_k A \phi_k) \psi_{n_k} \rangle \geq -C(\epsilon) \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0. \quad (\text{A.29})$$

The proof of Lemma 2.4 is now complete. \square

A.2. Technical Details for the Proof of Theorem 2.1.

Lemma A.1. *Let (ψ_n) satisfy the assumptions of Lemma 2.4. Furthermore, suppose that there exists a subsequence, (ψ_{n_k}) , that satisfies part ii) of Lemma 2.4. Then*

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_{n_k}|^2 \right) |\psi_{n_k}|^2 dx = 0.$$

Remark. A similar statement can be found in [12] in the context of other variational problems. For the sake of completeness, we present its proof for the situation at hand.

Proof (of Lemma A.1). Let (ψ_{n_k}) be a bounded sequence in $H^{1/2}(\mathbb{R}^3)$ such that

$$\int_{\mathbb{R}^3} |\psi_{n_k}|^2 dx = N, \quad \text{for all } k \geq 0, \quad (\text{A.30})$$

and assume that (ψ_{n_k}) satisfies part ii) in Lemma 2.4, i. e.,

$$\lim_{k \rightarrow \infty} \sup_{y \in \mathbb{R}^3} \int_{|x-y| < R} |\psi_{n_k}|^2 dx = 0, \quad \text{for all } R > 0. \quad (\text{A.31})$$

For simplicity, let $\psi_k := \psi_{n_k}$.

We introduce

$$f_\delta(x) := |x|^{-1} \chi_{\{|x|^{-1} \geq \delta\}}, \quad \text{with } \delta > 0, \quad (\text{A.32})$$

where χ_A denotes the characteristic function of the set $A \subset \mathbb{R}^3$. This definition leads to

$$\int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k|^2 \right) |\psi_k|^2 dx \leq \delta C + \int_{\mathbb{R}^3 \times \mathbb{R}^3} |\psi_k(x)|^2 |\psi_k(y)|^2 f_\delta(x-y) dx dy, \quad (\text{A.33})$$

where C is some constant. For $R > 0$ and $\delta > 0$, let

$$g_\delta^R(x) := \min\{f_\delta(x), R\}, \quad (\text{A.34})$$

$$f_\delta^R(x) := \max\{f_\delta(x) - R, 0\} \chi(x)_{\{|x| \leq R\}} + f_\delta(x) \chi(x)_{\{|x| > R\}}. \quad (\text{A.35})$$

Notice that $f_\delta \leq g_\delta^R \chi_{\{|x| \leq R\}} + f_\delta^R$ holds. In view of (A.33), this leads to

$$\begin{aligned} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k|^2 \right) |\psi_k|^2 dx &\leq \delta C + \int_{\mathbb{R}^3 \times \mathbb{R}^3} (|\psi_k(x)|^2 |\psi_k(y)|^2 \\ &\quad \times g_\delta^R(x-y) \chi(x-y)_{\{|x-y| \leq R\}}) dx dy \\ &\quad + \|\psi_k\|_{\delta/3}^4 \|f_\delta^R\|_2 \\ &=: \delta C + I + II, \end{aligned} \quad (\text{A.36})$$

using Young's inequality and that $f_\delta^R \in L^2(\mathbb{R}^3)$. By our assumption on (ψ_k) , we find that

$$I \leq R \int_{\mathbb{R}^3} |\psi_k(x)|^2 dx \int_{|x-y| \leq R} |\psi_k(y)|^2 dy \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

Furthermore, we have that

$$II \leq C \|f_\delta^R\|_2, \quad (\text{A.37})$$

by Sobolev's inequalities and the fact that (ψ_k) is bounded in $H^{1/2}(\mathbb{R}^3)$. Thus, we obtain

$$0 \leq \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k|^2 \right) |\psi_k|^2 dx \leq \delta C + C \|f_\delta^R\|_2 + r(k), \quad \text{for all } \delta, R > 0, \quad (\text{A.38})$$

where $r(k) \rightarrow 0$ as $k \rightarrow \infty$. Since $\|f_\delta^R\|_2 \rightarrow 0$ as $R \rightarrow \infty$, for each fixed $\delta > 0$, the assertion of Lemma A.1 follows by letting $R \rightarrow \infty$ and then sending δ to 0. \square

Lemma A.2. *Suppose that $\epsilon > 0$. Let (ψ_n) satisfy the assumptions of Lemma 2.4 and let (ψ_{n_k}) be a subsequence that satisfies part iii) with sequences (ψ_k^1) and (ψ_k^2) . Then, for k sufficiently large,*

$$\begin{aligned} - \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_{n_k}|^2 \right) |\psi_{n_k}|^2 dx &\geq - \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^1|^2 \right) |\psi_k^1|^2 dx \\ &\quad - \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^2|^2 \right) |\psi_k^2|^2 dx - r_1(k) - r_2(\epsilon), \end{aligned}$$

where $r_1(k) \rightarrow 0$ as $k \rightarrow \infty$ and $r_2(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

Proof (of Lemma A.2). Let $\epsilon > 0$ and suppose that (ψ_{n_k}) , (ψ_k^1) , and (ψ_k^2) satisfy the assumptions stated above. Introducing

$$\beta_k := \psi_{n_k} - (\psi_k^1 + \psi_k^2) \quad (\text{A.39})$$

and expanding the squares, we find that

$$\begin{aligned} \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_{n_k}|^2 \right) |\psi_{n_k}|^2 dx &= \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^1|^2 \right) |\psi_k^1|^2 dx \\ &+ \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^2|^2 \right) |\psi_k^2|^2 dx + \sum_{n=0}^4 I_n, \end{aligned} \quad (\text{A.40})$$

where

$$\begin{aligned} I_0 &= 2 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^1|^2 \right) |\psi_k^2|^2 dx + 4 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * (\operatorname{Re} \bar{\psi}_k^1 \psi_k^2) \right) (\operatorname{Re} \bar{\psi}_k^1 \psi_k^2) dx \\ &+ 4 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^1|^2 \right) (\operatorname{Re} \bar{\psi}_k^1 \psi_k^2) dx \\ &+ 4 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^2|^2 \right) (\operatorname{Re} \bar{\psi}_k^1 \psi_k^2) dx, \end{aligned} \quad (\text{A.41})$$

$$I_1 = 4 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^1 + \psi_k^2|^2 \right) (\operatorname{Re} \bar{\beta}_k (\psi_k^1 + \psi_k^2)) dx, \quad (\text{A.42})$$

$$\begin{aligned} I_2 &= 4 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * (\operatorname{Re} \bar{\beta}_k (\psi_k^1 + \psi_k^2)) \right) (\operatorname{Re} \bar{\beta}_k (\psi_k^1 + \psi_k^2)) dx \\ &+ 2 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi_k^1 + \psi_k^2|^2 \right) |\beta_k|^2 dx, \end{aligned} \quad (\text{A.43})$$

$$I_3 = 4 \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\beta_k|^2 \right) (\operatorname{Re} \bar{\beta}_k (\psi_k^1 + \psi_k^2)) dx, \quad (\text{A.44})$$

$$I_4 = \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\beta_k|^2 \right) |\beta_k|^2 dx. \quad (\text{A.45})$$

To estimate I_0 , we notice that if k is sufficiently large then ψ_k^1 and ψ_k^2 have disjoint supports receding from each other, i.e.,

$$d_k := \operatorname{dist}(\operatorname{supp} \psi_k^1, \operatorname{supp} \psi_k^2) \rightarrow \infty, \quad \text{as } k \rightarrow \infty; \quad (\text{A.46})$$

see the proof of Lemma 2.4 in Sect. A.1. Thus, the last three terms of the right-hand side in (A.41) equal 0 if k is large, since $\bar{\psi}_k^1 \psi_k^2 = 0$ a. e. if k is sufficiently large. Also by (A.46), we infer

$$\begin{aligned} &\left| \int_{\mathbb{R}^3 \times \mathbb{R}^3} |\psi_k^1(x)|^2 \frac{1}{|x-y|} |\psi_k^2(y)|^2 dx dy \right| \\ &= \left| \int_{\mathbb{R}^3 \times \mathbb{R}^3} |\psi_k^1(x)|^2 \frac{\chi_{\{|x-y| \geq d_k\}}(x-y)}{|x-y|} |\psi_k^2(y)|^2 dx dy \right| \\ &\leq \|\psi_k^1\|_2^2 \|\psi_k^2\|_2^2 \| |x|^{-1} \chi_{\{|x| \geq d_k\}}(x) \|_\infty \leq \frac{C}{d_k} \rightarrow 0, \quad \text{as } k \rightarrow \infty, \end{aligned} \quad (\text{A.47})$$

using Young's inequality. Thus we have shown that

$$|I_0| \leq r_1(k) \rightarrow 0, \quad \text{as } k \rightarrow \infty. \quad (\text{A.48})$$

The remaining terms I_1 – I_4 are controlled by the Hardy-Littlewood-Sobolev inequality and Hölder's inequality as follows:

$$|I_1| \leq C(\|\psi_k^1\|_{12/5}^3 + \|\psi_k^2\|_{12/5}^3)\|\beta_k\|_{12/5}, \quad (\text{A.49})$$

$$|I_2| \leq C(\|\psi_k^1\|_{12/5}^2 + \|\psi_k^2\|_{12/5}^2)\|\beta_k\|_{12/5}^2, \quad (\text{A.50})$$

$$|I_3| \leq C(\|\psi_k^1\|_{12/5} + \|\psi_k^2\|_{12/5})\|\beta_k\|_{12/5}^3, \quad (\text{A.51})$$

$$|I_4| \leq C\|\beta\|_{12/5}^4. \quad (\text{A.52})$$

We notice that $\|\psi_k^1\|_{12/5}$ and $\|\psi_k^2\|_{12/5}$ are uniformly bounded, by Sobolev's inequality and the $H^{1/2}$ -boundedness of these sequences. Furthermore, we have that

$$\|\beta_k\|_{12/5} \leq r_2(\epsilon) \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0, \quad (\text{A.53})$$

by part iii) of Lemma 2.4. Hence we conclude that

$$|I_1 + \cdots + I_4| \leq r_2(\epsilon) \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0, \quad (\text{A.54})$$

which proves Lemma A.2. \square

A.3. Commutator Estimate. An almost identical result is needed in [5], but we provide its proof again.

Lemma A.3. *Let $m \geq 0$, $v \in \mathbb{R}^3$, and define $A_v := \sqrt{-\Delta + m^2} + i(v \cdot \nabla)$. Furthermore, suppose that $f(x)$ is a locally integrable and that its distributional gradient, ∇f , is an $L^\infty(\mathbb{R}^3)$ vector-valued function. Then we have that*

$$\|[A_v, f]\|_{L^2 \rightarrow L^2} \leq C_v \|\nabla f\|_\infty,$$

for some constant C_v that only depends on v .

Remark. This result can be deduced by means of Calderón–Zygmund theory for singular integral operators and its consequences for pseudo-differential operators (see, e. g., [13, Sect. VII.3]). We give an elementary proof which makes good use of the spectral theorem, enabling us to write the commutator in a convenient way.

Proof (of Lemma A.3). Since $[i(v \cdot \nabla), f] = iv \cdot \nabla f$ holds, we have that

$$\|[i(v \cdot \nabla), f]\|_{L^2 \rightarrow L^2} \leq |v| \|\nabla f\|_\infty. \quad (\text{A.55})$$

Thus, it suffices to prove our assertion for $A := A_{v=0}$, i. e.,

$$A := \sqrt{p^2 + m^2}, \quad \text{where } p = -i\nabla. \quad (\text{A.56})$$

Since A is a self-adjoint operator on $L^2(\mathbb{R}^3)$ (with domain $H^1(\mathbb{R}^3)$), functional calculus (for measurable functions) yields the formula

$$A^{-1} = \frac{1}{\pi} \int_0^\infty \frac{1}{\sqrt{s}} \frac{ds}{A^2 + s}. \quad (\text{A.57})$$

Due to this fact and $A = A^2 A^{-1}$, we obtain the formula

$$[A, f] = \frac{1}{\pi} \int_0^\infty \frac{\sqrt{s}}{A^2 + s} [A^2, f] \frac{ds}{A^2 + s}. \quad (\text{A.58})$$

Clearly, we have that $[A^2, f] = [p^2, f] = p \cdot [p, f] + [p, f] \cdot p$, which leads to

$$[A, f] = \frac{1}{\pi} \int_0^\infty \frac{\sqrt{s}}{p^2 + m^2 + s} (p \cdot [p, f] + [p, f] \cdot p) \frac{ds}{p^2 + m^2 + s}. \quad (\text{A.59})$$

Moreover, since $[p, f] = -i \nabla f$ holds, we have that

$$\left\| \left[\frac{1}{p^2 + m^2 + s}, [p, f] \right] \right\|_{L^2 \rightarrow L^2} \leq \frac{2}{s} \|\nabla f\|_\infty. \quad (\text{A.60})$$

Hence we find, for arbitrary test functions $\xi, \eta \in C_c^\infty(\mathbb{R}^3)$, that

$$\begin{aligned} & \left| \langle \xi, \int_0^\infty \frac{\sqrt{s}}{p^2 + m^2 + s} ([p, f] \cdot p) \frac{ds}{p^2 + m^2 + s} \eta \rangle \right| \\ & \leq \left| \langle [p, f] \xi, p \int_0^\infty \frac{\sqrt{s} ds}{(p^2 + m^2 + s)^2} \eta \rangle \right| \\ & \quad + \left| \langle \xi, \int_0^\infty \left[\frac{1}{p^2 + m^2 + s}, [p, f] \right] \cdot \frac{p \sqrt{s} ds}{p^2 + m^2 + s} \eta \rangle \right| \\ & \leq \|[p, f] \xi\|_2 \left\| \int_0^\infty \frac{p \sqrt{s} ds}{(p^2 + m^2 + s)^2} \eta \right\|_2 \\ & \quad + 2 \|\xi\|_2 \|\nabla f\|_\infty \left\| \int_0^\infty \frac{p ds}{\sqrt{s}(p^2 + m^2 + s)} \eta \right\|_2. \end{aligned} \quad (\text{A.61})$$

Evaluation of the s -integrals yields

$$(\text{A.61}) \leq C \|\nabla f\|_\infty \|\xi\|_2 \left\| \frac{p}{\sqrt{p^2 + m^2}} \eta \right\|_2 \leq C \|\nabla f\|_\infty \|\xi\|_2 \|\eta\|_2. \quad (\text{A.62})$$

The same estimate holds if $[p, f] \cdot p$ is replaced by $p \cdot [p, f]$ in (A.61). Thus, we have found that

$$|\langle \xi, [A, f] \eta \rangle| \leq C \|\nabla f\|_\infty \|\xi\|_2 \|\eta\|_2, \quad \text{for } \xi, \eta \in C_c^\infty(\mathbb{R}^3), \quad (\text{A.63})$$

with some constant C independent of m . Since $C_c^\infty(\mathbb{R}^3)$ is dense in $L^2(\mathbb{R}^3)$, the assertion for the L^2 -boundedness of $[A, f]$ now follows. This completes the proof of Lemma A.3. \square

A.4. Lower Semicontinuity.

Lemma A.4. *Suppose that $m > 0$, $v \in \mathbb{R}^3$, with $|v| < 1$. Then the functional*

$$\mathcal{T}(\psi) := \langle \psi, (\sqrt{-\Delta + m^2}) \psi \rangle + \langle \psi, i(v \cdot \nabla) \psi \rangle$$

is weakly lower semicontinuous on $H^{1/2}(\mathbb{R}^3)$, i. e., if $\psi_k \rightharpoonup \psi$ weakly in $H^{1/2}(\mathbb{R}^3)$ as $k \rightarrow \infty$, then

$$\liminf_{k \rightarrow \infty} \mathcal{T}(\psi_k) \geq \mathcal{T}(\psi).$$

Moreover, if $\lim_{k \rightarrow \infty} \mathcal{T}(\psi_k) = \mathcal{T}(\psi)$ holds, then $\psi_k \rightarrow \psi$ strongly in $H^{1/2}(\mathbb{R}^3)$ as $k \rightarrow \infty$.

Proof (of Lemma A.4). Assume that $m > 0$, $v \in \mathbb{R}^3$, with $|v| < 1$ holds. By Fourier transform and Plancherel's theorem, we have that

$$\mathcal{T}(\psi) = \int_{\mathbb{R}^3} |\widehat{\psi}(k)|^2 \left(\sqrt{k^2 + m^2} - (v \cdot k) \right) dk. \quad (\text{A.64})$$

We notice that

$$c_1(|k| + m) \leq \sqrt{k^2 + m^2} - (v \cdot k) \leq c_2(|k| + m), \quad (\text{A.65})$$

for some suitable constants $c_1, c_2 > 0$, where the lower bound follows from the inequality $\sqrt{k^2 + m^2} \geq (1 - \delta)|k| + \delta m$, with $0 < \delta < 1$, and the fact that $|v| < 1$ holds. Thus,

$$\|\psi\|_{\mathcal{T}} := \sqrt{\mathcal{T}(\psi)} \quad (\text{A.66})$$

defines a norm that is equivalent to $\|\cdot\|_{\text{H}^{1/2}}$. Consequently, the notion of weak and strong convergence for these norms coincide. Finally, by (A.64), we identify $\|\psi\|_{\mathcal{T}}$ with the L^2 -norm of $\widehat{\psi}$ taken with respect to the integration measure

$$d\mu = \left(\sqrt{k^2 + m^2} - (v \cdot k) \right) dk. \quad (\text{A.67})$$

The assertion of Lemma A.4 now follows from corresponding properties of the $L^2(\mathbb{R}^3, \mu)$ -norm; see, e. g., [9, Theorem 2.11] for $L^p(\Omega, \mu)$ -norms, where Ω is a measure space with positive measure, μ , and $1 < p < \infty$. \square

B. Best Constant and Optimizers for Inequality (2.5)

Lemma B.1. *For any $v \in \mathbb{R}^3$ with $|v| < 1$, there exists an optimal constant, S_v , such that*

$$\int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx \leq S_v \langle \psi, (\sqrt{-\Delta} + iv \cdot \nabla) \psi \rangle \langle \psi, \psi \rangle \quad (\text{B.1})$$

holds for all $\psi \in \text{H}^{1/2}(\mathbb{R}^3)$. Moreover, we have that

$$S_v = \frac{2}{\langle Q_v, Q_v \rangle}, \quad (\text{B.2})$$

where $Q_v \in \text{H}^{1/2}(\mathbb{R}^3)$, $Q_v \neq 0$, is an optimizer for (B.1) and it satisfies

$$\sqrt{-\Delta} Q_v + i(v \cdot \nabla) Q_v - \left(\frac{1}{|x|} * |Q_v|^2 \right) Q_v = -Q_v. \quad (\text{B.3})$$

In addition, the following estimates hold:

$$S_{v=0} < \frac{\pi}{2} \quad \text{and} \quad S_{v=0} \leq S_v \leq (1 - |v|)^{-1} S_{v=0}. \quad (\text{B.4})$$

Proof (of Lemma B.1). Let $v \in \mathbb{R}^3$ with $|v| < 1$ be fixed and consider the unconstrained minimization problem

$$\frac{1}{S_v} := \inf_{\psi \in \text{H}^{1/2}(\mathbb{R}^3), \psi \neq 0} \frac{\langle \psi, (\sqrt{-\Delta} + iv \cdot \nabla) \psi \rangle \langle \psi, \psi \rangle}{\int_{\mathbb{R}^3} \left(|x|^{-1} * |\psi|^2 \right) |\psi|^2 dx}. \quad (\text{B.5})$$

For $v = 0$, a variational problem equivalent to (B.5) is studied in [11, App. B] by using strict rearrangement inequalities that allow restriction to radial functions. For $v \neq 0$, we

have to depart this line of argumentation and we employ (similarly to the discussion of (2.1) in Sect. 2) concentration-compactness-type methods.

By scaling properties of (B.5), it suffices to prove the existence of a minimizer with $\langle \psi, (\sqrt{-\Delta} + iv \cdot \nabla) \psi \rangle$ and $\langle \psi, \psi \rangle$ fixed. Thus, we introduce the constrained minimization problem, which is equivalent to (B.5), as follows:

$$I_v(\alpha, \beta) := \inf \left\{ - \int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx : \langle \psi, \psi \rangle = \alpha, \right. \\ \left. \langle \psi, (\sqrt{-\Delta} + iv \cdot \nabla) \psi \rangle = \beta \right\}, \quad (\text{B.6})$$

where $\alpha > 0$ and $\beta > 0$. In particular, it is sufficient to show that $I_v(\alpha = 1, \beta = 1)$ is finite and attained so that

$$S_v = -I_v(1, 1). \quad (\text{B.7})$$

In fact, we will show that all minimizing sequences for $I(1, 1)$ are relatively compact in $H^{1/2}(\mathbb{R}^3)$ up to translations. In turn, this relative compactness implies that all minimizing sequences for problem (B.5) are relatively compact in $H^{1/2}(\mathbb{R}^3)$ up to translations and rescalings: For any minimizing sequence, (ψ_n) , for (B.5), there exist sequences, $\{(y_k), (a_k), (b_k)\}$, with $y_k \in \mathbb{R}^3$, $0 \neq a_k \in \mathbb{C}$, $0 \neq b_k \in \mathbb{R}$, such that

$$a_k \psi_{n_k}(b_k(\cdot + y_k)) \rightarrow Q_v \quad \text{strongly in } H^{1/2}(\mathbb{R}^3) \text{ as } k \rightarrow \infty, \quad (\text{B.8})$$

along a suitable subsequence, (ψ_{n_k}) , and Q_v minimizes (B.5).

First we show that $I(\alpha, \beta)$ is indeed finite. The Hardy–Littlewood–Sobolev inequality implies

$$\int_{\mathbb{R}^3} \left(\frac{1}{|x|} * |\psi|^2 \right) |\psi|^2 dx \leq C \|\psi\|_{6/5}^2 = C \|\psi\|_{12/5}^4 \leq C \langle \psi, \sqrt{-\Delta} \psi \rangle \langle \psi, \psi \rangle, \quad (\text{B.9})$$

where we use Sobolev's inequality $\|\psi\|_3^2 \leq C \langle \psi, \sqrt{-\Delta} \psi \rangle$ in \mathbb{R}^3 and Hölder's inequality. Since $\langle \psi, \sqrt{-\Delta} \psi \rangle \leq (1 - |v|)^{-1} \langle \psi, (\sqrt{-\Delta} + iv \cdot \nabla) \psi \rangle$, we deduce that

$$I(\alpha, \beta) \geq -C\alpha\beta > -\infty, \quad (\text{B.10})$$

for some constant C . On the other hand, we have that

$$I(\alpha, \beta) < 0, \quad (\text{B.11})$$

since $\int_{\mathbb{R}^3} (|x|^{-1} * |\psi|^2) |\psi|^2 dx \neq 0$ when $\psi \not\equiv 0$.

Next, we show that $I_v(1, 1)$ is attained. Let (ψ_n) be a minimizing sequence for $I_v(1, 1)$. In order to invoke Lemma 2.4, we notice that $\int_{\mathbb{R}^3} |\psi_n|^2 dx = 1$ and that (ψ_n) is bounded in $H^{1/2}(\mathbb{R}^3)$, since $\langle \psi, (\sqrt{-\Delta} + iv \cdot \nabla) \psi \rangle$ is equivalent to $\langle \psi, \sqrt{-\Delta} \psi \rangle$ when $|v| < 1$, by (A.65) with $m = 0$.

Let us suppose now that case ii) of Lemma 2.4 occurs. Referring to Lemma A.1, we conclude that $I(1, 1) = 0$ holds, which contradicts (B.11). Next, let us assume that dichotomy occurs for a subsequence of (ψ_n) , i.e., property iii) of Lemma 2.4 holds. Using Lemma A.2 and the lim inf-estimate stated in iii) of Lemma 2.4 and by taking the limit $\epsilon \rightarrow 0$, we conclude that

$$I_v(1, 1) \geq I_v(\alpha, \beta) + I_v(1 - \alpha, 1 - \beta), \quad (\text{B.12})$$

for some $\alpha \in (0, 1)$ and $\beta \in [0, 1]$. On the other hand, we have the scaling behaviour

$$I_v(\alpha, \beta) = \alpha\beta I_v(1, 1) < 0, \quad (\text{B.13})$$

which follows from (B.6) and rescaling $\psi(x) \mapsto a\psi(bx)$ with $a, b > 0$. Combining (B.12) with (B.13) we get a contradiction. Therefore dichotomy for minimizing sequences is ruled out.

In summary, we see that any minimizing sequence, (ψ_n) , for $I_v(1, 1)$ contains a subsequence, (ψ_{n_k}) , with a sequence of translations, (y_k) , satisfying property i) of Lemma 2.4. Similarly to the proof of Theorem 2.1, we conclude that $\psi_{n_k}(\cdot + y_k) \rightarrow \tilde{Q}_v$ strongly in $H^{1/2}(\mathbb{R}^3)$ as $k \rightarrow \infty$, where $\tilde{Q}_v \in H^{1/2}(\mathbb{R}^3)$ is a minimizer for $I_v(1, 1)$.

To show that the best constant, S_v , is given by (B.2) with Q_v minimizing (B.5) and satisfying (B.3), let us denote the minimizer constructed above for $I(1, 1)$ by \tilde{Q}_v . Since \tilde{Q}_v also minimizes the unconstrained problem (B.5), it has to satisfy the corresponding Euler-Lagrange equation which reads as follows:

$$\sqrt{-\Delta} \tilde{Q}_v + iv \cdot \nabla \tilde{Q}_v - \frac{2}{S_v} \left(\frac{1}{|x|} * |\tilde{Q}_v|^2 \right) \tilde{Q}_v + \tilde{Q}_v = 0, \quad (\text{B.14})$$

where we use that $\langle \tilde{Q}_v, (\sqrt{-\Delta} + iv \cdot \nabla) \tilde{Q}_v \rangle = 1$ and $\langle \tilde{Q}_v, \tilde{Q}_v \rangle = 1$ holds. By putting $Q_v = \sqrt{2} S_v^{-1/2} \tilde{Q}_v$, we see that Q_v minimizes (B.5) and satisfies (B.2). Moreover, we have that $\langle Q_v, Q_v \rangle = 2/S_v$ holds.

Finally, we turn to the estimates for S_v stated in Lemma B.1. That $S_{v=0} < \pi/2$ holds follows from the appendices in [11, 7]. To see that $S_v \leq (1 - |v|)^{-1} S_{v=0}$ is true, we use the estimate $\sqrt{-\Delta} \leq (1 - |v|)^{-1} (\sqrt{-\Delta} + iv \cdot \nabla)$. Moreover, it is known from the discussion in [7] that if $v = 0$ the minimizer, $Q_{v=0}$, for (2.2) can be chosen to be radial (by symmetric rearrangement). This implies that $\langle Q_{v=0}, \nabla Q_{v=0} \rangle = 0$, which leads to $S_{v=0} \leq S_v$. \square

C. Exponential Decay

In this section, we prove pointwise exponential decay for solutions, $\varphi \in H^{1/2}(\mathbb{R}^3)$, of the nonlinear equation

$$(\sqrt{-\Delta + m^2} - m)\varphi + i(v \cdot \nabla)\varphi - \left(\frac{1}{|x|} * |\varphi|^2 \right)\varphi = -\mu\varphi. \quad (\text{C.1})$$

Clearly, $\varphi(x)$ is an eigenfunction for the Schrödinger type operator

$$H = H_0 + V, \quad (\text{C.2})$$

where

$$H_0 := (\sqrt{-\Delta + m^2} - m) + i(v \cdot \nabla) \quad \text{and} \quad V := -\left(\frac{1}{|x|} * |\varphi|^2 \right). \quad (\text{C.3})$$

By using the bootstrap argument for regularity (presented in the proof of Theorem 4.1)), we have that $\varphi \in H^s(\mathbb{R}^3)$ for all $s \geq 1/2$, which shows in particular that φ is smooth. Investigating the spectrum of H_0 we find that

$$\sigma(H_0) = \sigma_{\text{ess}}(H_0) = [\Sigma_v, \infty), \quad (\text{C.4})$$

where the bottom of the spectrum is given by

$$\Sigma_v = (\sqrt{1 - v^2} - 1)m. \quad (\text{C.5})$$

To see this, we remark that the function

$$f(k) = (k^2 + m^2)^{1/2} - m - v \cdot k \quad (\text{C.6})$$

obeys $f(k) \geq (\sqrt{1-v^2})m - m$ with equality for $k = (mv/\sqrt{1-v^2})$.

We have the following result.

Lemma C.1. *Suppose that $m > 0$, $v \in \mathbb{R}^3$, and $|v| < 1$. Furthermore, let $\varphi \in \mathbf{H}^{1/2}(\mathbb{R}^3)$ be a solution of (C.1) with $-\mu < \Sigma_v$. Then, for every $0 < \delta < \min\{m, \frac{\Sigma_v + \mu}{\sqrt{1-v^2}}\}$, there exists $0 < C(\delta) < \infty$ such that*

$$|\varphi(x)| \leq C e^{-\delta|x|}$$

holds for all $x \in \mathbb{R}^3$.

Proof (of Lemma C.1). We rewrite (C.1) as follows :

$$\varphi = -(H_0 + \mu)^{-1} V \varphi, \quad (\text{C.7})$$

where H_0 and V are defined in (C.3). Note that $(H_0 + \mu)^{-1}$ exists, since we have that $\mu \notin \sigma(H_0)$ holds, by the assumption that $-\mu < \Sigma_v$. We consider the Green's function, $G_\mu(x - y)$, given by

$$G_\mu(x - y) = \mathcal{F}^{-1} \left[\frac{1}{\sqrt{k^2 + m^2} - m - v \cdot k + \mu} \right] (x - y), \quad (\text{C.8})$$

where $\mathcal{F} : \mathcal{S}' \rightarrow \mathcal{S}'$ denotes the Fourier transform. Since $1/\sqrt{k^2 + m^2} \dots$ does not belong to $L^1(\mathbb{R}^3)$, we cannot use Payley–Wiener type theorems directly to deduce pointwise exponential decay for $G_\mu(z)$ in $|z|$. To overcome this difficulty, we first notice that

$$(H_0 + \mu)^{-1} = \int_0^\infty e^{-t\mu} e^{-tH_0} dt = \int_0^\infty e^{-t(\mu-m)} e^{-t(\sqrt{p^2+m^2}-v \cdot p)} dt, \quad (\text{C.9})$$

by self-adjointness of H_0 and functional calculus. Here and in what follows, we put $p = -i\nabla$ for convenience. By using the explicit formula for the Fourier transform of $\exp\{-t\sqrt{k^2 + m^2}\}$ (see e. g., [9]) in \mathbb{R}^3 and by analytic continuation, we obtain from (C.9) the formula

$$G_\mu(z) = A_m \int_0^\infty e^{-t(\mu-m)} \frac{t}{t^2 + (z + itv)^2} K_2(m\sqrt{t^2 + (z + itv)^2}) dt. \quad (\text{C.10})$$

Here $K_2(z)$ stands for the modified Bessel function of the third kind, and $A_m > 0$ denotes some constant. Notice that

$$w = t^2 + (z + itv)^2 = (1 - v^2)t^2 + z^2 + 2itv \cdot z \quad (\text{C.11})$$

is a complex number with $|\arg w| < \pi/2$.

Next we analyze $G_\mu(z)$ for $|z| \leq 1/m$ and for $|z| > 1/m$ separately. From [1] we recall the estimate

$$|K_2(mw)| \leq \frac{C}{|w|^2}, \quad \text{for } |\arg w| < \pi/2, \quad (\text{C.12})$$

which implies that $G_\mu(z)$ with $|z| \leq 1/m$ satisfies the bound

$$\begin{aligned} |G_\mu(z)| &\leq C \int_0^\infty e^{-t(\mu-m)} \left| \frac{t}{(1-v^2)t^2 + |z|^2 + 2itv \cdot z} \right. \\ &\quad \left. K_2(m\sqrt{(1-v^2)t^2 + |z|^2 + 2itv \cdot z}) \right| dt \\ &\leq C \int_0^\infty e^{-t(\mu-m)} \frac{t}{[(1-v^2)t^2 + |z|^2]^2} dt. \end{aligned} \quad (\text{C.13})$$

Since $\mu - m \geq 0$, the t -integral is finite for $z \neq 0$ and we obtain

$$|G_\mu(z)| \leq \frac{C}{|z|^2}, \quad \text{for } |z| \leq 1/m, \quad (\text{C.14})$$

where we use that $|a + ib| \geq |a|$ and $|\sqrt{a + ib}| \geq \sqrt{|a|}$ holds for $a, b \in \mathbb{R}$.

To estimate $G_\mu(z)$ for $|z| > 1/m$, we use the bound

$$|K_2(mw)| \leq C \left| \frac{e^{-mw}}{|w|^2} \right| \leq C \frac{e^{-m|\operatorname{Re} w|}}{|w|^2}, \quad \text{for } |\arg w| < \pi/2 \text{ and } |w| > 1, \quad (\text{C.15})$$

taken from [1]. By means of the inequality $\sqrt{a^2 + b^2} \geq (1 - \epsilon)|a| + \epsilon|b|$, for any $0 < \epsilon \leq 1$, we proceed to find that

$$|G_\mu(z)| \leq C e^{-\epsilon m|z|} \int_0^\infty e^{-t(\mu-m+(1-\epsilon)\sqrt{1-v^2}m)} \frac{t}{[(1-v^2)t^2 + |z|^2]^2} dt, \quad (\text{C.16})$$

for $|z| > 1/m$. The assumption on μ allows us to choose $\epsilon \in (0, 1]$ such that the exponent in the t -integral is nonpositive. The best ϵ is given by

$$\epsilon = \min \left\{ 1, \frac{\Sigma_v + \mu}{m\sqrt{1-v^2}} \right\} \in (0, 1], \quad (\text{C.17})$$

and hence

$$\begin{aligned} |G_\mu(z)| &\leq C e^{-\epsilon m|z|} \int_0^\infty \frac{t}{[(1-v^2)t^2 + |z|^2]^2} dt \\ &\leq C \frac{e^{-\epsilon m|z|}}{|z|^2}, \quad \text{for } |z| > 1/m. \end{aligned} \quad (\text{C.18})$$

Combining now (C.14) and (C.18), we see that

$$|G_\mu(z)| \leq C \frac{e^{-m\epsilon|z|}}{|z|^2}, \quad \text{for } z \in \mathbb{R}^3, \quad (\text{C.19})$$

where ϵ is given by (C.17) and C is some constant. This shows that $G_\mu(z)$ exhibits exponential decay; in particular, we have that $G_\mu \in L^p(\mathbb{R}^3)$ if $1 \leq p < 3/2$.

Returning to (C.7), we notice that

$$\varphi(x) = - \int_{\mathbb{R}^3} G_\mu(x-y)V(y)\varphi(y) dy. \quad (\text{C.20})$$

Moreover, the function $V(x) = -(|x|^{-1} * |\varphi|^2)(x)$ obeys

$$V \in C^0(\mathbb{R}^3) \quad \text{and} \quad \lim_{|x| \rightarrow \infty} V(x) = 0, \quad (\text{C.21})$$

since $f * g$ is a continuous function vanishing at infinity, provided that $f \in L^p$ and $g \in L^{p'}$ with $1/p + 1/p' = 1$ and $p > 1$; see, e. g., [9]. Here we note that, e. g., $|x|^{-1} \in L^2(\mathbb{R}^3) + L^4(\mathbb{R}^3)$ and in particular $|\varphi|^2 \in L^{4/3}(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$ since $\varphi \in H^s(\mathbb{R}^3)$ for all $s \geq 1/2$ (cf. beginning of App. C).

Using (C.20), (C.19) and (C.21), the claimed pointwise exponential decay of $\varphi(x)$ follows from a direct adaptation of an argument by Slaggie and Wichmann for exponential decay of eigenfunctions for Schrödinger operators; see, e. g., [6] for a convenient exposition of this method. This completes the proof of Lemma C.1. \square

Acknowledgement. The authors are grateful to I. M. Sigal and M. Struwe for useful discussions, as well as the referee for some helpful comments. Lastly, E. L. also thanks D. Christodoulou.

References

1. Abramowitz, M., Stegun, I.A.: ed.: *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. New York: Dover Publications Inc., 1992, Reprint of the 1972 edition
2. Cazenave, T., Lions, P.-L.: Orbital stability of standing waves for some nonlinear Schrödinger equations. *Commun. Math. Phys.* **85**, 549–561 (1982)
3. Elgart, A., Schlein, B.: Mean field dynamics of Boson Stars. *Comm. Pure Appl. Math.* **60**(4), 500–545 (2006)
4. Fröhlich, J., Jonsson, B.L.G., Lenzmann, E.: Effective dynamics for Boson stars. *Nonlinearity* **32**, 1031–1075 (2007)
5. Fröhlich, J., Lenzmann, E.: *Blow-up for nonlinear wave equations describing Boson Stars*. <http://arxiv.org/list/math-ph/0511003>, 2006 to appear in *Comm. Pure Appl. Math.*
6. Hislop, P.D.: Exponential decay of two-body eigenfunctions: A review. In: *Proceedings of the Symposium on Mathematical Physics and Quantum Field Theory (Berkeley, CA, 1999)*, Volume 4 of *Electron. J. Differ. Equ. Conf.*, pages 265–288 (electronic), San Marcos, TX, 2000. Southwest Texas State Univ.
7. Lenzmann, E.: *Well-posedness for semi-relativistic Hartree equations of critical type*. <http://arXiv.org/list/math.AP/0505456>, 2005, to appear in *Mathematical Physics, Analysis, and Geometry*
8. Lieb, E.H.: Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation. *Stud. Appl. Math.* **57**, 93–105 (1977)
9. Lieb, E.H., Loss, M.: *Analysis, Volume 14 of Graduate Studies in Mathematics*. A Providence, RI: Amer. Math. Soc., second edition, 2001
10. Lieb, E.H., Thirring, W.: Gravitational collapse in quantum mechanics with relativistic kinetic energy. *Ann. Physics* **155**(2), 494–512 (1984)
11. Lieb, E.H., Yau, H.-T.: The Chandrasekhar theory of stellar collapse as the limit of quantum mechanics. *Commun. Math. Phys.* **112**, 147–174 (1987)
12. Lions, P.-L.: The concentration-compactness principle in the calculus of variations. The locally compact case, Part I. *Ann. Inst. Henri Poincaré*, **1**(2), 109–145 (1984)
13. Stein, E.M.: *Harmonic Analysis*. Princeton, NJ: Princeton University Press, 1993

Communicated by H.-T. Yau