

## Boundary value problems for evolution inclusions

by NIKOLAOS S. PAPAGEORGIOU (Davis, California, U.S.A.)

**Abstract.** This paper examines boundary value problems for evolution inclusions, with nonlinear boundary conditions. Two existence theorems are proved. One for convex multivalued perturbations and the other for nonconvex ones. Finally an example from partial differential equations is presented.

**1. Introduction.** In this paper we study boundary value problems for evolution inclusions. Our work was motivated by the papers of Anichini [1], Kartsatos [6] and Zecca-Zezza [13]. Anichini [1], considered quasilinear differential equations in  $\mathbf{R}^n$ , with nonlinear boundary conditions and using a fixed point theorem due to Eilenberg-Montgomery, established the existence of solutions. Kartsatos [6] too considers boundary value problems for  $\mathbf{R}^n$ -valued differential equations, but over an unbounded time interval. Finally Zecca-Zezza [13] extend the work of Kartsatos to differential equations in Banach spaces.

In this note, the differential inclusion is defined on a compact time interval and this allows us to weaken considerably the hypotheses on the orientor field  $F(t, x)$ . Furthermore, contrary to Zecca-Zezza [13], here the linear operator is in general unbounded, covering this way the very important case of partial differential operators. Also we establish the existence of solutions for problems with nonconvex multivalued perturbations, a case which is not addressed in the paper of Zecca-Zezza [13]. Finally we present an application to partial differential equations.

**2. Preliminaries.** Let  $(\Omega, \Sigma)$  be a measurable space and  $X$  a separable Banach space. Throughout this paper we will be using the following notations:

$$P_{f(c)}(X) = \{A \subseteq X: \text{nonempty, closed, (convex)}\}$$

---

1980 AMS Subject Classification 35K35.

Research supported by N.S.F. Grant D.M.S. 8602313.

Work done while on leave at the University of Thessaloniki, School of Technology, Mathematics Division, Thessaloniki 54006, Greece.

and

$$P_{(w)k(c)}(X) = \{A \subseteq X : \text{nonempty, (weakly-)compact, (convex)}\}.$$

A multifunction  $F: \Omega \rightarrow P_f(X)$  is said to be *measurable* if, for all  $z \in X$ ,  $\omega \rightarrow d(z, F(\omega)) = \inf\{\|z-x\|: x \in F(\omega)\}$  is measurable. Other equivalent definitions of measurability of multifunctions can be found in Wagner [12]. By  $S_F^1$  we denote the set of  $L^1(X)$ -selectors of  $F(\cdot)$ , i.e.  $S_F^1 = \{f \in L^1(X): f(\omega) \in F(\omega) \text{ } \mu\text{-a.e.}\}$ . This set may be empty. It is nonempty if and only if  $\omega \rightarrow \inf\{\|x\|: x \in F(\omega)\}$  belongs in  $L^1_+$ . Using  $S_F^1$  we can define a set valued integral for  $F(\cdot)$ , by setting  $\int_{\Omega} F = \{\int_{\Omega} f: f \in S_F^1\}$ .

Next let  $Y, Z$  be Hausdorff topological spaces. Let  $G: Y \rightarrow 2^Z \setminus \{\emptyset\}$  be a multifunction. We say that  $G(\cdot)$  is *upper semicontinuous* (u.s.c.) (resp. *lower semicontinuous* (l.s.c.)) if for all  $U \subseteq Z$  open  $F^+(U) = \{y \in Y: F(y) \subseteq U\}$  is open in  $Y$  (resp.  $F^-(U) = \{y \in Y: F(y) \cap U \neq \emptyset\}$  is open in  $Y$ ).

**3. Existence result: convex case.** Let  $T = [0, b]$  and  $X$  be a separable Banach space. The multivalued boundary value problem under consideration is the following:

$$(*) \quad \dot{x}(t) \in A(t)x(t) + F(t, x(t)), \quad Lx = Mx.$$

We will assume that the family of linear operators  $\{A(t): t \in T\}$  generates a strongly continuous evolution operator  $S(t, s)$ ,  $0 \leq s \leq t \leq b$ . So by a solution of  $(*)$ , we will understand a mild solution. Thus we say that  $x(\cdot) \in C(T, X)$  solves  $(*)$  if

$$x(t) = S(t, 0)x(0) + \int_0^t S(t, s)f(s)ds \quad \text{for some } f \in S_{F(t, x(\cdot))}^1$$

and

$$Lx = Mx.$$

The full set of hypotheses on the data of problem  $(*)$  is the following:

$H(A)$ . The family  $\{A(t): t \in [0, b]\}$  generates a strongly continuous evolution operator  $S: A = \{0 \leq s \leq t \leq b\} \rightarrow \mathcal{L}(X)$ , which is compact for  $t-s > 0$ .

$H(F)$ .  $F: T \times X \rightarrow P_{wkc}(X)$  is a multifunction s.t.

$$(1) \quad (t, x) \rightarrow F(t, x) \quad \text{is measurable,}$$

$$(2) \quad x \rightarrow F(t, x) \quad \text{is u.s.c. from } X \text{ into } X_w,$$

$$(3) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \int_0^b \sup_{\|x\| \leq n} |F(t, x)| dt = 0.$$

$H(L)$ .  $L: C(T, X) \rightarrow X$  is continuous, linear. Also if  $V: C(T, X) \rightarrow C(T, X)$  is defined by  $(Vx)(\cdot) = x(\cdot) - S(\cdot, 0)x(0)$ , then there exists  $K: X \rightarrow \ker V$  continuous, linear s.t.  $(I - L_0 K)(Mx - L \int_0^t S(t, s) f(s) ds) = 0$  for all  $f \in S_{F(\cdot, x(\cdot))}^1$  and all  $x(\cdot) \in C(T, X)$ , with  $L_0 = L|_{\ker V}$ .

$H(M)$ .  $M: C(T, X) \rightarrow X$  is a generally nonlinear, completely continuous operator s.t.

$$\lim_{\|x\| \rightarrow x} \frac{\|Mx\|}{\|x\|} = 0.$$

Having these hypotheses, we can now state our first existence result concerning (\*).

**THEOREM 1.** *If hypotheses  $H(A)$ ,  $H(F)$ ,  $H(L)$  and  $H(M)$  hold, then (\*) admits a mild solution.*

**Proof.** For some  $x_0 \in \ker L_0$ , consider the multifunction  $R: C(T, X) \rightarrow 2^{C(T, X) \setminus \{\emptyset\}}$  defined by:

$$R(x) = \left\{ y \in C(T, X) : y(t) = x_0(t) + KMx - KL \int_0^t S(t, s) f(s) ds + \int_0^t S(t, s) f(s) ds, t \in T, f \in S_{F(\cdot, x(\cdot))}^1 \right\}.$$

Because of hypothesis  $H(L)$ , it is easy to check that a fixed point of  $R(\cdot)$  is the desired mild solution of (\*).

From the definition of  $R(\cdot)$  and the convexity of the values of  $F(\cdot, \cdot)$  (and so of  $S_{F(\cdot, x(\cdot))}^1$ ), we see that  $R(\cdot)$  is convex valued. We claim that the values of  $R(\cdot)$  are also closed. So let  $y_n \in R(x)$ ,  $y_n \rightarrow y$  in  $C(T, X)$ . We have:

$$y_n(t) = x_0(t) + KMx - KL \int_0^t S(t, s) f_n(s) ds + \int_0^t S(t, s) f_n(s) ds$$

with  $f_n \in S_{F(\cdot, x(\cdot))}^1$ . But from Proposition 3.1 of [9], we know that  $S_{F(\cdot, x(\cdot))}^1$  is weakly compact in  $L^1(X)$  and by the Eberlein–Shmul'yan theorem is weakly sequentially compact. Thus by passing to a subsequence if necessary, we may assume that  $f_n \rightharpoonup f \in S_{F(\cdot, x(\cdot))}^1$  in  $L^1(X)$ . Then exploiting the fact that a continuous, linear operator is also weakly continuous and that

$$\int_0^t S(t, s) f_n(s) ds \rightharpoonup \int_0^t S(t, s) f(s) ds,$$

we get that

$$y_n(t) \rightharpoonup x_0(t) + KMx - KL \int_0^t S(t, s) f(s) ds + \int_0^t S(t, s) f(s) ds, \quad t \in T$$

and  $f \in S_{F(\cdot, x(\cdot))}^1$ . So

$$y(t) = x_0(t) + KMx - KL \int_0^t S(t, s) f(s) ds + \int_0^t S(t, s) f(s) ds \Rightarrow y \in R(x).$$

Hence we conclude that  $R(x) \in P_{fc}(C(T, X))$ .

Now we will show that  $R(\cdot)$  has closed graph ( $\text{Gr} R = \{(x, y) \in C(T, X) \times C(T, X) : y \in R(x)\} = \text{graph of } R(\cdot)$ ). To this end let  $(x_n, y_n) \in \text{Gr} R$ ,  $(x_n, y_n) \rightarrow (x, y)$  in  $C(T, X) \times C(T, X)$ . Then we have:

$$y_n(t) = x_0(t) + KMx_n - KL \int_0^t S(t, s) f_n(s) ds + \int_0^t S(t, s) f_n(s) ds,$$

with  $f_n \in S_{F(\cdot, x_n(\cdot))}^1$ . Let  $G(t) = \overline{\text{conv}} \bigcup_{n \geq 1} F(t, x_n(t))$ . Since by hypothesis  $H(F)(2)$ ,  $F(t, \cdot)$  is u.s.c. from  $X$  into  $X_w$ , it maps compact sets in  $X$  into  $w$ -compact sets. Therefore  $\bigcup_{n \geq 1} \overline{F(t, x_n(t))}^w$  is  $w$ -compact and by the Krein-Shmul'yan theorem we have that  $\overline{\text{conv}} \bigcup_{n \geq 1} F(t, x_n(t))$  is  $w$ -compact. So for all  $t \in T$ ,  $G(t) \in P_{wkc}(X)$ .

Finally from hypothesis  $H(F)(3)$ , we see that  $G(\cdot)$  is integrably bounded (i.e.  $t \rightarrow |G(t)| = \sup\{\|z\| : z \in G(t)\} \in L^1_+$ ). Hence once again Proposition 3.1 of [9] tells us that  $S_G^1$  is  $w$ -compact in  $L^1(X)$ . So by passing to a subsequence if necessary, we may assume that  $f_n \xrightarrow{w} f$  in  $L^1(X)$ . From Theorem 3.1 of [10], we have

$$f(t) \in \overline{\text{conv}} w - \overline{\lim} f_n(t) \subseteq \overline{\text{conv}} w - \overline{\lim} F(t, x_n(t)) \subseteq F(t, x(t)) \quad \text{a.e.}$$

the last inclusion following from hypothesis  $H(F)(2)$ . So  $f \in S_{F(\cdot, x(\cdot))}^1$ . Also note that

$$x_0(t) + KMx_n - KL \int_0^t S(t, s) f_n(s) ds + \int_0^t S(t, s) f_n(s) ds$$

converges weakly to

$$\begin{aligned} x_0(t) + KMx - KL \int_0^t S(t, s) f(s) ds + \int_0^t S(t, s) f(s) ds &= y(t) \Rightarrow y \in R(x) \\ &\Rightarrow \text{Gr} R \text{ is closed.} \end{aligned}$$

Next, we claim that there exists  $r > 0$  s.t. for

$$\|x\|_\infty \leq r \Rightarrow |R(x)| = \sup\{\|y\|_\infty : y \in R(x)\} \leq r.$$

Suppose not. Then, we can find  $\{x_n\}_{n \geq 1} \subseteq C(T, X)$  s.t.  $\|x_n\|_\infty \leq n$  and  $|R(x_n)| > n$ . So we have  $1 < |R(x_n)|/n$ . But note that for  $y_n \in R(x_n)$  we have:

$$\begin{aligned} \|y_n(t)\| &< \|x_0\|_\infty + \|KMx_n\| + \left\| KL \int_0^t S(t, s) f_n(s) ds \right\| + \left\| \int_0^t S(t, s) f_n(s) ds \right\| \\ &\leq \|x_0\|_\infty + \|K\| \cdot \|Mx_n\| + \|KL\| \cdot N \int_0^t |F(s, x_n(s))| ds + N \int_0^t |F(s, x_n(s))| ds, \end{aligned}$$

where  $\|S(t, s)\| \leq N$ . So we have:

$$\begin{aligned} \frac{R(x_n)}{n} &\leq \frac{\|x_0\|_\infty}{n} + \|K\| \frac{\|Mx_n\|}{n} + N(\|KL\| + 1) \int_0^t \frac{|F(s, x_n(s))|}{n} ds \\ &\leq \frac{\|x_0\|_\infty}{n} + \|K\| \frac{\|Mx_n\|}{\|x_n\|} + N(\|KL\| + 1) \int_0^t \frac{|F(s, x_n(s))|}{n} ds. \end{aligned}$$

Using hypotheses  $H(F)$  (3) and  $H(M)$ , as well as the above inequality, we get that  $1 \leq \lim_{n \rightarrow \infty} R(x_n)/n = 0$ , a contradiction. So indeed  $R: B_r \rightarrow P_{fc}(B_r)$ , where  $B_r = \{x \in C(T, X): \|x\|_\infty \leq r\}$ .

Now we claim that  $R(B_r)$  is compact in  $C(T, X)$ . First note that for every  $t \in T$ , we have

$$R(B_r)(t) \subseteq x_0(t) + KM(B_r) - KL \int_0^t S(t, s) P(s) ds + \int_0^t S(t, s) P(s) ds,$$

where  $P(s) = \{y \in X: \|y\| \leq \sup\{|F(s, x)|: \|x\| \leq r\} = u_r(s)\}$ . But recall that  $M(\cdot)$  is completely continuous. So  $\overline{M(B_r)}$  is compact  $\Rightarrow \overline{KM(B_r)}$  is compact. Also since by hypothesis  $H(A)$ ,  $S(t, s)$  is compact for  $t - s > 0$ , we have that  $S(t, s)P(s) \in P_{kc}(X)$  and clearly  $s \rightarrow S(t, s)P(s)$  is measurable and integrably bounded. Hence using the Rådström embedding theorem (see Hiai–Umegaki [5], Theorem 4.5), we have that  $\int_0^t S(t, s)P(s)ds \in P_{kc}(X)$  (note that in the above mentioned result of Hiai–Umegaki [5], the RNP-hypothesis on  $X$  is superfluous, since by the corollary to Proposition 3.1 of [9],  $\int_0^t S(t, s)P(s)ds$  is closed). So for all  $t \in T$ ,  $\overline{R(B_r)(t)} \in P_k(X)$ . Now, let  $t', t \in T$ ,  $t < t'$ . For  $y \in R(B_r)$ , we have:

$$\begin{aligned} \|y(t') - y(t)\| &\leq \|S(t', 0)x_0 - S(t, 0)x_0\| \\ &+ \|KL\| \left\| \int_0^{t'} S(t', s) f(s) ds - \int_0^t S(t, s) f(s) ds \right\| + \left\| \int_0^{t'} S(t', s) f(s) ds - \int_0^t S(t, s) f(s) ds \right\|. \end{aligned}$$

Since  $S(t, s)$  is a strongly continuous evolution operator given  $\varepsilon > 0$ , there exists  $\delta_1(\varepsilon) > 0$  s.t. if  $\|t' - t\| < \delta_1$

$$\|S(t', 0)x_0 - S(t, 0)x_0\| < \varepsilon/3.$$

Also note that:

$$\begin{aligned} \left\| \int_0^{t'} S(t', s) f(s) ds - \int_0^t S(t, s) f(s) ds \right\| &\leq \left\| \int_0^{t-\delta_2} (S(t', s) f(s) - S(t, s) f(s)) ds \right\| \\ &+ \left\| \int_{t-\delta_2}^{t'} (S(t', s) - S(t, s)) f(s) ds + \int_t^{t'} S(t', s) f(s) ds \right\| \end{aligned}$$

Because of  $H(A)$ , from Proposition 2.1 of [11], we have that  $t \rightarrow S(t, s)$  is continuous in the uniform operator topology, uniformly in  $s$ , for  $t - s$  bounded away from 0. So by choosing  $\delta_2(\varepsilon) > 0$  appropriately small, we will have:

$$\begin{aligned} & \left\| \int_0^{t-\delta_2} (S(t', s) - S(t, s)) f(s) ds \right\| + \left\| \int_{t-\delta_2}^t (S(t', s) - S(t, s)) f(s) ds \right\| \\ & \qquad \qquad \qquad + \left\| \int_t^{t'} S(t', s) f(s) ds \right\| \\ & \leq \int_0^{t-\delta_2} \|S(t', s) - S(t, s)\| u_r(s) ds + 2N \int_{t-\delta_2}^t u_r(s) ds + N \int_t^{t'} u_r(s) ds < \varepsilon/3\alpha. \end{aligned}$$

where  $\alpha = \max(1, \|KL\|)$ . Thus for  $\delta = \min(\delta_1, \delta_2)$ , we have for  $|t - t'| < \delta$   
 $\|y(t') - y(t)\| < \varepsilon$  for all  $y(\cdot) \in R(B_r) \Rightarrow R(B_r)$  is equicontinuous.

Invoking the Arzela–Ascoli theorem, we deduce that  $\overline{R(B_r)}$  is compact in  $C(T, X)$ . Since  $R(\cdot)$  has closed graph and compact range when restricted to  $B_r$ , from Theorem 7.1.16 of Klein–Thompson [7]  $R(\cdot)$  is u.s.c. and so we can apply the Kakutani–Ky Fan fixed point theorem to get  $x \in B_r$  s.t.  $x \in R(x)$ . As we already indicated,  $x(\cdot)$  is a mild solution of (\*).  $\square$

**Remark.** If  $L_0$  has a continuous, linear inverse, then  $H(L)$  is satisfied.

**4. Existence result: nonconvex case.** We also have an existence result for the case where the multivalued perturbation  $F(t, x)$  is nonconvex valued. In this case the hypothesis about  $F(\cdot, \cdot)$  takes the following form:

$H(F)'$ .  $F: T \times X \rightarrow P_f(X)$  is a multifunction s.t.

- (1)  $(t, x) \rightarrow F(t, x)$  is measurable,
- (2)  $x \rightarrow F(t, x)$  is l.s.c. from  $X$  into  $X$ ,
- (3)  $\liminf_{n \rightarrow \infty} \frac{1}{n} \int_0^b \sup_{\|x\| \leq n} |F(t, x)| dt = 0$ .

**THEOREM 2.** *If hypotheses  $H(A)$ ,  $H(F)'$ ,  $H(L)$ ,  $H(M)$  hold with  $M$  linear, then (\*) admits a mild solution.*

**Proof.** We have already seen in the proof of Theorem 1, that  $R(\cdot)$  maps the ball  $B_r$  into itself and furthermore  $\hat{W} = \overline{\text{conv} R(B_r)}$  is compact in  $C(T, X)$ . Let  $H: \hat{W} \rightarrow P_f(L^1(X))$  be the multifunction defined by  $H(x) = S_{F(\cdot, x(\cdot))}^1$ . Let  $x_n \rightarrow x$  in  $C(T, X)$ . Then because of hypotheses  $H(F)'$  (1) and (3) we can apply Theorem 4.1 of [10] and get that  $H(x) \subseteq s\text{-}\liminf H(x_n) \Rightarrow H(\cdot)$  is l.s.c. (see Delahaye–Denel [3]). Apply Fryszkowski's selection theorem [4], to get  $h: \hat{W} \rightarrow L^1(X)$  continuous s.t.  $h(x) \in H(x)$  for all  $x \in \hat{W}$ . Then consider the following problem:

$$(*) (y) \quad \dot{x}(t) = A(t)x(t) + h(y)(t), \quad Lx = Mx.$$

Let  $Q: \hat{W} \rightarrow P_{fc}(W)$  be the multifunction defined by  $Q(y) = \{x(\cdot): x \text{ is a mild solution of } (*) (y)\}$ . It has nonempty values by Theorem 1 and since  $M$  is linear the values of  $Q$  are also convex.

Let  $(y_n, x_n) \in \text{Gr}Q$  s.t.  $(y_n, x_n) \rightarrow (y, x)$  in  $C(T, X) \times C(T, X)$ . We have:

$$x_n(t) = S(t, 0)x_n(0) + \int_0^t S(t, s)h(y_n)(s)ds, \quad Lx_n = Mx_n.$$

Passing to the limit as  $n \rightarrow \infty$  and exploiting the continuity of  $h(\cdot)$ , we get

$$\begin{aligned} x(t) &= S(t, 0)x(0) + \int_0^t S(t, s)h(y)(s)ds, \quad Lx = Mx, \\ &\Rightarrow x \in Q(y), \\ &\Rightarrow \text{Gr}Q \text{ is closed in } C(T, X) \times C(T, X) \text{ and } Q. \end{aligned}$$

Since  $\hat{W}$  is compact in  $C(T, X)$ , we conclude that  $Q(\cdot)$  is u.s.c. Apply the Kakutani-Ky Fan fixed point theorem to get  $y \in Q(y)$ . Clearly  $y(\cdot)$  solves  $(*)$ .  $\square$

**5. Application.** We consider the following multivalued boundary value problem.

$$\begin{aligned} \frac{\partial u(t, y)}{\partial t} &\in \sum_{k=1}^n \frac{\partial}{\partial y_k} p(y) \frac{\partial}{\partial y_k} u(t, y) + f(t, y, u(t, y)) \quad \text{on } T \times G, \\ (**) \quad u(t, y) &= 0, \quad (t, y) \in T \times \partial G, \\ u(0, y) - u(b, y) &= \int_0^b \int_G g(t, y, z, u(t, z)) dt dz. \end{aligned}$$

Here  $G \subseteq \mathbf{R}^n$  is an open, bounded domain with smooth boundary  $\partial G$ . Also  $T = [0, b]$ . We assume that  $g: T \times G \times G \times \mathbf{R} \rightarrow \mathbf{R}$  is a function satisfying the Carathéodory conditions, i.e.  $z \rightarrow g(t, y, z, r)$  is measurable and  $r \rightarrow g(t, y, z, r)$  is continuous. Moreover, for each  $k > 0$  there exist measurable functions

$$\beta_k: T \times G \times G \rightarrow \mathbf{R}_+ \quad \text{and} \quad \gamma_k: T \times G \times G \times \mathbf{R} \rightarrow \mathbf{R}_+ \quad \text{s.t.,}$$

$$|g(t, y, z, r)| \leq \beta_k(t, y, z) \quad \text{for } |r| \leq k \quad \text{and} \quad \int_0^b \int_G \beta_k(t, y, z) dt dz \leq M_k,$$

and

$$|g(t, y, z, r) - g(t, y', z, r)| \leq \gamma_k(t, y, y', z) \quad \text{for } |r| \leq k,$$

$$\lim_{y \rightarrow y'} \int_0^b \int_G \gamma_k(t, y, y', z) dt dz = 0 \quad \text{uniformly in } y'.$$

Finally, there exist  $p \geq 1$  and  $\beta < 2$  s.t.,  $|g(t, y, z, r)| \leq p(1 + |x|^\beta)$ . Also assume that  $f: T \times G \times \mathbf{R} \rightarrow P_{fc}(\mathbf{R})$  is a multifunction s.t.

- (a)  $(t, y) \rightarrow f(t, y, r)$  is measurable;  
 (b)  $r \rightarrow f(t, y, r)$  is  $d$ -continuous (i.e., for every  $z \in R$ ,  $r \rightarrow d(z, f(t, y, r))$  is continuous),  
 (c)  $|f(t, y, r)| \leq k(t)(1 + |r|^\alpha)$ ,  $0 < \alpha < 1$ .

Set  $X = L^2(G)$ ,  $D(A) = W_0^2(G)$  and on  $D(A)$  consider the operator

$$Au = \sum_{k=1}^n \frac{\partial}{\partial y_k} p(y) \frac{\partial}{\partial y_k} u(y).$$

So  $A(\cdot)$  is densely defined and it is well known (see for example Martin [8]), that it generates a compact semigroup  $S(t)$ ,  $t \in T$ .

Also let  $F: T \times X \rightarrow P_{fc}(X)$  be defined by  $F(t, u) = S_{f(t, \cdot, u(\cdot))}^2$ . Clearly because of the reflexivity of  $X = L^2(G)$ ,  $F(t, u) \in P_{wkc}(X)$ . Furthermore note that for all  $v \in X$  we have  $d(v, F(t, u)) = \int_F d(v(z), f(t, z, u(z))) dz$ . From the hypotheses on  $f(\cdot, \cdot, \cdot)$ ,  $(t, u) \rightarrow d(v, F(t, u))$  is measurable in  $t$ , continuous in  $u$ , hence jointly measurable and so  $(t, u) \rightarrow F(t, u)$  is measurable. Also from Theorem 4.2 of [10], we have that  $F(t, \cdot)$  is u.s.c. from  $X$  into  $X_w$ . Next let  $M: C(T, X) \rightarrow X$  be defined by  $(Mu)(y) = \int_0^b \int_G g(t, y, z, u(t)(z)) dt dz$ . From Proposition 4.2, p. 175 of Martin [8], we have that  $M(\cdot)$  is completely continuous. Also using the growth condition on  $g$ , we have

$$\lim_{\|u\| \rightarrow 0} \frac{\|Mu\|_{L^2(G)}}{\|u\|_{C(T, X)}} = 0.$$

Let  $L: C(T, X) \rightarrow X$  be defined by  $Lu = u(0, \cdot) - u(b, \cdot)$ . Clearly this is continuous, linear. Furthermore the only solution of  $\dot{u} = Au$ ,  $u(0) = u(b)$ , is  $u \equiv 0$ . Thus if  $\hat{L}x = L(S(\cdot)x)$ ,  $\hat{L}: X \rightarrow X$  is continuous, linear and  $\hat{L}x = (\text{Id} - S(b))x = 0$  has zero as its only solution. So  $\hat{L}^{-1}$  exists (Fredholm alternative)  $\Rightarrow H(L)$  is satisfied. So if we rewrite (\*\*) as the following evolution equation:

$$(**)' \quad \dot{u}(t) \in Au(t) + F(t, u(t)), \quad Lu = Mu,$$

we see that all hypotheses of Theorem 1 are satisfied and so we conclude the existence of a solution belonging in  $C(T, L^2(G))$ .

It is clear, that the general existence results proved here, can give us periodic solutions for problems of evolution inclusions, extending this way the work of Aubin-Cellina [2].

**Acknowledgment.** The author wishes to express his gratitude to the referee for his corrections and remarks.



## References

- [1] G. Anichini, *Nonlinear problems for systems of differential equations*, Nonlinear Anal. T.M.A. 1 (1977), 691–699.
- [2] J.-P. Aubin and A. Cellina, *Differential Inclusions*, Springer, Berlin 1984.
- [3] J.-P. Delahaye and J. Denel, *The continuities of the point to set maps, definitions and equivalences*, Math. Progr. Study 10 (1979), 3–12.
- [4] A. Fryszkowski, *Continuous selections for a class of nonconvex valued multivalued maps*, Studia Math. 76 (1983), 163–174.
- [5] F. Hiai and H. Umegaki, *Integrals, conditional expectations and martingales of multivalued functions*, J. Multiv. Anal. 7 (1977), 149–182.
- [6] A. Kartsatos, *Locally invertible operators and existence problems in differential systems*, Tôhoku Math. J. 28 (1976), 167–176.
- [7] E. Klein and A. Thompson, *Theory of Correspondences*, Wiley, New York 1984.
- [8] R. Martin, *Nonlinear Operators and Differential Equations in Banach Spaces*, Wiley, New York 1976.
- [9] N. S. Papageorgiou, *On the theory of Banach space valued multifunctions, Part 1. Integration and conditional expectation*, J. Multiv. Anal. 17 (1985), 185–207.
- [10] —, *Convergence theorems for Banach space valued integrable multifunctions*, Intern. J. Math. and Math. Sci. 10 (1987), 433–442.
- [11] —, *On multivalued evolution equations and differential inclusions in Banach spaces*, Comm. Math. Univ. Sancti Pauli 36 (1987), 21–39.
- [12] D. Wagner, *Survey of measurable selection theorems*, SIAM J. Control Optim. 15 (1977), 859–903.
- [13] P. Zecca and P. Zezza, *Nonlinear boundary value problems in Banach spaces for multivalued differential equations on a non-compact interval*, Nonlinear Anal. T.M.A. 3 (1979), 347–352.

UNIVERSITY OF CALIFORNIA  
DEPARTMENT OF MATHEMATICS  
DAVIS, CALIFORNIA, U.S.A.

Reçu par la Rédaction le 10.02.1988

---