

BOUNDARY VALUE PROBLEMS OF NONSINGULAR TYPE ON THE SEMI-INFINITE INTERVAL

RAVI P. AGARWAL AND DONAL O'REGAN

(Received April 15, 1998)

Abstract. Existence of a positive solution is established for second order boundary value problems on the semi-infinite interval.

1. Introduction. In this paper we discuss boundary value problems on the semi-infinite interval. In particular we examine

$$(1.1) \quad \begin{cases} y'' + \phi(t)f(t, y, y') = 0, & 0 < t < \infty \\ y(0) = 0, & y \text{ bounded on } [0, \infty), \end{cases}$$

$$(1.2) \quad \begin{cases} y'' + \phi(t)f(t, y, y') = 0, & 0 < t < \infty \\ y(0) = 0, & \lim_{t \rightarrow \infty} y(t) \text{ exists,} \end{cases}$$

and

$$(1.3) \quad \begin{cases} y'' + \phi(t)f(t, y, y') = 0, & 0 < t < \infty \\ y(0) = 0, & \lim_{t \rightarrow \infty} y'(t) = 0, \end{cases}$$

in Section 2; here $f : [0, \infty) \times [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$. By putting physically reasonable assumptions on f , we will show (1.1), (1.2) and (1.3) have solutions $y \in C^1[0, \infty) \cap C^2(0, \infty)$ with $y > 0$ on $(0, \infty)$ even if $y \equiv 0$ is also a solution. Problems of the above type have been discussed by many authors in the literature; we refer the reader to [2–11] and their references. The technique we use to establish existence is based on (i) establishing new results (see [1] also) on the finite interval $[0, n]$ for each $n \in N^+ = \{1, 2, \dots\}$ and (ii) a diagonalization argument. Consequently, the results of this paper are new and they extend and complement previously known results. We remark that the diagonalization argument applied in this paper has been used by many authors; see [2, 4, 6, 7].

To conclude this section we recall the following well-known existence principle [10] for

$$(1.4) \quad \begin{cases} y'' + \phi(t)F(t, y, y') = 0, & 0 < t < n \\ y(0) = a \geq 0 \\ y'(n) = b \geq 0; \text{ here } n \in \{1, 2, \dots\} \text{ is fixed.} \end{cases}$$

THEOREM 1.1. *Suppose*

$$(1.5) \quad \phi \in C(0, n) \text{ with } \phi > 0 \text{ on } (0, n) \text{ and } \phi \in L^1[0, n]$$

and

$$(1.6) \quad F : [0, n] \times \mathbf{R}^2 \rightarrow \mathbf{R} \text{ is continuous}$$

are satisfied. In addition, suppose there is a constant $M > a + bn$, independent of λ , with

$$|y|_1 = \max\{|y|_0, |y'|_0\} \neq M$$

for any solution $y \in C^1[0, n] \cap C^2(0, n)$ to

$$(1.7)_\lambda \quad \begin{cases} y'' + \lambda\phi(t)F(t, y, y') = 0, & 0 < t < n \\ y(0) = a, \quad y'(n) = b \end{cases}$$

for each $\lambda \in (0, 1)$; here $|u|_0 = \sup_{[0, n]} |u(t)|$. Then (1.4) has a solution $y \in C^1[0, n] \cap C^2(0, n)$ with $|y|_1 \leq M$.

2. Semi-infinite problem. In this section we discuss (1.1), (1.2) and (1.3). Throughout this section we will assume the following conditions hold:

$$(2.1) \quad \phi \in C(0, \infty) \text{ with } \phi > 0 \text{ on } (0, \infty),$$

$$(2.2) \quad \begin{cases} f : [0, \infty) \times [0, \infty) \times [0, \infty) \rightarrow [0, \infty) \text{ is continuous with} \\ f(t, u, p) > 0 \text{ for } (t, u, p) \in [0, \infty) \times (0, \infty) \times (0, \infty), \end{cases}$$

$$(2.3) \quad Q_\infty = \int_0^\infty \phi(s)ds < \infty,$$

$$(2.4) \quad R_\infty = \int_0^\infty s\phi(s)ds < \infty,$$

$$(2.5) \quad \begin{cases} f(t, u, p) \leq w(\max\{u, p\}) \text{ on } [0, \infty) \times (0, \infty) \times (0, \infty) \text{ with} \\ w \geq 0 \text{ continuous and nondecreasing on } [0, \infty), \end{cases}$$

$$(2.6) \quad \sup_{c \in (0, \infty)} \frac{c}{w(c) \max\{Q_\infty, R_\infty\}} > 1$$

and

$$(2.7) \quad \begin{cases} \text{for a constant } H > 0 \text{ there exists a function } \psi_H \text{ continuous} \\ \text{on } [0, \infty) \text{ and positive on } (0, \infty), \text{ and a constant } \gamma, 0 \leq \gamma < 1 \\ \text{with } f(t, u, p) \geq \psi_H(t)p^\gamma \text{ on } [0, \infty) \times [0, H]^2. \end{cases}$$

THEOREM 2.1. Suppose (2.1)–(2.7) hold. Then (1.1), (1.2) and (1.3) have solutions $y \in C^1[0, \infty) \cap C^2(0, \infty)$ with $y > 0$ on $(0, \infty)$.

PROOF. First fix $n \in N^+ = \{1, 2, \dots\}$ and consider the family of problems

$$(2.8) \quad \begin{cases} y'' + \phi(t)f(t, y, y') = 0, & 0 < t < n \\ y(0) = y'(n) = 0. \end{cases}$$

Choose $M > 0$ with

$$(2.9) \quad \frac{M}{w(M) \max\{Q_\infty, R_\infty\}} > 1.$$

Next choose $\varepsilon > 0$ with $\varepsilon < M/(n + 1)$ and

$$(2.10) \quad \frac{M}{w(M) \max\{Q_\infty, R_\infty\} + 2\varepsilon} > 1.$$

Let $n_0 \in N^+$ be chosen so that $n/n_0 < \varepsilon$ and let $N_0 = \{n_0, n_0 + 1, \dots\}$.

We first show that

$$(2.11)^m \quad \begin{cases} y'' + \phi(t)f^*(t, y, y') = 0, & 0 < t < n \\ y(0) = y'(n) = 1/m \end{cases}$$

has a solution for each $m \in N_0$; here

$$f^*(t, u, p) = \begin{cases} f(t, u, p), & u \geq 1/m, \quad p \geq 1/m \\ f(t, u, 1/m), & u \geq 1/m, \quad p < 1/m \\ f(t, 1/m, p), & u < 1/m, \quad p \geq 1/m \\ f(t, 1/m, 1/m), & u < 1/m, \quad p < 1/m. \end{cases}$$

To show that (2.11)^m has a solution, we consider the family of problems

$$(2.12)_\lambda^m \quad \begin{cases} y'' + \lambda\phi(t)f^*(t, y, y') = 0, & 0 < t < n \\ y(0) = y'(n) = 1/m, & m \in N_0 \end{cases}$$

for $0 < \lambda < 1$. Let $y \in C^1[0, n] \cap C^2(0, n)$ be any solution of (2.12)^m_λ. Then $y' \geq 1/m$ and $y \geq 1/m$ on $[0, n]$. Also from (2.5) we have

$$-y''(t) \leq \phi(t)w(|y|_1) \text{ for } t \in (0, n);$$

here $|y|_1 = \max\{|y|_0, |y'|_0\}$ and $|u|_0 = \sup_{[0, n]} |u(t)|$. Integrate from t to n to obtain

$$(2.13) \quad y'(t) \leq w(|y|_1) \int_t^n \phi(x)dx + \frac{1}{m} \text{ for } t \in [0, n].$$

In particular

$$(2.14) \quad y'(0) \leq w(|y|_1)Q_\infty + \varepsilon.$$

Also, by using (2.13) and the equality $\int_0^n s\phi(s)ds = \int_0^n \int_t^n \phi(x)dxdt$,

$$y(n) \leq \frac{n}{m} + \frac{1}{m} + w(|y|_1) \int_0^n s\phi(s)ds$$

and so

$$(2.15) \quad y(n) \leq 2\varepsilon + w(|y|_1)R_\infty.$$

Combine (2.14) and (2.15) to obtain

$$(2.16) \quad \frac{|y|_1}{w(|y|_1) \max\{Q_\infty, R_\infty\} + 2\varepsilon} \leq 1.$$

Now (2.10) together with (2.16) implies $|y|_1 \neq M$.

Thus Theorem 1.1 implies that (2.11)^m has a solution $y_{m,n}$ with $|y_{m,n}|_1 \leq M$. In fact

$$(2.17) \quad \frac{1}{m} \leq y_{m,n}(t) \leq M \quad \text{and} \quad \frac{1}{m} \leq y'_{m,n}(t) \leq M \quad \text{for } t \in [0, n]$$

and $y_{m,n}$ satisfies

$$\begin{cases} y'' + \phi(t)f(t, y, y') = 0, & 0 < t < n \\ y(0) = y'(n) = 1/m. \end{cases}$$

Now (2.7) guarantees the existence of a function $\psi_M(t)$ continuous on $[0, \infty)$ and positive on $(0, \infty)$, and a constant γ , $0 \leq \gamma < 1$, with $f(t, y_{m,n}(t), y'_{m,n}(t)) \geq \psi_M(t)[y'_{m,n}(t)]^\gamma$ for $(t, y_{m,n}(t), y'_{m,n}(t)) \in [0, n] \times [0, M]^2$. Of course, we have immediately that

$$(2.18) \quad y'_{m,n}(t) \geq \left((1-\gamma) \int_t^n \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} \quad \text{for } t \in [0, n]$$

and so

$$(2.19) \quad y_{m,n}(t) \geq \int_0^t \left((1-\gamma) \int_x^n \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} dx \quad \text{for } t \in [0, n].$$

It is also immediate that

$$(2.20) \quad \begin{cases} \{y_{m,n}^{(j)}\}_{m \in N_0} \text{ is a bounded, equicontinuous} \\ \text{family on } [0, n] \text{ for each } j = 0, 1. \end{cases}$$

The Arzelà-Ascoli Theorem guarantees the existence of a subsequence N of N_0 and a function $y_n \in C^1[0, n]$ with $y_{m,n}^{(j)}$ converging uniformly on $[0, n]$ to $y_n^{(j)}$ as $m \rightarrow \infty$ through N ; here $j = 0, 1$. Also $y_n(0) = 0 = y'_n(n)$ and

$$(2.21) \quad y_n(t) \geq \int_0^t \left((1-\gamma) \int_x^n \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} dx \quad \text{for } t \in [0, n],$$

in particular, $y_n > 0$ on $(0, n]$. Now $y_{m,n}$, $m \in N$, satisfies

$$(2.22) \quad y_{m,n}(t) = \frac{1}{m} + \frac{1}{m}t + \int_0^t s\phi(s)f(s, y_{m,n}(s), y'_{m,n}(s))ds + t \int_t^n \phi(s)f(s, y_{m,n}(s), y'_{m,n}(s))ds.$$

Fix $t \in [0, n]$ and let $m \rightarrow \infty$ through N in (2.22) to obtain

$$y_n(t) = \int_0^t s\phi(s)f(s, y_n(s), y'_n(s))ds + t \int_t^n \phi(s)f(s, y_n(s), y'_n(s))ds.$$

Consequently, $y \in C^2(0, n]$ with $y''_n + \phi(t)f(t, y_n, y'_n) = 0$ for $0 < t < n$. Also from (2.17) we have

$$(2.23) \quad 0 \leq y_n(t) \leq M \quad \text{and} \quad 0 \leq y'_n(t) \leq M \quad \text{for } t \in [0, n],$$

and the differential equation yields

$$(2.24) \quad 0 \leq -y''_n(t) \leq \phi(t)H_\infty \quad \text{for } t \in (0, n];$$

here $H_\infty = \sup\{f(t, u, p); (t, u, p) \in [0, \infty) \times [0, M]^2\}$. In addition we have

$$(2.25) \quad y'_n(t) \leq H_\infty \int_t^n \phi(x)dx \leq H_\infty \int_t^\infty \phi(x)dx \quad \text{for } t \in [0, n].$$

To show that (1.1), (1.2) and (1.3) have a solution, we will apply a diagonalization argument. Let

$$u_n(t) = \begin{cases} y_n(t), & t \in [0, n] \\ y_n(n), & t \in [n, \infty). \end{cases}$$

Notice that $u_n \in C^1[0, \infty)$ with

$$(2.26) \quad 0 \leq u_n(t) \leq M \quad \text{and} \quad 0 \leq u'_n(t) \leq M \quad \text{for } t \in [0, \infty),$$

and for $t, s \in [0, \infty)$ it is easy to see that

$$(2.27) \quad |u'_n(t) - u'_n(s)| \leq H_\infty \left| \int_s^t \phi(x)dx \right|.$$

In addition

$$(2.28) \quad u'_n(t) \leq H_\infty \int_t^\infty \phi(x)dx \quad \text{for } t \in [0, \infty),$$

and

$$(2.29) \quad u_n(t) \geq \int_0^t \left((1 - \gamma) \int_x^n \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} dx \quad \text{for } t \in [0, n].$$

Also notice for $n \in N^+$ that

$$(2.30) \quad u_n(t) \geq \int_0^t \left((1 - \gamma) \int_x^1 \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} dx \equiv a_1(t) \quad \text{for } t \in [0, 1].$$

The Arzelà-Ascoli Theorem guarantees the existence of a subsequence N_1 of N^+ and a function $z_1 \in C^1[0, 1]$ with $u_n^{(j)}$ converging uniformly on $[0, 1]$ to $z_1^{(j)}$ as $n \rightarrow \infty$ through N_1 ; here $j = 0, 1$. Also from (2.30), $z_1(t) \geq a_1(t)$ for $t \in [0, 1]$ (in particular, $z_1 > 0$ on $(0, 1]$).

Let $N_1^+ = N_1 \setminus \{1\}$. Notice from (2.29) that

$$(2.31) \quad u_n(t) \geq \int_0^t \left((1 - \gamma) \int_x^2 \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} dx \equiv a_2(t) \quad \text{for } t \in [0, 2].$$

The Arzelà-Ascoli Theorem guarantees the existence of a subsequence N_2 of N_1^+ and a function $z_2 \in C^1[0, 2]$ with $u_n^{(j)}$ converging uniformly on $[0, 2]$ to $z_2^{(j)}$ as $n \rightarrow \infty$ through N_2 ; here $j = 0, 1$. Also from (2.31), $z_2(t) \geq a_2(t)$ for $t \in [0, 2]$ (in particular, $z_2 > 0$ on $(0, 2]$). Note that $z_2 = z_1$ on $[0, 1]$, since $N_2 \subseteq N_1^+$. Let $N_2^+ = N_2 \setminus \{2\}$. Proceed inductively to obtain for $k = 1, 2, \dots$ a subsequence N_k of N_{k-1}^+ and a function $z_k \in C^1[0, k]$ with $u_n^{(j)}$ converging uniformly on $[0, k]$ to $z_k^{(j)}$ as $n \rightarrow \infty$ through N_k ; here $j = 0, 1$. Also

$$z_k(t) \geq a_k(t) = \int_0^t \left((1 - \gamma) \int_x^k \psi_M(s)\phi(s)ds \right)^{1/(1-\gamma)} dx \quad \text{for } t \in [0, k],$$

(so in particular, $z_k > 0$ on $(0, k]$). Note that $z_k = z_{k-1}$ on $[0, k - 1]$. Let $N_k^+ = N_k \setminus \{k\}$.

Define a function y as follows. Fix $t \in (0, \infty)$ and let $k \in N^+$ with $t \leq k$. Define $y(t) = z_k(t)$. Note that y is well-defined and $y(t) = z_k(t) > 0$. We can do this for each $t \in (0, \infty)$ and so $y \in C^1[0, \infty)$ with $y > 0$ on $(0, \infty)$. In addition, $0 \leq y(t) \leq M$, $0 \leq y'(t) \leq M$, and $y'(t) \leq H_\infty \int_t^\infty \phi(x) dx$ for $t \in [0, \infty)$.

Fix $x \in [0, \infty)$ and choose $k \geq x, k \in N^+$. Then, for $n \in N_k^+$, we have

$$y_n(x) = y'_n(k)x + \int_0^x s\phi(s)f(s, y_n(s), y'_n(s))ds + x \int_x^k \phi(s)f(s, y_n(s), y'_n(s))ds.$$

Let $n \rightarrow \infty$ through N_k^+ to obtain

$$z_k(x) = z'_k(k)x + \int_0^x s\phi(s)f(s, z_k(s), z'_k(s))ds + x \int_x^k \phi(s)f(s, z_k(s), z'_k(s))ds.$$

Thus

$$y(x) = y'(k)x + \int_0^x s\phi(s)f(s, y(s), y'(s))ds + x \int_x^k \phi(s)f(s, y(s), y'(s))ds.$$

Consequently, $y \in C^2(0, \infty)$ with $y'' + \phi(t)f(t, y, y') = 0$ for $0 < t < \infty$. Thus y is a solution of (1.1) with $y > 0$ on $(0, \infty)$. In addition, y is a solution of (1.2), since $y' \geq 0$ on $[0, \infty)$ and $0 \leq y \leq M$ on $[0, \infty)$. Finally, since $y'(t) \leq H_\infty \int_t^\infty \phi(x) dx$ for $t \in [0, \infty)$, we have that y is a solution of (1.3). □

EXAMPLE 2.1. The boundary value problem

$$(2.32) \quad \begin{cases} y'' + (y')^\beta e^{-t} = 0, & 0 < t < \infty \\ y(0) = 0, & \lim_{t \rightarrow \infty} y'(t) = 0, \end{cases}$$

with $0 \leq \beta < 1$, has a solution $y \in C^1[0, \infty) \cap C^2(0, \infty)$ with $y > 0$ on $(0, \infty)$.

REMARK 2.1. Notice that $y \equiv 0$ is also a solution of (2.32) if $\beta \neq 0$. Of course, one could construct explicitly a solution to (2.32).

We will apply Theorem 2.1 with $\phi(t) = e^{-t}$ and $w(x) = x^\beta$. Clearly (2.1)–(2.5) and (2.7) (with $\psi_H = 1$ and $\gamma = \beta$) hold. Also

$$\sup_{c \in (0, \infty)} \frac{c}{w(c) \max\{Q_\infty, R_\infty\}} = \sup_{c \in (0, \infty)} \frac{c}{c^\beta} = \infty,$$

so (2.6) is satisfied. Theorem 2.1 now guarantees that (2.32) has a solution $y \in C^1[0, \infty) \cap C^2(0, \infty)$ with $y > 0$ on $(0, \infty)$.

EXAMPLE 2.2. Consider the boundary value problem

$$(2.33) \quad \begin{cases} y'' + \mu(y^\alpha + \eta_0)((y')^\beta + \eta_1)e^{-t} = 0, & 0 < t < \infty \\ y(0) = 0, & \lim_{t \rightarrow \infty} y'(t) = 0, \end{cases}$$

with $\alpha \geq 0, 0 \leq \beta < 1, \eta_0 > 0, \eta_1 \geq 0$, and $\mu > 0$. If

$$(2.34) \quad \mu < \sup_{c \in (0, \infty)} \frac{c}{(c^\alpha + \eta_0)(c^\beta + \eta_1)},$$

then (2.33) has a solution $y \in C^1[0, \infty) \cap C^2(0, \infty)$ with $y > 0$ on $(0, \infty)$.

REMARK 2.2. Notice that $y \equiv 0$ is also a solution of (2.33), if $\eta_1 = 0$ and $\beta \neq 0$.

REMARK 2.3. If $\alpha + \beta < 1$, then (2.34) is satisfied for all $\mu > 0$.

We will apply Theorem 2.1 with $\phi(t) = \mu e^{-t}$ and $w(x) = (x^\alpha + \eta_0)(x^\beta + \eta_1)$. Clearly, (2.1)–(2.5) and (2.7) (with $\psi_H = \eta_0$ and $\gamma = \beta$) hold. Also

$$\sup_{c \in (0, \infty)} \frac{c}{w(c) \max\{Q_\infty, R_\infty\}} = \frac{1}{\mu} \sup_{c \in (0, \infty)} \frac{c}{(c^\alpha + \eta_0)(c^\beta + \eta_1)},$$

so (2.34) guarantees that (2.6) is true. Theorem 2.1 now establishes the result.

REFERENCES

- [1] R. P. AGARWAL AND D. O'REGAN, Second and higher order boundary value problems of nonsingular type, to appear in Acad. Roy. Belg. Bull. Cl. Sci.
- [2] J. W. BEBERNES AND L. K. JACKSON, Infinite interval problems for $y'' = f(t, y)$, Duke Math. J. 34 (1967), 39–47.
- [3] M. FURI AND P. PERA, A continuation method on locally convex spaces and applications to ordinary differential equations on noncompact intervals, Ann. Polon. Math. 47 (1987), 331–346.
- [4] A. GRANAS, R. B. GUENTHER, J. W. LEE AND D. O'REGAN, Boundary value problems on infinite intervals and semiconductor devices, J. Math. Anal. Appl. 116 (1986), 335–348.
- [5] W. OKRASINSKI, On a nonlinear ordinary differential equation, Ann. Polon. Math. 49 (1989), 237–245.
- [6] D. O'REGAN, Positive solutions for a class of boundary value problems on infinite intervals, NoDEA Nonlinear Differential Equations Appl. 1 (1994), 203–228.
- [7] D. O'REGAN, Nonnegative solutions to superlinear problems of generalized Gelfand type, J. Appl. Math. Stochastic Anal. 8 (1995), 275–290.
- [8] D. O'REGAN, Some fixed point theorems for concentrative mappings between locally convex linear topological spaces, Nonlinear Anal. 27 (1996), 1437–1446.
- [9] D. O'REGAN, Continuation fixed point theorems for locally convex linear topological spaces, Math. Comput. Modelling 24 (1996), 57–70.
- [10] D. O'REGAN, Existence theory for nonlinear ordinary differential equations, Mathematics and its Applications, 398, Kluwer Academic Publishers, Dordrecht, 1997.
- [11] K. SCHMITT AND R. THOMPSON, Boundary value problems for infinite systems of second-order differential equations, J. Differential Equations 18 (1975), 277–295.

RAVI P. AGARWAL
DEPARTMENT OF MATHEMATICS
NATIONAL UNIVERSITY OF SINGAPORE
10 KENT RIDGE CRESCENT
SINGAPORE 119260

DONAL O'REGAN
DEPARTMENT OF MATHEMATICS
NATIONAL UNIVERSITY OF IRELAND
GALWAY
IRELAND

