

Bounded, periodic and almost periodic classical solutions of some nonlinear wave equations with a dissipative term

By Mitsuhiro NAKAO

(Received Feb. 3, 1976)

(Revised June 20, 1977)

Introduction.

This paper deals with the questions of the existence and asymptotics of the bounded, periodic and almost periodic classical solutions for the equations

$$(E) \quad \frac{\partial^2}{\partial t^2} u + \sum_{|\alpha|, |\beta| \leq m} (-1)^{|\alpha|} D^\alpha (a_{\alpha, \beta}(x) D^\beta u(x, t)) + \nu \frac{\partial}{\partial t} u \\ + F(x, t, u) = 0 \quad \text{on } \Omega \times R \text{ (or } \Omega + R^+)$$

together with the boundary condition

$$(B) \quad D^\alpha u|_{\partial\Omega} = 0 \quad \text{for } |\alpha| \leq m-1,$$

where ν is a positive constant, Ω is a bounded domain in n -dimensional Euclidean space R^n and $\partial\Omega$ its boundary. We use the following notations

$$\alpha = (\alpha_1, \dots, \alpha_n), \quad |\alpha| = \sum_{i=1}^n |\alpha_i|, \quad D^\alpha = \prod_{i=1}^n \left(\frac{\partial}{\partial x_i} \right)^{\alpha_i}, \\ x = (x_1, \dots, x_n) \text{ etc.}$$

The functions to be considered are all real valued and throughout the paper we make the following assumptions:

$$H_1. \quad A = \sum_{|\alpha|, |\beta| \leq m} (-1)^{|\alpha|} D^\alpha (a_{\alpha, \beta}(x) D^\beta u)$$

is formally selfadjoint and coercive on \mathring{H}_m , i.e., there exist some positive constants c_0, c_1 such that

$$c_1 \|u\|_{\mathring{H}_m}^2 \geq \langle Au, u \rangle \geq c_0 \|u\|_{\mathring{H}_m}^2 \quad \text{for } u \in \mathring{H}_m$$

where we put

$$\langle Au, v \rangle = \sum_{|\alpha|, |\beta| \leq m} \int_{\Omega} a_{\alpha\beta}(x) D^\beta u(x) D^\alpha v(x) dx, \quad u, v \in \dot{H}_m$$

and

$$\|u\|_{\dot{H}_m} = \left(\sum_{|\alpha| \leq m} \int_{\Omega} |D^\alpha u|^2 dx \right)^{1/2}.$$

H₂. $F(x, t, u)$ is of the form

$$F(x, t, u) = g(x, u) + f(x, t)$$

and g, f satisfy the conditions:

$$g \in C^{\max\{[n/2]+1, 3\}}(\bar{\Omega} \times R)$$

and $\left| \frac{\partial^j}{\partial u^j} g(x, u) \right| \leq K_0 \sum_{i=1,2} |u|^{\max\{r_i+1-j, 0\}} \quad (j=0, 1, 2, 3)$

where $r_i (i=1, 2)$ are constants such that

$$0 < r_1 \leq r_2 < \infty \quad \text{if } 1 \leq n \leq 2m$$

and $0 < r_1 \leq r_2 \leq \frac{2m}{n-2m} \quad \text{if } n > 2m,$

and $f(\cdot, t) \in C^3(R; L^2) \cap C^0(R; H_{2m} \cap \dot{H}_m)$

where $C^i(I; V)$ denotes the class of functions from $I \subset R$ to V with uniformly continuous derivatives of order $\leq i$. Moreover we assume

$$M \equiv \sup_{\substack{t \in \bar{R} \\ i=0,1,2,3}} \max(\|D_t^i f(t)\|_{L^2}, \|Af(\cdot, t)\|_{L^2}, \|AD_t f(\cdot, t)\|_{L^2}) < +\infty.$$

H₃. $2m \geq \left[\frac{n}{2} \right] + 1, \quad m \geq \left[\frac{n}{3} \right], \quad a_{\alpha\beta} \in C^{2m+[n/2]+1+|\alpha|}(\bar{\Omega})$

and $\partial\Omega \in C^{2m+[n/2]+1}.$

Recently, Clements [6] and Biroli [5] have proved the existence of generalized periodic solution to the problem (E)-(B) in the case $F(x, t, u)$ is monotonically increasing in u . On the other hand the author [14] has treated a nonmonotonic case, proving the existence of bounded, periodic and almost periodic solutions for the problem (E)-(B) with $m=1$ and with $\nu \frac{\partial}{\partial t} u$ replaced by nonlinear term $\rho\left(x, \frac{\partial}{\partial t} u\right)$. But the solutions obtained are also generalized ones.

The object of this paper is to give existence theorems concerning bounded, periodic and almost periodic classical solutions to the problem (E)-(B) and to investigate some asymptotic properties of them.

In this paper we employ a Galerkin's method. In particular techniques are related to those in Sather [20], Ebihara-Nakao-Nanbu [7], Amerio-Prouse [3] and Nakao [14] [15] [16]. For other treatments of classical periodic solutions of the nonlinear wave equations with a dissipative term, see Rabinowitz [19] and Wahl [24], and for quasi-periodic solution see Yamaguchi [25].

The existence of a dissipative term is essential for our arguments. Indeed, the problem of the existence and regularity of bounded, periodic and almost periodic solutions for the nonlinear wave equations without dissipative term is extremely difficult and there have been researches only for periodic solutions of essentially one dimensional equations. For this special case see Vejvoda [23], Rabinowitz [18], Torelli [22], Hall [8], Nakao [11] [12] and the references cited there.

Similar problem for nonlinear parabolic equations has been treated by Biroli [4], Nakao [11] and Nakao-Nanbu [17] where differential inequalities concerning several norms of approximate solutions are used extensively, while here we use 'integral inequality' for the estimations of approximate solutions.

§ 1. Preliminaries.

We employ the usual notations for function spaces and for norms associated with them (see, e. g., Lions [9]). For $u \in \dot{H}_m$ we put $\|u\|_A \equiv \langle Au, u \rangle^{1/2}$ and define several functionals on \dot{H}_m as follows:

$$J_0(u) = \frac{1}{2} \|u\|_A^2 + \int_{\Omega} \int_0^{u(x)} g(x, s) ds dx,$$

$$J_1(u) = \|u\|_A^2 + \int_{\Omega} g(x, u(x)) u(x) dx,$$

$$\check{J}_0(u) = \frac{1}{2} \|u\|_A^2 - K_0 \sum_{i=1,2} S_{r_i+2}^{r_i+2} / (r_i+2) \cdot \|u\|_A^{r_i+2}$$

and

$$J_1(u) = \|u\|_A^2 - K_0 \sum_{i=1,2} S_{r_i+2}^{r_i+2} \|u\|_A^{r_i+2},$$

where S_q is the sobolev constant such that

$$\|u\|_{L^q} \leq S_q \|u\|_A \quad \text{for } u \in \dot{H}_m$$

with $0 < q \leq \frac{2n}{n-2m}$ if $n > 2m$ and $0 < q < \infty$ if $1 \leq n \leq 2m$.

Notice that $J_0(u) \geq \check{J}_0(u)$ and $J_1(u) \geq \check{J}_1(u)$ for $u \in \dot{H}_m$. For convenience we put, for $x \geq 0$,

$$J_0(x) = \frac{1}{2}x^2 - K_0 \sum_{i=1,2} S_{r_i+2}^{r_i+2} / (r_i+2) \cdot x^{r_i+2}$$

and

$$J_1(x) = x_2 - K_0 \sum_{i=1,2} S_{r_i+2}^{r_i+2} \cdot x^{r_i+2}.$$

Let D_1 , D_0 and x_0 be positive numbers such that

$$D_1 = \max_{x \geq 0} \tilde{J}_1(x) = \tilde{J}_1(x_0) \quad \text{and} \quad D_0 = \tilde{J}_0(x_0)$$

Then it is easy to see that

$$(1.1) \quad D_0 = \max_{0 \leq x \leq x_0} \tilde{J}_0(x) < \min(D_1, \max_{x \geq 0} \tilde{J}_0(x)).$$

Now, let $\{\phi_j\}_{j=1}^{\infty}$ be the set of eigen functions of the operator A in L^2 with the Dirichlet boundary condition:

$$A\phi_j = \mu_j \phi_j \quad \text{in } \Omega \quad \text{and} \quad D^\alpha \phi_j = 0 \quad \text{on } \partial\Omega \quad \text{for } |\alpha| \leq m-1.$$

As is well known, $\phi_j \in C^{4m+[\frac{n}{2}]+1}(\bar{\Omega})$ and $\{\phi_j\}$ is dense both in L_2 and \dot{H}_m (cf. Agmon [1]).

§ 2. Approximate solutions.

We employ the Galerkin's method for our purpose. Let $\{\phi_j\}$ be the basis of \dot{H}_m consisting of the eigen functions of A , and consider the system of ordinary differential equations

$$(2.1) \quad (u_r''(t), \phi_j) + (Au_r(t), \phi_j) + \nu(u_r'(t), \phi_j) + (F(\cdot, t, u_r), \phi_j) = 0, \\ j=1, 2, \dots, r,$$

with the initial condition

$$(2.2) \quad u_r'(0) = u_r(0) = 0,$$

where $u_r(t) = \sum_{j=1}^r \lambda_j^r(t) \phi_j$.

The equation (2.1) is equivalent to

$$(2.1)' \quad u_r''(t) + Au_r(t) + \nu u_r'(t) + P_r F(x, t, u_r) = 0,$$

where P_r is the projection of the space L_2 onto the r -dimensional subspace spanned by $\{\phi_j\}_{j=1}^r$.

Since $F(x, t, u)$ is locally Lipschitz continuous in u , $u_r(t)$ exists uniquely on some interval, say $[0, T_r)$. In what follows, we shall estimate $u_r(t)$ in various norms. First we show

LEMMA 2.1. *There exists a positive constant M_0 such that if $M < M_0$, then $u_r(t)$ exist on $[0, \infty)$ and the estimates*

$$(2.3) \quad \|u_r(t)\|_A \leq x_0(M) < x_0 \quad \text{and} \quad \|u'_r(t)\|_{L_2} \leq \sqrt{2k_0(M)} < \sqrt{2D_0}$$

hold for all r , where $x_0(M)$ and $k_0(M)$ are certain constants tending to 0 as $M \rightarrow 0$.

PROOF. Let $k_0(M)$ be a positive constant ($< D_0$) determined later, and $x_0(M)$ ($< x_0$) be the solution of the numerical equation in x :

$$(2.4) \quad \check{f}_0(x) = k_0(M), \quad x \geq 0.$$

In order to prove the lemma it suffices to show

$$(2.5) \quad \frac{1}{2} \|u'_r(t)\|_{L_2}^2 + J_0(u_r(t)) \leq k_0(M) \quad \text{for} \quad \forall t \in [0, T_r).$$

The proof of (2.5) can be carried out essentially in the same manner as the proof of Lemma 2.1 in [14], and we give it briefly. Suppose that the inequality (2.5) were false. There would then exist a time $\bar{t} \in [0, T_r)$ such that

$$(2.6) \quad \frac{1}{2} \|u'_r(t)\|_{L_2}^2 + J_0(u_r(t)) \leq k_0(M) \quad \text{for} \quad t \leq \bar{t}$$

and

$$(2.7) \quad \frac{1}{2} \|u'_r(t_s)\|_{L_2}^2 + J_0(u_r(t_s)) > k_0(M) \quad \text{for some} \quad t_s > \bar{t},$$

where t_s can be chosen as close to \bar{t} as wanted. For any $t_1, t_2 \in [0, T_r)$, we obtain by (2.1)

$$(2.8) \quad \begin{aligned} & \frac{1}{2} \|u'_r(t_2)\|_{L_2}^2 + J_0(u_r(t_2)) + \nu \int_{t_1}^{t_2} \|u'_r(t)\|_{L_2}^2 dt \\ &= \frac{1}{2} \|u'_r(t_1)\|_{L_2}^2 + J_0(u_r(t_1)) - \int_{t_1}^{t_2} (f(t), u'_r(t)) dt. \end{aligned}$$

Taking $t_2 = \bar{t}$ and $t_1 = 0$ in the above we have

$$(2.9) \quad \begin{aligned} & \frac{1}{2} \|u'_r(\bar{t})\|_{L_2}^2 + J_0(u_r(\bar{t})) + \nu \int_0^{\bar{t}} \|u'_r(t)\|_{L_2}^2 dt \\ & \leq \nu \int_0^{\bar{t}} \|u'_r(t)\|_{L_2}^2 dt + \frac{1}{4\nu} \int_0^{\bar{t}} \|f(t)\|_{L_2}^2 dt, \end{aligned}$$

and hence

$$4\nu k_0(M) \leq \int_0^{\bar{t}} \|f(t)\|_{L_2}^2 dt,$$

which implies $\bar{t} > 1$ if we choose $k_0(M)$ satisfying

$$(2.10) \quad k_0(M) > \frac{1}{4\nu} M^2.$$

Under the assumption (2.10), we take $t_2 = t_\varepsilon$ and $t_1 = t_\varepsilon - 1$ ($< \bar{t}$) in (2.8) to obtain

$$(2.11) \quad \int_{t_\varepsilon-1}^{t_\varepsilon} \|u_r'(t)\|_{L_2}^2 dt \leq (\nu^{-1} M)^2,$$

and hence there exist points $\bar{t}_1 \in [t_\varepsilon - 1, t_\varepsilon - \frac{3}{4}]$ and $\bar{t}_2 \in [t_\varepsilon - \frac{1}{4}, t_\varepsilon]$ such that

$$(2.12) \quad \|u_r'(\bar{t}_i)\|_{L_2} \leq 2\nu^{-1} M \quad (i=1, 2).$$

Therefore, multiplication (2.1) by $\lambda_j^r(t)$, summation over j and integration by parts yield

$$(2.13) \quad \begin{aligned} \int_{\bar{t}_1}^{\bar{t}_2} J_1(u_r(t)) dt &\leq |(u_r'(\bar{t}_1), u_r(\bar{t}_1))| + |(u_r'(\bar{t}_2), u_r(\bar{t}_2))| \\ &\quad + \int_{\bar{t}_1}^{\bar{t}_2} \{ \|u_r'(t)\|_{L_2}^2 + \nu |(u_r', u_r)| + |(f(t), u_r(t))| \} dt \\ &\leq (\nu^{-1} M)^2 + 4\nu^{-1} c_0^{-1} M x_0 + 2M c_0^{-1} \left(\int_{\bar{t}_1}^{\bar{t}_2} \|u_r(s)\|_{\mathcal{A}}^2 ds \right)^{1/2}. \end{aligned}$$

On the other hand we know $\|u_r(t)\|_{\mathcal{A}} \leq x_0$ if $t \leq t_\varepsilon$, and hence

$$(2.14) \quad \int_{\bar{t}_1}^{\bar{t}_2} J_1(u_r(t)) dt \geq \int_{\bar{t}_1}^{\bar{t}_2} \tilde{J}_1(u_r(t)) dt \geq c_2 \int_{\bar{t}_1}^{\bar{t}_2} \|u_r(t)\|_{\mathcal{A}}^2 dt,$$

where $c_2 = 1 - k_0 \sum_{i=1,2} S_{\tau_i+2}^{\tau_i+2} x_0^{\tau_i} > 0$.

From (2.13) and (2.14) we obtain

$$(2.15) \quad \int_{\bar{t}_1}^{\bar{t}_2} \|u_r(t)\|_{\mathcal{A}}^2 dt \leq c_2^{-1} \{ (2(\nu^{-1} M)^2 + 8\nu^{-1} c_0^{-1} M x_0 + 4M^2 c_0^{-2} c_2^{-1}) \}.$$

By (2.11) and (2.15) we see that there exists a point $t^* \in [\bar{t}_1, \bar{t}_2]$ such that

$$(2.16) \quad \frac{1}{2} (\|u_r'(t^*)\|_{L_2}^2 + \|u_r(t^*)\|_{\mathcal{A}}^2) \leq k_0'(M),$$

where

$$k'_0(M) \equiv (1 + 2c_2^{-1})(\nu^{-1}M)^2 + 4c_2^{-1}M(2\nu^{-1}c_0^{-1}x_0 + Mc_0^{-2}c_0^{-1}).$$

Therefore, by (2.8) with $t_2 = t_\varepsilon$ and $t_1 = t^*$, we obtain

$$(2.17) \quad \frac{1}{2} \|u'_r(t_\varepsilon)\|_{L_2}^2 + J_0(u_r(t_\varepsilon)) \leq \frac{1}{2} \|u'_r(t^*)\|_{L_2}^2 + J_0(u_r(t^*)) + \frac{1}{4\nu} M^2 \leq k_0(M),$$

where we define

$$k_0(M) = k'_0(M) + \frac{1}{4\nu} M^2 + k_0 \sum_{i=1,2} \frac{1}{r_i + 2} S_{r_i+2}^{r_i+2} (2k'_0(M))^{(r_i+2)/2}.$$

Then, (2.10) is automatically valid, and if we choose M_0 so that $k_0(M_0) = D_0$, the inequality (2.7) contradicts (2.17) for $M < M_0$. Q. E. D.

REMARK. As is easily seen from the proof, Lemma 1.1 is valid in fact even if we replace M by $\sup_t \left(\int_t^{t+1} \|f(s)\|_{L_2}^2 ds \right)^{1/2}$.

We proceed to further estimation of the approximate solutions. To do so it is convenient to prepare an estimate of the approximate solutions $\{U_r(t)\}$ of the linear equation (E) with $F(x, t, u) = F(x, t)$.

LEMMA 2.2. *Let $\{U_r(t)\}$ be the solution of the equation (2.1)-(2.2) with $F(x, t, u_r) = F(x, t)$. Then we have*

$$(2.18) \quad \max_{s \in [t, t+1]} \|U_r(s)\|_{E}^2 \leq c_3 (\|U_r(t)\|_{E}^2 - \|U_r(t+1)\|_{E}^2) + c_4 N(t)^2$$

for $t \in [0, \infty)$,

where

$$N(t) = \left(\int_t^{t+1} \|F(\cdot, s)\|_{L_2}^2 ds \right)^{1/2}, \quad \|U\|_{E}^2 = \|U'(t)\|_{L_2}^2 + \|U(t)\|_{A}^2,$$

$$c_3 = 2\nu^{-1} \{2\nu + 5 + 4(4 + \nu)^2 c_0^{-2}\} \quad \text{and} \quad c_4 = c_3 \nu^{-1} + 2 + 8c_0^{-2}.$$

PROOF. By (2.1) we have, for $0 \leq t_1 < t_2$,

$$(2.19) \quad \nu \int_{t_1}^{t_2} \|U'_r(s)\|_{L_2}^2 ds = \frac{1}{2} (\|U_r(t_1)\|_{E}^2 - \|U_r(t_2)\|_{E}^2) - \int_{t_1}^{t_2} (F(\cdot, s), U'_r(s)) ds,$$

and hence, taking $t_2 = t + 1$, $t_1 = t$,

$$(2.20) \quad \int_t^{t+1} \|U'_r(s)\|_{L_2}^2 ds \leq \nu^{-1} (\|U_r(t)\|_{E}^2 - \|U_r(t+1)\|_{E}^2) + (\nu^{-1} N)^2 \equiv D(t)^2.$$

It follows that there exist two points $t_1 \in [t, t + \frac{1}{4}]$ and $t_2 \in [t + \frac{3}{4}, t + 1]$ such that

$$\|U'_r(t_i)\|_{L_2} \leq 2D(t) \quad (i=1, 2),$$

and hence by (2.1) we have, as in (2.13),

$$(2.21) \quad \int_{t_1}^{t_2} \|U_r(s)\|_A^2 ds \leq D(t)^2 + ((4+\nu)D(t) + N) c_0^{-1} \max_{s \in [t, t+1]} \|U_r(s)\|_A.$$

From (2.20) and (2.21) we see that there exists a point $t^* \in [t_1, t_2]$ such that

$$(2.22) \quad \|U_r(t^*)\|_E^2 \leq 4D(t)^2 + 2((4+\nu)D(t) + N) c_0^{-1} \max_{s \in [t, t+1]} \|U_r(s)\|_A.$$

Using the estimate (2.22) we have from (2.19) with $t_1 = t^*$ (or $t_2 = t^*$)

$$\begin{aligned} \max_{s \in [t, t+1]} \|U_r(s)\|_E^2 &\leq 2\nu \int_t^{t+1} \|U'_r(s)\|_{L_2}^2 ds \\ &\quad + \|U_r(t^*)\|_E^2 + 2 \int_t^{t+1} \|F(s)\|_{L_2} \|U'_r(s)\|_{L_2} ds \\ &\leq 2\nu D(t)^2 + 4D(t)^2 + 2((4+\nu)D(t) + N) c_0^{-1} \\ &\quad \times \max_{s \in [t, t+1]} \|U_r(s)\|_E + 2ND(t) \\ &\leq \{5 + 2\nu + 4(4+\nu)^2 c_0^{-2}\} D(t)^2 + (1 + 4c_0^{-2}) N^2 \\ &\quad + \frac{1}{2} \max_{s \in [t, t+1]} \|U_r(s)\|_E^2, \end{aligned}$$

which proves the lemma.

Q. E. D.

We obtain further the following.

LEMMA 2.3. *Let $U_r(t)$ be as in Lemma 2.2. Then*

$$(2.23) \quad \|U_r(t)\|_E^2 \leq \max \left(\max_{s \in [0, 1]} \|U_r(s)\|_E^2, c_4 \bar{N}^2 \right) \\ \leq \|U_r(0)\|_E^2 + c_4 \bar{N}^2,$$

where $\bar{N} = \sup_{t \in \mathbb{R}} N(t)$.

PROOF. The former inequality in (2.23) is an immediate consequence of (2.18). For $t \in [0, 1]$ we have

$$\begin{aligned} \|U_r(t)\|_E^2 &\leq \|U_r(0)\|_E^2 + \frac{1}{2\nu} \int_0^t \|F(s)\|_{L_2}^2 ds \\ &\leq \|U_r(0)\|_E^2 + \frac{1}{2\nu} \bar{N}^2, \end{aligned}$$

which gives (2.23), because $c_4 > \frac{1}{2\nu}$.

Q. E. D.

Now we return to the estimation of $\{u_r(t)\}$.

LEMMA 2.4. Let M'_1 be the positive number such that

$$c_5(M'_1) \equiv 1 - 4c_4 \sum_{i=1,2} K_0^2 S_{2r_i+2}^{2r_i+2} x_0(M'_1)^{2r_i} = 0.$$

Then for $M < M_1 \equiv \min(M_0, M'_1)$ we have

$$(2.24) \quad \|D_t u_r(t)\|_E \leq k_1(M) \quad \text{for } t \in [0, \infty),$$

where $k_1(M) = (1 + 2c_4)^{1/2} M c_5(M)^{-1/2}$.

PROOF. By Lemma 2.3 with $U_r(t) = D_t u_r(t)$ and $F(x, t) = D_t F(x, t, u_r)$, we have

$$(2.25) \quad \max_{t \in \mathbb{R}^+} \|D_t u_r(t)\|_E^2 \leq \|D_t u_r(0)\|_E^2 + c_4 \max_{t \in \mathbb{R}^+} \|D_t \{g(\cdot, u_r(\cdot, t)) + f(t)\}\|_{L_2}^2 \cdot \|D_t \{g(\cdot, u_r(\cdot, t)) + f(t)\}\|_{L_2}$$

$$\begin{aligned} &\leq M + K_0 \left(\int_{\Omega} \left(\sum_{i=1,2} |u_r|^{r_i} |D_t u_r(x, t)| \right)^2 dx \right)^{1/2} \\ (2.26) \quad &\leq M + \sqrt{2} K_0 \left(\sum_{i=1,2} S_{2r_i+2}^{2r_i+2} x_0(M)^{2r_i} \right)^{1/2} \|D_t u_r(t)\|_E, \end{aligned}$$

and also

$$\begin{aligned} \|D_t u_r(0)\|_E^2 &= \|D_t^2 u_r(0)\|_{L_2}^2 + \|D_t u_r(0)\|_A^2 \\ &= \|-\nu u_r'(0) - A u_r(0) + P_r(g(\cdot, u_r(0)) + f(0))\|_{L_2}^2 \\ (2.27) \quad &\leq M^2. \end{aligned}$$

The inequalities (2.25) - (2.27) yield (2.24).

Q. E. D.

LEMMA 2.5. Let M_1 be the number given in Lemma 2.4. Then if $M < M_1$, we have

$$(2.28) \quad \|D_t^2 u_r(t)\|_E \leq k_2(M) \quad \text{for } t \in [0, \infty),$$

where

$$k_2(M) = \left\{ (2\nu^2 + 3)M^2 + 4c_4M^2 + 8c_4K_0^2 \sum_{i=1,2} S_{4+2\bar{r}_i}^{4+2\bar{r}_i} x_0(M)^{2\bar{r}_i} k_1(M)^4 \right\}^{1/2} \\ \times c_5(M)^{-1/2} \quad (\bar{r}_i = \max(r_i - 1, 0)).$$

PROOF.

$$\|D_i^2 F(\cdot, t, u_r(t))\|_{L_2}^2 \\ = \int_{\mathcal{Q}} \left| D_i^2 f(x, t) + \frac{\partial}{\partial u} g(x, u_r(t)) D_i^2 u_r + \left(\frac{\partial}{\partial u} \right)^2 g(x, u_r(t)) \right. \\ \left. \times (D_i u_r(x, t))^2 \right|^2 dx \\ \leq 4M^2 + 4K_0^2 \sum_{i=1,2} S_{2r_i+2}^{2r_i+2} \|u_r(t)\|_A^{2r_i} \|D_i^2 u_r(t)\|_A^2 \\ + 8K_0^2 \sum_{i=1,2} S_{4+2\bar{r}_i}^{4+2\bar{r}_i} \|u_r(t)\|_A^{2\bar{r}_i} \|D_i u_r(t)\|_A^4 \\ \leq 4M^2 + 4K_0^2 \sum_{i=1,2} S_{2r_i+2}^{2r_i+2} x_0(M)^{2r_i} \|D_i^2 u_r(t)\|_E^2 \\ + 8K_0^2 \sum_{i=1,2} S_{4+2\bar{r}_i}^{4+2\bar{r}_i} x_0(M)^{2\bar{r}_i} k_1(M)^4.$$

Also we have

$$\|D_i^2 u_r(\cdot, 0)\|_E^2 = \|-\nu D_i^2 u_r(0) - A D_i u_r(0) + P_r D_i F(\cdot, t, u_r(0))\|_{L_2}^2 \\ + \|-\nu D_i u_r(0) - A u_r(0) + P_r F(\cdot, t, u_r(0))\|_A^2 \\ = \|-\nu(-\nu D_i u_r(0) - A u_r(0) + P_r F(\cdot, t, u_r(0))) \\ + P_r D_i f(\cdot, 0)\|_{L_2}^2 + \|P_r f(\cdot, 0)\|_A^2 \\ \leq 2(\nu^2 + 1)M^2 + M \|A P_r f(\cdot, 0)\|_{L_2} \\ \leq (2\nu^2 + 3)M^2.$$

Here we have used the fact that A and P_r commute on $H_{2m} \cap \mathring{H}_m$. Thus by Lemma 2.3 with $U_r(t) = D_i^2 u_r(t)$ and with $F(x, t) = D_i^2 F(x, t, u_r(t))$, we obtain (2.28). Q. E. D.

LEMMA 2.6. *Let $M < M_1$. Then*

$$(2.29) \quad \|D_i^3 u_r(t)\|_E \leq k_3(M) \quad \text{for } t \in [0, \infty),$$

where

$$k_3(M) = \left[\{(\nu^2 + \nu + 2)^2 + (\nu + 1)^2\} M^2 + 6c_4 \{M^2 + 2K_0^2 \sum_{i=1,2} S_{6+2\tilde{r}_i}^{6+2\tilde{r}_i} x_0(M)^{2\tilde{r}_i} k_1(M)^6 + 6K_0^2 \sum_{i=1,2} S_{4+2\tilde{r}_i}^{4+2\tilde{r}_i} x_0(M)^{2\tilde{r}_i} k_1(M)^2 k_2(M)^2\} \right]^{1/2} c_5(M)^{-1/2}.$$

$$(\tilde{r}_i = \max(r_i - 2, 0).)$$

PROOF. Easy calculations give

$$\begin{aligned} \|D_t^3 F(\cdot, t, u_r(t))\|_{L_2}^2 &\leq 6 \left(M^2 + 2K_0^2 \sum_{i=1,2} S_{6+2\tilde{r}_i}^{6+2\tilde{r}_i} \|u_r(t)\|_A^{2\tilde{r}_i} \|D_t u_r(t)\|_A^6 \right. \\ &\quad + 18K_0^2 \sum_{i=1,2} S_{4+2\tilde{r}_i}^{4+2\tilde{r}_i} \|u_r(t)\|_A^{2\tilde{r}_i} \|D_t u_r(t)\|_A^2 \\ &\quad \left. \times \|D_t^2 u_r(t)\|_A^2 \right) \\ &\quad + 4K_0^2 \sum_{i=1,2} S_{2+2r_i}^{2+2r_i} \|u_r(t)\|_A^{2r_i} \|D_t^3 u_r(t)\|_A^2 \\ &\leq 6 \left(M^2 + 2K_0^2 \sum_{i=1,2} S_{6+2\tilde{r}_i}^{6+2\tilde{r}_i} x_0(M)^{2\tilde{r}_i} k_1(M)^6 \right. \\ &\quad \left. + 18K_0^2 \sum_{i=1,2} S_{4+2\tilde{r}_i}^{4+2\tilde{r}_i} x_0(M)^{2\tilde{r}_i} k_1(M)^2 k_2(M)^2 \right) \\ &\quad + 4K_0^2 \sum_{i=1,2} S_{2+2r_i}^{2+2r_i} x_0(M)^{2r_i} \|D_t^3 u_r(t)\|_A^2. \end{aligned}$$

Also we have

$$\begin{aligned} \|D_t^3 u_r(0)\|_{\mathbb{E}}^2 &= \{D_t(\nu D_t u_r(t) + Au_r(t) + P_r F(\cdot, t, u_r(t)))\}_{t=0}\|_{\mathbb{E}}^2 \\ &= \|\nu \{-\nu D_t u_r(t) - Au_r(t) - P_r F(\cdot, t, u_r(t))\}_{t=0} \\ &\quad + \{AD_t u_r(t) + P_r D_t F(\cdot, t, u_r(t))\}_{t=0}\|_{\mathbb{E}}^2 \\ &= \|\nu^2 D_t^2 u_r(0) + \nu P_r f(\cdot, 0) + A \{-\nu D_t u_r(t) - Au_r(t) \\ &\quad - P_r F(\cdot, t, u_r(t))\}_{t=0} + P_r D_t^2 f(\cdot, 0)\|_{L_2}^2 \\ &\quad + \|\nu P_r f(\cdot, 0) + P_r D_t f(\cdot, 0)\|_A^2 \end{aligned}$$

$$\begin{aligned}
&\leq \|\nu^2\{-\nu D_t u_r(0) - Au_r(0) - P_r F(\cdot, 0, u_r(0))\} \\
&\quad + \nu P_r f(\cdot, 0) - AP_r f(0) + P_r D_t^2 f(\cdot, 0)\|_{L_2}^2 \\
&\quad + \|\nu P_r f(\cdot, 0) + P_r D_t f(\cdot, 0)\|_{L_2} \|A(\nu P_r f(\cdot, 0) + P_r D_t f(\cdot, 0))\|_{L_2} \\
&\leq (\nu^2 + \nu + 2)^2 M^2 + (\nu + 1)^2 M^2.
\end{aligned}$$

(Note that $D_t f \in H_{2m} \cap \dot{H}_m$.)

Thus, by Lemma 2.3, the proof is completed.

Q. E. D.

Now we have finished the estimation of the approximate solutions.

§ 3. Passage to the limit.

In this section we shall study the convergency properties of the approximate solutions $u_r(t)$ as $r \rightarrow \infty$. Always we assume $M < M_1$. Then by the estimates given in the previous section we see first that there exists a subsequence of $\{u_r(t)\}$, which will be denoted by the same notation for simplicity, and a function $u(x, t)$ such that

$$(3.1) \quad D_t^i u_r(t) \longrightarrow D_t^i u(t) \quad \text{weakly* in } L^\infty(R^+; \dot{H}_m), \text{ a. e. on } \Omega \times R^+, \\ \text{and uniformly in } L_2 \text{ on each compact} \\ \text{set of } R^+ \text{ (} i=0, 1, 2, 3),$$

and

$$(3.2) \quad D_t^1 u_r(t) \longrightarrow D_t^1 u(t) \quad \text{weakly* in } L^\infty(R^+; L_2).$$

Moreover it is clear that the limit function $u(t) = u(x, t)$ is in $B^2(R^+; \dot{H}_m) \cap B^3(R^+; L_2)$ and satisfies the estimates (2.3), (2.24), (2.28) and (2.29) for $u_r = u$, where we put, for a Banach space V and an interval $I \subset R$,

$$B^i(I; V) = \{u \in C^i(I; V) \mid \sup_{\substack{t \in I \\ 0 \leq j \leq i}} \|D_t^j u(t)\|_V < +\infty\}.$$

Next, to show the uniform continuity of $D_t^3 u(t)$ in E , we shall prove the uniform equi-continuity of $\{D_t^3 u_r(t)\}$ with respect to the norm $\|\cdot\|_E$. For each $t \in R^+$, $u_r(t)$ is the solution of the elliptic equation

$$\begin{aligned}
Au_r(t) &= -u_r''(t) - \nu u_r'(t) + P_r F(x, t, u_r(t)) \\
&\equiv h_r(x, t) \in B(R^+; L_2)
\end{aligned}$$

and by the theory of elliptic boundary value problem (cf. Agmon-Douglis-Nirenberg [2]) we have for $\delta > 0$

$$\begin{aligned}
 |u_r(x, t + \delta) - u_r(x, t)| &\leq \text{const.} \|u_r(\cdot, t + \delta) - u_r(\cdot, t)\|_{H_{2m}} \\
 &\leq \text{const.} \|h_r(\cdot, t + \delta) - h_r(\cdot, t)\|_{L_2} \\
 (3.3) \qquad \qquad \qquad &\leq \phi_1(\delta) \longrightarrow 0 \quad (\text{as } \delta \rightarrow 0),
 \end{aligned}$$

where $\phi_i(\delta)$ ($i=1, 2, \dots$) denotes a function of δ independent of r, t , tending to 0 as $\delta \rightarrow 0$. Using the estimates (2.24), (2.28), (2.29), (3.3) and the hypothesis that $D_i^3 f$ is uniformly continuous in L_2 , we can show in a straight forward manner

$$(3.4) \qquad \|D_i^3 F(\cdot, t + \delta, u_r(t + \delta)) - D_i^3 F(\cdot, t, u_r(t))\|_{L_2}^2 \leq \phi_2(\delta)$$

and

$$(3.5) \qquad \|D_i^3 u_r(\delta) - D_i^3 u_r(0)\|_E^2 \leq \phi_3(\delta).$$

Therefore, applying Lemma 2.3 we have

$$(3.6) \qquad \max_{t \in R^+} \|D_i^3 u_r(t + \delta) - D_i^3 u_r(t)\|_E^2 \leq \phi_3(\delta) + c_4 \phi_2(\delta),$$

which implies that $\{D_i^3 u_r(t)\}$ is uniformly equicontinuous in the norm $\|\cdot\|_E$, and we can conclude by the resonance theorem

$$(3.7) \qquad D_i^3 u(t) \in B^0(R^+; E).$$

Now, the limit functions $D_i^i u(t)$ ($0 \leq i \leq 2$) satisfy the following elliptic equations in a generalized sense

$$\begin{aligned}
 (3.8) \qquad AD_i^i u(t) &= -(D_i^{2+i} u(t) + \nu D_i^{i+1} u(t) + D_i^i F(\cdot, t, u(t))) \\
 &\quad (\text{a. e. } t \in R^+).
 \end{aligned}$$

Since $D_i u, D_i^2 u, F(\cdot, t, u(t)) \in B^0(R^+; L_2)$, the equation (3.8) with $i=0$ implies

$$(3.9) \qquad u \in B^0(R^+; H_{2m} \cap \dot{H}_m) \subset C^0(\Omega \times R^+) \cap L^\infty(\Omega \times R^+)$$

and hence

$$D_i F(\cdot, t, u(t)) \in B^0(R^+; L^2(\Omega)).$$

Therefore (3.8) with $i=1$ implies

$$(3.10) \qquad D_i u \in B^0(R^+; H_{2m} \cap \dot{H}_m) \subset C^0(\bar{\Omega} \times R^+) \cap L^\infty(\Omega \times R^+).$$

By (3.1), (3.7) and (3.10) it is easy to see that the right hand side of (3.8) with $i=2$ belongs to $B^0(R^+; L_2)$, and we have

$$(3.11) \quad D_i^2 u \in B^0(R^+; H_{2m} \cap \dot{H}_m) \subset C^0(\bar{\Omega} \times R^+) \cap L^\infty(\Omega \times R^+).$$

Finally we note that $g(\cdot, u) \in B^0(R^+; H_{[n/2]+1})$ because $u(t) \in B^0(R^+; H_{2m}) \subset B^0(R^+; H_{[n/2]+1})$ (cf. Mizohata [Theorem 7.1; 10]). Thus by (3.10) and (3.11) the right hand side of the equation (3.8) with $i=0$ belongs to $B^0(R^+; H_{[n/2]+1})$ and by the theory of elliptic equation we have

$$(3.12) \quad u(t) \in B^0(R^+; H_{2m+[n/2]+1} \cap \dot{H}_m) \subset B^0(R^+; C^{2m}(\bar{\Omega})).$$

We summarize above results in the following

THEOREM 3.1. *Suppose H_1-H_3 and let $M < M_1$. Then the initial-boundary value problem (E)-(B) on $\Omega \times R^+$ with initial data $u(x, 0) \equiv u'(x, 0) \equiv 0$ admits a unique bounded classical solution u satisfying (2.3), (2.24), (2.28), (2.29) and*

$$(3.13) \quad \|u(t)\|_Q \equiv \max_{x \in \bar{\Omega}} \left(\sum_{|\alpha| \leq 2m} |D^\alpha u(x, t)| + \sum_{i=0}^2 \left| \frac{\partial^i}{\partial t^i} u(x, t) \right| \right) \leq k_4(M) \quad \text{for } t \in [0, \infty),$$

where $k_4(M)$ is a certain constant tending to 0 as $M \rightarrow 0$.

The proof of the uniqueness in Theorem 3.1 is trivial and omitted. Also it is easy from the procedure deriving the estimates (3.10)-(3.12) to see that the estimate (3.13) holds.

§ 4. Bounded, periodic and almost periodic solution.

Let us make the assumptions as in Theorem 3.1. Then the problem

$$(4.11) \quad \frac{\partial^2}{\partial t^2} u + \sum_{|\alpha|, |\beta| \leq m} (-1)^{|\alpha|} D^\alpha (a_{\alpha, \beta}(x) D^\beta u(x, t)) + \nu \frac{\partial}{\partial t} u + F(x, t, u) = 0$$

$$\text{on } \Omega \times (-r, \infty), \quad u(x, -r) = \frac{\partial}{\partial t} u(x, -r) = 0$$

and $D^\alpha u|_{\partial \Omega} = 0 \quad \text{for } |\alpha| \leq m-1 \quad (r=1, 2, 3, \dots)$

admits a unique bounded solution $u_r(x, t)$ satisfying (2.3), (2.24), (2.28), (2.29) and (3.11)-(3.13) for $t \geq -r$. Also it is easily seen from the proof of (3.13) that for each $T > 0$ the family $\{u_r(t)\} (r \geq T)$ is uniformly equicontinuous in the norm $\|\cdot\|_Q$ on $[-T, T]$. Since the imbedding maps from $H_{2m+[n/2]+1}$ and H_{2m} into $C^{2m}(\bar{\Omega})$ and $C(\bar{\Omega})$, respectively, are both compact, the vectorial

Ascoli-Arzelà's Lemma implies that $u_r(t)$ is convergent to a function $u(x, t)$ on $\bar{Q} \times R$ in the following sense; for each $T > 0$ and for $r > T$,

$$u_r(x, t) \longrightarrow u(x, t) \quad \text{in } C^{2m}(\bar{Q}) \quad \text{uniformly on } [-T, T]$$

and

$$\frac{\partial^i}{\partial t^i} u_r \longrightarrow \frac{\partial^i}{\partial t^i} u \quad \text{in } C(\bar{Q}) \quad \text{uniformly on } [-T, T]$$

($i=1, 2$).

Clearly $u(x, t)$ is a classical solution of the equation (1) on $\Omega \times R$, satisfying the same bounded properties as $u_r(x, t)$. Thus we have

THEOREM 4.1. *Under the same assumptions in Theorem 3.1 the problem (E)-(B) admits a unique bounded classical solution u satisfying the estimates (2.3), (2.24), (2.28), (2.29) and (3.13) for $t \in R$.*

PROOF. It remains to show the uniqueness of bounded solution satisfying such estimates. In fact this follows from the estimates (2.3) and (2.24) only. Indeed, let u and v be two bounded solutions satisfying (2.3) and (2.24) for $t \in R$. Then $w = u - v$ is a bounded solution of the equation

$$(4.2) \quad w'' + Aw + \nu w' = g(\cdot, u) - g(\cdot, v).$$

Now, using the mean value theorem, we can show easily

$$\|g(\cdot, u) - g(\cdot, v)\|_{L_2}^2 \leq 2K_0^2 \sum_{i=1,2} S_{2r_i+2}^{2r_i+2} x_0 (M)^{2r_i} \|w\|_A^2,$$

and hence by Lemma 2.2

$$\begin{aligned} \max_{s \in [t, t+1]} \|w(s)\|_E^2 &\leq c_3 (\|w(t)\|_E^2 - \|w(t+1)\|_E^2) \\ &\quad + 2c_4 K_0^2 \sum_{i=1,2} S_{2r_i+2}^{2r_i+2} x_0 (M)^{2r_i} \|w(t)\|_E^2. \end{aligned}$$

Thus, recalling the way of choice of the number M_1 , we have

$$\max_{s \in [t, t+1]} \|w(s)\|_E^2 \leq c_3 c_5^{-1} (M) (\|w(t)\|_E^2 - \|w(t+1)\|_E^2)$$

for $t \in R$, which together with the boundedness of $\|w(t)\|_E$ implies easily $w(t) \equiv 0$ (see [14]).

Q. E. D.

The following is an immediate consequence of the above theorem.

COROLLARY 4.1. *In addition to the assumption of Theorem 4.1, suppose that $f(\cdot, t)$ is ω -periodic in t . Then the bounded solution $u(x, t)$ in Theorem 4.1 is also ω -periodic.*

Now we shall study the almost periodicity of the bounded solution.

THEOREM 4.2. *In addition to the assumption of Theorem 4.1, suppose that $f(\cdot, t)$ is almost periodic as a function from R to $C(\bar{\Omega})$. Then the bounded solution $u(x, t)$ is almost periodic with respect to the norm $\|\cdot\|_Q$.*

PROOF. Let $\{t_r\}_{r=1}^\infty$ be any real sequence. By Bochner's criterion (see [3]) it suffices to show that $\{u(t+t_r)\}$ contains a subsequence which is convergent in the norm $\|\cdot\|_Q$ uniformly on R . Suppose that this were false. Then there would exist a positive constant ε_0 , two subsequences $\{t_{r_i}\}$ ($i=1, 2$) of $\{t_r\}$ and a real sequences s_r such that

$$(4.3) \quad \|u(s_r+t_{r_1})-u(s_r+t_{r_2})\|_Q \geq \varepsilon_0.$$

Consider the sequences of functions $\{u(t+t_{r_i}+s_r)\}$ ($i=1, 2$). They satisfy the estimate (2.3), (2.24), (2.28), (2.29) for $t \in R$, and by the same reason in the proof of Theorem 4.1 we may assume $u(t+t_{r_i}+s_r) \rightarrow U_i(t)$ in $\|\cdot\|_Q$ uniformly on each compact set in R as $r \rightarrow \infty$. $U_i(t)$ ($i=1, 2$) have, of course, the same boundedness properties as $u(x, t)$. Now, since $f(t)$ is almost periodic in $C(\bar{\Omega})$, we may assume $f(t+t_{r_i}+s_r)$ converge to a function $F(x, t)$ in the norm $C(\bar{\Omega})$ uniformly on R , and hence $U_i(t)$ ($i=1, 2$) are both bounded classical solutions of the problem (E)-(B) with $f(x, t)$ replaced by $F(x, t)$. Therefore the local uniqueness concerning bounded solution implies $U_1(t)=U_2(t)$. However by (4.3) we have

$$\|U_1(0)-U_2(0)\|_Q \geq \varepsilon_0 > 0$$

which is a contradiction.

Q. E. D.

§5. Asymptotic properties of bounded solutions.

In this section we shall investigate some asymptotic properties of the bounded solutions of the problem (E)-(B). For this purpose we let $u_i(t)$ ($i=1, 2$) be solutions on $\Omega \times (-r_0, \infty)$ of the equation (1) with $f=f_i$ ($i=1, 2$), respectively. Let us suppose that $u_i(t)$ fulfill the following boundedness conditions:

$$(5.1) \quad \|u_i(t)\|_A \leq a_0 \quad (< x_0(M_1)), \quad \|u_i'(t)\|_{L_2} \leq b_0 \quad (< k_0(M_1)) \\ \|D^j u_i(t)\|_E \leq a_j \quad (< k_j(M_1)) \quad \text{for } j=1, 2, 3, \quad t \in [-r_0, \infty)$$

with positive constants a_0, b_0, a_j .

From (5.1), we note, the same arguments as in the proof of Theorem 3.1 give

$$(5.2) \quad \|u_i(t)\|_Q \leq a_4 \quad \text{for } j=1, 2, \quad t \in [-r_0, \infty)$$

with some positive constant a_4 .

Assume that $f_i(t)$, $i=1, 2$, satisfy H_2 and we put

$$\delta_j(t) = \left(\int_t^{t+1} \|D^j_t(f_1(s) - f_2(s))\|_{L_2}^2 ds \right)^{1/2}$$

($j=0, 1, 2, 3$).

Then similar arguments as in the proofs of Lemmas 2.2-2.6 yield the following inequalities

$$(5.3) \quad \max_{s \in [t, t+1]} \|w(s)\|_E^2 \leq c_3 c'_5(a_0)^{-1} (\|w(t)\|_E^2 - \|w(t+1)\|_E^2) + 2c_4 c'_5(a_0)^{-1} \delta_0(t)^2,$$

$$(5.4) \quad \max_{s \in [t, t+1]} \|D_t w(s)\|_E^2 \leq c_3 c'_5(a_0)^{-1} (\|D_t w(t)\|_E^2 - \|D_t w(t+1)\|_E^2) + 4c_4 c'_5(a_0)^{-1} \left(2K_0^2 \sum_{i=1, 2} S_{4+2\bar{r}_i}^{4+2\bar{r}_i} a_0^{2\bar{r}_i} a_1^2 \|w(t)\|_E^2 + \delta_1(t)^2 \right)$$

and

$$(5.5) \quad \max_{s \in [t, t+1]} \|D_t^2 w(s)\|_E^2 \leq c_3 c'_5(a_0)^{-1} (\|D_t^2 w(t)\|_E^2 - \|D_t^2 w(t+1)\|_E^2) + 4c_4 c'_5(a_0)^{-1} \left\{ 2K_0^2 \sum_{i=1, 2} \left[S_{6+2\bar{r}_i}^{6+2\bar{r}_i} a_0^{2\bar{r}_i} a_1^4 \|w(t)\|_E^2 + S_{4+2\bar{r}_i}^{4+2\bar{r}_i} a_0^{2\bar{r}_i} (4a_1^2 \|D_t w\|_E^2 + a_2^2 \|w(t)\|_E^2) \right] + \delta_2(t)^2 \right\}$$

where $w(t) = u_1(t) - u_2(t)$ and

$$c'_5(a_0) = 1 - 4c_4 K_0^2 \sum_{i=1, 2} S_{2r_i+2}^{2r_i+2} a_0^{2r_i} \quad (>0).$$

Here we make an additional assumption;

H_2' . $g(x, u)$ is 4-times continuously differentiable in u .

Then we can easily show

$$\int_{\mathcal{Q}} |D_t^3 g(x, u_1(t)) - D_t^3 g(x, u_2(t)) + D_t^3 (f_1 - f_2)|^2 dx \leq c_6(a_4) \left(\sum_{i=0}^2 \|D_t^i w(t)\|_E^2 + \delta_3(t)^2 \right) + 4K_0^2 \sum_{i=1, 2} S_{2r_i+2}^{2r_i+2} a_0^{2r_i} \|D_t^3 w(t)\|_E^2$$

with a constant $c_6(a_4)$ depending on a_4 , and hence we have by Lemma 2.2

$$(5.6) \quad \max_{s \in [t, t+1]} \|D_t^3 w(s)\|_E^2 \leq c_3 c'_5 (a_0)^{-1} (\|D_t^3 w(t)\|_E^2 - \|D_t^3 w(t+1)\|_E^2) + 2c_4 c'_5 (a_0)^{-1} c_6 (a_4) \left(\delta_3(t)^2 + \sum_{i=0}^2 \|D_t^i w(t)\|_E^2 \right).$$

The following elementary lemma is useful.

LEMMA 5.1. *Let $\phi(t)$ be a bounded nonnegative function on $[-r_0, \infty)$ satisfying*

$$\max_{s \in [t, t+1]} \phi(s) \leq c_7 (\phi(t) - \phi(t+1)) + k(t)$$

with $c_7 > 0$ and $k(t) \leq c_8 e^{-\theta(t-r_0)}$ ($c_7, \theta > 0$).

Then we have

$$\phi(t) \leq c_9 e^{-\eta(t-r_0)} \quad (t \geq r_0)$$

where c_9 and η are positive constants depending on c_7, c_8 , and θ .

The above lemma is a special case of Lemma 2.1 in [16] and the proof is omitted (see also [15]).

Combining Lemma 5.1 with the estimates (5.3), (5.4), (5.5) and (5.6) we obtain that if $\delta_j(t) \leq K e^{-\theta(t-r_0)}$ ($j=0, 1, 2, 3$) with $\theta > 0$, then

$$(5.7) \quad \sum_{j=0}^3 \|D_t^j w(t)\|_E \leq K' e^{-\eta(t-r_0)} \quad \text{for } t \geq r_0$$

with some positive constants $K', \eta > 0$.

Thus combining (5.7) and the standard theory of elliptic equations as was used in the proof of Theorem 3.1 we can conclude

THEOREM 5.1. *Suppose H_1, H_2, H_2', H_3 and let*

$$\sum_{j=0}^3 \delta_j(t) + \left(\int_t^{t+1} \|f_1(s) - f_2(s)\|_{H_{2m}}^2 ds \right)^{1/2} \leq K_1 e^{-\theta(t-r_0)} \quad (\theta > 0).$$

Then for any solutions $u_i(t)$ ($i=1, 2$) on $[-r_0, \infty)$ of the problem (E)-(B) with $f(t)$ replaced by $f_i(t)$, which satisfy (5.1), we have

$$(5.8) \quad \|u_1(t) - u_2(t)\|_Q \leq K_2 e^{-\eta(t-r_0)} \quad \text{for } t \geq r_0$$

with some positive constants K_2, η .

The following is immediate from above.

COROLLARY 5.1. *The bounded solution in Theorem 4.1 is exponentially stable with respect to the norm $\|\cdot\|_Q$ in the class of solutions satisfying (5.1). And also if $\sum_{j=0}^3 \|D_t^j f(t)\|_{L_2} + \|f(t)\|_{H_{2m}}$ decays exponentially as $t \rightarrow \infty$, then the bounded solution decays exponentially in the norm $\|\cdot\|_Q$.*

The former part of the Corollary can be regarded as an extension in part of the result of Sattinger [21], where the existence of global classical solution and the asymptotic stability of the stationary solution have been investigated for the equation

$$u_{tt} - \sum_{i,j} \frac{\partial}{\partial x_j} \left(a^{ij} \frac{\partial}{\partial x_i} u(x,t) \right) + 2\alpha u_t = P(u), \quad \alpha > 0,$$

with linear homogeneous boundary condition $\mathfrak{B}u=0$ on $\partial\Omega$, when $n=1, 2, 3$. In [21], the method of developing the solutions in power series is employed and quite different from ours. Quite recently, A. Matsumura [26] has extended the result of [21] to the more general nonlinear equations of second order. The method of [26], however, is also different from ours.

References

- [1] S. Agmon, Lectures on elliptic boundary value problems, Van Nostrand C., 1965.
- [2] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary value conditions, I, *Comm. Pure Appl. Math.*, **12** (1959), 623-727.
- [3] L. Amerio and G. Prouse, Almost periodic functions and functional equations, Van Nostrand C., 1971.
- [4] M. Biroli, Sulla esistenza ed unicit  della soluzione limitata e della soluzione quasi periodica per una equazione parabolica con termine dissipativo non linear discontinuo, *Ricerche Mat.*, **19** (1970), 93-110.
- [5] M. Biroli, Sur l'equation des ondes avec un terme non lineaire monotone dans la fonction inconnue, *Nota 1, Atti. Accad. Naz. Lincei*, **53** (1972), 359-361.
- [6] J.C. Clements, Existence theorems for some non-linear equations of evolution, *Can. J. Math.*, **22-4** (1970), 726-745.
- [7] Y. Ebihara, M. Nakao and T. Nanbu, On the existence of global classical solution for $\square u - u^3 = f$, *Pacific J. Math.*, **61** (1975), 63-70.
- [8] W.S. Hall, On the existence of periodic solutions for the equations $D_{tt} + (-1)^p D_x^{2p} u = \epsilon f(\cdot, \cdot, u)$, *J. Differential Equations*, **7** (1970), 509-526.
- [9] J.L. Lions, *Quelques methodes de resolution des problemes aux limites nonlineaire*, Dunod Paris, 1969.
- [10] S. Mizohata, *Theory of partial differential equations*, (in Japanese), Iwanami, Tokyo, 1965, Cambridge UP., 1973.
- [11] M. Nakao, On some regularity of periodic solutions of nonlinear wave equations, *J. Differential Equations*, **17** (1975), 187-197.
- [12] M. Nakao, Periodic solutions of linear and nonlinear wave equations, *Arch. Rational Mech. Anal.*, **62** (1976), 87-98.
- [13] M. Nakao, On boundedness, periodicity and almost periodicity of solutions of some nonlinear parabolic equations, *J. Differential Equations*, **19** (1975), 371-385.
- [14] M. Nakao, Bounded, periodic or almost periodic solutions of nonlinear hyperbolic partial differential equations, *J. Differential Equations*, **23** (1977), 368-386.
- [15] M. Nakao, Asymptotic stability of the bounded or almost periodic solution of the wave equation with a nonlinear dissipative term, *J. Math. Anal. Appl.*, **58** (1977),

- 336-343.
- [16] M. Nakao, Convergence of solutions of the wave equation with a nonlinear dissipative term to the steady state, Mem. Fac. Sci. Kyushu Univ., **30** (1976), 257-265.
 - [17] M. Nakao and T. Nanbu, Bounded or almost periodic classical solutions for some non-linear parabolic equations, Mem. Fac. Sci. Kyushu Univ., **30** (1976), 191-211.
 - [18] P.H. Rabinowitz, Periodic solutions of nonlinear hyperbolic partial differential equations, Comm. Pure Appl. Math., **20** (1967), 145-205.
 - [19] P.H. Rabinowitz, Periodic solutions of nonlinear hyperbolic partial differential equations, II, Comm. Pure Appl. Math., **22** (1969), 15-39.
 - [20] J. Sather, The existence of a global classical solution of the initial boundary value problem for $\square u + u^3 = f$, Arch. Rational Mech. Anal., **22** (1966), 292-307.
 - [21] D.H. Sattinger, Stability of nonlinear hyperbolic equations, Arch. Rational Mech. Anal., **28** (1968), 226-244.
 - [22] G. Torelli, Soluzioni periodiche dell'equazione nonlineare $u_{tt} - u_{xx} + \varepsilon F(x, t, u) = 0$, Rend. Ist. Mat. Univ. Trieste, 1969, 123-137.
 - [23] O. Vejvoda, Periodic solutions of a linear and weakly nonlinear wave equation in one dimension, Czech. Math. J., **14** (1964), 341-382.
 - [24] W. V. Wahl, Periodic solutions of nonlinear wave equations with a dissipative term, Math. Ann., **190** (1971), 313-322.
 - [25] M. Yamaguchi, On the existence of quasiperiodic solutions of nonlinear partial differential equations, Proc. Fac. Sci. Tokai Univ., **10** (1974), 1-12.
 - [26] A. Matsumura, Global existence and asymptotics of the solutions of the second-order quasilinear hyperbolic equations with the first-order dissipation, Publ. Res. Inst. Math. Sci. A, **13** (1977), 349-379.

Mitsuhiro NAKAO
 Department of Mathematics
 Faculty of Science
 Kyushu University
 Fukuoka 812, Japan

Present address
 Department of Mathematics
 College of General Education
 Kyushu University
 Fukuoka 810, Japan