

Breeding for plant heat tolerance at vegetative and reproductive stages

Nicky Driedonks¹ · Ivo Rieu¹ · Wim H. Vriezen²

Received: 20 October 2015 / Accepted: 21 January 2016 / Published online: 13 February 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Key message Thermotolerant crop research.

Abstract Global warming has become a serious worldwide threat. High temperature is a major environmental factor limiting crop productivity. Current adaptations to high temperature via alterations to technical and management systems are insufficient to sustain yield. For this reason, breeding for heat-tolerant crops is in high demand. This review provides an overview of the effects of high temperature on plant physiology, fertility and crop yield and discusses the strategies for breeding heat-tolerant cultivars. Generating thermotolerant crops seems to be a challenging task as heat sensitivity is highly variable across developmental stages and processes. In response to heat, plants trigger a cascade of events, switching on numerous genes. Although breeding has made substantial advances in developing heat-tolerant lines, the genetic basis and diversity of heat tolerance in plants remain largely unknown. The development of new varieties is expensive and time-consuming, and knowledge of heat tolerance mechanisms would aid the design of strategies to screen germplasm for heat tolerance traits. However, gains in heat tolerance are limited by the often narrow genetic diversity. Exploration and use of wild relatives and landraces in breeding can increase useful

genetic diversity in current crops. Due to the complex nature of plant heat tolerance and its immediate global concern, it is essential to face this breeding challenge in a multidisciplinary holistic approach involving governmental agencies, private companies and academic institutions.

Keywords Heat tolerance · Plant breeding · Plant biotechnology · Crops · Knowledge transfer

Introduction

Ambient temperatures are rising at a considerable rate as part of the current global climate change. The last three decades are thought to be the warmest the earth has experienced in the past 1400 years in the Northern Hemisphere. Climate models predict that the global mean temperature will continue this trend, increasing by 1–4 °C by the end of the twenty-first century. Additionally, climatological extremes such as heat waves are likely to occur more frequently (IPCC 2013; Tebaldi et al. 2006; Hansen et al. 2015). While the above data refer to average global temperature increases, there are significant regional and seasonal differences with further potential impact on agriculture (IPCC 2007). The biggest temperature changes will be at higher latitudes (IPCC 2007). In these regions, the increase in temperature might benefit overall crop production by alleviating low-temperature growth inhibition at the start of the growing season, allowing earlier planting of crops, and the possibility of a longer growing season or more cropping cycles per year in the longer term (Gitay et al. 2001). Thus, a rise in temperature is expected to lead to expansion of areas suitable for crop production in the Russian Federation, North America and Northern Europe as well as in East Asia (Lotze-Campen and Schellnhuber 2009; Olesen and Bindi 2002). Offsetting these

Communicated by Enrico Schleiff.

A contribution to the special issue ‘Pollen development and stress response’.

✉ Ivo Rieu
i.rieu@science.ru.nl

¹ Department of Molecular Plant Physiology, Institute for Water and Wetland Research, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands

² Bayer CropScience Vegetable Seeds, PO Box 4005, 6080 AA Haelen, The Netherlands

benefits, however, are negative effects at lower latitudes, where temperatures are already at the higher end of the crops' optimal grow temperature ranges. Regions in Africa, for example, have been predicted to become (semi-)arid due to heat and water stress, resulting in significant yield losses (Fischer et al. 2002; Ortiz et al. 2008), and in Asia and the Middle East, crop yields are predicted to fall 15–35 % if the average temperature increases 3–4 °C (Ortiz et al. 2008; FAO 2009).

More than 200,000 plant species are estimated to exist globally, of which ~80,000 are edible to humans. Despite this huge variety, 95 % of the calories and protein intake of human and livestock are derived from only 20–25 species (Füleky 2009). For example, only three species, wheat, rice and maize, account for 75 % of global grain production (Bansal et al. 2014; Lobell and Gourdj 2012). Breeding and agronomic improvements of these species have resulted in increased production between 1985 and 2005, and previous IPCC projections assumed that this will continue in the future (Ainsworth and Ort 2010; Teixeira et al. 2013). However, based on an extended update of the IPCC projections, a new meta-study predicted a less optimistic scenario. Primarily due to a more negative effect of moderate warming on yield, worldwide yield reductions are now expected for wheat, rice and maize in both tropical and temperate regions under a scenario of 2 °C of local warming without adaptation (Challinor et al. 2014). Relative rates of yield increase for major cereal crops are already declining (Fischer and Edmeades 2010; Foley et al. 2011). However, to meet the demand for food from the population of an expected 9 billion people in 2050, a 70 % increase in food production is deemed necessary according to the Declaration of the World Summit on Food Security (FAO 2009). This means that the yearly increases in production for the coming 40 years need to be 38 % higher than those achieved historically (Tester and Langridge 2010).

Thus, the temperature changes associated with global warming have become a major challenge with respect to agricultural output. Here, we provide an overview of research that aims to support increasing heat tolerance of crops. We evaluate potential changes to crop management, the availability of genetic resources for breeding activities and the usefulness of (academic) research into the molecular and physiological basis of heat tolerance. Finally, it is discussed whether academic research and breeding activities are complementary with mutual benefits.

Effects of high temperature on crops

General physiological effects of high temperature

The intensity, duration and rate of temperature change together determine the impact of high temperature on plant

development and physiology (Wahid et al. 2007; Zinn et al. 2010). Air and soil temperature affect crop yield in different ways, and this should be considered when studying heat impact (Lobell and Gourdj 2012; Sharkey and Schrader 2006). Plant physiological responses to heat stress have been reviewed in great detail (Wahid et al. 2007; Bita and Gerats 2013; Boksztzanin et al. 2013; Mathur et al. 2014). In general, a moderate increase in air temperature leads to faster plant development and a shorter crop duration and consequently a reduction in cumulative light perception and assimilation over the plant's life cycle. In addition, disturbance of fundamental processes such as carbon assimilation, respiration and transpiration may reduce overall metabolic efficiency and result in vegetative developmental defects such as fewer, malformed and/or smaller organs (Takeoka et al. 1991; Maestri et al. 2002; Stone 2001). High air temperature can also negatively affect sexual reproduction and consequently fruit and seed yield (Peet et al. 1997; Erickson and Markhart 2002; Zinn et al. 2010). On the other hand, high soil temperature can reduce germination capability and plant emergence and can cause heat necrosis of roots (Stevenson et al. 2001).

Heat sensitivity in crop plants

Individual plant yield is a function of various components including plant architecture, photosynthetic efficiency, resource partitioning and reproductive success—each of these components may be vulnerable to heat. Optimum temperature range and consequently heat sensitivity vary among crop types, species and cultivars (Ulukan 2008; Levy and Veilleux 2007; Luo 2011; Saha et al. 2010). Heat sensitivity has been shown to cause yield reduction in species of both temperate and tropical zones, but in general, tropical varieties often tolerate higher temperatures better, compared to varieties of the same crop species grown in temperate zones, as was shown for yard-long bean, cucumber and radish (Wahid et al. 2007; Momonoki and Momonoki 1993; Yamamoto et al. 2011). Similarly, warm-season annuals usually cope better with high temperatures than cool-season annuals. For example, for the warm-season annuals cowpea and rice, the maximum temperature for emergence is 37 and 40 °C, respectively (Yoshida et al. 1981; Akman 2009), while cool-season crops such as chickpea, lentils and lettuce show decreased germination rate at soil temperatures above 33, 24 and 32 °C, respectively (Covell et al. 1986; Hall 2001).

Various temperature thresholds of a range of crops, including cereals, horticultural and legume crops, have been reviewed in detail (Luo 2011). Most crops suffer if high temperatures are encountered during the vegetative growth period, as has been documented for both cool-season annuals like wheat (Porter and Gawith 1999) and

Brassica juncea (Hayat et al. 2009) and for warm-season annuals, such as rice (Peng et al. 2004; Lyman et al. 2013), maize (Crafts-Brandner and Salvucci 2002), legumes (McDonald and Paulsen 1997) and tomato (Camejo et al. 2005).

However, for many crops, including rice, maize, soybean, legumes, rapeseed, sunflower and tomato, the reproductive stage appears to be even more vulnerable to temperature increase (Jagadish et al. 2014; Zinn et al. 2010; Barnabás et al. 2008; Hedhly et al. 2009). This is especially true during inflorescence/panicle development and during flowering, where heat may lead to flower abortion or reduced fertility, respectively (Maduraimuthu and Prasad 2014; Luo 2011).

Heat and fertility

Reduced fertility is a common problem associated with heat, and has been found to be caused by high temperatures during meiosis and fertilization in various species, e.g., *Arabidopsis*, tomato, rice, cowpea and barley (Bac-Moleenaar et al. 2015; Giorno et al. 2013; Jagadish et al. 2014; Ehlers and Hall 1998; Sakata and Higashitani 2008). The male reproductive organs and, in particular, pollen development are the most heat sensitive. Exposure to high temperature stress during flowering results in a reduction of viable and germinating pollen (Sato et al. 2006; Abiko et al. 2005; Prasad et al. 2006; Oshino et al. 2007; Jagadish et al. 2010; Zinn et al. 2010; Peet et al. 1998). For example, in rice, spikelet sterility occurs if temperatures exceed 35 °C for just 1 h (Yoshida et al. 1981; Endo et al. 2009). Tomato also has a dramatic decrease in fruit set in response to heat stress, especially when applied during microsporogenesis (Zhang and Yang 2014). During this stage, a short period at 40 °C or extended exposure just a few degrees above optimal temperature (32 °C rather than 26 °C during the day) results in male sterility (Sato et al. 2006; Giorno et al. 2013). Similarly, barley shows vulnerability to chronic mild heat stress (30 °C/25 °C day/night, for 5 days), by failure of tapetum differentiation and injuries to the microsporogenesis process (Sakata and Higashitani 2008).

Maternal tissues of the pistil and the female gametophyte have traditionally been considered to be more thermotolerant. However, malformations of the female tissues can occur in some species when subjected to heat. Embryo sac malformations have been reported in peach developed above 25 °C, in wheat at 30 °C and rapeseed at 32 °C, which consequently reduced the seed set in the latter two species (Hedhly 2011). In apricot, even a mild increase of 3 °C above control conditions during the last week of flower development resulted in shortening of the style and abnormal ovaries (Rodrigo and Herrero 2002). In addition

to effects on male tissue, stigma receptivity is shortened by heat in cherry and peach, and ovule longevity is reduced in cherry and plum (Endo et al. 2009; Hedhly 2011). Such alterations result in a lack of synchrony between male and female reproductive tissues, ultimately leading to reduced fertilization efficiency. However, timely pollination does not guarantee fruit or seed set, as post-pollination processes such as pollen tube growth, fertilization, formation of the endosperm and embryo development were also shown to be heat sensitive (Peet et al. 1997; Erickson and Markhart 2002; Barnabás et al. 2008).

Adaptation of cultivation methods to avoid heat stress

As part of the plant's phenotype, yield is the result of the expression of the genotype (G), the environment (E) and their interaction (G × E). In the field of agriculture, management practices (M) are often included as a separate third factor, leading to the G × E × M model. Thus, yield improvement can in principle be achieved by adapting the genotype, as discussed later, the environment, or the management practices.

At the level of farming, a few technical and management adjustments may contribute to an increased ability of crops to cope with temperature changes. Firstly, assuming concomitant higher winter temperatures, the dates of planting can be adapted to avoid heat stress later in the growing season (Olesen and Bindi 2002; Easterling 1996; Rosenzweig and Tubiello 2007; Lotze-Campen and Schellnhuber 2009). Crop planting and harvesting dates from around the world have been recorded and used to make a so-called crop calendar. This calendar contains ~1300 planting and harvesting date observations for 19 crops, allowing estimation of the effects of planting time patterns at many geographic locations (Sacks et al. 2010). For example, in the US Midwest, early planting seems to be a successful strategy to avoid summer heat for maize and spring wheat (Reilly et al. 2003). Secondly, improvements in water management can alleviate heat stress in agriculture, as plants transpire to keep foliage temperature under control. One option is shifting from rain-fed to irrigated agriculture, including low-cost “rainwater harvesting” practices. Additionally, adjusting the timing of irrigation may ensure a crop's water supply at critical, temperature-sensitive stages (Easterling 1996; Smithers and Blay-Palmer 2001; Smit and Skinner 2002; Lotze-Campen and Schellnhuber 2009).

Another strategy of avoiding heat stress is to change the “environment” factor of the GxExM model, by shifting the geographical location of crop cultivation. Although this strategy is drastic, it is already occurring, for instance, in

Australia's wine industry, where several large wine producers have bought new properties in cooler regions to maintain vineyards in the future (Chapman et al. 2012; Park et al. 2012). It is also occurring for other crops, like maize and rice (Kenny et al. 1993; Duzheng 2003; Tchebakova et al. 2011).

While adapting to climatic changes via alterations in cultivation may be possible for some crops, heat stress cannot be avoided by this approach alone (Reilly et al. 2003; Tubiello et al. 2002). For example, in areas where water is a scarce and valuable resource, improving water supply might not be an option. Similarly, moving cultivation areas geographically might be a solution for Australia's wine industry, but is not for Australia's wheat farmers, because winter temperatures at lower latitudes near the sea are also too high (Chapman et al. 2012). Because of these limitations, the introduction of more heat-tolerant cultivars or shifting to other crops is essential to maintain food production in areas with increasing temperatures.

Conventional breeding for heat tolerance

Increasing temperature tolerance by conventional breeding is an obvious approach to reduce the negative effects of heat on crop yield. Usually, breeding programs are carried out in a climacteric region similar to that where the crop will be produced eventually. Thus, the selection of breeding lines for relatively hot regions takes place under hot conditions (Mickelbart et al. 2015). This implies that in hot regions, thermotolerance traits are “passively” selected for by locally operating breeders. Considering that cultivars from warmer regions are often more heat-tolerant than those from cooler regions, it seems that this technique has been rewarding (Tonsor et al. 2008; Smillie and Nott 1979; Yamamoto et al. 2011; Momonoki and Momonoki 1993; Kugblenu et al. 2013). Conventional breeding has also been used to intentionally develop new heat-tolerant crop genotypes. For example, a variety of broccoli has an improved head quality thanks to early maturation, because this trait prevents hot days later in season to affect the heat-sensitive flower initiation developmental stage (Farnham and Bjorkman 2011). In addition, new varieties of cowpea showed higher average grain yield when grown under hot and long days during reproduction (Ehlers and Hall, 1998) and recurrent selection has also been successful for improving wheat yield using ancestor *T. tauschii* as a gene donor, leading to increased rates of grain filling and larger grains in BC1F6 plants (Gororo et al. 2002). Finally, in potato breeding a genetic gain was obtained after three cycles of recurrent selection for heat tolerance leading to

strong increase in yield up to 37.8 % (Benites and Pinto 2011).

Although conventional “yield” breeding has succeeded in developing heat-tolerant lines, the ultimate genetic and physiological bases of the improvements remain unclear. This prevents the development of molecular or other biomarkers, which would assist germplasm screening for improved heat tolerance and allow for efficient breeding of the complex trait. Another drawback of conventional breeding is that the programs are often based on crossing relatively advanced starting material, which has already been used in the particular breeding areas specifically related to the market segment that is targeted. This implies that the potential gain in heat tolerance level is limited by the low genetic diversity (Ladizinsky 1985; Paran and Van Der Knaap 2007).

Advanced breeding for heat tolerance

Intra-specific QTL discovery

Heat tolerance seems to be polygenic, which might explain why the genetic basis of heat stress tolerance in plants is poorly understood (Wahid et al. 2007; Ainsworth and Ort 2010; Collins et al. 2008). In order to improve knowledge about thermotolerance at the genetic level, many efforts have been made to identify quantitative trait loci (QTL) in segregating mapping populations. Jha et al. (2014) recently listed QTLs associated with heat tolerance in various plants, including *Arabidopsis*, azuki bean, barley, brassica, cowpea, maize, potato, rice, sorghum, tomato and wheat. In this paper, the authors showed several types of genetic markers linked to different traits of interest which spanned the various aspects of a plant's vulnerability to heat. This included QTLs for yield traits, such as fruit set or grain filling rate, under heat. Also, QTLs for several heat tolerance-related traits have been discovered, such as for lower canopy temperature during vegetative and reproductive stages and higher chlorophyll fluorescence in wheat (Pinto et al. 2010; Lopes et al. 2013; Vijayalakshmi et al. 2010). High chlorophyll fluorescence represents heat-tolerant photosynthesis, and lower canopy temperature reflects efficient water uptake which has been associated with deep rooting (Pinto and Reynolds 2015). A major QTL for high-temperature seed germination capacity in lettuce, *Htg6.1*, colocalizes with a temperature-sensitive gene encoding an abscisic acid biosynthesis enzyme (*LsNCED4*) (Argyris et al. 2008, 2011). In potato, nine QTLs for internal heat necrosis in tubers were detected that each explain between 4.5 and 29.4 % of the phenotypic variation (McCord et al. 2011). Many studies have focused on the effect of high

temperature on reproductive characteristics, including pollen germinability, pollen tube growth, grain weight, days to heading, grain filling and post-anthesis leaf senescence, fruit set and quality traits such as white-back kernels in rice. In maize, five and six QTLs for pollen quality and tube growth have been identified with a high heritability of 0.64 and 0.68, respectively. However, the pollen tests were performed in vitro and might not be representative of the situation in vivo (Frova and Sari-Gorla 1994). Lastly, in tomato, six QTLs were identified that explain 33 % of the phenotypic variation related to fruit set at high temperature (Ventura et al. 2007). A recent QTL study in rice focussed on spikelet fertility under high temperature (Ye et al. 2015). This study followed up previous work (Ye et al. 2012) and confirmed the presence of a recessive QTL on chromosome 4 which contributes 15 % higher rice spikelet fertility under heat stress compared to plants without the QTL (Ye et al. 2015). Heat tolerance QTLs on this chromosome have been identified in different populations of heat-tolerant rice varieties (Ye et al. 2012, 2015; Xiao et al. 2010). Ye et al. (2015) showed that the QTL is located in a highly conserved chromosomal region. Unfortunately, this limits mapping resolution and causal gene identification. More advanced approaches such as the use of multi-parent advanced generation inter-cross (MAGIC) populations were suggested as a way to introduce more genetic variation and determine the genes involved in thermotolerance of spikelet fertility (Ye et al. 2015).

Besides dedicated mapping populations, QTLs can be detected via exploration of natural populations. As noted previously, linkage mapping can be considered as being useful for identification of major genes and QTLs. However, due to the limited number of generations and thus recombination events, those QTLs cover a relatively large region and gene identification requires time-consuming fine-mapping processes. Exploiting natural diversity panels avoids these hurdles. Using a genome-wide association approach, the linkage decay is fast, therefore providing a much higher resolution. Consequently, fine mapping is often not necessary for identification of candidate genes (Bergelson and Roux 2010). So far, genome-wide association study (GWAS) panels have been established in *Arabidopsis* and several crops including maize, rice, sorghum and foxtail millet (Buckler et al. 2009; Huang et al. 2011; Jia et al. 2013; Morris et al. 2013). Maize and rice seem to be the two major models for crop GWAS, considering the magnitude of resources already developed and published for these species (Huang and Han 2014). So far, QTLs explaining transition of the vegetative to generative stage have been established in these crops (Buckler et al. 2009; Huang et al. 2011). Although those QTLs are not related to heat tolerance directly, QTLs explaining flowering time are of interest for heat tolerance breeding, as

early flowering might enable a plant to complete the heat-sensitive reproductive processes before late-season heat episodes (Ishimaru et al. 2010). QTLs directly associated with high-temperature-induced reduction in fertility have recently been revealed in *Arabidopsis*. An *Arabidopsis* inflorescence can have flowers at many different developmental stages, and by measurement of the plant silique length after short-term heat stress, sensitivity of the different reproductive stages was determined. Meiosis, fertilization and early embryogenesis were shown to be most vulnerable to heat. GWAS study of this experimental setup revealed four QTLs related to specific developmental stages. Three QTLs were responsible for sensitivity of pre-anthesis reproductive processes including male and female meiosis, while one QTL explained population variation of early embryogenesis heat sensitivity. A strong negative correlation between flowering time and silique length was detected, which were strongly and moderately associated with the same SNP, respectively. Interestingly, this SNP has been linked to the flowering time repressor *FLC*, suggesting a role for the regulation of flowering time in the heat stress response (Bac-Molenaar et al. 2015). As *Arabidopsis* is a member of the *Brassicaceae* family, the results might provide insights for breeding within this family.

Altogether, the QTL studies in different crops all identified multiple QTLs per trait, varying from two in rice and azuki bean (enhancing spikelet and pollen viability under heat stress, respectively) up to 34 in barley for several heat-related traits (e.g., number of spikes per plant and days until heading). This demonstrates that heat tolerance is dependent on a range of factors and QTLs which seem to vary between crops (reviewed in Jha et al. 2014).

Expanding genetic diversity with crop-related wild species

Crop domestication may be regarded as the first stage of plant breeding, resulting in dramatic morphological and physiological modifications to meet human needs, including seed and fruit size and number, seed shattering, seed dormancy, photoperiod and flowering time, taste, nutrition and overall plant architecture (Meyer and Purugganan 2013; Gross and Olsen 2010). Domestication inevitably involves a genetic bottleneck due to selection and breeding of similar lines with favorable traits, after which only a subset of the genes and alleles available in the wild progenitor gene pool are present among crop cultivars (Godfray et al. 2010; Ladizinsky 1985; Olsen and Wendel 2013). This reduction in diversity seems to have led to a loss of abiotic stress tolerance traits, since many wild relatives and landraces are more tolerant to stresses compared to domesticated crops (Dolferus 2014; Maduraimuthu and

Prasad 2014). Therefore, the identification of superior wild alleles that are lacking in cultivated germplasm has become of great interest (Tanksley and McCouch 1997; Grandillo et al. 2007; Lippman et al. 2007; Feuillet et al. 2008). For example, introgression of wild alleles has resulted in crop improvement in several cereals such as rice (Atwell et al. 2014), wheat (Pradhan et al. 2012) and maize (Prasanna 2012). However, exploiting wild relatives as a source of novel alleles has been hindered by the introduction of linked, undesirable traits, compounded by a lack of molecular markers for precision breeding. (Dolferus 2014). Only recently, new sequencing methods have made it cost-effective to re-sequence complete genomes, as has been done for wild tomato (Aflitos et al. 2014), cucumber (Qi et al. 2013), sorghum (Mace et al. 2013), grape (Lijavetzky et al. 2007), soybean (Li et al. 2013), rice (Xu et al. 2010) and maize (www.panzea.org). Difficulties still arise when QTLs are crossed in a particular genetic background and show a smaller or no effect at all. Due to the potential for unfavorable epistatic interactions, it is difficult or even impossible to predict in advance whether a QTL might be transferable to elite backgrounds (Podlich et al. 2004; Collins et al. 2008).

Despite difficulties associated with fine mapping, identification of the causal gene was successful for a major QTL for seedling survival rates under high temperature in African rice (*Oryza glaberrima*) (Li et al. 2015). This species has several advantageous traits, including tolerance to drought, salinity and heat (Sakai et al. 2011). The heat tolerance QTL was associated with a single gene, *Thermotolerance 1 (TTI)*, coding for an $\alpha 2$ subunit of the 26S proteasome which is involved in the degradation of ubiquitinated proteins. This gene is thought to protect cells from heat stress by enhancing efficient elimination of cytotoxic denatured proteins and maintaining heat-response processes (Li et al. 2015). Comparison of the sequence of *TTI* from the Asian and African cultivated parental lines, *Oryza sativa* ssp. *japonica* and *Oryza glaberrima*, revealed three exonic SNPs, one of which resulted in an amino acid substitution. Interestingly, *TTI* was suggested to be a major determinant for variation in thermotolerance among *O. sativa* varieties: geographical distribution of three *TTI* haplotypes showed that environmental pressure was responsible for the selection of the *TTI* heat tolerance locus. In addition to a higher seedling survival, near-isogenic lines (NILs) containing the *O. glaberrima TTI* allele showed a higher thermotolerance at flowering and filling stages compared to NILs with an *O. sativa TTI* allele. Overexpression of this gene not only led to enhanced seedling thermotolerance in other rice species, but also in *Arabidopsis* and *Festuca elata* (Li et al. 2015), showing the enormous potential of this allele to enhance crop productivity under high-temperature stresses.

Together, these studies indicate that QTL analysis and subsequent fine mapping and cloning are promising ways to identify loci and genes for heat tolerance. Several candidate genes have been proposed, but characterizing the causal gene underlying a heat tolerance QTL remains challenging. However, for breeding purposes, the exact underlying genes do not have to be known. Using molecular markers based on linked flanking polymorphisms of a QTL, a QTL can still be successfully introduced into crossable breeding germplasm.

Discovery of thermotolerance genes

In response to high-temperature stress, plants modulate the expression of a plethora of genes. These genes and their annotation could help to identify the processes that are induced or repressed such as those involved in acclimation and protection to heat stress.

Transcriptional profiling has been performed during the onset and recovery of heat stress, between stressed and unstressed plants, or between heat-tolerant and heat susceptible variants. Such analyses have been performed in many crops, e.g., rice, tomato, barley, brassica and grape (Frank et al. 2009; Sarkar et al. 2014; Frey et al. 2015; Bitá et al. 2011; Liu et al. 2012; Mangelsen et al. 2011; Dong et al. 2015). An extensive summary of recent transcriptomic analysis in plant species was published by Lavania et al. (2015). This summary revealed that plants reprogram their signal transduction pathway, transcription factors and proteins associated with metabolism in a conserved manner. Although the studies were performed in different crops which were exposed to different heat regimes, there was considerable similarity in the heat stress-responsive genes.

For example, there is a conserved induction of genes encoding for enzymes that govern the fluidity of membranes upon heat stress. In agreement with this, overexpression of one of the enzymes: *glycerol-3-phosphate acyltransferase*, resulted in increased saturation of the thylakoid membrane lipids of transgenic tobacco plants, showed a faster recovery after heat stress compared to wild-type plants (Yan et al. 2008). When plants are exposed to heat, reactive oxygen species (ROS) are formed as a by-product in various aerobic metabolic pathways in different cellular compartments (Miller et al. 2009; Wang et al. 2014a; Chou et al. 2012; Dat et al. 1998; Volkov et al. 2006; Wu et al. 2012; Vacca et al. 2004; Mostofa et al. 2013), and cause cellular damage to membranes, proteins, lipids and DNA (Volkov et al. 2006; Wu et al. 2012; Boksaczanin et al. 2013; Baker and Orlandi 1995; Giardi et al. 1997; O’Kane et al. 1996; Larkindale and Knight 2002). In order to prevent damage to the cell and regain redox homeostasis, a typical response to heat is hyper-

activation of the ROS scavenging machinery. The transcript and protein level of genes responsible for ROS scavenging are increased under heat stress in many different plant species (Chou et al. 2012; Chao et al. 2009; Mittal et al. 2012; Suzuki et al. 2013) and this has been associated with basal heat tolerance (Almeselmani et al. 2006; Bhattacharjee 2012; Gupta et al. 1993; Kang et al. 2009; Rui et al. 1990; Wang et al. 2014b; Sairam et al. 2000; Badiani et al. 1993). A ROS scavenging-related gene that seems to be important for thermotolerance is glutaredoxin (GRX). This small ubiquitous protein is a regulator in diverse cellular processes and oxidative stress response, and its function is conserved in prokaryotes and eukaryotes (Lillig et al. 2008; Cheng et al. 2009; Wu et al. 2012). As a critical component of ROS metabolism, *Arabidopsis AtGRXS17* may be crucial for temperature-dependent postembryonic growth and development (Cheng et al. 2011). Indeed, improvement of plant heat stress tolerance has been achieved by increasing antioxidant enzyme and GRX activities (Almeselmani et al. 2006; Badiani et al. 1993; Gupta et al. 1993; Rui et al. 1990; Sairam et al. 2000; Wu et al. 2012; Chen et al. 2013). Probably, the best-studied mechanism in response to heat stress is the production of heat shock proteins (HSPs) upon exposure to high temperature (Wang et al. 2004). By acting as molecular chaperones, HSPs prevent deleterious protein conformations and eliminate non-native aggregations formed during stress (Morimoto 1998; Boston et al. 1996; Vierling 1991). Strong transcriptional up-regulation of a number of HSPs by heat stress has been shown in plants and many other organisms. The expression of HSPs and various other heat-responsive genes is controlled by heat shock transcription factors (HSFs) (Kotak et al. 2007). Experiments in *Arabidopsis*, rice, tobacco and tomato have shown that enhanced thermotolerance can be gained by overexpressing HSPs or HSFs (reviewed in Grover et al. 2013).

Conclusion

Despite the urgent need to improve crop heat tolerance, a very limited number of heat-tolerant varieties have been developed. The development of new varieties through plant breeding is expensive and time-consuming (Lotze-Campen and Schellnhuber 2009; Rosenzweig and Tubiello 2007)—for annual crops, it may take 10–30 years to introduce specific adaptations (Chapman et al. 2012; Rosenzweig and Tubiello 2007; Smit and Skinner 2002; Olesen and Bindi 2002). Therefore, it is very important that genetic variation for the trait can be identified and characterized efficiently in order to introduce it in a breeding program. At this point, fundamental research plays an important role as knowledge on molecular physiology of the plant heat response can

speed up the cloning of causal genes after QTL identification. Furthermore, fundamental knowledge may be used to generate leads for biotechnological modification of heat tolerance traits. Although genetic modification is controversial in some parts of the world, the products generated with new gene editing techniques may be in the near future classified as non-transgenic in the EU (<https://www.euroseeds.eu/new-plant-breeding-techniques>). Gene editing techniques involve several site-directed nuclease techniques such as SDN-1 and SDN-2 using zinc-finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), meganucleases and the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated Protein9 (Cas9) system (CRISPR/CAS9) (Sander and Joung 2014; Lusser et al. 2012; Hartung and Schiemann 2014; Mahfouz et al. 2014). Several heat-tolerant transgenic plant species have been generated already. However, a major finding of fundamental research is that the plant heat stress response is highly complex, with challenges that may be tissue, developmental stage and even species specific (Hedhly 2011; Maduraimuthu and Prasad 2014). Thus, heat tolerance should not be regarded as a single trait, and as such, it is unlikely that a general strategy can be developed to generate heat tolerance. For the near future, it will be important to evaluate to what extent the current research data obtained from model species such as *Arabidopsis* is translatable to crop species. Fortunately, genomes of most of the important crop species have been sequenced and annotated, making it possible to transfer technologically advanced methods to the crop species themselves. A remaining limitation is the space and expertise necessary to grow a crop under controlled, representative conditions and geographic location. Herein lies an opportunity for academic research groups to closely work together with breeding companies so that each can benefit from the other's expertise.

Despite widely being regarded as essential, academic research is expensive and companies are often unwilling or unable to subsidize research when no short-term payback is foreseen. At the same time, research with a high certainty of application may not be suitable or challenging for academics, many who strive to publish in higher-ranked scientific journals. A solution to this problem may lie in tripartite collaboration between academia, the private sector and governments, based on the shared aim of contributing to sustainable food production for a growing population in a warming world. Well-balanced investments may have synergistic effects on academic research output and the potential for application of findings.

In conclusion, in order to achieve success, combined efforts of plant physiologist, molecular biologists and crop breeders are required. Given the importance of global food security, the need for a versatile and linked global strategy

and multidisciplinary collaboration involving governmental agencies, companies and academics is particularly evident.

Author contribution statement N.D., W.V. and I.R. wrote the manuscript. All authors read and approved the manuscript.

Acknowledgments This work was supported by the Dutch Topsector Horticulture and Starting Materials (Grant Number 2013-H320) and the European Commission (Marie Curie Initial Training Network: Solanaceae Pollen Thermotolerance/SPOT-ITN, Grant Number 289220). We are thankful to Jacob Monash (The New Zealand Institute for Plant and Food Research, Auckland, New Zealand) for his kind help in proofreading the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abiko M, Akibayashi K, Sakata T, Kimura M, Kihara M, Itoh K, Asamizu E, Sato S, Takahashi H, Higashitani A (2005) High-temperature induction of male sterility during barley (*Hordeum vulgare* L.) anther development is mediated by transcriptional inhibition. *Sex Plant Reprod* 18:91–100. doi:10.1007/s00497-005-0004-2
- Aflitos S, Schijlen E, de Jong H, de Ridder D, Smit S, Finkers R, Wang J, Zhang G, Li N, Mao L et al (2014) Exploring genetic variation in the tomato (*Solanum section Lycopersicon*) clade by whole-genome sequencing. *Plant J* 80:136–148. doi:10.1111/tj.12616
- Ainsworth EA, Ort DR (2010) How do we improve crop production in a warming world? *Plant Physiol* 154:526–530. doi:10.1104/pp.110.161349
- Akman Z (2009) Comparison high temperature stress in maize, rice and sorghum by plant growth regulators. *J Anim Vet Adv* 8:358–361
- Almeselmani M, Deshmukh PS, Sairam RK, Kushwaha SR, Singh TP (2006) Protective role of antioxidant enzymes under high temperature stress. *Plant Sci* 171:382–388. doi:10.1016/j.plantsci.2006.04.009
- Argyris J, Dahal P, Hayashi E, Still DW, Bradford KJ (2008) Genetic variation for lettuce seed thermoinhibition is associated with temperature-sensitive expression of abscisic acid, gibberellin, and ethylene biosynthesis, metabolism, and response genes. *Plant Physiol* 148:926–947. doi:10.1104/pp.108.125807
- Argyris J, Truco MJ, Ochoa O, McHale L, Dahal P, Van Deynze A, Michelmore RW, Bradford KJ (2011) A gene encoding an abscisic acid biosynthetic enzyme (*LsNCED4*) colocalizes with the high temperature germination locus *Htg6.1* in lettuce (*Lactuca* sp.). *Theor Appl Genet* 122:95–108. doi:10.1007/s00122-010-1425-3
- Atwell BJ, Wang H, Scafaro AP (2014) Could abiotic stress tolerance in wild relatives of rice be used to improve *Oryza sativa*? *Plant Sci* 215–216:48–58. doi:10.1016/j.plantsci.2013.10.007
- Bac-Molenaar JA, Fradin EF, Becker FFM, Rienstra JA, van der Schoot J, Vreugdenhil D, Keurentjes JJB (2015) Genome-wide association mapping of fertility reduction upon heat stress reveals developmental stage-specific QTLs in *Arabidopsis thaliana*. *Plant Cell* 27:1857–1874. doi:10.1105/tpc.15.00248
- Badiani M, Schenone G, Paolacci AR, Fumagalli I (1993) Daily fluctuations of antioxidants in bean (*Phaseolus vulgaris* L.) leaves as affected by the presence of ambient air pollutants. *Plant Cell Physiol* 34:271–279
- Baker CJ, Orlandi EW (1995) Active oxygen in plant pathogenesis. *Annu Rev Phytopathol* 33:299–321. doi:10.1146/annurev.py.33.090195.001503
- Bansal KC, Lenka SK, Mondal TK (2014) Genomic resources for breeding crops with enhanced abiotic stress tolerance. *Plant Breed* 133:1–11. doi:10.1111/pbr.12117
- Barnabás B, Jäger K, Fehér A (2008) The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell Environ* 31:11–38. doi:10.1111/j.1365-3040.2007.01727.x
- Benites FRG, Pinto CABP (2011) Genetic gains for heat tolerance in potato in three cycles of recurrent selection. *Crop Breed Appl Biotechnol* 11:133–140. doi:10.1590/S1984-70332011000200005
- Bergelson J, Roux F (2010) Towards identifying genes underlying ecologically relevant traits in *Arabidopsis thaliana*. *Nat Rev Genet* 11:867–879. doi:10.1038/nrg2896
- Bhattacharjee S (2012) An inductive pulse of hydrogen peroxide pretreatment restores redox-homeostasis and oxidative membrane damage under extremes of temperature in two rice cultivars. *Plant Growth Regul* 68:395–410. doi:10.1007/s10725-012-9728-9
- Bitá CE, Gerats T (2013) Plant tolerance to high temperature in a changing environment: scientific fundamentals and production of heat stress-tolerant crops. *Front Plant Sci* 4:273. doi:10.3389/fpls.2013.00273
- Bitá CE, Zenoni S, Vriezen WH, Mariani C, Pezzotti M, Gerats T (2011) Temperature stress differentially modulates transcription in meiotic anthers of heat-tolerant and heat-sensitive tomato plants. *BMC Genom* 12:384. doi:10.1186/1471-2164-12-384
- Bokszczanin KL, Solanaceae PollenThermotolerance Initial Training Network (SPOT-ITN), C, Fragkostefanakis S (2013) Perspectives on deciphering mechanisms underlying plant heat stress response and thermotolerance. *Front Plant Sci* 4:315. doi:10.3389/fpls.2013.00315
- Boston RS, Viitanen PV, Vierling E (1996) Molecular chaperones and protein folding in plants. In: Post-transcriptional control of gene expression in plants, pp 191–222
- Buckler ES, Holland JB, Bradbury PJ, Acharya CB, Brown PJ, Browne C, Ersoz E, Flint-Garcia S, Garcia A, Glaubitz JC et al (2009) The genetic architecture of maize flowering time. *Science* 325:714–718. doi:10.1126/science.1174276
- Camejo D, Rodríguez P, AngelesMorales M, Miguel Dell'Amico J, Torrecillas A, Alarcón JJ (2005) High temperature effects on photosynthetic activity of two tomato cultivars with different heat susceptibility. *J Plant Physiol* 162:281–289. doi:10.1016/j.jplph.2004.07.014
- Challinor AJ, Watson J, Lobell DB, Howden SM, Smith DR, Chhetri N (2014) A meta-analysis of crop yield under climate change and adaptation. *Nat Clim Change* 27:1–5. doi:10.1038/NCLIMATE2153
- Chao YY, Hsu YT, Kao CH (2009) Involvement of glutathione in heat shock- and hydrogen peroxide-induced cadmium tolerance of rice (*Oryza sativa* L.) seedlings. *Plant Soil* 318:37–45. doi:10.1007/s11104-008-9815-x
- Chapman SC, Chakraborty S, Dreccer MF, Howden SM (2012) Plant adaptation to climate change—opportunities and priorities in breeding. *Crop Pasture Sci* 63:251–268. doi:10.1071/CP11303

- Chen S, Liu A, Zhang S, Li C, Chang R, Liu D, Ahammed GJ, Lin X (2013) Overexpression of mitochondrial uncoupling protein conferred resistance to heat stress and *Botrytis cinerea* infection in tomato. *Plant Physiol Biochem* 73:245–253. doi:10.1016/j.plaphy.2013.10.002
- Cheng L, Zou Y, Ding S, Zhang J, Yu X, Cao J, Lu G (2009) Polyamine accumulation in transgenic tomato enhances the tolerance to high temperature stress. *J Integr Plant Biol* 51:489–499. doi:10.1111/j.1744-7909.2009.00816.x
- Cheng NH, Liu JZ, Liu X, Wu Q, Thompson SM, Lin J, Chang J, Whitham SA, Park S, Cohen JD et al (2011) Arabidopsis monothiol glutaredoxin, AtGRXS17, is critical for temperature-dependent postembryonic growth and development via modulating auxin response. *J Biol Chem* 286:20398–20406. doi:10.1074/jbc.M110.201707
- Chou TS, Chao YY, Kao CH (2012) Involvement of hydrogen peroxide in heat shock- and cadmium-induced expression of ascorbate peroxidase and glutathione reductase in leaves of rice seedlings. *J Plant Physiol* 169:478–486. doi:10.1016/j.jplph.2011.11.012
- Collins NC, Tardieu F, Tuberosa R (2008) Quantitative trait loci and crop performance under abiotic stress: Where do we stand? *Plant Physiol* 147:469–486. doi:10.1104/pp.108.118117
- Covell S, Ellis RH, Roberts EH, Summerfield RJ (1986) The influence of temperature on seed germination rate in grain legumes: a comparison of chickpea, lentil, soybean, and cowpea at constant temperatures. *J Exp Bot* 37:705–715
- Crafts-Brandner SJ, Salvucci ME (2002) Sensitivity of photosynthesis in a C4 plant, maize, to heat stress. *Plant Physiol* 129:1773–1780. doi:10.1104/pp.002170
- Das K, Roychoudhury A (2014) Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Front Environ Sci* 2:53. doi:10.3389/fenvs.2014.00053
- Dat JF, Foyer CH, Scott IM (1998) Changes in salicylic acid and antioxidants during induced thermotolerance in mustard seedlings. *Plant Physiol* 118:1455–1461. doi:10.1104/pp.118.4.1455
- del Río LA, Sandalio LM, Corpas FJ, Palma JM, Barroso JB (2006) Reactive oxygen species and reactive nitrogen species in peroxisomes. Production, scavenging, and role in cell signaling. *Plant Physiol* 141:330–335. doi:10.1104/pp.106.078204
- Dolferus R (2014) Plant science to grow or not to grow: a stressful decision for plants. *Plant Sci* 229:247–261. doi:10.1016/j.plantsci.2014.10.002
- Dong X, Yi H, Lee J, Nou I-S, Han C-T, Hur Y (2015) Global gene-expression analysis to identify differentially expressed genes critical for the heat stress response in *Brassica rapa*. *PLoS One* 10:e0130451. doi:10.1371/journal.pone.0130451
- Duzheng YE (2003) The northward shift of climatic belts in China during the last 50 years and the corresponding seasonal responses. *Adv Atmos Sci* 20:959–967
- Easterling WE (1996) Adapting North American agriculture to climate change in review. *Agric For Meteorol* 80:1–53. doi:10.1016/0168-1923(95)02315-1
- Ebrahim MK, Zingsheim O, El-Shourbagy MN, Moore PH, Komor E (1998) Growth and sugar storage in sugarcane grown at temperatures below and above optimum. *J Plant Physiol* 153:593–602. doi:10.1016/S0176-1617(98)80209-5
- Ehlers JD, Hall AE (1998) Heat tolerance of contrasting cowpea lines in short and long days. *Field Crop Res* 55:11–21. doi:10.1016/S0378-4290(97)00055-5
- Endo M, Tsuchiya T, Hamada K, Kawamura S, Yano K, Ohshima M, Higashitani A, Watanabe M, Kawagishi-Kobayashi M (2009) High temperatures cause male sterility in rice plants with transcriptional alterations during pollen development. *Plant Cell Physiol* 50:1911–1922. doi:10.1093/pcp/pcp135
- Erickson AN, Markhart AH (2002) Flower developmental stage and organ sensitivity of bell pepper (*Capsicum annuum* L.) to elevated temperature. *Plant Cell Environ* 25:123–130. doi:10.1046/j.0016-8025.2001.00807.x
- FAO (2009) Food and Agriculture Organisation (FAO) of the United Nations, Declaration of the World Summit on Food Security, Rome, 16–18 Nov 2009. www.fao.org/wsfs/world-summit/en/
- Farnham MW, Bjorkman T (2011) Breeding vegetables adapted to high temperatures: a case study with broccoli. *HortScience* 46:1093–1097
- Feuillet C, Langridge P, Waugh R (2008) Cereal breeding takes a walk on the wild side. *Trends Genet* 24:24–32. doi:10.1016/j.tig.2007.11.001
- Fischer RA, Edmeades GO (2010) Breeding and cereal yield progress. *Crop Sci* 50:S85–S98. doi:10.2135/cropsci2009.10.0564
- Fischer G, Shah M, Van Velthuizen H (2002) Climate change and agricultural vulnerability, world summit on sustainable development. Johannesburg
- Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, Mueller ND, O’Connell C, Ray DK, West PC et al (2011) Solutions for a cultivated planet. *Nature* 478:337–342. doi:10.1038/nature10452
- Frank G, Pressman E, Ophir R, Althan L, Shaked R, Freedman M, Shen S, Firon N (2009) Transcriptional profiling of maturing tomato (*Solanum lycopersicum* L.) microspores reveals the involvement of heat shock proteins, ROS scavengers, hormones, and sugars in the heat stress response. *J Exp Bot* 60:3891–3908. doi:10.1093/jxb/erp234
- Frey FP, Urbany C, Hüttel B, Reinhardt R, Stich B (2015) Genome-wide expression profiling and phenotypic evaluation of European maize inbreds at seedling stage in response to heat stress. *BMC Genomics* 16:123. doi:10.1186/s12864-015-1282-1
- Frova C, Sari-Gorla M (1994) Quantitative trait loci (QTLs) for pollen thermotolerance detected in maize. *Mol Gen Genet* 245:424–430. <http://www.ncbi.nlm.nih.gov/pubmed/7808391>
- Füleky G (2009) Cultivated plants, primarily as food resources. In: *Encyclopedia of life support systems (EOLSS)*, vol I
- Galani S, Wahid A, Arshad M (2013) Tissue-specific expression and functional role of dehydrins in heat tolerance of sugarcane (*Saccharum officinarum*). *Protoplasma* 250:577–583. doi:10.1007/s00709-012-0443-1
- Gechev TS, Hille J (2005) Hydrogen peroxide as a signal controlling plant programmed cell death. *J Cell Biol* 168:17–20. doi:10.1083/jcb.200409170
- Giardi MT, Masojidek J, Godde D (1997) Minireview Effects of abiotic stresses on the turnover of the D1 reaction centre II protein. *Physiol Plant* 101:635–642
- Giorno F, Wolters-Arts M, Mariani C, Rieu I (2013) Ensuring reproduction at high temperatures: the heat stress response during anther and pollen development. *Plants* 2:489–506. doi:10.3390/plants2030489
- Gitay H, Brown S, Easterling W, Jallow B (2001) Ecosystems and their goods and services. In: *Climate change 2001 impacts, adapt. Vulnerability*, pp 237–342. doi:10.1387/ijdb.113386mm
- Godfray C, Beddington J, Crute I, Haddad L, Lawrence D, Muir JF, Pretty J, Robinson S, Thomas SM, Toulmin C (2010) Food security: the challenge of feeding 9 billion people. *Science* 327:812–818. doi:10.1126/science.1185383
- Gororo NN, Eagles HA, Eastwood RF, Nicolas ME, Flood RG (2002) Use of *Triticum tauschii* to improve yield of wheat in low-yielding environments. *Euphytica* 123:241–254. doi:10.1023/A:1014910000128
- Grandillo S, Tanksley SD, Zamir D (2007) Exploitation of natural biodiversity through genomics. In: *Genomics-assisted crop improvement: vol 1: genomics approaches and platforms*, pp 121–150. doi:10.1007/978-1-4020-6295-7_6

- Gross BL, Olsen KM (2010) Genetic perspectives on crop domestication. *Trends Plant Sci* 15:529–537. doi:[10.1016/j.tplants.2010.05.008](https://doi.org/10.1016/j.tplants.2010.05.008)
- Grover A, Mittal D, Negi M, Lavania D (2013) Generating high temperature tolerant transgenic plants: achievements and challenges. *Plant Sci* 205–206:38–47. doi:[10.1016/j.plantsci.2013.01.005](https://doi.org/10.1016/j.plantsci.2013.01.005)
- Gupta AS, Webb RP, Holaday AS, Allen RD (1993) Overexpression of superoxide dismutase protects plants from oxidative stress (induction of ascorbate peroxidase in superoxide dismutase-overexpressing plants). *Plant Physiol* 103:1067–1073. doi:[10.2307/4275509](https://doi.org/10.2307/4275509)
- Hall AE (2001) Heat stress and its impact, crop responses to environment. CRC Press LLC, Boca Raton
- Hansen J, Sato M, Hearty P, Ruedy R, Kelley M, Masson-Delmotte V, Russell G, Tselioudis G, Cao J, Rignot E et al (2015) Ice melt, sea level rise and superstorms: evidence from paleoclimate data, climate modeling, and modern observations that 2°C global warming is highly dangerous. *Atmos Chem Phys Discuss* 15:20059–20179. doi:[10.5194/acpd-15-20059-2015](https://doi.org/10.5194/acpd-15-20059-2015)
- Hartung F, Schiemann J (2014) Precise plant breeding using new genome editing techniques: opportunities, safety and regulation in the EU. *Plant J* 78:742–752. doi:[10.1111/tbj.12413](https://doi.org/10.1111/tbj.12413)
- Hasanuzzaman M, Nahar K, Alam MM, Roychowdhury R, Fujita M (2013) Physiological, biochemical, and molecular mechanisms of heat stress tolerance in plants. *Int J Mol Sci* 14:9643–9684. doi:[10.3390/ijms14059643](https://doi.org/10.3390/ijms14059643)
- Hayat S, Masood A, Yusuf M, Fariduddin Q, Ahmad A (2009) Foliar application of Burkholderia sp. strain TNAU-1 leads to activation of defense responses in chilli (*Capsicum annuum* L.). *Braz J Plant Physiol* 23:261–266
- Hedhly A (2011) Sensitivity of flowering plant gametophytes to temperature fluctuations. *Environ Exp Bot* 74:9–16. doi:[10.1016/j.envexpbot.2011.03.016](https://doi.org/10.1016/j.envexpbot.2011.03.016)
- Hedhly A, Hormaza JI, Herrero M (2009) Global warming and sexual plant reproduction. *Trends Plant Sci* 14:30–36. doi:[10.1016/j.tplants.2008.11.001](https://doi.org/10.1016/j.tplants.2008.11.001)
- Hossain MA, Bhattacharjee S, Armin S-M, Qian P, Xin W, Li H-Y, Burritt DJ, Fujita M, Tran L-SP (2015) Hydrogen peroxide priming modulates abiotic oxidative stress tolerance: insights from ROS detoxification and scavenging. *Front Plant Sci* 6:1–19. doi:[10.3389/fpls.2015.00420](https://doi.org/10.3389/fpls.2015.00420)
- Huang X, Han B (2014) Natural variations and genome-wide association studies in crop plants. *Annu Rev Plant Biol* 65:531–551. doi:[10.1146/annurev-arplant-050213-035715](https://doi.org/10.1146/annurev-arplant-050213-035715)
- Huang X, Zhao Y, Wei X, Li C, Wang A, Zhao Q, Li W, Guo Y, Deng L, Zhu C et al (2011) Genome-wide association study of flowering time and grain yield traits in a worldwide collection of rice germplasm. *Nat Genet* 44:32–39. doi:[10.1038/ng.1018](https://doi.org/10.1038/ng.1018)
- IPCC (2007) Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Solomon S, Qin D, Manning M, Alley RB, Berntsen T, Bindoff NL, Chen ZC. Cambridge University Press, Cambridge, UK and New York, NY, USA. http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_wg1_report_the_physical_science_basis.htm
- IPCC (2013) Summary for policymakers climate change 2013: The physical science basis Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Stocker, Qin TF, Plattner D, Tignor G-K, Allen M, Boschung SK, Naue J. Cambridge, United Kingdom and New York, NY, USA. <https://www.ipcc.ch/report/ar5/>
- Ishimaru T, Hirabayashi H, Ida M, Takai T, San-Oh YA, Yoshinaga S, Ando I, Ogawa T, Kondo M (2010) A genetic resource for early-morning flowering trait of wild rice *Oryza officinalis* to mitigate high temperature-induced spikelet sterility at anthesis. *Ann Bot* 106:515–520. doi:[10.1093/aob/mcq124](https://doi.org/10.1093/aob/mcq124)
- Jagadish SVK, Muthurajan R, Oane R, Wheeler TR, Heuer S, Bennett J, Craufurd PQ (2010) Physiological and proteomic approaches to address heat tolerance during anthesis in rice (*Oryza sativa* L.). *J Exp Bot* 61:143–156. doi:[10.1093/jxb/erp289](https://doi.org/10.1093/jxb/erp289)
- Jagadish B, Craufurd PA, Shi W, Oane AR (2014) A phenotypic marker for quantifying heat stress impact during microsporogenesis in rice (*Oryza sativa* L.). *Funct Plant Biol* 41:48–55. doi:[10.1071/FP13086](https://doi.org/10.1071/FP13086)
- Jha UC, Bohra A, Singh NP (2014) Heat stress in crop plants: its nature, impacts and integrated breeding strategies to improve heat tolerance. *Plant Breed* 133:679–701. doi:[10.1111/pbr.12217](https://doi.org/10.1111/pbr.12217)
- Jia G, Huang X, Zhi H, Zhao Y, Zhao Q, Li W, Chai Y, Yang L, Liu K, Lu H et al (2013) A haplotype map of genomic variations and genome-wide association studies of agronomic traits in foxtail millet (*Setaria italica*). *Nat Genet* 45:957–961. doi:[10.1038/ng.2673](https://doi.org/10.1038/ng.2673)
- Kang N, Kang Y, Kang K, ByoungRyong J (2009) Induction of thermotolerance and activation of antioxidant enzymes in H2O2 pre-applied leaves of cucumber and tomato seedlings. *J Jpn Soc Hortic Sci* 78:320–329
- Kenny GJ, Harrison PA, Olesen JE, Parry ML (1993) The effects of climate change on land suitability of grain maize, winter wheat and cauliflower in Europe. *Eur J Agron* 2:325–338
- Kotak S, Larkindale J, Lee U, von Koskull-Döring P, Vierling E, Scharf KD (2007) Complexity of the heat stress response in plants. *Curr Opin Plant Biol* 10:310–316. doi:[10.1016/j.pbi.2007.04.011](https://doi.org/10.1016/j.pbi.2007.04.011)
- Kugblenu YO, Oppong Danso E, Ofori K, Andersen MN, Abenney-Mickson S, Sabi E, Plauborg F, Abekoe MK, Ortiz R, Jørgensen ST (2013) Screening tomato genotypes in Ghana for adaptation to high temperature. *Acta Agric Scand Sect B Soil Plant Sci* 63:516–522. doi:[10.1080/09064710.2013.813062](https://doi.org/10.1080/09064710.2013.813062)
- Ladizinsky G (1985) Founder effect in crop-plant evolution. *Econ Bot* 39:191–199
- Larkindale J, Knight MR (2002) Protection against heat stress-induced oxidative damage in Arabidopsis involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiol* 128:682–695. doi:[10.1104/pp.010320.682](https://doi.org/10.1104/pp.010320.682)
- Levy D, Veilleux RE (2007) Adaptation of potato to high temperatures and salinity—a review. *Am J Potato Res* 84:487–506. doi:[10.1007/BF02987885](https://doi.org/10.1007/BF02987885)
- Li Y, Zhao S, Ma J, Li D, Yan L, Li J, Qi X, Guo X, Zhang L, He W et al (2013) Molecular footprints of domestication and improvement in soybean revealed by whole genome re-sequencing. *BMC Genom* 14:579. doi:[10.1186/1471-2164-14-579](https://doi.org/10.1186/1471-2164-14-579)
- Li X-M, Chao D-Y, Wu Y, Huang X, Chen K, Cui L-G, Su L, Ye W-W, Chen H, Chen H-C et al (2015) Natural alleles of a proteasome $\alpha 2$ subunit gene contribute to thermotolerance and adaptation of African rice. *Nat Genet* 47:827–833. doi:[10.1038/ng.3305](https://doi.org/10.1038/ng.3305)
- Lijavetzky D, Cabezas J, Ibáñez A, Rodríguez V, Martínez-Zapater JM (2007) High throughput SNP discovery and genotyping in grapevine (*Vitis vinifera* L.) by combining a re-sequencing approach and SNPlex technology. *BMC Genomics* 8:424. doi:[10.1186/1471-2164-8-424](https://doi.org/10.1186/1471-2164-8-424)
- Lillig CH, Berndt C, Holmgren A (2008) Glutaredoxin systems. *Biochim Biophys Acta Gen Subj* 1780:1304–1317. doi:[10.1016/j.bbagen.2008.06.003](https://doi.org/10.1016/j.bbagen.2008.06.003)
- Lippman ZB, Semel Y, Zamir D (2007) An integrated view of quantitative trait variation using tomato interspecific introgression lines. *Curr Opin Genet Dev* 17:545–552. doi:[10.1016/j.gde.2007.07.007](https://doi.org/10.1016/j.gde.2007.07.007)
- Liu G-T, Wang L, Li S, Wang J, Cramer G, Dai Z, Duan W, Xu H, Wu B, Fan P (2012) Transcriptomic analysis of grape (*Vitis*

- vinifera* L.) leaves during and after recovery from heat stress. BMC Plant Biol 12:174. doi:10.1186/1471-2229-12-174
- Lobell D, Gourdji S (2012) The influence of climate change on global crop productivity. Plant Physiol 160:1686–1697. doi:10.1104/pp.112.208298
- Lopes MS, Reynolds MP, McIntyre CL, Mathews KL, Jalal Kamali MR, Mossad M, Feltaous Y, Tahir ISA, Chatrath R, Ogbonaya F et al (2013) QTL for yield and associated traits in the Seri/Babax population grown across several environments in Mexico, in the West Asia, North Africa, and South Asia regions. Theor Appl Genet 126:971–984. doi:10.1007/s00122-012-2030-4
- Lotze-Campen H, Schellnhuber H-J (2009) Climate impacts and adaptation options in agriculture: what we know and what we don't know. J für Verbraucherschutz und Leb 4:145–150. doi:10.1007/s00003-009-0473-6
- Luo Q (2011) Temperature thresholds and crop production: a review. Clim Change 109:583–598. doi:10.1007/s10584-011-0028-6
- Lusser M, Parisi C, Plan D, Rodríguez-Cerezo E (2012) Deployment of new biotechnologies in plant breeding. Nat Biotechnol 30:231–239. doi:10.1038/nbt.2142
- Lyman NB, Jagadish KSV, Nalley LL, Dixon BL, Siebenmorgen T (2013) Neglecting rice milling yield and quality underestimates economic losses from high-temperature stress. PLoS One 8:e72157. doi:10.1371/journal.pone.0072157
- Mace ES, Tai S, Gilding EK, Li Y, Prentis PJ, Bian L, Campbell BC, Hu W, Innes DJ, Han X et al (2013) Whole-genome sequencing reveals untapped genetic potential in Africa's indigenous cereal crop sorghum. Nat Commun 4:2320. doi:10.1038/ncomms3320
- Maduraimuthu D, Prasad PVV (2014) High temperature stress. In: Jackson M, Ford-Lloyd BV, Perry ML (eds) Plant genetic resources and climate change (CABI), pp 201–220
- Maestri E, Klueva N, Perrotta C, Gulli M, Nguyen HT, Marmiroli N (2002) Molecular genetics of heat tolerance and heat shock proteins in cereals. Plant Mol Biol 48:667–681
- Mahfouz MM, Piatek A, Stewart CN (2014) Genome engineering via TALENs and CRISPR/Cas9 systems: challenges and perspectives. Plant Biotechnol J 12:1006–1014. doi:10.1111/pbi.12256
- Mangelsen E, Kilian J, Harter K, Jansson C, Wanke D, Sundberg E (2011) Transcriptome analysis of high-temperature stress in developing barley caryopses: early stress responses and effects on storage compound biosynthesis. Mol Plant 4:97–115. doi:10.1093/mp/ssq058
- Mathur S, Agrawal D, Jajoo A (2014) Photosynthesis: response to high temperature stress. J Photochem Photobiol B 137:116–126. doi:10.1016/j.jphotobiol.2014.01.010
- McCord PH, Sosinski BR, Haynes KG, Clough ME, Yencho GC (2011) QTL mapping of internal heat necrosis in tetraploid potato. Theor Appl Genet 122:129–142. doi:10.1007/s00122-010-1429-z
- McDonald GK, Paulsen GM (1997) High temperature effects on photosynthesis and water relations of grain legumes. Plant Soil 196:47–58
- Meyer RS, Purugganan MD (2013) Evolution of crop species: genetics of domestication and diversification. Nat Rev Genet 14:840–852. doi:10.1038/nrg3605
- Mickelbart MV, Hasegawa PM, Bailey-Serres J (2015) Genetic mechanisms of abiotic stress tolerance that translate to crop yield stability. Nat Rev Genet 16:237–251. doi:10.1038/nrg3901
- Miller G, Schlauch K, Tam R, Cortes D, Torres MA, Shulaev V, Dangi JL, Mittler R (2009) The plant NADPH oxidase RBOHD mediates rapid systemic signaling in response to diverse stimuli. Sci Signal 2:ra45. doi:10.1126/scisignal.2000448
- Mittal D, Madhyastha DA, Grover A (2012) Gene expression analysis in response to low and high temperature and oxidative stresses in rice: combination of stresses evokes different transcriptional changes as against stresses applied individually. Plant Sci 197:102–113. doi:10.1016/j.plantsci.2012.09.008
- Momonoki YS, Momonoki T (1993) Changes in acetylcholine-hydrolyzing activity in heat-stressed plant cultivars. Biosci Biotechnol Biochem 62:438–446. doi:10.1248/cpb.37.3229
- Morgan RW, Christman MF, Jacobson FS, Storz G, Ames BN (1986) Hydrogen peroxide-inducible proteins in *Salmonella typhimurium* overlap with heat shock and other stress proteins. Proc Natl Acad Sci USA 83:8059–8063. doi:10.1073/pnas.83.21.8059
- Morimoto RI (1998) Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. Genes Dev 12:3788–3796. doi:10.1101/gad.12.24.3788
- Morris GP, Ramu P, Deshpande SP, Hash CT, Shah T, Upadhyaya HD, Riera-Lizarazu O, Brown PJ, Acharya CB, Mitchell SE et al (2013) Population genomic and genome-wide association studies of agroclimatic traits in sorghum. Proc Natl Acad Sci USA 110:453–458. doi:10.1073/pnas.1215985110
- Mostofa MG, Yoshida N, Fujita M (2013) Spermidine pretreatment enhances heat tolerance in rice seedlings through modulating antioxidative and glyoxalase systems. Plant Growth Regul 73:31–44. doi:10.1007/s10725-013-9865-9
- Navrot N, Rouhier N, Gellhaye E, Jacquot JP (2007) Reactive oxygen species generation and antioxidant systems in plant mitochondria. Physiol Plant 129:185–195. doi:10.1111/j.1399-3054.2006.00777.x
- O'Kane D, Gill V, Boyd P, Burdon R (1996) Chilling, oxidative stress and antioxidant responses in *Arabidopsis thaliana* callus. Planta 198:371–377. doi:10.1007/BF00195186
- Olesen JE, Bindi M (2002) Consequences of climate change for European agricultural productivity, land use and policy. Eur J Agron 16:239–262. doi:10.1016/S1161-0301(02)00004-7
- Olsen KM, Wendel JF (2013) A bountiful harvest: genomic insights into crop domestication phenotypes. Annu Rev Plant Biol 64:47–70. doi:10.1146/annurev-arplant-050312-120048
- Ortiz R, Braun HJ, Crossa J, Crouch JH, Davenport G, Dixon J, Dreisigacker S, Duveiller E, He Z, Huerta J et al (2008) Wheat genetic resources enhancement by the International Maize and Wheat Improvement Center (CIMMYT). Genet Resour Crop Evol 55:1095–1140. doi:10.1007/s10722-008-9372-4
- Oshino T, Abiko M, Saito R, Ichiishi E, Endo M, Kawagishi-Kobayashi M, Higashitani A (2007) Premature progression of anther early developmental programs accompanied by comprehensive alterations in transcription during high-temperature injury in barley plants. Mol Genet Genomics 278:31–42. doi:10.1007/s00438-007-0229-x
- Paran I, Van Der Knaap E (2007) Genetic and molecular regulation of fruit and plant domestication traits in tomato and pepper. J Exp Bot 58:3841–3852. doi:10.1093/jxb/erm257
- Park SE, Marshall NA, Jakku E, Dowd AM, Howden SM, Mendham E, Fleming A (2012) Informing adaptation responses to climate change through theories of transformation. Glob Environ Change 22:115–126. doi:10.1016/j.gloenvcha.2011.10.003
- Peet MM, Willits DH, Gardner R (1997) Response of ovule development and post-pollen production processes in male-sterile tomatoes to chronic, sub-acute high temperature stress. J Exp Bot 48:101–111
- Peet MM, Sato S, Gardner RG (1998) Comparing heat stress effects on male-fertile and male-sterile tomatoes. Plant Cell Environ 21:225–231
- Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X, Centeno GS, Khush GS, Cassman KG (2004) Rice yields decline with higher night temperature from global warming. Proc Natl Acad Sci USA 101:9971–9975. doi:10.1073/pnas.0403720101

- Pinto RS, Reynolds MP (2015) Common genetic basis for canopy temperature depression under heat and drought stress associated with optimized root distribution in bread wheat. *Theor Appl Genet* 128:575–585. doi:10.1007/s00122-015-2453-9
- Pinto RS, Reynolds MP, Mathews KL, McIntyre CL, Olivares-Villegas J-J, Chapman SC (2010) Heat and drought adaptive QTL in a wheat population designed to minimize confounding agronomic effects. *Theor Appl Genet* 121:1001–1021. doi:10.1007/s00122-010-1351-4
- Podlich DW, Winkler CR, Cooper M (2004) Mapping as you go. An effective approach for marker assisted selection of complex traits. *Crop Sci* 44:1560–1571
- Porter JR, Gawith M (1999) Temperatures and the growth and development of wheat: a review. *Eur J Agron* 10:23–36. doi:10.1016/S1161-0301(98)00047-1
- Pradhan GP, Prasad PVV, Fritz AK, Kirkham MB, Gill BS (2012) High temperature tolerance in *Aegilops* species and its potential transfer to wheat. *Crop Sci* 52:292–304. doi:10.2135/cropsci2011.04.0186
- Prasad PVV, Boote KJ, Allen LH, Sheehy JE, Thomas JMG (2006) Species, ecotype and cultivar differences in spikelet fertility and harvest index of rice in response to high temperature stress. *Field Crop Res* 95:398–411. doi:10.1016/j.fcr.2005.04.008
- Prasanna BM (2012) Diversity in global maize germplasm: characterization and utilization. *J Biosci* 37:843–855. doi:10.1007/s12038-012-9227-1
- Qi J, Liu X, Shen D, Miao H, Xie B, Li X, Zeng P, Wang S, Shang Y, Gu X et al (2013) A genomic variation map provides insights into the genetic basis of cucumber domestication and diversity. *Nat Genet* 45:1510–1515. doi:10.1038/ng.2801
- Reilly JM, Tubiello FN, McCarl BA, Abler D, Darwin R, Fuglie K, Hollinger S, Izaurrealde RC, Jagtap S, Jones J et al (2003) U.S. agriculture and climate change: new results. *Clim Change* 57:43–67. doi:10.1023/A:1022103315424
- Rodrigo J, Herrero M (2002) Effects of pre-blossom temperatures on flower development and fruit set in apricot. *Sci Hortic (Amsterdam)* 92:125–135. doi:10.1016/S0304-4238(01)00289-8
- Rosenzweig C, Tubiello FN (2007) Adaptation and mitigation strategies in agriculture: an analysis of potential synergies. *Mitig Adapt Strateg Glob Change* 12:855–873. doi:10.1007/s11027-007-9103-8
- Rui R, Nie Y, Tong H (1990) SOD activity as a parameter for screening stress-tolerant germplasm resources in sweet potato (*Ipomoea batatas* L.). *J Agric Sci* 6:52–56
- Sacks WJ, Deryng D, Foley JA, Ramankutty N (2010) Crop planting dates: an analysis of global patterns. *Glob Ecol Biogeogr* 19:607–620. doi:10.1111/j.1466-8238.2010.00551.x
- Saha SR, Hossain MM, Rahman MM, Kuo CG, Abdullah S (2010) Effect of high temperature stress on the performance of twelve sweet pepper genotypes. *Bangladesh J Agric Res* 35:525–534
- Sairam RK, Srivastava GC, Saxena DC (2000) Increased antioxidant activity under elevated temperatures: a mechanism of heat stress tolerance in wheat genotypes. *Biol Plant* 43:245–251
- Sakai H, Ikawa H, Tanaka T, Numa H, Minami H, Fujisawa M, Shibata M, Kurita K, Kikuta A, Hamada M et al (2011) Distinct evolutionary patterns of *Oryza glaberrima* deciphered by genome sequencing and comparative analysis. *Plant J* 66:796–805. doi:10.1111/j.1365-3113X.2011.04539.x
- Sakata T, Higashitani A (2008) Male sterility accompanied with abnormal anther development in plants—genes and environmental stresses with special reference to high temperature injury. *Int J Plant Dev Biol* 2:42–51. [http://www.globalsciencebooks.info/JournalsSup/images/0806/IJPDB_2\(1\)42-51o.pdf](http://www.globalsciencebooks.info/JournalsSup/images/0806/IJPDB_2(1)42-51o.pdf)
- Sander JD, Joung JK (2014) CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat Biotechnol* 32:347–355. doi:10.1038/nbt.2842
- Sarkar NK, Kim YK, Grover A (2014) Coexpression network analysis associated with call of rice seedlings for encountering heat stress. *Plant Mol Biol* 84:125–143. doi:10.1007/s11103-013-0123-3
- Sato S, Kamiyama M, Iwata T, Makita N, Furukawa H, Ikeda H (2006) Moderate increase of mean daily temperature adversely affects fruit set of *Lycopersicon esculentum* by disrupting specific physiological processes in male reproductive development. *Ann Bot* 97:731–738. doi:10.1093/aob/mcl037
- Sharkey TD, Schrader SM (2006) High temperature stress. *Physiology and molecular biology of stress tolerance*. Springer, Dordrecht, pp 101–130
- Smillie RM, Nott R (1979) Heat injury in leaves of alpine, temperate and tropical plants. *Funct Plant Biol* 6:135–141. <http://www.publish.csiro.au/?paper=PP9790135>
- Smit B, Skinner MW (2002) Adaptation options in agriculture to climate change: a typology. *Mitig Adapt Strateg Glob Change* 7:85–114. doi:10.1023/A:1015862228270
- Smithers J, Blay-Palmer A (2001) Technology innovation as a strategy for climate adaptation in agriculture. *Appl Geogr* 21:175–197. doi:10.1016/S0143-6228(01)00004-2
- Stevenson WR, Loria R, Franc GC, Weingartner DP (2001) Physiological disorders of tubers: internal symptoms. In: *Compendium of potato diseases*. American Phytopathological Society, St. Paul
- Stone P (2001) The effects of heat stress on cereal yield and quality. In: Basra AS (ed) *Crop responses and adaptations to temperature stress*. Food Products Press, Binghamton, pp 243–291
- Sundaram S, Rathinasabapathi B (2010) Transgenic expression of fern *Pteris vittata* glutaredoxin PvGrx5 in *Arabidopsis thaliana* increases plant tolerance to high temperature stress and reduces oxidative damage to proteins. *Planta* 231:361–369. doi:10.1007/s00425-009-1055-7
- Suzuki N, Miller G, Sejima H, Harper J, Mittler R (2013) Enhanced seed production under prolonged heat stress conditions in *Arabidopsis thaliana* plants deficient in cytosolic ascorbate peroxidase 2. *J Exp Bot* 64:253–263
- Takeoka Y, Hiroi K, Kitano H, Wada T (1991) Pistil hyperplasia in rice spikelets as affected by heat stress. *Sex Plant Reprod* 4:39–43. doi:10.1007/BF00194570
- Tanksley SD, McCouch SR (1997) Seed banks and molecular maps: unlocking genetic potential from the wild. *Science* 277:1063–1066. doi:10.1126/science.277.5329.1063
- Tchebakova NM, Parfenova EI, Lysanova GI, Soja AJ (2011) Agroclimatic potential across central Siberia in an altered twenty-first century. *Environ Res Lett* 6:045207. doi:10.1088/1748-9326/6/4/045207
- Tebaldi C, Hayhoe K, Arblaster JM, Meehl GA (2006) Going to the extremes: an intercomparison of model-simulated historical and future changes in extreme events. *Clim Change* 79:185–211. doi:10.1007/s10584-006-9051-4
- Teixeira EI, Fischer G, Van Velthuisen H, Walter C, Ewert F (2013) Global hot-spots of heat stress on agricultural crops due to climate change. *Agric For Meteorol* 170:206–215. doi:10.1016/j.agrformet.2011.09.002
- Tester M, Langridge P (2010) Breeding technologies to increase crop production in a changing world. *Science* 327:818–822. doi:10.1126/science.1183700
- Tonsor SJ, Scott C, Boumaza I, Liss TR, Brodsky JL, Vierling E (2008) Heat shock protein 101 effects in *A. thaliana*: genetic variation, fitness and pleiotropy in controlled temperature conditions. *Mol Ecol* 17:1614–1626. doi:10.1111/j.1365-294X.2008.03690.x
- Torres MA, Dangl JL (2005) Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development.

- Curr Opin Plant Biol 8:397–403. doi:[10.1016/j.pbi.2005.05.014](https://doi.org/10.1016/j.pbi.2005.05.014)
- Tubiello FN, Rosenzweig C, Goldberg RA, Jagtap S, Jones JW (2002) Effects of climate change on US crop production: simulation results using two different GCM scenarios. Part I: wheat, potato, maize, and citrus. *Clim Res* 20:259–270. doi:[10.3354/cr020259](https://doi.org/10.3354/cr020259)
- Ulukan H (2008) Agronomic adaptation of some field crops: a general approach. *J Agron Crop Sci* 194:169–179. doi:[10.1111/j.1439-037X.2008.00306.x](https://doi.org/10.1111/j.1439-037X.2008.00306.x)
- Vacca RA, De Pinto MC, Valenti D, Passarella S, Marra E, Nazionale CAVA, Vegetale P, Orabona V, Bari I et al (2004) Production of reactive oxygen species, alteration of cytosolic ascorbate peroxidase, and impairment of mitochondrial metabolism are early events in heat shock-induced programmed cell death in tobacco bright-yellow 2 cells 1. *Plant Physiol* 134:1100–1112. doi:[10.1104/pp.103.035956.persensitive](https://doi.org/10.1104/pp.103.035956.persensitive)
- Ventura G, Grilli G, Braz LT, Gertrudes E, Lemos M (2007) QTL identification for tolerance to fruit set in tomato by fAFLP markers. *Crop Breed Appl Biotechnol* 7:234–241
- Vierling E (1991) The roles of heat shock proteins in plants. *Annu Rev Plant Physiol Plant Mol Biol* 42:579–620. doi:[10.1146/annurev.pp.42.060191.003051](https://doi.org/10.1146/annurev.pp.42.060191.003051)
- Vijayalakshmi K, Fritz AK, Paulsen GM, Bai G, Pandravada S, Gill BS (2010) Modeling and mapping QTL for senescence-related traits in winter wheat under high temperature. *Mol Breed* 26:163–175. doi:[10.1007/s11032-009-9366-8](https://doi.org/10.1007/s11032-009-9366-8)
- Volkov RA, Panchuk II, Mullineaux PM, Schöffl F (2006) Heat stress-induced H₂O₂ is required for effective expression of heat shock genes in Arabidopsis. *Plant Mol Biol* 61:733–746. doi:[10.1007/s11103-006-0045-4](https://doi.org/10.1007/s11103-006-0045-4)
- Wahid A, Close TJ (2007) Expression of dehydrins under heat stress and their relationship with water relations of sugarcane leaves. *Biol Plant* 51:104–109. doi:[10.1007/s10535-007-0021-0](https://doi.org/10.1007/s10535-007-0021-0)
- Wahid A, Gelani S, Ashraf M, Foolad M (2007) Heat tolerance in plants: an overview. *Environ Exp Bot* 61:199–223. doi:[10.1016/j.envexpbot.2007.05.011](https://doi.org/10.1016/j.envexpbot.2007.05.011)
- Wang W, Vinocur B, Shoseyov O, Altman A (2004) Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. *Trends Plant Sci* 9:244–252. doi:[10.1016/j.tplants.2004.03.006](https://doi.org/10.1016/j.tplants.2004.03.006)
- Wang L, Guo Y, Jia L, Chu H, Zhou S, Chen K, Wu D, Zhao L (2014a) Hydrogen peroxide acts upstream of nitric oxide in the heat shock pathway in arabidopsis seedlings. *Plant Physiol* 164:2184–2196. doi:[10.1104/pp.113.229369](https://doi.org/10.1104/pp.113.229369)
- Wang Y, Zhang J, Li J-L, Ma X-R (2014b) Exogenous hydrogen peroxide enhanced the thermotolerance of *Festuca arundinacea* and *Lolium perenne* by increasing the antioxidative capacity. *Acta Physiol Plant* 36:2915–2924. doi:[10.1007/s11738-014-1661-2](https://doi.org/10.1007/s11738-014-1661-2)
- Wu Q, Lin J, Liu J-Z, Wang X, Lim W, Oh M, Park J, Rajashekar CB, Whitham SA, Cheng N-H et al (2012) Ectopic expression of Arabidopsis glutaredoxin AtGRXS17 enhances thermotolerance in tomato. *Plant Biotechnol J* 10:945–955. doi:[10.1111/j.1467-7652.2012.00723.x](https://doi.org/10.1111/j.1467-7652.2012.00723.x)
- Xiao Y, Pan Y, Luo L, Zhang G, Deng H, Dai L, Liu X, Tang W, Chen L, Wang G-L (2010) Quantitative trait loci associated with seed set under high temperature stress at the flowering stage in rice (*Oryza sativa* L.). *Euphytica* 178:331–338. doi:[10.1007/s10681-010-0300-2](https://doi.org/10.1007/s10681-010-0300-2)
- Xu J, Zhao Q, Du P, Xu C, Wang B, Feng Q, Liu Q, Tang S, Gu M, Han B et al (2010) Developing high throughput genotyped chromosome segment substitution lines based on population whole-genome re-sequencing in rice (*Oryza sativa* L.). *BMC Genom* 11:656. doi:[10.1186/1471-2164-11-656](https://doi.org/10.1186/1471-2164-11-656)
- Yamamoto K, Sakamoto H, Momonoki YS (2011) Maize acetylcholinesterase is a positive regulator of heat tolerance in plants. *J Plant Physiol* 168:1987–1992. doi:[10.1016/j.jplph.2011.06.001](https://doi.org/10.1016/j.jplph.2011.06.001)
- Yan K, Chen N, Qu YY, Dong XC, Meng QW, Zhao SJ (2008) Overexpression of sweet pepper glycerol-3-phosphate acyltransferase gene enhanced thermotolerance of photosynthetic apparatus in transgenic tobacco. *J Integr Plant Biol* 50:613–621. doi:[10.1111/j.1744-7909.2008.00652.x](https://doi.org/10.1111/j.1744-7909.2008.00652.x)
- Ye C, Argayoso MA, Redoña ED, Sierra SN, Laza MA, Dilla CJ, Mo Y, Thomson MJ, Chin J, Delaviña CB et al (2012) Mapping QTL for heat tolerance at flowering stage in rice using SNP markers. *Plant Breed* 131:33–41. doi:[10.1111/j.1439-0523.2011.01924.x](https://doi.org/10.1111/j.1439-0523.2011.01924.x)
- Ye C, Tenorio FA, Argayoso MA, Laza MA, Koh H-J, Redoña ED, Jagadish KS, Gregorio GB (2015) Identifying and confirming quantitative trait loci associated with heat tolerance at flowering stage in different rice populations. *BMC Genet* 16:41. doi:[10.1186/s12863-015-0199-7](https://doi.org/10.1186/s12863-015-0199-7)
- Yoshida S, Satake T, Mackill DS (1981) High-temperature stress in rice. *IRRI Research Paper Series SB206 A32*, 1–26
- Zhang D, Yang L (2014) Specification of tapetum and microsporocyte cells within the anther. *Curr Opin Plant Biol* 17:49–55. doi:[10.1016/j.pbi.2013.11.001](https://doi.org/10.1016/j.pbi.2013.11.001)
- Zinn KE, Tunc-Ozdemir M, Harper JF (2010) Temperature stress and plant sexual reproduction: uncovering the weakest links. *J Exp Bot* 61:1959–1968. doi:[10.1093/jxb/erq053](https://doi.org/10.1093/jxb/erq053)