

UC Berkeley

UC Berkeley Previously Published Works

Title

Bright, Months-long stellar outbursts announce the explosion of interaction-powered supernovae

Permalink

<https://escholarship.org/uc/item/0mw4x6xc>

Journal

Astrophysical Journal, 907(2)

ISSN

0004-637X

Authors

Strotjohann, NL
Ofek, EO
Gal-Yam, A
[et al.](#)

Publication Date

2021-02-01

DOI

10.3847/1538-4357/abd032

Peer reviewed

Bright, months-long stellar outbursts announce the explosion of interaction-powered supernovae

NORA L. STROTJOHANN,¹ ERAN O. OFEK,¹ AVISHAY GAL-YAM,¹ RACHEL BRUCH,¹ STEVE SCHULZE,¹ NIR SHAVIV,² JESPER SOLLERMAN,³
ALEXEI V. FILIPPENKO,^{4,5} OFER YARON,¹ CHRISTOFFER FREMLING,⁶ JAKOB NORDIN,⁷ ERIK C. KOOL,³ DAN A. PERLEY,⁸ ANNA Y. Q. HO,^{6,4,9} YI YANG,¹
YUHAN YAO,⁶ MAAYANE T. SOUMAGNAC,⁴ MELISSA L. GRAHAM,¹⁰ CRISTINA BARBARINO,³ LEONARDO TARTAGLIA,^{3,11} KISHALAY DE,⁶
DANIEL A. GOLDSTEIN,⁶ DAVID O. COOK,¹² THOMAS G. BRINK,⁴ KIRSTY TAGGART,⁸ LIN YAN,¹³ RAGNHILD LUNNAN,³ MANSI KASLIWAL,⁶
SHRI R. KULKARNI,⁶ PETER E. NUGENT,^{4,14} FRANK J. MASCI,¹² PHILIPPE ROSNET,¹⁵ SCOTT M. ADAMS,⁶ IGOR ANDREONI,⁶ ASHOT BAGDASARYAN,⁶
ERIC C. BELLM,¹⁰ KEVIN BURDGE,⁶ DMITRY A. DUEV,⁶ ALISON DUGAS,^{6,16} SARA FREDERICK,¹⁷ SAMANTHA GOLDWASSER,¹ MATTHEW HANKINS,⁶
IDO IRANI,¹ VIRAJ KARAMBELKAR,⁶ THOMAS KUPFER,¹⁸ JINGYI LIANG,¹ JAMES D. NEILL,⁶ MICHAEL PORTER,¹³ REED L. RIDDLE,¹⁹ YASHVI SHARMA,⁶
PHIL SHORT,²⁰ FRANCESCO TADDIA,³ ANASTASIOS TZANIDAKIS,⁶ JAN VAN ROESTEL,⁶ RICHARD WALTERS,⁶ AND ZHUYUN ZHUANG⁶

¹*Benozio Center for Astrophysics, The Weizmann Institute of Science, Rehovot 76100, Israel*

²*The Racah Institute of Physics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel*

³*Department of Astronomy, The Oskar Klein Centre, Stockholm University, AlbaNova, 10691 Stockholm, Sweden*

⁴*Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA*

⁵*Miller Senior Fellow, Miller Institute for Basic Research in Science, University of California, Berkeley, CA 94720, USA*

⁶*Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA*

⁷*Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489, Berlin, Germany*

⁸*Astrophysics Research Institute, Liverpool John Moores University, Liverpool Science Park, 146 Brownlow Hill, Liverpool L35RF, UK*

⁹*Miller Institute for Basic Research in Science, University of California, Berkeley, CA 94720, USA*

¹⁰*DiRAC Institute, Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA*

¹¹*INAF - Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy*

¹²*IPAC, California Institute of Technology, 1200 E. California Blvd, Pasadena, CA 91125, USA*

¹³*Caltech Optical Observatories, California Institute of Technology, Pasadena, CA 91125, USA*

¹⁴*Computational Science Department, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, MS 50B-4206, Berkeley, CA 94720, USA*

¹⁵*Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*

¹⁶*Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA*

¹⁷*Department of Astronomy, University of Maryland, College Park, MD 20742, USA*

¹⁸*Texas Tech University, Department of Physics & Astronomy, Box 41051, 79409, Lubbock, TX, USA*

¹⁹*Caltech Optical Observatories, California Institute of Technology, 1200 E. California Blvd, Pasadena, CA 91125, USA*

²⁰*Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK*

(Received 2020 October 27; Revised 2020 December 1; Accepted 2020 December 1)

ABSTRACT

Interaction-powered supernovae (SNe) explode within an optically-thick circumstellar medium (CSM) that could be ejected during eruptive events. To identify and characterize such pre-explosion outbursts we produce forced-photometry light curves for 196 interacting SNe, mostly of Type IIn, detected by the Zwicky Transient Facility between early 2018 and June 2020. Extensive tests demonstrate that we only expect a few false detections among the 70,000 analyzed pre-explosion images after applying quality cuts and bias corrections. We detect precursor eruptions prior to 18 Type IIn SNe and prior to the Type Ibn SN 2019uo. Precursors become brighter and more frequent in the last months before the SN and month-long outbursts brighter than magnitude -13 occur prior to 25% (5–69%, 95% confidence range) of all Type IIn SNe within the final three months before the explosion. With radiative energies of up to 10^{49} erg, precursors could eject $\sim 1 M_{\odot}$ of material. Nevertheless, SNe with detected precursors are not significantly more luminous than other SNe IIn and the characteristic narrow hydrogen lines in their spectra typically originate from earlier, undetected mass-loss events. The long precursor durations require ongoing energy injection and they could, for example, be powered by interaction or by a continuum-driven wind. Instabilities during the neon and oxygen burning phases are predicted to launch

precursors in the final years to months before the explosion; however, the brightest precursor is 100 times more energetic than anticipated.

Keywords: Eruptive phenomena (475) – Stellar mass loss (1613) – Circumstellar matter (241) – Late stellar evolution (911) – Stellar flares (1603) – Core-collapse supernovae (304)

1. INTRODUCTION

Despite the detection of more than 2000 core-collapse supernovae (SNe) per year, the processes leading to their explosions are still not entirely understood (see, e.g., Janka et al. 2016; Müller 2016; Glas et al. 2019) and remain unobservable as they happen deep within the cores of stars in distant galaxies. However, at least for some progenitor stars, the impending core collapse seems to have direct implications for the stellar envelope. Bright optical flares have been observed in the years leading up to the SN explosion and may offer another means of probing the conditions near the surface of progenitor stars, which are, with exception of the nearest events, too faint to be detected by any telescope.

The first pre-explosion outburst was detected two years prior to the explosion of the Type Ibn SN 2006jc (Pastorello et al. 2007; Foley et al. 2007). Most precursors were observed prior to Type IIn SNe (see, e.g., Ofek et al. 2013b; Mauerhan et al. 2013; Fraser et al. 2013; Margutti et al. 2013; Tartaglia et al. 2016; Elias-Rosa et al. 2016; Ofek et al. 2016; Thöne et al. 2017; Nyholm et al. 2017; Pastorello et al. 2018; Reguitti et al. 2019), as well as prior to a broad-lined Type Ic SN (Ho et al. 2019), and possibly a SN Iib (Strotjohann et al. 2015). This suggests that numerous types of progenitor stars can produce such flares. However, since most flares have been detected prior to Type IIn SNe, the progenitors of these relatively rare explosions are either more likely to generate such flares, or the generated flares are brighter. A systematic study by Ofek et al. (2014a) showed that precursor eruptions prior to Type IIn SNe are the rule rather than the exception. In a similar search, Bilinski et al. (2015) did not find any precursors and claim that the rate is lower, but they relied on a small SN sample.

Type IIn SNe are characterized by relatively narrow hydrogen emission lines (see, e.g., Filippenko 1997; Gal-Yam 2017; Smith 2017), which indicate the presence of a slowly moving circumstellar medium (CSM) surrounding the SN ejecta. This material originates from the star itself and is expelled in the years to decades before the explosion either during precursor eruptions or by a stellar wind. The SN ejecta crash into the CSM and a fraction of the ejecta kinetic energy is converted to high-energy photons (see, e.g., Katz et al. 2011; Murase et al. 2011, 2014). If the CSM is optically thick to gamma-ray and X-ray photons, a part of the radiation may be converted to the UV-optical regime. Some Type IIn SNe therefore reach much brighter optical peak magnitudes than noninteracting SNe (Kiewe et al. 2012; Stritzinger et al.

2012; Gal-Yam 2019) and their diverse light-curve shapes can be explained by different CSM geometries, which might consist of several shells (see, e.g., Margutti et al. 2013; Nyholm et al. 2017) or be aspherical (Patat et al. 2011; Soumagnac et al. 2019, 2020), as is commonly observed for planetary nebulae in our Galaxy.

The progenitor stars of a few nearby Type IIn SNe were identified in archival images and are consistent with being luminous blue variables (LBVs; see, e.g., Gal-Yam et al. 2007; Gal-Yam & Leonard 2009; Foley et al. 2011; Kochanek & Szczygiel 2011). These bright and massive stars are named after their hot surface temperatures and their high-amplitude luminosity variability. They launch strong winds, which remove part of their hydrogen envelope. They were therefore traditionally considered stars in a transitional phase which evolve from a main-sequence star into a hydrogen-stripped Wolf-Rayet star (Humphreys & Davidson 1994). Another possibility is that they develop from a main-sequence star that gains mass and angular momentum from a binary companion, which turns it into an LBV rather than a red supergiant (Smith 2017 see also Justham et al. 2014). However, there is also a class of lower-luminosity transients with Type-IIn-like spectra that originate from heavily obscured stars with masses of $\sim 10 M_{\odot}$ (see e.g. Prieto et al. 2008; Kochanek 2011; Szczygiel et al. 2012). It is hence possible that not all Type IIn SNe originate from massive LBVs.

In addition to Type IIn SNe, evidence for interaction has also been observed for several other SN classes. Type Ibn SNe explode within a helium-rich CSM and their rapid light-curve evolution might indicate that the CSM is confined to a small radius (see, e.g., Pastorello et al. 2016; Gal-Yam 2017; Hosseinzadeh et al. 2017). The spectra of Type II superluminous supernovae (SLSNe-II) often look similar to the ones of Type IIn SNe and their large radiative energy is usually attributed to strong CSM interaction (see, e.g., Gal-Yam 2019). So called flash-spectroscopy SNe exhibit narrow emission features during for the first few days after their explosion; these could originate from a confined CSM shell that is flash ionized by radiation from the shock breakout and is the quickly swept up by the expanding ejecta (see, e.g., Gal-Yam et al. 2014; Khazov et al. 2016; Yaron et al. 2017; Smith 2017; Bruch et al. 2020). Type Ia-CSM SNe are thermonuclear explosions of white dwarfs that explode inside a hydrogen-rich CSM, potentially produced by a binary companion star (Hamuy et al. 2003; Dilday et al. 2012; Silverman et al. 2013; Gal-Yam 2017). In the following, we use the ex-

pression “interaction-powered SNe” to refer to all these subclasses.

A first systematic search for precursor eruptions was done by Ofek et al. (2014a) for a sample of 16 nearby Type II SNe using data from the Palomar Transient Factory (PTF; Law et al. 2009; Rau et al. 2009). It established that most Type II progenitor stars undergo one or several precursor eruptions in the last 2.5 yr before the SN and that the rate increases in the last 4 months before the explosion. However, the study was limited by the small SN sample and by the relatively sparse sampling of the pre-explosion light curves. The majority of the observations were obtained in the Mould-*R* band, such that the precursor colors could not be determined.

Here, we build on the work by Ofek et al. (2014a) and use data from the Zwicky Transient Facility (ZTF; Bellm et al. 2019; Graham et al. 2019) to systematically search for precursor eruptions prior to interacting SNe, mostly of Type II SNe. Compared to PTF, ZTF has a ~ 15 times faster survey speed: with its large field of view of 47 deg^2 , it monitors nearly the complete sky at declinations larger than -30° and smaller than 80° (Bellm et al. 2019). Since the commissioning of the ZTF camera in fall 2017, the survey has detected more than 200 interacting SNe for which nearly 10^5 pre-explosion images are available in the *g*, *r*, and *i* bands. We here search unbinned and binned light curves for pre-explosion activity. Owing to the abundant photometric data provided by the ZTF survey and the larger SN sample, we expect to detect more precursor eruptions and measure the precursor rate more precisely. Thus, we extend the previous search to fainter, shorter, and less-common precursors and expect that the eruptions are better observed with data in multiple bands.

The paper is structured as follows. Section 2 describes the analysis and quality cuts which allow us to reduce the rate of false-positive detections. The detected precursors are described in Sec. 3 and the luminosity-dependent precursor rates are measured in Sec. 4. In Sec. 5, we show that the material ejected during most of the detected precursors cannot account for the characteristic narrow hydrogen lines in the spectra of Type II SNe. One exception is the Type Ib SN 2019uo, described Sec. 5.3, for which the observed interaction can be explained by the precursor 320 days before the explosion. In Sec. 6 we consider which mechanisms might power the precursor luminosity and whether wave-driven mass loss could launch the observed precursors. Our findings are summarized in Sec. 7.

2. METHODS

The following subsections introduce the sample selection (Sec. 2.1), the forced photometry pipeline (Sec. 2.2), and the tests we perform on the pipeline (Sec. 2.3). Next, we explain how images with astrometric errors are rejected (Sec. 2.4)

and how we correct the baseline offsets and rescale underestimated error bars (Sec. 2.5). Finally, we describe how observations are combined in bins (Sec. 2.6) and estimate the expected number of false detections in Sec. 2.7.

2.1. Sample Selection

The ZTF survey produces about 1 million alerts per night (Patterson et al. 2019) which are then scored by a deep-learning algorithm to identify genuine astrophysical transients (Duv et al. 2019). The resulting alert stream is filtered either by the AMPEL broker (Nordin et al. 2019; Soumagnac & Ofek 2018) or the GROWTH “Marshal” (Kasliwal et al. 2019) based on different science goals, such as the detection of young SNe (Gal-Yam 2019; Bruch et al. 2020) or bright transients (Fremling et al. 2020). In most science programs potentially interesting objects are identified by astronomers who request spectroscopy or other follow-up observations. Transients brighter than magnitude ~ 19 are usually first classified based on spectra from the SED Machine (Ben-Ami et al. 2012; Blagorodnova et al. 2018; Rigault et al. 2019) and higher-resolution spectra might be obtained later.

The commissioning phase of the ZTF survey started in fall 2017, while the survey officially began in spring 2018 after commissioning and building reference images. To select a sample of interaction-powered SNe with ZTF pre-explosion observations, we query both the Transient Name Server (TNS¹) and the private ZTF database, the GROWTH Marshal using the ZTFquery code (Rigault 2018), for transients discovered since 2018 January 1 and until 2020 June 24. We only consider SNe at locations that are observable by ZTF, with declinations larger than -30° . Our sample includes all objects that are classified as SNe of Type II, Ib, Ia-CSM, or SLSNe-II by members of the ZTF team or on TNS (see, e.g., Perley et al. 2020 for details). In addition, we include objects that show flash-spectroscopy features in early-time spectra, which were identified by Bruch et al. (2020). This brings the total sample to 239 SNe.

An accurate localization is required to perform forced photometry (see, e.g., Yao et al. 2019), and we therefore only consider objects with at least ten ZTF detections. We find that this ensures that the position is within $0.15''$ of the best position for 90% of the SNe in the ZTF coordinate system². Out of 239 SNe, 12 objects have fewer than ten ZTF detections and are discarded. The remaining 227 SNe are listed in the online version of Table 1.

To confirm both the SN classification and the redshift, we visually inspect spectra from the ZTF Marshal as well as the TNS. We discard in total 18 objects which we cannot verify

¹ <https://wis-tns.weizmann.ac.il/>

² A precision of $\lesssim 0.15''$ is the required threshold for forced photometry (Frank Masci, priv. comm.).

Table 1. SNe with detected pre-explosion activity

IAU name	ZTF name	SN Type	R.A. (J2000)	Dec. (J2000)	z	t_0	Separation	Comment
			(deg)	(deg)		(JD)	(arcsec)	
SN 2018eru	ZTF 18ablqehq	IIn	185.115828	41.79289029	0.03069	2458316.6	11	
SN 2018gho	ZTF 18abucxcj	IIn	246.8412533	39.1091986	0.033	2458366.4	2.8	
SN 2018hxe	ZTF 18abwlupf	IIn	221.0426466	62.89518	0.134	2458370.7	0.37	
SN 2018kag	ZTF 18acwzyor	IIn	133.951979	3.64152020	0.02736	2458466.5	6.3	
SN 2019uo	ZTF 19aadnxbh	Ibn	180.6525136	41.0616364	0.020454	2458501.2	27	
SN 2019bxq	ZTF 19aamkmxv	IIn	254.4938	78.6037	0.0139	2458555.8	0.01	
SN 2019cmy	ZTF 19aanpcep	IIn	227.2118369	40.7137261	0.0314	2458567.9	5.8	
SN 2019iay	ZTF 19abandzh	IIn	200.27061809	8.1684897	0.0406	2458656.6	7.9	
SN 2019meh	ZTF 19abclykm	SLSN-II	321.8227253	64.4164373	0.0935	2458657.3	0.04	bg AGN ^a
SN 2019gjs	ZTF 19abiszoe	IIn	224.7381972	20.0529308	0.043	2458690.7	10	
SN 2019mom	ZTF 19ablojrw	IIn	28.9021955	53.5918978	0.0488	2458690.9	–	
SN 2019njv	ZTF 19abpidqn	IIn	304.98829689	15.37745280	0.01458	2458706.9	2.9	
SN 2019fmb	ZTF 19aavyvbn	IIn	186.68196	56.0757834	0.016	2458715.8	17	t_0 uncertain
SN 2019sae	ZTF 19acahbxd	IIn	41.2693283	26.0714348	0.048	2458728.8	9.9	
SN 2019aafe	ZTF 19abzfxel	IIn	349.0686877	48.4284178	0.075	2458740.8	–	
SN 2019vkl	ZTF 19acukucu	IIn	29.1283206	18.4399406	0.064	2458808.6	–	
SN 2019vts	ZTF 19acxmnkc	IIn	98.7642676	50.434783	0.0395	2458816.9	24	
SN 2019qny	ZTF 19adannbl	IIn	53.24978869	–2.778198	0.048	2458827.0	5.7	
SN 2020iq	ZTF 20aabcemq	IIn	43.8321616	–11.4134991	0.096	2458832.6	2.5	
SN 2019yzx	ZTF 19adcxbkw	Ia-CSM	142.6721432	21.4558832	0.057	2458840.0	2.0	
SN 2019zrk	ZTF 20aacbyec	IIn	174.9475073	19.9296524	0.0362	2458889.0	14	
SN 2020dcs	ZTF 20aacqkr	IIn	183.3561586	37.6993902	0.023958	2458894.9	3.0	
SN 2020dfh	ZTF 20aasivpe	IIn	265.506694	3.2008709	0.0293	2458903.1	6.3	
SN 2020edh	ZTF 20aaswzdm	IIn	259.0999066	40.8081331	0.033	2458914.9	3.4	

NOTE—The R.A. and Dec. values represent the median coordinates of at least 10 ZTF detections. The discovery time t_0 is either the first detection time announced on TNS or a smaller value if the transient flux is visible earlier in ZTF data. The penultimate column lists the separation from the center of the host galaxy, to judge whether active galactic nucleus (AGN) activity might contribute to the pre-explosion variability. Here, we only list SNe for which pre-explosion activity is detected (see Sec. 3). The full table, containing all 227 considered SNe described in Sec. 2.1, is available online.

^aThe detected variability likely originates from AGN activity in the center of the host galaxy and not from the progenitor star (see Sec. 2.7 and 3.1 for details).

are interacting transients. For most of these objects no good spectra are available or the observed narrow lines might originate from the host galaxy. For objects that are classified as SLSNe-II, we check whether they surpass a peak magnitude of -21 in any band. SNe with fainter peak magnitudes are here considered regular Type IIn SNe.

Forced photometry is obtained for all 209 remaining SNe and we apply the quality cuts as described in the following sections. After all cuts, pre-explosion observations are available for 196 SNe. This remaining sample consists of 131 Type IIn SNe, 26 SLSNe-II, 20 SNe with flash-spectroscopy signatures, 12 Type Ibn SNe, and 7 SNe Ia-CSM. Table 1 lists the SNe for which pre-explosion activity is detected (see

Sec. 3) and a full version of this table containing all initially considered 227 SNe is available online.

2.2. The Forced Photometry Pipeline

We perform forced photometry using the pipeline described by Yao et al. (2019) on difference images obtained from IPAC via IRSA³. Details of the ZTF image reduction are given by Masci et al. (2019) and image subtraction is based on the method developed by Zackay et al. (2016). We have access to images from the ZTF partnership survey (40% of the observation time) and images that became available

³ <https://irsa.ipac.caltech.edu/Missions/ztf.html>

during the third data release⁴, which includes images from the public survey (also 40% of the time) until December 2019 and Caltech data (20% of the time) until December 2018. Forced photometry on more recent public or Caltech data cannot be done as the full images are not yet available.

The forced-photometry pipeline was implemented by Yao et al. (2019). It relies on the IPAC difference images and the measured point-spread functions (PSFs). An image cutout around the SN position is produced and the background is measured within an annulus with an inner radius of 10 pixels and an outer radius of 15 pixels, where 1 pixel corresponds of 1.01'' on the sky. The median background flux is subtracted from the cutout and the 7×7 pixels around the SN position are used for the PSF fit. To quantify the uncertainty in the flux, the normalization of the PSF is fitted with a Markov chain Monte Carlo algorithm. While Yao et al. (2019) used 250 random walkers for the fit, we lower the number to 50 walkers to reduce the computation time. For 50 walkers the fitting algorithm introduces an uncertainty that is smaller than 2% of the typical error in the measured flux. We hence find that 50 walkers provide sufficient accuracy.

Based on the procedure of Yao et al. (2019) as well as our own findings, we exclude some data points from the light curves. Our exclusion criteria are as follows.

1. Images obtained early in the survey with an unknown quadrant ID for which the reference image cannot be identified.
2. Flagged difference images which might suffer from issues during the image subtraction.
3. Observations with seeing > 4''. The PSF fit is only done on the inner 7×7 pixels and might not be accurate for a very broad PSF.
4. Images affected by bad pixels at the SN position (inner 7 × 7 pixels).
5. Early *g*-band observations obtained between JD 2458120 and 2458140, which are not well calibrated.
6. Difference images with a background standard deviation > 25 in units of detector data number (see Yao et al. 2019) which indicate problems during the image subtraction.
7. Data points with flux errors that are seven times larger than the median flux error for this SN to remove images for which the PSF fit did not converge.

These initial quality cuts remove ~ 10% of the data (see also Sec. 3). We are left with 85,333 pre-explosion data points

which are listed in Table 2. All fluxes are corrected for Milky Way extinction using the python package *sfdmap*, which is based on the dust map of Schlegel et al. (1998) recalibrated to the values of Schlafly & Finkbeiner (2011) and the Cardelli et al. (1989) extinction law.

2.3. Background Samples

We quantify the expected rate of false detections by performing forced photometry in locations where no precursors are expected. The four background samples are

1. empty positions in the sky close to the SN position, but outside of the host galaxy;
2. faint *Gaia* stars with *g*-band magnitudes between 20.5 and 18.5 close to the SN position, to identify misaligned images;
3. the SN position mirrored across the center of its host galaxy; and
4. the positions of Type II_n SNe discovered during the PTF survey before 2015.

The tests are designed such that they start from a case for which image subtraction is easy (an empty position in the image) and progress to increasingly more realistic, but challenging environments for our pipeline. The first three tests are done for the exact same images that also contain the SN positions; hence, they have the same observing conditions, reference images, and subtractions. The two last tests are considered the most realistic ones as they are performed in host galaxies or locations where Type II_n SNe explode. The second background sample is used to identify and exclude images with astrometric errors.

The positions for the background samples are generated as follows. For empty locations we randomly pick several locations at a distance of 50 pixels (i.e., 50.6'') from the SN position. Faint stars or unresolved galaxies are selected from the *Gaia* catalog. To reject extended sources we require an astrometric excess noise of less than 1 milliarcsec and the *g*-band magnitude is limited to values between 20.5 to 18.5 to ensure that the luminosity is similar to that of a faint precursor. Moreover, the separation from the SN position is required to be at least 20 pixels, such that the SN light does not fall within the annulus region for which the background level is calculated (see Sec. 2.2). To identify the SN host galaxies, we query the NED database for objects close to the SN position. We reject those identified as stars, the SN itself, and infrared sources, many of which are also stars (Cutri et al. 2013).

The selected empty locations, faint stars, and host-galaxy candidates are then displayed on top of the reference image for visual inspection. When selecting empty positions and

⁴ <https://www.ztf.caltech.edu/page/dr3>

Table 2. Forced photometry pre-explosion light curves

SN name	ZTF name	JD	band	ref. im.	flux	flux err.	sys. err.	red. χ^2	red. χ^2_{star}
					10^{-10}	10^{-10}	10^{-10}		
SN 2018atq	ZTF 18aahmhxu	2458076.93147	<i>r</i>	5751232	-0.0448	8.38	1.26	0.56	0.85
SN 2018atq	ZTF 18aahmhxu	2458079.03350	<i>r</i>	5751232	6.49	11.7	1.26	0.96	1.33
SN 2018atq	ZTF 18aahmhxu	2458089.03796	<i>r</i>	5751232	6.66	8.41	1.26	1.15	1.30
SN 2018atq	ZTF 18aahmhxu	2458091.02603	<i>r</i>	5751232	-39.4	24.4	1.26	0.44	1.25
SN 2018atq	ZTF 18aahmhxu	2458091.04729	<i>r</i>	5751232	11.5	29.3	1.26	0.87	1.32

NOTE—The fifth column specifies which reference image was used (e.g. for the first rows the image for the ZTF field 575, CCD 12, quadrant 3 and filter 2, the *r* band; see also Yao et al. 2019). All fluxes have been corrected for the zeropoint and are given as a dimensionless ratio (see Eq. 8 in Yao et al. 2019). This flux ratio is also known as "maggie" (Finkbeiner et al. 2004). The third to last column lists the noise level in the reference image which is a systematic error on the measured flux. The two last columns show the reduced χ^2 of the PSF fit at the SN location as well at the location of a nearby faint star (see Sect. 2.4). The full version of the table is available online.

stars, we check that they are isolated, located outside of the host galaxy, and are not affected by artefacts in the reference images, such as dead columns, stellar spikes, or the edge of the image. Among the host-galaxy candidates we select the most likely host. For most images, a known galaxy is consistent with the visible center of the host in the reference image, but for a few objects we select a UV source. If several NED sources are close to the center of the host we compare with multicolor SDSS images to identify the most likely center. We caution that we might not identify the true host center in all cases. These positions are primarily used to build a background sample, so we do not require a high accuracy. With this method we locate the presumable centers of 160 host galaxies. The hosts of the remaining SNe are not listed in the NED database, mostly because they are faint. Some of them are even undetected in the ZTF reference images, especially for SLSNe.

The SN position is then mirrored on the location of the identified host galaxy and we verify that the two positions are sufficiently separated. The PSF fit is done for the inner 7×7 pixels — that is, the pixel containing the SN position and the three neighboring ones. However, if the seeing disk is large, the PSF of the SN could be broader. We therefore require a separation of at least 10 pixels between the actual and mirrored positions. Only 59 out of 160 SNe with identified host galaxies show a sufficiently large separation (see also Table 1). To increase the sample size, we select in addition SNe of Type II_n that were discovered during the PTF survey. We query the TNS database for publicly available SNe detected prior to 2015. A slowly developing Type II_n SN might still be detectable after ~ 3 yr, but an inspection of the ZTF light curves shows that this is not the case for any of the selected objects. Moreover, we add six objects analyzed by Ofek et al. (2014a) for which the SN was not observed by

PTF. This brings the sample to a total of 104 objects out of which ZTF data are available for 100.

We produce forced photometry light curves for all selected positions to test the pipeline. The sample of *Gaia* stars is used in Sec. 2.4 to reject misaligned images with astrometric residuals produced during the image subtraction. The other samples are used in Sec. 2.5 to inspect the data quality and in Sec. 2.7 to estimate the rate of false-positive detections. Table 3 shows the impact of the derived cuts and corrections on the number of (false) detections and on the total number of data points. The sample of *Gaia* stars is omitted in the table, because variable stars may result in actual detections.

2.4. Astrometric Errors

The large number of analyzed observations requires tight cuts on the data quality to avoid false-positive detections. Some of the reference or difference images might suffer from misalignments such that residuals are created in the image-subtraction process. Alignment errors result from several factors, including atmospheric scintillations (e.g., Osborn et al. 2015; Ofek 2019). To identify and remove affected images, we perform forced photometry at the position of a relatively faint star or an unresolved galaxy close to the SN position as described in Sec. 2.3. We choose faint stars because they roughly represent the surface brightness of bright galaxies. If the images are well aligned, no detection is expected for a nonvariable star, or for a variable star the residual should be well described by the PSF.

Images with astrometric residuals are identified via the reduced χ^2 of the PSF fit. We find that requiring a reduced $\chi^2 < 1.4$ at the position of the star removes most false detections. The corresponding difference images are flagged and are not used when searching for precursors at the SN position. In addition to misalignments, there could be more

Table 3. Cuts on data quality

step	# precursors / # data points			
	empty pos.	mirrored pos.	PTF SNe	real data
0 before cuts	4 / 176815	3 / 45092	116 / 48250	415 / 95442
1 known reference image	4 / 175888	3 / 44491	116 / 48067	415 / 94515
2 difference image not flagged	4 / 169522	3 / 43007	94 / 46328	399 / 91000
3 seeing $\leq 4''$	3 / 166119	3 / 42155	82 / 45292	382 / 88850
4 no bad pixels within 7×7 pixels	3 / 166119	3 / 42155	82 / 45292	382 / 88850
5 no early <i>g</i> -band images	3 / 165351	3 / 41988	75 / 45038	365 / 88078
6 std. of bkg. < 25	3 / 163808	3 / 40877	75 / 44349	362 / 86141
7 err. on flux < 7 times median err.	3 / 162485	3 / 40575	75 / 44054	361 / 85333
8 red. $\chi^2 < 1.4$ for nearby star	3 / 150637	3 / 37774	73 / 40740	265 / 78946
9 red. $\chi^2 < 1.4$ at SN position	2 / 148884	3 / 36888	5 / 37058	204 / 73105
10 ≥ 20 pre-expl. observations	2 / 136338	3 / 33600	4 / 36300	189 / 70420
11 offset correction	2 / 136338	9 / 33600	11 / 36300	189 / 70420
12 error-bar scaling	2 / 136338	1 / 33600	3 / 36300	136 / 70420
13 ref. sys. error / final unbinned	2 / 136338	0 / 33600	3 / 36300	152 / 70420
14 1-day bins	1 / 63791	1 / 15682	4 / 16979	124 / 32993
15 7-day bins	0 / 25528	2 / 6456	4 / 7616	84 / 14193
16 90-day bins	0 / 3983	2 / 1045	2 / 1281	37 / 2093

NOTE—Number of remaining data points and (false) 5σ detections after each step of the analysis, as described in Sec. 2.2 (steps 1 to 7), Sec. 2.4 (step 8 and 9), Sec. 2.5 (steps 10 to 12) and in Sec. 2.6 (step 13 and 14). Our actual search (last column) yields a much larger number of precursor detections than the three background samples. The initially large number of detections for the PTF sample is due to AGN activity in the host galaxy of SN 2011cc (see Sec. 2.7) and a few detections of this AGN persist after all cuts. The false detections for the empty and mirrored positions are all caused by a faulty reference image.

localized residuals or artefacts. We therefore require that the reduced χ^2 at the SN position is also smaller than 1.4. As shown in Table 3, these two cuts remove in total 12% of the data. The reduced χ^2 values for each data point are given in Table 2.

2.5. Offsets and Rescaling Flux Errors

As a next step, we verify that the pre-explosion light curves are centered around zero flux and that the estimated flux errors account for the observed flux scatter. When visually inspecting pre-explosion light curves, we find that the median fluxes are sometimes systematically offset from zero. In some cases, the offset could be due to light in the reference image either from the transient or from a precursor. However, we also see such offsets for the background samples. They can be as large as the typical error bar of the unbinned fluxes. We therefore do a baseline correction for all light curves. Consequently, we cannot identify precursors during the reference period or very long-lasting precursors that affect all data points (see also Appendix B). Moreover, we find that the size of the error bars is overestimated or underesti-

mated by typically 10–20%. For a handful of locations, the errors even have to be increased by as much as 50% to account for the observed scatter.

These biases are corrected for each reference image separately. To do this precisely, we require at least 20 pre-explosion observations with the same reference image. If fewer observations are available, the corresponding data points are discarded (step 10 in Table 3). After applying all cuts, we find that no pre-explosion observations are left for 13 out of 209 SNe (see Sec. 2.1); most of them were detected in the beginning of the survey (see online version of Table 1). Our final sample hence consists of 196 SNe.

We find that using the median pre-explosion flux to correct the baseline does not work for all SNe, because some of them have long-lasting precursors that contain close to half of the data points. We therefore calculate the *iterative median* which is more robust. We first combine same-night observations in bins to avoid individual nights with many observations dominating the result. Next, the median is calculated for the binned data points and the data point with the largest deviation from the median (regardless of the size of the error

bars) is removed from the sample. This last step is repeated; we recalculate the median for the remaining points and remove the most distant data point, until only 30%, but at least 20, of the data points are left. The median of these remaining points is used as the baseline correction. We find that this algorithm reliably identifies the zero flux level and removes the impact of any precursors during the reference period.

When searching for precursors at the SN positions, we select all objects with positive or negative 5σ detections and check whether we can redo the baseline correction for a time range that excludes the potential precursor, preferentially after the SN has faded. If this is possible we recalculate the baseline correction, this time using a simple median. This step leads to additional precursor detections for SNe with few pre-explosion observations (e.g., SN2018eru and SN2018kag) and improves the baseline correction for SNe for which a large fraction of the data points are part of the precursor, such as SN2019fmb (see Sec. 3.1). We also find that observations obtained after SN2019cmy had faded are systematically lower than pre-explosion observations. As discussed in Appendix B, we are not sure whether this drop in flux is due to a systematic error or an extremely bright progenitor star. In this paper, we exclude the late-time observations and only discuss the short precursor detected relative to the flux level of the pre-explosion light curve (see Sec. 3.1).

Next, we scale up the flux errors if they are underestimated, which is again done for every reference image separately. As before, the result might be biased by precursors which can inflate the error bars and remain undetected as a consequence. We therefore split the pre-explosion light curves for each reference image into equal segments of 15 or more data points. We calculate the *local robust standard deviation* for each segment by determining the 15.9% and 84.1% percentile and dividing its difference by 2. The median standard deviation for all segments is used to judge whether the error bars are sufficiently large to account for the observed noise level. If the standard deviation is larger than 1 (i.e., the error bars cannot fully account for the observed size of the 1σ -region), the error bars are multiplied with the robust standard deviation of the median segment. No scaling is done if the standard deviation is smaller than 1 (i.e., the errors are overestimated compared to the observed scatter).

2.6. Binned Light Curves and the Systematic Error of the Reference Image

To increase our sensitivity to faint precursors we also search binned light curves. The bins are chosen such that same-night observations are always combined in the same bin and the edge of the last pre-explosion bin is at the end of the night in which the SN is discovered. We ensure that data points before and after the estimated explosion date are never combined in the same bin by binning the two parts of




Figure 1. Flux residuals for the background sample of mirrored positions (see Sec. 2.3) compared to a normal distribution. The baselines have been centered around zero and underestimated flux errors have been rescaled as described in Sec. 2.5.

the light curve separately. For each bin, we use the median observation date as the observation time of the bin and calculate the weighted mean flux and its uncertainty.

When combining a large number of observations in one bin, the uncertainty in the flux can become very small. However, the ZTF reference images only consist of about 15 coadded observations; hence, the noise level in the reference image has to be considered. For this purpose we convert the limiting magnitude of the reference image to a flux which is given in Table 2. This systematic error is added in quadrature to the uncertainty of the unbinned or binned fluxes. It is typically ten times smaller than the uncertainty in the flux measured in a single image and thus only becomes relevant if many observations are coadded in a bin. When combining flux measurements that have different reference images we use the median systematic error.

We verify that the flux residuals indeed follow a normal distribution with a width of 1 by showing the flux residuals (i.e., the flux divided by its uncertainty) in Fig. 1. Except for statistical fluctuations, the residuals roughly follow a normal distribution. We expect some deviations from a normal distribution, for example because we do not reduce the size of overestimated flux errors (see Sec. 2.5). For 15-day bins the systematic error in the reference image also becomes relevant, such that we expect a slightly narrower distribution. The distributions in Fig. 1 illustrate that only very few data points deviate from 0 by more than 3σ . This indicates that the forced-photometry pipeline and our cleaning process work well in locations where interacting SNe explode and that the error bars have an appropriate size after the scaling described in Sec. 2.5. For the precursor search we use a 5σ threshold.

2.7. Expected Number of False Detections and Astrophysical Backgrounds

The empty positions as well as the mirrored positions and historic SNe serve as a quality check of the forced-photometry pipeline. We do not expect any astrophysical precursors at these positions and can therefore use these to calculate the false-alarm rate. Table 3 shows that our actual search (last column) yields 152 5σ detections for unbinned light curves even though only a few false detections are expected. This gives us confidence that the majority of the detected precursors are astrophysical. In addition, these precursors are almost exclusively detected among Type II SNe and they prefer low-redshift objects, as expected.

Nevertheless, a small number of false detections persist after all cuts. We inspect them to identify possible reasons. For empty and mirrored positions all false detections occur in the g -band images that contain SN 2018bih. A visual check shows that the reference image contains structures that are not astrophysical. In the actual search for this SN, we do not find any precursor candidates, potentially because only few pre-explosion images are available owing to its explosion date in May 2018. Another notable issue is the large number of detections at locations where PTF SNe exploded prior to 2015 (penultimate column in Table 3). Most of them (97 out of 116 detections before cuts) are at the position of SN 2011cc and are likely due to AGN activity close to the SN position (see below). Our cuts remove most detections at this position, but a few remain. We conclude that our precursor search might yield a few false detections, for example owing to faulty reference images (1 out of 860 reference images affected) and background AGNs (1 out of 100 SN positions affected).

A large number of false detections is found at the position of SN 2011cc, which exploded at a distance of $2.4''$ from the center of its host galaxy. The host, IC 4612, is classified as a star-forming galaxy in the SDSS catalog (Ahumada et al. 2019) and as a narrow-line AGN by Liu et al. (2011). We therefore hypothesize that the variability observed in ZTF data is caused by AGN activity. Requiring a reduced χ^2 of < 1.4 at the SN location removes most detections, because the background AGN is slightly offset from the SN position. No variability or precursors were detected in the PTF pre-explosion light curve of SN 2011cc (Ofek et al. 2014a), likely owing to the relatively small number of observations. Another case of apparent variability due to potential background AGN activity is detected in the pre-explosion light curve of SN 2019meh when searching the actual SN locations (see Sec. 3.1).

In addition, we find that light from the Type Ia SN SN 2018big contaminated the pre-explosion light curve of the flash-spectroscopy object SN 2019nvm. Both SNe happened in the same host galaxy with a separation of $3.6''$, so SN 2018big is just at the edge of the 7×7 pixel region for which the PSF fit is done (see Sec. 2.2). Since we re-

quire a small reduced χ^2 at the SN position (step 9 in Table 3), all detections of SN 2018big are rejected, such that the object does not show up as a potential precursor in the search described in Sec. 3.1. These coincidences serve as reminders that pre-explosion activity does not necessarily originate from the progenitor star, but could be related to bright, variable objects within $\lesssim 4''$.

It is also possible that a different star close to the progenitor produces precursor eruptions. It could even be the progenitor of a SN that might explode at a later time. We consider this scenario relatively rare, as no further precursors are detected in ZTF data at 100 positions where PTF detected Type II SNe before 2015 (see Sec. 2.7). Nevertheless, this possibility cannot be ruled out in individual cases.

Another challenge is distinguishing between a precursor and the rising SN light curve. Double peaks or early plateaus, likely powered by shock cooling (see e.g. Sapir & Waxman 2017), have been observed for several SNe of Type Ib, Ibn and Iib (see e.g. Gal-Yam 2017). Piro & Nakar (2013) estimate that SNe powered by radioactivity can undergo a dark phase of up to several days. After this time emission from centrally located radioactive nickel-56 is able to diffuse outwards and the SN starts to rise to its main peak. We find such early detections for several SNe (e.g. for SN 2019fci). If the detection is separated by less than a week from the observed rise, we assume conservatively that the SN has already exploded at this time and adjust the discovery date t_0 accordingly. As a consequence, we might miss short-lived precursors immediately prior to the SN detection. This is especially true for objects for which the rise of the light curve is not well observed.

3. PRECURSOR PROPERTIES

After developing and testing our analysis in the previous section, here we apply it to the actual data. The detected precursors and additional tests are described in Sec. 3.1, the precursor absolute magnitude light curves and radiated energies are calculated in Sec. 3.2, and their $g - r$ colors are presented in Sec. 3.3.

3.1. Detected Precursors

To search for precursors, we produce forced photometry light curves at the SN positions and apply the cuts and corrections described in Sec. 2. Any pre-explosion data points that are significant at the 5σ level are considered detections. To gain sensitivity to fainter precursors, we search in addition the binned light curves (see Sec. 2.6). The precursor durations are unknown and moreover depend on the detection threshold. To cover a wide range of timescales we use six different bin sizes with lengths of 1, 3, 7, 15, 30, and 90 days. The bin sizes are chosen such that the amount of

SN2018eru (SN IIn)				1	2	2	2
SN2018gho (SN IIn)	4	5	5	5	3	2	1
SN2018hxe (SN IIn)	1	1	1				
SN2018kag (SN IIn)	4	7	5	3	3	3	2
SN2019uo (SN Ibn)				1	1	1	1
SN2019bxq (SN IIn)	2	1	2	1	2	2	2
SN2019cmy (SN IIn)	1	2	1	2	1	1	1
SN2019iay (SN IIn)	1	1	1	1	2	2	2
SN2019meh (SLSN-II)	1	1	3	5	5	3	2
SN2019gjs (SN IIn)	5	5	5	6	8	7	3
SN2019mom (SN IIn)	1	2	1	3	1		
SN2019njv (SN IIn)		1	1				
SN2019fmb (SN IIn)	50	34	29	20	15	8	5
SN2019sae (SN IIn)							1
SN2019aafe (SN IIn)	5	5	6	4	3	3	2
SN2019vkl (SN IIn)	13	11	13	10	9	5	2
SN2019vts (SN IIn)					3	3	2
SN2019qny (SN IIn)	2	2	2	2	3	3	3
SN2020iq (SN IIn)			1		1	1	1
SN2019yzx (SN Ia-CSM)		1	1				1
SN2019zrk (SN IIn)	57	38	21	12	6	4	2
SN2020dcs (SN IIn)	4	5	4	4	3	2	1
SN2020dfh (SN IIn)	1		1	1	1	1	1
SN2020edh (SN IIn)		2	3	3	2	2	
	unbinned	1 day	3 days	7 days	15 days	30 days	90 days

Figure 2. Number of pre-explosion detections above the 5σ threshold for each bin size. Observations in different photometric bands are binned separately and the white numbers indicate the total number of detections in all three bands. Dark-green fields represent securely detected precursors, while light-green fields mark unconfirmed precursors that are only seen in a single bin and band (see text). Orange fields signal that the flux variability is likely caused by AGN activity in the host galaxy and not by the progenitor star.

data approximately doubles or triples when going to the next larger bin size.

In our search of the pre-explosion data of 196 SNe, we find precursor candidates prior to 24 SNe, mostly of Type IIn; Fig. 2 indicates the number of 5σ detections in each search channel. Most precursors are detected using several different bin sizes, indicating that they are both bright and long-lasting. The precursor light curves in 1-day bins are shown in Figs. 3 and 4, and their properties are summarized in Table 4. In addition, we show coadded difference images of the precursors in Appendix A. They demonstrate that the detections are indeed due to point sources at the SN location with the exception of SN 2019sae, which might be spurious.

Marginally detected precursor candidates are inspected in more detail, to test whether they are genuine. For precursors that are only detected in a single bin, we check whether fluxes in the three bins before or after the detection surpass the 3σ significance threshold or whether reducing the bin size leads to at least two data points above the 3σ threshold. If we do not find any additional 3σ detections, we conclude that the detection is driven by data collected within a single night and band and refer to these detections as *unconfirmed precursors*. The light-green color in Fig. 2 highlights the four precursors that do not pass this test (the Type IIn SN 2018hxe, SN 2019njv, and SN 2019sae, and the Type Ia-

CSM SN 2019yzx). The fact that we only found few false 5σ detections when searching the background samples in Sec. 2.3 suggests that at least some of the unconfirmed precursors are astrophysical nonetheless. In the following we only focus on the 19 securely detected precursors.

One pre-explosion light curve, prior to SN 2019meh, shows long-term up-and-down fluctuations as expected for AGNs (see Fig. 3). Indeed, the SN is located within $1''$ of the center of its host galaxy and we therefore conclude that the variability is likely due to nuclear activity and is not caused by the progenitor star, as described for SN 2011cc in Sec. 2.7. SN 2019meh is a SLSN of Type II located at a relatively high (for our sample) redshift of 0.0935. Such a distant progenitor star would have to reach an extreme luminosity to be detectable, which supports the hypothesis that we are seeing AGN activity rather than stellar flares. We therefore remove this object from the sample.

We also check whether shifting the bin positions leads to the detection of additional precursors. For this purpose, we repeat the search with 7-day bins six times while moving the bin edges by one day for each new search. We detect precursors prior to a few SNe that are not found with the original 7-day bins, but all additional SNe already have detected precursors when using smaller or larger bins (see Fig. 2). We thus conclude that the bin positions only have a minor influence on the results.

We summarize that we securely detect pre-explosion outbursts prior to 18 different SNe of Type IIn and prior to the Type Ibn SN 2019uo (see Fig. 2). Figures 3 and 4 show that some SNe, such as SN 2018eru, SN 2019bxq, SN 2019mom, or SN 2020edh, might undergo several separate precursor eruptions. It is, however, also possible that the detections are part of a single flaring episode that lasts for several hundred days.

3.2. Precursor Energy

To put the precursor eruptions into context, we calculate the absolute magnitude light curves and estimate the radiated energies of the securely detected precursors found in Sec. 3.1. Fluxes are converted to “asinh magnitudes,” also called “luptitudes,” with a softening parameter of 10^{-10} as defined by Lupton et al. (1999). Magnitude errors are given as $\sigma_{\text{mag}} = 1.0857 \log(f/\sigma_{\text{flux}})$, where f and σ_{flux} are the dimensionless normalized fluxes and uncertainties shown in Figs. 3 and 4. The 5σ limiting magnitude is calculated as $m_{\text{lim}} = -2.5 \log(5 \times \sigma_{\text{flux}})$ and the significance of a detection is given as f/σ_{flux} . All calculations are here done for 7-day bins and we consider 3σ detections significant if they are part of a previously detected precursor. For data points that do not reach the 3σ threshold we calculate 5σ upper limits.

Figure 5 shows the resulting absolute magnitude r -band light curves (dashed lines indicate that the g band was




Figure 3. Precursor light curves in 1-day bins. Solid circles mark 5σ detections, circles filled with a lighter shade and without a black edge have significance between 3σ and 5σ , and open points are less significant. Green, red, and black data points were obtained in the g , r , and i bands, respectively. The vertical black line indicates t_0 , a rough estimate for the explosion date, and colored areas identify precursors that are detected significantly when using larger bins. The flux f is given as a unitless ratio relative to the zeropoint which is equivalent to the unit “maggie” used in SDSS catalogs (Finkbeiner et al. 2004). Corresponding AB magnitudes are given on the right-hand ordinate axis and are calculated as $m_{AB} = -2.5 \log_{10}(f)$.

used instead for SN 2018gho, SN 2018kag, SN 2019bxq, SN 2020edh, and the early detection of SN 2018eru). For clarity we omit nondetections that do not directly constrain the precursor duration. Most precursors are detectable for several weeks and some of them start more than 100 days before the explosion. The peak magnitudes vary between -13 and -17.5 as also summarized in Table 4. For comparison we add the r -band light curve measured for the 2012a event observed immediately prior to the likely final explosion of SN 2009ip (data taken from Margutti et al. 2013, Prieto et al. 2013, and Pastorello et al. 2013). Its duration, peak magnitude, and shape are similar to those of several of the less energetic precursors found in this search. We hence conclude that bright and long-lasting precursors are common in the last months before the explosion of Type II_n SNe. Their rate is quantified in Sec. 4.

Next, we calculate the precursor energies by integrating the fluxes per bin from the first to the last detection, even if individual data points in between are not significant at the 3σ level. The calculation is done for each band separately and gaps in the data are interpolated if one or two 7-day bins are empty, such as for SN 2018gho (see Fig. 3). This interpolation increases the total energy by at most 30% and thus does not have a major impact on the results. We obtain similar results for 3-day bins and thus conclude that the energy estimates in Table 4 roughly describe the observed precursor energy. These are lower limits on the true radiated energies of the precursors, which are often only partially detected, and also radiate outside of the visible-light bands which we cover. The brightest precursors reach radiative energies close to 10^{49} ergs, about 10% of the total radiative energy in a typical SN explosion.

3.3. Precursor Colors

Here, we calculate the $g - r$ color index for precursors that have observations in both bands. For this purpose we select all bins in which a significance of 3σ is reached in at least one band. If the detection in the second band is less significant we quote lower or upper limits accordingly. The resulting colors are shown in Fig. 6. Compared to young SNe, the precursors exhibit quite red colors, which correspond to lower effective temperatures. We caution, however, that the $H\alpha$ line falls within the r band. The spectrum of the precursor prior to PTF 13efv showed relatively strong, narrow hydrogen lines (Ofek et al. 2016), and the same is true for Type II_n SNe and LBV outbursts. A red color could therefore be mimicked by a blue continuum flux with a strong $H\alpha$ line. The precursor prior to SN 2019fmt is also detected in the i band and shows a mean $g - i$ color of 1.1 mag. This corresponds to an effective temperature of ~ 4300 K, similar to the result from the $g - r$ color index shown in Fig. 6. For this object, at least,

we conclude that the rather low effective temperature is not primarily due to a strong $H\alpha$ line.

We also show the effective temperatures of the 2012a outburst of SN 2009ip in Fig. 6. They were obtained by fitting a blackbody continuum to the multiband photometry (Margutti et al. 2013) and are therefore less susceptible to line fluxes. The precursor of SN 2009ip is slightly hotter than most precursors observed in our sample, and we find that the precursors detected here typically do not cool down as observed for the 2012a event prior to the final explosion of SN 2009ip (Margutti et al. 2013).

If the precursor's bolometric luminosity L and temperature T are known, photospheric radii can be estimated via the Stefan-Boltzmann law $R = (L/4\pi\sigma_B)^{0.5}T^{-2}$, where σ_B is the Stefan-Boltzmann constant. A faint and hot precursor (with a temperature of 8,000 K and a bolometric magnitude of -13) would have a photosphere with a small radius of $\sim 4 \times 10^{13}$ cm, while a bright and cool precursor (4,000 K and a magnitude of -17) would have a radius of $\sim 10^{15}$ cm. When using r -band luminosities (shown in Fig. 5) as order of magnitude estimates for the precursor bolometric luminosity and the $g - r$ color index as a crude temperature estimate, we find that most detected precursors have photospheric radii of a few times 10^{14} cm. These large radii suggest that we cannot see down to the surface of the progenitor star.

4. PRECURSOR RATES

Here we focus on the whole sample of pre-explosion light curves and use it to calculate precursor rates. Except for one, all confirmed precursors are found prior to Type II_n SNe and we therefore first describe the rate for this SN class in Sec. 4.1 and Sec. 4.2. Precursor rates for other types of possibly interacting SNe are presented in Sec. 4.4.

4.1. Precursor Rates for Type II_n SNe

The rate calculation is done for 7-day bins, because this search channel is sensitive to faint precursors without losing short precursors (see Fig. 2). Another advantage of using 7-day bins is that they partly compensate for differences between light curves obtained by the private and public surveys, which have typical cadences of one day and three days, respectively. None of the unconfirmed precursors is detected for 7-day bins, so they do not enter the rate calculation.

The precursor rate is here defined as the fraction of time during which precursors are observed above a certain limiting magnitude. As a result, we do not distinguish between two 1-week long precursors and a single precursor that lasts for two weeks. The rate depends on the absolute magnitude of the precursors and we calculate it in steps of 0.25 mag. For each absolute magnitude we select all pre-explosion bins with a deeper limiting magnitude. We then calculate which fraction of these bins have precursor detections. Consequently, precursors detected with a high significance (i.e., a

Table 4. Detected precursors

	band	start phase (days)	end phase (days)	median flux (mag)	energy (10^{46} ergs)	v_{CSM} (km s^{-1})	$\epsilon M_{\text{CSM, pre.}}$ (M_{\odot})	t_{rise} (days)	$M_{\text{CSM, diff.}}$ (M_{\odot})
SN 2018eru	<i>g</i>	−202.2	−195.2	−13.3	4	1100	0.02	–	–
	<i>g</i>	−69.2	−62.2	−13.3	4				
	<i>r</i>	−76.2	−34.2	−13.4	23				
SN 2018gho	<i>g</i>	−63.0	−14.0	−15.2	160	210	4	13	< 1.3
SN 2018kag	<i>g</i>	−48.1	−20.1	−14.3	40	1100	0.04	–	–
	<i>r</i>	−41.1	−20.1	−14.7	50				
SN2019uo	<i>r</i>	−342.8	−307.8	−13.0	17	880	0.007	8	< 0.8
SN 2019bxq	<i>g</i>	−349.4	−342.4	−13.0	3	330	0.06	18	< 1.8
	<i>g</i>	−188.4	−181.4	−12.9	3				
SN 2019cmv	<i>g</i>	−13.4	−6.4	−13.9	7	150	1.1	8	< 0.8
	<i>r</i>	−13.4	−6.4	−14.6	30				
SN 2019fmb	<i>g</i>	−125.3	−27.3	−12.9	40	990	0.08	–	–
	<i>r</i>	−174.3	−20.3	−13.3	80				
	<i>i</i>	−111.3	−41.3	−14.1	80				
SN 2019iay	<i>g</i>	−34.2	−13.2	−14.7	40	340	0.4	9	< 0.9
	<i>r</i>	−34.2	−13.2	−14.8	50				
SN 2019gjs	<i>g</i>	−97.2	−20.2	−15.3	300	320	3	7	< 0.7
	<i>r</i>	−104.2	−20.2	−15.6	300				
	<i>i</i>	−55.2	−48.2	−15.5	30				
SN 2019mom	<i>g</i>	−342.5	−335.5	−14.7	13	590	0.19	–	–
	<i>r</i>	−342.5	−335.5	−15.1	19				
	<i>g</i>	−6.5	0	−16.0	50				
	<i>r</i>	−6.5	0	−16.0	50				
SN 2019aafe	<i>g</i>	−20.4	−6.4	−16.9	300	1100	0.9	4	< 0.4
	<i>r</i>	−97.4	−6.4	−16.1	1000				
SN 2019vkl	<i>g</i>	−62.2	−6.2	−16.0	500	770	1.3	10	< 1.0
	<i>r</i>	−83.2	−6.2	−16.1	800				
SN 2019vts	<i>g</i>	−41.5	−20.5	−14.6	40	340	1.2	–	–
	<i>r</i>	−62.5	−6.5	−14.9	140				
SN 2019qny	<i>g</i>	−104.6	−90.6	−15.4	50	350	2	25	< 2.5
	<i>r</i>	−125.6	−62.6	−15.4	300				
SN 2020iq	<i>r</i>	−482.2	−454.2	−15.9	170	160	7	–	–
SN 2019zrk	<i>g</i>	−55.5	−6.5	−16.2	500	350	5	7	< 0.7
	<i>r</i>	−111.5	−6.5	−16.1	700				
SN 2020dcs	<i>r</i>	−34.4	−6.4	−15.1	90	180	3	14	< 1.4
SN 2020dfh	<i>r</i>	−34.6	−20.6	−15.2	40	200	1.1	–	–
SN 2020edh	<i>g</i>	−692.5	−545.5	−13.0	80	600	0.2	–	–

NOTE—Properties of the detected precursors and the SNe. The first columns list the beginning and end of precursors with respect to t_0 , the median magnitudes and precursor energies. The CSM velocity is derived from the median width of narrow lines and P Cygni profiles (see Sec. 5.1) and is used to estimate the CSM mass multiplied by an unknown efficiency factor ϵ . t_{rise} quantifies how many days it takes the SN to rise by a factor of e (1.086 mag) to its peak in the r band (g band used for SN 2018gho and SN 2019iay). The rise time provides a rough upper limit on the total CSM mass given in the last column (see Sec. 5.2).




Figure 4. Precursor light curves – continuation of Fig. 3




Figure 5. Absolute magnitude precursor light curves in 7-day bins. All SNe are of Type II_n, except for the Type Ib_n SN 2019uo (open markers). If available, we show the *r*-band light curve, and dotted lines indicate that the *g*-band light curves are depicted for SN 2018gho, SN 2018kag, SN 2019bxq, SN 2020edh, and the early detection of SN 2018eru. For clarity, only the most relevant nondetections are displayed. The 2012a event prior to the likely final explosion of SN 2009ip is shown as a black dashed line for comparison (light curve taken from Margutti et al. 2013, including data from Prieto et al. 2013 and Pastorello et al. 2013).




Figure 6. Color index of precursors observed in the *g* and *r* bands. The light curves are binned in 7-day bins and we quote 3σ lower or upper limits instead of detections if the *g*-band or *r*-band flux is not significant at the 3σ level.

large difference between its magnitude and the limiting magnitude of the bin) may contribute in several magnitude bins. On the other hand, detections just at the 5σ threshold may not contribute at all, if they fall in between the magnitude

steps⁵. The resulting rate is cumulative, as we search for precursors that are brighter than the corresponding magnitude threshold. The 95% uncertainty associated with the rate is calculated using the Wilson binomial confidence interval

⁵ For example, a precursor detected with an absolute magnitude of -14.2 and with a limiting magnitude of -14.1 would not count as a detection in the bin at magnitude -14.25 because it is not bright enough. In the next fainter bin at a magnitude -14 it also does not contribute because the limiting magnitude is not sensitive enough.




Figure 7. Fraction of time during which precursors brighter than the respective absolute magnitude are observed for Type IIIn SNe. The lines show the measured rates and the shaded area indicates the 95% confidence region. If it reaches down to zero, its upper edge should be interpreted as an upper limit; otherwise it corresponds to the 95% error bar on the rate. The calculation was done for 7-day bins; the green, red, and black colors show the rates for the g , r , and i bands, respectively. The i -band precursor rate is only measured at magnitude -14 , as indicated by the black dot. In the main panel, the median phase of the observations is ~ 6 months prior to the explosion, while it is 1.4 months and 10.4 months (respectively) for the two right panels. The r -band precursor rate measured in the last 90 days before the SN explosion (upper small panel) is typically 6 times larger than the 95% upper limit on the rate measured at earlier times (lower small panel).

(Wilson 1927; Wallis 2013) as implemented in the *astropy* package (Robitaille et al. 2013; Price-Whelan et al. 2018).

The main panel of Fig. 7 shows the fraction of time during which precursors are detected for Type IIIn SNe as a function of the absolute magnitude. The green, red, and gray shaded regions correspond to the parameter space that is allowed at the 95% confidence level for the g , r , and i bands (respectively), and the solid lines depict the cumulative precursor rate. For bins without detections (e.g., for bright absolute magnitudes) the colored area reaches down to zero and its upper edge corresponds to a 95% upper limit.

As shown in the main panel of Fig. 7, g -band and r -band precursors are detected with absolute magnitudes ranging from -17 to -12 . The rate is slightly lower in the g band, because of the red precursor colors observed in Sec. 3.3. The fraction of time during which we observe bright precursors with an absolute r -band magnitude of -16 or brighter is $\sim 0.7\%$ with a 95% confidence range of 0.4% to 1.2%. For fainter magnitudes, the rate increases and reaches $\sim 12\%$ (6–23%) for precursors brighter than magnitude -13 . The measured rates are also summarized in Table 5.

We caution that r -band precursors fainter than magnitude -14 are only detected for SN 2019fmb, so the rate of such faint precursors is determined by this object and by the fact

that few other SNe have as constraining observations. The g -band rate is more robust, since such faint precursors are detected for four different SNe. The dip in the r -band rate at magnitude -14 is likely a statistical fluctuation caused by the relatively small number of SNe with precursors. The gray shaded region indicates that the i -band observations are typically not sensitive enough to detect precursors. The reason is that fewer observations were obtained and they have in addition larger error bars, in part owing to the lower quantum efficiency in this wavelength range for the ZTF CCD (Bellm et al. 2019). A black dot marks the only i -band detection, a precursor with magnitude -14 prior to SN 2019fmb.

4.2. Time Dependence of the Precursor Rate for SNe IIIn

The rate calculation in the left-hand panel of Fig. 7 was done using all pre-explosion data that were collected over a period of up to 2.5 yr before each SN explosion. The median phase of the pre-explosion light curves is 267 days (nearly nine months) before the discovery date t_0 . The precursor light curves in Fig. 5 show that most precursors are detected in the final few months before the SN explosion. To quantify the time dependency, we split the dataset into two parts: observations collected within 90 days before the estimated explosion date (with a median of 42 days) and observations collected

Table 5. Precursor rates

sample	band	number of SNe	median phase	rate of bright pre. (≤ -16 mag)	rate of faint pre. (≤ -13 mag)
			(months)	(%)	(%)
Type IIIn, all data	<i>g</i>	122	-8.7	0.5 (0.3 – 1.1)	6 (2 – 15)
	<i>r</i>	126	-8.8	0.7 (0.4 – 1.2)	12 (6 – 23)
	<i>i</i>	49	-9.0	< 0.9	–
Type IIIn, ≤ 90 days before SN	<i>r</i>	107	-1.3	3 (1.7 – 6)	26 (12 – 49)
Type IIIn, > 90 days before SN	<i>r</i>	121	-10.4	< 0.2	5 (1.4 – 17)
bright Type IIIn (peak mag. < -18.5)	<i>r</i>	84	-8.5	2 (1.2 – 4)	< 73
faint Type IIIn (peak mag. > -18.5)	<i>r</i>	33	-8.4	< 0.3	12 (6 – 12)
Type Ibn	<i>r</i>	11	-8.6	< 1.0	< 31
SLSNe-II	<i>r</i>	24	-8.4	< 72	–
flash-spectroscopy SNe	<i>r</i>	20	-4.4	< 0.5	< 2.7
Type Ia-CSM	<i>r</i>	7	-7.2	< 5.0	–

NOTE—Fraction of time during which bright or faint precursors are observed with the 95% confidence range given in parentheses. If no precursors are detected the 95% upper limit is quoted instead. The calculation was done for 7-day bins and the numbers are taken from Figs. 7, 9, and 10. The number of SNe with data is given in the third column, and the fourth column lists the median phase of the pre-explosion observations which is close to nine months for most subsamples. Dashes indicate that no data are available, so the rate remains unconstrained (e.g., the rate of faint precursors in the *i* band).

earlier (at a median time of 317 days before the SN). We then repeat the rate calculation and display the results in the two side panels of Fig. 7.

The *r*-band 95% confidence regions in the two smaller panels of Fig. 7 do not overlap for absolute magnitudes > -14.5 and the precursor rate is significantly larger in the final 90 days before the explosion. The measured rate in the final months before the explosion is up to 16 times larger than the 95% upper limit on the precursor rate before that. In the *r* band, the median difference for all magnitude bins between magnitude -13 and -17.5 is a factor of 6 (i.e., the precursor rate at early times is typically more than 6 times smaller). The difference would be even larger when dividing the dataset at 120 days, because several of the detections in the lower-right panel of Fig. 7 are part of the > 100 -day long precursors (e.g., prior to SN 2019fmb and SN 2019gjs; see Fig. 5). The large number of precursor detections shortly prior to the explosion is hence not caused by the larger amount of data available at these times, but is a genuine and significant difference.

In the three months before the explosion, faint precursors with an *r*-band magnitude of > -13 are observed 26% of the time (with a 95% confidence range of 12–49%; see also Table 5), while the rate is 5% (1.4–17%) at earlier times. The time dependence of the rate is even stronger for brighter precursors with absolute magnitudes > -16 : their rate is 3% (1.7–6%) in the three months before explosion, while it is




Figure 8. Fraction of Type IIIn SNe with long-lasting (≥ 30 days) precursors within 90 days prior to their explosion. The lines mark measured rates and the shaded area indicates the 95% confidence regions.

$< 0.2\%$ prior to that. We conclude that precursors become brighter and more frequent in the final months leading up to the explosion.

Early precursors are only observed prior to five SNe (see Fig. 5) and they appear to be fainter and short-lived compared to the precursors immediately before the explosion that typically last for several months. The rate calculation in Fig. 7 shows that this effect is real and not caused by a smaller number of observations at early times. The luminosity increase

likely continues within the last three months before the explosion as shown in Fig. 5.

While we so far constrained the fraction of time during which precursors are observed, we here calculate in addition the fraction of progenitor stars that undergo long-lasting precursors within 90 days before the SN explosion. For this purpose, we compile a sample of SNe for which such precursors are detectable. We bin the light curves in 30-day bins and require that at least two bins contain data (i.e., that observations are available in two out of three months). If this condition is met, we estimate down to which limiting magnitude a precursor can be detected. For this purpose we use the second lowest limiting magnitude — that is, the median for three data points or the least constraining bin for two data points. The rate is then calculated for each magnitude bin by dividing the number of detected precursors by the number of light curves for which such a precursor would have been detectable.

The fraction of SNe with long-lasting precursors in the last three months before the explosion is shown in Fig. 8. Long-lasting precursors brighter than magnitude -16 occur for about 4% (1.1–14%, 95% confidence range) of the SNe in the r band, while fainter precursors with an absolute magnitude brighter than -13 occur for 25% (5–69%) of the Type IIIn SNe. The i -band precursor rate is unity at magnitude -14 , but it is purely determined by SN 2019fmb as no other SN has as constraining observations. The rate is here detected in two magnitude bins because the 30-day-long light curve bins yield deeper limiting magnitudes than the 7-day-long bins used in Fig. 7.

We hence conclude that precursor eruptions brighter than magnitude -13 occur prior to many, but not all Type IIIn SNe. This result is in tension with some of the findings by Ofek et al. (2014a), who calculate that the average Type IIIn progenitor undergoes several precursors brighter than magnitude -14 in the last year before its explosion. Based on this they estimate that $> 52\%$ of all Type IIIn SNe exhibit at least one bright precursor in the final four months before the explosion at a confidence level of 99%. Ofek et al. (2014a) calculate the precursor rate by dividing the number of precursors by the time during which such precursors are detectable, the so-called “control time.” However, if the light curve has gaps, the control time (and thus the rate) depends on the bin size while the number of precursors does not change, as long as the bin size is smaller than their duration. To avoid such a dependence on the bin size, we calculate instead the fraction of bins with precursors or the fraction of well-observed SNe with precursors. The rate calculation used by Ofek et al. (2014a) and Strotjohann et al. (2015) are thus only valid if each light-curve bin contains observations.

Our results are likely consistent with the findings of Bilinski et al. (2015), who did not detect any precursors for a sam-

ple of five Type IIIn SNe and one SN imposter. They report that a precursor similar to the 2012a event prior to the explosion of SN 2009ip would have been detectable for two of their objects. We measure that $\sim 10\%$ of all Type IIIn SNe have precursors as bright as magnitude -15 (see Fig. 8) which is consistent with their non-detections. Bilinski et al. (2015) do not quote a control time, so we cannot compare to all of their results.

We conclude that the precursor rate increases by a factor of more than six within the last three months before the SN explosion compared to earlier observations obtained on average ten months before the SN. While the rate of faint precursors (with an r -band magnitude of -13) increases by a factor of ~ 5 , the difference is more than a factor of 10 for bright precursors with an r -band magnitude of brighter than -16 . Our observations do not constrain the rate of long-lasting precursors that are fainter than magnitude -13.5 . It is hence possible that all progenitors of Type IIIn SNe exhibit precursors if at least one third of them are fainter than this threshold.

4.3. Precursor Rates for Faint and Bright SNe IIIn

Type IIIn SNe can have diverse peak luminosities and SN energies. Here, we split the sample of Type IIIn SNe into bright and faint subsamples to test whether they have similar precursor rates. We consider a SN bright if it reaches an absolute magnitude of -18.5 in any ZTF band. This threshold is chosen such that the measured precursor rates are relatively well constrained in both subsamples. Detections in all three bands are considered, because some SN light curves only have sparse observations, especially if their peak occurred in the year 2020, for which part of the data has not yet been released (see Sec. 2.2).

We compare the r -band precursor rates for bright and faint SNe in Fig. 9 and the subsample of bright SNe has a higher rate of bright precursors. The rate of faint precursors is not well constrained for the bright SN sample, because most objects in this subsample are located at large distances. The rates could therefore agree below an absolute magnitude of -14 (see also Table 5). The difference between the bright and faint sample is relatively strong in the left-hand panel of Fig. 9, which shows the rate as the fraction of time during which the progenitor stars undergo precursors and thus depends on the precursor duration (see also Sec. 4.1). In the right-hand panel of the figure, we show instead the fraction of SNe that undergo a long-lasting precursor immediately before the explosion (like in Fig. 8) and the difference is not significant any more. A possible explanation for this change could be that bright precursors have longer durations. Indeed, the three brightest precursors in Fig. 5 are all observed for ~ 100 days. We hence find indications that luminous SNe typically undergo brighter and longer-lasting precursors. This correlation is quantified in Sec. 5.2.




Figure 9. r -band precursor rates for faint and bright (peak magnitude < -18.5 in the g , r , or i band) SNe of Type IIb. The left plot displays the fraction of light-curve bins in which precursors are detected, which means that the rate depends on the precursor duration (like Fig. 7). The right figure is only based on well-observed SNe and the rate indicates whether a long-lasting precursor is detected within the last three months before the explosion (like in Fig. 8).

4.4. Precursor Rates for Different Interacting SNe

As described in Sec. 2.1, our full sample also contains interacting SNe that do not belong to the class of Type IIb SNe. Here, we present precursor rates for SNe of Type Ib_n (based on 12 objects for which pre-explosion observations are available; see the online version of Table 1), SLSNe-II (26 objects after excluding SN 2019meh which falls on top of a background AGN), flash-spectroscopy SNe (20 objects), and Type Ia-CSM SNe (7 objects). The number given in Table 5 can be lower as not all SNe have pre-explosion data in the r -band. Flash-spectroscopy events are here defined as objects showing narrow He II lines in their early-time spectra up to a week after the discovery. There is some overlap between flash-spectroscopy SNe and the other classes: Some flash-spectroscopy SNe show narrow hydrogen lines for several weeks and are here included in the sample of Type IIb SNe (such as SN 2019cmy). SN 2019uo, is considered a Type Ib_n SN, even though it might show flash-spectroscopy lines at early times (Gangopadhyay et al. 2020).

We calculate the fraction of time during which precursors are observed in the same way as in Sec. 4.1 and show the results for each subsample in Fig. 10 for 7-day bins. The precursor detected prior to the Type Ib_n SN 2019uo (described in more detail in Sec. 5.3), does not appear because it is marginally above the 5σ threshold. An unconfirmed precursor is detected 680 days before the explosion of the Type Ia-CSM SN 2019yzx, as shown in Fig. 4. However, its significance is purely driven by observations in a single night while the two neighboring data points are consistent with zero. The location is observed relatively sparsely, so we cannot confirm whether the detection is real. We here conservatively assume that the detection is not astrophysical.

For comparison, the measured r -band precursor rate for Type IIb SNe (from the main panel of Fig 7) is shown as a

dashed red line in Fig. 10. The Type IIb rate is nearly always in the allowed region of parameter space, which means that we do not expect to detect any precursors even if the rates are as high as for Type IIb SNe. The lower sensitivity is due to the small sample size, or in the case of SLSNe to the fact that the objects are located at large distances (see also Table 5). The only region where the Type IIb SN rate is higher than the upper limit is for the sample of flash-spectroscopy SNe at faint precursor magnitudes of > -14 . However, in this region the Type IIb SN rate is completely dominated by SN 2019fmb and we therefore consider it less reliable.

Thus, we conclude that we only observe a single precursor that was not associated with a Type IIb SN, but with the Type Ib_n SN 2019uo. However, this small number of detections is expected owing to the small sample sizes of the subclasses and to the large distances of SLSNe.

5. IMPACT OF THE PRECURSORS ON THE SNE

In this section we explore the impact of the observed precursors on the SN spectra and light curves. In Sec. 5.1 we consider whether the narrow lines in the SN spectra can be produced by material emitted during the observed precursors. Next, in Sec. 5.2 we test whether SNe with observed precursors are brighter than other SNe in our sample. Finally, in Sec. 5.3 we describe how the precursor prior to the Type Ib_n SN 2019uo could account for both the SN light curve and the spectral evolution of this object.

5.1. Progenitor Mass-Loss History

A massive star of $100 M_{\odot}$ reaches its Eddington luminosity when it becomes brighter than 1.3×10^{40} ergs s^{-1} . For a hot LBV star with a temperature of 20,000 K (see, e.g., Smith et al. 2004) this luminosity corresponds to an absolute r -band magnitude of -8.1 , while it is -10.0 for a temperature of 5,000 K which is more similar to the temperatures observed




Figure 10. 95% upper limits on the precursor rates for 7-day bins for Type Ibn SNe, SLSNe-II, flash-spectroscopy events, and Type Ia-CSM SNe. The shaded areas indicate the 95% confidence area and the area above it excluded by the nondetection of precursors. The g and r -band observations are nearly equally sensitive such that the upper limits fall on top of each other, but the i -band is less constraining. The red dashed line indicates the r -band rate that was measured for Type IIc SNe in the main panel of Fig. 7.

for the precursors in Fig. 6. Figure 5 shows that the luminosities of all detected precursors are clearly above this threshold, so the outbursts are likely accompanied by strong mass-loss events.

By detecting the precursor, we measure the time of the mass ejection; however, the velocity of the material is uncertain (see also Sec. 6.1.2). The gray regions in Fig. 11 indicate out to which radii the material has expanded at the time of the SN explosion, depending on its velocity. CSM velocities between 0 and $1,500 \text{ km s}^{-1}$ are shown on the linear ordinate axis. We here assume that the material was ejected from a radius of $100 R_{\odot}$, $7 \times 10^{12} \text{ cm}$. Using a radius that is a factor of a few larger or smaller does not have a major impact on the results, as spectra are usually obtained after the SN ejecta have expanded far beyond this radius.

Most detected precursors occur within the last few months before the explosion, so any ejected material is still located within a radius of 10^{15} cm even if it has a velocity of $\lesssim 1,000 \text{ km s}^{-1}$. Earlier precursors are only observed for six SNe (SN 2018eru, SN 2019uo, SN 2019bxq, SN 2019mom, SN 2020iq, and SN 2019edh; see Figs. 3, 4, and 5). The ma-

terial ejected in these precursors might be located at radii of a few $\times 10^{15} \text{ cm}$, but likely below 10^{16} cm .

Next, we consider the narrow emission lines in the spectra of the SNe. To estimate CSM velocities, we measure the full width at half-maximum intensity (FWHM) of the narrow component of the $H\alpha$ line. We subtract the approximate resolution of the spectrograph in quadrature or quote upper limits if the result is smaller than half of the resolution. In addition, we look for narrow P Cygni features in the $H\alpha$ line (He lines for the Type Ibn SN 2019uo), as their minimum indicates the typical velocity of material moving toward the observer. The results for all spectra with clear narrow features are listed in Table 6. The quoted velocities are only order-of-magnitude estimates as we do not fit line profiles, measure the actual resolution of the spectra, or subtract host-galaxy contributions.

The exact location of the material that produces the narrow features is unknown, but the time when the spectrum was obtained provides an order-of-magnitude lower limit on its radius. Narrow features can only originate from unshocked material, which must be located at larger radii than the SN ejecta. In order to estimate these radii, we adopt a fiducial average ejecta velocity of 10^4 km s^{-1} , which is close to the




Figure 11. Radial CSM distribution for SNe with precursors. The gray regions indicate the possible locations of material ejected during the observed precursors depending on its velocity (shown on a linear scale). Data points correspond to spectra in which we observe narrow lines (blue points) or narrow P Cygni profiles (orange points). The ordinate represents the measured velocity while the abscissa is a lower limit on the radius of the material. The unshocked material must be located above the ejecta for which we assume a fiducial mean velocity of 10^4 km s^{-1} . For four SNe, black lines indicate that broad features from shocked material or the SN ejecta emerge, which means that the unshocked CSM is no longer optically thick. All other SNe are still optically thick at the time when the last spectrum is obtained and the shaded area represents a lower limit on the extension of the CSM.

width of the broad hydrogen features observed in the late-time spectra of SN 2018kag, SN 2019cmy, and SN 2019zrk. To estimate out to which radius the ejecta have approximately expanded we multiply this velocity by the time since the explosion. The resulting distances and CSM velocities are represented by the data in Fig. 11, where blue points indicate velocities measured from the line width while orange points indicate the velocities of narrow P Cygni profiles. We emphasize that both the radii and velocities are rough estimates.

For most SNe, the data points are located below or to the right of the gray shaded region which indicates the location of the CSM produced during the observed precursor. This implies that the material ejected during the precursor cannot account for the observed narrow emission lines, because it would be located at smaller radii if it propagates with the observed velocity. Instead, it is more likely that the emission lines are produced by slow-moving material that was expelled earlier. This conclusion is exclusively based on the distance out to which the SN ejecta have expanded at a certain time and is therefore also valid for aspherical CSM distributions (see, e.g., Soumagnac et al. 2020), as long as the SN ejecta expand with an average velocity of at least 10^4 km s^{-1} in all directions. The only SNe for which the narrow features might originate from CSM produced during the precursor are SN 2019uo, SN 2019mom (material from the early precursor), SN 2019aafe, SN 2019vkl, and SN 2020edh. In all other cases, material ejected during the precursor is swept up quickly if it has a low velocity or, if it is faster, it cannot account for the low line velocities.

The measured CSM velocities and the lower limits on the radius allow us to roughly estimate when the material that produces the narrow lines was ejected. In half of the spectra we see matter that was presumably ejected at least 1 yr before the explosion, while 10% of the spectra show signatures of material ejected 2.5 yr or more before the SN. Additional material could be ejected earlier and the resulting CSM shells at larger distances can lead to rebrightenings or bumps in the SN light curve, as observed for example in SN 2009ip (Margutti et al. 2013), PTF 10tel (Ofek et al. 2013b), or iPTF 13z (Nyholm et al. 2017).

We typically observe similar line velocities in spectra of the same SNe, perhaps with the exceptions of SN 2018kag and SN 2019uo, where the scatter is larger. One explanation is that the narrow lines are produced by the same material that is located at a large radius above the photosphere. Another option is that progenitor stars eject material with a characteristic velocity (see, e.g., Owocki et al. 2019, who find an equipartition between the gravitational and kinetic energy of material ejected from the surface of an LBV). If the CSM velocities are indeed determined by the surface gravity of the progenitor stars, the escape velocity of the progenitor of SN 2019cmy (and maybe SN 2020dcs and SN 2020dfh,

for which we only have lower resolution spectra) are relatively low as shown in Table 6. The escape velocity is determined by the stellar mass and radius, and is given by $v_{\text{escape}} = (2 \times G \times M/R)^{0.5}$. For a stellar mass of $30 M_{\odot}$, the stars would have large radii of 300 to $500 R_{\odot}$. The highest escape velocities are observed for SN 2019uo, SN 2019fmb, and SN 2019aafe, which would yield radii of only $10 R_{\odot}$ to $15 R_{\odot}$, again assuming a stellar mass of $30 M_{\odot}$. Especially for the Type Ibn SN 2019uo, this interpretation seems appropriate: the star has already stripped its hydrogen envelope and is therefore likely much more compact than a typical LBV star.

For four SNe, broad emission lines or broad P Cygni features become visible a few weeks or months after the SN explosion. This suggests that the ejecta have reached the radius where the CSM is optically thin. The corresponding radii are marked by black lines for SN 2019uo, SN 2019cmy, SN 2019aafe, and SN 2019zrk. The late-time spectra of the first three SNe continue to exhibit narrow features on top of the broad line, indicating that unshocked, optically thin material is still located above the ejecta. Spectroscopic monitoring of SN 2019zrk continued and about one month after the broad features first emerged, it turned into a Type II SN without any narrow components (as will be described by Fransson et al., in prep.). For all other SNe, the CSM is still optically thick at the time when the last spectrum was obtained, meaning that the dense CSM extends to larger radii as indicated by the shaded area.

We conclude that the material ejected during the observed precursors typically cannot account for the narrow emission lines in the SN spectra (see also Moriya et al. 2014). The narrow lines that are observed while the SN is bright are instead produced by slow-moving material ejected years before the observed precursors and SN explosion.

5.2. Correlations with SN Properties

Here, we test whether the observed precursors increase the SN peak brightness or prolong the rise time. Ofek et al. (2014a) found several marginally significant and weak correlations between the CSM mass estimate and the SN peak luminosity, rise time, and SN energy. All of these correlations are based on a small sample of precursors and require confirmation. Figure 12 shows all SNe with and without precursors and their peak magnitudes. Precursors are detected for many nearby, faint Type II_n SNe, but not for nearby SNe of other types with the exception of the Type Ibn SN 2019uo. Bright precursors are rare, as demonstrated in Sec. 4, so fewer precursors are detected for distant SNe. The correlation between the redshift and the SN luminosity in Fig. 12 is due to the Malmquist bias (Malmquist 1922), which describes that faint objects are undetectable at large distances.

To quantify whether SNe with precursors of any luminosity tend to be more luminous, we calculate a partial correlation




Figure 12. SN peak magnitudes for SNe with and without precursors. We use the brightest detection in the g , r , or i band as a proxy for the peak magnitude. The apparent correlation between the redshift and peak magnitude is caused by the Malmquist bias (Malmquist 1922). Most precursors are observed for relatively nearby SNe as they are faint. We find that SNe with detected precursors are not significantly more luminous than the complete sample. We do not apply K corrections.

between the SN peak magnitude and an array which specifies whether or not a precursor is detected. The distance modulus is used as a control variable to correct for the impact of the Malmquist bias. The distance modulus is chosen rather than the redshift or distance, because it is proportional to the apparent SN magnitude and hence to the detection probability. The partial correlation is calculated for 116 Type IIn SNe with r -band pre-explosion observations and with measured peak magnitudes, and we find a Pearson correlation coefficient of 0.06 which corresponds to a p -value of 0.53. We thus do not detect a correlation between the SN peak magnitude and the detection of a precursor in our search. This might indicate that both groups of SNe have massive CSM shells.

In Sec. 4.3 we found indications that luminous SNe tend to have more luminous precursors (see Fig. 9). To quantify the significance of this observation we calculate a partial correlation between the precursor and the SN peak luminosity while again using the distance modulus as a control variable. The calculation is done for the r -band precursor magnitudes of 12 SNe that have long-lasting precursors in the final three months before the SN explosion, the same objects that contribute to the rate measurement in the right-hand panel of Fig. 9. We calculate the Spearman rank coefficient, which measures whether brighter precursors are associated with brighter SNe without assuming a linear relation between the two luminosities. We find a positive correlation with a correlation coefficient of 0.84 and a p -value of 0.001 or a significance of 3.1σ . We can hence confirm that more luminous precursors occur prior to more luminous SNe even after

correcting for the impact of the Malmquist bias. There could be many possible explanations for the detected correlation: A more massive CSM might boost both the luminosity of the precursor and of the SN, or an energetic precursor could create a more massive CSM shell which results in stronger interaction and a more luminous SN (see also the simple exploration by Ofek et al. 2014a). Alternatively, stars with certain properties might produce more violent precursors and more energetic SN explosions. We also caution that less luminous SNe tend to be nearby while more luminous SNe are located at larger redshifts. While we corrected for the fact that distant SNe appear fainter on Earth, there could be many other differences between these objects which all might play a role.

A massive, optically thick CSM slows down the rise of the SN light curve because the photons diffuse out to the photosphere, and we thus expect a correlation between the SN rise time and the estimated precursor mass. To estimate the diffusion time we inspect the SN light curves in 1-day bins and quantify how many days it takes the light curve to reach its peak from a flux level that is 1.086 mag lower, which corresponds to a luminosity increase by a factor of e . This is done in the r band if available and in the g band for SN 2018gho. The g band typically peaks earlier than the r band, so the numbers might not be directly comparable. The rise time should be measured for the bolometric light curve, which typically rises more quickly as a large part of the energy is emitted in the UV. Our estimates are hence upper limits on the actual rise time. The estimated rise times are given in the penultimate column in Table 4, if the rise and peak are well observed.

With rise times of 4–25 days, all SNe with detected precursors are part of the fast rising subgroup identified by Nyholm et al. (2020)⁶ which includes approximately two thirds of the Type IIn SNe in their sample. A likely explanation is that fast rising SNe IIn are typically fainter (Nyholm et al. 2020) and our precursor search is most sensitive to nearby, faint SNe (see Fig. 12). We here assume that energetic precursors eject more material and search for a correlation between the radiative precursor energy and the estimated SN rise time. With a Pearson correlation coefficient -0.37 with a p -value of 0.25, no significant correlation is found. It is either possible that a relation is washed out owing to the large uncertainties on both quantities or the observed precursors do not dominate the total CSM mass.

In conclusion, we do observe that more luminous precursors are detected prior to more luminous SNe even after cor-

⁶ While we define the rise time as the time it takes the SN to rise by a factor of e to its peak, Nyholm et al. (2020) determine the rise time from a power-law fit. Thus, the quoted numbers might not be directly comparable, but the bottom panel of their Fig. 7 allows us to read off rise times that are consistent with our definition.

recting for the impact of the SN distance. However, SNe with detected precursors are not systematically more luminous than SNe without detected precursors, and we do not measure a correlation between the precursor energy and the SN rise time. Thus, we do not find that the observed precursors have a major impact on the SN light curve. This seems to fit with our results from Sec. 5.1, where we find that the narrow hydrogen features in the spectra of Type II_n SNe typically do not originate from material ejected during the observed precursor. Together, both results might indicate that a large fraction of the CSM is ejected in earlier mass-loss events that we do not detect here.

5.3. The Type Ib_n SN 2019uo

One of the precursors discovered in this study occurs prior to SN 2019uo, making it the second Type Ib_n with a detected precursor after SN 2006jc (Pastorello et al. 2007; Foley et al. 2007). The coadded difference images in Appendix A confirm that a point source is visible ~ 320 days before the explosion of SN 2019uo. As shown in Fig. 13, the precursor starts ~ 340 days before the explosion of SN 2019uo and is observed over 35 days (see also Fig. 3 and Table 4). In addition to the 5σ detection, another two data points surpass the 3σ threshold when using 7-day bins, and in total 24 individual images contribute to the detection. The mean precursor magnitude is -13 , making it one of the faintest precursors detected in this search. The observed radiative energy in the r band is 1.7×10^{47} ergs. However, the observations shortly before and after the precursor are not very constraining, so it could last longer.

With a redshift of 0.020, SN 2019uo is the closest Type Ib_n SN out of the 12 objects in our sample. It was classified as a Type Ib_n based on a spectrum showing narrow helium P Cygni features that was obtained 9 days after the estimated explosion date (Fremming et al. 2019). SN 2019uo was studied by Gangopadhyay et al. (2020) in detail, and they find that SN 2019uo is slightly underluminous and evolves quickly, typical for Type Ib_n SNe. They model the bolometric light curve with CSM interaction and their best-fitting models require the presence of $0.4\text{--}0.7 M_{\odot}$ of material located at a radius of $\gtrsim (0.2\text{--}2) \times 10^{15}$ cm.

SN 2019uo is one of the few objects for which the narrow P Cygni profiles in the spectra might originate from the CSM ejected during the observed precursor (see Fig. 11). We measure typical velocities of $\sim 900 \text{ km s}^{-1}$ (see Table 6), which would imply that the CSM has propagated to a radius of $\sim 2 \times 10^{15}$ cm when the SN explodes 320 days after the precursor. This radius is approximately consistent with the inner CSM radius required by the modelling of Gangopadhyay et al. (2020). Moreover, it roughly coincides with the radius at which the CSM turns optically thin shown in Fig. 11: broad features, likely from the shocked CSM, first appear in a




Figure 13. Light curve of SN 2019uo in 7-day bins before the estimated explosion date (JD 2,458,501.3) and in 1-day bins afterward. Solid data points with black edges are 5σ detections, transparently colored data points have significances larger than 3σ , and upper limits are less significant. Green, red, and black data points correspond to observations in the g , r , and i bands, respectively. A line connects the three detections that we consider to be part of the precursor. The 3σ detection at ~ 30 days could be a statistical fluctuation and is not discussed here.

spectrum obtained 18 days after the explosion date (see Fig. 3 by Gangopadhyay et al. 2020). A CSM shell located at a radius of 2×10^{15} cm is swept up by the SN ejecta if they have an average velocity of $13,000 \text{ km s}^{-1}$. Narrow helium features remain visible in later spectra and might originate from an optically thin wind located above the CSM. The same material could also produce the flash-ionization features detected by Gangopadhyay et al. (2020) before the SN reaches its peak.

The observed precursor energy can be produced via interaction if the mass lost during the precursor is $\gtrsim 0.007 M_{\odot}$ (see Fig. 14 or Table 4). However, the SN light curve modelled by Gangopadhyay et al. (2020) requires a CSM mass a factor of 50–100 times larger. A possible explanation is that the material ejected during the precursor propagates through a low-density environment such that only 1–2% of its kinetic energy is converted to radiation in the r band. This would imply that the progenitor star was not surrounded by a massive CSM before the observed precursor. Alternatively, only a small fraction of the CSM is emitted during the observed precursor and the rest of the required material is expelled during earlier or fainter precursors which we do not detect. In this case the radiative efficiency would be larger. We conclude that a relatively simple picture can explain the characteristics of the precursor and subsequent SN: the observed precursor could produce the complete CSM of $\sim 0.5 M_{\odot}$ if its radiative efficiency is low with $\epsilon \approx 1\%$. The resulting dense CSM shell is confined to a radius of $\sim 2 \times 10^{15}$ cm and can account for the SN bolometric light curve (shown by Gangopadhyay et al. 2020) as well as for the spectroscopic development.

Until now, the only precursor observed prior to a Type Ibn SN was observed for SN 2006jc (Pastorello et al. 2007; Foley et al. 2007). The 9-day-long precursor with a peak magnitude of -14.1 was detected 2 yr before the SN explosion. The two precursors are hence similar to each other, as they are both relatively faint and happen hundreds of days before the SN explosion. In both cases, the CSM is helium-rich, so we do not witness the stripping of the hydrogen envelope. Based on 11 Type Ibn SNe with pre-explosion data in the r band, we show in Fig. 10 and Table 4 that precursors with an r -band magnitude of < -16 happen $< 1.0\%$ of the time (95% confidence level), while faint precursors with magnitudes brighter than -13 might happen up to 31% of the time. These limits are based on observations collected up to 2.5 yr before the SN explosion and the median observation time is 8.6 months before the explosion. We thus do not have very strong constraints on faint outbursts and they might be relatively common.

Thus, we confirm that the progenitor stars of some Type Ibn SNe are able to produce relatively bright flares in the last years before their explosion. Except for the two precursor detections, no Type Ibn SN progenitor has been identified in archival observations and their nature is debated. The most commonly suggested progenitors are Wolf-Rayet stars that have shed their hydrogen envelopes or massive stars that are stripped by a binary partner (see, e.g., Smith 2017; Hosseinzadeh et al. 2017). Alternatively, the progenitors of some Type Ibn SNe were hypothesized to be white dwarf binaries (Sanders et al. 2013; Hosseinzadeh et al. 2019), or very massive stars that undergo pulsational pair instability events (Woosley 2017; Karamahmetoglu et al. 2019) which would imply that no core collapse occurs and the star is likely still present. Contrary to LBVs, classical Wolf-Rayet stars are not known to undergo giant eruptions, so bright flares cannot be common during the lifetime of the star. There is an intermediate class of Ofpe/WN9 stars which have stripped most of their hydrogen envelope, but undergo LBV-like outbursts (see, e.g., Smith et al. 2020). Such stars have been suggested as progenitors for two Type Ibn SNe that also show relatively strong hydrogen lines (Smith et al. 2012; Kool et al. 2020). However, the hydrogen features in the spectra of SN 2006jc are much weaker and SN 2019uo might not show any hydrogen. It is therefore unclear under which conditions stripped-envelope stars can produce as bright eruptions. If such flares are related to the late stages of nuclear burning (see Sec. 6.2), they would only occur shortly before the SN explosion, which would explain why no such flares are observed for Wolf-Rayet stars in the Milky Way or in nearby galaxies.

6. NATURE OF THE PRECURSORS

We emphasize that the nature of the observed transients is ambiguous. Owing to the optically thick CSM, we cannot directly observe the expanding SN ejecta, but only see light diffusing out through the photosphere. We therefore cannot determine whether core collapse occurred or at what time it happened (see, e.g., Moriya 2015; Tartaglia et al. 2016; Woosley 2017 for possible scenarios). It is conceivable that some of the precursors are already part of the SN light curve, rather than preceding the explosion. Nevertheless, in the following we adopt the interpretation that core collapse occurs shortly before the SN rises to its main peak and that the precursors are produced by the progenitor star before it explodes.

In Sec. 6.1 we explore whether the precursor luminosity could be produced via interaction or by a continuum wind. The underlying energy source is unknown and suggested mechanisms include unstable nuclear burning phases (Smith & Arnett 2014), shell burning (Arnett & Meakin 2011a,b), interaction with a binary companion (Smith & Arnett 2014; McIcley & Soker 2014; Danieli & Soker 2019; Owocki et al. 2019), or reduced gravity due to high neutrino luminosities (Moriya 2014). Specific predictions exist for wave-driven mass loss triggered by instabilities during the neon and oxygen burning phases, and we compare our observations to the model described by Shiode & Quataert (2014) in Sec. 6.2.

6.1. What Powers the Precursor Luminosity?

In this section we explore possible mechanisms that could produce the observed precursor luminosity. First, we point out in Sec. 6.1.1 that the long precursor durations likely require a persistent energy source. This could, for example, be interaction of the ejected material with pre-existing CSM (described in Sec. 6.1.2) or a brightening of the star (discussed in Sec. 6.1.3). In Sec. 6.1.4, we measure the SN rise times to derive upper limits on the CSM mass.

6.1.1. Precursor Durations

The densely sampled light curves of the ZTF survey allow us to measure the durations for a sample of precursors. As shown in Fig. 5, the outbursts are typically observed over several months. Their true durations are likely even longer, if the fainter parts of the precursors remain undetected. With rise times of only ~ 1 mag over ~ 50 – 100 days, most precursors develop much more slowly than the subsequent SNe. This implies that the diffusion time in the CSM does not dominate the precursor duration. Consequently, the precursors are likely not powered by a single short-lived eruption from the stellar surface, but require a long-lived energy source, such as ongoing interaction, a stellar wind, potentially a series of explosions, or maybe a short-lived event deep within the stellar envelope, where the diffusion time is much larger than within the CSM. We thus conclude that the long duration of several detected precursors is likely intrinsic and not due to diffusion.

6.1.2. Interaction-Powered Precursors

The light curves of Type II_n SNe are mainly powered by interaction between the ejecta and the circumstellar material surrounding the star. It thus might seem logical that the same is true for the precursors. In this scenario, a large amount of material is ejected from the stellar surface and a fraction of the kinetic energy $\epsilon < 1$ is converted to radiative energy when the ejected material is slowed down by pre-existing CSM. If the velocity of the newly ejected material is known, its mass can be estimated using

$$E_{\text{rad}} = \epsilon E_{\text{kin}} = \epsilon \frac{1}{2} M_{\text{CSM, pre.}} v_{\text{CSM, pre.}}^2, \quad (1)$$

where $M_{\text{CSM, pre.}}$ and $v_{\text{CSM, pre.}}$ are respectively the mass and velocity of the material ejected during the precursor.

Equation 1 shows that the required CSM mass strongly depends on the velocity. CSM velocities can be estimated from the narrow hydrogen (or helium) features, however; as demonstrated in Sec. 5.1, these lines are likely produced by material ejected at earlier times. It is hence possible that the matter expelled during the observed precursors has larger velocities which would result in lower CSM mass estimates. The observed velocities and corresponding CSM masses are given in the fourth and third to last columns of Table 4 (the quoted velocity is the median of all spectra given in Table 6). The required CSM masses for different velocities are illustrated in Fig. 14 for some of the precursors.

Most observed precursors develop quite slowly as pointed out in Sec. 6.1.1. Nyholm et al. (2020) measured the rise and decline rates for ~ 30 Type II_n SNe from the PTF/iPTF sample and the precursors we detect rise more slowly than any Type II_n SNe in their sample. A possible explanation could be that the precursor shock front propagates with a substantially lower velocity compared to the SN ejecta. As a consequence, the rate at which CSM is swept up is lower and kinetic energy is converted to radiation more slowly. According to the model by Svirski et al. (2012) (see also Ofek et al. 2014b), the radiated luminosity produced during CSM interaction is $L_{\text{pre.}} \propto r^2 \rho v_{\text{CSM, pre.}}^3$, which simplifies to $L \propto v_{\text{CSM, pre.}}^3$ for a wind-like density profile with $\rho \propto r^{-2}$. The luminosity difference between precursors and SNe on the order of 100 suggests that the average shock velocity is a factor of ~ 5 lower (i.e., $v_{\text{CSM, pre.}} \approx 2000 \text{ km s}^{-1}$ compared to $\sim 10^4$ for SN ejecta). However, this assumes that the radiative efficiency ϵ (i.e., the fraction of kinetic energy that is converted to radiation) is similar for the precursor and the SN. These higher velocities would reduce the amount of ejected material by about one order of magnitude as shown in Fig. 14.

Interaction stops either when the shock reaches the edge of the CSM or when it is slowed down, which happens when the mass of the swept-up CSM is comparable to the mass in the




Figure 14. CSM masses required to produce the observed precursor energies, if they are powered by interaction. The black contour lines show the amount of CSM, $M_{\text{CSM, pre.}}$ in solar masses, ejected during the precursor if the kinetic energy is completely converted to radiation (i.e., $\epsilon = 1$). The observed precursor energies (taken from Table 4) are indicated by vertical lines. They are lower limits on the actual radiative energy as we only detect the brightest part of the precursors and because we did not apply a bolometric correction when calculating the energy. CSM velocities measured from the spectra are marked by dots, but they do not necessarily correspond to the velocities of the precursor material. The gray horizontal line shows the approximate amount of CSM that a continuum-driven wind would eject (see Sec. 6.1.3).

shock front. Several precursors (e.g., prior to SN 2019vkl, SN 2019aafe, or SN 2019zrk) continue to rise for more than 100 days. If their luminosity is dominated by interaction, the associated mass loss must be substantial, such that the ejected material is not slowed down considerably within this time (i.e., $\epsilon \ll 1$). If it expands with an average velocity of $2,000 \text{ km s}^{-1}$ it reaches a radius of $\sim 2 \times 10^{15} \text{ cm}$ within 100 days. Several other precursors (e.g., prior to SN 2019fmb, SN 2019gjs, or SN 2020dcs) fade in the weeks to months before the SN explosion, as was also observed for the 2012a event prior to the final explosion of SN 2009ip. This might indicate that the material has slowed down or that it only collides with a thin CSM shell and then continues to expand through a lower density environment. This qualitative description assumes that each precursor is associated with a single short-lived mass-loss event. It is, however, also possible that material is ejected in a series of eruptions from the stellar surface. If the energy of these eruptions changes with time, this could account for both rising or falling precursor light curves.

As noted in Sec. 3.2 precursors with detections in both the g and r band appear rather red compared to young Type II_n SNe. We suggest that these low temperatures are a consequence of the low precursor luminosity in combination with a relatively large photospheric radius due to the extended CSM.

We conclude that the observed precursors could be powered by interaction if the star undergoes major mass-loss events. The precursor luminosities require that the ejected material has velocities on the order of $\sim 2,000 \text{ km s}^{-1}$. Such low-velocity shock fronts could explain the long duration of the precursors compared to the SN light curve. Figure 14 shows that the most energetic precursor requires the ejection of $\sim 0.3 M_{\odot}$ of material for a CSM velocity of $2,000 \text{ km s}^{-1}$. However, a more realistic radiative efficiency of $\epsilon \lesssim 0.3$ would bring the required mass back to $\sim 1 M_{\odot}$. CSM envelopes of several solar masses have been observed for some Type II_n SNe, so high-mass estimates are not necessarily unrealistic, and for less energetic precursors the required masses are lower by a factor of up to 100.

6.1.3. Wind-Driven Precursors

In an alternative scenario the observed luminosity originates from the star itself. Shaviv (2001a) showed that the outer part of the stellar envelope (from the radius out to which convection is efficient to the “hydrostatic surface” of the star) becomes unstable when the stellar luminosity approaches the Eddington limit. Local density differences reduce the effective opacity of the star, so it remains quasistable even when it exceeds the Eddington luminosity. However, the instabilities only have an effect as long as the atmosphere is optically thick over a scale height. An optically thick continuum-driven wind is therefore accelerated from this region, with the actual photosphere sitting farther out. The resulting mass loss is smaller than can be expected without the lower effective opacity. The resulting wind has a typical velocity of $(L/L_{\text{Eddington}})^{0.5} v_{\text{escape}}$ (Shaviv 2001b) — that is, usually $v_{\text{CSM, pre.}} \gtrsim 1000 \text{ km s}^{-1}$, which is larger than most line velocities measured in Sec. 5.1.

Contrary to the interaction scenario a fraction of the radiative energy is converted into kinetic energy (i.e., $\epsilon > 1$). The mass loss can be calculated using

$$E_{\text{rad}} = \frac{1}{W} M_{\text{CSM, pre.}} c_s c = \epsilon E_{\text{kin.}} \quad , \quad (2)$$

where $W \approx 5$ is an empirical factor, $c_s \approx 60 \text{ km s}^{-1}$ is the speed of sound at the base of the optically thick wind (Shaviv 2000, 2001b), and c is the speed of light. We note that the results of Eq. 2 are equal to Eq. 1 for a CSM velocity of $v_{\text{CSM, pre.}} = 2,700 \text{ km s}^{-1}$ and $\epsilon = 1$. As a result the mass loss is lower by a factor of ~ 10 compared to the numbers given in the third to last column of Table 4. The most energetic precursor with a radiative energy of 10^{49} ergs only results in a mass loss of $0.1 M_{\odot}$ as shown in Fig. 14.

Once the wind reaches an equilibrium state, it forms a photosphere. Owocki & Shaviv (2016) show that a wide range of mass-loss rates and wind velocities result in photospheric temperatures between 5,000 K and 6,000 K, the temperature

at which hydrogen recombines and the opacity drops. These temperatures are consistent with most observed precursor temperatures in Sec. 3.3. Larger temperatures are expected before the wind reaches its equilibrium state, but lower temperatures are more difficult to explain. We note that the precursor temperatures are also similar to the temperature of the giant eruption of η Carina (Rest et al. 2012).

6.1.4. Constraints on the CSM Mass

We also estimate upper limits on the total CSM mass (here material located within $5 \times 10^{15} \text{ cm}$) based on the SN rise time. A massive, optically thick CSM slows down the rise of the SN light curve, because the photons diffuse out to the photosphere. Therefore, a quickly rising SN is inconsistent with a large CSM mass, while a slow rise could be intrinsic and does not necessarily imply a massive CSM. SN rise times were estimated in Sec. 5.2 and are given in Table 4. For an infinite wind-like CSM profile (i.e., with a density that decreases with the radius like $\rho \propto r^{-2}$), the diffusion time t_{diff} is given by

$$t_{\text{diff.}} \approx \frac{\kappa K}{c} (\ln(c/v_{\text{shock}}) - 1) \quad , \quad (3)$$

where κ is the opacity, v_{shock} is the velocity of the shock front, and the mass-loading factor K is defined as $K = \dot{M}/(4\pi v_{\text{CSM}})$ for the mass-loss rate \dot{M} and the CSM velocity v_{CSM} (Ginzburg & Balberg 2012).

The CSM mass between two radii R_{inner} and R_{outer} can be obtained by integrating the CSM density profile:

$$\begin{aligned} M_{\text{CSM, diff.}} &= \int_{R_{\text{inner}}}^{R_{\text{outer}}} 4\pi r^2 K r^{-2} dr = 4\pi K (R_{\text{outer}} - R_{\text{inner}}) \\ &\approx 4\pi R_{\text{outer}} \frac{t_{\text{diff}} c}{\kappa (\ln(c/v_{\text{shock}}) - 1)} \\ &\approx 0.10 R_{\text{outer, } 5e15 \text{ cm}} t_{\text{diff, 1d}} M_{\odot} \quad . \end{aligned} \quad (4)$$

In the second line, we assume that the inner radius is much smaller than the outer one and insert Eq. 3. We then adopt a typical value for $\kappa = 0.34 \text{ cm}^2 \text{ g}^{-1}$, appropriate for a medium that consists of 70% hydrogen (see, e.g., Ginzburg & Balberg 2012; Ofek et al. 2013a) and a shock velocity of $v_{\text{shock}} = 10^4 \text{ km s}^{-1}$. The CSM mass is hence proportional to the diffusion time as well as to the outer radius of the CSM.

We integrate the CSM density out to a radius of $5 \times 10^{15} \text{ cm}$, which is approximately consistent with the observed spectral evolution shown in Fig. 11. The resulting upper limits on the CSM masses are given in the last column of Table 4. While many of the mass upper limits are consistent with the mass estimates for interaction-powered precursors, there are discrepancies for a few objects (e.g., SN 2019zrk, SN 2019gjs, or SN 2018gho). We emphasize that a difference of a factor of a few does not necessarily imply an inconsistency, as both CSM masses are order-of-magnitude estimates owing to the assumed wind-like density profile, the adopted shock

velocity, and the outer CSM radius of 5×10^{15} cm. Furthermore, the diffusion time might be different for nonspherical CSM geometries (Soumagnac et al. 2020). Another alternative would be considerably higher CSM velocities of a few $\times 1,000 \text{ km s}^{-1}$ (see Sec. 6.1.2). We summarize that several SNe, such as SN 2019zrk, rise quickly even though they experienced powerful precursor eruptions shortly before the explosion. This might imply that the precursors are not associated with mass-loss events of several solar masses. Careful modeling of a well-observed SN with an energetic precursor would be required to establish whether fast SN rise times can be reconciled with extensive mass-loss episodes in the last months before the explosion.

6.2. Can Wave-Driven Mass Loss Trigger the Detected Precursors?

Unbinding a solar mass of material requires a substantial energy deposition in the stellar envelope. The fact that the precursor rate increases in the last months before the SN explosion could imply that the precursors are associated with late nuclear burning stages which last from a few years to a few months (see, e.g., Shiode & Quataert 2014).

When fusing carbon or heavier elements, both the energy production and neutrino cooling rates increase dramatically in the stellar core (see, e.g., Woosley et al. 2002; Arnett & Meakin 2011a; Quataert & Shiode 2012). While the two processes are in equilibrium on average, local imbalances cause vigorous convection within the core and Shiode & Quataert (2014) estimate that $\lesssim 10\%$ of the fusion energy is carried by convection. The convection excites gravity waves which typically remain confined to the core (Meakin & Arnett 2006; Quataert & Shiode 2012) and do not affect the envelopes of most massive stars. However, the internal structure of some progenitor stars may allow part of the wave energy to tunnel out of the core and excite acoustic waves in the stellar envelope. The energy deposited in the envelope can be as large as a few times $10^{40} \text{ ergs s}^{-1}$ or $\sim 10^{47} \text{ ergs}$ over a year. It might trigger strong adiabatic mass-loss events or inflate the stellar envelope (Quataert & Shiode 2012; Shiode & Quataert 2014; Quataert et al. 2016; Fuller 2017; Fuller & Ro 2018).

The neon and oxygen burning phases last a few years to months with more massive cores burning out more quickly (see, e.g., Shiode & Quataert 2014). Silicon fusion only occurs in the last few days to hours before the explosion and therefore cannot account for the observed months-long precursors. Shiode & Quataert (2014) modelled different progenitor stars with the *MESA* stellar code (Paxton et al. 2011) and found that it takes the waves about several weeks to about a year to reach the stellar surface and that progenitor stars with a wide range of initial conditions can fulfill the requirements for matter outflow. We observe that the brightest precursors start several months before the SN explosion. They

hence might be powered by energy produced during the neon or oxygen burning phases. According to the model by Shiode & Quataert (2014), the time of the precursor eruption is inversely related to the core mass. A precursor that occurs one month before the SN explosion would imply a helium core mass of $\lesssim 15 M_{\odot}$, while a precursor one year before the explosion requires a lower core mass of $\lesssim 5 M_{\odot}$.

The most energetic precursor, detected prior to the explosion of SN 2019aafe, released an *r*-band energy of 10^{49} ergs over 100 days. If this luminosity is interaction-powered, the kinetic energy would be even larger. For progenitors with $\sim 15 M_{\odot}$ helium cores, Shiode & Quataert (2014) calculate wave energies of up to a few times 10^{47} ergs, about 50 times lower than the observed radiative energy. Ten times higher energies could be reached by stars with $\gtrsim 30 M_{\odot}$ helium cores, but their oxygen burning phase only lasts for about a month, such that the produced waves only reach the surface days before the SN explosion (similar precursor energies and timescales were calculated by Wu & Fuller 2020). Fusing $\sim 1 M_{\odot}$ of material releases an energy of $\sim 10^{51}$ ergs during the neon and oxygen burning phases each (Quataert & Shiode 2012), so the wave transport would have to be extremely efficient to produce as energetic precursors as observed. The precursors observed prior to other SNe, such as SN 2019fmb or SN 2020dcs, are at least 10 times less energetic and are more easily explained by the model.

We summarize that wave-driven mass loss, powered by instabilities during the neon and oxygen burning phases, could explain why precursor eruptions occur in the last few months before the SN explosion. According to the model by Shiode & Quataert (2014), less energetic early precursors are produced by lower mass stars, while stars with massive cores produce more powerful precursors that occur only days before the explosion. This could at least qualitatively explain our observation that brighter precursors become more common in the last months before the SN explosion (see Sec. 4.2). However, the brightest detected precursor is about two orders of magnitude more energetic than predicted by Shiode & Quataert (2014), and it is unclear whether the model could account for such events. We emphasize that our observations cannot confirm that wave-driven mass loss triggers precursor eruptions, and that other mechanisms such as interaction with a binary companion star might also be able to explain the observations (see, e.g., Quataert et al. 2016; Owocki et al. 2019).

7. CONCLUSIONS

One main finding of this study is that bright precursors are relatively common immediately before the explosion of Type II_n SNe (see Fig. 5) and that most of them last for one or several months. Long-lasting precursors that are brighter than -13 mag in the *r* band are observed immediately prior to

the explosion of about 25% of all Type II_n SNe (with a 95% confidence range of 5–69%). Some of the brightest precursors are better described as a continuous brightening rather than a discrete flare (see, e.g., the precursor light curves prior to SN 2019zrk or SN 2019aafe in Fig. 5). The most powerful precursor found here releases an energy of 10^{49} ergs over 100 days ($\sim 10\%$ of the radiative energy released in a typical SN explosion), but most precursors are an order of magnitude less energetic.

All precursors are much brighter than the Eddington luminosity of a massive star, such that they likely involve extensive mass-loss events. The mass of the ejected CSM is difficult to quantify: if the kinetic energy of the CSM is similar to the radiative energy of the precursor, the mass loss would typically amount to one or a few solar masses with large uncertainties owing to the unknown CSM velocity. For wind-driven precursors, the expected mass loss is typically ten times lower. Several SNe with detected precursors rise to their peak luminosity within a few days, which might imply that the mass of their CSM shell is typically $\lesssim 1 M_{\odot}$.

To our knowledge, such bright and long-lasting precursors have so far only been detected prior to Type II_n SNe and precursors prior to other SNe are typically less energetic. Nevertheless, material ejected during these events cannot account for the characteristic narrow hydrogen emission lines in the spectra of these SNe. Within the short time before the SN explosion the ejected material cannot expand to radii larger than 10^{14} cm, if its velocity is as low as the line widths indicate. Hydrogen emission lines are, on the other hand, observed for many weeks or even years and are thus produced by material located at larger radii. For the average spectrum, this material must have been ejected at least ~ 1 yr before the explosion. We only find few, rather faint and short, precursors at these times. This might indicate that earlier mass-loss events are likely substantially fainter than the observed precursors and therefore remain undetected. SNe with precursors are not significantly brighter than SNe without observed precursors at similar redshifts, which likely supports our hypothesis that the precursors detected immediately before the explosion do not dominate the total CSM mass.

The Type Ib_n SN 2019uo is an exception: the observed precursor might have ejected $\sim 0.5 M_{\odot}$ of helium ~ 320 days before the SN explosion. The low precursor luminosity is expected if the material propagates through a low-density environment. If the material propagates with the CSM velocity observed in the spectra it would remain confined to a rather small radius. This would explain both the fast light-curve evolution and the appearance of broad spectral features only 18 days after the explosion that indicate that the ejecta have reached the edge of the optically thick CSM. It is hence possible that the complete CSM of SN 2019uo is created during the observed precursor, while the spectra and light curves of

the Type II_n SNe require earlier mass-loss events (see also Moriya & Maeda 2016).

Prior to Type II_n SNe, we only detect five precursors that happen more than three months before core collapse (at phases of 700 to 180 days before the explosion). At these earlier times the rate (or duration) of bright precursors with r -band magnitudes < -16 is lower by a factor of more than 10 and the rate of precursors brighter than magnitude -13 is a factor of 5 lower. The increasing rate of bright precursors could be explained if the precursors are powered by wave-driven mass loss triggered by instabilities during the neon and oxygen burning phases. Shiode & Quataert (2014) argue that stars with lower core masses undergo fainter precursors about a year before the SN explosion, while more massive stellar cores produce brighter flares only weeks or days before core collapse, because of their shorter nuclear burning phases. However, the most energetic precursor is 100 times more energetic than predicted by Shiode & Quataert (2014), and it is not clear whether energy transport to the stellar envelope could be as efficient.

The bright outbursts shortly before the SN explosion open the door to the possibility of predicting SN explosions. Four of the precursors reach an apparent magnitude of < 20 (see Figs. 3 and 4) and are potentially detectable with the ZTF discovery pipeline. Based on these numbers, we estimate that 1–2 precursors per year are bright enough to allow the prediction of an imminent SN explosion. Indeed, the precursor prior to SN 2019fmb was reported as a transient by the Pan-STARRS collaboration (Chambers et al. 2019), but it was not realized at the time that this was a pre-explosion outburst. We conclude that the ZTF survey has the potential to predict SN explosions if a dedicated search is implemented.

ACKNOWLEDGEMENTS

We thank A. Nyholm for his comments on the manuscript. This work would not have been possible without the spectroscopic follow-up observations carried out by S. Anand, D. Bektsev, N. Blagorodnova, M. Bulla, S. B. Cenko, W. Chen, P. Chinchilla, R. Clavero Jimenez, C. Cunningham, A. Dahiwal, L. Dominguez, A. J. Drake, C. Frohmaier, F. J. Galindo-Guil, E. Hammerstein, T. Hung, N. Janssen, J. Jencson, R. Karjalainen, H. Ko, M. Kuhn, E. McEwen, A. Miller, S. Moran, M. C. Ramirez-Tannus, A. Smith, E. Swann, K. Teet, J. Vinko, and J. Viuho. We would like to thank participating observers on the UW APO ZTF follow-up team, including Brigitta Spiócz, Eric Bellm, Zach Golkhou, Keaton Bell, and James Davenport. In addition, we thank A. Gangopadhyay, H. Ko, and S. Prentice for reducing optical spectra and for sharing their data.

Based on observations obtained with the 48-inch Samuel Oschin Telescope and the 60-inch Telescope at Palomar Observatory as part of the Zwicky Transient Facility project.

Table 6. SN Spectra

SN	obs. JD	instrument	time after t_0 (days)	line width (km s ⁻¹)	inst. res. (km s ⁻¹)	velocity (km s ⁻¹)	comment
SN 2018eru	2458351.7	P200/DBSP	35.0	1150	130	1143, 1000*	
SN 2018gho	2458372.7	P200/DBSP	6.2	400	130	378	
	2458383.4	LT/SPRAT	16.9	800	830	< 415	
SN 2018kag	2458470.5	LT/SPRAT	4.0	1150	830	796	
	2458502.5	NOT/ALFOSC	35.9	750	640	391	
	2458535.5	WHT/ACAM	68.9	1500	670	1342	broad
SN 2019uo	2458505.0	FTN/FLOYDS	3.7	–	–	990*	flash, He line
	2458506.9	FTN/FLOYDS	5.6	–	–	750*	
	2458508.3	LJT/YFOSC	7.0	–	–	850*	
	2458509.9	P200/DBSP	8.6	–	–	550*	
	2458519.9	FTN/FLOYDS	18.6	–	–	1200*	broad
	2458525.7	NOT/ALFOSC	24.4	–	–	1000*	broad
	2458529.2	LJT/YFOSC	27.9	–	–	1000*	broad
	2458579.9	Keck-1/LRIS	78.6	–	–	1000*	broad
SN 2019bxq	2458596.7	LT/SPRAT	40.8	900	830	< 415	
	2458598.0	P200/DBSP	42.1	330	130	303	
SN 2019cmy	2458572.8	APO/DIS	4.9	400	370	< 185	flash
	2458577.0	Keck-1/LRIS	9.1	350	320	< 160	
	2458580.9	APO/DIS	13.0	400	370	< 185	
	2458637.9	Keck-1/LRIS	70.0	400	320	240	broad
SN 2019iay	2458668.5	Lick 3-m/Kast	11.8	500	300	400	
	2458668.7	P200/DBSP	12.1	300	130	270	
SN 2019gjs	2458705.7	P200/DBSP	15.0	350	130	325	
SN 2019mom	2458705.9	P200/DBSP	14.9	450	130	431, 750*	
SN 2019fmb	2458792.0	APO/DIS	76.2	900	370	820	
	2458793.0	P200/DBSP	77.2	1000	130	992, 1050*	
SN 2019aafe	2458750.9	Keck-1/LRIS	10.1	1100	320	1052	
SN 2019vkl	2458812.1	Lijiang-2.4m/YFOSC	3.4	850	350	775	
SN 2019vts	2458834.8	APO/DIS	17.8	500	370	336	
SN 2019qny	2458853.4	LT/SPRAT	26.3	900	830	< 415	
SN 2020iq	2458898.8	Keck-1/LRIS	66.1	200	320	< 160 ^h	
SN 2019zrk	2458893.8	P200/DBSP	4.8	350	130	325 ^h	
	2458896.9	APO/DIS	7.8	400	370	< 185 ^h	
	2458896.6	LT/SPRAT	7.6	750	830	< 415 ^h	
	2458905.6	NOT/ALFOSC	16.6	800	640	480 ^h	
	2458930.9	Keck-1/LRIS	41.9	500	320	384 ^h	broad
SN 2020dcs	2458897.6	LT/SPRAT	2.7	850	830	< 415	
SN 2020dfh	2458906.9	ESO-NTT/EFOSC2	3.8	1000	980	< 490	
SN 2020edh	2458919.0	FTN/FLOYDS N-SIRAH	4.1	1000	800	600	

NOTE—List of SN spectra with narrow lines or P Cygni features. The typical resolution of the spectrograph (third column from the end) is subtracted from the measured width in quadrature, or we quote an upper limit if the result would be smaller than half the resolution. For most SNe, additional low-resolution spectra were obtained with the SED Machine.

Asterisks (*) mark velocities measured from P Cygni profiles and not from line widths. Spectra marked with an ^h have strong host lines such that the measured velocities are less reliable.

ZTF is supported by the National Science Foundation (NSF) under grant AST-1440341 and a collaboration including Caltech, IPAC, the Weizmann Institute for Science, the Oskar Klein Centre at Stockholm University, the University of Maryland, the University of Washington, Deutsches Elektronen-Synchrotron and Humboldt University, Los Alamos National Laboratories, the TANGO Consortium of Taiwan, the University of Wisconsin at Milwaukee, and the Lawrence Berkeley National Laboratory. Operations are conducted by COO, IPAC, and UW. The SED Machine is based upon work supported by NSF grant 1106171. This work was supported by the GROWTH project funded by the NSF under PIRE grant 1545949. Partially based on observations made with the Nordic Optical Telescope, operated by the Nordic Optical Telescope Scientific Association at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias. Some of the data presented herein were obtained with ALFOSC. Some of the data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and NASA; the observatory was made possible by the generous financial support of the W. M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain. The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with finan-

cial support from the UK Science and Technology Facilities Council. Research at Lick Observatory is partially supported by a generous gift from Google. The ztfquery code was funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement n759194 - USNAC, PI Rigault). E.O.O. is grateful for the support by grants from the Israel Science Foundation, Minerva, Israeli Ministry of Technology and Science, the US-Israel Binational Science Foundation, Weizmann-UK, Weizmann-Yale, and the Weizmann-Caltech grants. A.G.Y.'s research is supported by the EU via ERC grant 725161, the ISF GW excellence center, an IMOS space infrastructure grant and BSF/Transformative and GIF grants, as well as The Benozziyo Endowment Fund for the Advancement of Science, the Deloro Institute for Advanced Research in Space and Optics, The Veronika A. Rabl Physics Discretionary Fund, Paul and Tina Gardner, Yeda-Sela and the WIS-CIT joint research grant; A.G.Y. is the recipient of the Helen and Martin Kimmel Award for Innovative Investigation. N.J.S. is grateful for the support by the ISF (grant 1770/19). A.V.F. acknowledges support from the Christopher R. Redlich Fund, the TABASGO Foundation, and the Miller Institute for Basic Research in Science. L.T. acknowledges support from MIUR (PRIN 2017 grant 20179ZF5KS). R.L. is supported by a Marie Skłodowska-Curie Individual Fellowship within the Horizon 2020 European Union (EU) Framework Programme for Research and Innovation (H2020-MSCA-IF-2017-794467). M.L.G. acknowledges support from the DiRAC Institute in the Department of Astronomy at the University of Washington. The DiRAC Institute is supported through generous gifts from the Charles and Lisa Simonyi Fund for Arts and Sciences, and the Washington Research Foundation.

REFERENCES

- Ahumada, R., Prieto, C. A., Almeida, A., et al. 2019, The Sixteenth Data Release of the Sloan Digital Sky Surveys: First Release from the APOGEE-2 Southern Survey and Full Release of eBOSS Spectra
- Arnett, W. D. & Meakin, C. 2011a, *ApJ*, 733, 78
- Arnett, W. D. & Meakin, C. 2011b, *ApJ*, 741, 33
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002
- Ben-Ami, S., Konidaris, N., Quimby, R., et al. 2012, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 8446, *Ground-based and Airborne Instrumentation for Astronomy IV*, 844686
- Bertin, E., Mellier, Y., Radovich, M., et al. 2002, in *Astronomical Society of the Pacific Conference Series*, Vol. 281, *Astronomical Data Analysis Software and Systems XI*, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 228
- Bilinski, C., Smith, N., Li, W., et al. 2015, *MNRAS*, 450, 246
- Blagorodnova, N., Neill, J. D., Walters, R., et al. 2018, *PASP*, 130, 035003
- Bruch, R. J., Gal-Yam, A., Schulze, S., et al. 2020, *arXiv e-prints*, arXiv:2008.09986
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Chambers, K. C., Boer, T. D., Bulger, J., et al. 2019, *Transient Name Server Discovery Report*, 2019-796, 1
- Chevalier, R. A. & Irwin, C. M. 2011, *ApJL*, 729, L6
- Cutri, R. M., Wright, E. L., Conrow, T., et al. 2013, *Explanatory Supplement to the AllWISE Data Release Products*, *Explanatory Supplement to the AllWISE Data Release Products*
- Danieli, B. & Soker, N. 2019, *MNRAS*, 482, 2277
- Dilday, B., Howell, D. A., Cenko, S. B., et al. 2012, *Science*, 337, 942

- Duev, D. A., Mahabal, A., Masci, F. J., et al. 2019, *MNRAS*, 489, 3582
- Elias-Rosa, N., Pastorello, A., Benetti, S., et al. 2016, *MNRAS*, 463, 3894
- Filippenko, A. V. 1997, *ARA&A*, 35, 309
- Finkbeiner, D. P., Padmanabhan, N., Schlegel, D. J., et al. 2004, *AJ*, 128, 2577
- Foley, R. J., Berger, E., Fox, O., et al. 2011, *ApJ*, 732, 32
- Foley, R. J., Smith, N., Ganeshalingam, M., et al. 2007, *ApJL*, 657, L105
- Fraser, M., Magee, M., Kotak, R., et al. 2013, *ApJL*, 779, L8
- Fremling, C., Dugas, A., & Sharma, Y. 2019, *Transient Name Server Classification Report*, 2019-188, 1
- Fremling, C., Miller, A. A., Sharma, Y., et al. 2020, *ApJ*, 895, 32
- Fuller, J. 2017, *MNRAS*, 470, 1642
- Fuller, J. & Ro, S. 2018, *MNRAS*, 476, 1853
- Gal-Yam, A. 2017, *Handbook of Supernovae*, 195–237
- Gal-Yam, A. 2019, in *American Astronomical Society Meeting Abstracts*, Vol. 233, American Astronomical Society Meeting Abstracts #233, 131.06
- Gal-Yam, A. 2019, *ARA&A*, 57, 305–333
- Gal-Yam, A., Arcavi, I., Ofek, E. O., et al. 2014, *Nature*, 509, 471
- Gal-Yam, A. & Leonard, D. C. 2009, *Nature*, 458, 865
- Gal-Yam, A., Leonard, D. C., Fox, D. B., et al. 2007, *ApJ*, 656, 372
- Gangopadhyay, A., Misra, K., Hiramatsu, D., et al. 2020, *ApJ*, 889, 170
- Ginzburg, S. & Balberg, S. 2012, *ApJ*, 757, 178
- Glas, R., Just, O., Janka, H. T., & Obergaulinger, M. 2019, *ApJ*, 873, 45
- Graham, M. J., Kulkarni, S. R., Bellm, E. C., et al. 2019, *PASP*, 131, 078001
- Hamuy, M., Phillips, M. M., Suntzeff, N. B., et al. 2003, *Nature*, 424, 651
- Ho, A. Y. Q., Goldstein, D. A., Schulze, S., et al. 2019, *ApJ*, 887, 169
- Hosseinzadeh, G., Arcavi, I., Valenti, S., et al. 2017, *ApJ*, 836, 158
- Hosseinzadeh, G., McCully, C., Zabludoff, A. I., et al. 2019, *ApJ*, 871, L9
- Humphreys, R. M. & Davidson, K. 1994, *PASP*, 106, 1025
- Janka, H.-T., Melson, T., & Summa, A. 2016, *Annual Review of Nuclear and Particle Science*, 66, 341
- Justham, S., Podsiadlowski, P., & Vink, J. S. 2014, *ApJ*, 796, 121
- Karamehmetoglu, E., Fransson, C., Sollerman, J., et al. 2019, *arXiv e-prints*, arXiv:1910.06016
- Kasliwal, M. M., Cannella, C., Bagdasaryan, A., et al. 2019, *PASP*, 131, 038003
- Katz, B., Sapir, N., & Waxman, E. 2011, *arXiv e-prints*, arXiv:1106.1898
- Khazov, D., Yaron, O., Gal-Yam, A., et al. 2016, *ApJ*, 818, 3
- Kiewe, M., Gal-Yam, A., Arcavi, I., et al. 2012, *ApJ*, 744, 10
- Kochanek, C. S. 2011, *ApJ*, 741, 37
- Kochanek, C. S. & Szczygiel, D. M. 2011, *ApJ*, 737, 76
- Kool, E. C., Karamehmetoglu, E., Sollerman, J., et al. 2020, *arXiv e-prints*, arXiv:2008.04056
- Law, N. M., Kulkarni, S. R., Dekany, R. G., et al. 2009, *PASP*, 121, 1395
- Liu, X., Shen, Y., Strauss, M. A., & Hao, L. 2011, *ApJ*, 737, 101
- Lupton, R. H., Gunn, J. E., & Szalay, A. S. 1999, *AJ*, 118, 1406
- Malmquist, K. G. 1922, *Meddelanden fran Lunds Astronomiska Observatorium Serie I*, 100, 1
- Margutti, R., Milisavljevic, D., Soderberg, A. M., et al. 2013, *ApJ*, 780, 21
- Masci, F. J., Laher, R. R., Rusholme, B., et al. 2019, *PASP*, 131, 018003
- Mauerhan, J. C., Smith, N., Filippenko, A. V., et al. 2013, *MNRAS*, 430, 1801
- Mcley, L. & Soker, N. 2014, *MNRAS*, 445, 2492
- Meakin, C. A. & Arnett, D. 2006, *ApJL*, 637, L53
- Moriya, T. J. 2014, *A&A*, 564, A83
- Moriya, T. J. 2015, *The Astrophysical Journal*, 803, L26
- Moriya, T. J. & Maeda, K. 2016, *ApJ*, 824, 100
- Moriya, T. J., Maeda, K., Taddia, F., et al. 2014, *MNRAS*, 439, 2917
- Müller, B. 2016, *PASA*, 33, e048
- Murase, K., Thompson, T. A., Lacki, B. C., & Beacom, J. F. 2011, *PhRvD*, 84, 043003
- Murase, K., Thompson, T. A., & Ofek, E. O. 2014, *MNRAS*, 440, 2528
- Nordin, J., Brinnel, V., van Santen, J., et al. 2019, *A&A*, 631, A147
- Nyholm, A., Sollerman, J., Taddia, F., et al. 2017, *Astronomy & Astrophysics*, 605, A6
- Nyholm, A., Sollerman, J., Tartaglia, L., et al. 2020, *Astronomy & Astrophysics*, 637, A73
- Ofek, E. O. 2019, *PASP*, 131, 054504
- Ofek, E. O., Cenko, S. B., Shaviv, N. J., et al. 2016, *ApJ*, 824, 6
- Ofek, E. O., Lin, L., Kouveliotou, C., et al. 2013a, *ApJ*, 768, 47
- Ofek, E. O., Rabinak, I., Neill, J. D., et al. 2010, *ApJ*, 724, 1396
- Ofek, E. O., Sullivan, M., Cenko, S. B., et al. 2013b, *Nature*, 494, 65
- Ofek, E. O., Sullivan, M., Shaviv, N. J., et al. 2014a, *ApJ*, 789, 104
- Ofek, E. O., Zoglauer, A., Boggs, S. E., et al. 2014b, *ApJ*, 781, 42
- Osborn, J., Föhrling, D., Dhillon, V. S., & Wilson, R. W. 2015, *MNRAS*, 452, 1707
- Owocki, S. P., Hirai, R., Podsiadlowski, P., & Schneider, F. R. N. 2019, *MNRAS*, 485, 988
- Owocki, S. P. & Shaviv, N. J. 2016, *MNRAS*, 462, 345
- Pastorello, A., Cappellaro, E., Inzerra, C., et al. 2013, *ApJ*, 767, 1
- Pastorello, A., Kochanek, C. S., Fraser, M., et al. 2018, *MNRAS*, 474, 197

- Pastorello, A., Smartt, S. J., Mattila, S., et al. 2007, *Nature*, 447, 829
- Pastorello, A., Wang, X. F., Ciabattari, F., et al. 2016, *MNRAS*, 456, 853
- Patat, F., Taubenberger, S., Benetti, S., Pastorello, A., & Harutyunyan, A. 2011, *A&A*, 527, L6
- Patterson, M. T., Bellm, E. C., Rusholme, B., et al. 2019, *PASP*, 131, 018001
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, 192, 3
- Perley, D. A., Fremling, C., Sollerman, J., et al. 2020, *ApJ*, 904, 35
- Piro, A. L. & Nakar, E. 2013, *ApJ*, 769, 67
- Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., et al. 2018, *AJ*, 156, 123
- Prieto, J. L., Brimacombe, J., Drake, A. J., & Howerton, S. 2013, *ApJ*, 763, L27
- Prieto, J. L., Kistler, M. D., Thompson, T. A., et al. 2008, *ApJL*, 681, L9
- Quataert, E., Fernández, R., Kasen, D., Klion, H., & Paxton, B. 2016, *MNRAS*, 458, 1214
- Quataert, E. & Shiode, J. 2012, *MNRAS*, 423, L92
- Rau, A., Kulkarni, S. R., Law, N. M., et al. 2009, *PASP*, 121, 1334
- Reguitti, A., Pastorello, A., Pignata, G., et al. 2019, *MNRAS*, 482, 2750
- Rest, A., Prieto, J. L., Walborn, N. R., et al. 2012, *Nature*, 482, 375
- Rigault, M. 2018, *ztfquery*, a python tool to access ZTF data
- Rigault, M., Neill, J. D., Blagorodnova, N., et al. 2019, *A&A*, 627, A115
- Robitaille, T. P., Tollerud, E. J., Greenfield, P., et al. 2013, *A&A*, 558, A33
- Sanders, N. E., Soderberg, A. M., Foley, R. J., et al. 2013, *ApJ*, 769, 39
- Sapir, N. & Waxman, E. 2017, *ApJ*, 838, 130
- Schlafly, E. F. & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Shaviv, N. J. 2000, *ApJL*, 532, L137
- Shaviv, N. J. 2001a, *ApJ*, 549, 1093
- Shaviv, N. J. 2001b, *MNRAS*, 326, 126
- Shiode, J. H. & Quataert, E. 2014, *ApJ*, 780, 96
- Silverman, J. M., Nugent, P. E., Gal-Yam, A., et al. 2013, *ApJS*, 207, 3
- Smith, N. 2017, *Handbook of Supernovae*, 403–429
- Smith, N. 2017, *Philosophical Transactions of the Royal Society of London Series A*, 375, 20160268
- Smith, N. & Arnett, W. D. 2014, *ApJ*, 785, 82
- Smith, N., E Andrews, J., Moe, M., et al. 2020, *MNRAS*, 492, 5897
- Smith, N., Mauerhan, J. C., Silverman, J. M., et al. 2012, *MNRAS*, 426, 1905
- Smith, N., Vink, J. S., & de Koter, A. 2004, *ApJ*, 615, 475
- Soumagnac, M. T. & Ofek, E. O. 2018, *PASP*, 130, 075002
- Soumagnac, M. T., Ofek, E. O., Gal-Yam, A., et al. 2019, *ApJ*, 872, 141
- Soumagnac, M. T., Ofek, E. O., Liang, J., et al. 2020, *ApJ*, 899, 51
- Stritzinger, M., Taddia, F., Fransson, C., et al. 2012, *ApJ*, 756, 173
- Strotjohann, N. L., Ofek, E. O., Gal-Yam, A., et al. 2015, *ApJ*, 811, 117
- Svirski, G., Nakar, E., & Sari, R. 2012, *ApJ*, 759, 108
- Szczygieł, D. M., Kochanek, C. S., & Dai, X. 2012, *ApJ*, 760, 20
- Tartaglia, L., Pastorello, A., Sullivan, M., et al. 2016, *MNRAS*, 459, 1039
- Thöne, C. C., de Ugarte Postigo, A., Leloudas, G., et al. 2017, *A&A*, 599, A129
- Wallis, S. 2013, *Journal of Quantitative Linguistics*, 20, 178
- Wilson, E. B. 1927, *Journal of the American Statistical Association*, 22, 209
- Woosley, S. E. 2017, *ApJ*, 836, 244
- Woosley, S. E., Heger, A., & Weaver, T. A. 2002, *Reviews of Modern Physics*, 74, 1015
- Wu, S. & Fuller, J. 2020, *arXiv e-prints*, arXiv:2011.05453
- Yao, Y., Miller, A. A., Kulkarni, S. R., et al. 2019, *ApJ*, 886, 152
- Yaron, O., Perley, D. A., Gal-Yam, A., et al. 2017, *Nature Physics*, 13, 510–517
- Zackay, B., Ofek, E. O., & Gal-Yam, A. 2016, *ApJ*, 830, 27

APPENDIX

A. PRECURSOR IMAGES

To test whether the precursor candidates are real, we inspect the difference images. We select either the g or r band and coadd the difference images that yield the most significant fluxes in our search (see Figs. 3 and 4). Before the coaddition we subtract the median pixel flux from the image and divide all fluxes by the robust standard deviation (half of the difference between the 15.9% and 84.1% percentile). The images are then coadded using the *SWarp* software (Bertin et al. 2002). The resulting skymaps are shown in Fig. 15. Clear point sources are visible for most SNe.

To quantify the significance of the detections in Fig. 15, we calculate the S-image as described by Zackay et al. (2016). For this purpose, we cross correlate the difference image with the PSF of the difference image and coadd the resulting significance maps again using *SWarp*. To normalize the significances we divide each image by the robust standard deviation and show the significance maps in Fig. 16.

Most precursors are visible as point sources, with exception of SN 2019sae. This precursor candidate is considered unconfirmed as only a single bin drives the significance (compare Sec. 3.1) and we conclude that it is likely spurious. The other three unconfirmed precursors (marked by red titles in Figs. 15 and 16) appear as point sources, so we cannot rule out that they are astrophysical. As described in Sec. 3.1 we neglect these marginal detections here and only focus on the 19 confirmed precursors. In some cases astrometric residuals are visible in the cutouts close to the precursors. Especially affected are SN 2018gho and SN 2020dfh. We nevertheless consider them true detections as they also appear to be present in single images without astrometric residuals. We caution however that their light curves could be affected by the missubtractions. They are therefore less reliable than the light curves of other precursors.

B. BASELINE CORRECTION OF SN 2019CMY

When using late-time observations (> 380 days after the explosion) for the baseline correction of SN 2019cmy, we find that the complete pre-explosion light curve becomes positive as shown in Fig. 17. The late-time observations include > 60 observations per filter collected in > 30 different nights over a period of 50 days. They are consistent with a constant flux, which is however $\sim 10^{-9}$ lower than the median flux before the explosion. To probe whether a systematic error could explain this deviation we perform a series of tests: The hour angle and air mass are on average similar before and after the explosion. The seeing is better during the last block of observations collected since March 2020, potentially because of reduced air pollution as a consequence of the lockdown measures due to the Corona pandemic. However, other SNe in our sample do not show a similarly strong flux change, implying that we do not see a global effect caused by changes in the atmosphere.

The pixel coordinates of the SN position drift slowly by in total 100 pixels over three years, but there is no abrupt change consistent with the two different flux levels. We also inspect the mirrored SN position, located at a distance of 11.2 arcsec on the other side of the host galaxy's center. We observe a constant flux level for the complete light curve, i.e. a similar flux reduction is not observed at late times. Finally, we use the ZUDS pipeline to construct a different reference image, redo the image subtraction and produce a forced photometry light curve. The flux difference between the observations before and after the explosion is similarly large. It is therefore likely not induced by the image subtraction or forced photometry pipeline.

After we could not identify any systematic effect responsible for the lower fluxes after the SN has faded, we here discuss an astrophysical interpretation. If real, the flux drop would imply that the progenitor star is detected for 400 days before the SN explosion. A flux of 10^{-9} corresponds to a magnitude of 22.5, i.e. an absolute magnitude of -14.1 for the SN redshift of 0.0314. The detected progenitor star prior to SN 2005gl had a V band magnitude of only -10.4 (Gal-Yam & Leonard 2009), but we cannot rule out that the progenitor underwent a more than year-long outburst with an approximately constant luminosity. Similar outbursts were observed several years prior to the explosion of SN 2015bh and reached typical r -band magnitudes between -9 and -13 (Elias-Rosa et al. 2016; Thöne et al. 2017). It is thus not excluded that we indeed detect the progenitor star.

Since we are not sure whether or not the flux reduction after the SN explosion is astrophysical, we here exclude the late-time observations and do the baseline correction using pre-explosion data. r -band observations in two nights, 7 and 6 days before the estimated explosion date surpass the 5σ detection as shown in Fig. 3. They could either be a precursor eruption or a wind-shock breakout peak (Ofek et al. 2010; Chevalier & Irwin 2011; Piro & Nakar 2013).




Figure 15. Coadded ZTF difference images showing the precursors. Red captions indicate that the corresponding precursor is only detected in a single bin and is therefore considered unconfirmed (see Sec. 3.1). No point source is visible for SN 2019sae and we conclude that this precursor candidate is not real. A negative source is detected at the position of SN 2019meh due to the AGN variability in the host galaxy. SN 2019zrk and SN 2020dcs are located close to the edge of the CCD.




Figure 16. Significance maps of the precursor candidates.




Figure 17. Complete light curve for the position of SN2019cmy when using the last observations (yellow region) for the baseline correction. The peak of the SN is cut for better visibility. The complete pre-explosion light curve (gray region) is positive, i.e. the fluxes before and after the SN cannot be zero at the same time. We could not identify a systematic effect that might explain this discrepancy. If the effect is astrophysical, it would imply that the progenitor star is detected with an absolute magnitude of -14.1 . In this paper, we use pre-explosion observations for the baseline correction and only discuss the precursor detected one week before the SN (red region).