



## Broad host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial community

Klümper, Uli; Riber, Leise; Dechesne, Arnaud; Sannazzaro, Analia; Hansen, Lars H.; Sørensen, Søren J.; Smets, Barth F.

*Published in:*  
I S M E Journal

*Link to article, DOI:*  
[10.1038/ismej.2014.191](https://doi.org/10.1038/ismej.2014.191)

*Publication date:*  
2014

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Klümper, U., Riber, L., Dechesne, A., Sannazzaro, A., Hansen, L. H., Sørensen, S. J., & Smets, B. F. (2014). Broad host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial community. *I S M E Journal*, 1-12. <https://doi.org/10.1038/ismej.2014.191>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Title:**

Broad host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial community

**Short title:**

Plasmids invade surprisingly diverse bacteria

5

**Author affiliations:**

Uli Klümper<sup>a</sup>, Leise Riber<sup>b</sup>, Arnaud Dechesne<sup>a</sup>, Analia Sannazzarro<sup>b</sup>, Lars H. Hansen<sup>b</sup>, Søren J. Sørensen<sup>b</sup>,  
Barth F. Smets<sup>a\*</sup>

<sup>a</sup> Technical University of Denmark, Department of Environmental Engineering, Miljøvej 113, 2840 Kgs.

10 Lyngby, Denmark

<sup>b</sup> University of Copenhagen, Department of Biology, Microbiology, Universitetsparken 15, 2100

Copenhagen Ø, Denmark

\*corresponding author

15 **Subject category:**

Evolutionary genetics

**Corresponding author:**

Barth F. Smets

Technical University of Denmark

20 DTU Miljø

Miljøvej 113

2840 Kgs. Lyngby

Denmark

Ph. +45 45 25 22 30

25 Fax +45 45 93 28 50

E-mail bfm@env.dtu.dk

**Abstract:**

Conjugal plasmids can provide microbes with full complements of new genes and constitute potent vehicles for horizontal gene transfer. Conjugal plasmid transfer is deemed responsible for the rapid spread of antibiotic resistance among microbes. While broad host range plasmids are known to transfer to diverse hosts in pure culture, the extent of their ability to transfer in the complex bacterial communities present in most habitats has not been comprehensively studied. Here, we isolated and characterized transconjugants with a degree of sensitivity not previously realized in order to investigate the transfer range of IncP-type broad host range plasmids from three proteobacterial donors to a soil bacterial community. We identified transfer to many different recipients belonging to 11 different bacterial phyla. The prevalence of transconjugants belonging to diverse Gram-positive Firmicutes and Actinobacteria suggests that inter-Gram plasmid transfer of IncP-1 and IncPromA-type plasmids is a frequent phenomenon. While the plasmid receiving fractions of the community were both plasmid- and donor- dependent, we identified a core super-permissive fraction that could take up different plasmids from diverse donor strains. This fraction, comprising 80% of the identified transconjugants, thus has the potential to dominate IncP- and IncPromA - type plasmid transfer in soil. Our results demonstrate that these broad host range plasmids have a hitherto unrecognized potential to readily transfer to very diverse bacteria and can, therefore, directly connect large proportions of the soil bacterial gene pool. This finding reinforces the evolutionary and medical significances of these plasmids.

**Keywords:** Broad host range/Conjugation/Gene transfer/Plasmid/Transfer range

## Introduction

50 Conjugal plasmid transfer is a process by which bacteria horizontally transfer complete sets of genes to other, potentially distantly related, organisms. Conjugal plasmids frequently carry accessory genes, often encoding antibiotic or metal resistances, catabolic pathways, or virulence factors. They are often implicated in the evolution of pathogenic bacteria and the rapid spread of antibiotic resistance, likely fostering the rise of multiple-resistant microbes in hospitals (Levy & Marshall, 2004) and animal husbandries (Zhu *et al.*,  
55 2013). While the relevance of plasmid transfer has become very acute in this age of massive antibiotic usage, plasmids have been exchanged for much longer and many prokaryotic genomes present signs of intense past horizontal gene transfer (Ochman *et al.*, 2000).

Plasmids present different abilities to transfer into, and be maintained in, distantly related bacterial hosts and are loosely categorized as having a narrow or broad host range. The transfer of narrow host range  
60 plasmids is limited at one of the steps required for successful transfer, such as the formation of mating pairs, the avoidance of the recipient's restriction system, or the correct expression of its replication and maintenance systems in the recipient (Thomas & Nielsen, 2005). Some broad host range plasmids can transfer across bacterial phyla and even across domains of life (Waters, 2001; Heinemann & Sprague, 1989), and several genetic determinants conferring broad host transfer capability have been identified (Jain  
65 & Srivastava, 2013).

The host range is thus a key parameter that controls the ecology and fate of plasmids. The evaluation of host range has traditionally been conducted using few individual pure strains as recipients, a situation that contrasts with the fact that most bacteria - and thus most plasmids - exist within complex communities of hundreds to thousands of species (Hong *et al.*, 2006; Brown Kav *et al.*, 2012). Among these diverse  
70 communities, all strains are obviously not equally permissive towards plasmid receipt, even for broad host range plasmids. This notion was supported when studying plasmid transfer to a range of strains isolated

from marine water or wastewater treatment bioreactors (Sørensen, 1993; Inoue *et al.*, 2005). With the use of fluorescent reporter genes to track plasmids, which reduces the need for selection and cultivation steps to identify transconjugants, it has become apparent that, in complex communities, broad host range plasmids can indeed be received by bacteria distantly related to the donor, even in the absence of selective pressure for plasmid carriage (Musovic *et al.*, 2014, 2006; Shintani *et al.*, 2014; De Gelder *et al.*, 2005). However these efforts, limited to inspection of a few hundred transconjugants at best, most likely underestimate the true diversity of transconjugal pools and do not accurately describe how plasmid permissiveness may vary across taxa in complex microbial communities.

Horizontal gene transfer between different species has been recognized as a common and major evolutionary process (Zhaxybayeva & Doolittle, 2011), most acutely demonstrated in the heavy interconnection between the resistome of soil dwelling bacteria and human pathogens (Forsberg *et al.*, 2012). The behavior of this environmental resistome may, thus, govern the spread of antibiotic resistance genes to pathogens (Finley *et al.*, 2013). Plasmids serve as main vessels of gene flow in microbial communities, linking distinct genetic pools (Halary *et al.*, 2010; Norman *et al.*, 2009). The *in situ* host range of plasmids may, then, well govern the taxonomic breadth across which gene flow occurs.

Here, taking advantage of high throughput cell sorting and next-generation sequencing technologies, we map for the first time the intrinsic diversity of the bacterial recipients of broad host range plasmids in a microbial community extracted from soil, under conditions where cell-to-cell contacts are maximized. We analyzed matings initiated with combinations of three plasmid donors and three plasmids to identify, how permissiveness towards broad host range plasmids is distributed across taxa among the recipient community.

## 95 **Material & Methods**

### **Donor strain construction**

Soil bacterial communities were challenged with various plasmid/donor combinations through solid surface filter matings. The plasmids were marked with a genetic tag encoding conditionally expressible green fluorescent proteins. The used entranceposon (Bahl *et al.*, 2009) carries a *lacI<sup>q</sup>* repressible promoter upstream the *gfpmut3* gene, encoding for the green fluorescent protein (GFP). Plasmid donor strains were all chromosomally tagged with a gene cassette encoding constitutive red fluorescence and constitutive *lacI<sup>q</sup>* production. As a result, there is no *gfp* expression in the donor strains, but upon plasmid transfer to a soil bacterium, *gfp* expression is possible, resulting in green fluorescent cells or microcolonies, which can be detected and sorted by fluorescence microscopy or FACS, respectively (Figure 1) (Sørensen *et al.*, 2005).

105 *Pseudomonas putida* KT2440, *Escherichia coli* and *Kluyvera* sp. served as donor strains, and were each electroporated with the plasmid pGRG36-*lacI<sup>q</sup>*-Km-Lpp-mCherry carrying both the transposase genes and the Tn7 *lacI<sup>q</sup>*-Lpp-mCherry-Km<sup>R</sup> region for specific integration of the *lacI<sup>q</sup>*-Lpp-mCherry-Km<sup>R</sup> gene cassette into the chromosomal *attTn7* site. Colonies were selected for Km<sup>R</sup> on LB agar plates at 30 degrees. Colonies were restreaked on selective LB agar plates at 30 °C, incubated in liquid LB overnight culture without antibiotics at 30 °C and finally streaked on LB agar plates without selection at 37 °C for integration of the gene cassette and subsequent loss of the Tn7 helper plasmid. Colonies were tested for successful loss of helper plasmid and chromosomal integration of gene cassette by PCR (McKenzie & Craig, 2006). The same colonies were also phenotypically verified to be bright red fluorescent using stereo microscopy.

### **Construction of *gfpmut3*-tagged plasmid pKJK5**

115 Plasmids RP4 and pIPO2tet have been constructed earlier (Musovic *et al.*, 2010, 2014). The 54 kbp IncP-1 plasmid, pKJK5, originally isolated from a soil/manure environment, harbors a tetracycline and a trimethoprim resistance determinant as well as a class 1 integron (Sengeløv *et al.*, 2001). The

Entranceposon [ $Km^R$ , PA10403-*gfpmut3*], carrying a kanamycin resistance determinant and a  $\text{LacI}^q$  repressible promoter upstream the *gfpmut3* gene, encoding the Green Fluorescent Protein (*gfp*), was derived from pEntranceposon [ $Km^R$ ] (Finnzymes, F-766) and randomly inserted into the plasmid pJKK5 using the artificial Mu transposon in vitro delivery system as described previously (Bahl *et al.*, 2009). Transformed *E. coli* GeneHogs single colonies were selected for resistance towards trimethoprim and kanamycin and screened for sensitivity towards tetracycline in order to select for plasmid derivatives with an Entranceposon insert location directed to an accessory element (the tetracycline resistance determinant), thereby excluding any potential impacts on conjugation transfer ability. The exact insert location of [ $Km^R$ , PA10403-*gfpmut3*] in the selected pJKK5 derivative of this study was determined by sequencing out from the inserted fragment in one direction using primer Seq\_Bw\_Ent\_gfp: 5'-GCCAGAACCGTTATGATGTCGG-3'. The insertion mapped to position 30.614 bp in the *tetA* gene (30.435-31.634 bp) of plasmid pJKK5 (accession no. AM261282). The selected *gfpmut3*-tagged pJKK5 plasmid was finally introduced into *E. coli* MG1655:: $Km^R$ -Lpp-mCherry, *P. putida* KT2440:: $Km^R$ -Lpp-mCherry and *Kluyvera* spp.:: $Km^R$ -Lpp-mCherry cells by transformation.

### Soil sampling & community extraction

Soil samples were taken at the annually tilled CRUCIAL (Closing the Rural Urban Nutrient Cycle) agricultural field site (Taastrup, Denmark) from a plot subjected to no further agricultural treatment (Magid *et al.*, 2006). Soil samples were collected in late fall 2012. Samples were taken from three different plots of this treatment. Each plot was sampled for 1 kg of soil at 5 locations. The resulting soil volume was sieved and homogenized to obtain a representative sample. From a total of 30 g of the homogenized chosen soils, indigenous bacterial communities were isolated by Nycodenz<sup>®</sup>-extraction (Musovic *et al.*, 2010) and used as recipients in the mating assay. Donor strains were grown overnight in LB-medium supplement to the plasmid specific antibiotics (Table 1), harvested by centrifugation.

### Solid surface filter mating assay

The extracted recipient community was challenged with exogenous plasmids via solid-surface filter matings (Musovic *et al.*, 2010) modified to an initial ratio of donor to recipient bacteria of 1:1 at a density of approximately 30,000 bacteria/mm<sup>2</sup> on the filter. As a growth medium we used a 10% soil extract medium as described by Musovic *et al.* (2010) buffered at pH 7.2 with 5mM MOPS and supplemented with 20 µg/mL Nystatin to avoid fungal growth. Unlike in Musovic *et al.* (2010), we did not use additional nutrient additions, but only relied on soil extracted nutrients to support activity during the mating incubations. Successful conjugation was checked after 48 hours by epifluorescence stereo microscopy and confocal laser scanning microscopy (CLSM) (Figure 1) (Musovic *et al.*, 2010).

#### 150 **Cell collection and triple gated fluorescence activated cell sorting of transconjugants**

Cells from 5 filters per mating combination replicate were harvested in 2mL 0.9% NaCl-solution by vortexing for 3 minutes. Flow cytometric detection of cells was carried out using a FACSAria IIIu (Becton Dickinson Biosciences, San Jose, CA). The following settings and voltages were used during analysis: forward scatter (FSC) = 505 V, side scatter (SSC) = 308 V, and detectors for green (BP filter 530/30 nm) and red fluorescence (BP filter 610/20 nm) were set at 508 V and 500 V, respectively. A 70 µm nozzle was used at a sheath fluid pressure of 70 psi. The BD FACSDiva™ software v6.1.3 was used for both operating and analyzing results. Sorting was performed using a 488 nm (20 mW) laser connected to the green fluorescence detector at 515-545 nm and a 561 nm (50 mW) laser connected to the red fluorescence detector at 600-620 nm. Three gates were defined in bivariate plots to sort for transconjugants. On the SSC-A vs FSC-A plot a gate for only particles of bacterial size was used. On the FITC-A vs SSC-A plot a gate was set that covered all green fluorescent particles, while using an additional non-red gate on the PE-Texas Red-A vs SSC-A plot excluded all small autofluorescent particles from soil or leaking donors (Figure 2) to sort out only transconjugants. All samples were diluted in 0.9% NaCl to approximately 2000 counting events s<sup>-1</sup> before fluorescent activated cell sorting to assure for optimal sorting. Transconjugants that originally made up for less than 0.1% of the total community in the filter matings and were enriched to up



to 82% in a first fast sorting step, before isolating over 10,000 transconjugants per sample in a second purification step, leading to 100% purity of green cells as observed by fluorescent counting in the flow cytometer. Plating of more than 200 isolated transconjugants on 10% soil extract medium (Musovic *et al.*, 2010) resulted in detection of green fluorescence in all colonies, additionally verifying purification of gfp-  
170 expressing transconjugants. Of the isolated transconjugants, twenty were subject to 16S rRNA gene sequences; the recovery of proteobacterial, sphingobacterial and actinobacterial phylotypes indicated diversity among transconjugants.

### **Bacterial cell lysis, amplification and sequencing**

Bacterial transconjugal cells from the second sort, initially collected in 5 mL sterile polystyrene round-  
175 bottom Falcon™ tubes (BD Biosciences, San Jose, CA) with 0.5 mL 0.9% NaCl solution, were transferred to 1.5 mL Eppendorf tubes and centrifuged at 10,000 x g for 30 min to collect the cell pellets. The supernatant was carefully removed, the cell pellet suspended in 20 µL of Lyse and Go PCR Reagent (Thermo Scientific, Waltham, MA, USA) and the lysis mixtures transferred to 0.2 mL amplification tubes. Cell lysis was subsequently performed in an Arktik™ Thermal Cycler (Thermo Scientific, Waltham, MA, USA) using the  
180 program: one initial cycle at 57 °C for 30 s, a second cycle at 8 °C for 30 s, a third cycle at 65 °C for 90 s, a fourth cycle with heating to 97 °C for 3 min, a fifth cycle with cooling to 8 °C for 60 sec, a sixth cycle with heating to 65 °C for 3min followed by additional heating to 97 °C for 60 s and cooling to 65 °C for 60 s with a final end-step at 80 °C. DNA-containing cell lysis products were immediately put on ice and used directly for subsequent PCR. Then, 5 µL of the cell lysis product from the previous step was used directly for  
185 sequencing library preparation. Tag-encoded 16S rRNA gene pyrosequencing was carried out after amplification of the V3 and V4 region (Primers: 341F: 5'-CCTAYGGGRBGCASCAG-3 and 806R 5'-GGACTACNNGGTATCTAAT-3) using the PCR procedures and GS FLX Titanium chemistry as described previously (Hansen *et al.*, 2012).

### **Sequence analysis and tree construction**

190 Sequence analysis was carried out using mothur v.1.32.1 (Schloss *et al.*, 2009) and the 454 SOP (Schloss *et al.*, 2011) as accessed on 01.11.2013 on [http://www.mothur.org/wiki/454\\_SOP](http://www.mothur.org/wiki/454_SOP). Sequences were classified based on the RDP classifier (Wang *et al.*, 2007). Phylogenetic trees were constructed using iTOL (<http://itol.embl.de/>) (Letunic & Bork, 2007). All sequences have been submitted to NCBI and can be accessed under number XXX.

## 195 **Results and Discussion**

### **High throughput isolation and sequencing of transconjugants**

We explored the ability of a bacterial community extracted from soil to engage in horizontal gene transfer and receive one of three *gfp*-tagged broad host-range plasmids from three different red fluorescent-tagged donor strains in which plasmid-mediated *gfp* expression is repressed (Table 1). In soil, physical barriers limit contact between freshly introduced plasmid donors and potential recipients (Dechesne *et al.*, 2005); here we maximized cell-to-cell contact in a gene transfer assay (Musovic *et al.*, 2010) to study the intrinsic permissiveness of the recipient community. All three plasmids (RP4, pIPO2tet, and pKJK5) were introduced to the soil community in matings with a *Pseudomonas putida* donor strain, while plasmid pKJK5 was also introduced via *Escherichia coli* and *Kluyvera* spp. donors (Supplementary Table 1). After mating, the *gfp*-expressing transconjugant cells (Figure 1) were isolated from the mixed community by fluorescent activated cell sorting (FACS). A novel triple gated FACS approach based on size, green fluorescence, and lack of red fluorescence, allowed specific isolation of large numbers of transconjugant cells, in spite of their low relative abundance (less than 0.1%) in the mating mixture (Figure 2). At least 14,000 transconjugant cells were obtained for each mating, corresponding to 28,000 – 116,500 transconjugants per donor/plasmid combination, depending on the number of replicate matings. The eleven pools of sorted transconjugants as well as the total soil recipient community were then subjected to deep amplicon sequencing of 16S rRNA genes, resulting in 29,894 to 50,398 sequences per sample after processing with the mothur pipeline

(Schloss *et al.*, 2009). This corresponds to more sequences than sorted transconjugants for most samples (Supplementary Table 1), providing an adequate picture of the observed plasmid transfer range.

#### 215 **Transconjugal pools are plasmid- and donor- specific**

The phylogenetic structure of the transconjugal pools was compared after clustering the partial 16S rRNA gene sequences in OTUs at 97% similarity. The eleven transconjugal pools clustered clearly and significantly apart from the recipient community, as shown by PCoA (Figure 3) and AMOVA (Excoffier *et al.*, 1992) ( $p=0.028$ ). Mating plates contained soil extracts as nutrient sources and growth on filter did not significantly modify the soil community structure ( $p=0.797$ ) based on UNIFRAC comparisons (Lozupone *et al.*, 2011), in spite of a diversity reduction by 72%. The transconjugal pools were clearly distinct from the recipient community, but also differed from each other based on plasmid or donor. Considering different plasmids in an identical donor strain (*P. putida*) and providing the same plasmid (pKJK5) in different donor strains revealed phylogenetically distinct transconjugal pools (AMOVA,  $p<0.001$ ). Hence, plasmid acquisition is not a stochastic process, even for broad host range plasmids. While replicates of the same donor/ plasmid combinations differed based on weighted UNIFRAC comparisons ( $p<0.05$ ), the average interreplicate dissimilarity ( $W=0.36$ ) was clearly less than dissimilarity between different plasmid/donor combinations ( $W=0.49$ ) or between transconjugal pools and the soil community ( $W=0.60$ ). Slight differences between the replicates can also be seen in their phylum level distribution of transconjugants (Supplementary Figure 1). This dissimilarity between replicates can most likely be decreased through sorting of higher numbers of transconjugants per replicate, since replicates from the same donor-plasmid combinations grouped significantly together in PCoA ( $p<0.01$ ) (Figure 3). Based on this PCoA grouping and because the numbers of replicates per combination differed (Supplementary Table 1), replicates were pooled for subsequent phylogenetic analysis.

#### 235 **Transconjugal pools span most of the major bacterial phyla**

More than 300 transconjugant OTUs were detected across all plasmid/donor combinations (Figure 4, Figure 5), a large expansion over the low number of distinct bacterial isolates identified previously from matings in complex environmental communities (De Gelder *et al.*, 2005; Musovic *et al.*, 2014, 2010; Shintani *et al.*, 2014). As expected, Proteobacteria, known to be the main hosts for the studied broad-host-range plasmids (Suzuki *et al.*, 2010), were represented. Unlike in previous studies (Musovic *et al.*, 2010; Shintani *et al.*, 2014), all five classes ( $\alpha$ - $\epsilon$ ) of Proteobacteria were identified among the transconjugants. More strikingly, the diversity of transconjugants extended much beyond the proteobacterial phylum, and included diverse members of ten additional phyla including Verrucomicrobia, Bacteroidetes and Actinobacteria, some of which are known as poorly cultivable (Joseph *et al.*, 2003). The IncP transfer apparatus is known to build conjugative bridges between a huge variety of organisms (Grahn *et al.*, 2000; Thomas & Nielsen, 2005). Shuttle vectors for gene transfer from Proteobacteria to distantly related recipients such as Cyanobacteria (Wolk *et al.*, 1984) or gram-positive bacteria and yeast (Heinemann & Sprague, 1989; Samuels *et al.*, 2000) have, indeed, been built using the RP4 transfer system, a IncP-1 $\alpha$  subgroup plasmid. While the wide transfer potential of the RP4 conjugation system has therefore been known in artificial constructs under laboratory conditions, we are the first to prove that a large proportion of this transfer potential can be realized in nature. A similarly huge transfer potential is demonstrated for pKJK5, a plasmid closely related to RP4 belonging to the IncP-1 $\epsilon$  subgroup and for pIPO2tet, a currently unclassified, phylogenetically more distant, cryptic plasmid that merely seems to provide plasmid mobilization capability to its host (Figure 4). We thus show that a variety of broad host range plasmids can effectively be transferred to, and encoded genes can be expressed in large proportions of the bacterial tree of life, much beyond the limited transfer range identified so far (Shintani *et al.*, 2014). The realized transfer range in the soil community under natural conditions might be even higher taking into account that Nycodenz extraction might not be able to recover all bacterial phyla from the soil sample (Holmsgaard *et al.*, 2011). Of the total extractable soil microbial community only the phyla Chloroflexi, Deinococcus-Thermus, Nitrospira and SR1, were not represented in the transconjugal pools in our experiments.

In particular, we identified transfer from the used Gram-negative donor strains to a wide variety of gram positive bacteria (Figure 4, Figure 5). Over 15 OTUs within the Actinobacteria phylum and more than 10 OTUs belonging to 6 different orders of Bacilli and Clostridia within the Firmicutes phylum were identified as transconjugants. Inter Gram conjugal gene transfer has been shown with vectors consisting partly of the broad host range transfer machinery of RP4 recombined with the *sacB* gene from Gram-positive *Bacillus subtilis* (Schäfer *et al.*, 1994), but has only exceptionally been identified in natural habitats (Musovic *et al.*, 2006). Our observations suggest that it may be a more common process than previously considered.

### **Abundance in recipient community and phylogenetic distance to the donor do not explain the composition of transconjugal pools**

In spite of the large diversity within the transconjugal pools, not all OTUs of the recipient community were represented in each pool and the relative abundance of OTUs in transconjugal pools was very heterogeneous. Our method cannot distinguish between original horizontal plasmid transfer events from subsequent vertical plasmid transfer through growth of transconjugants on the mating filter. Therefore, relative abundance in the transconjugal pools can be influenced by the relative growth rate of recipients. However, the fact that OTU abundance in the transconjugal pools is not explained by their abundance in the reference soil recipient community (Figure 4, Figure 5, and Supplementary Table 2) indicates that plasmid transfer occurs preferentially to some recipients and that transconjugal pools are not simply determined by the recipient's growth ability.

Next, we tested whether phylogenetic distance between donor and recipient, calculated based on the Sogin distance algorithm (Sogin *et al.*, 2006), influenced the abundance of individual OTUs among the transconjugal pools. We found no significant correlation between phylogenetic distance to the donor and recipient frequency in the transconjugal pools ( $p=0.09-0.94$ ) for any of the donor plasmid combinations (Supplementary Figure 2). For example, the most abundant OTUs in soil that do not appear in the transconjugal pools (Supplementary Table 2) are Gammaproteobacteria; they display more than 90% 16S

285 rRNA gene sequence similarity to the donor strains, while other OTUs with less than 70% sequence  
similarity to donor cells, such as several members of the Flavobacterium phylum, did receive at least one of  
the plasmids. Transfer of an IncP-1 plasmid from *E.coli* to phylogenetically distant Flavobacteria was  
detected in soil microcosms (Pukall *et al.*, 1996), indicating that transfer to distant nodes of the  
phylogenetic tree is not only possible, but also realized in undisturbed soil environments. In pure culture,  
290 permissiveness towards broad host range plasmids of isolates that are indistinguishable by 16S rRNA gene  
analysis can differ by more than 100-fold (Heuer *et al.*, 2010). Here we confirm that inferring plasmid  
uptake and transfer frequency cannot be predicted based on the phylogenetic identity of an OTU.

However, we confirm the role of donors in defining the plasmid transfer host range (De Gelder *et al.*, 2005),  
and show that this effect is significant even for two donors belonging to the same family of  
295 Enterobacteriales (*E. coli* & *Kluyvera* sp.) and thus sharing a high genomic similarity. The reasons behind this  
are uncertain, but certain strains might have distinct abilities to achieve efficient cell-to-cell contact with a  
specific recipient, for example through specific mating mediating pheromones (Hirt, 2002). Earlier studies  
have shown that plasmid exchange between two taxonomically different species can exceed intraspecies  
transfer frequencies (Bingle *et al.*, 2003), proving that the regulatory interactions of donor, recipient, and  
300 plasmid can influence transfer efficiency.

Similarly, three broad host range plasmids, all carried by the same *P. putida* strain, were transferred to  
distinct pools of recipients. Yano *et al.* (2013) hypothesized that, genetic differences appearing among  
closely related IncP-1 plasmids through plasmid backbone evolution can result in significant diversities in  
host range efficiency without affecting their broad host range nature. Such backbone alterations exist  
305 between the IncP-1 $\alpha$  (RP4) and IncP-1 $\epsilon$  (pKJK5) core regulatory proteins such as *KorB*, *TrfA*, *TrbA* and *Ssb*  
(Bahl *et al.*, 2007). Although these two plasmids are incompatible (both IncP-1), differences in gene  
silencing and expression of the different core proteins could explain the different transconjugal patterns.  
Since already minor differences in regulation between two IncP-1 plasmids lead to distinct transconjugal

pools, it is coherent that the unrelated transfer machinery of plasmid pIPO2tet caused significantly ( $p < 0.05$ )  
310 dissimilar transconjugal pools when compared to the IncP ones.

### **A core super-permissive community fraction dominates gene transfer**

Out of 281 OTUs identified in the transconjugal pools with the three different broad host range plasmids  
and *P. putida* as donor, 74 OTUs were common to all three pools (Figure 6A). A similar observation (46 out  
of 279 OTUs shared) held when comparing the transconjugal pools for plasmid pKJK5 introduced via three  
315 different donors (Figure 6B). Therefore, the majority of transconjugant OTUs were only identified in single  
donor/plasmid combinations. This might result from mating pair combinations that each favor or reduce  
gene transfer abilities (Bingle *et al.*, 2003; Thomas & Nielsen, 2005; Yano *et al.*, 2013).

While only 74 and 46 OTUs are shared among the compared transconjugal pools, these OTUs represent  
over 80% of the transconjugal sequences (Figure 6C&D). This core super-permissive community fraction  
320 shared by all five transconjugal pools is able to take up diverse broad host range plasmids from diverse  
donor strains at high frequencies. The presence of this shared core in each analyzed transconjugal pool is  
the crucial discriminant that groups transconjugal pools apart from the original soil community (Figure 3).  
The core super-permissive community consists mainly of diverse Proteobacteria like Enterobacteriales ( $\gamma$ ),  
Burkholderiales ( $\beta$ ), Pseudomonadales ( $\gamma$ ) and Rhizobiales ( $\alpha$ ) (Figure 4). In addition, within this core super-  
325 permissive fraction, several OTUs that are rare in the recipient community ( $< 0.001\%$ ) are more than 20-fold  
overrepresented in transconjugal pools (Figure 4). The participation of these rare community members in  
gene transfer might play a crucial role in increasing the communal gene pool through rapid recombination  
with plasmids, since the rare biosphere can harbor a great reservoir of genes (Sogin *et al.*, 2006).

### **Medical relevance**

330 The large realized transfer potential of newly introduced plasmids in soil may be of medical importance. In  
recent EAHEC outbreaks in Germany, recombination of a pathogenic with the plasmid of a non-pathogenic  
*E. coli* strain increased the pathogenic potential to cause a deadly combination (Brzuszkiewicz *et al.*, 2011).

Soil borne antibiotic resistance has been found to be shared with human pathogens (Benveniste & Davies, 1973; Forsberg *et al.*, 2012). Several organisms among the identified transconjugants belong to groups  
335 known to contain opportunistic human pathogens, providing a direct link between plasmid encoded soil  
resistome and opportunistic pathogens. These groups include the proteobacterial *Enterobacteria*,  
*Pseudomonas* or *Campylobacter* but also groups from other phyla such as *Fusobacterium*, *Streptococcus*  
and *Staphylococcus*, most of which are treated with antibiotic therapy. Especially the acquisition of new  
antibiotic resistance genes through plasmid mediated gene transfer may push the pathogenic potential of  
340 *Staphylococcus*, originating from rapid evolution of virulence and drug resistance (Holden *et al.*, 2004),  
even further.

The observed transfer of broad-host-range IncP-1 type plasmids between Gram-negative and Gram-positive  
bacteria might lead to a reassessment of the potential of soil bacterial communities to spread antibiotic  
resistance genes. Indeed, Actinobacteria, the origin of many soil-borne resistance genes (D'Costa *et al.*,  
345 2006) which are sometimes identified in clinical isolates of Gram negative antibiotic-resistant bacteria  
(Benveniste & Davies, 1973), are frequent among the transconjugants we identified. Broad host range  
plasmids of the IncP-1 and IncPromA group can thus provide a direct link between diverse bacterial groups.  
Especially IncP-1 $\epsilon$  plasmids such as pJK5 have been identified as vectors of antibiotic resistance genes  
transfer among Proteobacteria by additionally hosting Class 1 integron gene cassettes (Heuer *et al.*, 2012).  
350 These Class 1 integrons may not only spread in their originally identified Gram-negative *Enterobacteriaceae*  
hosts but can also be found among many Gram-positive bacteria (Nandi *et al.*, 2004) . Here we  
demonstrated a possible direct way of accession of these Class 1 integrons in Gram-positive bacteria  
through IncP-1 $\epsilon$  plasmid transfer from Proteobacteria.

### **Ecological & Evolutionary Relevance**

355 Plasmid host range can be defined in several ways depending on the duration and intimacy of the  
considered plasmid-host relationship, including the transfer host range, the replication and maintenance



host range, or the evolutionary host range (Suzuki *et al.*, 2010). We show here that the immediate transfer range for IncP plasmids is much wider than previously reported, proving that in absence of physical barriers to cell-to-cell contact, broad host range plasmids have a high likelihood to be, hosted by very diverse  
360 bacteria, at least transiently.

However, comparative analysis of plasmid sequences has indicated that the evolutionary host range of IncP plasmids seems to be mostly limited to Proteobacterial classes (Suzuki *et al.*, 2010). This suggests that these plasmids are not maintained long enough outside of this phylum to be significantly affected by non-Proteobacterial genomes. Long-term evolutionary adaptation of the plasmid backbone to the new host, as  
365 known for IncP plasmids (Norberg *et al.*, 2011), might therefore also not take place. Poor maintenance of these plasmids in non-Proteobacterial hosts is the likely bottleneck explaining the difference between the very wide realized transfer range and the narrower evolutionary range. Mating pair formation and conjugation systems in these plasmids are evolutionary adapted to connect and span Gram-negative membranes. The observed transfer to Gram-positive bacteria might therefore become a dead end in many  
370 cases for Gram-negative associated plasmids if the Type IV coupling and secretion system cannot efficiently spread the plasmid to other neighboring bacteria. However, an actinobacterial *Mycobacterium* strain has been shown to host and transfer a IncP type plasmid indicating that maintenance and transfer is possible across the Gram border (Leão *et al.*, 2013). Also, the transient presence of a plasmid can provide the new host with a punctual adaptive gene pool and result in a short-term, but highly significant, fitness gain.

375 Accessory genes on plasmids are mostly arranged in transposons flanked by insertion sequence (IS) elements, which can recombine with the recipient bacterial chromosomes (e.g. Class 1 integron of pKJK5) delivering packages of fitness altering DNA without the need for plasmid replication. Additionally, transient hosts can increase the transfer range further by allowing transfer to organisms that had a lower transfer potential from the original donor strain (Yano *et al.*, 2013).

380 We show within a bacterial community that there is a high variability in permissiveness to broad-host range  
plasmids that cannot be explained by the phylogeny of the potential recipient. The ability to take up diverse  
broad host range plasmids from different hosts at high frequencies as represented by the super permissive  
fraction of the community has not previously been described. We do not know if it is a strain-specific trait  
and how environmental conditions affect its manifestation. Also, we do not know to what extent the  
385 employed mating conditions might have biased the observed pattern of super permissive plasmid  
recipients. But, if strain-specific, these super-permissive strains would be expected to play a  
disproportionate role as central nodes in networks of lateral gene acquisitions (Popa *et al.*, 2011). Most  
gene acquisitions occur between donors and recipients residing in the same habitat (Popa & Dagan, 2011)  
and while gene acquisition in nature mainly occurs within taxonomically homogenous groups, the  
390 heterogeneous soil community provides a hot-spot for gene acquisition from phylogenetically distant  
groups (Popa *et al.*, 2011). In soil a few strains build the core nodes of a heavily connected network of  
lateral gene acquisition (Popa *et al.*, 2011), which could be a possible indication of being part of the super-  
permissive fraction. These species are mainly found within Enterobacteriales (Gammaproteobacteria),  
Burkholderiales (Betaproteobacteria), and Staphylococci (Bacilli), groups that contain most of our super-  
395 permissive OTUs. Finding the same group of bacteria as central nodes in lateral gene transfer networks  
(Popa *et al.*, 2011) and as main contributors to plasmid flow in soil suggests that there is a indeed a link  
between increased plasmid uptake ability and long-term gene acquisition potential.

**Supplementary information is available at ISMEJ's website**

#### **Acknowledgements**

400 We thank S. M. Milani for assistance in FACS sorting and method development; J. Magid for access to the  
CRUCIAL field plot. This work was funded by the Villum Kann Rasmussen Foundation Center of Excellence  
CREAM (Center for Environmental and Agricultural Microbiology).

#### **Competing Financial Interest Statement**

All authors declare no competing financial interest.

## 405 **References**

Bahl MI, Hansen LH, Goesmann A, Sørensen SJ. (2007). The multiple antibiotic resistance IncP-1 plasmid pJK5 isolated from a soil environment is phylogenetically divergent from members of the previously established alpha, beta and delta sub-groups. *Plasmid* **58**:31–43.

410 Bahl MI, Oregaard G, Sørensen SJ, Hansen LH. (2009). Construction and use of flow cytometry optimized plasmid-sensor strains. *Methods Mol Biol* **532**:257–68.

Benveniste R, Davies J. (1973). Aminoglycoside Antibiotic-Inactivating Enzymes in Actinomycetes Similar to Those Present in Clinical Isolates of Antibiotic-Resistant Bacteria. *Proc Natl Acad Sci U S A* **70**:2276–2280.

415 Bingle LEH, Zatyka M, Manzoor SE, Thomas CM. (2003). Co-operative interactions control conjugative transfer of broad host-range plasmid RK2: full effect of minor changes in TrbA operator depends on KorB. *Mol Microbiol* **49**:1095–1108.

Brown Kav A, Sasson G, Jami E, Doron-Faigenboim A, Benhar I, Mizrahi I. (2012). Insights into the bovine rumen plasmidome. *Proc Natl Acad Sci U S A* **109**:5452–7.

420 Brzuszkiewicz E, Thürmer A, Schuldes J, Leimbach A, Liesegang H, Meyer F-D, *et al.* (2011). Genome sequence analyses of two isolates from the recent Escherichia coli outbreak in Germany reveal the emergence of a new pathotype: Entero-Aggregative-Haemorrhagic Escherichia coli (EAHEC). *Arch Microbiol* **193**:883–91.

D’Costa VM, McGrann KM, Hughes DW, Wright GD. (2006). Sampling the antibiotic resistome. *Science* **311**:374–7.

425 Dechesne A, Pallud C, Bertolla F, Grundmann GL. (2005). Impact of the microscale distribution of a Pseudomonas strain introduced into soil on potential contacts with indigenous bacteria. *Appl Environ Microbiol* **71**:8123–31.

Excoffier L, Smouse PE, Quattro JM. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**:479–491.

430 Finley RL, Collignon P, Larsson DGJ, McEwen SA, Li X-Z, Gaze WH, *et al.* (2013). The scourge of antibiotic resistance: the important role of the environment. *Clin Infect Dis* **57**:704–10.

Forsberg KJ, Reyes A, Wang B, Selleck EM, Sommer MOA, Dantas G. (2012). The shared antibiotic resistome of soil bacteria and human pathogens. *Science* **337**:1107–11.

435 De Gelder L, Vandecasteele FPJ, Brown CJ, Forney LJ, Top EM. (2005). Plasmid donor affects host range of promiscuous IncP-1beta plasmid pB10 in an activated-sludge microbial community. *Appl Environ Microbiol* **71**:5309–17.

- Grahn AM, Haase J, Bamford DH, Lanka E. (2000). Components of the RP4 conjugative transfer apparatus form an envelope structure bridging inner and outer membranes of donor cells: implications for related macromolecule transport systems. *J Bacteriol* **182**:1564–1574.
- 440 Halary S, Leigh JW, Cheaib B, Lopez P, Baptiste E. (2010). Network analyses structure genetic diversity in independent genetic worlds. *Proc Natl Acad Sci U S A* **107**:127–32.
- Hansen CHF, Krych L, Nielsen DS, Vogensen FK, Hansen LH, Sørensen SJ, *et al.* (2012). Early life treatment with vancomycin propagates *Akkermansia muciniphila* and reduces diabetes incidence in the NOD mouse. *Diabetologia* **55**:2285–94.
- 445 Heinemann JA, Sprague GF. (1989). Bacterial conjugative plasmids mobilize DNA transfer between bacteria and yeast. *Nature* **340**:205–9.
- Heuer H, Binh CTT, Jechalke S, Kopmann C, Zimmerling U, Krögerrecklenfort E, *et al.* (2012). IncP-1 $\epsilon$  Plasmids are Important Vectors of Antibiotic Resistance Genes in Agricultural Systems: Diversification Driven by Class 1 Integron Gene Cassettes. *Front Microbiol* **3**:2.
- 450 Heuer H, Ebers J, Weinert N, Smalla K. (2010). Variation in permissiveness for broad-host-range plasmids among genetically indistinguishable isolates of *Dickeya* sp. from a small field plot. *FEMS Microbiol Ecol* **73**:190–6.
- Hirt H. (2002). In Vivo Induction of Virulence and Antibiotic Resistance Transfer in *Enterococcus faecalis* Mediated by the Sex Pheromone-Sensing System of pCF10. *Infect Immun* **70**:716–723.
- 455 Holden MTG, Feil EJ, Lindsay JA, Peacock SJ, Day NPJ, Enright MC, *et al.* (2004). Complete genomes of two clinical *Staphylococcus aureus* strains: evidence for the rapid evolution of virulence and drug resistance. *Proc Natl Acad Sci U S A* **101**:9786–91.
- Holmsgaard PN, Norman A, Hede SC, Poulsen PHB, Al-Soud WA, Hansen LH, *et al.* (2011). Bias in bacterial diversity as a result of Nycodenz extraction from bulk soil. *Soil Biol Biochem* **43**:2152–2159.
- 460 Hong S-H, Bunge J, Jeon S-O, Epstein SS. (2006). Predicting microbial species richness. *Proc Natl Acad Sci U S A* **103**:117–22.
- Inoue D, Sei K, Soda S, Ike M, Fujita M. (2005). Potential of predominant activated sludge bacteria as recipients in conjugative plasmid transfer. *J Biosci Bioeng* **100**:600–605.
- Jain A, Srivastava P. (2013). Broad host range plasmids. *FEMS Microbiol Lett* **348**:87–96.
- 465 Joseph SJ, Hugenholtz P, Sangwan P, Osborne CA, Janssen PH. (2003). Laboratory Cultivation of Widespread and Previously Uncultured Soil Bacteria. *Appl Environ Microbiol* **69**:7210–7215.
- Leão SC, Matsumoto CK, Carneiro A, Ramos RT, Nogueira CL, Lima JD, *et al.* (2013). The detection and sequencing of a broad-host-range conjugative IncP-1 $\beta$  plasmid in an epidemic strain of *Mycobacterium abscessus* subsp. *bolletii*. Ahmed, N (ed). *PLoS One* **8**:e60746.
- 470 Letunic I, Bork P. (2007). Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and annotation. *Bioinformatics* **23**:127–8.

- Levy SSB, Marshall B. (2004). Antibacterial resistance worldwide: causes, challenges and responses. *Nat Med* **10**:S122–S129.
- Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. (2011). UniFrac: an effective distance metric for microbial community comparison. *ISME J* **5**:169–72.
- 475 Magid J, Luxhøi J, Jensen LS, Møller J. (2006). Establishment of a long-term field trial with urban fertilizers. In: Bruun, S., Raupp, J., Pekrun, C., Oltmanns, M. and Köpke, U. (eds.). *Is recycling of nutrients from urban areas to peri-urban organic farms feasible?*, ISOFAR Scientific Series, Bonn, Germany, Verlag Dr. H. J. Köster, pp. 59–78.
- 480 McKenzie GJ, Craig NL. (2006). Fast, easy and efficient: site-specific insertion of transgenes into enterobacterial chromosomes using Tn7 without need for selection of the insertion event. *BMC Microbiol* **6**:39.
- Musovic S, Dechesne A, Sørensen J, Smets BF. (2010). Novel assay to assess permissiveness of a soil microbial community toward receipt of mobile genetic elements. *Appl Environ Microbiol* **76**:4813–8.
- 485 Musovic S, Klümper U, Dechesne A, Magid J, Smets BF. (2014). Long-term manure exposure increases soil bacterial community potential for plasmid uptake. *Environ Microbiol Rep* **6**:125–30.
- Musovic S, Oregaard G, Kroer N, Sørensen SJ. (2006). Cultivation-independent examination of horizontal transfer and host range of an IncP-1 plasmid among gram-positive and gram-negative bacteria indigenous to the barley rhizosphere. *Appl Environ Microbiol* **72**:6687–92.
- 490 Nandi S, Maurer JJ, Hofacre C, Summers AO. (2004). Gram-positive bacteria are a major reservoir of Class 1 antibiotic resistance integrons in poultry litter. *Proc Natl Acad Sci U S A* **101**:7118–22.
- Norberg P, Bergström M, Jethava V, Dubhashi D, Hermansson M. (2011). The IncP-1 plasmid backbone adapts to different host bacterial species and evolves through homologous recombination. *Nat Commun* **2**:268.
- 495 Norman A, Hansen LH, Sørensen SJ. (2009). Conjugative plasmids: vessels of the communal gene pool. *Philos Trans R Soc Lond B Biol Sci* **364**:2275–89.
- Ochman H, Lawrence JG, Groisman EA. (2000). Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**:299–304.
- Popa O, Dagan T. (2011). Trends and barriers to lateral gene transfer in prokaryotes. *Curr Opin Microbiol* **14**:615–623.
- 500 Popa O, Hazkani-Covo E, Landan G, Martin W, Dagan T. (2011). Directed networks reveal genomic barriers and DNA repair bypasses to lateral gene transfer among prokaryotes. *Genome Res* **21**:599–609.
- Pukall R, Tschäpe H, Smalla K. (1996). Monitoring the spread of broad host and narrow host range plasmids in soil microcosms. *FEMS Microbiol Ecol* **20**:53–66.
- 505 Samuels AL, Lanka E, Davies JE. (2000). Conjugative Junctions in RP4-Mediated Mating of Escherichia coli. *J Bacteriol* **182**:2709–2715.

- Schloss PD, Gevers D, Westcott SL. (2011). Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PLoS One* **6**:e27310.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, *et al.* (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* **75**:7537–41.
- Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. (1994). Small mobilizable multi-purpose cloning vectors derived from the Escherichia coli plasmids pK18 and pK19: selection of defined deletions in the chromosome of Corynebacterium glutamicum. *Gene* **145**:69–73.
- Sengeløv G, Kristensen KJ, Sørensen AH, Kroer N, Sørensen SJ. (2001). Effect of Genomic Location on Horizontal Transfer of a Recombinant Gene Cassette Between Pseudomonas Strains in the Rhizosphere and Spermosphere of Barley Seedlings. *Curr Microbiol* **42**:160–167.
- Shintani M, Matsui K, Inoue J-I, Hosoyama A, Ohji S, Yamazoe A, *et al.* (2014). Single-cell analyses revealed transfer ranges of IncP-1, IncP-7, and IncP-9 plasmids in a soil bacterial community. *Appl Environ Microbiol* **80**:138–45.
- Sogin ML, Morrison HG, Huber JA, Mark Welch D, Huse SM, Neal PR, *et al.* (2006). Microbial diversity in the deep sea and the underexplored “rare biosphere”. *Proc Natl Acad Sci U S A* **103**:12115–20.
- Suzuki H, Yano H, Brown CJ, Top EM. (2010). Predicting plasmid promiscuity based on genomic signature. *J Bacteriol* **192**:6045–55.
- Sørensen SJ. (1993). Transfer of plasmid RP4 from Escherichia coli K-12 to indigenous bacteria of seawater. *Microb releases viruses, Bact fungi* **2**:135–41.
- Sørensen SJ, Bailey M, Hansen LH, Kroer N, Wuertz S. (2005). Studying plasmid horizontal transfer in situ: a critical review. *Nat Rev Microbiol* **3**:700–10.
- Thomas CM, Nielsen KM. (2005). Mechanisms of, and barriers to, horizontal gene transfer between bacteria. *Nat Rev Microbiol* **3**:711–21.
- Wang Q, Garrity GM, Tiedje JM, Cole JR. (2007). Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* **73**:5261–7.
- Waters VL. (2001). Conjugation between bacterial and mammalian cells. *Nat Genet* **29**:375–6.
- Wolk CP, Vonshak A, Kehoe P, Elhai J. (1984). Construction of shuttle vectors capable of conjugative transfer from Escherichia coli to nitrogen-fixing filamentous cyanobacteria. *Proc ...* **81**:1561–1565.
- Yano H, Rogers LM, Knox MG, Heuer H, Smalla K, Brown CJ, *et al.* (2013). Host range diversification within the IncP-1 plasmid group. *Microbiology* **159**:2303–15.
- Yue JC, Clayton MK. (2005). A Similarity Measure Based on Species Proportions. *Commun Stat - Theory Methods* **34**:2123–2131.
- Zhaxybayeva O, Doolittle WF. (2011). Lateral gene transfer. *Curr Biol* **21**:R242–6.

- 540 Zhu Y-G, Johnson TA, Su J-Q, Qiao M, Guo G-X, Stedtfeld RD, *et al.* (2013). Diverse and abundant antibiotic resistance genes in Chinese swine farms. *Proc Natl Acad Sci U S A* **110**:3435–40.

## Titles and legends to figures

545 **Figure 1:** Typical transconjugal microcolonies for plasmid pKJK5::*gfp* introduced through *E. coli* MG1655::*Km<sup>R</sup>-Lpp-mCherry*. Observation was carried out with the confocal laser scanning microscope (CLSM). Transconjugants are green fluorescent, due to *gfp*-expression. Gfp-repressing donor strains are red fluorescent through chromosomal *mCherry* tagging. Black background represents soil bacteria not taking part in gene transfer events.

550 **Figure 2:** FACS sorting of transconjugal cells from a mating mixture initiated with soil bacteria and *E. coli* carrying pKJK5. The procedure consists in three successive gates (marked by pink stars in line A): Gate I sorts for bacterial size based on front and side scatter; Gate II sorts for green fluorescent cells; Gate III selects only those green cells that possess no red fluorescence. Line A shows the sorting of the initial soil bacterial recipient community in absence of any donor strain and proves that the presence of green autofluorescent particles (A-II) does not yield false positive as they are excluded at the third gate, due to their red fluorescence (A-III). The sorting of a pure culture of the donor strain is shown in line B, where, again, no false positive events are recorded at the final gate. Line C represents the analysis of the mating mixture before sorting. Line D shows the enrichment of transconjugants after the first fast enrichment sorting step to over 80% transconjugal cells, with minor contamination by donor or soil particles. Line E shows how only pure transconjugants are obtained after the second purification sorting step.

**Figure 3:** Principal Coordinate Analysis of individual transconjugal pools, as well as of the extracted soil community (Soil\_Extract) and the reference soil community as grown on filters (Soil\_Filter) based on the ThetaYC algorithm (Yue & Clayton, 2005). Each axis explains a certain fraction of dissimilarity according to the axis loading given in brackets. The three different plasmids are represented by color. The three different donor strains are named next to the data points.

**Figure 4:** Phylogenetic tree showing all identified transconjugant OTUs for three different plasmids (pKJK5, RP4, pIPO2tet) from the same donor (*P. putida*). The colors of the branches mark different phylogenetic




groups. The three donor strains are shown in white letters in the trees. Green heatmap-circle around the tree represents the log transformed relative OTU abundance in the soil reference recipient community.

570 Three heatmap-circles in blue and red display the x-fold over- and underrepresentation of the OTU in the respective transconjugal pool in comparison to the abundance in the reference soil sample. Stars mark the shared (present in all 3 transconjugal pools) and abundant (present at more than 1% relative sequence abundance) transconjugant OTUs, which constitute the core super-permissive community fraction. Sample size was normalized to 30000 sequences per transconjugal pool.

575 **Figure 5:** Phylogenetic tree showing all identified transconjugant OTUs for the same plasmid (pKJK5) introduced through 3 different donor strains (*P.putida*; *Kluyvera sp.*; *E.coli*)(B). Colors of the branches mark different phylogenetic groups. The three donor strains are shown in white letters in the trees. Green heatmap-circle around the tree represents the log transformed relative OTU abundance in the soil reference recipient community. Three heatmap-circles in blue and red display the x-fold over- and  
580 underrepresentation of the OTU in the respective transconjugal pool in comparison to the abundance in the reference soil sample. Stars mark the shared (present in all 3 transconjugal pools) and abundant (present at more than 1% relative sequence abundance) transconjugant OTUs, which constitute constituting to the core super-permissive community fraction. Sample size was normalized to 30000 sequences per transconjugal pool.

585 **Figure 6:** Venn diagram of transconjugal pools for plasmid pKJK5 transferred from three different donor strains (*E. coli*, *P. putida* & *Kluyvera sp.*) (A&B) and for three different plasmids (pKJK5, RP4, pIPO2tet) introduced through *P. putida* into the soil community. Venn diagrams are presented for OTU incidence (C&D) and for OTU relative abundance (right, 100% represents the total number of transconjugal sequences). OTUs were defined at 97% sequence similarity and sequence sample size was normalized to  
590 30000 per transconjugal pool.

Figure 1:



595 Figure 2:











Figure 6:



610 **Supplementary Information File**

**Title:**

Deep sequencing of transconjugal pools reveals unexpectedly diverse bacterial community fraction receiving broad host range plasmids

615 **Author affiliations:**

Uli Klümper<sup>a</sup>, Leise Riber<sup>b</sup>, Arnaud Dechesne<sup>a</sup>, Analia Sannazzarro<sup>b</sup>, Lars H. Hansen<sup>b</sup>, Søren J. Sørensen<sup>b</sup>, Barth F. Smets<sup>a\*</sup>

<sup>a</sup> Technical University of Denmark, Department of Environmental Engineering, Miljøvej 113, 2840 Kgs. Lyngby, Denmark

620 <sup>b</sup> University of Copenhagen, Department of Biology, Microbiology, Universitetsparken 15, 2100 Copenhagen Ø, Denmark

\*corresponding author

**Subject category:**

Evolutionary genetics

625 **Corresponding author:**

Barth F. Smets

Technical University of Denmark

DTU Miljø

Miljøvej 113

630 2840 Kgs. Lyngby

Denmark

Ph. +45 45 25 22 30

Fax +45 45 93 28 50

E-mail bfm@env.dtu.dk

635



**Supplementary Information content:**

Supplementary Table 1: Number of FACS sorted transconjugant cells; raw and quality checked sequences for each mating combination and its replicates.

Supplementary Table 2: OTUs with more than 0.01% sequence representation in the soil reference  
640 community that were not represented in the transconjugal pools. Sequences were classified using the RDP classifier. Numbers behind the name indicate % sequence similarity to the closest representative.

Supplementary Figure 1: Phylum level distribution of the isolated and sequenced transconjugants in each of the 11 samples including all replicates per donor and plasmid combination.

Supplementary Figure 2: Relative over-/underrepresentation of OTUs in transconjugal pools compared with  
645 the soil reference recipient community as a function of the OTU's Sogin phylogenetic dissimilarity to the respective plasmid donor strain. Overrepresentation is calculated as relative abundance of the OTU in the transconjugal pool divided by relative abundance in soil reference community subtracted by 1.

Underrepresentation is calculated as the inverse of overrepresentation. Values given in the figure display the square root of the absolute over or underrepresentation.

650

Supplementary Table 1:

Donor	Plasmid	Replicate	Sorted Transconjugants	Raw reads	Sequences after mothur processing	Sequences/ Transconjugant sorted
<i>E.coli</i>	pKJK5	A	19000	65683	43227	2.28
<i>E.coli</i>	pKJK5	B	14500	57849	45870	3.16
<i>E.coli</i>	pKJK5	C	16500	60608	45292	2.74
<i>Kluyvera</i>	pKJK5	A	34500	52319	41510	1.20
<i>P.putida</i>	pKJK5	A	14000	47482	38931	2.78
<i>P.putida</i>	pKJK5	B	14000	63181	29894	2.14
<i>P.putida</i>	pIPO2tet	A	36500	55632	44156	1.21
<i>P.putida</i>	pIPO2tet	B	35500	65931	50398	1.42
<i>P.putida</i>	pIPO2tet	C	44500	57482	40027	0.90
<i>P.putida</i>	RP4	A	24000	52095	39972	1.67
<i>P.putida</i>	RP4	B	20000	61081	48986	2.45

Number of FACS sorted transconjugant cells, raw and quality checked sequences for each mating combination and its replicates.

Supplementary Table 2:

Phylum		Class		Order		Family		Genus	
Actinobacteria	100	Actinobacteria	100	Actinomycetales	100	unclassified	100	unclassified	100
Bacteroidetes	100	Sphingobacteria	100	Sphingobacteriales	100	Sphingobacteriaceae	100	Pedobacter	100
Bacteroidetes	100	Sphingobacteria	100	Sphingobacteriales	100	Sphingobacteriaceae	100	Sphingobacterium	100
Proteobacteria	100	Alphaproteobacteria	100	Rhizobiales	67	unclassified	67	unclassified	67
Proteobacteria	100	Alphaproteobacteria	100	unclassified	100	unclassified	100	unclassified	100
Proteobacteria	100	Betaproteobacteria	100	Burkholderiales	100	Burkholderiaceae	100	unclassified	100
Proteobacteria	100	Betaproteobacteria	100	Burkholderiales	100	Comamonadaceae	100	Delftia	100
Proteobacteria	100	Betaproteobacteria	100	Burkholderiales	100	Comamonadaceae	100	unclassified	100
Proteobacteria	100	Betaproteobacteria	100	Burkholderiales	100	Comamonadaceae	100	Variovorax	100
Proteobacteria	100	Deltaproteobacteria	100	Myxococcales	100	Myxococcaceae	100	Corallococcus	100
Proteobacteria	100	Deltaproteobacteria	100	Myxococcales	100	unclassified	100	unclassified	100
Proteobacteria	100	Gammaproteobacteria	100	Pseudomonadales	100	Pseudomonadaceae	100	Pseudomonas	93
Proteobacteria	100	Gammaproteobacteria	100	Pseudomonadales	100	Pseudomonadaceae	100	Pseudomonas	100
Proteobacteria	100	Gammaproteobacteria	100	Pseudomonadales	100	Pseudomonadaceae	100	Pseudomonas	93
Proteobacteria	100	Gammaproteobacteria	100	unclassified	100	unclassified	100	unclassified	100
Proteobacteria	100	Gammaproteobacteria	100	Xanthomonadales	100	Xanthomonadaceae	100	unclassified	61
Proteobacteria	100	unclassified	100	unclassified	100	unclassified	100	unclassified	100

OTUs with more than 0.01% sequence representation in the soil reference community that were not able to take up any of the introduced plasmids. Sequences were classified using the RDP classifier. Numbers behind the name displays percent of sequence similarity to the closest representative.

