

Burnout and Self-Reported Quality of Care in Community Mental Health

Michelle P. Salyers, Ph.D.

Associate Professor, Department of Psychology, Indiana University Purdue University Indianapolis (IUPUI); Co-Director, ACT Center of Indiana

Sadaaki Fukui, Ph.D.

Director of Research, The University of Kansas School of Social Welfare Office of Mental Health Research and Training

Angela L. Rollins, Ph.D.

Research Scientist, VA HSR&D Center on Implementing Evidence-based Practice, Roudebush VAMC; Research Director, ACT Center of Indiana; Assistant Research Professor, Department of Psychology, IUPUI

Ruth Firmin, B.A.

Doctoral Student in Clinical Psychology, Department of Psychology, Indiana University Purdue University Indianapolis (IUPUI)

Timothy Gearhart, MSW, LCSW

Vice President of Child and Adolescent Services, Four County Counseling Center, Logansport, IN

James P. Noll, Ph.D., HSPP

Director of Client Experience and Training, Four County Counseling Center, Logansport, IN

Stacy Williams, MBA

Business Office Manager, Four County Counseling Center, Logansport, IN

C.J. Davis, Psy.D.

Chief Executive Officer, Four County Counseling Center, Logansport, IN

Accepted for publication in *Administration and Policy in Mental Health and Mental Health Services Research*

Acknowledgements: This study was funded by an IP-RISP grant from the National Institute of Mental Health (R24 MH074670; Recovery Oriented Assertive Community Treatment). The content is solely the responsibility of the authors and does not represent the official views of the National Institutes of Health.

Address correspondence to:

Michelle P. Salyers, Ph.D.

Department of Psychology, IUPUI

402 N. Blackford Street

Indianapolis, IN 46202

(317) 274-2904 (tel.); Email: mpsalyer@iupui.edu

Accepted version; Final version published as Salyers, M. P., Fukui, S., Rollins, A. L., Firmin, R. L., Gearhart, T., Noll, J. P., Williams, S., & Davis, C.J. (2014). Burnout and Self-Reported Quality of Care in Community Mental Health. *Administration and Policy in Mental Health and Mental Health Services Research*.

Abstract

Staff burnout is widely believed to be problematic in mental healthcare, but few studies have linked burnout directly with quality of care. The purpose of this study was to examine the relationship between burnout and a newly developed scale for quality of care in a sample of community mental health workers (N=113). The Self-Reported Quality of Care scale had three distinct factors (Client-Centered Care, General Work Conscientiousness, and Low Errors), with good internal consistency. Burnout, particularly personal accomplishment, and to a lesser extent depersonalization, were predictive of overall self-reported Quality of Care, over and above background variables.

Key words: Burnout, quality of care, community mental health

Burnout and Self-Reported Quality of Care in Community Mental Health

Professionals working in community mental health are at high risk for experiencing staff burnout (Oddie & Ousley, 2007; Rohland, 2000; Siebert, 2005), which is frequently conceptualized as high levels of emotional exhaustion and depersonalization (negative or cynical attitudes), along with a diminished sense of personal achievement (Maslach, 1993). Burnout has been associated with negative consequences for mental health providers, including physical health problems (e.g., insomnia, headaches, poor overall health), relationship problems, reduced job satisfaction, and increased mental health problems (e.g., depression, anxiety, substance abuse) (Carson et al., 1999; Heaney, Price, & Rafferty, 1995; Killian, 2008; Long, Blackwell, & Midgley, 1990; Perrone, Aegisdottir, Webb, & Blalock, 2006; Rohland, 2000; Smoot & Gonzolas, 1995; Stalker & Harvey, 2002). Burnout is also problematic for organizations, and literature reviews frequently report increased absenteeism, tardiness, reduced job commitment, and, in some studies, poor job performance and increased turnover (Burke & Richardsen, 1993; Schwab, Jackson, & Schuler, 1986; Smoot & Gonzolas, 1995; Stalker & Harvey, 2002). Staff burnout is widely believed to impact quality of care; however, little direct research has examined burnout in relation to mental health services.

Quality of care provided in healthcare settings may involve a number of possible domains. Specifically in mental health, measurement of quality has been identified as “particularly challenging,” because of a lack of measurement tools, as well as a lack of consistent infrastructure and strategy for assessing quality

(Kilbourne, Keyser, & Pincus, 2010). While consensus on specific assessment approaches is lacking, the Institute of Medicine has highlighted six domains of quality, specifying that care should be safe, effective, patient-centered, timely, efficient, and equitable (Institute of Medicine, 2001). Quality of care could also be conceptualized as the vehicle by which mental health clinicians impact consumer outcomes. As an example, seminal meta-analytic work by Horvath and Symonds (1991) and later updated by others (Lambert & Barley, 2001; Martin, Garske, & Davis, 2000) have indicated a link between the quality of the therapeutic relationship (e.g., patient-centeredness and therapeutic alliance) and psychotherapy outcomes, over and above the use of specific therapeutic techniques. Thus, any influences of staff burnout on quality of care could be particularly important in impacting consumer outcomes.

Burnout could impact quality of care in a number of different ways. At the organizational level, burned out workers who take excessive time off or leave their jobs will disrupt the continuity of mental health care (Boyer & Bond, 1999). Further, studies have shown that burnout among some workers is likely to have a contagion effect, damaging the morale of other employees and leading to further staff turnover (Bakker & Demerouti, 2007; Bakker, Le Blanc, & Schaufeli, 2005). At the individual level, burnout has been associated with cognitive impairments, including decreased attention (van der Linden, Keijsers, Eling, & van Schaijk, 2005), which can lead to errors and less engagement. Burnout has been associated with decreased empathy (Astrom, Nilsson, Norberg, Sandman, & Winblad, 1991), and in a recent study of medical residents, reduced empathy mediated the relationship between burnout and

lower self-ratings of patient-centered care (Passalacqua & Segrin, 2012). Burnout has been associated with negative feelings about mental health consumers (Holmqvist & Jeanneau, 2006) and negative expectations about consumer recovery (Salyers, Brennan, & Kean, 2013). Negative staff attitudes, in turn, have been linked with poorer outcomes among consumers with serious mental illness, for example, in the area of employment (Gowdy, Carlson, & Rapp, 2003).

One indirect indicator of high quality care may be satisfaction with care, and several studies have examined the links between staff burnout and patient satisfaction in healthcare settings. Higher levels of burnout among doctors (Argentero, Dell'Olivo, & Ferretti, 2008; Halbesleben & Rathert, 2008) and nurses (Argentero et al., 2008; Leiter, Harvie, & Frizzell, 1998; McHugh, Kutney-Lee, Cimiotti, Sloane, & Aiken, 2011; Vahey, Aiken, Sloane, Clarke, & Vargas, 2004) have been associated with lower patient satisfaction. We found one study specifically examining burnout and satisfaction in community mental health. Garman and colleagues (Garman, Corrigan, & Morris, 2002) surveyed clinicians and consumers from 31 psychosocial rehabilitation programs. Using team-level aggregation, higher levels of emotional exhaustion were associated with less consumer satisfaction related to the environment, treatment, and autonomy; depersonalization was associated with less satisfaction with treatment; and low personal accomplishment was associated with less satisfaction with therapists.

A more rigorous indicator of high quality of care is improved consumer outcomes, and a few studies have gone beyond satisfaction to assess outcomes. Priebe and colleagues (Priebe et al., 2004) studied 24 assertive outreach teams

working with adults with severe mental illness in London. Team level burnout was associated with increased hospitalizations 9 months later; however, multivariate tests showed different findings, indicating other factors may moderate/mediate these relationships. Halbesleben and Rathert (2008) surveyed 178 matched pairs of physicians and patients who had been hospitalized in the prior year. Using path analysis, physician depersonalization was related to lower patient satisfaction and longer patient reported recovery times. More recently, a cross-sectional study of 89 outpatient substance abuse treatment facilities aggregated staff and consumer surveys at the level of the facility, finding burnout not directly related to treatment participation (Landrum, Knight, & Flynn, 2012). While the relationship between burnout and consumer satisfaction is generally consistent, research linking burnout and consumer engagement and outcomes are few and have mixed findings.

Another approach to assessing quality of care is by asking providers their perceptions of care, an alternative that may be less time and labor-intensive. In addition, staff-reported quality of care may be a good proxy measure when actual client outcomes or satisfaction data are not readily available. In studies of health care professions that empirically examine the relationship between burnout and self-reported quality of care, physicians reporting higher levels of burnout (emotional exhaustion and/or depersonalization) tend to report lower levels of quality of care (Gopal et al., 2005; Shanafelt, Bradley, Wipt, & Back, 2002; Williams, Manwell, Konrad, & Linzer, 2007). One study found that burnout mediated the relationship between work load and poor self-reported quality of care (Shirom, Nirel, & Vinokur, 2006). Nurses with high levels of burnout also tend to report lower quality of care

(Halbesleben et al., 2008; Van Bogaert, Meulemans, Clarke, Vermeyen, & Van de Heyning, 2009). We found only one study examining staff-reported quality of care and staff morale in the mental health field. Van Bogaert and colleagues (Van Bogaert, Clarke, Willems, & Mondelaers, 2012) surveyed psychiatric nurses in inpatient settings and found that better work engagement (conceptualized as the opposite of burnout) was associated with higher self-reported quality of care. To our knowledge, no studies have examined burnout in relation to staff-reported quality of care in community mental health settings.

In a related area of work, several studies have found higher levels of burnout were correlated with more self-reported errors among physicians (Fahrenkopf et al., 2008; Hayashino, Utsugi-Ozaki, Feldman, & Fukuhara, 2012; Prins et al., 2009; Shanafelt et al., 2010; Williams et al., 2007). One longitudinal study found a prospective relationship; residents with high levels of burnout were more likely to report having made an error 3 months later (West, 2006). The relationship between burnout and self-reported errors in nurses, however, is mixed. Laschinger and Leiter (2006) found that nurses with high levels of burnout reported more patient safety adverse events. In another study of nurses, burnout was not related to self-reported errors, but was related to frequency of reporting near misses -- though in the opposite direction; nurses with high emotional exhaustion were *less likely* to report near misses (Halbesleben et al., 2008).

We were interested in examining the relationship between burnout and self-reported quality of care among community mental health workers. Prior studies have used a variety of methods to assess self-reported quality of care, none specific

to community mental health care. For example, Van Bogaert and colleagues (Van Bogaert et al., 2012) asked three global indices of care (e.g., “The quality of care on the unit is fair or poor”). Shanafelt and colleagues (Shanafelt et al., 2002) developed an 8-item scale of more specific behaviors (e.g., “I found myself discharging patients to make the service ‘manageable’ because the team was so busy”) and attitudes (“I had little emotional reaction to the death of one of my patients”). Items from this measure have been used in other studies of medical doctors (Gopal et al., 2005; Williams et al., 2007). However, we know little about what constitutes quality of care as perceived by community mental health workers and how the constructs are related to burnout. Building on this prior work, we developed a scale that could be used to examine the relationship between quality of care and burnout in community mental health.

Because a new scale was created, we first examined psychometric properties through factor analysis, internal consistency, and correlations with related constructs. We then tested the relationship between burnout and quality. We hypothesized that community mental health staff with higher levels of emotional exhaustion and depersonalization, and lower levels of personal accomplishment, would report lower levels of quality of care.

Method

Participants

Employees at a Community Mental Health Center (CMHC) in a Midwestern city were invited to participate. As part of an ongoing program evaluation, participants completed a semi-annual staff survey that included scales to assess

burnout, job satisfaction, intentions to turnover, and expectations of consumers. We added a new scale to assess quality of care specifically developed for this study. All staff were invited to participate, not just direct clinical care providers. Of approximately 156 possible staff, 113 (72%) responded with complete data. Participants were primarily female (83%) and white (96%). Staff education levels included 36% with less than a bachelor's degree, 38% with a bachelor's, and 26% with a graduate degree. Staff reported spending an average of 50.2% (SD= 32.7) of their time in direct care, ranging from 0 to 100%. Staff also reported spending an average of 21.7% (SD = 28.2) of their time in administrative duties, ranging from 0 to 100%. The mean length of time at the agency was 6.7 years (SD = 6.2), ranging from a month to 29 years, and the mean length of time in the mental health field was 10.5 years (SD = 8.2), ranging from one month to 38 years.

Measures

The survey included limited demographic information, our primary independent variable (burnout), our primary dependent variable (Quality of Care), and other job-related variables of job satisfaction, intentions to turnover, and expectations of consumers.

Burnout was assessed with an adapted version of the Human Service Provider version of the Maslach Burnout Inventory (Maslach, 1996) that used 7-point anchors ranging from 1= 'strongly disagree' to 7 = 'strongly agree'. This widely-used scale measures three subscales: emotional exhaustion, depersonalization, and personal accomplishment. The subscales have shown good internal consistency, stability over time, and convergent validity with related constructs (Maslach, 1996).

Self-Reported Quality of Care. We created a quality of mental healthcare scale for this study. Items were generated by input from clinicians and administrators from the participating CMHC, listing areas of care that they thought could be impacted by provider well-being. We combined clinical input with ideas from scales that have been developed in other healthcare settings (Gopal et al., 2005; Shanafelt et al., 2002; Williams et al., 2007). Items were vetted and refined with input from four other mental health service researchers (all clinical psychologists), supporting face validity of the scale. The scale was pared to 25 items (see Table 1), and respondents were asked to report on the frequency of each occurring in the past month (0 = not at all to 5 = very often). Ten items were reverse scored so that higher numbers indicate better quality.

Job Satisfaction was assessed with 5 items from the Job Diagnostic Survey (Hackman & Oldham, 1974). The scale has good internal consistency (Hackman & Oldham, 1974) and evidence of convergent and divergent validity (Fried, 1991).

Intentions to turnover were assessed by two individual items: “How often have you seriously considered leaving your job in the past six months?” (1=never to 6=several times a week) and “How likely are you to leave your job in the next six months?” (1=not likely at all to 4=very likely). These have been used in other studies of mental health providers (Salyers, Hudson, et al., 2011; Salyers, Rollins, Kelly, Lysaker, & Williams, 2011).

Expectations of consumer recovery were measured with the Provider Expectations for Recovery Scale, a 10-item scale adapted from the 16-item Consumer Optimism Scale (Salyers, Tsai, & Stultz, 2007). Respondents are asked

to think about consumers they currently work with and to answer questions about how many consumers they expect to have specific outcomes (e.g., in housing, employment) on a 5-point scale ranging from 1 = almost all to 5 = none. The scale has been shortened to 10 items through Rasch analyses and has strong internal consistency ($\alpha = .91$) (Salyers et al., 2013).

Procedures

The research team attended an all-staff meeting in November 2012 and distributed surveys to staff members as they arrived. Participation was voluntary and anonymous, and staff could return completed surveys directly to the team throughout the morning. We randomly selected five participants to receive a \$25 gift card. The study and procedures were approved by [the university's] Institutional Review Board.

Analyses

An exploratory factor analysis with a principal axis factoring method and promax rotation was conducted to uncover the underlying structure of the Quality of Care (QOC) scale. Convergent validity of the scale was also examined via correlations with related constructs, including job satisfaction, turnover intentions (past six months and next six months), and provider expectations for consumer recovery. Finally, in order to test our main hypothesis that burnout is associated with lower QOC, we conducted a multiple regression analysis predicting total QOC scores and the three QOC subscale scores from burnout (emotional exhaustion, depersonalization, and personal accomplishment) controlling for demographics (gender, education level, length of time at the agency, length of time in the mental

health field, and job role [time spent in direct care, administration]). In order to account for missing data, a covariance matrix obtained by an Expectation-Maximization algorithm was used for the factor analysis and multiple imputation (30 imputed datasets) was utilized for the correlation and regression analyses (Enders, 2010). SAS ver. 9.2 was used for the analyses.

Results

The exploratory factor analysis retained 17 items and extracted three factors, which we labeled Client-Centered Care, General Work Conscientiousness, and Low Errors. See Table 1 for item descriptives and factor loadings. The number of factors were determined based on scree tests, Kaiser's criterion (i.e., eigenvalues greater than one), and interpretability. Eight items eliminated were those with either poor factor loading for all factors (i.e., $<.30$), cross-loadings, freestanding, or high correlations with other items. Kaiser-Meyer-Olkin measure of sampling adequacy was $.77$. The internal consistency (Cronbach's alphas) for each factor was $.88$, $.72$, and $.71$, respectively, and the total QOC was $.84$. As shown in Table 2, correlations between factors were low to moderate, ranging from $.05$ to $.46$. Client-Centered Care appeared relatively independent of the other two domains of General Work Conscientiousness ($r=.16$, $p=0.12$) and Low Errors ($r=.05$, $p=0.63$).

As hypothesized, burnout was associated with self-reported quality of care (also shown in Table 2). Total QOC was related to higher personal accomplishment and less depersonalization. Total QOC was also related to more positive expectations of consumer recovery. At the factor level, Client-Centered Care was positively correlated with more positive personal accomplishment and expectations of consumer recovery ($r=.29$, $p<.01$), but had almost no relationship to emotional exhaustion, job satisfaction, or turnover intentions. In contrast, General Work Conscientiousness was positively correlated with job satisfaction ($r=.24$, $p<.01$), and negatively correlated with emotional exhaustion, depersonalization, and turnover intentions in the past six months ($r=-.30$, $p<.01$) and the next six months ($r=-.29$,

$p < .01$). Errors were not significantly related to burnout, job satisfaction, turnover intentions, or expectations of consumer recovery.

The multiple regression analysis (Table 3) revealed that participants with higher personal accomplishment ($p < .001$), lower depersonalization ($p < .05$), and higher education (more than a bachelor's degree) ($p < .05$) had higher total QOC. Similarly, participants with higher personal accomplishment ($p < .001$) and higher education ($p < .05$) had higher Client-Centered Care. The overall regression models for two of the QOC subscales (i.e., General Work Conscientiousness and Low Errors) were not statistically significant ($p > .05$).

Discussion

This was the first study to assess self-reported quality of care (behaviors and attitudes) in community mental health and to link it with staff burnout. Our newly developed scale demonstrated good internal consistency and correlated with other job related variables (including job satisfaction, turnover intentions, and providers' expectations of consumer recovery). In addition, burnout was related to quality of care over and above background variables.

Quality of care factored into three distinct areas: client-centeredness, conscientiousness, and reporting low errors. Client-centeredness was comprised of items tapping into how the clinician is working directly with consumers, including attending to, interacting in a collaborative and compassionate way, and helping consumers with goals and treatment plans. In terms of the IOM definitions of quality (Institute of Medicine, 2001), this domain may most tap into being patient-centered,

effective, and equitable. The second factor of conscientiousness related to aspects outside of direct client care (missing deadlines or appointments, being late) and may fit best into areas of timeliness and efficiency. The third factor, assessing self-reported errors, aligns most closely with safety concerns.

The first two factors appeared to be the most robust, each with different patterns of relationships to other job-related variables. Conscientiousness was moderately related to a range of other variables including job satisfaction, turnover intentions, as well as emotional exhaustion and depersonalization. Conscientiousness may reflect a level of organizational commitment (i.e., to their employer) (Mowday, Steers, & Porter, 1979), whereas client-centered care is more indicative of a clinician's commitment to their consumers. Staff who are dissatisfied with their jobs and are considering leaving the organization, are not putting in as much effort into meeting deadlines and punctuality. Although we can't determine direction of relationships, this pattern would be consistent with the Conservation of Resources theory of burnout -- that being exhausted leads to a reduction of investment to conserve one's energy (Hobfoll, 1989). Conversely, client-centeredness was relatively independent of job satisfaction, turnover, and exhaustion. Even when experiencing work-related stress, mental health workers may be able to (or choose to) protect the quality of their direct clinical interactions.

The third factor (low errors) did not significantly correlate with other variables in this study. This factor was comprised of only two items, one of which had floor effects (reporting major mistakes in one's work). Although similar items have been used in other samples (Shanafelt et al., 2010; Shanafelt et al., 2002), it may be that

errors look different in a community mental health context or for different roles (e.g., case manager versus prescriber), or that reporting behavior differs in these settings. Alternatively, mistakes and errors reported in global self-ratings could simply be more vulnerable to biased reporting. Further work is needed to better assess errors or even “near misses” as they constitute important domains of care quality (Kohn, Corrigan, & Donaldson, 2000). Institutional-level incident reports of mistakes or near misses could be a better method for assessing safety aspects of care quality, although even those are reliant on clinician reports of incidents in most cases.

The findings linking burnout to quality of care in our sample show some consistency with prior work on burnout and self-reported quality of care in medical residents. For example, our bivariate correlations showed significant relationships between both emotional exhaustion and depersonalization with one aspect of quality -- lower general work conscientiousness (e.g., missing appointments, being late). This is similar to work showing depersonalization and/or emotional exhaustion are related to self-reported quality of care in residents (Gopal et al., 2005; Shanafelt et al., 2002; Williams et al., 2007). In our study, although depersonalization was related to overall quality of care, emotional exhaustion was not, and neither emotional exhaustion nor depersonalization were related to client-centered aspects of care quality (e.g., spending time with consumers, helping them meet goals).

Instead, personal accomplishment was the aspect of burnout most closely associated with both overall self-reported quality and client-centered care. Even after controlling for background variables, staff with higher levels of personal accomplishment reported higher levels of client-centered care. This is consistent

with a recent qualitative study of physicians who described the important role of patient-doctor relationships and feeling effective in their work as two key areas of gratification in their work (Zwack & Schweitzer, 2013). The direction of these relationships, however, is not clear. It may be that staff who have a sense of pride and accomplishment in their work are then more dedicated to the consumers they serve. But it may also be that providing client-centered care is rewarding in itself, leading to a greater sense of personal accomplishment. The other studies examining self-reported quality of care and burnout, however, did not examine personal accomplishment (Gopal et al., 2005; Shanafelt et al., 2002; Williams et al., 2007). Replication and further modeling in longitudinal studies is needed to sort out these relationships over time.

Taken together, an intriguing implication from our findings suggests different routes to improving quality of care through addressing staff burnout. Increasing positive affect (e.g., feelings of personal accomplishment) of staff may be more important for improving client-centered quality than decreasing negative affect (e.g., addressing emotional exhaustion and depersonalization). Conversely, for issues of conscientiousness, it may be more important to find ways to decrease the negative affect. These are hypotheses worth testing in the context of future burnout interventions. For example, one recent study found that a one-day stress management workshop (BREATHE) was associated with reduced emotional exhaustion and depersonalization, but not increased personal accomplishment (Salyers, Hudson, et al., 2011). It may be that the BREATHE intervention would

have more impact on conscientiousness, and other interventions may be needed to improve client-centeredness.

Our study used self-reported quality of care, and we are limited in our ability to comment on more objective indicators of care quality. For example, supervisor or consumer ratings of staff behaviors may be helpful in assessing quality of care being provided. In addition to perceived quality of care (by staff, supervisors, or consumers), actual consumer outcomes will be important to examine as well. There are many theoretical and practical reasons to believe that staff well-being and consumer well-being are related. However, research linking burnout with patient reported outcomes (Halbesleben & Rathert, 2008; Landrum et al., 2012) or more objective indicators such as hospital use (Priebe et al., 2004) is in the early stages. In addition, these types of indicators could also be used to better validate the new measure of quality of care. Finally, our study is limited in the sampling frame. We created the scale and tested it in one site that was comprised of predominantly white women, which limits generalizability of the findings. Additional studies are needed in other settings, with a more diverse sample of participants.

Although preliminary, the Self-Reported Quality of Care scale could be useful in future efforts to measure quality of care and understand how burnout impacts quality in mental health organizations. A strength of the scale is that it was developed in conjunction with stakeholders, as recommended by Kilbourne and colleagues (2010) as one step to overcome challenges in quality assessment in the mental health field. The scale could be administered as a user-friendly and economical tool for program evaluation intending to monitor and improve mental

health care quality at both individual and agency levels. In addition, the scale revealed different aspects of quality of care, which were associated with different aspects of burnout and other job-related constructs (including job satisfaction, turnover intentions, and expectations of consumer recovery). These are often concerns for administrators as indicators that may damage organizational climates. Future research is of course needed to validate the scale in other samples, and with other indicators of quality, but this study provides an initial foray into an important domain of inquiry.

Table 1. Descriptive information and factor loadings for Quality of Care items (N=113).

Item #	Label	Factor1	Factor2	Factor3	M	SD
<u>Client-Centered Care (Cronbach's alpha = 0.88)</u>						
19	I was able to help a client meet a personal or treatment goal.	.81	-.15	.03	3.98	0.92
21	I was able to come up with a creative intervention that helped a client.	.73	-.24	.23	3.51	0.97
23	I was able to develop a meaningful assessment or treatment plan in collaboration with a client.	.71	.00	.09	3.98	1.01
2	I spent extra time with a client who needed me.	.69	.04	-.25	4.08	0.97
18	I have involved clients in decision-making about their care.	.68	-.09	-.01	4.48	0.74
24	I went "above and beyond the normal call of duty" to serve a client.	.64	.12	.07	3.77	0.85
22	I felt like I was able to really show compassion to a client	.60	.08	.04	4.47	0.72
17	I have felt connected to the clients I am working with.	.54	.02	.00	4.30	0.87
25	The care I gave to my clients is as good or better than what they would get anywhere.	.52	.11	-.03	4.51	0.65
1	I collaborated with another staff person to help a client.	.50	.28	-.28	4.37	0.82
8	I have given my full attention to conversations with clients.	.42	.31	-.01	4.68	0.53
<u>General Work Conscientiousness (Cronbach's alpha = 0.72)</u>						
R_11	I have missed deadlines at work.	.00	.73	.16	4.23	0.74
R_10	I have missed appointments or meetings.	-.10	.63	.01	4.50	0.61
R_6	I have been late for work.	.05	.56	-.02	4.08	0.90
R_16	I have had difficulty prioritizing what needs to be done.	.11	.47	.13	3.98	0.85
<u>Low Errors (Cronbach's alpha = 0.71)</u>						
R_13	I have made minor mistakes in my work.	.00	.06	.67	3.51	0.60
R_14	I have made major mistakes in my work.	.01	.17	.65	4.44	0.61
<u>Items not included in final factors</u>						
3	I helped another staff member who needed me.				4.25	0.77
R_4	I have been distracted by my work.				3.14	0.98
R_5	I used paid time off.				3.15	0.93
7	I have met my collaborative service goals (e.g., productivity).				4.04	1.09
R_9	I have had a high rate of clients not showing up for appointments.				2.97	0.93
12	I have seen positive changes in my clients.				3.79	0.94
R_15	I have been directive with clients (telling them what to do).				3.20	1.15
20	I have been recognized/appreciated for good work.				3.06	1.11

Note: Items with R_ are reverse scored so that higher scores indicate higher quality.

Table 2. Correlations among Quality of Care, burnout, and other job-related variables (N=113)

	1	2	3	4	5	6	7	8	9	10	11
1. Total QOC	1										
2. Client-centered care	.94 ***	1									
3. General Work Conscientiousness	.46 ***	.16	1								
4. Low Errors	.25 **	.05	.26 **	1							
5. Job satisfaction	.09	.03	.24 **	-.05	1						
6. Emotional exhaustion	-.05	.02	-.22 *	-.01	-.62 ***	1					
7. Personal accomplishment	.50 ***	.53 ***	.09	.03	.22 *	-.25 **	1				
8. Depersonalization	-.25 **	-.18	-.24 *	-.13	-.44 ***	.50 ***	-.32 ***	1			
9. Turnover intent past 6 months	-.03	.09	-.30 **	-.12	-.63 ***	.58 ***	-.14	.42 ***	1		
10. Turnover intent next 6 months	-.05	.05	-.29 **	-.07	-.60 ***	.51 ***	-.05	.35 ***	.66 ***	1	
11. Provider expectations	.25 *	.29 **	-.04	-.01	.11	-.15	.15	-.12	-.11	-.01	1

Table 3. Summary of multiple regression results predicting Total QOC scores and the three subscale scores across 30 imputed datasets (N=113)

	TOTAL QOC Scores			Client-Centered Care			General Work Conscientiousness			Low Errors		
	B	SE	p	B	SE	p	B	SE	p	B	SE	p
<u>Demographic</u>												
Gender	0.13	0.08	0.13	0.10	0.08	0.21	0.11	0.11	0.30	0.05	0.11	0.68
Education	0.22	0.11	<.05*	0.23	0.10	<.05*	0.06	0.13	0.66	0.02	0.13	0.86
Tenure at Agency	-0.17	0.10	0.10	-0.18	0.10	0.06	-0.10	0.13	0.42	0.17	0.13	0.21
Tenure in Mental Health	0.03	0.10	0.75	0.07	0.09	0.48	<.01	0.12	0.99	-0.21	0.13	0.11
Time in Direct Care	0.20	0.12	0.10	0.22	0.12	0.06	-0.03	0.15	0.86	0.13	0.14	0.34
Time in Admin	0.06	0.12	0.61	0.03	0.11	0.76	0.07	0.13	0.62	0.08	0.12	0.53
<u>Burnout</u>												
Emotional Exhaustion	0.04	0.11	0.72	0.08	0.10	0.42	-0.14	0.13	0.29	0.04	0.14	0.77
Personal Accomplishment	0.36	0.10	<.001***	0.41	0.09	<.001***	0.02	0.11	0.89	<.01	0.11	0.97
Depersonalization	-0.19	0.09	<.05*	-0.14	0.09	0.13	-0.16	0.12	0.17	-0.16	0.12	0.19
intercept	<.01	0.08	0.98	<.01	0.07	0.98	<.01	0.09	0.99	<.01	0.10	0.99
	$F(9,103)=9.51$ [range 8.12-11.11], $p<.001$			$F(9,103)=11.56$ [range 10.14-14.64], $p<.001$			$F(9,103)=1.44$ [range 1.13-1.73], $n.s.$			$F(9,103)=0.80$ [range 0.67-1.04], $n.s.$		
	Ave. R2 = 45% (range 42%-49%)			Ave. R2 = 50% (range 47%-56%)			Ave. R2 = 11% (range 9%-13%)			Ave. R2 = 7% (range 6%-8%)		

References

- Argentero, P., Dell'Olivo, B., & Ferretti, M. S. (2008). Staff burnout and patient satisfaction with the quality of dialysis care. *American Journal of Kidney Disease, 51*(1), 80-92.
- Astrom, S., Nilsson, M., Norberg, A., Sandman, P. O., & Winblad, B. (1991). Staff burnout in dementia care--relations to empathy and attitudes. *International Journal of Nursing Studies, 28*(1), 65-75.
- Bakker, A. B., & Demerouti, E. (2007). The job demands-resources model: State of the art. *Journal of Managerial Psychology, 22*(3), 309-328.
- Bakker, A. B., Le Blanc, P. M., & Schaufeli, W. B. (2005). Burnout contagion among intensive care nurses. *Journal of Advanced Nursing, 51*(3), 276-287.
- Boyer, S. L., & Bond, G. R. (1999). Does assertive community treatment reduce burnout? A comparison with traditional case management. *Mental Health Services Research, 1*(1), 31-45.
- Burke, R. J., & Richardsen, A. M. (1993). Psychological burnout in organizations. In R. T. Golembiewski (Ed.), *Handbook of organizational behavior*. New York: Marcel Dekker.
- Carson, J., Cavagin, J., Bunclark, J., Maal, S., Gournay, K., & Kuipers, E., et al. (1999). Effective communication in mental health nurses; Did social support save the psychiatric nurse? *Nursing Times Research, 4*(1), 31-42.
- Enders, C. K. (2010). *Applied missing data analysis*. New York, NY: Guilford press.
- Fahrenkopf, A. M., Sectish, T. C., Barger, L. K., Sharek, P. J., Lewin, D., Chiang, V. W., . . . Landrigan, C. P. (2008). Rates of medication errors among depressed

- and burnt out residents: prospective cohort study. *BMJ: British Medical Journal (International Edition)*, 336(7642), 488-491.
- Fried, Y. (1991). Meta-analytic comparison of the Job Diagnostic Survey and Job Characteristics Inventory as correlates of work satisfaction and performance. *Journal of Applied Psychology*, 76(5), 690-697.
- Garman, A. N., Corrigan, P. W., & Morris, S. (2002). Staff burnout and patient satisfaction: evidence of relationships at the care unit level. *Journal of Occupational Health Psychology*, 7(3), 235-241.
- Gopal, R. K., Carreira, F., Baker, W. A., Glasheen, J. J., Miyoshi, T. J., Crane, L. A., & Prochazka, A. V. (2005). Does internal medicine resident burnout affect self-reported quality of care? *Journal of General Internal Medicine*, 20, 153-154.
- Gowdy, E. A., Carlson, L. S., & Rapp, C. A. (2003). Practices differentiating high-performing from low-performing supported employment programs. *Psychiatric Rehabilitation Journal*, 26, 232-239.
- Hackman, J. R., & Oldham, G. R. (1974). The Job Diagnostic Survey: An Instrument for the Diagnosis of Jobs and the Evaluation of Job Redesign Projects.
- Halbesleben, J. R., & Rathert, C. (2008). Linking physician burnout and patient outcomes: exploring the dyadic relationship between physicians and patients. *Health Care Management Review*, 33(1), 29-39.
- Halbesleben, J. R., Wakefield, B. J., Wakefield, D. S., Cooper, L. B., Halbesleben, J. R. B., Wakefield, B. J., . . . Cooper, L. B. (2008). Nurse burnout and patient

- safety outcomes: nurse safety perception versus reporting behavior. *Western Journal of Nursing Research*, 30(5), 560-577.
- Hayashino, Y., Utsugi-Ozaki, M., Feldman, M. D., & Fukuhara, S. (2012). Hope modified the association between distress and incidence of self-perceived medical errors among practicing physicians: Prospective cohort study. *PLoS ONE*, 7(4), 1-7.
- Heaney, C. C., Price, R. H., & Rafferty, J. (1995). The caregiver support program: An intervention to increase employee coping resources and enhance mental health. In L. R. Murphy & J. J. J. Hurrell (Eds.), *Job stress interventions* (pp. 93-108). Washington, DC: American Psychological Association.
- Hobfoll, S. E. (1989). Conservation of resources: A new attempt at conceptualizing stress. *American Psychologist*, 44(3), 513-524.
- Holmqvist, R., & Jeanneau, M. (2006). Burnout and psychiatric staff's feelings towards patients. *Psychiatry Research*, 145(2-3), 207-213.
- Horvath, A. O., & Symonds, B. D. (1991). Relation between working alliance and outcome in psychotherapy: A meta-analysis. *Journal of Counseling Psychology*, 38(2), 139.
- Institute of Medicine. (2001). *Crossing the Quality Chasm. Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington, D.C: National Academy Press.
- Kilbourne, A., Keyser, D., & Pincus, H. (2010). Challenges and opportunities in measuring the quality of mental health care. *Canadian Journal of Psychiatry*, 55(9), 549-557.

- Killian, K. D. (2008). Helping till it hurts? A multimethod study of compassion fatigue, burnout, and self-care in clinicians working with trauma survivors. *Traumatology, 14*, 32-44.
- Kohn, L. T., Corrigan, J. M., & Donaldson, M. S. (Eds.). (2000). *To err is human: building a safer health system*. (Vol. 627). Washington DC: National Academies Press.
- Lambert, M. J., & Barley, D. E. (2001). Research summary on the therapeutic relationship and psychotherapy outcome. *Psychotherapy: Theory, Research, Practice, Training, 38*(4), 357-336.
- Landrum, B., Knight, D. K., & Flynn, P. M. (2012). The impact of organizational stress and burnout on client engagement. *Journal of Substance Abuse Treatment, 42*(2), 222-230.
- Laschinger, H. K. S., & Leiter, M. P. (2006). The impact of nursing work environments on patient safety outcomes: The mediating role of burnout engagement. *Journal of Nursing Administration, 36*(5), 259-267.
- Leiter, M. P., Harvie, P., & Frizzell, C. (1998). The correspondence of patient satisfaction and nurse burnout. *Social Science & Medicine, 47*(10), 1611-1617.
- Long, C. G., Blackwell, C. C., & Midgley, M. (1990). An evaluation of two systems of in-patient care in a general hospital psychiatric unit II: Measure of staff and patient performance. *Journal of Advanced Nursing, 15*, 1436-1442.

- Martin, D. J., Garske, J. P., & Davis, M. K. (2000). Relation of the therapeutic alliance with outcome and other variables: a meta-analytic review. *Journal of Consulting and Clinical Psychology, 68*(3), 438-450.
- Maslach, C. (1993). Burnout: A multidimensional perspective. In W. B. Schaufeli, C. Maslach & T. Marek (Eds.), *Professional burnout: Recent developments in theory and research* (pp. 19-32). Washington, DC: Taylor & Francis.
- Maslach, C., Jackson, S. E., Leiter, M. P. (1996). *Maslach Burnout Inventory Manual* (3 ed.). Palo Alto, California: Consulting Psychologists Press.
- McHugh, M. D., Kutney-Lee, A., Cimiotti, J. P., Sloane, D. M., & Aiken, L. H. (2011). Nurses' widespread job dissatisfaction, burnout, and frustration with health benefits signal problems for patient care. *Health Affairs (Millwood), 30*(2), 202-210.
- Mowday, R. T., Steers, R. M., & Porter, L. W. (1979). The measurement of organizational commitment. *Journal of Vocational Behavior, 14*(2), 224-247.
- Oddie, S., & Ousley, L. (2007). Assessing burn-out and occupational stressors in a medium secure service. *The British Journal of Forensic Practice, 9*(2), 32-48.
- Passalacqua, S. A., & Segrin, C. (2012). The effect of resident physician stress, burnout, and empathy on patient-centered communication during the long-call shift. *Health Communication, 27*(5), 449-456.
- Perrone, K. M., Aegisdottir, S., Webb, L. K., & Blalock, R. H. (2006). Work-family interface: Commitment, conflict, coping and satisfaction. *Journal of Career Development, 32*, 286-300.

- Priebe, S., Fakhoury, W., White, I., Watts, J., Bebbington, P., Billings, J., . . . Wright, C. (2004). Characteristics of teams, staff and patients: associations with outcomes of patients in assertive outreach. *British Journal of Psychiatry, 185*, 306-311.
- Prins, J. T., van der Heijden, F., Hoekstra-Weebers, J., Bakker, A. B., van de Wiel, H. B. M., Jacobs, B., & Gazendam-Donofrio, S. M. (2009). Burnout, engagement and resident physicians' self-reported errors. *Psychology Health & Medicine, 14*(6), 654-666.
- Rohland, B. M. (2000). A survey of burnout among mental health center directors in a rural state. *Administration & Policy in Mental Health, 27*(4), 221-237.
- Salyers, M. P., Brennan, M., & Kean, J. (2013). Provider Expectations for Recovery Scale: Refining a measure of provider attitudes. *Psychiatric Rehabilitation Journal, 36*(3), 153-159.
- Salyers, M. P., Hudson, C., Morse, G., Rollins, A. L., Monroe-DeVita, M., Wilson, C., & Freeland, L. (2011). BREATHE: A pilot study of a one-day retreat to reduce burnout among mental health professionals. *Psychiatric Services, 62*(2), 214-217.
- Salyers, M. P., Rollins, A. L., Kelly, Y.-F., Lysaker, P. H., & Williams, J. R. (2011). Job satisfaction and burnout among VA and community mental health workers. *Administration and Policy in Mental Health and Mental Health Services Research, 40*(2), 1-7.
- Salyers, M. P., Tsai, J., & Stultz, T. A. (2007). Measuring recovery orientation in a hospital setting. *Psychiatric Rehabilitation Journal, 31*(2), 131-137.

- Schwab, R. L., Jackson, S. E., & Schuler, R. S. (1986). Educator burnout: Sources and consequences. *Educational Research Quarterly, 10*, 15-30.
- Shanafelt, T. D., Balch, C. M., Bechamps, G., Russell, T., Dyrbye, L., Satele, D., . . . Freischlag, J. (2010). Burnout and medical errors among American surgeons. *Annals Of Surgery, 251*(6), 995-1000.
- Shanafelt, T. D., Bradley, K. A., Wipt, J. E., & Back, A. L. (2002). Burnout and Self-Reported Patient Care in an Internal Medicine Residency Program. *Annals of Internal Medicine, 136*(5), 358.
- Shirom, A., Nirel, N., & Vinokur, A. D. (2006). Overload, autonomy, and burnout as predictors of physicians' quality of care. *Journal of Occupational Health Psychology, 11*(4), 328-342.
- Siebert, D. C. (2005). Personal and occupational factors in burnout among practicing social workers: Implications for research, practitioners, and managers. *Journal of Social Service Research, 32*(2), 25-44.
- Smoot, S. L., & Gonzolas, J. L. (1995). Cost-effective communication skills training for state hospital employees. *Psychiatric Services, 46*, 819-822.
- Stalker, C., & Harvey, C. (2002). Professional burnout: A review of theory, research, and prevention *Partnerships for Children and Families Project*. Waterloo: Wilfrid Laurier University.
- Vahey, D. C., Aiken, L. H., Sloane, D. M., Clarke, S. P., & Vargas, D. (2004). Nurse burnout and patient satisfaction. *Med Care, 42*(2 Suppl), I157-66.
- Van Bogaert, P., Clarke, S., Willems, R., & Mondelaers, M. (2012). Staff engagement as a target for managing work environments in psychiatric

- hospitals: implications for workforce stability and quality of care. *Journal of Clinical Nursing*, 22(11), 1717-1728.
- Van Bogaert, P., Meulemans, H., Clarke, S., Vermeyen, K., & Van de Heyning, P. (2009). Hospital nurse practice environment, burnout, job outcomes and quality of care: test of a structural equation model. *Journal Of Advanced Nursing*, 65(10), 2175-2185.
- van der Linden, D., Keijsers, G. P. J., Eling, P., & van Schaijk, R. (2005). Work stress and attentional difficulties: An initial study on burnout and cognitive failures. *Work & Stress*, 19(1), 23-36.
- West, C. P., Huschka M. M., Novotny P. J., Sloan, J. A., Kolars, J. C., Habermann, T. M. and Shanafelt, T. D. (2006). Association of perceived medical errors with resident distress and empathy: A prospective longitudinal study. *Journal of the American Medical Association*, 296(9), 1071-1078.
- Williams, E. S., Manwell, L. B., Konrad, T. R., & Linzer, M. (2007). The relationship of organizational culture, stress, satisfaction, and burnout with physician-reported error and suboptimal patient care: Results from the MEMO study. *Health Care Management Review July/September*, 32(3), 203-212.
- Zwack, J., & Schweitzer, J. (2013). If every fifth physician is affected by burnout, what about the other four? Resilience strategies of experienced physicians. *Academic Medicine*, 88(3), 382-389.