

OPEN

Cadmium and Mercury phytostabilization from soil using *Miscanthus* × *giganteus*

Zeljka Zgorelec^{1*}, Nikola Bilandzija^{2*}, Kristina Knez¹, Marija Galic¹ & Silva Zuzul³

The determination of the effects of cadmium and mercury on the growth, biomass productivity and phytoremediation potential of *Miscanthus* × *giganteus* (MxG) grown on contaminated soil was the main aim of this paper. The use of bioenergy plants as an innovative strategy in phytotechnology gives additional benefits, including mitigation and adaptation to climate change, and soil remediation without affecting soil fertility. An experiment was set up as a randomized complete block design with the treatments varied in concentrations of Cd (0, 10 and 100 mg kg⁻¹ soil) and Hg (0, 2 and 20 mg kg⁻¹ soil) added to the soil. Three vegetative years were studied. Yield values ranged from 6.3–15.5 t_{DM} ha⁻¹, cadmium concentration in plants varied from 45–6758 µg kg⁻¹ and Hg varied from 8.7–108.9 µg kg⁻¹. Values between treatments and years were significantly different. MxG can accumulate and remove very modest amount (up to 293.8 µg Cd and 4.7 µg Hg) per pot per year in aboveground biomass. Based on this data it can be concluded that MxG, as a valuable energy crop, is a potential candidate for the phytostabilization and biomass production on soils contaminated with Cd and Hg moderately.

Phytoremediation is considered as a simple and a natural technology that uses plants which can be utilize for efficient absorption of pollutants from contaminated soils^{1,2}. Generally, remediation of heavy metal polluted soils could be classified as physico-chemical and biological remediation techniques³. Compared to physico-chemical techniques (vitrification, soil washing, solidification and stabilization), phytoremediation technology could reduce dust emission, risk of exposure to soil, provide erosion control and prevent runoff⁴. Unlike physical and chemical treatments that irreversibly alter soil properties, phytoremediation generally improves physical, chemical, and biological quality of contaminated soils, improving soil quality and functionality and carbon sequestration^{5,6}. Phytoremediation is suitable for different types of contaminants (organic, metals and radionuclides), with relatively low financial costs, does not require additional energy delivery (energy is obtained from solar radiation) and with minimally influence to the site destruction and destabilization. Additionally, it contributes to the improvement of the visual aspect of the landscape, provides habitats for animals, and reduces leaching and mobilization of contaminants in soil⁷. Disadvantages of phytoremediation include: long remediation time requirement (the process is slow and requires 3–20 growing seasons to achieve remediation goals); relatively shallow cleaning depths; potentially contamination of the food chain; a site-specific technology (structure of the soil profile, pH, presence of salt or other toxins, water quality including nutrients, oxygen content) with a choice of plants being critical, and the fate of contaminants often unclear (the technology may relocate contaminants from the subsurface to the plant, creating residual waste to be disposed of); groundwater contamination possibility and climate dependence^{4,6–8}. The technology is applicable only to moderately contaminated land, it cannot compete with conventional remediation on heavily polluted sites. There is also concern about the content of toxic elements in the biomass of energy crops produced on a contaminated land that may generate hazardous emissions⁷. Based on removal mechanisms and type of pollutants, phytoremediation techniques can be categorized as phytoextraction, phytovolatilization, phytofiltration, phytostabilisation, phytodegradation and rhizosphere bioremediation⁹. Phytoremediation is technique that can be used for the reduction of bioavailability and mobility of heavy metals in soils^{1,2,10,11} and is defined as possibility of plants to increase retention of specific metals in root zone¹². Phytoextraction and phytostabilization are the most wide used remediation techniques, and differ in accumulation of elements in above ground or root parts of plants. Generally, advantages of phytostabilisation are effective

¹University of Zagreb Faculty of Agriculture, Agroecology Unit, Svetosimunska c. 25, 10000, Zagreb, Croatia.

²University of Zagreb Faculty of Agriculture, Agricultural Engineering and Technology Unit, Svetosimunska c. 25, 10000, Zagreb, Croatia. ³Institute for Medical Research and Occupational Health, Ksaverska c. 2, 10000, Zagreb, Croatia. *email: zzgorelec@agr.hr; nbilandzija@agr.hr

rapid immobilization and no need for biomass disposal, while major disadvantage is a fact that pollutants still remain in the soil¹³ or in the root system, generally in the rhizosphere. In comparison to phytoextraction, where pollutants are accumulated in the biomass, which is considered a possible problem afterwards. Phytostabilisation has proved to be useful for the treatment of Pb, As, Cd, Cr, Cu and Zn contaminated soils⁹. Phytoremediation is a very complex biotechnology, which is under the significant influence of (I) plant morphology (growth rate, biomass yields) and physiology (accumulation potential, stress tolerance)^{14,15}, (II) agro-ecological conditions of cultivated land (soil type and environment)^{16–18}, (III) agronomy practices (cultivars, planting density, soil amendments application)^{19–21}, (IV) origin of contamination^{22–24}. To increase the potential of phytoremediation, which is usually limited by low above ground biomass and/or a shallow root system^{25–27} more and more attention has been given to perennial plants with high biomass yield potential^{20,28}. The use of fast growing energy crops for purifying polluted lands is an innovative strategy to derive additional benefits from such remediation activities^{29–31}, it may have important role from ecological and energy point of view⁷. Due to the competition for arable land, water and nutrient resources, an implementation of energy crops in the phytoremediation strategy directly avoids the potential conflict between food and fuel production^{31–33}. The potential soil amendments and phytoremediation stimulants, including (I) mineral fertilizers²⁸, (II) farm manure³⁴, (III) organic wastes/biosolids (sewage sludge, compost)^{35,36}, (IV) solid bioproducts (biochar)¹⁹, (V) organic substances/biostimulants (mycorrhizal fungi)²¹ can be used to increase the biomass yield, the absorption potential of the plants, the amount of soil organic matter and to immobilize the metals in soil. Due to its morphological and physiological characteristics, one of the most investigated bioenergy plant for the purposes of remediation is *Miscanthus × giganteus* (MxG). Soils that have suffered from physical, biological and/or chemical degradation (i.e. soils contaminated with trace elements), or are uncultivated or/and adversely affected by climate conditions could be defined as marginal lands³⁷. Elbersen *et al.* mapped 29% of agricultural land in EU being marginal³⁸. Production of large quantities of biomass, thus providing the effective phytoremediation showed good potential of using *Miscanthus sp.* commercially on marginal sites in the regions of Central and Eastern Europe, and United States^{39,40}. Khalid reported that the most efficient remediation could be achieved with high biomass plants utilization¹¹. Nsanganwimana *et al.* highlighted MxG as a promising crop for the phytostabilization²¹, while Kerr characterized it as a tolerant phytoremediator in terms of growth on heavily polluted soils contaminated with Cu, Cd, Cr, Pb, Ni and Zn⁴¹. Fernando and Oliveira investigated the influence of heavy metals (Cd, Cu, Cr, Hg, Ni, Pb and Zn) on plant growth and productivity of MxG⁴². Although above mentioned authors recommend application of MxG plant species in phytoremediation, the higher heavy metal concentrations could negatively affect plant growth and productivity. Barbosa *et al.* noted that Zn contamination only reduced significantly MxG production, while not *M. sinensis* or *M. floridulus* yields and concluded that *Miscanthus* genotypes have shown different phytoremediation potential^{43,44}. Some authors^{21,40,45,46} classify MxG as excluder. Barbu *et al.* investigated the possibility of using MxG for Cd uptake and reported accumulation of 35–55 g Cd per hectare per year⁴⁷, and Nsanganwimana *et al.* revealed that majority of Cd accumulation was in roots of the plant²¹. Furthermore, on soils contaminated with 0.01 g Cd m⁻², *Miscanthus* accumulated 0.013 mg kg⁻¹ in its shoots⁴². Rosikon *et al.* observed positive influence of sewage sludge on Cd bioavailability by MxG²⁰ by investigating the influence of sewage sludge fertilization applied at different rates to *M. sacchariflorus* and compared with plant treatment by mineral fertilizers for uptake of different metals into stems and leaves during two years of observation. Cadmium was not detected in *M. sacchariflorus* biomass in the first year whereas large amounts of the metal were recorded in the second year (6–9 mg kg⁻¹)^{39,48–50}. The increased accumulation of Cd in MxG shoots with increasing Cd concentrations in the soil induced a reduction in plant height and shoot dry weight⁴⁰. *Miscanthus* showed low tolerance to Hg toxicity in terms of biomass productivity. It can be grown in fields contaminated with Hg only for soil remediation purposes, since economically might not be feasible due to decreased productivity. Effectively, with the increased Hg concentrations in the soil, biomass showed significantly higher accumulation of Hg, with lower biomass production, in comparison with the control⁴⁹. The average typical value of Cd in *Miscanthus* crops was reported 0.1 mg kg⁻¹ and for Hg was 0.03 mg kg⁻¹ and for grass in general 0.2 mg Cd kg⁻¹ and <0.02 mg Hg kg⁻¹⁵¹. The main aim of this paper was to determine the effects of cadmium and mercury on the growth, biomass productivity and phytoremediation potential of MxG grown on contaminated soil.

Materials and Methods

Experimental plot. The experiment was set up on 4th of March in 2014, in an open greenhouse in plastic experimental pots (EP). Weather conditions (sunlight duration, temperature and precipitation) were natural and soil moisture was controlled on daily basis and if necessary maintained to the field water capacity. According to the data of Croatian Meteorological and Hydrological Service (station Maksimir, Zagreb) mean annual values of precipitation (887.1 mm; 858.6 mm and 897.0 mm) and temperatures (12.6 °C; 12.2 °C and 12.6 °C) were noted for 3 studied vegetative years (2015, 2016 and 2017). The experiment was set up according to the completely randomized design and MxG was planted in four treatments in three replications during three years. High quality rhizomes (15 × 15 × 15 cm, cube of plant/soil) of three-year-old *Miscanthus* from Bistra field in the Republic of Croatia were planted in truncated cone plastic pot EP ($\varnothing_T = 28$ cm; $\varnothing_B = 19$ cm; h = 29 cm) where \varnothing_T and \varnothing_B present a top and bottom diameter of EP, respectively.

Preparation of contaminated soil. Four treatments (C, L₁, L₁ + SS, L₂) varied in concentration levels of Cd (0, 10 and 100 mg kg⁻¹) and Hg (0, 2 and 20 mg kg⁻¹) were applied to the soil. The first control group (C) consisted of pure soil. The soil in the second group was treated with lower level (L₁) of contaminants: 10 mg Cd kg⁻¹ (in CdO (s) form) and 2 mg Hg kg⁻¹ (in HgCl₂ (s) form). A third group (L₁ + SS) was treated with identical concentrations of Cd and Hg applied to soil as in L₁, but with an addition of sewage sludge in an equivalent of maximal 1.66 t_{DM} ha⁻¹ according to Croatian legislative⁵². Soil in the fourth group was treated with a higher level (L₂) of contaminants: 100 mg Cd kg⁻¹ and 20 mg Hg kg⁻¹ of soil. Contaminants were applied as p.a. salts in solid

Parameter	Protocol/Norm
Drying/grinding/milling/homogenizing	HRN ISO 11464:2009 ⁸⁰ at 60 °C to constant mass -> plant powder
w(ST), w(H ₂ O) [%]	HRN ISO 11465:2004 ⁶²
Cd and Hg extraction	HNO ₃ :H ₂ O ₂ = 5:1 (v/v) digestion in ratio 1:30 (m/v) 0.2 g -> 6 mL -> 50 mL
Cd [µg kg ⁻¹]	HRN ISO 11047:2004 ⁸¹ ISO/TS 16965:2013 ⁸²
Hg [µg kg ⁻¹]	HRN ISO 16772:2009 ⁸³

Table 1. Parameters and methods used in biomass analysis.

phase to dry soil before first vegetative year. Subsamples of clean soil were mixed with adequate amounts of salts to achieve homogeneity and then were vigorously mixed for a long time with the whole pot volume mass (~18 kg).

Soil characteristics. Soil used in the experiment was characterized as silt-loam texture (66.3% silt, 21.3% sand and 12.4% clay; sieving and sedimentation method were used⁵³) with acid reaction ($\text{pH}_{\text{KCl}} = 5.12$; obtained in 1 M KCl in 1:2.5 (m/v)⁵⁴) and having low content of organic matter (OM = 2.26%; determined by wet combustion method with sulphochromic oxidation⁵⁵). Soil was classified as Stagnosol⁵⁶. Soil was well supplied with total nitrogen (0.12%, determined by dry combustion (Dumas) method⁵⁷). All light elements (C, H, N and S) were analyzed by dry combustion method on Vario Macro CHNS analyzer, Elementar, 2006. Soil was low supply with plant available potassium (74 mg kg⁻¹) and phosphorous (26 mg kg⁻¹); (AL method; extraction with ammonium lactate acetic acid in 1:20 (m/v) ratio⁵⁸). CEC was 18.4 cmol⁺ kg⁻¹ (determined using barium chloride method in 1:40 (m/v) ratio⁵⁹). Total Cd and Hg in soil were measured in aqua regia extract⁶⁰ on AAS equipped with graphite and hydride technique (SOLAR AA Spectrometer M Series, Thermo Scientific, 2008 with Graphite Furnace and Cold Vapour System; see plant analysis, Table 1). Measured Cd concentration was 119 µg kg⁻¹, which was far below MAC (maximal allowable concentration) for agricultural soils (MAC = 1500 µg Cd kg⁻¹ for soils with pH value between 5 and 6) and measured Hg concentration was 66 µg kg⁻¹, which was also far below MAC value for agricultural soils (MAC = 1000 µg Hg kg⁻¹) according to Croatian legislative⁶¹.

Sewage sludge characteristics. Wastewater sewage sludge (SS) was characterized having neutral pH value ($\text{pH}_{\text{KCl}} = 7.54$ in 1 M KCl in 1:2.5 (m/v) ratio⁵⁴) and 63% of water⁶². Total carbon content was 22.7% (determined by dry combustion⁶³). Content of hydrogen was 9.52%, nitrogen 2.24%⁵⁷ and sulphur 0.36%⁶⁴. A total phosphorous content was 1.35% (extraction in aqua regia⁶⁰; and detection by ICP-OES⁶⁵, ICP-OES, Vista MPX Axial, Varian, 2004). The concentration of total cadmium in municipal waste water sewage sludge was 349 µg kg⁻¹ which was approximately 3 times higher than in soil, and concentration of mercury was 299 µg kg⁻¹, which was 4.5 times higher than in soil. Still, concentrations of Cd and Hg in sewage sludge were far below the permitted content of heavy metals in the sludge prescribed by Croatian law while used in agriculture (5000 µg kg⁻¹ for Cd and Hg⁵²).

Biomass sampling and growth parameters. The sampling of MxG was conducted at the beginning of March in 2015, 2016 and 2017 for each experimental pot. The biomass harvest was carried out by manual cutting of the plants at the height of 5 cm from the soil. Whole above ground biomass represents the sample with all dead leaves which were collected, if were any. Yield parameters, including plant height, shoot numbers per rhizome and mass of biomass with natural moisture content were determined on the site. Afterwards samples were cut to smaller pieces and transported to the Lab. The dry matter yield was determined gravimetrically after drying at 60 °C to the constant mass. Afterwards samples were milled to a powder and proceeded to digestion and metals analysis.

Biomass analysis. Table 1 shows methods used in the analysis of biomass. Aliquots of dried and homogenized plants (about 0.2 g) were wet digested with combination of HNO₃ and H₂O₂ (5:1 = (v/v)). All chemicals used in digestion were high purity (p.a.). Wet digestion was conducted in the digestion block (Velp, 2007) using the programme for Cd, 1 h at 65 °C and 3 h at 150 °C and for Hg, 4 h at 55 °C. The digest was diluted up to 50 mL with Milli-Q water. Mercury and cadmium detection in plants harvested in 2015 were determined with AAS (SOLAR AA Spectrometer M Series, Thermo Scientific, Autosampler, 2008). For Cd detection, graphite technique was used (Graphite Furnace, GF 95 + FS % Furnace) and for Hg detection hydride technique was used (Vapour System, VP100). In plants harvested in 2016 and 2017, Hg concentrations were detected with a PerkinElmer Flow Injection Mercury Hydride System, FIMS 400 with autosampler AS-91, 2006 and Cd concentrations (isotope 111) were determined by inductively coupled plasma mass spectrometry (ICP-MS 7500 cx, Agilent Technologies, Waldbronn, Germany) with rhodium as internal standard. The ICP-MS was tuned so that the oxides and doubly charged ions were less than 2% and each solution was analyzed in triplicate, in full quant mode, with helium as collision gas.

Statistical analysis and quality control. Statistical analysis was done with the use of statistical software SAS 9.1 (SAS Inst. Inc.), One-Way ANOVA and post-hoc (Fisher LSD) test were used for processing of data. The threshold of significance was 5% for all tests. Quality control was included. Measurement accuracy and method precision for Cd and Hg determination were checked using reference materials (IPE 171 and IPE 186 for plant and ISE 865 for soil, Wageningen University) and were satisfactory. Absolute error for Cd measurements was up




Figure 1. Yield of MxG ($t_{DM} ha^{-1}$) according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatment are not statistically significant (Fisher test).




Figure 2. Length of MxG plants according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatments are not statistically significant (Fisher test).

to maximal 8% and for Hg up to 5%, respectively. Relative standard deviation (RSD) or repeatability of measurement for Cd was up to maximum 7% and for Hg 8%, respectively.

Results and Discussion

Growth parameters. Results of the study including yield, length of plants, number of shoots regarding the treatment options and vegetative years are presented in Figs. 1 to 3. Statistical analysis of the results shows the influence of vegetative years on the yield for MxG by different treatment. The significant difference for C, L₁ + SS and L₂ treatments has been determined between the years of investigation (Fig. 1). In comparison to the first year of the study, the decrease of the MxG's yield was found in the second and third year, both on the control treatment and on the contaminated soil, in the range of 37% up to 55%. It is highly unlikely that the shoots will accumulate any significant amount of heavy metals from the soil in the first year of growth and have a significant impact on the yield, thus obtained values were expected. However, only in the control treatment the increase in the yield was observed in the third year. An interaction between MxG yield and treatments displays the statistically significant difference only in the third year of the research. It can be noticed that the highest yield was determined




Figure 3. Number of MxG shoots according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatments are not statistically significant (Fisher test).

in the control treatment C (7.3–15.5 $t_{DM} ha^{-1}$), and the lowest in the treatment L₂ (6.3–11.4 $t_{DM} ha^{-1}$). Even application of municipal sewage sludge, as a soil amendment was not resulting in biomass increase because the yield on sewage sludge treatment compared to L₁ was not statistically significant. Thus, no impact on soils slightly contaminated with Cd and Hg during three-year investigation period was observed. The negative impact of Cd and/or Hg on biomass yield was also determined by Arduini *et al.* and Fernando and Oliveira^{42,45}. Antonkiewicz *et al.* found a positive impact of sewage sludge on the yield, in five years long research⁶⁶. However, they applied up to 36 times higher doses of municipal sewage sludge (0–60 $t_{DM} ha^{-1}$) in comparison to our study. The average yield in their study was reported 15.3 $t_{DM} ha^{-1}$ in control treatment and 16.6 $t_{DM} ha^{-1}$ in treatment with sewage sludge, respectively. The length of the plant was not statistically influenced by the treatments, while significant differences can be seen between years of research for treatments C, L₁ + SS and L₂ (Fig. 2). If we compare first and third year, the increasing length of the plant in treatment C could be noticed; while this was not the case for other treatments. Length of the plant in this study ranged from 103 cm up to 172 cm. Fernando and Oliveira, and Arduini *et al.* determined the reduction of plant length in relation to the increase of Cd concentration in the soil^{42,67}. Zhang *et al.* noted that growth of *Miscanthus sacchariflorus* was significantly inhibited when Cd concentration in the soil was above 50 $mg kg^{-1}$ compared with control⁶⁸. Kocóń and Jurga were investigating shoot numbers and shoot length of MxG and *Sida hermaphrodita* on two different soil textures, including sandy and loamy soil contaminated with Cd, Cu, Ni, Pb and Zn of the second year of cultivation. They determined that MxG had greater number of shoots and lower shoot length compared to *Sida hermaphrodita* regardless of soil texture²⁸. However, Fernando and Oliveira did not observe the negative impact of Hg on the length of MxG plant⁴². There were no statistically significant differences between treatments and years for the number of MxG shoots (Fig. 3). The number of shoots per plant in this research ranged from 6 up to 11 and these observed values are lower compared to research conducted by Arduini *et al.* and Pogrzeba *et al.*^{45,69}. Like with plant height, Fernando and Oliveira observed that treatments with Hg in the soil did not have an impact on the number of shoots of MxG⁴².

Cadmium and mercury concentration in plants. Concentrations of cadmium and mercury in the aboveground biomass of MxG, in relation to treatments and years of research are shown in Figs. 4 and 5. Statistically significant influence of years and treatments on concentration of Cd and Hg in aboveground biomass is revealed. Cadmium concentration in plants varied from 45 $\mu g kg^{-1}$ (C, 2015) up to 6758 $\mu g kg^{-1}$ (L₂, 2016), and Hg varied from 8.7 $\mu g kg^{-1}$ (C and L₁) measured in 2015 up to 108.9 $\mu g kg^{-1}$ (L₂) observed 2016 (Figs. 4 and 5). Values of Cd in the first investigated year were low in all treatments, much lower than typical value (100 $\mu g kg^{-1}$) of Cd in *Miscanthus* according to HRN ISO 17225-1:2014⁵¹ (Solid Biofuels-Fuel specifications and Classes-Part 1: General requirements) while, in the second and third year of investigation the concentration of Cd in MxG were higher in C and much higher in all other contaminated treatments. In the second and third year of investigation, significantly higher concentrations of Cd in MxG are observed regarding rising doses of Cd in the soil, and correlation are completely positive⁷⁰ (Fig. 4), while, according to Pogrzeba *et al.*⁶⁹ Cd concentration where at similar levels during all three years of investigation. Arduini *et al.* investigated Cd in MxG stems, in relation to different concentrations of Cd in soil (0.75–3.00 $mg L^{-1}$), and revealed 275 up to 1237 μg of Cd per plant⁴⁵. In relation to *Sida hermaphrodita*, the MxG accumulate better Zn, Cu, Pb, Ni compared to Cd³⁹. Antonkiewicz *et al.* studied an application of different sludge doses (0–60 $t_{DM} ha^{-1}$) to the soil and reported the average (5 years) concentration of Cd in aboveground biomass from 90 to 180 $\mu g kg^{-1}$ ⁶⁶. Zhang *et al.* measured Cd concentration in *M. sacchariflorus* in aboveground part of 0.92, 2.28, 4.41, 6.26 and 18.36 $mg kg^{-1}$ on soil contaminated with 1, 5, 25, 50 and 100 $mg Cd kg^{-1}$, respectively⁶⁸. According to Rosikon *et al.*, different fertilize treatments have positive




Figure 4. Cd concentrations in MxG according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatments are not statistically significant (Fisher test).




Figure 5. Hg concentrations in MxG according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatments are not statistically significant (Fisher test).

influence on Cd accumulation by the MxG and negative by the *Phalaris arundinacea* L. Furthermore, same authors observed that MxG and *Phalaris arundinacea* L. have lower possibility of Cd accumulation in the second year of cultivation compared to the first year, what is opposite to the results of our investigation²⁰. The same as, for Cd, the highest concentrations of Hg in the MxG have been determined in the second year of research (Fig. 5). Statistically significant differences of Hg in the MxG are observed between years. For the first and second year of investigation, we noted statistically significant differences of Hg in the MxG between treatments too. All measured values of Hg in the MxG except those revealed at L₂, 2016 (108.9 µg kg⁻¹) were below typical (30 µg kg⁻¹) of Hg in the MxG according to HRN ISO 17225-1:2014⁵¹. Fernando and Oliveira investigated Hg concentration in the MxG aboveground biomass, cultivated on soils with two different levels of contamination (5000 and 6700 µg Hg m⁻²), and determined that contamination with lower Hg dose resulted with Hg in biomass below limit of detection of method (<LOD), while contamination of soil with higher Hg dose resulted with Hg concentration in the biomass of 4 µg kg⁻¹⁴². Pérez-Sanz *et al.* investigated mercury uptake by *Silene vulgaris*, grown on contaminated (5.5 mg Hg kg⁻¹) spiked soils (alkali and neutral pH) and observed that *S. vulgaris* retains more Hg in the root (3700 and 2900 µg kg⁻¹) than in aerial part (550 and 980 µg kg⁻¹)⁷¹. Still, plants grew healthy and showed good appearance throughout the study without significantly decrease in the biomass production. Hg values in the aboveground biomass of the MxG in this study are considerably lower than those expected. However, it is not surprising due to Hg as a specific element and its behavior being a bit different than all other heavy metals. Mercury has been known as an environmental pollutant for over a century and it is well known that it may evaporate (volatilization) from some compounds and be released to various ecosystems. When added to the soil, whether in elemental, inorganic or organic form, it is likely to be strongly bond. Generally, 97–99% of total Hg is in complex form, and behavior of Hg species in the soil is controlled by soil factors, especially temperature, pH,




Figure 6. Cd biomass removals by MxG according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatments are not statistically significant (Fisher test).




Figure 7. Hg biomass removals by MxG according to treatments and studied years. Mean values marked with the same capital letters between the years and the same small letters between treatments are not statistically significant (Fisher test).

texture, organic matter content but also the concentration of all other ions. Phytoavailability and toxicity of Hg in the soil-plant system depend on the forms in soil⁷². Lomonte *et al.* study has shown that Hg accumulated by *C. zizanioides* via root uptake is mainly present in the root epidermis and exodermis and its translocation to the aerial parts is insignificant⁷³. In contrast, another study done by Lomonte *et al.* shows that some species (*Atriplex conodocarpa* and *Australodanthonia caespitose*) can be good candidates for mercury phytoextraction because of their ability to translocate mercury from roots to the aboveground tissues⁷⁴. Lomonte *et al.* also study Hg behavior in soils and Hg efflux to atmosphere⁷⁵. They applied biosolids (3.5–8.4 mg Hg kg⁻¹) from waste water treatment plant to soil and investigated potential for Hg remediation. They observed that 59% of the total mercury was complexed with organic ligands and that the influence of water content and irradiation on the emission of gaseous elemental mercury are the main factors affecting this emission with flux values up to 132 ngm⁻² h⁻¹. Lomonte *et al.* revealed that some ions mobilize Hg in the soil, creating chelate-assisted phytoextraction for some species and increase its uptake in the plant shoots⁷⁶. Pogrzeba *et al.* also studied Hg behavior in contaminated soil with addition of granular sulphur and observed that in Hg stressed environment, plant (*Poa pratensis*) developed the defense mechanism resulting in the reduction of Hg evaporation and higher S content in plant tissue⁷⁷. Those authors recommended this technology for soil remediation heavily contaminated with mercury.

Cadmium and mercury biomass removal. Cadmium and Hg biomass removals by MxG according to treatments and studied years are presented in Figs. 6 and 7. Statistically significant interaction between years of research and Cd removal is determined for all treatments, with the highest biomass removal determined in the second year. In the first year of investigation, Cd biomass removal between the treatments is not statistically significant and values determined below 10 µg pot⁻¹. In the second and third year of investigation, significant increase in Cd biomass removal is noted and is in a complete positive correlation with Cd concentration in soil (Fig. 6)⁷⁰. Values of Cd removal ranged from 70.8 up to 293.8 µg per pot (11.8–49.0 g ha⁻¹) observed on contaminated treatments (L₁ and L₂), in the second and third year of investigation, respectively. This is in accordance

to Barbu *et al.* who determined an uptake of 35–55 g Cd ha⁻¹⁴⁷. Bang *et al.* noted limited Cd accumulation by *Miscanthus* in a marginally contaminated ecosystem, although they observed 100% of Cd removal from contaminated water after 40 days⁷⁸. Yao *et al.* observed Cd accumulation in plant tissue of 2.2 mg Cd m⁻² (22 g ha⁻¹) in *Miscanthus sacchariflorus* and 700 kg Cd per year where it was accumulated by aboveground organs and removed from the lake (Dongting Lake wetlands, China) through harvesting for paper manufacture⁷⁹. Hg biomass removals by the MxG according to treatments and studied years varied up to 4.7 µg per pot (0.79 g ha⁻¹) in our study. Statistically significant influence of years on Hg removal has been determined for treatments L₁ + SS and L₂. In terms of interaction between treatments, significant Hg removal is observed only in the second year of research in treatment L₂ (Fig. 7).

Conclusion

Values of Cd in the MxG in first investigated year were low in all treatments, much lower than the typical value when MxG is used as a biofuel (100 µg kg⁻¹). The significantly higher concentrations of Cd in the MxG were observed in the second and third year of investigation due to increased doses of Cd in the soil, and correlations were completely positive.

The detected mercury concentration in MxG plants was very low. The whole measured values of Hg in MxG except those of treatment L₂, 2016 (108.9 µg kg⁻¹) were below typical for MxG used as biofuel (30 µg kg⁻¹).

Finally, it can be concluded that MxG, as a valuable energy crop, could be a good candidate for the Hg and Cd phytostabilization, due to the low metal accumulation in aboveground biomass. This could be benefit for biomass production of MxG on soils moderately contaminated with Cd and Hg, where contamination still not significantly affected the yields amounts.

Except phytostabilization of the contamination, the MxG can also be used in locations where some other remediation strategies of ecosystem/agroecosystem need to be achieved, like prevention of soil erosion due to the high biomass above but also below ground.

Data availability

The raw datasets generated and/or analyzed during this study are available from the corresponding authors upon request.

Received: 23 October 2019; Accepted: 1 April 2020;

Published online: 21 April 2020

References

- Raskin, I., Smith, R. D. & Salt, D. E. Phytoremediation of metals: using plants to remove pollutants from the environment. *Curr. Opin. Biotechnol.* **8**, 221–226 (1997).
- Shin, M. N. *et al.* Characterization of lead resistant endophytic *Bacillus* sp. MN3-4 and its potential for promoting lead accumulation in metal hyperaccumulator *Alnus firma*. *J. Hazard. Mater.* **199**, 314–320 (2012).
- Ashraf, S., Ali, Q., Zahir, Z. A., Ashraf, S. & Asghar, H. N. Phytoremediation: Environmentally sustainable way for reclamation of heavy metal polluted soils. *Ecotoxicol. Environ. Saf.* **174**, 714–727 (2019).
- Green, C. & Hoffnagle, A. Phytoremediation field studies database for chlorinated solvents, pesticides, explosives and metals. <http://www.cluin.org/download/studentpapers/hoffnagle-phytoremediation.pdf> (2019).
- Liu, L., Li, W., Song, W. & Guo, M. Remediation techniques for heavy metal-contaminated soils: Principles and applicability. *Sci. Tot. Environ.* **633**, 206–219 (2018).
- Kisic, I., Zgorelec, Z. & Percin, A. Soil treatment engineering. *Phy. Sci. Rev.* **2**, 1–26 (2017).
- Hauptvogel, M. *et al.* Phytoremediation Potential of Fast-Growing Energy Plants: Challenges and Perspectives – a Review. *Pol. J. Environ. Stud.* **29**, 505–516 (2020).
- Yeung, A. T. Remediation Technologies for Contaminated Sites. *Advances in Environmental Geotechnics. Proc. of Int. Symp. on Geoenvironmental Eng.* **8–10**, 328–369 (2010).
- Kumar Yadav, K. *et al.* Mechanistic understanding and holistic approach of phytoremediation: A review on application and future prospects. *Ecol. Eng.* **120**, 274–298 (2018).
- Sylvain, B. *et al.* Phytostabilization of As, Sb and Pb by two willow species (*S. viminalis* and *S. purpurea*) on former mine technosols. *Catena*. **136**, 44–52 (2016).
- Khalid, S. *et al.* A comparison of technologies for remediation of heavy metal contaminated soils. *J. Geochem. Explor.* **182**, 247–268 (2017).
- Mahar, A. *et al.* Challenges and opportunities in the phytoremediation of heavy metals contaminated soils: A review. *Ecotoxicol. Environ. Saf.* **126**, 111–121 (2016).
- Ghosh, M. & Singh, S. P. A review on phytoremediation of heavy metals and utilization of its byproducts. *Appl. Ecol. Environ. Res.* **3**, 1–18 (2005).
- He, Z. L., Yang, X. E. & Stoffella, P. J. Trace elements in agroecosystems and impacts on the environment. *J. Trace. Elem.* **19**, 125–140 (2005).
- Malik, R. N., Husain, S. Z. & Nazir, I. Heavy metal contamination and accumulation in soil and wild plant species from industrial area of Islamabad, Pakistan. *Pak. J. Bot.* **42**, 291–301 (2010).
- Robson, D. B., Knight, J. D., Farrell, R. E. & Germida, J. J. Ability of cold-tolerant plants to grow in hydrocarbon-contaminated soil. *Intern. J. Phytoremediation.* **5**, 105–113 (2003).
- Sun, T. R. *et al.* Roles of abiotic losses, microbes, plant roots, and root exudates on phytoremediation of PAHs in a barren soil. *J. Hazard. Mater.* **176**, 919–925 (2010).
- Guala, S. D., Flora, A. V. & Covelo, E. F. The dynamics of heavy metals in plant–soil interactions. *Ecol. Modell.* **22**, 1148–1152 (2010).
- Hartley, W., Dickinson, N. M., Riby, P. & Lepp, N. W. Arsenic mobility in brownfield soils amended with green waste compost or biochar and planted with *Miscanthus*. *Environ. Pollut.* **157**, 2654–2662 (2009).
- Rosikon, K., Fijalkowski, K. & Kacprzak, M. Phytoremediation Potential of Selected Energetic Plants (*Miscanthus giganteus* L. and *Phalaris arundinacea* L.) in Dependence on Fertilization. *J. Environ. Eng. Sci.* **4**, 587–595 (2015).
- Nsanganwimana, F. *et al.* Metal accumulation and shoot yield of *Miscanthus × giganteus* growing in contaminated agricultural soils: Insights into agronomic practices. *Agric. Ecosyst. Environ.* **213**, 61–71 (2015).
- Alexander, M. *Biodegradation and Bioremediation*. 2nd ed. Academic Press, San Diego (1999).

23. Binet, P., Portal, J. M. & Leval, C. Dissipation of 3–6 ring polycyclic aromatic in the rhizosphere of ryegrass. *Soil. Biol. Biochem.* **32**, 2011–2017 (2000).
24. Hechmi, N., Nadhira, B. A., Hassen, A. & Naceur, J. Phytoremediation potential of maize (*Zea mays* L.) In co-contaminated soils with pentachlorophenol and cadmium. *Int. J. Phytoremediation.* **15**, 703–713 (2013).
25. Dushenkov, V., Kumar, P. B., Motto, H. & Raskin, I. Rhizofiltration: The Use of Plants to Remove Heavy Metals from Aqueous Streams. *Environ. Sci. Technol.* **29**, 1239–1245 (1995).
26. Pilon-Smits, E. & Pilon, M. Phytoremediation of Metals Using Transgenic Plants. *Crit. Rev. Plant. Sci.* **21**, 439–456 (2002).
27. Pulford, I. D. & Watson, C. Phytoremediation of Heavy Metal-Contaminated Land by Trees: A Review. *Environ. Int.* **29**, 529–540 (2003).
28. Kocoń, A. & Jurga, B. The evaluation of growth and phytoextraction potential of *Miscanthus × giganteus* and *Sida hermaphrodita* on soil contaminated simultaneously with Cd, Cu, Ni, Pb, and Zn. *Environ. Sci. Pollut. Res.* **24**, 4990–5000 (2017).
29. Figala, J., Vranova, V., Rejsek, K. & Formanek, P. Giant *Miscanthus* (*Miscanthus × Giganteus* Greef et. Deu.) – A Promising Plant for Soil Remediation: A Mini Review. *Acta. Univ. Agric. Silv. Mendel. Brun.* **63**, 2241–2246 (2015).
30. Prelac, M., Bilandzija, N. & Zgorelec, Z. The phytoremediation potential of heavy metals from soil using Poaceae energy crops: A review. *J. Cent. Eur. Agr.* **17**, 901–916 (2016).
31. Tripathi, V., Edrisi, S. A. & Abhilash, P. C. Towards the coupling of phytoremediation with bioenergy production. *Renew. Sust. Energ. Rev.* **57**, 1386–1389 (2016).
32. Edrisi, S. A. & Abhilash, P. C. Exploring marginal and degraded lands for biomass and bioenergy production: An Indian scenario. *Renew. Sust. Energ. Rev.* **54**, 1537–1551 (2016).
33. Pandey, V. C., Bajpai, O. & Singh, N. Energy crops in sustainable phytoremediation. *Renew. sust. energy. rev.* **54**, 58–73 (2016).
34. Walker, D. J., Clemente, R., Roig, A. & Pilar Bernal, M. The effects of soil amendments on heavy metal bioavailability in two contaminated Mediterranean soils. *Environ. Pollut.* **122**, 303–312 (2003).
35. Kolodziej, B. *et al.* Use of sewage sludge in bioenergy production – a case study on the effects on sorghum biomass production. *Eur. J. Agron.* **69**, 63–74 (2015).
36. Grobelak, A. Organic Soil Amendments in the Phytoremediation Process. *Phytorem.* 21–39 (2016).
37. Blanco-Canqui, H. Growing Dedicated Energy Crops on Marginal Lands and Ecosystem Services. *Soil Sci. Soc. Am. J.* **80**, 845–858 (2016).
38. Elbersen, B. *et al.* Mapping Marginal land potentially available for industrial crops in Europe. Paper presented at 26th European Biomass Conference & Exhibition, Wageningen (2019).
39. Pidlisnyuk, V., Stefanovska, T., Lewis, E. E., Erickson, L. E. & Davis, L. C. *Miscanthus* as a Productive Biofuel Crop for Phytoremediation. *Crit. Rev. Plant. Sci.* **33**, 1–19 (2014).
40. Korzeniowska, J. & Stanisławska-Głubiak, E. Phytoremediation potential of *Miscanthus × giganteus* and *Spartina pectinata* in soil contaminated with heavy metals. *Environ. Sci. Pollut. Res.* **22**, 11648–11657 (2015).
41. Kerr, J. Phytoremediation of metal-contaminated soils by industrial crops. Dissertation, University of Glasgow (2003).
42. Fernando, A. L. & Oliveira, J. S. Effects on growth, productivity, and biomass quality of *Miscanthus × giganteus* of soils contaminated with heavy metals. Paper presented at 2nd World Congress on Biomass for Energy, Industry, and Climate Protection, Rome, Italy (2004).
43. Barbosa, B. *et al.* Phytoremediation of Heavy Metal-Contaminated Soils Using the Perennial Energy Crops *Miscanthus* spp. and *Arundo donax* L. *Bioenerg. Res.* **8**, 1500–1511 (2015).
44. Krzyżak, J. *et al.* Heavy Metal Uptake by Novel *Miscanthus* Seed-Based Hybrids Cultivated in Heavy Metal Contaminated Soil. *Civ. environ. eng.* **26**, 121–132 (2017).
45. Arduini, I., Ercoli, L., Mariotti, M. & Masoni, A. Response of *Miscanthus* to toxic cadmium applications during the period of maximum growth. *Environ. Exper. Bot.* **55**, 29–40 (2006).
46. Pavel, P. B., Puschenreiter, M., Wenzel, W. W., Diacu, E. & Barbu, C. H. Aided phytostabilization using *Miscanthus sinensis × giganteus* on heavy metal-contaminated soils. *Sci. Total. Environ.* **479**, 125–131 (2014).
47. Barbu, C. H., Pavel, P. B., Sand, C. & Pop, M. R. *Miscanthus sinensis* behaviour on soil polluted with heavy metals: Metal Elements in Environment, Medicine and Biology. Paper presented at 9th International Symposium of Romanian Academy Branch, Cluj-Napoca, Romania (2009).
48. Kalembsa, D. & Malinowska, E. Influence of sewage sludge fertilization on heavy metal content in biomass of *Miscanthus sacchariflorus* during field experiment. *Environ. Protect. Engin.* **35**, 149–155 (2009a).
49. Kalembsa, D. & Malinowska, E. The yield and content of trace elements in biomass of *Miscanthus sacchariflorus* (Maxim.) Hack. and in soil in the third year of a pot experiment. *J. Elementology.* **14**, 685–691 (2009b).
50. Kalembsa, D. & Malinowska, E. Influence of after effect of waste activated sludge used in soil of pot experiment on heavy metals content in *Miscanthus sacchariflorus*. *Acta. Agrophys.* **13**, 377–384 (2009c).
51. HRN ISO 17225-1 Solid biofuels - Fuel specifications and classes - Part 1: General requirements. International Organization for Standardization, Geneva (2014).
52. Official Gazette NN 38. Ministry of Environment and Energy. Regulation on the protection of agricultural land from pollution (in Croatian). 1307, Zagreb, Croatia. https://narodne-novine.nn.hr/clanci/sluzbeni/2008_04_38_1307.html (2018).
53. HRN ISO 11277 Soil quality - Determination of particle size distribution in mineral soil material — Method by sieving and sedimentation. International Organization for Standardization, Geneva (2009).
54. Modified HRN ISO 10390 Soil quality - Determination of pH. International Organization for Standardization, Geneva (2004).
55. Modified HRN ISO 14235 Soil Quality - Determination of organic carbon by sulfochromic oxidation. International Organization for Standardization, Geneva (2004).
56. World reference base for soil resources (WRB) IUSS Working Group 2014 (Update 2015). International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106, (FAO, Rome, 2015).
57. HRN ISO 13878 Soil quality - Determination of total nitrogen content by dry combustion (“elemental analysis”). International Organization for Standardization, Geneva (2004).
58. Škorić, A. Handbook of Soil Science Research. University of Zagreb Faculty of Agriculture, Zagreb, in Croatian, (1982).
59. HRN ISO 11260 Soil quality - Determination of effective cation exchange capacity and base saturation level using barium chloride solution. International Organization for Standardization, Geneva (2004).
60. HRN ISO 11466 Soil quality. Extraction of trace elements soluble in aqua regia. International Organization for Standardization, Geneva (2004).
61. Official Gazette NN 71. Ministry of Agriculture. Regulation on the protection of agricultural land from pollution (in Croatian). 1507, Zagreb, Croatia. https://narodne-novine.nn.hr/clanci/sluzbeni/2019_07_71_1507.html (2019).
62. HRN ISO 11465 Soil quality. Determination of dry matter and water content on a mass basis - Gravimetric method. International Organization for Standardization, Geneva (2004).
63. HRN ISO 10694 Soil quality - Determination of organic and total carbon after dry combustion (elementary analysis). International Organization for Standardization, Geneva (2004).
64. HRN ISO 15178 Soil quality - Determination of total sulfur by dry combustion. International Organization for Standardization, Geneva (2005).

65. HRN ISO 22036 Soil quality - Determination of trace elements in extracts of soil by inductively coupled plasma - atomic emission spectrometry (ICP - AES). International Organization for Standardization, Geneva (2011).
66. Antonkiewicz, J., Kołodziej, B. & Bielińska, E. J. The use of reed canary grass and giant miscanthus in the phytoremediation of municipal sewage sludge. *Environ. Sci. Pollut. Res.* **23**, 9505–9517 (2016).
67. Arduini, L., Masoni, A., Mariotti, M. & Ercoli, L. Low cadmium application increase miscanthus growth and cadmium translocation. *Environ. Exper. Bot.* **52**, 89–100 (2004).
68. Zhang, J., Yang, S., Huang, Y. & Zhou, S. The Tolerance and Accumulation of Miscanthus Sacchariflorus (Maxim.) Benth., an Energy Plant Species, to Cadmium. *Int. J. Phytoremediation.* **17**, 538–545 (2015).
69. Pogrzeba, M., Krzyżak, J. & Sas-Nowosielska, A. Environmental hazards related to Miscanthus × giganteus cultivation on heavy metal contaminated soil. *E3S Web of Conferences.* **1**, 1–4 (2013).
70. Vasilj, D. Biometrics Experimentation in Plant Growth. University of Zagreb: Croatian Agronomy Society PRESS (In Croatian) (2000).
71. Pérez-Sanz, A. *et al.* Mercury uptake by Silene vulgaris grown on contaminated spiked soils. *J. Environ. Manage.* **95**, 233–237 (2012).
72. Kabata-Pendias, A. & Mukherjee, A. B. Trace elements from Soil to Human, Springer (2007).
73. Lomonte, C. *et al.* Study of the Spatial Distribution of Mercury in Roots of Vetiver Grass (Chrysopogon zizanioides) by Micro-Pixe Spectrometry. *Int. J. Phytoremediation.* **16**, 1170–1182 (2014).
74. Lomonte, C., Doronila, A. I., Gregory, D., Baker, A. J. M. & Kolev, S. D. Phytotoxicity of biosolids and screening of selected plant species with potential for mercury phytoextraction. *J. Hazard. Mater.* **173**, 494–501 (2010a).
75. Lomonte, C. *et al.* Assessment of the pollution potential of mercury contaminated biosolids. *Environ. Chem.* **7**, 146–152 (2010b).
76. Lomonte, C., Doronila, A., Gregory, D., Baker, A. J. M. & Kolev, S. D. Chelate-assisted phytoextraction of mercury in biosolids. *Sci. Tot. Environ.* **409**, 2685–2692 (2011).
77. Pogrzeba, M., Ciszek, D., Galimska-Stypa, R., Nowak, B. & Sas-Nowosielska, A. Ecological strategy for soil contaminated with mercury. *Plant. Soil.* **49**, 371–387 (2016).
78. Bang, J. *et al.* Phytoremediation of Heavy Metals in Contaminated Water and Soil Using Miscanthus sp. Goedae-Uksae 1. *Int. J. Phytoremediation.* **17**, 515–520 (2015).
79. Yao, X. *et al.* Heavy metal bioaccumulation by Miscanthus sacchariflorus and its potential for removing metals from the Dongting Lake wetlands, China. *Environ. Sci. Pollut. R.* **25**, 20003–20011 (2018).
80. HRN ISO 11464 Soil quality - Pretreatment of samples for physico-chemical analysis. International Organization for Standardization, Geneva (2009).
81. HRN ISO 11047 Soil quality. Determination of cadmium, chromium, cobalt, copper, lead, manganese, nickel and zinc - Flame and electrothermal atomic absorption spectrometric methods. International Organization for Standardization, Geneva (2004).
82. ISO/TS 16965 Soil quality. Determination of trace elements using inductively coupled plasma mass spectrometry (ICP-MS). International Organization for Standardization, Geneva (2013).
83. HRN ISO 16772 Soil quality. Determination of mercury in aqua regia soil extracts with cold-vapour atomic spectrometry or cold-vapour atomic fluorescence spectrometry. International Organization for Standardization, Geneva (2009).

Acknowledgements

Publication was supported by the OpenAccess Publication Fund of the University of Zagreb Faculty of Agriculture.

Author contributions

Zeljka Zgorelec conceived and designed the experiment, contributed to data analysis (statistic), data interpretation, and wrote the manuscript. Bilandzija Nikola conceived and designed the experiment, data interpretation, wrote the manuscript. Knez Kristina collected the data, performed the analysis (sample preparation). Galic Marija collected the data, performed the soil analysis (soil agrochemical and physical analysis), technical editing of the manuscript (figures and references). Zuzul Silva performed the analysis (metals in soil and plant samples). All authors reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Z.Z. or N.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020