

## RESEARCH ARTICLE

**Caffeic Acid Inhibited Receptor Activator of Nuclear Factor  $\kappa$ B Ligand (RANKL)-Tumor Necrosis Factor (TNF)  $\alpha$ -TNF Receptor Associated Factor (TRAF) 6 induced Osteoclastogenesis Pathway**Ferry Sandra<sup>1,2,\*</sup>, Toshio Kukita<sup>3</sup>, Tatsushi Muta<sup>4</sup>, Tadahiko Iijima<sup>5</sup><sup>1</sup>Department of Biochemistry and Molecular Biology, Faculty of Dentistry, Trisakti University, Jl. Kyai Tapa No.260, Jakarta, Indonesia<sup>2</sup>Prodia Clinical Laboratory, Jl. Kramat Raya No.150, Jakarta, Indonesia<sup>3</sup>Laboratory of Oral Cellular and Molecular Biology, Division of Oral Biological Sciences, Faculty of Dental Science, Kyushu University, Fukuoka, Japan<sup>4</sup>Department of Molecular and Cellular Biochemistry, Graduate School of Medical Science, Kyushu University, Fukuoka, Japan<sup>5</sup>Laboratory of Oral and Maxillofacial Morphology, Division of Oral Biological Sciences, Faculty of Dental Science, Kyushu University, Fukuoka, Japan

\*Corresponding author. E-mail: ferrysandra@gmail.com

## Abstract

**BACKGROUND:** Caffeic acid was reported in our previous study to have potential in inhibiting osteoclastogenesis through inhibition of nuclear factor  $\kappa$ B (NF $\kappa$ B). Here in our current study, we would like to investigate further the caffeic acid-affected signaling pathway leading to NF $\kappa$ B inhibition. Since tumor necrosis factor (TNF) receptor-associated factor 6 (TRAF6) plays important role in osteoclastogenesis, we applied TRAF6-transfected RAW264 cells D-Clone (RAW-D) cells as model in this study.

**METHODS:** Caffeic acid in various concentrations was added to in vitro osteoclastogenesis of receptor activator nuclear factor  $\kappa$ B ligand (RANKL)-TNF $\alpha$ -induced TRAF6-transfected RAW-D cells. Cells were collected, lysed and immunoblotted to detect TRAF6 expression. To detect tartrate resistant acid phosphatase (TRAP)<sup>+</sup> polynucleated cells (PNCs), TRAP staining was performed. Meanwhile, to measure NF $\kappa$ B Activity, cells were transfected with pNF $\kappa$ B-TA-Luc and subjected to Dual Luciferase Reporter Assay System.

**RESULTS:** Caffeic acid did not influence TRAF6 expression of RANKL-TNF $\alpha$ -induced TRAF6-transfected RAW-D cells. Caffeic acid diminished NF $\kappa$ B activity of RANKL-TNF $\alpha$ -induced TRAF6-transfected RAW-D cells in a concentration dependent manner. Significant NF $\kappa$ B activity inhibitions were seen under treatment of 1 and 10  $\mu$ g/ml caffeic acid. By adding 10  $\mu$ g/ml caffeic acid in

## Abstrak

**LATAR BELAKANG:** Potensi *caffeic acid* dalam menghambat osteoklastogenesis dengan inhibisi *nuclear factor  $\kappa$ B* (NF $\kappa$ B) telah dilaporkan pada penelitian kami sebelumnya. Pada penelitian kali ini, kami meneliti pengaruh *caffeic acid* pada jalur persinyalan yang mengarah pada inhibisi NF $\kappa$ B. Oleh karena *tumor necrosis factor* (TNF) *receptor-associated factor 6* (TRAF6) berperan penting dalam osteoklastogenesis, kami menggunakan sel *RAW264 D-Clone* (RAW-D) yang ditransfeksi dengan TRAF6 sebagai model penelitian.

**METODE:** *Caffeic acid* dalam berbagai konsentrasi ditambahkan pada osteoklastogenesis *in vitro* dengan menggunakan induksi *receptor activator nuclear factor  $\kappa$ B ligand* (RANKL)-TNF $\alpha$  pada sel RAW-D. Sel dikoleksi dan dilisiskan, serta dilakukan imunoblot untuk mendeteksi ekspresi TRAF6. Untuk mendeteksi sel polinuklear (SPN) *tartrate resistant acid phosphatase* (TRAP)<sup>+</sup>, dilakukan pengecatan TRAP. Sedangkan untuk memeriksa aktivitas NF $\kappa$ B, sel ditransfeksi dengan *pNF $\kappa$ B-TA-Luc* dan diperiksa menggunakan *Dual Luciferase Reporter Assay System*.

**HASIL:** *Caffeic acid* tidak mempengaruhi ekspresi TRAF6 pada sel RAW-D yang diinduksi dengan RANKL-TNF $\alpha$  dan ditransfeksi dengan TRAF6. Aktivitas NF $\kappa$ B pada sel RAW-D yang diinduksi dengan RANKL-TNF $\alpha$  dan ditransfeksi dengan TRAF6, dihambat oleh *caffeic acid* dalam ketergantungan dengan konsentrasi. Inhibisi aktivitas NF $\kappa$ B yang bermakna terlihat pada pemberian *caffeic acid*

RANKL-TNF $\alpha$ -induced TRAF6-transfected RAW-D cells, TRAP<sup>+</sup> PNCs number was significantly suppressed.

**CONCLUSION:** Caffeic acid inhibited RANKL-TNF $\alpha$ -TRAF6-induced osteoclastogenesis pathway. Since caffeic acid did not influence TRAF6 expression, TRAF6-RANK interactions and/or TRAF6 downstream signaling pathway should be further pursued to disclose inhibition mechanism of caffeic acid.

**KEYWORDS:** caffeic acid, osteoclastogenesis, TRAF6, RANKL, TNF $\alpha$ , NF $\kappa$ B, RAW-D

*Indones Biomed J. 2013; 5(3): 173-8*

dengan konsentrasi 1 dan 10  $\mu$ g/ml. Dengan menambahkan *caffeic acid* 10  $\mu$ g/ml pada sel RAW-D yang diinduksi dengan RANKL-TNF $\alpha$  dan ditransfeksi dengan TRAF6, jumlah SPN TRAP<sup>+</sup> berkurang secara bermakna.

**KESIMPULAN:** *Caffeic acid* menghambat jalur osteoklastogenesis yang diinduksi oleh RANKL-TNF $\alpha$ -TRAF6. Oleh karena *caffeic acid* tidak mempengaruhi ekspresi TRAF6, sebaiknya dilakukan penelitian lebih lanjut pada interaksi TRAF6-RANK dan/atau jalur persinyalan di bawah TRAF6, sehingga mekanisme hambatan *caffeic acid* dapat diketahui.

**KATA KUNCI:** *caffeic acid*, osteoklastogenesis, TRAF6, RANKL, TNF $\alpha$ , NF $\kappa$ B, RAW-D

## Introduction

Osteoclast is a unique bone-resorbing cell derived from the pluripotent hematopoietic stem cell, which gives rise to a myeloid stem cell that can further differentiate into megakaryocytes, granulocytes, monocytes/macrophages and osteoclasts.(1) Loss of osteoclast function or differentiation will inhibit bone marrow cavity formation, an osteopetrotic phenotype.(2) However, osteoclasts is dispensable for HSC mobilization and may function as negative regulators in the hematopoietic system.(2)

Osteoclastogenesis is a complicated procedure that includes many stages, such as commitment, differentiation, multinucleation, and activation of immature osteoclasts.(1) Receptor activator of nuclear factor  $\kappa$ B ligand (RANKL), a member of the tumor necrosis factor (TNF) superfamily, is a type 2 transmembrane residing protein, which activates receptor activator of nuclear factor  $\kappa$ B (RANK) on osteoclast precursors in a trimeric symmetric complex.(1,3) RANKL induces osteoclast differentiation(3), interacting with an adaptor molecule TNF receptor-associated factor 6 (TRAF6)(1,4). Meanwhile TNF $\alpha$ , which is a member of the TNF ligand superfamily and a multifunctional cytokine, induces osteoclast differentiation and bone resorption by activating osteoclast progenitors.(5) TNF $\alpha$  did not effectively induce osteoclast differentiation from osteoclast progenitor cells derived from TRAF6-deficient mice into mature multinucleated osteoclasts, although c-jun N-terminal kinase (JNK) and TNF $\alpha$  activation was observed in osteoclast progenitor cells.(5) Thus, TRAF6 is involved in TNF $\alpha$ -induced osteoclastogenesis.(5)

TRAF6 plays essential roles in both the differentiation and maturation of osteoclasts by activating various kinases via its multiple domains.(6) TRAF6-deficient mice show severe osteopetrosis due to impaired osteoclastogenesis.(1) Upon recruitment of TRAF6, nuclear factor  $\kappa$ B (NF $\kappa$ B) and c-Jun N-terminal kinase (JNK) pathways will be further activated.(4) Mapping of the structural requirements for TRAF/RANK interaction revealed multiple TRAF binding sites clustered in two distinct domains in the RANK cytoplasmic tail, which are functionally important for the RANK-dependent induction of NF $\kappa$ B and JNK activities.(7)

In our attempt to search for a natural agent having potent activity in inhibiting osteoclastogenesis, we found that sweet potato leaf extract could inhibit osteoclastogenesis.(8) The extract was then further investigated and we found some potential contents including caffeic acid, a major representative of hydroxycinnamic acids.(8) Since caffeic acid potentially inhibited osteoclastogenesis, we performed a study using a synthetic high purified caffeic acid.(9) Then we found the synthetic one could inhibit osteoclastogenesis as well. The caffeic acid osteoclastogenesis inhibition studies were performed in RANKL-TNF $\alpha$ -macrophage colony stimulating factor (M-CSF)-induced bone marrow-derived monocyte/macrophage precursor cells (BMMs) as well as RANKL-TNF $\alpha$ -induced RAW264 cells D-Clone (RAW-D) cells.(9) Later we found that caffeic acid inhibited osteoclastogenesis of RAW-D cells by suppressing NF $\kappa$ B activity.(9) Since TRAF6 plays important role in osteoclastogenesis, we further investigated the inhibition effect of caffeic acid on osteoclastogenesis pathway of TRAF6-transfected RAW-D cells.

## Methods

### Cell Cultures


RAW-D cells, a high RANKL responsive osteoclast precursor cell line clone (10), were cultured in  $\alpha$ -MEM (GIBCO-BRL, Grand Island, NY) with 10% FBS (Biosource, Camarillo, CA) at 37 °C in a humidified incubator with 5% CO<sub>2</sub>.

### Immunoblotting

Cells were harvested and incubated with lysis buffer containing 20 mM Tris buffer (pH 7.4), 5 mM EDTA, 1% Triton-X, 50 mM sodium fluoride, 2 mM sodium orthovanadate, 10 mM sodium pyrophosphate, 1mM p-amidinophenyl methanesulfonyl fluoride hydrochloride and protease inhibitor cocktail (Sigma-Aldrich, St.Louis, MO). Samples were separated by SDS-polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane (Biorad, Richmond, CA). After blocking with 5% skim milk in phosphate buffer saline (PBS) (pH 7.4), the sheets were probed with the following 1:1000 diluted antibodies: mouse monoclonal anti-TRAF6 (D-10, Santa Cruz Biotechnology, Santa Cruz, CA) and rabbit polyclonal anti-Actin (H-300, Santa Cruz Biotechnology). The secondary antibody was horseradish peroxidase-conjugated 1:2000 diluted sheep anti-mouse (Amersham, Buckinghamshire, UK) or 1:1000 diluted donkey anti-rabbit (Amersham) IgG antibody. The bound antibodies were visualized using the ECL system (Amersham).

### Tartrate Resistant Acid Phosphatase (TRAP) Staining

TRAP staining using Leukocyte Acid Phosphatase Kit (Sigma-Aldrich) was performed according to kit protocol. TRAP<sup>+</sup> polynucleated cells (PNCs) (>3 nuclei) were counted.



**Figure 1. TRAF6 expression of RANKL-TNF $\alpha$ -caffeic acid-treated TRAF6-transfected RAW-D cells.** One hundred thousand RAW-D cells were transfected with/without TRAF6 or pcDNA3 vector only. After 30 hours, transfected cells were treated with 20 ng/ml RANKL, 1 ng/ml TNF $\alpha$  and 0-10  $\mu$ g/ml caffeic acid as indicated in the panel for 12 hours. Cells were lysed and subjected to immunoblotting assay using anti-TRAF6 and anti-Actin antibodies. Data represent a typical result from 3 independent experiments. V: pcDNA3 (empty vector of TRAF6).

### Cell Transfection and NF $\kappa$ B Activity Luciferase Assay

RAW-D cells were transfected with pNF $\kappa$ B-TA-Luc (Clontech, Palo Alto, CA) and/or pcDNA3 or pcDNA vector only in a serum free  $\alpha$ -MEM using FuGene6 (Roche Molecular Biochemicals, Mannheim, Germany). For internal control, pRLTK (an internal control plasmid containing the herpes simplex thymidine kinase promoter linked to a constitutively active Renilla luciferase reporter gene) was also transfected. After transfection, cells were treated with 20 ng/ml RANKL, 1 ng/ml TNF $\alpha$  and 0-10  $\mu$ g/ml caffeic acid for 12 hours. Luciferase assay was performed using the Dual Luciferase Reporter Assay System (Promega, Madison, WI). Briefly, after rinse with PBS, 1X passive lysis buffer (PLB) was added to cells. Cells were rocked for 15 minutes at room temperature. Then 20  $\mu$ l of each cell lysate was immediately applied to 96-well plate for measuring NF $\kappa$ B luciferase activity using TR717 Microplate Luminometer (Tropix, Bedford, MA).

### Statistical Analysis

Analyses were performed using IBM SPSS for Windows version 19.0 (IBM Corp., Armonk, NY). Student's t-test was used to determine the statistical differences between the means of experiments. A probability value <0.05 was considered to be statistically significant.

## Results


### Caffeic acid did not influence TRAF6 expression of TRAF6-transfected RAW-D cells.

As shown in Figure 1, successful RAW-D cells transfection was achieved. Elevated TRAF6 expression of TRAF6-transfected RAW-D cells was observed (lane 3). Meanwhile, TRAF6 expression of pcDNA3 vector control-transfected RAW-D cells (lane 2) was similar to endogenous TRAF6 expression of wild type RAW-D cells (lane 1). Under induction of 20 ng/ml RANKL, 1 ng/ml TNF $\alpha$  and 0-10  $\mu$ g/ml caffeic acid, expressions of TRAF6 did not show distinct differences.

### Caffeic acid inhibited RANKL-TNF $\alpha$ -induced NF $\kappa$ B luciferase activity in TRAF6-transfected RAW-D cells.

By induction of 20 ng/ml RANKL and 1 ng/ml TNF $\alpha$  for 12 hours, NF $\kappa$ B activity of wild type RAW-D cells was elevated (Figure 2). The NF $\kappa$ B activity elevation was further escalated in TRAF6-transfected RAW-D cells under the same induction of RANKL and TNF $\alpha$ . Caffeic acid, which was shown previously to inhibit osteoclastogenesis(8), was added in this assay. The results showed that caffeic

acid diminished NFκB activity of RANKL-TNFα-induced TRAF6-transfected RAW-D cells in a concentration dependent manner. Significant NFκB activity inhibitions



**Figure 2. NFκB luciferase activity of RANKL-TNFα-caffeic acid-treated TRAF6-transfected RAW-D cells.** One hundred thousand RAW-D cells were transfected with pNFκB-TA-Luc and/or pcDNA-TRAF6 or pcDNA3 vector only. After 30 hours, transfected cells were treated with 20 ng/ml RANKL, 1 ng/ml TNFα and 0-10 μg/ml caffeic acid as indicated in the panel for 12 hours. Cells were lysed and subjected to NFκB Luciferase assay. Data were analyzed by Student’s t-test, \*p<0.05, \*\*p<0.01. Data represent a typical result from 3 independent experiments. V: pcDNA3 (empty vector of TRAF6).



were seen under treatment of 1 and 10 μg/ml caffeic acid with p<0.05 and p<0.01, respectively.

**Caffeic acid inhibited RANKL-TNFα-induced osteoclastogenesis in TRAF6-transfected RAW-D cells.**

Under induction of 20 ng/ml RANKL and 1 ng/ml TNFα, wild type RAW-D cells were successfully differentiated into TRAP+ osteoclast-like PNCs (Figure 3A-C). Meanwhile, under the same induction, significantly increased PNCs number was observed in TRAF6-transfected RAW-D cells (Figure 3D). However, by adding 10 μg/ml caffeic acid in RANKL-TNFα-induced TRAF6-transfected RAW-D cells, TRAP+ PNCs number was significantly suppressed.

**Discussion**

In this study, we demonstrated that TRAF6 introduction could escalate RANKL-TNFα-induced NFκB activation. As being reported, TRAF6 plays essential roles in both the differentiation and maturation of osteoclast. Osteoclast differentiation from osteoclast progenitor cells into mature



**Figure 3. Expression and number of TRAP+ PNCs of RANKL-TNFα-caffeic acid-treated TRAF6-transfected RAW-D cells.** Six thousand RAW-D cells were transfected with pcDNA3 vector (A) or pcDNA3-TRAF6 (B&C). After 30 hours, transfected cells were treated with 20 ng/ml RANKL, 1 ng/ml TNFα and 0 (B) or 10 (C) μg/ml caffeic acid as indicated in the panel for 3 days. Cells were fixed and stained with TRAP. D: TRAP+ PNCs were counted and analyzed by Student’s t-test, \*p<0.05. Data represent a typical result from 3 independent experiments. Bar: 100 μm. V: pcDNA3 (empty vector of TRAF6).

multinucleated osteoclasts was disrupted in TRAF6-deficient mice(5) as well as in Si-RNA TRAF6-transfected human macrophages and Si-RNA TRAF6-transfected RAW 264.7 cells(11). By addition of caffeic acid in the model, we observed that NF $\kappa$ B activity was then diminished. This suggests that inhibition of caffeic acid could affect TRAF6-dependent osteoclastogenesis pathway.

Inhibition of TRAF6 could be achieved by ubiquitination, for example carboxyl terminus of heat shock protein (Hsp)70-interacting protein (CHIP) was reported to promote TRAF6 ubiquitination and proteasome degradation.(12) This CHIP-mediated TRAF6 degradation plays an important role in osteoclastogenesis inhibition. However, in our current study we found that TRAF6 expression was not disrupted by caffeic acid. This suggests that caffeic acid could possibly play a role in TRAF6 and RANK/TNF Receptor 1 interactions(13), or in TRAF6 downstream signaling pathway. TRAF6 downstream signaling includes activation of inhibitor of  $\kappa$ B (I $\kappa$ B) kinase (IKK), mitogen-activated protein kinase (MAPK) kinase kinase (MKK) and sarcoma viral oncogene homolog c-src-dependent phosphatidylinositol 3-kinase (PI3K).(1,4,11,14-18). Activation of IKK leads to NF $\kappa$ B activation, while MKK activates subsequent stimulation of p38 MAPK, extracellular signal-regulated kinase (Erk), and c-Jun N-terminal kinase (JNK) leading to nuclear translocation of the transcription factors: c-Fos, c-Jun, and nuclear factor of activated T-cells c1 (NFATc1).(1,4,11,14,16) In the PI3K pathway, resulted phosphatidylinositol-3,4,5-triphosphate (PIP3), activates Akt, which then phosphorylates and inactivates the pro-apoptotic protein Bad.(1,15-18)

Under induction of RANKL and TNF $\alpha$ , differentiation of RAW-D cells into TRAP+ osteoclast-like PNCs was well reported.(8,19,20) A significant increase of TRAP+ osteoclast-like PNCs was observed in RANKL-TNF $\alpha$ -induced TRAF6-transfected RAW-D cells. These results are in accordance with previous report showing that increased numbers of TRAF6-binding sites in the cytoplasmic tails leading to a dose-dependent increase and more pronounced osteoclastogenesis (21). When caffeic acid was added in the model of RANKL-TNF $\alpha$ -induced TRAF6-transfected RAW-D cells, significant lower number of osteoclast-like PNCs was observed.

## Conclusion

Taken together, our results showed that caffeic acid inhibited RANKL-TNF $\alpha$ -TRAF6-induced osteoclastogenesis pathway. However, further investigation should be pursued to disclose further the inhibition mechanism of caffeic acid.

## References

1. Yavropoulou MP, Yovos JG. Osteoclastogenesis - Current knowledge and future perspectives. *J Musculoskelet Neuronal Interact.* 2008; 8: 204-16.
2. Miyamoto K, Yoshida S, Kawasumi M, Hashimoto K, Kimura T, Sato Y, et al. Osteoclasts are dispensable for hematopoietic stem cell maintenance and mobilization. *J Exp Med.* 2011; 208:2175-81.
3. Miyamoto T, Suda T. Differentiation and function of osteoclasts. *Keio J Med.* 2003; 52: 1-7.
4. Takayanagi H, Kim S, Taniguchi T. Signaling crosstalk between RANKL and interferons in osteoclast differentiation. *Arthritis Res* 2002, 4(suppl 3): S227-32.
5. Kaji K, Katogi R, Azuma Y, Naito A, Inoue JI, Kudo A. Tumor necrosis factor alpha-induced osteoclastogenesis requires tumor necrosis factor receptor-associated factor 6. *J Bone Miner Res.* 2001;16: 1593-9.
6. Kobayashi N, Kadono Y, Naito A, Matsumoto K, Yamamoto T, Tanaka S, et al. Segregation of TRAF6-mediated signaling pathways clarifies its role in osteoclastogenesis. *EMBO J.* 2001; 20: 1271-80.
7. Galibert L, Tometsko ME, Anderson DM, Cosman D, Dougall WC. The involvement of multiple tumor necrosis factor receptor (TNFR)-associated factors in the signaling mechanisms of receptor activator of NF-kappaB, a member of the TNFR superfamily. *J Biol Chem.* 1998; 273: 34120-7.
8. Tang QY1, Kukita T, Ushijima Y, Kukita A, Nagata K, Sandra F, et al. Regulation of osteoclastogenesis by Simon extracts composed of caffeic acid and related compounds: successful suppression of bone destruction accompanied with adjuvant-induced arthritis in rats. *Histochem Cell Biol.* 2006; 125: 215-25.
9. Sandra F, Kukita T, Tang QY, Iijima T. Caffeic Acid Inhibits NF $\kappa$ B Activation of Osteoclastogenesis Signaling Pathway. *Indones Biomed J.* 2011; 3: 216-22.
10. Watanabe T, Kukita T, Kukita A, Wada N, Toh K, Nagata K, et al. Direct stimulation of osteoclastogenesis by MIP-1 $\alpha$ : evidence obtained from studies using RAW264 cell clone highly responsive to RANKL. *J Endocrinol.* 2004; 180: 193-201.
11. Yen ML, Hsu PN, Liao HJ, Lee BH, Tsai HF. TRAF-6 dependent signaling pathway is essential for TNF-related apoptosis-inducing ligand (TRAIL) induces osteoclast differentiation. *PLoS One.* 2012; 7: e38048.
12. Li S1, Shu B, Zhang Y, Li J, Guo J, Wang Y, et al. CHIP regulates osteoclast formation through promoting TRAF6 protein degradation. *Arthritis Rheumatol.* 2014; doi: 10.1002/art.38521.
13. Armstrong AP, Tometsko ME, Glaccum M, Sutherland CL, Cosman D, Dougall WC. A RANK/TRAF6-dependent signal transduction pathway is essential for osteoclast cytoskeletal organization and resorptive function. *J Biol Chem.* 2002; 277: 44347-56.
14. Lamothe B, Webster WK, Gopinathan A, Besse A, Campos AD, Darnay BG.
15. Son A, Kim MS, Jo H, Byun HM, Shin DM. Effects of Inositol 1,4,5-triphosphate on Osteoclast Differentiation in RANKL-induced Osteoclastogenesis. *Korean J Physiol Pharmacol.* 2012; 16: 31-6.
16. Ha H, Kwak HB, Lee SK, Na DS, Rudd CE, Lee ZH, et al. Membrane rafts play a crucial role in receptor activator of nuclear factor kappaB signaling and osteoclast function. *J Biol Chem.* 2003; 278: 18573-80.
17. Liu H, Tamashiro S, Baritaki S, Penichet M, Yu Y, Chen H, et al. TRAF6 activation in multiple myeloma: a potential therapeutic target. *Clin Lymphoma Myeloma Leuk.* 2012; 12: 155-63.
18. Tu Q, Zhang J, Dong LQ, Saunders E, Luo E, Tang J, et al. Adiponectin

- inhibits osteoclastogenesis and bone resorption via APPL1-mediated suppression of Akt1. *J Biol Chem.* 2011; 286: 12542-53.
19. Tang QY, Kukita T, Ushijima Y, Kukita A, Nagata K, Sandra F, et al. Regulation of osteoclastogenesis by Simon extracts composed of caffeic acid and related compounds: successful suppression of bone destruction accompanied with adjuvant-induced arthritis in rats. *Histochem Cell Biol.* 2006; 125: 215-25.
20. Kukita T, Wada N, Kukita A, Kakimoto T, Sandra F, Toh K, et al. RANKL-induced DC-STAMP is a key transmembrane molecule in osteoclastogenesis. *J Exp Med.* 2004; 200: 941-6.
21. Kadono Y, Okada F, Perchonock C, Jang HD, Lee SY, Kim N, et al. Strength of TRAF6 signalling determines osteoclastogenesis. *EMBO Rep.* 2005; 6(2): 171-6.