



Calcium-Mediated Abiotic Stress Signaling in Roots

Katie A. Wilkins, Elsa Matthus, Stéphanie M. Swarbreck and Julia M. Davies*

Department of Plant Sciences, University of Cambridge, Cambridge, UK

Roots are subjected to a range of abiotic stresses as they forage for water and nutrients. Cytosolic free calcium is a common second messenger in the signaling of abiotic stress. In addition, roots take up calcium both as a nutrient and to stimulate exocytosis in growth. For calcium to fulfill its multiple roles must require strict spatio-temporal regulation of its uptake and efflux across the plasma membrane, its buffering in the cytosol and its sequestration or release from internal stores. This prompts the question of how specificity of signaling output can be achieved against the background of calcium's other uses. Threats to agriculture such as salinity, water availability and hypoxia are signaled through calcium. Nutrient deficiency is also emerging as a stress that is signaled through cytosolic free calcium, with progress in potassium, nitrate and boron deficiency signaling now being made. Heavy metals have the capacity to trigger or modulate root calcium signaling depending on their dose and their capacity to catalyze production of hydroxyl radicals. Mechanical stress and cold stress can both trigger an increase in root cytosolic free calcium, with the possibility of membrane deformation playing a part in initiating the calcium signal. This review addresses progress in identifying the calcium transporting proteins (particularly channels such as annexins and cyclic nucleotide-gated channels) that effect stress-induced calcium increases in roots and explores links to reactive oxygen species, lipid signaling, and the unfolded protein response.

OPEN ACCESS

Edited by:

Janin Riedelsberger,
University of Talca, Chile

Reviewed by:

Girdhar Kumar Pandey,
University of Delhi, India
Joachim Krebs,
Max Planck Institute for Biophysical
Chemistry, Germany

*Correspondence:

Julia M. Davies
jmd32@cam.ac.uk

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 24 May 2016

Accepted: 12 August 2016

Published: 29 August 2016

Citation:

Wilkins KA, Matthus E,
Swarbreck SM and Davies JM (2016)
Calcium-Mediated Abiotic Stress
Signaling in Roots.
Front. Plant Sci. 7:1296.
doi: 10.3389/fpls.2016.01296

Keywords: abiotic stress, calcium, heavy metal, hypoxia, nutrition, salinity, signaling


INTRODUCTION

Plant roots are exposed to a variety of abiotic stresses as they navigate the soil, foraging for nutrients and water. Cytosolic free calcium ($[Ca^{2+}]_{cyt}$) is central to the response to these stresses, acting as a second messenger but also driving exocytosis (Carroll et al., 1998). Specificity of $[Ca^{2+}]_{cyt}$ signaling is determined by the amplitude and duration (and possible oscillation) of the $[Ca^{2+}]_{cyt}$ increase, often referred to as the “signature” (McAinsh and Pittman, 2009), that is elicited by the stimulus. This signature would be driven by the opening of plasma membrane (PM) and endomembrane Ca^{2+} -permeable channels and terminated by the activity of Ca^{2+} efflux transporters in those membranes, plus Ca^{2+} -binding proteins, to restore the resting $[Ca^{2+}]_{cyt}$ of 100–200 nM. Use of organelle-targeted Ca^{2+} reporting proteins has shown that the Ca^{2+} content of the endoplasmic reticulum (ER) and Golgi increases after stress-induced transient increases in $[Ca^{2+}]_{cyt}$, strongly suggesting that Ca^{2+} is sequestered there to terminate the $[Ca^{2+}]_{cyt}$ signal (Ordenes et al., 2012; Bonza et al., 2013). Transport of Ca^{2+} into organelles is catalyzed by Ca^{2+} -ATPases. There are two distinct families: The Auto-inhibited C Ca^{2+} -ATPases, ACA (that also

operate at the PM) and the ER Ca^{2+} -ATPases, ECA; reviewed by Bonza et al., 2016). The lower affinity CAX (Cation/ H^+ Exchangers) appear to be restricted to endomembranes but also facilitate Ca^{2+} sequestration (Connorton et al., 2012). Changes in organelle free Ca^{2+} in roots could also play a part in signaling, most notably in the formation of symbioses and cell death (Stael et al., 2012; Zhao et al., 2013; Wagner et al., 2015). Decoding the $[\text{Ca}^{2+}]_{\text{cyt}}$ signature will be effected by specific Ca^{2+} -binding proteins. Calmodulins (CaMs) and Calmodulin-like proteins (CMLs) are encoded by multi-gene families in plants. They lack kinase domains, suggesting these proteins must target others with enzymatic activity. CaMs modulate transcription by binding to Calmodulin-binding Transcription Activators (CAMTAs) (Viridi et al., 2015). Other multi-gene families are also evident for Ca^{2+} -Dependent Protein Kinases (CPKs) and Calcineurin-B Like proteins (CBLs). The latter target CBL-Interacting Protein Kinases (CIPKs) to effect cellular responses (Thoday-Kennedy et al., 2015). Changes in $[\text{Ca}^{2+}]_{\text{cyt}}$ also have the potential to activate lipid signaling pathways. A somewhat forgotten aspect of Ca^{2+} signaling is the Ca^{2+} activation of members of the Phospholipase C and Phospholipase D families (Qin et al., 1997; Hunt et al., 2004; Dressler et al., 2014; Ruelland et al., 2015; Hou et al., 2016). Phospholipase C catalyses production of diacylglycerol and inositol trisphosphate (InsP_3) while Phospholipase D catalyses production of phosphatidic acid, thus $[\text{Ca}^{2+}]_{\text{cyt}}$ would have the capacity to trigger distinct lipid signals depending on the location and Ca^{2+} -sensitivity of the phospholipases. Targets of lipid signals have been reviewed recently by Hou et al. (2016).

The vast majority of $[\text{Ca}^{2+}]_{\text{cyt}}$ measurements are from *Arabidopsis thaliana* seedlings and guard cells, achieved using the luminescent Ca^{2+} -interacting aequorin protein. Far fewer studies have focused specifically on roots or utilized the greater sensitivity and spatial resolution of ratiometric fluorescent dyes. The genetically encoded YC3.6 Ca^{2+} reporter is now being used for both *Arabidopsis* and rice roots (Behera et al., 2015), holding much promise for the future. It is now clear that an identical stimulus can elicit markedly different root $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures depending on genus. So far, rice root $[\text{Ca}^{2+}]_{\text{cyt}}$ signals have been found to be lower in amplitude but of longer duration than those of *Arabidopsis* (Behera et al., 2015).

Electrophysiological studies of root cell plasma membrane (PM) have advanced our understanding of the Ca^{2+} influx routes that could generate $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures. There is a central role for PM voltage in $[\text{Ca}^{2+}]_{\text{cyt}}$ signaling, as individual stresses can hyperpolarize (render it more negative) or depolarize (render it less negative). Manipulating PM voltage elicits distinct $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures and resultant transcriptional responses (Whalley et al., 2011; Whalley and Knight, 2013). Studies on root epidermal and root hair PM have shown that this membrane harbors channels that are activated by hyperpolarized voltage (Hyperpolarization-Activated Ca^{2+} Channels (HACCs); Véry and Davies, 2000; Demidchik et al., 2002, 2009; Ma et al., 2012), Depolarization-Activated Ca^{2+} Channels (DACCS); Demidchik et al., 2002; Miedema et al., 2008) and Voltage-Independent Ca^{2+} Channels (VICCs) (Demidchik et al., 2002). Thus changes in voltage would activate specific suites of channels to generate a signature. An



additional tier of regulation of the PM Ca^{2+} influx routes is afforded by reactive oxygen species (ROS) that are produced during development and stress responses (Figure 1). This regulation depends on the specific ROS, its position, the cell type and the cell's developmental state. In *Arabidopsis* roots, sensitivity of PM Ca^{2+} channel activation by extracellular H_2O_2 decreases as epidermal cells mature but is still greater than that of the cortex (Demidchik et al., 2007). A similar picture emerges for extracellular hydroxyl radicals, which elicit greater PM Ca^{2+} influx currents in the epidermis and root hairs than the pericycle (Demidchik et al., 2003; Foreman et al., 2003). In epidermal PM of the elongation zone, extracellular hydroxyl radicals elicit different Ca^{2+} channel activity to extracellular H_2O_2 (Demidchik et al., 2003, 2007). Thus, ROS will play a significant part of generating cell-specific $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures in response to stress.


Stress-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation in roots remains poorly understood in terms of the genes encoding the PM or endomembrane Ca^{2+} channels involved. Plants have multi-gene families of Glutamate Receptor-Like channels (GLR; activated by a range of extracellular nitrogenous ligands) and Cyclic Nucleotide-Gated channels (CNGC; activated by intracellular cyclic nucleotides), with each gene encoding a potential subunit of a potentially tetrameric channel. Some members have been characterized as having Ca^{2+} channel forming ability (reviewed by Swarbreck et al., 2013 and Weiland et al., 2016). Membrane residency has yet to be determined for all proteins and while the majority tested are in the PM, in *Arabidopsis* GLR3.5 has been localized to both mitochondria and chloroplast, depending on its splicing variant (Teardo et al., 2015), CNGC19 to the vacuole (Yuen and Christopher, 2013) and CNGC20 potentially to both PM and vacuole (Fischer et al., 2013; Yuen and Christopher,

2013). Mechanosensitive Ca^{2+} channels of the PM have also been identified (Nakagawa et al., 2007; Hou et al., 2014; Yuan et al., 2014; Kamano et al., 2015), as has a vacuolar Ca^{2+} efflux channel TPC1 (Two Pore Channel1; Peiter et al., 2005). Root cells will express specific complements of these genes and their transcription can change under abiotic stress (Dinneny et al., 2008; Roy et al., 2008), with the implication that stress resets the $[\text{Ca}^{2+}]_{\text{cyt}}$ signaling system.

The threat of abiotic stress is global. Drought threatens plant productivity across continents, with water shortage not only imposing an osmotic challenge but also leading to soil hardness that roots must overcome. Changing weather patterns are bringing greater rainfalls to some areas (particularly Northern Europe) thus leading to the threat of hypoxic challenge from waterlogged soil (Shabala et al., 2014). Salinity stress arising from sodic soils is made worse by irrigation and counteracting nutritional deprivation by fertilizer application comes with an increasing economic and environmental cost. In this review, the effects of salinity, water availability (including soil hardness), nutritional deprivation, heavy metals and cold on root $[\text{Ca}^{2+}]_{\text{cyt}}$ will be addressed. The candidate channels for elevating $[\text{Ca}^{2+}]_{\text{cyt}}$ in roots will be introduced and the downstream consequences of the signal will be reviewed.

SALINITY STRESS FROM CHANNEL TO TRANSCRIPTION

The transporters for Na^{+} influx into the root are not fully known but include the PM cyclic nucleotide-gated channels CNGC3 (Gobert et al., 2006) and CNGC10 (Guo et al., 2008; Jin et al., 2015) in *Arabidopsis*. Na^{+} ingress is opposed by the Annexin1 protein and the AGB1 heterotrimeric G protein subunit in *Arabidopsis* roots (Laohavisit et al., 2013; Yu and Assmann, 2015). The mechanisms for sensing the increase in cytosolic Na^{+} remain obscure (Maathuis, 2014; Shabala et al., 2015) however what is clear is that Na^{+} entry depolarizes the root epidermal PM voltage (Maathuis, 2014). This is significant in that it implicates depolarization-activated and voltage-independent PM Ca^{2+} -permeable channels in generating the transient $[\text{Ca}^{2+}]_{\text{cyt}}$ increases observed in roots in response to NaCl (Figure 2). Both channels types are present in *Arabidopsis* root epidermal PM (Demidchik et al., 2002). However, involvement of hyperpolarization-activated PM Ca^{2+} channels should not be dismissed because increasing $[\text{Ca}^{2+}]_{\text{cyt}}$ shifts their activation voltage to more depolarized values (Véry and Davies, 2000; Demidchik et al., 2002) and they are implicated in the *Arabidopsis* root response to NaCl (Ma et al., 2012; Laohavisit et al., 2013).



Critically, hyperosmotic stress hyperpolarizes root epidermal PM voltage (Maathuis, 2014) and this would potentially be a basis for generating a component of the $[Ca^{2+}]_{cyt}$ signal specific to the hyperosmotic component of NaCl stress.

Ca²⁺ Influx across the PM

Application of NaCl can cause a heterogeneous increase in *Arabidopsis* and rice root $[Ca^{2+}]_{cyt}$ that depends on cell type, external $[Ca^{2+}]$, and the bathing medium's effect on PM voltage (Kiegle et al., 2000; Tracy et al., 2008; Laohavisit et al., 2013; Choi et al., 2014; Zhang Y. et al., 2015). Pericycle cells of *Arabidopsis* have a lower amplitude of $[Ca^{2+}]_{cyt}$ increase and a more variable recovery phase than the surrounding tissues (Kiegle et al., 2000), suggesting a cell-specific transporter complement. Block of NaCl-induced $[Ca^{2+}]_{cyt}$ increase by lanthanides implicates PM Ca²⁺ influx channels in the response of both *Arabidopsis* and rice roots (Tracy et al., 2008; Zhang Y. et al., 2015). The anti-apoptotic protein Bcl-2 mimics lanthanides in that its overexpression impairs the NaCl-induced $[Ca^{2+}]_{cyt}$ increase in rice roots (Kim et al., 2014). Clearly, the $[Ca^{2+}]_{cyt}$ increase could lead to cell death if it were great enough and Bcl-2 may interact directly with PM channels. The genetic identities of the PM Ca²⁺ influx channels initiating the NaCl-induced $[Ca^{2+}]_{cyt}$ increase remain unknown. It has been suggested that they may be Ca²⁺-permeable CNGCs (Shabala et al., 2015) as salinity stress increased cytosolic cGMP in *Arabidopsis* seedlings within seconds (Donaldson et al., 2004). The source of cGMP may prove to be critical as only the $[Ca^{2+}]_{cyt}$ response to low NaCl (50 mM) was sensitive to inhibition of soluble guanylyl cyclase activity and the signal evoked by an equivalent osmotic stress was insensitive (Donaldson et al., 2004). It remains feasible that neither the ionic nor osmotic components of the $[Ca^{2+}]_{cyt}$ increase in response to high [NaCl] require cGMP. The osmotic component of NaCl stress could increase $[Ca^{2+}]_{cyt}$ in *Arabidopsis* roots through the PM mechanosensitive Ca²⁺-permeable channel OSCA1 (Reduced Hyperosmolality-Induced $[Ca^{2+}]_I$ increase). This channel mediates the $[Ca^{2+}]_{cyt}$ response to hyperosmotic stress (Yuan et al., 2014). Its discovery through a screen of aequorin-expressing mutants (Yuan et al., 2014) highlights the potential of this approach for identification of channels (and their families) involved in stress-induced $[Ca^{2+}]_{cyt}$ elevation.

The initial increase in $[Ca^{2+}]_{cyt}$ could be amplified by the production of ROS sourced ultimately by PM NADPH oxidases (encoded by *Respiratory Burst Oxidase Homolog* genes), with $[Ca^{2+}]_{cyt}$ activating these enzymes through their EF hands (Figure 2). In accordance with this, *Arabidopsis* root cortical cells lacking RBOHD and F have much lower PM hyperpolarization-activated Ca²⁺ activity in response to NaCl challenge than wild type (Ma et al., 2012). Mutant seedlings have an impaired $[Ca^{2+}]_{cyt}$ response. *Arabidopsis* RBOHD can be activated by CIPK26/CBL1/9 (Drerup et al., 2013) but a direct link to stress has not been shown. Extracellular hydroxyl radicals are likely to be the ROS involved in $[Ca^{2+}]_{cyt}$ elevation (Chung et al., 2008; Demidchik et al., 2010; Laohavisit et al., 2013; Richards et al., 2015). The extracellular superoxide anions produced by NADPH oxidases are readily converted to H₂O₂ and wall Fe/Cu act as Fenton catalysts to generate hydroxyl radicals (Richards

et al., 2015). Their production by *Arabidopsis* roots is significantly increased under NaCl stress (Demidchik et al., 2010) and RBOHC is implicated as a driver for production (Chung et al., 2008). Extracellular hydroxyl radicals activate a PM Ca²⁺ influx in *Arabidopsis* root epidermis (Demidchik et al., 2003; Foreman et al., 2003) that has now been shown to be mediated by Annexin1 (Laohavisit et al., 2012).

Annexins are Ca²⁺-binding proteins that can bind to or insert into membranes and are implicated in stress reactions (Laohavisit and Davies, 2011; Davies, 2014). Critically, when root epidermal protoplasts are challenged with NaCl, the resultant PM hyperpolarization-activated Ca²⁺ influx is lost in the *annexin1* loss of function mutant and the $[Ca^{2+}]_{cyt}$ signal is impaired (Laohavisit et al., 2013). Hydroxyl radicals are potent but short-lived so their effects must be close to the site of production (Richards et al., 2015). As NADPH oxidases can be held in lipid rafts it may be that hydroxyl radicals target co-resident channels. Finally, extracellular ATP levels of *Arabidopsis* roots increase in response to NaCl (Dark et al., 2011). Extracellular ATP activates root epidermal PM hyperpolarization-activated Ca²⁺ influx channels *via* RBOHC (Demidchik et al., 2009), suggesting the involvement of extracellular ROS in channel activation. Whether these channels involve Annexin1 remains to be determined but both the *annexin1* and *rbohC* loss of function mutants are impaired in $[Ca^{2+}]_{cyt}$ -dependent transcriptional responses under NaCl stress (Chung et al., 2008; Laohavisit et al., 2013).

Calcium Release from Stores

Although not demonstrated in roots, the *Arabidopsis* Actin-Related Protein2/3 (ARP2/3) acts to limit NaCl-induced $[Ca^{2+}]_{cyt}$ increase, partly by limiting Ca²⁺ release from mitochondria. In the *arp2/3* mutant, the $[Ca^{2+}]_{cyt}$ increase is greater than wild type and so is the extent of mitochondrial-driven cell death (Zhao et al., 2013). Release of vacuolar Ca²⁺ to the cytosol in *Arabidopsis* roots may be by a Na⁺/Ca²⁺ exchanger encoded by *AtNCL (Na⁺/Ca²⁺ Exchanger-Like)*; Wang et al., 2012; Li P.H. et al., 2016). This tonoplast protein is thought to sequester Na⁺ into the vacuole, coupled to the release of vacuolar Ca²⁺. The concomitant increase in $[Ca^{2+}]_{cyt}$ could provide a negative feedback mechanism to limit further transport as Ca²⁺ binding to the exchanger's EF hands has been shown to be inhibitory *in vitro* (Li P.H. et al., 2016). Pharmacological approaches have also implicated internal stores in the NaCl-induced $[Ca^{2+}]_{cyt}$ increase in roots of both *Arabidopsis* and rice. Inhibitors of store release of Ca²⁺ by cADPR (cyclic ADP ribose) and inositol trisphosphate (InsP₃) suggested involvement of the InsP₃ pathway in *Arabidopsis* roots (Tracy et al., 2008). Moreover, salt and hyperosmotic stress in *Arabidopsis* roots caused an InsP₃ accumulation that correlated well with $[Ca^{2+}]_{cyt}$ increase (DeWald et al., 2001). The rice $[Ca^{2+}]_{cyt}$ signal was also sensitive to disruption of putative InsP₃-gated store release whilst impairment by thapsigargin implicated the ER as a participating store (Zhang Y. et al., 2015). However, using ER-targeted YC3.6, Bonza et al. (2013) detected an increase in *Arabidopsis* root ER $[Ca^{2+}]$ in response to salt stress. This followed the salt-induced $[Ca^{2+}]_{cyt}$ increase and critically, a drop in ER $[Ca^{2+}]$ prior to

the $[Ca^{2+}]_{cyt}$ increase was never observed. Therefore, with the level of resolution available, it appears that in this system the ER does not contribute to the $[Ca^{2+}]_{cyt}$ signal through store release but acts to return $[Ca^{2+}]_{cyt}$ to resting levels. Studies on *Populus euphratica* cultured cells identified the vacuole as the site of $InsP_3$ and cADPR action (Zhang X. et al., 2015). Certainly, Phospholipase C isoforms (as the source of $InsP_3$) are firmly implicated in salt stress responses (reviewed by Ruelland et al., 2015).

The genetic identities of the endomembrane Ca^{2+} -permeable channels implicated by pharmacological studies remain elusive. GLRs have recently been postulated to be involved in ER Ca^{2+} release (Weiland et al., 2016). The TPC1 vacuolar channel of *Arabidopsis* would be capable of releasing Ca^{2+} to the cytosol and recent analyses of its crystal structure has thrown greater light on its regulation by voltage and EF hands (Guo et al., 2016; Kintzer and Stroud, 2016). At current resolution afforded by YC3.6, its loss does not appear to have a significant impact on the magnitude of the *Arabidopsis* root $[Ca^{2+}]_{cyt}$ increase to salt, rather it slightly delays the response (Choi et al., 2014). However, TPC1 has a significant part to play in the propagation of a $[Ca^{2+}]_{cyt}$ wave that travels from the root apex through the cortical and endodermal tissue to signal the NaCl challenge to the shoots and elicit a transcriptional response. ROS also relay a salinity stress signal (reliant on RBOHD) in *Arabidopsis* leaves (Miller et al., 2009) but whether this occurs in roots and is involved in propagating the $[Ca^{2+}]_{cyt}$ wave remains to be tested.

Decoding, Na^+ Clearance, Transcription and the Unfolded Protein Response

Salinity-induced $[Ca^{2+}]_{cyt}$ elevation in roots drives a transcriptional response (Laohavisit et al., 2013; Zhang Y. et al., 2015) and post-translational modifications. The proteins sensing the NaCl-induced $[Ca^{2+}]_{cyt}$ increase are now being elucidated. For example, the *Arabidopsis* vacuolar Two Pore K^+ channel 1 (TPK1) would bind Ca^{2+} , and open to release K^+ to the cytosol to maintain a favorable Na^+/K^+ ratio (Latz et al., 2013). This could be further enhanced by phosphorylation by CPK3, which requires micromolar $[Ca^{2+}]$ for activity. CPK3 is present at both the PM and vacuole. It does not appear to contribute to a transcriptional response under salt stress but has a discrete set of protein targets to phosphorylate (Mehlmer et al., 2010). On prolonged salt stress, CPK29 expression is induced. This protein can phosphorylate TPK1 at sub-micromolar $[Ca^{2+}]$ and is envisaged to be part of longer-term K^+ homeostasis in adapted roots (Latz et al., 2013). Also in *Arabidopsis*, CPK27 (present at the root PM) acts to promote Na^+ efflux (Zhao et al., 2015). Intriguingly, CPK7 acts to limit water transport in *Arabidopsis* roots through lowering PIP1 aquaporin abundance (Li et al., 2015) but whether this is relevant to salinity or osmotic stress is not yet known. It can be readily envisaged that calmodulins will bind Ca^{2+} and as these are negative regulators of CNGC channels (Hua et al., 2003), would act to limit further Na^+ or Ca^{2+} influx at the PM. Further, CaM activation of ACAs (Autoinhibited Ca^{2+} ATPases) would restore $[Ca^{2+}]_{cyt}$ to pre-stimulus levels (Boursiac and Harper, 2007). Expression

of ACAs varies with salt stress and, as shown by rice roots, can relate to salt tolerance (Yamada et al., 2014).

The Salt Overly Sensitive (SOS) pathway lies downstream of the root $[Ca^{2+}]_{cyt}$ increase. Delineated in *Arabidopsis* and now acknowledged as operating in crops and trees (Thoday-Kennedy et al., 2015), the SOS pathway leads to Na^+ efflux from the cytosol. Efflux across the PM is mediated by the SOS1 Na^+/H^+ antiporter. Salt stress induction of SOS1 transcription lies downstream of Annexin1 in *Arabidopsis* roots and as SOS1 is required for adaptive adventitious root formation, the *annexin1* loss of function mutant accordingly produces fewer of these than wild type (Laohavisit et al., 2013). Additionally, the stability of salt stress-induced SOS1 transcript requires RBOHC (Chung et al., 2008), further suggesting that this NADPH oxidase and Annexin1 may be in the same pathway. In *Arabidopsis* roots, increased $[Ca^{2+}]_{cyt}$ is sensed at the PM by CBL4 (SOS3) which can then react with the serine/threonine protein kinase CIPK24 (SOS2) (Liu et al., 2000). The resultant CBL/CIPK complex phosphorylates the PM Na^+/H^+ antiporter SOS1 at its auto-inhibitory C-terminus to achieve activation of Na^+ efflux (Quintero et al., 2011). Activation of SOS1 can also be achieved in *Arabidopsis* by Mitogen Activated Protein Kinase6 (MPK6) and loss of MPK6 function impairs root growth under salt stress (Yu et al., 2010). MPK6 is activated by a phosphatidic acid produced under salt stress by Phospholipase $D\alpha 1$ ($PLD\alpha 1$) (Yu et al., 2010). As this PLD isoform contains a Ca^{2+} -binding C2 domain for its activation it is feasible that salt-induced $[Ca^{2+}]_{cyt}$ increase could activate SOS1 through this lipid-mediated pathway via MPK6 (Yu et al., 2015). This may help explain the importance of activation of PLDs in salt-stressed crop roots such as barley (Meringer et al., 2016). SOS2 may also promote activity of the vacuolar V-type H^+ -ATPase to provide the driving force for Na^+ sequestration (Batelli et al., 2007). The $NHX1$ Na^+-H^+ exchanger may also be regulated by SOS2 to aid vacuolar Na^+ sequestration (Qiu et al., 2004) and SOS2 activation of the vacuolar $CAX1$ Ca^{2+}/H^+ antiporter would help terminate a $[Ca^{2+}]_{cyt}$ signature through Ca^{2+} sequestration (Cheng et al., 2004). Other CAX are involved in root tolerance of salt stress and the pathways to their induction and regulation now need to be identified (Yamada et al., 2014).

Salt exposure puts the plant's ER under stress, leading to an accumulation of unfolded or misfolded proteins that could lead to cell death (Liu et al., 2007, 2011). Such ER stress triggers upregulation of a suite of responses termed the "Unfolded Protein Response" (UPR), in which folding capacity is upregulated (including by Ca^{2+} -regulated chaperones), translation is curtailed and the ER-associated degradation pathway acts to lower the aberrant protein load (Deng et al., 2013; Ruberti et al., 2015; Hossain et al., 2016; Wan and Jiang, 2016). Expression of the ER Ca^{2+} -binding chaperones Calnexin and Calreticulin has been shown to upregulated in the *Arabidopsis* UPR (Christensen et al., 2008; Liu et al., 2011) and it would now be interesting to test whether these are involved in regulating levels of Ca^{2+} in the ER under stress. However, expression of rice's only calnexin gene is decreased under salt stress (Sarwat and Naqvi, 2013). It is not yet clear whether the salt-induced $[Ca^{2+}]_{cyt}$ signal in roots (or other parts

of the plant) helps initiate or regulate the UPR. It is feasible that the salt-induced $[Ca^{2+}]_{cyt}$ increase activates the root PM Phospholipase C2 (Hunt et al., 2004) because the *Arabidopsis* loss of function mutant is hypersensitive to tunicamycin, which can induce the UPR (Kanehara et al., 2015). With IP_3 as the product of Phospholipase C activity, this would implicate IP_3 -mediated release of Ca^{2+} from stores. Indeed, for *Arabidopsis* seedlings the application of 2-aminoethoxydiphenyl borate as an inhibitor of IP_3 -mediated Ca^{2+} release prevented salt stress-induced transcription of *BIP1/2* (encoding an ER chaperone) as a diagnostic of a UPR response (Liu et al., 2011). Intriguingly, application of La^{3+} as a blocker of PM Ca^{2+} influx prevented the upregulation of UPR gene expression which was produced by application of spermine to *Arabidopsis* seedlings (Sagor et al., 2015). Exogenous spermine (as with salt) can depolarise the root epidermal PM (Pottosin et al., 2014) and can also attenuate hydroxyl radical-induced cation fluxes at root epidermal PM (Zepeda-Jazo et al., 2011). Whether spermine and salt stress share a common pathway to the UPR response merits further investigation.

WATER AVAILABILITY IS SIGNALLED THROUGH $[Ca^{2+}]_{cyt}$

The hyperosmotic challenge in $[Ca^{2+}]_{cyt}$ determinations is acute and does not mimic the chronic, progressive drought conditions that roots may face. Nevertheless, such studies have proved fruitful. As described above, the *Arabidopsis* OSCA1 PM Ca^{2+} influx channel drives the root's initial hyperosmotic stress $[Ca^{2+}]_{cyt}$ signal (Yuan et al., 2014). In an elegant study, heterologous expression of *Arabidopsis* genes in Chinese Hamster Ovary (CHO) cells containing the Fura-2 Ca^{2+} -reporting dye lead to the identification of Calcium-permeable Stress-gated channel1 (CSC1; Hou et al., 2014), a close relative of OSCA1. CSC1 has been characterized in CHO cells as a PM Ca^{2+} -permeable channel that is activated by hyperosmotic stress. It is resident in plant PM (Hou et al., 2014) and is expressed in roots but to date an *in planta* role remains unreported. Targeting aequorin to the cytosolic face of the *Arabidopsis* vacuolar membrane has revealed the capacity of the vacuole to release Ca^{2+} in response to acute hyperosmotic stress, with pharmacological intervention suggesting an involvement of $InsP_3$ (Knight et al., 1997). As with salt stress, hyperosmotic stress-induced $[Ca^{2+}]_{cyt}$ increase could activate Phospholipases and initiate lipid signaling. Osmotic stress activates PLD in barley roots (Meringer et al., 2016). Although not tested directly, phosphatidic acid downstream of PLD could be involved in the activation of two sucrose non-fermenting-1 related protein kinase 2 proteins (SnRK2.4 and 2.10) in *Arabidopsis* roots under hyperosmotic stress. These SnRK2s are also activated under salt stress and relocate from the root epidermal cytosol to the PM; loss of function impairs root growth (McLoughlin et al., 2012). Work using Golgi-targeted aequorin in *Arabidopsis* seedlings has shown that an increase in Golgi $[Ca^{2+}]$ follows the $[Ca^{2+}]_{cyt}$ increase induced by hyperosmotic stress, suggesting that this organelle helps terminate the $[Ca^{2+}]_{cyt}$ signal (Ordenes et al., 2012).

The ABA produced under drought stress inhibits primary root growth and $[Ca^{2+}]_{cyt}$ is likely to play a role in the signaling pathway as exogenous ABA elevates root $[Ca^{2+}]_{cyt}$, which in *Arabidopsis* roots is controlled by the PM Proline-rich Extensin-like Receptor Kinase4 (PERK4) (Bai et al., 2009). PERK4's extracellular domain is wall associated and its intracellular domain has kinase activity. PM HACCs lie downstream of ABA and PERK4. The *perk4* mutant is not only impaired in ABA-induced HACC activity and $[Ca^{2+}]_{cyt}$ activation but also in ABA inhibition of primary root elongation (Bai et al., 2009). This implies that $[Ca^{2+}]_{cyt}$ elevation acts to arrest growth. RBOHD/F may lie downstream of PERK4 and upstream of HACCs in the root ABA pathway if the HACCs were ROS-activated. The *rbohdf* mutant is impaired in both ABA-induced HACC activation and $[Ca^{2+}]_{cyt}$ elevation (Jiao et al., 2013). Another possibility is that CNGCs are involved as ABA can increase cGMP levels (Isner et al., 2012). GLRs are firmly implicated in the drought response; overexpression of rice GLR1 and GLR2 enhances drought tolerance of both rice and *Arabidopsis* (Lu et al., 2014). The $[Ca^{2+}]_{cyt}$ increase is likely to lead to a transcriptional response as Ca^{2+} -sensing proteins are activated. ABA increases abundance and activity of *Arabidopsis* CPK4 and CPK11 leading to phosphorylation of the ABA-responsive transcription factors ABF1 and ABF4 and induction of drought stress genes (Zhu et al., 2007). Drought leads to CAMTA1 activity and the regulation of discrete gene sets including those for drought recovery (Pandey et al., 2013). In maize roots, drought stress triggers activity of CIPK8 (which interacts with CBL1, 4, and 9) and likely leads to adaptive transcription (Tai et al., 2016).

Hypoosmotic stress also elevates $[Ca^{2+}]_{cyt}$ and is relevant to waterlogged soils. In *Arabidopsis*, root PM harbors two mechanosensitive Ca^{2+} -permeable channels, MCA1 and MCA2 (Mid1-Complementing Activity; Kamano et al., 2015). MCA1 responds to hypoosmotic stress to elevate $[Ca^{2+}]_{cyt}$ (Nakagawa et al., 2007). Rice only harbors an MCA1 in the PM. It is present in the root and mediates hypoosmotic shock-induced $[Ca^{2+}]_{cyt}$ expression in cultured cells, probably lying upstream of NADPH oxidase activity (Kurusu et al., 2012). Abundant water not only exposes roots to potential hypoosmotic stress but also risks limiting their oxygen supply, the consequences of which are reviewed in the following section.

MECHANISTIC BASIS OF $[Ca^{2+}]_{cyt}$ RESPONSE TO O_2 DEFICIENCY REMAINS POORLY UNDERSTOOD

Oxygen deficiency (hypoxia) or absence (anoxia) causes transient increases in root $[Ca^{2+}]_{cyt}$ but the signature is organ- and species-dependent (reviewed by Shabala et al., 2014). For example, when challenged by anoxia, root protoplasts from hypoxia-tolerant rice display a greater $[Ca^{2+}]_{cyt}$ signature than hypoxia-intolerant wheat root protoplasts (Yemelyanov et al., 2011). The location of the Ca^{2+} influx also varies; use of pharmacological blockers showed that the rice signature was generated by both PM influx and store release whilst wheat

appeared to rely solely on stores (Yemelyanov et al., 2011). The types of Ca^{2+} channels mediating the $[\text{Ca}^{2+}]_{\text{cyt}}$ increase have yet to be identified in any species. However, those activated by PM voltage depolarization are implicated. As O_2 deficiency lessens ATP production, activity of the PM H^+ -ATPase can be compromised thus resulting in a less negative (depolarized) PM voltage as H^+ efflux is curtailed. The extent and duration of membrane depolarization varies with sensitivity to O_2 deprivation and cell type. Values of -70 to -80 mV have been reported for O_2 -deprived barley root cells (Zeng et al., 2014) and theoretically these would be sufficient to activate PM depolarization-activated Ca^{2+} influx channels (Miedema et al., 2008) to contribute to an hypoxia/anoxia $[\text{Ca}^{2+}]_{\text{cyt}}$ signature. Downstream of the $[\text{Ca}^{2+}]_{\text{cyt}}$ signature, it is likely that CaM and ROPs (Rho GTPase) are activated (reviewed by Shabala et al., 2014). In *Arabidopsis* roots, hypoxia causes rapid upregulation of *CML38* expression and this protein appears to require Ca^{2+} to associate with the cytosolic stress granules that form and store messenger RNA ribonucleoproteins (Lokdarshi et al., 2016).

RBOH activity is firmly implicated in the response to low O_2 . In *Arabidopsis*, *RBOHD* expression is induced by hypoxia and is required for transcription of hypoxia-induced genes (Yang and Hong, 2015). Hypoxia also induces ethylene production and in wheat roots this causes RBOH induction (Yamauchi et al., 2014). A further level of regulation has been found in *Arabidopsis*; the Hypoxia Responsive Universal Stress Protein 1 (HRU1) interacts with GTP-bound ROP2 and RBOHD (Gonzali et al., 2015). HRU1 is one of 44 putative universal stress proteins in *Arabidopsis*. It exists as a cytosolic dimer but anoxia promotes monomer formation and increased association with the PM. There it is thought to be part of a mobile complex with ROP2 and ATRBOHD leading to activation of that NADPH oxidase (Gonzali et al., 2015). Although RBOH activity has been linked to PM Ca^{2+} channel activation in several abiotic stress scenarios, hypoxia and anoxia have yet to be tested.

Ethylene has been shown to activate PM Ca^{2+} -permeable channels (with a weak voltage dependence) in tobacco suspension cells (Zhao et al., 2007) and it remains a possibility that these may play a part in the root hypoxia $[\text{Ca}^{2+}]_{\text{cyt}}$ signal with RBOH as an intermediary. Oxygen deprivation (and also sulfur and phosphate, Pi, deprivation) triggers programmed cell death (PCD) in mid-cortical cells for aerenchyma formation (Fagerstedt, 2010). This PCD is stimulated by ethylene and ROS and involves Ca^{2+} (Xu et al., 2013; Petrov et al., 2015). In wheat roots, anoxia causes mitochondria to release their Ca^{2+} and high $[\text{Ca}^{2+}]_{\text{cyt}}$ causes cytochrome c release (Virolainen et al., 2002). The abnormal mitochondrial ultrastructure in *Arabidopsis* caused by hypoxia is partially phenocopied by loss of GLR3.5 from the inner mitochondrial membrane, suggesting that this channel must be deactivated during PCD (Teardo et al., 2015). Further channels must now be identified to understand the hypoxic/anoxic response. Recent work on *Arabidopsis* roots has shown distinct, cell-specific levels of $[\text{Ca}^{2+}]_{\text{cyt}}$ after 24 h of hypoxia and highlighted the importance of CAX11 in controlling meristem $[\text{Ca}^{2+}]_{\text{cyt}}$ (Wang et al., 2016). Loss of CAX4 function resulted in lower tolerance of hypoxia thus further demonstrating

the importance of Ca^{2+} sequestration in this stress response (Wang et al., 2016).

CA²⁺ SIGNALING IN NUTRIENT DEPRIVATION IS AN EMERGING AREA

Investigating $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation in response to nutrient deprivation or resupply is technically challenging, particularly if using aequorin. Nevertheless it is now clear that nutrient levels can induce $[\text{Ca}^{2+}]_{\text{cyt}}$ changes and that downstream Ca^{2+} sensors regulate appropriate responses. The nutritional status of the root will have a part to play in determining the $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures, particularly of the endodermis as the extent of suberization is set by nutrition (Barberon et al., 2016) and suberin lamellae determine endodermal $[\text{Ca}^{2+}]_{\text{cyt}}$ responses (Moore et al., 2002). To date, potassium, nitrate and boron have been studied.

Potassium

Plants must maintain cytosolic K^+ at around 80 mM for optimal growth even though soil concentration may be sub-millimolar and they deploy a multigene family of K^+ transporters in homeostatic control (Shabala and Pottosin, 2014). As extracellular K^+ decreases, the root epidermal PM hyperpolarizes and $[\text{Ca}^{2+}]_{\text{cyt}}$ increases (Demidchik et al., 2002). This increase can be abolished by gadolinium, implicating PM Ca^{2+} influx. The genetic identities of the PM Ca^{2+} -permeable channels that would effect the $[\text{Ca}^{2+}]_{\text{cyt}}$ increase remain unknown. The hyperpolarised PM correlates with induction of *HAK5* expression in tomato root (Nieves-Cordones et al., 2008). *HAK5* encodes the PM High Affinity K⁺ transporter that facilitates K^+ uptake from low external concentrations that are thermodynamically unfavorable for channel-mediated influx. In *Arabidopsis* roots, low K^+ -induction of *HAK5* expression (and other K^+ -deprivation genes) has been shown to depend on RBOHC activity and ROS (Shin and Schachtman, 2004). At present it is unclear whether the hyperpolarization of the PM directly activates RBOHC (which as an electron exporter may be voltage-dependent) or whether ROS-activated PM Ca^{2+} channels are involved in the K^+ -deprivation $[\text{Ca}^{2+}]_{\text{cyt}}$ signal. *HAK5* activity in *Arabidopsis* roots is regulated by CIPK23-mediated phosphorylation, downstream of CBL1, CBL8, CBL9, and CBL10 (Ragel et al., 2015). The extent to which the transcriptional response to low K^+ availability is governed by $[\text{Ca}^{2+}]_{\text{cyt}}$ is unclear but deficiency does result in upregulation of transcripts of Ca^{2+} signaling proteins (CaM, CBL, CIPK) in *Arabidopsis* seedlings (Armengaud et al., 2004) and sugarcane roots (Zeng et al., 2015). At higher external $[\text{K}^+]$, the AKT1 channel (*Arabidopsis* K⁺ Transporter1) facilitates uptake and its activity is promoted by CIPK23-mediated phosphorylation, downstream of CBL1 and CBL9 (Li et al., 2006; Cheong et al., 2007). This regulation is recapitulated in rice roots where CIPK23/CBL1 activate AKT1 (Li et al., 2014).

Nitrate

Nitrate is the most important form of nitrogen for agriculture and deprivation triggers significant transcriptional and

developmental responses. The effect of nitrate withdrawal on $[Ca^{2+}]_{cyt}$ has yet to be reported but recently it was shown that nitrate-starved *Arabidopsis* roots responded to nitrate resupply with a rapid, monophasic transient increase in $[Ca^{2+}]_{cyt}$ that was sensitive to lanthanides and phospholipase C (PLC) inhibition (Riveras et al., 2015). Lanthanum also blocked nitrate-induced $InsP_3$ production, suggesting that Ca^{2+} influx across the PM activated a PLC. The $[Ca^{2+}]_{cyt}$ and $InsP_3$ increases were entirely dependent on the PM nitrate influx transporter NRT1.1 (Nitrate Transporter1.1; Riveras et al., 2015). By using the *nrt1.1* mutant and pharmacological blockers, nitrate-induced gene transcription was also found to lie downstream of NRT1.1, and $[Ca^{2+}]_{cyt}$ elevation from PM influx and $InsP_3$ -gated store release. Calcium is also key to the regulation of nitrate uptake capacity as CIPK23, which is activated by CBL9 and CBL1, and dephosphorylated by ABI2 (a member of the PP2C protein phosphatase family; Lérán et al., 2015), phosphorylates NRT1.1 under low nitrate condition, thus converting it from a low to high affinity transporter (Ho et al., 2009). By contrast CIPK8 positively regulates the low-affinity phase of the nitrate primary response which includes transcriptional regulation, but its regulation of primary root elongation is concentration independent in *Arabidopsis* (Hu et al., 2009). CBL7, which is upregulated under nitrate deprivation, positively regulates the nitrate-dependent induction of *NTR2.4* and *NTR2.5* gene expression (Ma Q. et al., 2015). Given the lack of a $[Ca^{2+}]_{cyt}$ reporter line available in crops up until recently for rice, little is known about nitrate deficiency-induced $[Ca^{2+}]_{cyt}$ signaling but CaM protein abundance of wheat roots declines under nitrate deficiency, suggesting a remodeling of signaling systems (Jiang et al., 2015).

Boron

Boron deficiency is widespread worldwide and particularly prevalent in China (Shorrocks, 1997). As B plays a dominant role in co-ordinating cell wall structure (Kobayashi et al., 1996), changes in cell wall stability are likely to influence the signal relayed into the cell upon B deprivation and indeed a rapid change in cell wall modulus has been observed (Goldbach et al., 2001). Early work in *Vicia faba* showed a release of membrane-bound Ca^{2+} into the apoplast (Mühling et al., 1998), raising the possibility of Ca^{2+} signaling during the early stages of B deficiency. Increased levels of both $[Ca^{2+}]_{cyt}$ and ROS have been suggested to lead to the increased root hair growth known to occur under B deprivation (González-Fontes et al., 2016). In cultured tobacco cells, transcriptional changes in response to short-term B deprivation (1 h) were abolished when withdrawing Ca^{2+} from the growth medium or upon treatment with the Ca^{2+} channel blocker lanthanum, thus implicating PM Ca^{2+} influx channels in generating a $[Ca^{2+}]_{cyt}$ signal (Koshiba et al., 2010). However, a transient $[Ca^{2+}]_{cyt}$ signal in direct response to B withdrawal has yet to be reported. Rather, work has focused on the possible remodeling of Ca^{2+} transport and signaling as a consequence of B deprivation.

Challenging cultured tobacco cells with Ca^{2+} resulted in a higher amplitude of $[Ca^{2+}]_{cyt}$ transient in B-deprived cells (1 h deprivation) than those grown under replete conditions

(Koshiba et al., 2010). This suggests that B deprivation rapidly “resets” the PM's Ca^{2+} transport systems to generate altered $[Ca^{2+}]_{cyt}$ responses. The $[Ca^{2+}]_{cyt}$ response of B-deprived cells was sensitive to lanthanum and diphenylpicrylhydrazolium, pointing to the involvement of PM Ca^{2+} channels and NADPH oxidases respectively (Koshiba et al., 2010). *Arabidopsis* roots expressing the YC3.6 $[Ca^{2+}]_{cyt}$ reporter exhibited higher levels of $[Ca^{2+}]_{cyt}$ at the apex than controls after 6 and 24 h of B deprivation (Quiles-Pando et al., 2013). This time course of B deprivation also resulted in significant upregulation of *CNGC19* (encoding a vacuolar channel), four genes of the ACA family of P_{HIB} -type Ca^{2+} -ATPases (*ACA1,10,12,13*) and *CAX3* encoding a vacuolar cation- H^+ antiporter (Quiles-Pando et al., 2013). This suite of transporters could effect Ca^{2+} efflux from the vacuole (*CNGC19*) to increase $[Ca^{2+}]_{cyt}$ with clearance to the apoplast by *ACA10-13* and sequestration to the vacuole by *CAX3*. How they are regulated remains to be determined, as does the involvement of the structurally compromised wall and the consequence of this higher level of apical $[Ca^{2+}]_{cyt}$. The area of higher $[Ca^{2+}]_{cyt}$ reported appears to correspond with the zone of inhibition of primary root elongation and the induction of cell death (Oiwa et al., 2013; Camacho-Cristobal et al., 2015). Once again, ethylene production appears to be upstream of NADPH oxidase-driven ROS production in growth arrest and death (Oiwa et al., 2013; Camacho-Cristobal et al., 2015).

Eight *CML* genes were also significantly upregulated after a day's B deprivation of *Arabidopsis* roots (*CML11,12,23,24,30.37,45.47*), as were three *CPK* genes (*CPK1,28,29*) all suggesting a distinct change in intracellular Ca^{2+} signaling (Quiles-Pando et al., 2013). This B deprivation also caused upregulation of *WRKY* transcription factors (TF) (*WRKY38,40,46*), two three *MYB* family TF (*MYB14,15,78*) and downregulation of two *BZIP* family TFs (*bZIP34,61*) (González-Fontes et al., 2016). B deprivation has also been found to promote the senescence-associated *WRKY6* TF in the root tip (Kasajima et al., 2010). It has been suggested that the *CMLs* and *CPKs* upregulated by B deprivation are part of the chain of events leading to TF activation in the nucleus (González-Fontes et al., 2016) and this now requires direct testing. CIPK8 also positively regulates nitrate induced up-regulation of *BOR1*, a gene encoding a boron transporter responsible for xylem loading (Hu et al., 2009), suggesting that root signaling results in preservation of shoot B supply.

HEAVY METAL STRESS HAS THE CAPACITY TO DISTORT Ca^{2+} SIGNALING

At the opposite end of nutritional deprivation is heavy metal stress. Industrial activity, mining and modern agricultural practices can lead to soil contamination by heavy metals (defined here as 7 g/cm³ and above). Although some of these metals (such as Zn, Cu) are required as micronutrients they can be damaging in excess whilst others (such as Cd) have no physiological role and can be deleterious even at low concentrations, often impairing mineral nutrition. The consequences of heavy metal

exposure have been reviewed recently by Singh et al. (2016) and these authors explore signaling pathways (although not explicitly addressing Ca^{2+}), intersects with hormonal responses and detoxification.

Cadmium

Cadmium is a particular threat to Ca^{2+} -based processes because of its similar size. A recent review by Chmielowska-Bak et al. (2014) summarized the effects of Cd on ROS accumulation, NO accumulation, MAP kinase activation and downstream responses in a wide range of plant systems and, importantly, did this as a function of Cd exposure. With regards to Ca^{2+} a key question is whether Cd (as a Ca^{2+} “substitute”) generates a signal in its own right or whether its impairment of Ca^{2+} homeostasis will be, in effect, the signal. Certainly Cd depolarises the rice root epidermal PM, which would impair Ca^{2+} influx and results in inhibition of root elongation (Li et al., 2012). Cd can permeate guard cell PM Ca^{2+} channels (Perfus-Barbeoch et al., 2002) and may compete with Ca^{2+} for entry into rice root hairs through PM HACC, thus limiting Ca^{2+} influx (Li et al., 2012). Competitive effects appear likely given the ability of exogenous Ca^{2+} addition to alleviate Cd inhibition of root growth of both terrestrial and aquatic plants, implying competition for uptake (Zhang et al., 2012; Rodriguez-Hernandez et al., 2015). Additionally, competitive inhibition by Cd of Ca^{2+} uptake through the wheat LCT1 transporter expressed in yeast has been observed (Clemens et al., 1998). Root hair growth in *Arabidopsis* is inhibited by Cd; it inhibits Ca^{2+} influx and so dissipates the apical $[\text{Ca}^{2+}]_{\text{cyt}}$ gradient needed for growth (Fan et al., 2011). This could reflect block of root hair Ca^{2+} influx channels or Cd outcompeting Ca^{2+} for that influx pathway. However, Yeh et al. (2007) observed an increase in rice root $[\text{Ca}^{2+}]_{\text{cyt}}$ following 15 min of Cd exposure, with a more tolerant variety sustaining a greater increase than a sensitive variety. In roots of the aquatic plant *Typha latifolia*, Cd exposure increases transcription of *TPC1* which suggests that vacuoles might release Ca^{2+} to the cytosol (Rodriguez-Hernandez et al., 2015). Inhibitor treatments indicated involvement of NADPH oxidases, hydroxyl radicals and CIPK upstream of a MAP kinase induction that linked to tolerance (Yeh et al., 2007). Cd stimulates expression and activity of NADPH oxidases in cucumber roots but it is not known if these responses are Ca^{2+} -dependent (Jakubowska et al., 2015). It is worth noting that it is not only an increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ that could stimulate NADPH oxidase activity but that also the restriction of Ca^{2+} entry to the cytosol could cause stimulation (Mortimer et al., 2008). It seems likely that in *Arabidopsis* roots it would be CAXs predominating in terminating a Cd-induced increase in $[\text{Ca}^{2+}]_{\text{cyt}}$. CAX4 mediates Cd and Ca^{2+} transport into *Arabidopsis thaliana* vacuoles and is needed for correct growth under Cd stress (Mei et al., 2009). *CAX1* expression is higher in roots of the Cd-tolerant *A. halleri* compared to the sensitive *A. thaliana* (Baliardini et al., 2015).

A key point for future studies is the intersect between Cd and hormones in relation to $[\text{Ca}^{2+}]_{\text{cyt}}$. Exogenous Ca^{2+} can ameliorate Cd's inhibition of *Arabidopsis* root growth by counteracting effects of NO on auxin homeostasis (Hu et al., 2013; Li P. et al., 2016; Yuan and Huang, 2016). Cd also interferes

with auxin homeostasis in barley roots (Zelinova et al., 2015). Auxin itself can increase *Arabidopsis* root $[\text{Ca}^{2+}]_{\text{cyt}}$ and this increase is mediated by CNGC14 at the PM, downstream of an unidentified auxin receptor (Shih et al., 2015). This begs the question of whether CNGC14 is an entry route for Cd and the pathway to disrupted auxin homeostasis. Additionally, Cd has been described as a “metallohormone” in that it triggers expression of brassinosteroid-regulated genes in *Arabidopsis* roots (Villiers et al., 2012). Brassinosteroids are themselves capable of transiently elevating $[\text{Ca}^{2+}]_{\text{cyt}}$ in *Arabidopsis* roots (through PM Ca^{2+} influx) and activate a possible DACC in wheat root PM (Straltsova et al., 2015). Again this raises the question of channel identity to help elucidate the relationship between Cd and brassinosteroid signaling and combat the effects of this potent soil contaminant.

Copper, Gadolinium and Lead

In contrast to Cd, transition heavy metals could be capable of generating ROS directly and so perturb $[\text{Ca}^{2+}]_{\text{cyt}}$ signaling. Transition metals can catalyze production of hydroxyl radicals from superoxide anion and hydrogen peroxide through the Haber-Weiss reaction, with Cu^+ and Fe^{2+} catalyzing hydroxyl radical production from hydrogen peroxide through the Fenton reaction (Richards et al., 2015). Unless levels of these catalytic metals are tightly controlled, production of hydroxyl radicals (the most potent of the ROS) could inflict significant oxidative damage. Taking Cu as the exemplar, it plays a positive role in hydroxyl radical-activated cell wall loosening for root elongation (Fry et al., 2002) and this could be coupled in *Arabidopsis* roots to hydroxyl radical-activation of PM Annexin1-mediated Ca^{2+} influx to stimulate exocytosis and growth (Foreman et al., 2003; Laohavisit et al., 2012). This model proposes regulation by extracellular hydroxyl radicals while Rodrigo-Moreno et al. (2013) have proposed that Cu^+ binding at the intracellular face of the PM in *Arabidopsis* root tips catalyzes hydroxyl radical production to regulate ion flux. This would allow coupling of Cu transport into the root to regulation of PM Ca^{2+} influx channels. Certainly, low levels of Cu can promote elongation of *Arabidopsis* primary root and this effect is diminished by blocking PM Ca^{2+} influx channels (Demidchik et al., 2003). The effects of extracellular Cu on PM Ca^{2+} channels and therefore on $[\text{Ca}^{2+}]_{\text{cyt}}$ signaling are likely to be complex. Electrophysiological analysis of *Arabidopsis* root epidermal PM has shown that Cu not only activates Ca^{2+} channels through ROS production (Annexin 1; Laohavisit et al., 2012) but also blocks channels, the molecular identity of which remains unknown (Demidchik et al., 2003).

Catalytic production of ROS by Cu is not the only route to modulating $[\text{Ca}^{2+}]_{\text{cyt}}$. Longer-term exposure to Cu can stimulate exocytosis-mediated ROS production. Lin et al. (2013) found that inhibiting vesicle traffic with brefeldin also inhibited Cu-stimulated ROS production in rice roots. Whether this involved NADPH oxidase or other ROS generators remains to be determined. Rice root $[\text{Ca}^{2+}]_{\text{cyt}}$ increases in response to Cu addition and could link to NADPH oxidases and CIPK activity leading to MAP kinase activation (Yeh et al., 2007). It triggers oxidative stress in *Populus* roots that leads to regulation of

CaM genes, implicating perturbation of $[Ca^{2+}]_{cyt}$ (Guerra et al., 2009).

In common with Cd, excess Cu also alters auxin homeostasis in roots and interferes with NO signaling (Lequeux et al., 2010; Kolbert et al., 2012). In excess, Cu stunts *Arabidopsis* root and root hair elongation and can inhibit lateral root outgrowth (Kolbert et al., 2012). Accumulation of lignin has been observed in both *Arabidopsis* and rice roots (Lequeux et al., 2010; Liu et al., 2015) and in the former accumulation is associated with the endodermis. It would now be interesting to ascertain whether such wall modification changes $[Ca^{2+}]_{cyt}$ signals in central cells or affects $[Ca^{2+}]_{cyt}$ propagative signaling to the shoot. Root tip cell death has also been observed (Huang et al., 2007; Lequeux et al., 2010; Rodrigo-Moreno et al., 2013). Cu-induced cell death in rice roots was attenuated by chelating extracellular Ca^{2+} thus implicating Ca^{2+} influx across the PM (Huang et al., 2007) while in *Arabidopsis* roots cell death was attenuated by addition of Gd^{3+} or verapamil as PM Ca^{2+} channel blockers (Rodrigo-Moreno et al., 2013).

Gd^{3+} is routinely used as a PM Ca^{2+} channel blocker, for example with proven efficacy against *Arabidopsis* root epidermal and root hair HACC, DACC and VICC (Véry and Davies, 2000; Demidchik et al., 2002; Miedema et al., 2008). It is also effective against the root PM hydroxyl radical-activated and H_2O_2 -activated Ca^{2+} influx channels (Demidchik et al., 2003, 2007) and the Annexin1-mediated pathway (Laohavisit et al., 2012). As a consequence Gd^{3+} lowers Ca^{2+} influx (measured with ^{45}Ca ; Demidchik et al., 2002). From this it can be anticipated that Gd^{3+} would have the capacity to dampen stress-induced $[Ca^{2+}]_{cyt}$ signaling in roots, including that mediated by oxidative stress (Demidchik et al., 2003, 2007) and elicited by NaCl (Laohavisit et al., 2013). Pb is less well studied in terms of $[Ca^{2+}]_{cyt}$. Exogenous Ca^{2+} can protect against root Pb accumulation, suggesting competitive uptake (Rodríguez-Hernández et al., 2015). However, Pb was found to evoke Ca^{2+} accumulation and diphenylpicrylhydrazyl-sensitive ROS increase in rice roots, the latter leading to MAPK activation (Huang and Huang, 2008). Entry of Pb into *Arabidopsis* roots involves CNGC1 as its deletion confers greater tolerance but whether Pb permeates this channel is unknown (Sunkar et al., 2000). Pb has also been shown to increase expression of *TPC1* in *Typha* roots (Rodríguez-Hernández et al., 2015) but whether this results in increased $[Ca^{2+}]_{cyt}$ in this or other roots is not known yet.

MECHANICAL STRESS

Roots experience a range of mechanical stimuli, induced as they encounter soil particles or neighboring roots. An increase in $[Ca^{2+}]_{cyt}$ together with an apoplastic alkalinisation are early effects induced by mechanical stimuli. One of the known downstream events from the increase in $[Ca^{2+}]_{cyt}$ is the upregulation of touch-sensitive genes such as *CML12* and *CML24* (Braam, 1992). Mechanically triggered $[Ca^{2+}]_{cyt}$ changes are dependent on the type of stimulus and the responding tissue (Legue et al., 1997; Monshausen et al., 2009). For example, manually bending an *Arabidopsis* root induces a rapid, biphasic increase in $[Ca^{2+}]_{cyt}$

on the convex side of the roots where cells are stretched (Monshausen et al., 2009) while previously compressed cells on the concave side of the roots show a monophasic and less intense increase in $[Ca^{2+}]_{cyt}$ upon return to their resting position. The source of Ca^{2+} is likely to be extracellular (Monshausen et al., 2009; Richter et al., 2009; Kurusu et al., 2012); with internal stores being mobilized to amplify the response (Chehab et al., 2009; Toyota and Gilroy, 2013).

Over-expressing *MCA1* in *Arabidopsis* lead to an enhanced $[Ca^{2+}]_{cyt}$ transient post mechanical stimulation by the addition of a membrane crenator, however, the *mca1* mutant showed no difference from the wild type (Nakagawa et al., 2007). Accordingly, Shih et al. (2014) showed no change in *mca1* apoplastic alkalinisation following root bending, which is closely related to the $[Ca^{2+}]_{cyt}$ signature. Thus some levels of compensation occur in mutants deficient in mechanosensitive Ca^{2+} channels. In contrast to the MCAs, MSLs (*MscS-Like*) were identified in *Arabidopsis* due to their sequence similarity to the bacterial *Mechanosensitive channels of small conductance* (*MscS*) (Haswell et al., 2008). These are encoded by a multiple gene family of 10 members, with MSL9 and MSL10 found in the PM of root cells and required for a mechanosensitive channel activity. They are predominantly anion channels and their relationship to mechano-stimulated $[Ca^{2+}]_{cyt}$ increase has yet to be shown but may be through PM voltage regulation. At this point, it is unclear whether these channels represent mechano-sensors or act downstream of yet unknown mechano-sensors.

Recently, the PM receptor-like kinase FERONIA has been implicated in regulating the *Arabidopsis* mechano-stimulated $[Ca^{2+}]_{cyt}$ increase and downstream transcriptional regulation (Shih et al., 2014). *feronia* mutants lack the second peak of the biphasic increase in $[Ca^{2+}]_{cyt}$ elicited in stretched cells by manual bending. However, the mechanism by which FERONIA can regulate $[Ca^{2+}]_{cyt}$ remains unclear as its kinase activity is not essential and targets are unknown. As far as we know, this study was the first in which a mutant with an aberrant mechano-stimulated $[Ca^{2+}]_{cyt}$ increase also showed a root skewing phenotype. Root skewing (deviation from the vertical when grown on vertical or inclined agar) may be influenced by mechano-sensing. At this point, it is unclear whether FERONIA also plays a role in the mechanical induction of lateral root formation (Richter et al., 2009).

Cold Stress

Cold plays a key role in the regulation of physiology and development; the signaling processes relaying non-stressful temperatures ($12^{\circ}C$ and above) have been reviewed by Wigge (2013). The signaling cascades activated by cold stress (typically $4^{\circ}C$ experimentally) and their relations with hormonal signaling have been reviewed by Knight and Knight (2012), Jeon and Kim (2013), Shi et al. (2015) and Eremina et al. (2016). Most studies on the effect of cold stress on $[Ca^{2+}]_{cyt}$ report on seedlings or leaves. From these it is well established that $[Ca^{2+}]_{cyt}$ elevation involves both Ca^{2+} influx across the PM and release from predominantly vacuolar stores (e.g., Knight et al., 1996; Mazars et al., 1997; Knight and Knight, 2000; Zhu et al., 2013). By

using an extracellular Ca^{2+} -reporting microelectrode, Sulaiman et al. (2012) confirmed that Ca^{2+} influx from the extracellular space is a component of the *Arabidopsis* root response to cold stress. For *Arabidopsis* roots, the $[\text{Ca}^{2+}]_{\text{cyt}}$ response to cold stress measured using aequorin is lower in amplitude and of much shorter duration than leaves (Zhu et al., 2013). Using aequorin Plieth et al. (1999) determined that the faster the rate of cooling *Arabidopsis* roots, the greater the $[\text{Ca}^{2+}]_{\text{cyt}}$ response. Sensitivity and magnitude of the $[\text{Ca}^{2+}]_{\text{cyt}}$ cooling response were enhanced by low temperature but repeated exposure to cold lead to desensitizing of the response. These results imply the existence of Ca^{2+} transport systems that could be regulated at the post-translational and possibly transcriptional levels. In *Arabidopsis*, cold-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation would activate a root PM calcium/calmodulin-regulated receptor-like kinase (CRLK1; Yang et al., 2010). This would then activate a specific mitogen-activated protein kinase kinase kinase (MEKK1) that then targets protein kinase kinase (MKK2; Furuya et al., 2013). This pathway leads to gene activation and freezing tolerance (Furuya et al., 2014). $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation would also activate CIPK3, that acts an intersect with ABA signaling (Kim et al., 2003). Part of the transcriptional response to cold in *Arabidopsis* could be activated by increased $[\text{Ca}^{2+}]_{\text{cyt}}$ through CAMTA3 regulation of the transcriptional regulator CBF2 (C-Repeat/Dehydration Responsive Element Binding Factor2) (Doherty et al., 2009).

Cold stress has been found to depolarise the PM of root cells from cucumber and *Triana bogotensis* (Lyalin and Ktitorova, 1969; Minorsky and Spanswick, 1989), consistent with the effect of Ca^{2+} influx across the PM. A modeling exercise revealed the possible importance of a PM DACC in cold stress-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation of roots (White, 2009). Cold has been shown to activate a PM DACC in leaf protoplasts (Carpaneto et al., 2007) but this has not yet been shown for roots. As microtubule destabilization activates the *Arabidopsis* root PM DACC (Thion et al., 1998), this could link to the increased cold-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ signal observed when microtubules are disrupted in *Nicotiana plumbaginifolia* leaf protoplasts (Mazars et al., 1997). Actin depolymerization is implicated in cold-induced Ca^{2+} influx across the PM into *Medicago sativa* suspension cells (Orvar et al., 2000). This in turn could relate to the cold-induced activation of pear pollen PM HACC that involves actin depolymerization (Wu et al., 2012). It would be timely therefore to investigate root PM for cold effects on HACC and DACC, exploring also the involvement of the cytoskeleton. In common with mechanical stress, cold stress could perturb the PM bilayer sufficiently to change Ca^{2+} channel activity. Work on *M. sativa* suspension cells showed that increasing membrane fluidity at 4°C prevented the Ca^{2+} influx across the PM that is necessary to trigger gene expression for freezing tolerance (Orvar et al., 2000). Conversely, imposing membrane rigidity at normal temperature triggers activation of the Ca^{2+} -dependent MEKK1-MKK2-MPK4 pathway in *Arabidopsis* seedlings (Furuya et al., 2014). The possibility that mechanosensitive channels are involved in cold-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ signaling merits investigation. However, recent work on rice roots indicates another route to Ca^{2+} influx. COLD1 has been identified as a PM activator of the

heterotrimeric G protein subunit α (Ma Y. et al., 2015). It is required for cold-activated Ca^{2+} influx to roots (measured using extracellular Ca^{2+} -reporting microelectrode) and elevation of root $[\text{Ca}^{2+}]_{\text{cyt}}$ (aequorin and cameleon determinations). This indicates that heterotrimeric G proteins lie upstream of PM Ca^{2+} channels in the cold response. In common with other stress responses, NADPH oxidase activity also appears in cold stress and needs to be placed in relation to Ca^{2+} channels. *Arabidopsis* roots exposed to cold stress use AtSRC2 (Soybean gene Regulated by Cold2) to enhance Ca^{2+} activation of the AtRBOHF NADPH Oxidase (Kawarazaki et al., 2013). This suggests that NADPH oxidase activity can lie downstream of Ca^{2+} channel activity. Curtailling cold-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation could involve vacuolar CAX1 activity in *Arabidopsis* (Catala et al., 2003).

CONCLUSION AND FUTURE PROSPECTS

A repeated message from this review is how incomplete our knowledge is of the channels mediating stress-induced $[\text{Ca}^{2+}]_{\text{cyt}}$ increases, and by extension those of organelles. Many members of channel gene families still await characterization. The identification of new families of channels is challenging and will require different approaches linking forward and reverse genetics to electrophysiology. Targets of Ca^{2+} -binding and interacting proteins also require further study. Components common to different abiotic stresses are emerging such as *Arabidopsis* CIPK23 in K^+ and nitrate deprivation. These common regulatory components are likely to represent critical steps where complex stress signals encountered in the soil are integrated in unified responses. Receptor like kinases such as FERONIA or PERK4 have emerged as new components in $[\text{Ca}^{2+}]_{\text{cyt}}$ signaling and perhaps other related proteins will be found to have a role in abiotic stress signaling. Remodeling of calcium signaling machinery after stress is also apparent with the possibility of components common to different stresses. For example, *Arabidopsis* CNGC19 is upregulated under B limitation and salinity stress (Kugler et al., 2009). Finally, $[\text{Ca}^{2+}]_{\text{cyt}}$ -dependent transcriptional responses can be delineated and future work could include the impact of stress-induced calcium signaling on epigenetic inheritance (Sani et al., 2013; Probst and Scheid, 2015).

AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication. KW, EM, SS, and JD wrote the review. KW produced the figures.

FUNDING

Funding for this work was from the BBSRC (BB/K009869/1 and Doctoral Training Programme) and the University of Cambridge Broodbank Trust.

REFERENCES

- Armengaud, P., Beiting, R., and Amtmann, A. (2004). The potassium-dependent transcriptome of *Arabidopsis* reveals a prominent role of jasmonic acid in nutrient signaling. *Plant Physiol.* 136, 2556–2576. doi: 10.1104/pp.104.046482
- Bai, L., Zhang, G., Zhou, Y., Zhang, Z., Wang, W., Du, Y., et al. (2009). Plasma membrane-associated proline-rich extension-like receptor kinase 4, a novel regulator of Ca²⁺ signaling, is required for abscisic acid responses in *Arabidopsis thaliana*. *Plant J.* 60, 314–327. doi: 10.1111/j.1365-313X.2009.03956.x
- Baliardini, C., Meyer, C.-L., Salis, P., Saumitou-Laprade, P., and Verbruggen, N. (2015). CATION EXCHANGER1 cosegregates with cadmium tolerance in the metal hyperaccumulator *Arabidopsis halleri* and plays a role in limiting oxidative stress in *Arabidopsis* Spp. *Plant Physiol.* 169, 549–559. doi: 10.1104/pp.15.01037
- Barberon, M., Vermeer, J. E. M., De Bellis, D., Wang, P., Naseer, S., Anderson, T. G., et al. (2016). Adaptation of root function by nutrient-induced plasticity of endodermal differentiation. *Cell* 164, 447–459. doi: 10.1016/j.cell.2015.12.021
- Batelli, G., Verslues, P. R., Agius, F., Qiu, Q., Fujii, H., Pan, S., et al. (2007). SOS2 promotes salt tolerance in part by interacting with the vacuolar H⁺-ATPase and upregulating its transport activity. *Mol. Cell Biol.* 27, 7781–7790. doi: 10.1128/MCB.00430-07
- Behera, S., Wang, N., Zhang, C., Schmitz-Thom, I., Strohkamp, S., Schuelte, S., et al. (2015). Analyses of Ca²⁺ dynamics using a ubiquitin-10 promoter-driven Yellow Cameleon 3.6 indicator reveal reliable transgene expression and differences in cytoplasmic Ca²⁺ responses in *Arabidopsis* and rice (*Oryza sativa*) roots. *New Phytol.* 206, 751–760. doi: 10.1111/nph.13250
- Bonza, M. C., Loro, G., Behera, S., Wong, A., Kudla, J., and Costa, A. (2013). Analyses of Ca²⁺ accumulation and dynamics in the endoplasmic reticulum of *Arabidopsis* root cells using a genetically encoded cameleon sensor. *Plant Physiol.* 163, 1230–1241. doi: 10.1104/pp.113.226050
- Bonza, M. C., Luoni, L., Olivari, C., and DeMichelis, M. I. (2016). Plant type 2B Ca²⁺-ATPases: the diversity of isoforms of the model plant *Arabidopsis thaliana*. *Adv. Biochem. Health Dis.* 14, 227–241.
- Boursiac, Y., and Harper, J. F. (2007). The origin and function of calmodulin regulated Ca²⁺ pumps in plants. *J. Bioen. Biomem.* 39, 409–414. doi: 10.1007/s10863-007-9104-z
- Braam, J. (1992). Regulated expression of the calmodulin-related TCH genes in cultured *Arabidopsis* cells – induction by calcium and heat shock. *Proc. Natl. Acad. Sci. U.S.A.* 89, 3213–3216. doi: 10.1073/pnas.89.8.3213
- Camacho-Cristobal, J. J., Martin-Rejano, E. M., Begona Herrera-Rodriguez, M., Teresa Navarro-Gochicoa, M., Rexach, J., and Gonzalez-Fontes, A. (2015). Boron deficiency inhibits root cell elongation via an ethylene/auxin/ROS-dependent pathway in *Arabidopsis* seedlings. *J. Expt. Bot.* 66, 3831–3840. doi: 10.1093/jxb/erv186
- Carpaneto, A., Ivashikina, N., Levchenko, V., Krol, E., Jeworutzki, E., Zhu, J.-K., et al. (2007). Cold transiently activates calcium-permeable channels in *Arabidopsis* mesophyll cells. *Plant Physiol.* 143, 487–494. doi: 10.1104/pp.106.090928
- Carroll, A. D., Moyon, C., Van Kesteren, P., Tooke, F., Battay, N. H., and Brownlee, C. (1998). Ca²⁺, annexins, and GTP modulate exocytosis from maize root cap protoplasts. *Plant Cell* 10, 1267–1276. doi: 10.1105/tpc.10.8.1267
- Catala, R., Santos, E., Alonso, J. M., Ecker, J. R., Martinez-Zapater, J. M., and Salinas, J. (2003). Mutations in the Ca²⁺/H⁺ transporter CAX1 increase CBF/DREB1 expression and the cold-acclimation response in *Arabidopsis*. *Plant Cell* 15, 2940–2951. doi: 10.1105/tpc.015248
- Chehab, E. W., Eich, E., and Braam, J. (2009). Thigmomorphogenesis: a complex plant response to mechanostimulation. *J. Expt. Bot.* 60, 43–56. doi: 10.1093/jxb/ern315
- Cheng, N. H., Pittman, J. K., Zhu, J. K., and Hirschi, K. D. (2004). The protein kinase SOS2 activates the *Arabidopsis* H⁺/Ca²⁺ antiporter CAX1 to integrate calcium transport and salt tolerance. *J. Biol. Chem.* 279, 2922–2926. doi: 10.1074/jbc.M309084200
- Cheong, Y. H., Pandey, G. K., Grant, J. J., Baticic, O., Li, L., Kim, B.-G., et al. (2007). Two calcineurin B-like calcium sensors, interacting with protein kinase CIPK23, regulate leaf transcription and root potassium uptake in *Arabidopsis*. *Plant J.* 52, 223–239. doi: 10.1111/j.1365-313X.2007.03236.x
- Chmielowska-Bak, J., Gryl, J., Rucinska-Sobkowiak, R., Arasimowicz-Jelonek, M., and Deckert, J. (2014). The new insights into cadmium sensing. *Front. Plant Sci.* 5:245. doi: 10.3389/fpls.2014.00245
- Choi, W.-G., Toyota, M., Kim, S.-H., Hilleary, R., and Gilroy, S. (2014). Salt stress-induced Ca²⁺ waves are associated with rapid, long-distance root-to-shoot signaling in plants. *Proc. Natl. Acad. Sci. U.S.A.* 111, 6497–6502. doi: 10.1073/pnas.1319955111
- Christensen, A., Svensson, K., Persson, S., Jung, J., Michalak, M., Widell, S., et al. (2008). Functional characterization of *Arabidopsis* calreticulin1a: A key alleviator of endoplasmic reticulum. *Plant Cell Physiol.* 49, 912–924. doi: 10.1093/pcp/pcn065
- Chung, J. S., Zhu, J. K., Bressan, R. A., Hasagawa, P. M., and Shi, H. (2008). Reactive oxygen species mediate Na⁺-induced SOS1 mRNA stability in *Arabidopsis*. *Plant J.* 53, 554–565. doi: 10.1111/j.1365-313X.2007.03364.x
- Clemens, S., Antosiewicz, D. M., Ward, J. M., Schachtman, D. B., and Schroeder, J. I. (1998). The plant cDNA LCT1 mediates the uptake of calcium and cadmium in yeast. *Proc. Natl. Acad. Sci. U.S.A.* 95, 12043–12048. doi: 10.1073/pnas.95.20.12043
- Connorton, J. M., Webster, R. E., Cheng, N. H., and Pittman, J. K. (2012). Knockout of multiple *Arabidopsis* cation/H⁺ exchangers suggests isoform-specific roles in metal stress response, germination and see mineral nutrition. *PLoS ONE* 7:e47455. doi: 10.1371/journal.pone.0047455
- Dark, A. M., Demidchik, V., Richards, S. L., Shabala, S. N., and Davies, J. M. (2011). Release of extracellular purines from plant roots and effect on ion fluxes. *Plant Signal. Behav.* 6, 1855–1857. doi: 10.4161/psb.6.11.17014
- Davies, J. M. (2014). Annexin-mediated calcium signaling in plants. *Plants* 3, 128–140. doi: 10.3390/plants3010128
- Demidchik, V., Bowen, H. C., Maathuis, F. J. M., Shabala, S. N., Tester, M. A., White, P. J., et al. (2002). *Arabidopsis thaliana* root non-selective cation channels mediate calcium uptake and are involved in growth. *Plant J.* 32, 799–808. doi: 10.1046/j.1365-313X.2002.01467.x
- Demidchik, V., Cuin, T. A., Svistunenko, D., Smith, S. J., Miller, A. J., Shabala, S., et al. (2010). *Arabidopsis* root K⁺-efflux conductance activated by hydroxyl radicals: single-channel properties, genetic basis and involvement in stress-induced death. *J. Cell Sci.* 123, 1468–1479. doi: 10.1242/jcs.064352
- Demidchik, V., Shabala, S. N., Coutts, K. B., Tester, M. A., and Davies, J. M. (2003). Free oxygen radicals regulate plasma membrane Ca²⁺ and K⁺-permeable channels in plant root cells. *J. Cell Sci.* 116, 81–88. doi: 10.1242/jcs.00201
- Demidchik, V., Shabala, S. N., and Davies, J. M. (2007). Spatial variation in H₂O₂ response of *Arabidopsis thaliana* epidermal Ca²⁺ flux and plasma membrane Ca²⁺ channels. *Plant J.* 49, 377–386. doi: 10.1111/j.1365-313X.2006.02971.x
- Demidchik, V., Shang, Z., Shin, R., Thompson, E. P., Rubio, L., Laohavisit, A., et al. (2009). Plant extracellular ATP signalling by plasma membrane NADPH oxidase and Ca²⁺ channels. *Plant J.* 58, 903–913. doi: 10.1111/j.1365-313X.2009.03830.x
- Deng, Y., Srivastava, R., and Howell, S. H. (2013). Protein kinase and ribonuclease domains of IRE1 confer stress tolerance, vegetative growth and reproductive development. *Proc. Natl. Acad. Sci. U.S.A.* 110, 19633–19638. doi: 10.1073/pnas.1314749110
- DeWald, D. B., Torabinejad, J., Jones, C. A., Shope, J. C., Cangelosi, A. R., Thompson, J. E., et al. (2001). Rapid accumulation of phosphatidylinositol 4,5-bisphosphate and inositol 1,4,5-trisphosphate correlates with calcium mobilization in salt-stressed *Arabidopsis*. *Plant Physiol.* 126, 759–769. doi: 10.1104/pp.126.2.759
- Dinnyen, J. R., Long, T. A., Wang, J. Y., Jung, J. W., Mace, D., Pointer, S., et al. (2008). Cell identity mediates the response of *Arabidopsis* roots to abiotic stress. *Science* 320, 942–945. doi: 10.1126/science.1153795
- Doherty, C. J., Van Buskirk, H. A., Myers, S. J., and Thomashow, M. F. (2009). Roles for *Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. *Plant Cell* 21, 972–984. doi: 10.1105/tpc.108.063958
- Donaldson, L., Ludidi, N., Knight, M. R., Gehring, C., and Denby, K. (2004). Salt and osmotic stress cause rapid increases in *Arabidopsis thaliana* cGMP levels. *FEBS Letts.* 569, 317–320. doi: 10.1016/j.febslet.2004.06.016
- Drerup, M. M., Schluucking, K., Hashimoto, K., Manishankar, P., Steinhorst, L., Kuchitsu, K., et al. (2013). The Calcineurin B-Like calcium sensors CBL1 and CBL9 together with their interacting protein kinase CIPK26 regulate the *Arabidopsis* NADPH oxidase RBOHF. *Mol. Plant* 6, 559–569. doi: 10.1093/mp/sst009

- Dressler, L., Golbik, R., and Ulbrich-Hofmann, R. (2014). Lanthanides as substitutes for calcium ions in the activation of plant alpha-type phospholipase D. *Biol. Chem.* 395, 791–799. doi: 10.1515/hsz-2014-0112
- Eremina, M., Rozhon, W., and Poppenberger, B. (2016). Hormonal control of cold stress responses in plants. *Cell. Mol. Life Sci.* 73, 797–810. doi: 10.1007/s00018-015-2089-6
- Fagerstedt, K. V. (2010). “Programmed cell death and aerenchyma formation under hypoxia,” in *Waterlogging Signalling and Tolerance in Plants*, eds S. Mancuso and S. Shabala (Berlin: Springer), 99–118.
- Fan, J.-L., Wei, X.-Z., Wan, L.-C., Zhang, L.-Y., Zhao, X.-Q., Liu, W.-Z., et al. (2011). Disarrangement of actin filaments and Ca^{2+} gradient by $CdCl_2$ alters cell wall construction in *Arabidopsis thaliana* root hairs by inhibiting vesicular trafficking. *J. Plant Physiol.* 168, 1157–1167. doi: 10.1016/j.jplph.2011.01.031
- Fischer, C., Kugler, A., Hoth, S., and Dietrich, P. (2013). An IQ domain mediates the interaction with calmodulin in a plant cyclic nucleotide-gated channel. *Plant Cell Physiol.* 54, 573–584. doi: 10.1093/pcp/pct021
- Foreman, J., Demidchik, V., Bothwell, J. H. F., Mylona, P., Miedema, H., Torres, M. A., et al. (2003). Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* 422, 442–446. doi: 10.1038/nature01485
- Fry, S. C., Miller, J. G., and Dumville, J. C. (2002). A proposed role for copper ions in cell wall loosening. *Plant Soil* 247, 57–67. doi: 10.1023/A:1021140022082
- Furuya, T., Matsuoka, D., and Nanmori, T. (2013). Phosphorylation of *Arabidopsis thaliana* MEKK1 via Ca^{2+} signaling as a part of the cold stress response. *J. Plant Res.* 126, 833–840. doi: 10.1007/s10265-013-0576-0
- Furuya, T., Matsuoka, D., and Nanmori, T. (2014). Membrane rigidification functions upstream of the MEKK1-MKK2-MPK4 cascade during cold acclimation in *Arabidopsis thaliana*. *FEBS Lett.* 588, 2025–2030. doi: 10.1016/j.febslet.2014.04.032
- Gobert, A., Park, G., Amtmann, A., Sanders, D., and Maathuis, F. J. M. (2006). *Arabidopsis thaliana* cyclic nucleotide-gated channel 3 forms a non-selective ion transporter involved in germination and cation transport. *J. Expt. Bot.* 57, 791–800. doi: 10.1093/jxb/erj064
- Goldbach, H. E., Yu, Q., Wiegand, R., Schulz, M., Wimmer, M., Findelee, P., et al. (2001). Rapid response reactions of roots to boron deprivation. *J. Plant Nutr. Soil Sci.* 164, 173–181. doi: 10.1002/1522-2624(200104)164:2<173::AID-JPLN173>3.3.CO;2-6
- González-Fontes, A., Herrera-Rodriguez, M. B., Martín-Rejano, E. M., Navarro-Góchicoa, M. T., Rexach, J., and Camacho-Cristobal, J. J. (2016). Root responses to boron deficiency mediated by ethylene. *Front. Plant Sci.* 6:1103. doi: 10.3389/fpls.2015.01103
- Gonzali, S., Loreti, E., Cardarelli, F., Novi, G., Parlanti, S., Pucciariello, C., et al. (2015). Universal stress protein HRU1 mediates ROS homeostasis under anoxia. *Nat. Plants* 1, 15151. doi: 10.1038/NPLANTS.2015.151
- Guerra, F., Duplessis, S., Kohler, A., Martin, F., Tapia, J., Lebed, P., et al. (2009). Gene expression analysis of *Populus deltoides* roots subjected to copper stress. *Environ. Exp. Bot.* 67, 335–344. doi: 10.1016/j.envexpbot.2009.08.004
- Guo, J., Zeng, W., Chen, Q., Lee, C., Chen, L., Yang, Y., et al. (2016). Structure of the voltage-gated two-pore channel TPC1 from *Arabidopsis thaliana*. *Nature* 531, 196–201. doi: 10.1038/nature16446
- Guo, K.-M., Babourina, O., Christopher, D. A., Borsics, T., and Rengel, Z. (2008). The cyclic nucleotide-gated channel, AtCNGC10, influences salt tolerance in *Arabidopsis*. *Physiol. Plant.* 134, 499–507. doi: 10.1111/j.1399-3054.2008.01157.x
- Haswell, E. S., Peyronnet, R., Barbier-Brygoo, H., Meyerowitz, E. M., and Frachisse, J. M. (2008). Two MscS homologs provide mechanosensitive channel activities in the *Arabidopsis* root. *Curr. Biol.* 18, 730–734. doi: 10.1016/j.cub.2008.04.039
- Ho, C.-H., Lin, S.-H., Hu, H.-C., and Tsay, Y.-F. (2009). CHL1 functions as a nitrate sensor in plants. *Cell* 138, 1184–1194. doi: 10.1016/j.cell.2009.07.004
- Hossain, M. A., Henriquez-Valencia, C., Gómez-Páez, M., Medina, J., Orellana, A., Vicente-Carbajosa, J., et al. (2016). Identification of novel components of the unfolded protein response in *Arabidopsis*. *Front. Plant Sci.* 7:650. doi: 10.3389/fpls.2016.00650
- Hou, C. C., Tian, W., Kleist, T., He, K., Garcia, V., Bai, F., et al. (2014). DUF221 proteins are a family of osmosensitive calcium-permeable cation channels conserved across eukaryotes. *Cell Res.* 24, 632–635. doi: 10.1038/cr.2014.14
- Hou, Q., Ufer, G., and Bartels, D. (2016). Lipid signalling in plant responses to abiotic stress. *Plant Cell Environ.* 39, 1029–1048. doi: 10.1111/pce.12666
- Hu, H. C., Wang, Y. Y., and Tsay, Y. F. (2009). AtCIPK8, a CBL-interacting protein kinase, regulates the low-affinity phase of the primary nitrate response. *Plant J.* 57, 264–278. doi: 10.1111/j.1365-313X.2008.03685.x
- Hu, Y. F., Zhou, G., Na, X. F., Yang, L., Nan, W. B., Liu, X., et al. (2013). Cadmium interferes with maintenance of auxin homeostasis in *Arabidopsis* seedlings. *J. Plant Physiol.* 170, 965–975. doi: 10.1016/j.jplph.2013.02.008
- Hua, B. G., Mercier, R. W., and Zielinski, R. E. (2003). Functional interaction of calmodulin with a plant cyclic nucleotide gated cation channel. *Plant Physiol. Biochem.* 41, 945–954. doi: 10.1016/j.plaphy.2003.07.006
- Huang, T.-L., and Huang, H.-J. (2008). ROS and CDPK-like kinase-mediated activation of MAP kinase in rice roots exposed to lead. *Chemosphere* 71, 1377–1385. doi: 10.1016/j.chemosphere.2007.11.031
- Huang, W.-C., Huang, D.-D., Chien, P.-S., Yeh, C.-M., Chen, P.-Y., Chi, W.-C., et al. (2007). Protein tyrosine dephosphorylation during copper-induced cell death in rice roots. *Chemosphere* 69, 55–62. doi: 10.1016/j.chemosphere.2007.04.073
- Hunt, L., Otterhag, L., Lee, J. C., Lasheen, T., Hunt, J., Seki, M., et al. (2004). Gene-specific expression and calcium activation of *Arabidopsis thaliana* phospholipase C isoforms. *New Phytol.* 162, 643–654. doi: 10.1111/j.1469-8137.2004.01069.x
- Isner, J. C., Nühse, T., and Maathuis, F. J. M. (2012). The cyclic nucleotide cGMP is involved in plant hormone signalling and alters phosphorylation of *Arabidopsis thaliana* root proteins. *J. Expt. Bot.* 63, 3199–3205. doi: 10.1093/jxb/ers045
- Jakubowska, D., Janicka-Russak, M., Kabala, K., Migocka, M., and Reda, M. (2015). Modification of plasma membrane NADPH oxidase activity in cucumber seedling roots in response to cadmium stress. *Plant Sci.* 234, 50–59. doi: 10.1016/j.plantsci.2015.02.005
- Jeon, J., and Kim, J. (2013). Cold stress signaling networks in *Arabidopsis*. *J. Plant Biol.* 56, 69–76. doi: 10.1007/s12374-013-0903-y
- Jiang, H. B., Wang, S. F., Yang, F., Zhang, Z. H., Qiu, H. C., Yi, Y., et al. (2015). Plant growth, nitrate content and Ca signaling in wheat (*Triticum aestivum* L.) roots under different nitrate supply. *Plant Sci. J.* 33, 362–368.
- Jiao, Y., Sun, L., Song, Y., Wang, L., Liu, L., Zhang, L., et al. (2013). AtrbohD and AtrbohF positively regulate abscisic acid-inhibited primary root growth by affecting Ca^{2+} signalling and auxin response of roots in *Arabidopsis*. *J. Expt. Bot.* 64, 4183–4192. doi: 10.1093/jxb/ert228
- Jin, Y. K., Jing, W., Zhang, Q., and Zhang, W. H. (2015). Cyclic nucleotide gated channel 10 negatively regulates salt tolerance by mediating Na^+ transport in *Arabidopsis*. *J. Plant Res.* 128, 211–220. doi: 10.1007/s10265-014-0679-2
- Kamano, S., Kume, S., Iida, K., Lei, K.-J., Nakano, M., Nakayama, Y., et al. (2015). Transmembrane topologies of Ca^{2+} -permeable mechanosensitive channels MCA1 and MCA2 in *Arabidopsis thaliana*. *J. Biol. Chem.* 290, 30901–30909. doi: 10.1074/jbc.M115.692574
- Kanehara, K., Yu, C.-Y., Cho, Y., Cheong, W.-F., Torta, F., Shui, G., et al. (2015). *Arabidopsis* AtPLC2 is a primary phosphoinositide-specific phospholipase C in phosphoinositide metabolism and the endoplasmic reticulum stress response. *PLOS Genet.* 11:e1005511. doi: 10.1371/journal.pgen.1005511
- Kasajima, I., Ide, Y., Hirai, M. Y., and Fujiwara, T. (2010). WRKY6 is involved in the response to boron deficiency in *Arabidopsis thaliana*. *Physiol. Plant.* 139, 80–92. doi: 10.1111/j.1399-3054.2010.01349.x
- Kawarazaki, T., Kimura, S., Iizuka, A., Hanamata, S., Nibori, H., Michikawa, M., et al. (2013). A low temperature-inducible protein AtSRC2 enhances the ROS-producing activity of NADPH oxidase AtrbohF. *Biochim. Biophys. Acta Mol. Cell Res.* 1833, 2775–2780. doi: 10.1016/j.bbamcr.2013.06.024
- Kiegle, E., Moore, C. A., Haseloff, J., Tester, M. A., and Knight, M. R. (2000). Cell-type-specific calcium responses to drought, salt and cold in the *Arabidopsis* root. *Plant J.* 23, 267–278. doi: 10.1046/j.1365-313x.2000.00786.x
- Kim, K. N., Cheong, Y. H., Grant, J. J., Pandey, G. K., and Luan, S. (2003). CIPK3, a calcium sensor-associated protein kinase that regulates abscisic acid and cold signal transduction in *Arabidopsis*. *Plant Cell* 15, 411–423. doi: 10.1105/tpc.006858
- Kim, Y., Wang, M., Bai, Y., Zeng, Z., Guo, F., Han, N., et al. (2014). Bcl-2 suppresses activation of VPEs by inhibiting cytosolic Ca^{2+} level with elevated K^+ efflux in NaCl-induced PCD in rice. *Plant Physiol. Biochem.* 80, 168–175. doi: 10.1016/j.plaphy.2014.04.002

- Kintzer, A. F., and Stroud, R. M. (2016). Structure, inhibition and regulation of two-pore channel TPC1 from *Arabidopsis thaliana*. *Nature* 531, 258–264. doi: 10.1038/nature17194
- Knight, H., and Knight, M. R. (2000). Imaging spatial and cellular characteristics of low temperature calcium signature after cold acclimation in *Arabidopsis*. *J. Expt. Bot.* 51, 1679–1686. doi: 10.1093/jexbot/51.351.1679
- Knight, H., Trewavas, A. J., and Knight, M. R. (1996). Cold calcium signaling in *Arabidopsis* involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* 8, 489–503. doi: 10.1105/tpc.8.3.489
- Knight, H., Trewavas, A. J., and Knight, M. R. (1997). Calcium signalling in *Arabidopsis thaliana* responding to drought and salinity. *Plant J.* 12, 1067–1078. doi: 10.1046/j.1365-313X.1997.12051067.x
- Knight, M. R., and Knight, H. (2012). Low-temperature perception leading to gene expression and cold tolerance in higher plants. *New Phytol.* 195, 737–751. doi: 10.1111/j.1469-8137.2012.04239.x
- Kobayashi, B., Matoh, T., and Azuma, J. (1996). Two chains of rhamnogalacturonan II are cross-linked by borate-diol ester bonds in higher plant cell walls. *Plant Physiol.* 110, 1017–1020.
- Kolbert, Z., Pető, A., Lehotai, N., Feigl, G., and Erdei, L. (2012). Long-term copper (Cu^{2+}) exposure impacts on auxin, nitric oxide (NO) metabolism and morphology of *Arabidopsis thaliana* L. *Plant Growth Regul.* 68, 151–159. doi: 10.1007/s10725-012-9701-7
- Koshiba, T., Ishihara, A., and Matoh, T. (2010). Boron nutrition of cultured Tobacco BY-2 Cells. VI. Calcium is involved in early responses to boron deprivation. *Plant Cell Physiol.* 51, 323–327. doi: 10.1093/pcp/pcp179
- Kugler, A., Koehler, B., Palme, K., Wolff, P., and Dietrich, P. (2009). Salt-dependent regulation of a CNG channel subfamily in *Arabidopsis*. *BMC Plant Biol.* 9:140. doi: 10.1186/1471-2229-9-140
- Kurusu, T., Nishikawa, D., Yamazaki, Y., Gotoh, M., Nakano, M., Hamada, H., et al. (2012). Plasma membrane protein OsMCA1 is involved in regulation of hypo-osmotic shock-induced Ca^{2+} influx and modulates generation of reactive oxygen species in cultured rice cells. *BMC Plant Biol.* 12:11. doi: 10.1186/1471-2229-12-11
- Laohavisit, A., and Davies, J. M. (2011). Annexins. *New Phytol.* 189, 40–53. doi: 10.1111/j.1469-8137.2010.03533.x
- Laohavisit, A., Richards, S. L., Shabala, L., Chen, C., Colaço, R., Swarbrick, S. M., et al. (2013). Salinity-induced calcium signaling and root adaptation in *Arabidopsis* require the regulatory protein annexin1. *Plant Physiol.* 163, 253–262. doi: 10.1104/pp.113.217810
- Laohavisit, A., Shang, Z.-L., Rubio, L., Cuin, T. A., Véry, A. A., Wang, A. H., et al. (2012). *Arabidopsis* annexin 1 mediates the radical-activated plasma membrane Ca^{2+} and K^{+} -permeable conductance in root cells. *Plant Cell* 24, 1522–1533. doi: 10.1105/tpc.112.097881
- Latz, A., Mehlmer, N., Zapf, S., Mueller, T. D., Wurzinger, B., Pfister, B., et al. (2013). Salt stress triggers phosphorylation of the *Arabidopsis* vacuolar K^{+} channel TPK1 by calcium-dependent protein kinases (CDPKs). *Mol. Plant* 6, 1274–1289. doi: 10.1093/mp/sss158
- Legue, V., Blancaflor, E., Wymer, C., Perbal, G., Fahtin, D., and Gilroy, S. (1997). Cytoplasmic free calcium in *Arabidopsis* roots changes in response to touch but not gravity. *Plant Physiol.* 114, 789–800. doi: 10.1104/pp.114.3.789
- Lequeux, H., Hermans, C., Lutts, S., and Verbruggen, N. (2010). Response to copper excess in *Arabidopsis thaliana*: impact on the root system architecture, hormone distribution, lignin accumulation and mineral profile. *Plant Physiol. Biochem.* 48, 673–682. doi: 10.1016/j.plaphy.2010.05.005
- Léran, S., Edel, K. H., Pervent, M., Hashimoto, K., Corratge-Faillie, C., Offenborn, J. N., et al. (2015). Nitrate sensing and uptake in *Arabidopsis* are enhanced by ABI2, a phosphatase inactivated by the stress hormone abscisic acid. *Sci. Signal.* 8, ra43. doi: 10.1126/scisignal.aaa4829
- Li, G., Boudsocq, M., Hem, S., Vialaret, J., Rossignol, M., Maurel, C., et al. (2015). The calcium-dependent protein kinase CPK7 acts on root hydraulic conductivity. *Plant Cell Environ.* 38, 1312–1320. doi: 10.1111/pce.12478
- Li, J., Long, Y., Qi, G. N., Xu, Z. J., Wu, W. H., and Wang, Y. (2014). The OsAKT1 channel is critical for K^{+} uptake in rice roots and is modulated by the rice CBL1-CIPK23 complex. *Plant Cell* 26, 3387–3402. doi: 10.1105/tpc.114.123455
- Li, L., Kim, B.-G., Cheong, Y. H., Pandey, G. K., and Luan, S. (2006). A Ca^{2+} signalling pathway regulates a K^{+} channel for low K^{+} response in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 103, 12625–12630. doi: 10.1073/pnas.0605129103
- Li, P., Zhao, C. Z., Zhang, Y. Q., Wang, X. M., Wang, X. Y., Wang, J. F., et al. (2016). Calcium alleviates cadmium-induced inhibition on root growth by maintaining auxin homeostasis in *Arabidopsis* seedlings. *Protoplasma* 253, 185–200. doi: 10.1007/s00709-015-0810-9
- Li, P. H., Zhang, G. Y., Gonzales, N., Guo, Y. Q., Hu, H. H., Park, S. H., et al. (2016). Ca^{2+} -regulated and diurnal rhythm-regulated $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger AtNCL affects flowering time and auxin signalling in *Arabidopsis*. *Plant Cell Environ.* 39, 377–392. doi: 10.1111/pce.12620
- Li, S., Yu, J., Zhu, M., Zhao, F., and Luan, S. (2012). Cadmium impairs ion homeostasis by altering K^{+} and Ca^{2+} channel activities in rice root hairs. *Plant Cell Environ.* 35, 1998–2013. doi: 10.1111/j.1365-3040.2012.02532.x
- Lin, C. Y., Trinh, N. N., Fu, S. F., Hsiung, Y. C., Chia, L. C., Lin, C. W., et al. (2013). Comparison of early transcriptome responses to copper and cadmium in rice roots. *Plant Mol. Biol.* 81, 507–522. doi: 10.1007/s11103-013-0020-9
- Liu, J. P., Ishitani, M., Halfter, U., Kim, C. S., and Zhu, J. K. (2000). The *Arabidopsis thaliana* SOS2 gene encodes a protein kinase that is required for salt tolerance. *Proc. Natl. Acad. Sci. U.S.A.* 97, 3730–3734. doi: 10.1073/pnas.97.7.3730
- Liu, J.-X., Srivastava, R., Che, P., and Howell, S. H. (2007). Salt stress responses in *Arabidopsis* utilize a signal transduction pathway related to endoplasmic reticulum stress signaling. *Plant J.* 51, 897–909. doi: 10.1111/j.1365-313X.2007.03195.x
- Liu, L., Li, Q., Yin, B., Zhang, H., Lin, B., Wu, Y., et al. (2011). The endoplasmic reticulum-associated degradation is necessary for plant salt tolerance. *Cell Res.* 21, 957–969. doi: 10.1038/cr.2010.181
- Liu, Q., Zheng, L., He, F., Zhao, F.-J., Shen, Z., and Zheng, L. (2015). Transcriptional and physiological analyses identify a regulatory role for hydrogen peroxide in the lignin biosynthesis of copper-stressed rice roots. *Plant Soil* 387, 323–336. doi: 10.1007/s11104-014-2290-7
- Lokdarshi, A., Conner, W. C., McClintock, C., Li, T., and Roberts, D. M. (2016). *Arabidopsis* CML38, a calcium sensor that localizes to ribonucleoprotein complexes under hypoxia stress. *Plant Physiol.* 170, 1046–1059. doi: 10.1104/pp.15.01407
- Lu, G., Wang, X., Liu, J., Yu, K., Gao, Y., Liu, H., et al. (2014). Application of T-DNA activation tagging to identify glutamate receptor-like genes that enhance drought tolerance in plants. *Plant Cell Rep.* 33, 617–631. doi: 10.1007/s00299-014-1586-7
- Lyalin, O. O., and Kitorova, I. N. (1969). Resting potential of the root hair of *Trianea bogotensis*. *Sov. Plant Phys.* 16, 214–221.
- Ma, L., Zhang, H., Sun, L., Jiao, Y., Zhang, G., Miao, C., et al. (2012). NADPH oxidase AtrbohD and AtrbohF function in ROS-dependent regulation of $\text{Na}^{+}/\text{K}^{+}$ homeostasis in *Arabidopsis* under salt stress. *J. Expt. Bot.* 63, 305–317. doi: 10.1093/jxb/err280
- Ma, Q., Tang, R. J., Zheng, X. J., Wang, S. M., and Luan, S. (2015). The calcium sensor CBL7 modulates plant responses to low nitrate in *Arabidopsis*. *Biochem. Biophys. Res. Commun.* 468, 59–65. doi: 10.1016/j.bbrc.2015.10.164
- Ma, Y., Dai, X. Y., Xu, Y. Y., Luo, W., Zheng, X. M., Zeng, D. L., et al. (2015). COLD1 confers chilling tolerance in rice. *Cell* 160, 1209–1221. doi: 10.1016/j.cell.2015.01.046
- Maathuis, F. J. M. (2014). Sodium in plants: perception, signalling and regulation of sodium fluxes. *J. Expt. Bot.* 65, 849–858. doi: 10.1093/jxb/ert326
- Mazars, C., Thion, L., Thuleau, P., Graziana, A., Knight, M. R., Moreau, M., et al. (1997). Organization of cytoskeleton controls the changes in cytosolic calcium of cold-shocked *Nicotiana plumbaginifolia* protoplasts. *Cell Calcium* 22, 413–420. doi: 10.1016/S0143-4160(97)90025-7
- McAinsh, M. R., and Pittman, J. K. (2009). Shaping the calcium signature. *New Phytol.* 181, 275–294. doi: 10.1111/j.1469-8137.2008.02682.x
- McLoughlin, F., Galvan-Ampudia, C. S., Julkowska, M. M., Caarls, L., van der Does, D., Lauriere, C., et al. (2012). The Snf1-related protein kinases SnRK2.4 and SnRK2.10 are involved in maintenance of root system architecture during salt stress. *Plant J.* 72, 436–449. doi: 10.1111/j.1365-313X.2012.05089.x
- Mehlmer, N., Wurzinger, B., Stael, S., Hofmann-Rodrigues, D., Csaszar, E., Pfister, B., et al. (2010). The Ca^{2+} -dependent protein kinase CPK3 is required for MAPK-independent salt-stress acclimation in *Arabidopsis*. *Plant J.* 63, 484–498. doi: 10.1111/j.1365-313X.2010.04257.x
- Mei, H., Cheng, N. H., Zhao, J., Park, S., Escareno, R. A., Pittman, J. K., et al. (2009). Root development under metal stress in *Arabidopsis thaliana* requires

- the H⁺/cation antiporter CAX4. *New Phytol.* 183, 95–105. doi: 10.1111/j.1469-8137.2009.02831.x
- Meringer, M. V., Villasuso, A. L., Peppino, M. M., Usorach, J., Pasquare, S. J., Giusto, N. M., et al. (2016). Saline and osmotic stresses stimulate PLD/diacylglycerol kinase activities and increase the level of phosphatidic acid and proline in barley roots. *Environ. Expt. Bot.* 128, 69–78. doi: 10.1016/j.envexpbot.2016.03.011
- Miedema, H., Demidchik, V., Véry, A.-A., Bothwell, J. H. F., Brownlee, C., and Davies, J. M. (2008). Two voltage-dependent calcium channels co-exist in the apical plasma membrane of *Arabidopsis thaliana* root hairs. *New Phytol.* 179, 378–385. doi: 10.1111/j.1469-8137.2008.02465.x
- Miller, G., Schlauch, K., Tam, R., Cortes, D., Torres, M. A., Shulaev, V., et al. (2009). The plant NADPH oxidase RBOHD mediates rapid systemic signaling in response to diverse stimuli. *Sci. Signal.* 2, ra45. doi: 10.1126/scisignal.2000448
- Minorsky, P. V., and Spanswick, R. M. (1989). Electrophysiological evidence for a role for calcium in temperature sensing by roots of cucumber seedlings. *Plant Cell Environ.* 12, 137–143. doi: 10.1111/j.1365-3040.1989.tb01925.x
- Monshausen, G. B., Bibikova, T. N., Weisenseel, M. H., and Gilroy, S. (2009). Ca²⁺ regulates reactive oxygen species production and pH during mechanosensing in *Arabidopsis* roots. *Plant Cell* 21, 2341–2356. doi: 10.1105/tpc.109.068395
- Moore, C. A., Bowen, H. C., Scrase-Field, S., Knight, M. R., and White, P. J. (2002). The deposition of suberin lamellae determines the magnitude of cytosolic Ca²⁺ elevations in root endodermal cells subjected to cooling. *Plant J.* 30, 457–465. doi: 10.1046/j.1365-313X.2002.01306.x
- Mortimer, J. C., Laohavisit, A., Miedema, H., and Davies, J. M. (2008). Voltage, reactive oxygen species and the influx of calcium. *Plant Sig. Behav.* 3, 698–699. doi: 10.4161/psb.3.9.6405
- Mühling, K. H., Wimmer, M., and Goldbach, H. E. (1998). Apoplastic and membrane associated Ca²⁺ in leaves and roots as affected by boron deficiency. *Physiol. Plant.* 102, 179–184. doi: 10.1034/j.1399-3054.1998.1020204.x
- Nakagawa, Y., Katagiri, T., Shinozaki, K., Qi, Z., Tatsumi, H., Furuichi, T., et al. (2007). *Arabidopsis* plasma membrane protein crucial for Ca²⁺ influx and touch sensing in roots. *Proc. Natl. Acad. Sci. U.S.A.* 104, 3639–3644. doi: 10.1073/pnas.0607703104
- Nieves-Cordones, M., Miller, A. J., Aleman, F., Martinez, V., and Rubio, F. (2008). A putative role for the plasma membrane potential in the control of the expression of the gene encoding the tomato high-affinity potassium transporter HAK5. *Plant Mol. Biol.* 68, 521–532. doi: 10.1007/s11103-008-9388-3
- Oiwa, Y., Kitayama, K., and Matoh, T. (2013). Boron deprivation immediately causes cell death in growing roots of *Arabidopsis thaliana* (L.) Heynh. *Soil Sci. Plant Nutr.* 59, 621–627. doi: 10.1080/00380768.2013.813382
- Ordenes, V. R., Moreno, I., Maturana, D., Norambuena, L., Trewavas, A. J., and Orellana, A. (2012). In vivo analysis of the calcium signature in the plant Golgi apparatus reveals unique dynamics. *Cell Calcium* 52, 397–404. doi: 10.1016/j.ceca.2012.06.008
- Orvar, B. L., Sangwan, V., Omann, F., and Dhindsa, R. S. (2000). Early steps in cold sensing by plant cells: the role of actin cytoskeleton and membrane fluidity. *Plant J.* 23, 785–794. doi: 10.1046/j.1365-313x.2000.00845.x
- Pandey, N., Ranjan, A., Pant, P., Tripathi, R. K., Ateek, F., Pandey, H. P., et al. (2013). CAMTA 1 regulates drought responses in *Arabidopsis thaliana*. *BMC Genomics* 14:216. doi: 10.1186/1471-2164-14-216
- Peiter, E., Maathuis, F. J. M., Mills, L. N., Knight, H., Pelloux, J., Hetherington, A. M., et al. (2005). The vacuolar Ca²⁺-activated channel TPC1 regulates germination and stomatal movement. *Nature* 434, 404–408. doi: 10.1038/nature03381
- Perfus-Barbeoch, L., Leonhardt, N., Vavasseur, A., and Forestier, C. (2002). Heavy metal toxicity: cadmium permeates through calcium channels and disturbs the plant water status. *Plant J.* 32, 539–548. doi: 10.1046/j.1365-313X.2002.01442.x
- Petrov, V., Hille, J., Mueller-Roeber, B., and Gechev, T. S. (2015). ROS-mediated abiotic stress-induced programmed cell death in plants. *Front. Plant Sci.* 6:69. doi: 10.3389/fpls.2015.00069
- Plieth, C., Hansen, U. P., Knight, H., and Knight, M. R. (1999). Temperature sensing by plants: the primary characteristics of signal perception and calcium response. *Plant J.* 18, 491–497. doi: 10.1046/j.1365-313X.1999.00471.x
- Pottosin, I., Maria Velarde-Buendia, A., Bose, J., Fuglsang, A. T., and Shabala, S. (2014). Polyamines cause plasma membrane depolarization, activate Ca²⁺-, and modulate H⁺-ATPase pump activity in pea roots. *J. Expt. Bot.* 65, 2463–2472. doi: 10.1093/jxb/eru133
- Probst, A. V., and Scheid, O. M. (2015). Stress-induced structural changes in plant chromatin. *Curr. Opin. Plant Biol.* 27, 8–16. doi: 10.1016/j.pbi.2015.05.011
- Qin, W. S., Pappan, K., and Wang, X. M. (1997). Molecular heterogeneity of phospholipase D (PLD) – Cloning of PLD gamma and regulation of plant PLD gamma, -beta, and -alpha by polyphosphoinositides and calcium. *J. Biol. Chem.* 272, 28267–28273. doi: 10.1074/jbc.272.45.28267
- Qiu, Q. S., Guo, Y., Quintero, F. J., Pardo, J. M., Schumaker, K. S., and Zhu, J. K. (2004). Regulation of vacuolar Na⁺/H⁺ exchange in *Arabidopsis thaliana* by the salt-overly-sensitive (SOS) pathway. *J. Biol. Chem.* 279, 207–215. doi: 10.1074/jbc.M307982200
- Quiles-Pando, C., Rexach, J., Navarro-Gochicoa, M. T., Camacho-Cristóbal, J. J., Herrera-Rodríguez, M. B., and González-Fontes, A. (2013). Boron deficiency increases the level of cytosolic Ca²⁺ and expression of Ca²⁺-related genes in *Arabidopsis thaliana* roots. *Plant Physiol. Biochem.* 65, 55–60. doi: 10.1016/j.plaphy.2013.01.004
- Quintero, F. J., Martinez-Atienza, J., Villalta, I., Jiang, X., Kim, W. Y., Ali, Z., et al. (2011). Activation of the plasma membrane Na/H antiporter Salt-Overly-Sensitive 1 (SOS1) by phosphorylation of an auto-inhibitory C-terminal domain. *Proc. Natl. Acad. Sci. U.S.A.* 108, 2611–2616. doi: 10.1073/pnas.1018921108
- Ragel, P., Rodenas, R., Garcia-Martin, E., Andres, Z., Villalta, I., Nieves-Cordones, M., et al. (2015). SOS the CBL-interacting protein kinase CIPK23 regulates HAK5-mediated high-affinity K⁺ uptake in *Arabidopsis* roots. *Plant Physiol.* 169, 2863–2873. doi: 10.1104/pp.15.01401
- Richards, S. L., Wilkins, K. A., Swarbreck, S. M., Anderson, A., Habib, N., McAinsh, M., et al. (2015). The hydroxyl radical; from seed to seed. *J. Expt. Bot.* 66, 37–46. doi: 10.1093/jxb/eru398
- Richter, G. L., Monshausen, G. B., Krol, A., and Gilroy, S. (2009). Mechanical stimuli modulate lateral root organogenesis. *Plant Physiol.* 151, 1855–1866. doi: 10.1104/pp.109.142448
- Rivera, E., Alvarez, J. M., Vidal, E. A., Oses, C., Vega, A., and Gutierrez, R. A. (2015). The calcium ion is a second messenger in the nitrate signaling pathway of *Arabidopsis*. *Plant Physiol.* 169, 1397–1404. doi: 10.1104/pp.15.00961
- Rodrigo-Moreno, A., Andres-Colas, N., Poschenrieder, C., Gunse, B., Penarrubia, L., and Shabala, S. (2013). Calcium- and potassium-permeable plasma membrane transporters are activated by copper in *Arabidopsis* root tips: linking copper transport with cytosolic hydroxyl radical production. *Plant Cell Environ.* 36, 844–855. doi: 10.1111/pce.12020
- Rodriguez-Hernandez, M. C., Bonifas, I., Alfaro-De la Torre, M. C., Flores-Flores, J. I., Bañuelos-Hernández, B., and Patiño-Rodríguez, O. (2015). Increased accumulation of cadmium and lead under Ca and Fe deficiency in *Typha latifolia*: a case study of two pore channel (TPC1) gene. *Environ. Exp. Bot.* 115, 38–48. doi: 10.1016/j.envexpbot.2015.02.009
- Roy, S. J., Gilligham, M., Berger, B., Essah, P. A., Cheffings, C., Miller, A. J., et al. (2008). Investigating glutamate receptor-like gene co-expression in *Arabidopsis thaliana*. *Plant Cell Environ.* 31, 861–871. doi: 10.1111/j.1365-3040.2008.01801.x
- Ruberti, C., Kim, S. J., Stefano, G., and Brandizzi, F. (2015). Unfolded protein responses in plants: one master, many questions. *Curr. Opin. Plant Biol.* 27, 59–66. doi: 10.1016/j.pbi.2015.05.016
- Ruelland, E., Kravets, V., Derevyanchuk, M., Martinec, J., Zachowski, A., and Pokotylo, I. (2015). Role of phospholipid signalling in plant environmental responses. *Environ. Exp. Bot.* 114, 129143. doi: 10.1016/j.envexpbot.2014.08.009
- Sagor, G. H. M., Chawla, P., Kim, D. W., Berberich, T., Kojima, S., Niitsu, M., et al. (2015). The polyamine spermine induces the unfolded protein response via the MAPK cascade in *Arabidopsis*. *Front. Plant Sci.* 6:687. doi: 10.3389/fpls.2015.00687
- Sani, E., Herzyk, P., Perrella, G., Colot, V., and Amtmann, A. (2013). Hyperosmotic priming of *Arabidopsis* seedlings establishes a long-term somatic memory accompanied by specific changes of the epigenome. *Genome Biol.* 14, R59. doi: 10.1186/gb-2013-14-6-r59
- Sarwat, M., and Naqvi, A. R. (2013). Heterologous expression of rice calnexin (OsCNX) confers drought tolerance. *Mol. Biol. Rep.* 40, 5451–5464. doi: 10.1007/s11033-013-2643-y

- Shabala, S., and Pottosin, I. (2014). Regulation of potassium transport in plants under hostile conditions: implications for abiotic and biotic stress tolerance. *Physiol. Plant* 151, 257–279. doi: 10.1111/pp.12165
- Shabala, S., Shabala, L., Barcelo, J., and Poschenrieder, C. (2014). Membrane transporters mediating root signalling and adaptive responses to oxygen deprivation and soil flooding. *Plant Cell Environ.* 37, 2216–2233. doi: 10.1111/pce.12339
- Shabala, S., Wu, H., and Bose, J. (2015). Salt stress sensing and early signalling in plant roots: current knowledge and hypothesis. *Plant Sci.* 241, 109–119. doi: 10.1016/j.plantsci.2015.10.003
- Shi, Y. T., Ding, Y. L., and Yang, S. H. (2015). Cold signal transduction and its interplay with phytohormones during cold acclimation. *Plant Cell Physiol.* 56, 7–15. doi: 10.1093/pcp/pcu115
- Shih, H. W., Depew, C. L., Miller, N. D., and Monshausen, G. B. (2015). The cyclic nucleotide-gated channel CNGC14 regulates root gravitropism in *Arabidopsis thaliana*. *Curr. Biol.* 25, 1–7. doi: 10.1016/j.cub.2015.10.025
- Shih, H.-W., Miller, N. D., Dai, C., Spalding, E. P., and Monshausen, G. B. (2014). The receptor-like kinase FERONIA is required for mechanical signal transduction in *Arabidopsis* seedlings. *Curr. Biol.* 24, 1887–1892. doi: 10.1016/j.cub.2014.06.064
- Shin, R., and Schachtman, D. P. (2004). Hydrogen peroxide mediates plant root cell response to nutrient deprivation. *Proc. Natl. Acad. Sci. U.S.A.* 101, 8827–8832. doi: 10.1073/pnas.0401707101
- Shorrocks, V. M. (1997). The occurrence and correction of boron deficiency. *Plant Soil* 193, 121–148. doi: 10.1023/A:1004216126069
- Singh, S., Parihar, P., Singh, R., Singh, V. P., and Prasad, S. M. (2016). Heavy metal tolerance in plants: role of transcriptomics, proteomics, metabolomics, and ionomics. *Front. Plant Sci.* 6:1143. doi: 10.3389/fpls.2015.01143
- Stael, S., Wurzing, B., Mair, A., Mehler, N., Vothknecht, U. C., and Teige, M. (2012). Plant organellar calcium signalling: an emerging field. *J. Expt. Bot.* 63, 1525–1542. doi: 10.1093/jxb/err394
- Straltsova, D., Chykun, P., Subramanian, S., Sosan, A., Kolbanov, D., Sokolik, A., et al. (2015). cation channels are involved in brassinosteroid signalling in higher plants. *Steroids* 97, 98–106. doi: 10.1016/j.steroids.2014.10.008
- Sulaiman, Y., Knight, M. R., and Katakya, R. (2012). Non-invasive monitoring of temperature stress in *Arabidopsis thaliana* roots, using ion amperometry. *Anal. Meth.* 4, 1656–1661. doi: 10.1039/c2ay05747f
- Sunkar, R., Kaplan, B., Bouche, N., Arazi, T., Dolev, D., Talke, I. N., et al. (2000). Expression of a truncated tobacco NtCBP4 channel in transgenic plants and disruption of the homologous *Arabidopsis* CNGC1 gene confer Pb²⁺ tolerance. *Plant J.* 24, 533–542. doi: 10.1046/j.1365-313x.2000.00901.x
- Swarbreck, S. M., Colaço, R., and Davies, J. M. (2013). Update on plant calcium-permeable channels. *Plant Physiol.* 163, 1514–1522. doi: 10.1104/pp.113.220855
- Tai, F., Yuan, Z., Li, S. P., Wang, Q., Liu, F., and Wang, W. (2016). ZmCIPK8, a CBL-interacting protein kinase, regulates maize response to drought stress. *Plant Cell Tiss. Org. Cul.* 124, 459–469. doi: 10.1007/s11240-015-0906-0
- Teardo, E., Carraretto, L., De Bortoli, S., Costa, A., Behera, S., and Wagner, R. (2015). Alternative splicing-mediated targeting of the *Arabidopsis* GLUTAMATE RECEPTOR3.5 to mitochondria affects organelle morphology. *Plant Physiol.* 167, 216–227. doi: 10.1104/pp.114.242602
- Thion, L., Mazars, C., Nacry, P., Bouchez, D., Moreau, M., Ranjeva, R., et al. (1998). Plasma membrane depolarization-activated calcium channels, stimulated by microtubule-depolymerizing drugs in wild-type *Arabidopsis thaliana* protoplasts, display constitutively large activities and a longer half-life in ton 2 mutant cells affected in the organization of cortical microtubules. *Plant J.* 13, 603–610.
- Thoday-Kennedy, E. L., Jacobs, A. J., and Roy, S. J. (2015). The role of the CBL-CIPK calcium signalling network in regulating ion transport in response to abiotic stress. *Plant Growth Regul.* 76, 3–12. doi: 10.1007/s10725-015-0034-1
- Toyota, M., and Gilroy, S. (2013). Gravitropism and mechanical signalling in plants. *Am. J. Bot.* 100, 111–125. doi: 10.3732/ajb.1200408
- Tracy, F. E., Gilliam, M., Dodd, A. N., Webb, A. A. R., and Tester, M. (2008). NaCl-induced changes in cytosolic free Ca²⁺ in *Arabidopsis thaliana* are heterogeneous and modified by external ionic composition. *Plant Cell Environ.* 31, 1063–1073. doi: 10.1111/j.1365-3040.2008.01817.x
- Véry, A. A., and Davies, J. M. (2000). Hyperpolarization-activated calcium channels at the tip of *Arabidopsis* root hairs. *Proc. Natl. Acad. Sci. U.S.A.* 97, 9801–9806. doi: 10.1073/pnas.160250397
- Villiers, F., Jourdain, A., Bastien, O., Leonhardt, N., Fujioka, S., Tichtinck, G., et al. (2012). Evidence for functional interaction between brassinosteroids and cadmium response in *Arabidopsis thaliana*. *J. Expt. Bot.* 63, 1185–1200. doi: 10.1093/jxb/err335
- Virdi, A. S., Singh, S., and Singh, P. (2015). Abiotic stress responses in plants: roles of calmodulin-regulated proteins. *Front. Plant Sci.* 6:809. doi: 10.3389/fpls.2015.00809
- Virolainen, E., Blokhina, O., and Fagerstedt, K. (2002). Ca²⁺-induced high amplitude swelling and cytochrome c release from wheat (*Triticum aestivum* L.) mitochondria under anoxic stress. *Ann. Bot.* 90, 509–516. doi: 10.1093/aob/mcf221
- Wagner, S., Behera, S., De Bortoli, S., Logan, D. C., Fuchs, P., and Carraretto, L. (2015). The EF-Hand Ca²⁺-binding protein MICU choreographs mitochondrial Ca²⁺ dynamics in *Arabidopsis*. *Plant Cell* 27, 3190–3212. doi: 10.1105/tpc.15.00509
- Wan, S., and Jiang, L. (2016). Endoplasmic reticulum (ER) stress and the unfolded protein response (UPR) in plants. *Protoplasma* 273, 753–764. doi: 10.1007/s00709-015-0842-1
- Wang, F. F., Chen, Z. H., Liu, X. H., Colmer, T. D., Zhou, M. X., and Shabala, S. (2016). Tissue-specific root ion profiling reveals essential roles of the CAX and ACA calcium transport systems in response to hypoxia in *Arabidopsis*. *J. Expt. Bot.* 67, 3747–3762. doi: 10.1093/jxb/erw034
- Wang, P., Li, Z. W., Wei, J. S., Zhao, Z. L., Sun, D. Y., and Cui, S. J. (2012). A Na⁺/Ca²⁺ exchanger-like protein (AtNCL) involved in salt stress in *Arabidopsis*. *J. Biol. Chem.* 287, 44062–44070. doi: 10.1074/jbc.M112.351643
- Weiland, M., Mancuso, S., and Baluska, F. (2016). Signalling via glutamate and GLRs in *Arabidopsis thaliana*. *Funct. Plant Biol.* 43, 1–25.
- Whalley, H. J., and Knight, M. R. (2013). Calcium signatures are decoded by plants to give specific gene responses. *New Phytol.* 197, 690–693. doi: 10.1111/nph.12087
- Whalley, H. J., Sargeant, A. W., Steele, J. F., Lacoere, T., Lamb, R., Saunders, N. J., et al. (2011). Transcriptomic analysis reveals calcium regulation of specific promoter motifs in *Arabidopsis*. *Plant Cell* 23, 4079–4095. doi: 10.1105/tpc.111.090480
- White, P. J. (2009). Depolarization-activated calcium channels shape the calcium signatures induced by low-temperature stress. *New Phytol.* 183, 7–8. doi: 10.1111/j.1469-8137.2009.02857.x
- Wigge, P. A. (2013). Ambient temperature signalling in plants. *Curr. Opin. Plant Biol.* 16, 661–666. doi: 10.1016/j.pbi.2013.08.004
- Wu, J. Y., Jin, C., Qu, H. Y., Tao, S. T., Xu, G. H., Wu, J., et al. (2012). Low temperature inhibits pollen viability by alteration of actin cytoskeleton and regulation of pollen plasma membrane ion channels in *Pyrus pyrifolia*. *Environ. Expt. Bot.* 78, 70–75. doi: 10.1016/j.envexpbot.2011.12.021
- Xu, Q. T., Fan, H. Y., Jiang, Z., Zhou, Z. Q., Yang, L., Mei, F. Z., et al. (2013). Cell wall degradation and the dynamic changes of Ca²⁺ and related enzymes in the developing aerenchyma of wheat (*Triticum aestivum* L.) under waterlogging. *Acta Biol. Hung.* 64, 328–340. doi: 10.1556/ABiol.64.2013.3.6
- Yamada, N., Theerawitaya, C., Ca-Um, S., Kirdmanee, C., and Takabe, T. (2014). Expression and functional analysis of putative vacuolar Ca²⁺-transporters (CAXs and ACAs) in roots of salt tolerant and sensitive rice cultivars. *Protoplasma* 251, 1067–1075. doi: 10.1007/s00709-014-0615-2
- Yamauchi, T., Watanabe, K., Fukazawa, A., Mori, H., Abe, F., Kawaguchi, K., et al. (2014). Ethylene and reactive oxygen species are involved in root aerenchyma formation and adaptation of wheat seedlings to oxygen-deficient conditions. *J. Expt. Bot.* 65, 261–273. doi: 10.1093/jxb/ert371
- Yang, C.-Y., and Hong, C.-P. (2015). The NADPH oxidase RbohD is involved in primary hypoxia signalling and modulates expression of hypoxia-inducible genes under hypoxic stress. *Env. Expt. Bot.* 115, 63–72. doi: 10.1016/j.envexpbot.2015.02.008
- Yang, T. B., Chaudhuri, S., Yang, L. H., Du, L. Q., and Poovaiah, B. W. (2010). A calcium/calmodulin-regulated member of the receptor-like kinase family confers cold tolerance in plants. *J. Biol. Chem.* 285, 7119–7126. doi: 10.1074/jbc.M109.035659
- Yeh, C. M., Chien, P. S., and Huang, H. J. (2007). Distinct signalling pathways for induction of MAP kinase activities by cadmium and copper in rice roots. *J. Expt. Bot.* 58, 659–671. doi: 10.1093/jxb/erl240
- Yemelyanov, V. V., Shishova, M. F., Chirkova, T. V., and Lindberg, S. M. (2011). Anoxia-induced elevation of cytosolic Ca²⁺ concentration depends on

- different Ca^{2+} sources in rice and wheat protoplasts. *Planta* 234, 271–280. doi: 10.1007/s00425-011-1396-x
- Yu, H. Q., Yong, T. M., Li, H. J., Liu, Y. P., Zhou, S. F., Fu, F. L., et al. (2015). Overexpression of a phospholipase D alpha gene from *Ammopiptanthus nanus* enhances salt tolerance of phospholipase D alpha 1-deficient *Arabidopsis* mutant. *Planta* 242, 1495–1509. doi: 10.1007/s00425-015-2390-5
- Yu, L. J., Nie, J. N., Cao, C. Y., Jin, Y. K., Yan, M., Wang, F. Z., et al. (2010). Phosphatidic acid mediates salt stress response by regulation of MPK6 in *Arabidopsis thaliana*. *New Phytol.* 188, 762–773. doi: 10.1111/j.1469-8137.2010.03422.x
- Yu, Y. Q., and Assmann, S. M. (2015). The heterotrimeric G-protein beta subunit, AGB1, plays multiple roles in the *Arabidopsis* salinity response. *Plant Cell Environ.* 38, 2143–2156. doi: 10.1111/pce.12542
- Yuan, F., Yang, H., Xue, Y., Kong, D., Ye, R., Li, C., et al. (2014). OSCA1 mediates osmotic-stress-evoked Ca^{2+} increases vital for osmosensing in *Arabidopsis*. *Nature* 514, 367–371. doi: 10.1038/nature13593
- Yuan, H.-M., and Huang, X. (2016). Inhibition of root meristem growth by cadmium involves nitric oxide-mediated repression of auxin accumulation and signalling in *Arabidopsis*. *Plant Cell Environ.* 39, 120–135. doi: 10.1111/pce.12597
- Yuen, C. C. Y., and Christopher, D. A. (2013). The group IV-A cyclic nucleotide-gated channels, CNGC19 and CNGC 20, localize to the vacuole membrane in *Arabidopsis thaliana*. *AoB Plants* 5, plt012. doi: 10.1186/1471-2164-15-853
- Zelinova, V., Alemayehu, A., Bocova, B., Huttova, J., and Tamas, L. (2015). Cadmium-induced reactive oxygen species generation, changes in morphogenic responses and activity of some enzymes in barley root tip are regulated by auxin. *Biologia* 70, 356–364. doi: 10.1515/biolog-2015-0035
- Zeng, F., Konnerup, D., Shabala, L., Zhou, M., Colmer, T. D., Zhang, G., et al. (2014). Linking oxygen availability with membrane potential maintenance and K^+ retention of barley roots: implications for waterlogging stress tolerance. *Plant Cell Environ.* 37, 2325–2338. doi: 10.1111/pce.12422
- Zeng, Q., Ling, Q., Fan, L., Li, Y., Hu, F., Chen, J., et al. (2015). Transcriptome profiling of sugarcane roots in response to potassium stress. *PLoS ONE* 10:e0126306. doi: 10.1371/journal.pone.0126306
- Zepeda-Jazo, I., Maria Velarde-Buendia, A., Enriquez-Figueroa, R., Bose, J., Shabala, S., Muniz-Murguía, J., et al. (2011). Polyamines interact with hydroxyl radicals in activating Ca^{2+} and K^+ transport across the root epidermal plasma membranes. *Plant Physiol.* 157, 2167–2180. doi: 10.1016/j.plaphy.2012.09.002
- Zhang, L., Chen, Z., and Zhu, C. (2012). Endogenous nitric oxide mediates alleviation of cadmium toxicity induced by calcium in rice seedlings. *J. Environ. Sci.* 24, 940–948. doi: 10.1016/S1001-0742(11)60978-9
- Zhang, X., Shen, Z. D., Sun, J., Yu, Y. C., Deng, S. R., and Li, Z. Y. (2015). NaCl-elicited, vacuolar Ca^{2+} release facilitates prolonged cytosolic Ca^{2+} signaling in the salt response of *Populus euphratica* cells. *Cell Calcium* 57, 348–365. doi: 10.1016/j.ceca.2015.03.001
- Zhang, Y., Wang, Y., Taylor, J. L., Jiang, Z., Zhang, S., Mei, F., et al. (2015). Aequorin-based luminescence imaging reveals differential calcium signalling responses to salt and reactive oxygen species in rice roots. *J. Expt. Bot.* 66, 2535–2545. doi: 10.1093/jxb/erv043
- Zhao, M. G., Tian, Q. Y., and Zhang, W. H. (2007). Ethylene activates a plasma membrane Ca^{2+} -permeable channel in tobacco suspension cells. *New Phytol.* 174, 507–515. doi: 10.1111/j.1469-8137.2007.02037.x
- Zhao, R., Sun, H. M., Zhao, N., Jing, X. S., Shen, X., and Chen, S. (2015). The *Arabidopsis* Ca^{2+} -dependent protein kinase CPK27 is required for plant response to salt-stress. *Gene* 563, 203–214. doi: 10.1016/j.gene.2015.03.024
- Zhao, Y., Pan, Z., Zhang, Y., Qu, X. L., Zhang, Y. G., and Yang, Y. G. (2013). The Actin-Related Protein2/3 complex regulates mitochondrial-associated calcium signaling during salt stress in *Arabidopsis*. *Plant Cell* 25, 4544–4559. doi: 10.1105/tpc.113.117887
- Zhu, S.-Y., Yu, X.-C., Wang, X.-J., Zhao, R., Li, Y., Fan, R.-C., et al. (2007). Two calcium-dependent protein kinases, CPK4 and CPK11, regulate abscisic acid signal transduction in *Arabidopsis*. *Plant Cell* 19, 3019–3036. doi: 10.1105/tpc.107.050666
- Zhu, X. H., Feng, Y., Liang, G. M., Liu, N., and Zhu, J. K. (2013). Aequorin-based luminescence imaging reveals stimulus- and tissue-specific Ca^{2+} dynamics in *Arabidopsis* plants. *Mol. Plant* 6, 444–455. doi: 10.1093/mp/sst013

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Wilkins, Matthus, Swarbreck and Davies. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.