

Article

Calculated Third Order Rate Constants for Interpreting the Mechanisms of Hydrolyses of Chloroformates, Carboxylic Acid Halides, Sulfonyl Chlorides and Phosphorochloridates

T. William Bentley

Chemistry Unit, Grove Building, School of Medicine, Swansea University,
Swansea SA2 8PP, Wales, UK; E-Mail: t.w.bentley@swansea.ac.uk;
Tel.: +44-1792-295-262; Fax: +44-1792-295-554

Academic Editor: Malcolm D'Souza

Received: 7 April 2015 / Accepted: 30 April 2015 / Published: 8 May 2015

Abstract: Hydrolyses of acid derivatives (e.g., carboxylic acid chlorides and fluorides, fluoro- and chloroformates, sulfonyl chlorides, phosphorochloridates, anhydrides) exhibit pseudo-first order kinetics. Reaction mechanisms vary from those involving a cationic intermediate (S_N1) to concerted S_N2 processes, and further to third order reactions, in which one solvent molecule acts as the attacking nucleophile and a second molecule acts as a general base catalyst. A unified framework is discussed, in which there are two reaction channels—an S_N1 - S_N2 spectrum and an S_N2 - S_N3 spectrum. Third order rate constants (k_3) are calculated for solvolytic reactions in a wide range of compositions of acetone-water mixtures, and are shown to be either approximately constant or correlated with the Grunwald-Winstein Y parameter. These data and kinetic solvent isotope effects, provide the experimental evidence for the S_N2 - S_N3 spectrum (e.g., for chloro- and fluoroformates, chloroacetyl chloride, *p*-nitrobenzoyl *p*-toluenesulfonate, sulfonyl chlorides). Deviations from linearity lead to U- or V-shaped plots, which assist in the identification of the point at which the reaction channel changes from S_N2 - S_N3 to S_N1 - S_N2 (e.g., for benzoyl chloride).

Keywords: solvolysis; substituent effects; solvent effects; acid chlorides; reaction mechanisms

1. Introduction

Solvolysees involve reactions of a solvent nucleophile (e.g., water or alcohol) with a suitable substrate (e.g., of general formula RX , where R is an alkyl residue and X is an electronegative group) [1]. To aid

the solubility of RX in water, an organic cosolvent (e.g., acetone) is often added. An understanding of the factors influencing the reactivity of RX can be gained by studying the dependence of logarithms of observed pseudo-first order rate constant ($\log k_{\text{obs}}$) on empirical parameters for substituent and/or solvent effects [2]. These are examples of linear free energy relationships (LFER), from which reaction mechanisms can be proposed and effects of changing reaction conditions (e.g., cosolvents) could be predicted [3,4].

A useful concept is the idea that solvolyses proceed via a spectrum of reaction mechanisms. An S_N1 - S_N2 spectrum of mechanisms is well established for many solvolyses (e.g., solvolyses of secondary alkyl sulfonates [5]). Density functional (DFT) calculations for benzoyl chlorides [6] were consistent with concerted S_N2 reactions, with “loose” (cationic) transition states for electron-donating substituents and “tight” (associative) transition states for electron withdrawing substituents (Figure 1). Very recent DFT calculations [7] further illustrated the S_N1 - S_N2 spectrum (e.g., for benzyl halides), but also provided support for the proposal [8] that there was also an S_N2 - S_N3 spectrum of mechanisms for solvolyses of aromatic sulfonyl chlorides (**2**).




Figure 1. Possible reaction mechanisms for solvolyses of *p*-Z-substituted benzoyl chlorides (**1**): (a) The S_N1 - S_N2 (cationic) reaction channel in which one or two molecules undergo covalency change to varying extents; (b) The S_N2 - S_N3 third order (addition-elimination) reaction channel in which three molecules undergo covalency change to varying extents.

Electrophilic solvation of the developing chloride anion and general solvation are excluded from the assessment of molecularity, and are not shown in Figure 1. At the S_N1 extreme of the S_N1 - S_N2 spectrum, there may be evidence for a cationic intermediate and reactions are unimolecular (e.g., solvolyses of *p*-dimethylaminobenzoyl fluoride (**3**, $Z = \text{NMe}_2$) show common ion rate depression [9]).

The key feature of the S_N3 mechanism is that the reactions are termolecular: a solvent nucleophile attacks the substrate (e.g., RX or *p*-Z-substituted benzoyl chlorides as in Figure 1b), and a second solvent molecule assists by partially deprotonating the nucleophile (three molecules undergo covalency change, but they could be preassembled by hydrogen bonding).

Although the kinetic order is usually first order (rate constant k_{obs}), it has proved useful in some cases to calculate various third order rate constants (k_3); as shown below, values of $\log k_3$ may be

approximately constant or may give good linear plots *vs.* solvent polarity parameters for solvolyses of acid derivatives. Furthermore, deviations from linearity provide some of the clearest evidence for the relatively subtle change in mechanism from the end of the S_N1-S_N2 spectrum to the beginning of the S_N2-S_N3 spectrum.

2. Results and Discussion

Initial results were obtained for the range of substrates shown in Figure 2.




Figure 2. The substrates (1–7) are: named as follows: (1) *p*-Z-substituted benzoyl chlorides; (2) *p*-Z-substituted sulfonyl chlorides; (3) *p*-Z-substituted benzoyl fluorides; (4) *p*-Z-substituted benzoyl tosylates, where tosylate (OTs) is *p*-toluenesulfonate); (5) alkyl chloroformates; (6) *p*-Z-substituted phenyl chloroformates; (7) alkyl fluoroformates.

2.1. Choice of Equation for Correlating Solvolysis Rates

The simplest (but not reliable) equation for obtaining the kinetic order of a pseudo-first order solvolysis is Equation (1), where *n* is the kinetic order, *k_n* is the *n*th order rate constant and [water] refers to the molar concentration of water in for example a binary mixture such as acetone-water. Taking logarithms, a plot of log *k_{obs}* *vs.* log [water] would have a slope of *n* (Equation (2)). To illustrate the lack of reliability of Equation (2), two examples using kinetic data from the recent literature (but not the published interpretation of the data) are: (a) solvolyses of benzoyl fluoride (3, Z = H) in acetone–water, which would have a kinetic order of 3.3 (kinetic data from Reference [10]); (b) solvolyses of *p*-nitrobenzoyl tosylate (4, Z = NO₂), which would have a kinetic order of 0.3 (kinetic data from Reference [11]).

$$k_{\text{obs}} = k_n[\text{water}]^n \quad (1)$$

$$\log k_{\text{obs}} = n \times \log [\text{water}] + \log k_n \quad (2)$$

An alternative, more credible explanation is that solvolyses of both 3, Z = H and 4, Z = OTs are third order. Solvolyses of 3, Z = H in water and D₂O give a kinetic solvent isotope effect (KSIE) of 2.0 [9], and the KSIE for 4, Z = OTs in methanol and MeOD varies from 1.84 at –10 °C to 1.59 at 25 °C [11]. These values are consistent with some cleavage of O-H or O-D bonds in the rate determining step, as shown in Figure 1b. Various reactions of acid derivatives including esters are thought to react by such general base-catalysed processes [12].

There are many publications in which the KSIE for MeOH and MeOD were obtained (see below), and a wide range of KSIE data have been published by Koo *et al.* [13,14]; two of these were included in the recent DFT calculations [7]: values of KSIE in water are: for 2, Z = OMe (calc. 1.64, expt. 1.37);

for **2**, $Z = \text{NO}_2$ (calc. 1.97, expt. 1.76). Both $\text{S}_{\text{N}}1$ [15] and $\text{S}_{\text{N}}2$ [16] reactions typically have KSIEs < 1.3 (see also Table 5 of Reference [8]), so hydrolyses of **2**, $Z = \text{OMe}$ are close to those expected for $\text{S}_{\text{N}}2$ reactions, but those for **2**, $Z = \text{NO}_2$ are significantly higher.

Calculated third order rate constants Equation (3) give satisfactory plots vs. the Grunwald-Winstein (GW) parameter (Y) for solvent ionizing power [4,17,18] (a measure of solvent polarity). The plots (Figure 3) show that calculated third order rate constants for solvolyses of benzoyl fluoride (**3**, $Z = \text{H}$) increase as Y increases ($\text{slope} = 0.27 \pm 0.02$, $R^2 = 0.979$, std. error = 0.09, $n = 9$), but those for p -nitrobenzoyl tosylate (**4**, $Z = \text{NO}_2$) decrease ($\text{slope} = -0.40 \pm 0.02$, $R^2 = 0.994$, std. error = 0.06, $n = 7$).

$$\log k_3 = \log \{k_{\text{obs}}/[\text{water}]^2\} = \log k_{\text{obs}} - 2 \log [\text{water}] = \text{slope} \times Y + \text{intercept} \quad (3)$$




Figure 3. Plots of logarithms of third order rate constants (Equation (3)) vs. Grunwald-Winstein Y values [4,17] for acetone-water mixtures; kinetic data for benzoyl fluoride (**3**, $Z = \text{H}$) at 25 °C are from Reference [10]; kinetic data for p -nitrobenzoyl tosylate (**4**, $Z = \text{NO}_2$) at -10 °C are from Reference [11].

It should not be inferred that the linearity of the plots (Figure 3) is by itself evidence for third order reactions, because a plot (not shown) of $\log [\text{water}]^2$ vs. Y is close to linear, but curves downwards for 80% and 90% acetone-water; this curvature can lead to a “kink” in Equation (3) plots. Consequently, the main difference between Equation (3) and a GW plot of $\log k_{\text{obs}}$ vs. Y is that the slopes of Equation (3) are lower, but in some cases (see below) mechanistic changes are indicated much more clearly using Equation (3); note however that for bimolecular reactions in 80% and 90% acetone-water, there may be a change in slope which is not due to a mechanistic change.

As with other mechanistic indicators, it is always desirable to provide independent evidence, and KSIE data will be quoted extensively during the following discussion. Multi-parameter equations provide a more comprehensive account of solvent effects, and these are discussed in Section 2.4.

Also, there are various examples of chloroformates (**5,6**) where the slope based on Equation (3) is close to zero, implying that a plot of $\log k_{\text{obs}}$ vs. $\log [\text{H}_2\text{O}]$ (Equation (2)) would give a value of n (the kinetic order with respect to water) of close to 2 (e.g., see Section 2.2.1).

An important advantage of Equation (3) over Equation (2) is that trends in slopes can be interpreted with the benefit of long experience of applications of the original GW equation [18]. The selection of Y rather

than one of the other Y_X parameters [19] for solvent ionizing power ensures that none of the Y values are based on extrapolated data, and that slopes for all substrates are comparable (for further discussion on this choice, see Reference [20]). For many aqueous binary mixtures (including acetone-water), to a good approximation $Y = Y_{Cl} \times 0.75$ (see Table 8 of Reference [19]). The factors influencing the slopes of correlations based on Equation (3) were investigated for a wide range of substrates.

2.2. Evidence for the Third Order (Addition-Elimination) Reaction Channel from Correlations Using Equation (3) and Kinetic Solvent Isotope Effects

2.2.1. Chloroformates

In general, chloroformates (**5,6**) are much more likely than the corresponding carboxylic acid chlorides (**2**) to react via higher order (associative) reactions. Recent G3 calculations of gas phase heterolytic bond dissociation energies (HBDE) for the C–Cl bonds show for example that PhCOCl (**1**, $Z = H$) has an HBDE of 150.1 kcal/mol whereas the corresponding chloroformate (**6**, $Z = H$) has a much higher HBDE of 166.5 kcal/mol [21]. Consequently, the electrophilic “pull” is less for chloroformates, and a greater nucleophilic “push” might be expected [22].

Inspection of each column of Table 1, shows that logarithms of the calculated third order rate constants ($\log k_3$) are almost independent of solvent composition for solvolyses ethyl chloroformate (**5**, $R = Et$), phenyl- and substituted phenyl chloroformates (**6**, $Z = H, Me, OMe$), and trichloroethyl chloroformate (**5**, $R = CH_2CCl_3$) in a wide range of acetone-water mixtures. Equation (3) gives slightly negative slopes for solvolyses of: (**5**, $R = C(Me)_2CCl_3$) for which $slope = -0.26 \pm 0.02$, $R^2 = 0.969$, std. error = 0.08; **6**, $Z = Cl$ ($slope = -0.08 \pm 0.01$, $R^2 = 0.946$, std. error = 0.03) and **6**, $Z = NO_2$ ($slope = -0.19 \pm 0.01$, $R^2 = 0.994$, std. error = 0.02). All three are plots may alternatively be interpreted as slightly curved (Figure 4).

Table 1. Approximately constant values of logarithms of calculated third order rate constants (Equation (3)) for solvolyses of chloroformates in acetone-water.

Acetone (% v/v)	5 , $R = Et^a$ 24.2 °C	6 , $Z = H^b$ 25 °C	6 , $Z = Me^b$ 25 °C	6 , $Z = OMe^b$ 25 °C	5 , $R = CH_2CCl_3$ Reference ^c , 35 °C
95	−7.01				
90	−7.01	−5.11 ^d			−4.62
80	−7.11	−5.20	−5.43	−5.47	−4.82
70		−5.35	−5.54	−5.55	−4.83
60	−7.13	−5.40	−5.59	−5.60	−4.86
50		−5.43	−5.59	−5.58	−4.82
40	−7.06	−5.41	−5.56	−5.53	−4.78
30		−5.36	−5.51	−5.47	−4.67
20	−6.96	−5.34	−5.46	−5.43	−4.62
10		−5.34	−5.45	−5.39	
water	−6.96	−5.37	−5.46	−5.55	

^a Data from Reference [23]; ^b Data from Reference [14], unless stated otherwise; ^c Data from Reference [24];

^d Data from Reference [25].




Figure 4. Plots of logarithms of third order rate constants (Equation (3)) vs. Grunwald-Winstein Y values [4,17] for acetone-water mixtures; kinetic data: for **5**, $R = C(\text{Me})_2\text{CCl}_3$ at 25 °C from Reference [26]; for *p*-chlorophenyl chloroformate (**6**, $Z = \text{Cl}$) at 25 °C from Reference [14]; for *p*-nitrophenyl chloroformate (**6**, $Z = \text{NO}_2$) at 25 °C from Reference [27].

Data for solvolyses of many other chloroformates [28] in a limited range of acetone-water solvent compositions, usually within the range 90%–60% show the same trends as those discussed above. Exceptions to these trends indicate the possibility of mechanistic changes, as discussed in Section 2.3.

2.2.2. Fluoroformates (**7**)

One of the first extensive study of solvent effects on these solvolyses was for *n*-octyl fluoroformate (**7**, $R = n\text{-octyl}$); data in 95%, 90%, 80% and 60% *v/v* acetone-water [29] give a positive *slope* based on Equation (3) of 0.22 ± 0.02 . Positive *slopes* are observed for many other fluoroformates, as well as for benzoyl fluoride (**3**, $Z = \text{H}$)—see Figure 3. Details are not given here because of the limited range of acetone-water solvent compositions (usually between 90% and 60% *v/v* [28]). Instead, typical studies involve a much wider range of pure solvents and binary solvent mixtures (Section 2.4), and mechanistic assignments are supported by studies of KSIEs in MeOH [28].

Usually solvolyses of fluoroformates in MeOH give KSIEs >2 , consistent with general base catalysis and the mechanism shown in Figure 1b. However, *t*-butyl fluoroformate (**7**, $R = \text{Bu}^t$) gives a KSIE in MeOH of 1.26 [30], characteristic of the cationic reaction channel (Figure 1a). Also based on Equation (3), the *slope* for **7**, $R = \text{Bu}^t$ is 0.33 ± 0.02 in 90%, 80%, 70% and 60% acetone-water. Consequently, even if a wider range of data was available, changes in mechanism may not lead to convincing changes in slopes for solvolyses of fluoroformates in acetone-water mixtures. In contrast, carboxylic acid and sulfonyl chlorides do show clear changes in slopes (see Section 3).

When a wide range of solvents of varying ionizing power and nucleophilicity are investigated, it is possible to separate the two mechanisms; the extended Grunwald-Winstein equation (Section 2.4) can then be used to correlate the kinetic data and to obtain m values for each mechanism are obtained. It was found that the slopes (m values) for the cationic channel were lower than for the associative

channel; both channels were observed for 1-adamantyl fluoroformates [31], and the ionization channel for *t*-butyl fluoroformate gave an *m* value of 0.41 [30], slightly lower than typical values for the associative channel (see Table 4 of Reference [28]). This is the opposite trend to that observed for chloroformates [32] and other acid chlorides (see below).

2.2.3. Carboxylic Acid Halides

Solvolyses of many carboxylic acid chlorides in a single binary solvent mixture (such as acetone-water) would give linear plots of $\log k_{\text{obs}}$ or $\log k_3$ vs. *Y*. Examples discussed in this section will be restricted to those for which KSIE data indicates the possibility of an S_N3 mechanism.

Good linear correlations of $\log k_3$ vs. *Y* (Equation (3)) for 90%–20% acetone-water have already been published for solvolyses of *p*-nitrobenzoyl chloride (**1**, Z = NO₂) and chloroacetyl chloride (ClCH₂COCl): *slope* = −0.18, see Figure 2 of Reference [33] and Figure 3 of Reference [34] respectively. In each case the KSIE in MeOH is >2. Extrapolation using equation (4) of the 90%–20% acetone plot [33] to pure water gives a predicted value of $k = 0.068 \pm 0.004$, in satisfactory agreement with the experimentally observed values of $k_{\text{obs}} = 0.062$ [9] and 0.056 [35] published subsequently. Also the KSIE in water is approximately 2, although published values vary from 1.75 to 2.3 [35].

$$\log k_3 \text{ (for } \mathbf{1}, Z = \text{NO}_2) = -0.176Y - 4.046 \quad (4)$$

More highly chlorinated substrates such as CHCl₂COCl would also be expected to fit the associative mechanism, but experimental data are limited by their high reactivity; this is overcome in the derivative (**8**, Figure 5), deactivated by the substituents [36]; values of $\log k_3$ are approximately constant from 80% acetone to water (the slope of a plot based on Equation (3) is 0.08 ± 0.03), and also $k_{\text{MeOH}}/k_{\text{MeOD}} = 2.4$ [36].




Figure 5. Structures of additional substrates, named as follows: α -Methoxy- α -(trifluoromethyl) phenylacetylchloride (**8**); *trans*- β -styrenylsulfonyl chloride (**9**); 4-substituted-2,6-dimethylbenzene-sulfonylchlorides (**10**); Phenyl-*N*-phenylphosphoramidochloridate (**11**); 2-chloro-1,3,2-dioxaphospholane-2-oxide (**12**); 2-chloro-5,5-dimethyl-1,3,2-dioxaphosphorinane-2-oxide (**13**).

Solvolyses of **1**, Z = NO₂ from 90% acetone to water [33] and of **8** from 80% acetone to water [36] fit all of the suggested criteria for a third order associative mechanism (*i.e.*, KSIE in MeOH and water >1.5, and a linear plot Equation (3), characteristic of the third order reaction channel (Figure 1b). Also, logarithms of observed rate constants for solvolyses of **1**, Z = NO₂ and methyl chloroformate (MeOCOCl) correlate linearly with those for chloroacetyl chloride (see Figure 1 of Reference [34]), showing strong similarities between solvolyses of chloroformates and other acid chlorides.

Based on the observations [9,10] for benzoyl fluoride (Figure 3), and the reduced “pull” of fluoride as a leaving group, it would be expected that solvolyses of typical acid fluorides would react via the third order mechanism. Supporting evidence includes: (i) the solvent effect, $k(\text{water})/k(50\% \text{ acetone-water})$, of 14 for spontaneous hydrolyses of acetyl fluoride can be compared with that of 22.6 [10] for benzoyl fluoride—hydrolyses are also acid and base-catalysed [37]; (ii) observed rates of solvolyses of *p*-dimethylaminobenzoyl fluoride (**3**, $Z = \text{NMe}_2$) are almost independent of solvent composition from 80% to 60% ethanol, before a mechanistic change occurs to a cationic pathway [9].

2.2.4. Sulfonyl Chlorides

Sulfonyl chlorides have relatively high heterolytic bond dissociation energies [21], implying that formation of sulfonyl cations is unfavourable. Even when there is an adjacent nitrogen lone pair, as in $\text{Me}_2\text{NSO}_2\text{Cl}$, solvolyses do not show common-ion effects [38]. Comparison with reactions of various sulfonyl chlorides in acetone-water by plotting $\log k_{\text{obs}}$ vs. Y gave a relatively high value of $m = 0.69$ for $\text{Me}_2\text{NSO}_2\text{Cl}$, leading to the conclusion that the transition state was more ionic (looser) than for other sulfonyl chlorides, but the mechanism was not $\text{S}_{\text{N}}1$ as suggested by others [39]; the KSIE in water and D_2O of 1.3 (Table 2, References [38–45]) is compatible with either $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$ mechanisms. In the terminology of Figure 1, the mechanism of solvolysis of $\text{Me}_2\text{NSO}_2\text{Cl}$, is $\text{S}_{\text{N}}2$, and within the $\text{S}_{\text{N}}2$ – $\text{S}_{\text{N}}1$ spectrum throughout the wide range of Y values, but is not $\text{S}_{\text{N}}1$ even in water. Recent more extensive studies of solvent effects also led to the proposal of an $\text{S}_{\text{N}}2$ mechanism [46].

Table 2. Correlations Equation (3) for acetone-water (% *v/v*) and kinetic solvent isotope effects (KSIE) in water and methanol for solvolyses of sulfonyl chlorides at 25 °C (unless stated otherwise).

Substrate	Slope	Intercept	R^2	Std. Error	Range (%)	KSIE ^a	KSIE ^b
$\text{Me}_2\text{NSO}_2\text{Cl}$ ^c	0.36 ± 0.01	-7.16 ± 0.03	0.990	0.06	80–0	1.33 ^c , 1.29 ^d	
9 ^{ef}	0.16 ± 0.02	-6.05 ± 0.04	0.882	0.11	90–0 ^g	1.46	1.76
<i>i</i> -PrSO ₂ Cl ^h	0.07 ± 0.02	-8.18 ± 0.03	0.770	0.06	80–0	1.66	2.54
MeSO ₂ Cl ⁱ	-0.07 ± 0.02	-6.81 ± 0.03	0.788	0.06	80–0	1.81	1.62
PhCH ₂ SO ₂ Cl ^j	-0.07 ± 0.02	-6.25 ± 0.04	0.688	0.08	90–20		2.34
2 , $Z = \text{NO}_2$ ^k	-0.11 ± 0.01	-5.56 ± 0.03	0.922	0.06	90–0	1.76 ^l	2.31 ^l
2 , $Z = \text{Cl}$ ^m	0.04 ± 0.02	-6.32 ± 0.03	0.461	0.07	90–0	1.65 ^l	1.89 ^l
2 , $Z = \text{H}$ ^m	0.13 ± 0.03	-6.56 ± 0.06	0.781	0.13	90–0	1.59 ^l	1.79 ^l
CF ₃ SO ₂ Cl ^{en}	-0.42 ± 0.03	-6.14 ± 0.06	0.971	0.11	80–0	2.24	3.08

^a Refers to $k_{\text{water}}/k_{\text{D}_2\text{O}}$; ^b Refers to $k_{\text{MeOH}}/k_{\text{MeOD}}$; ^c Reference [38]; ^d Reference [39]; ^e At 45 °C; ^f Reference [40];

^g If the data point for 90% acetone-water is omitted: slope = 0.20 ± 0.02 ; intercept = -6.14 ± 0.04 ; $R^2 = 0.953$; std. error = 0.07; ^h Reference [41]; ⁱ Reference [42]; ^j At 35 °C, Reference [43]; ^k Reference [44]; ^l Reference [13];

^m Reference [8]; ⁿ Reference [45], including curved plots of $\log k_{\text{obs}}$ vs. Y_{Cl} and linear plots of $\log k_3$ vs. Y_{Cl} .

Doubts about the interpretation of KSIE data in terms of general base catalysis were based [40,47] on the anomalously high value (2.54) for the KSIE of *i*-PrSO₂Cl in MeOH (Table 2). Also the low value of 1.62 for MeSO₂Cl does not fit the general trend. These outliers obscure the general trend of higher values for KSIE and very low and/or negative Equation (3) slopes for typical sulfonyl chlorides (Table 2). Apart from the first and perhaps second entries in Table 2, the slopes and particularly the

KSIE values of the remaining substrates are consistent with mechanistic changes within the S_N2 - S_N3 spectrum (Figure 1b) throughout the range of acetone-water mixtures. As the concentration of acetone increases, ΔS^\ddagger becomes progressively more negative for solvolyses of **2**, $Z = \text{NO}_2$, Br, H and Me [48,49], indicating more highly ordered transition states.

Most of the slopes shown in Table 2 are so low that the correlation with Y is poor. Typical plots have a very shallow U-shape. Arenesulfonyl chlorides (**2,10**) permit the systematic introduction of more electron donating groups, and U-shaped curvature of the Equation (3) plots becomes progressively more pronounced. These results are discussed in Section 2.3.3.

2.2.5. Anhydrides

Data [11] for solvolyses of *p*-nitrobenzoyl tosylate (**4**, $Z = \text{NO}_2$) show a linear plot (Figure 3) based on Equation (3) and a KSIE of *ca.* 1.7 in methanol [11], consistent with a third order mechanism (Section 2.1). Other anhydrides show lower KSIE values: e.g., in methanol for acetyl tosylate (CH_3COOTs , KSIE = 0.99 at -39.6°C [50]), benzoyl tosylate (PhCOOTs , KSIE = 1.1 at -10°C [11]), and other anhydrides ($\text{PhSO}_2)_2\text{O}$, Ts_2O , Ms_2O) KSIE = 1.35–1.4 at -10°C [51,52]); in mixed organic/water mixtures of 1.2–1.3 for solvolyses of various aromatic sulfonic acid anhydrides at 22.5°C [53]. These results are consistent with changes within the S_N2 - S_N1 spectrum. A rationalization is that the electrophilic “pull” of sulfonates is much larger than for chlorides (e.g., $k_{\text{OMs}}/k_{\text{Cl}} \sim 10^3$ for methanesulfonyl derivatives [52]), so the extra “push” of a third order process is not required.

For solvolyses *p*-bromophenylsulfonic acid anhydride in acetone-water mixtures, an Equation (2) slope of 1.75 was reported [54]. However, these data do not provide sufficient support for a third order mechanism, because of the unreliability of Equation (2)—see Section 2.1. Slopes of plots *vs.* Y depend on the size of the leaving group, probably due to the extent of charge delocalisation in the developing anion; for S_N1 solvolyses of 1-adamantyl substrates an order of slopes $\text{Cl} > \text{Br} > \text{I} > \text{OTs}$ was established [55]; tosylates have relatively low slopes, and given the trend, it would be expected that fluorides would have the highest slopes (e.g., Figure 3).

2.2.6. Phosphorus Halides

KSIE data are few (Table 3, References [56–63]), although there is an additional value of 1.25 for $(\text{Pr}^i\text{O})_2\text{POCl}$ in $k_{\text{water}}/k_{\text{D}_2\text{O}}$ [64]. $(\text{PhO})_2\text{POCl}$ shows approximately constant values of $\log k_3$ and KSIE of **3**, consistent with third order kinetics and general base catalysis by a second water molecule [61]; Hartree-Fock and DFT levels calculations [65] for $(\text{RO})_2\text{POX}$ ($\text{R} = \text{H}$ or Me , $\text{X} = \text{F}$ or Cl) supported this proposal [65]. Other substrates show low and usually positive slopes, due to small changes in $\log k_3$, and hence a poor correlation with Y . The largest positive slope is 0.28 for Ph_2PSCl , but the data refer only to 95%–50% acetone-water [63]. Emphasising the importance of nucleophilic attack, it is usually stated that the hydrolyses are bimolecular [57–65]; it is now suggested that additional KSIE data would confirm the role of catalysis by a second solvent molecule.

Table 3. Results (Equation (3)) for acetone-water (% v/v) and kinetic solvent isotope effects (KSIE) in water and methanol for solvolyses of phosphorus chlorides at 25 °C (unless stated otherwise).

Substrate	Slope	Intercept	R^2	Std. Error	Range (%)	KSIE ^a	KSIE ^b
(Me ₂ N) ₂ POCl ^c	0.17 ± 0.03	-6.28 ± 0.06	0.930	0.09	80–0	1.34 ^d	
11 ^e	0.15 ± 0.06 ^f	-6.55 ± 0.12	0.463	0.30	90–0		
(MeO) ₂ POCl ^g	0.07 ± 0.01	-5.08 ± 0.01	0.979	0.02	90–60		
12 ^h	0.16 ± 0.03	-4.22 ± 0.04	0.884	0.10	90–20		
13 ^{h,i}	-0.06 ± 0.02	-6.22 ± 0.04	0.597	0.08	80–0		
(PhO) ₂ POCl ^j	0 ^k				95–0	2.9 ^l	3.06
Ph ₂ POCl ^m	0 ⁿ				95–50		
Ph ₂ PSCl ^o	0.28 ± 0.03	-4.90 ± 0.04	0.967	0.09	95–50		1.83
(MeO) ₂ PSCl ^g	0.08 ± 0.03	-6.67 ± 0.06	0.503	0.15	90–0		

^a Refers to $k_{\text{water}}/k_{\text{D}_2\text{O}}$; ^b Refers to $k_{\text{MeOH}}/k_{\text{MeOD}}$; ^c Reference [56]; ^d Data from Reference [57] at 10 °C; ^e Reference [58]; ^f Shallow-U-shaped plot; ^g Reference [59]; ^h Reference [60]; ⁱ At 50 °C; ^j Reference [61]; ^k Very shallow U-shaped plot, with all values of $\log k_3$ within the narrow range -4.8 to -5.26; ^l At 5 °C; ^m Reference [62]; ⁿ Values of $\log k_3$ vary within a very narrow range (-2.87 to -2.97); ^o Data from Reference [63] at 55 °C.

2.3. Deviations from Equation (3) as Evidence for Mechanistic Changes

2.3.1. Carboxylic Acid Chlorides

From the results given in Section 2.2.3 (supported by the more extensive data for chloroformates in Section 2.2.1), the normal expectation is that third order solvolyses in acetone-water will fit Equation (3) with zero or negative slopes (mechanism shown in Figure 1b). In contrast, when the cationic mechanism (Figure 1a) operates, there is a much higher and positive slope (e.g., see Figure 1 of Reference [66]). Consequently, it is possible to detect a change in mechanism from third order to cationic by plotting $\log k_{\text{obs}}$ vs. Y ; e.g., for solvolyses of benzoyl chloride (**1**, Z = H) in a wide range of acetone-water mixtures [67]. In addition, a much clearer change in slopes is obtained by plotting $\log k_3$ vs. Y (Equation (3)) instead of $\log k_{\text{obs}}$; the two plots are compared in Figure 6.




Figure 6. Comparison of plots of logarithms of first order rate constants ($\log k_{\text{obs}}$) and third order rate constants ($\log k_3$, Equation (3)) vs. Grunwald-Winstein Y values [4,17] for solvolyses of benzoyl chloride (**1**, Z = H) in acetone-water mixtures at 25 °C; kinetic data from Reference [68].

Close to the point where the two lines intersect (Figure 6, around $Y = 1$, ~60% acetone-water), the two mechanisms are assumed to be concurrent, and so values of $\log k_{\text{obs}}$ and $\log k_3$ are corrected by subtracting $\log 2 = 0.3$ from both $Y = 1$ values (these points are then fitted to both relevant correlation lines).

Solvolyses of *p*-chlorobenzoyl chloride (**1**, $Z = \text{Cl}$) and phenylacetyl chloride in acetone water also show sharp changes in slope when $\log k_3$ is plotted vs. Y for acetone-water mixtures (see Figure 4 of Reference [69] and Figure 3 of Reference [34]). In contrast, the same plots for *p*-nitrobenzoyl chloride (**1**, $Z = \text{NO}_2$) and chloroacetyl chloride are linear throughout the range of acetone-water mixtures (see Section 2.2.3). Solvolyses of trimethylacetyl chloride give a similar plot to Figure 6 when $\log k_{\text{obs}}$ is plotted vs. Y (see Figure 2 of Reference [70]). For solvolyses of *p*-dimethylaminobenzoyl fluoride (**3**, $Z = \text{NMe}_2$) in ethanol-water mixtures, a sharp change in slope is readily apparent even in plots [9] of first order rate constants ($\log k_{\text{obs}}$).

The above results are significant because they show that a change in hydrolysis mechanism can occur when the % water in a binary mixture such as acetone-water is altered.

2.3.2. Chloroformates

Solvolyses of isopropyl chloroformate (**5**, $R = ^i\text{Pr}$) show a KSIE in water at 25 °C of only 1.25 [71], in contrast to values of 1.8–2.2 for methyl, ethyl and aryl derivatives [72]; also, values of $\log k_3$ in water for **5**, $R = ^i\text{Pr}$ deviate substantially from the downward trend set by 90%–70% acetone-water compositions (Figure 7). These results are in marked contrast to solvolyses of ethyl chloroformate (**5**, $R = \text{Et}$), which shows a close to constant value of $\log k_3$ from 95% v/v acetone-water to 100% water (Table 1). It can be concluded that both **5**, $R = \text{Et}$ and **5**, $R = ^i\text{Pr}$ are hydrolysed by the third order mechanism (Figure 1b) in 90% and 80% acetone-water mixtures, but the mechanism changes to cationic for hydrolysis of **5**, $R = ^i\text{Pr}$ in water. The minimum at $Y \sim 0$ corresponds to 70% acetone, and $\log 2 = 0.3$ was subtracted from this data point before fitting it to both correlation lines (Figure 7).




Figure 7. Changes in mechanism based on plots of $\log k_3$ vs. Y Equation (3) for solvolyses in acetone-water of **5**, $R = ^i\text{Pr}$ at 40 °C and **5**, $R = \text{CH}_2\text{CH}(\text{CH}_3)_2$ at 45 °C; kinetic data from References [71–73].

The above conclusion is in good agreement with published work [72], using an independent approach. From the extended GW equation (Section 2.4) for only seven solvents clearly favouring the cationic mechanism (water, formic acid, and various $\text{CF}_3\text{CH}_2\text{OH}$ -water and $(\text{CF}_3)_2\text{CHOH}$ -water mixtures), it was calculated [72] that the mechanistic changeover for **5**, $R = ^i\text{Pr}$ occurred between

70 and 80% acetone-water. The prediction in Figure 7 that the mechanistic change occurs in 70% acetone-water is based on deviations from an independent correlation for the third order mechanism.

The plot (red in Figure 7) of $\log k_3$ vs. Y for solvolyses of isobutyl chloroformate (**5**, $R = \text{CH}_2\text{CH}(\text{CH}_3)_2$) show very similar values to those for **5**, $R = i\text{Pr}$ in 90%–60% acetone-water (first 4 data points); note that the temperatures are not identical (40 and 45 °C). Compared with the line drawn for **5**, $R = i\text{Pr}$, additional data in 50%–10% acetone for **5**, $R = \text{CH}_2\text{CH}(\text{CH}_3)_2$ show a smaller change in slope; also the KSIE in water of 1.54 [73] is larger. The KSIE in methanol of 2.03 [73] and the low slope of the plot for 90%–60% acetone (Y values from -2 to 1) are consistent with third order reactions, and a change to a bimolecular mechanism was suggested for reaction in water [73]. Comparing slopes and KSIE values in water, it appears that the change in mechanism begins slightly later and is more gradual for **5**, $R = \text{CH}_2\text{CH}(\text{CH}_3)_2$ than for **5**, $R = i\text{Pr}$.

Both 9-fluorenylmethyl chloroformate (**14**, Figure 8) and **5**, $R = \text{CH}_2\text{CH}(\text{CH}_3)_2$ have a β,β -dialkyl group and solvolyses of **14** in acetone-water [74] also give shallow U-shaped plot (not shown); values of $\log k_3$ at 45 °C vary from -5.75 (90% acetone) to -6.11 (50% acetone) and then up to -5.70 (20% acetone); the KSIE in methanol is 2.2, but there are no KSIE data for water [74].




Figure 8. Structures of additional substrates: 9-fluorenylmethyl chloroformate (**14**); phenyl chlorothionoformate (**15**); 5-dimethylamino-naphthalene-1-sulfonyl (dansyl) chloride (**16**).

An almost identical pattern of results is observed (Figure 9) for solvolyses of phenyl chlorothionoformate (**15**); $\log k_3$ values are close to constant from $Y = -1$ to $+2$ (80%–40% acetone); then there is an increase in slope (Figure 9); the KSIE value are 2.02 for methanol, but only 1.45 for water [75], so a gradual change to a bimolecular mechanism may be underway.




Figure 9. Plots of $\log k_3$ vs. Y (Equation (3)) for solvolyses in acetone-water of phenylthiono-formate (**15**) at 25 °C, and isopropenyl chloroformate (**5**, $R = \text{CH}_3\text{C}=\text{CH}_2$) at 35 (upper plot) and 10 °C (middle plot); kinetic data from References [75–77].

Equation (3) plots for isopropenyl chloroformate (**5**, R = CH₃C=CH₂) at 10 °C show close to constant values of log k_3 from 90% acetone to water (Figure 9, middle), whereas the upper plot at 35 °C is U-shaped. Solvent effects and KSIE values in water and methanol of >2.0 at 10 °C are very similar to those for **1**, Z = NO₂ [76], so there is strong evidence for the third order mechanism in acetone-water. At 35 °C [77], there may be an increase in slope from 50% to 20% acetone-water ($Y > 1.4$, Figure 9) and a bimolecular cationic mechanism (Figure 1a) having a higher activation energy may start to be competitive. Evidence for a change to the cationic reaction mechanism was later obtained by comparing solvolyses in fluorinated alcohols with those for phenyl chloroformate [78].

2.3.3. Sulfonyl Chlorides

Although most sulfonyl chlorides react by the third order mechanism within the S_N3-S_N2 spectrum (Section 2.2.4), there is evidence for a mechanistic changeover to the S_N2-S_N1 spectrum for electron-rich substrates; however, formation of sulfonyl cations is relatively unfavourable [21], and there is no evidence for an S_N1 mechanism. Evidence for the changeover will be based on KSIE data (Table 4, References [79–81]) and non-linear Equation (3) plots (Figures 10 and 11), but there is also support from independent evidence on alcohol/water selectivities [79,81,82].

In all four examples (Table 4), the KSIE decreases from >1.5 in methanol (S_N3-S_N2 spectrum) to reach a value of 1.3–1.4 more characteristic of the S_N2-S_N1 spectrum. Also, the substituent effect for 4-substituted 2,6-dimethyl substrates (**10**) changes from $k_{\text{OMe}} < k_{\text{Me}}$ on the left side of Figure 11 (S_N3-S_N2 spectrum) to $k_{\text{OMe}} > k_{\text{Me}}$ on the right side (consistent with relatively more bond breaking for the S_N2-S_N1 spectrum).

Table 4. Mechanistic changeover for solvolyses of electron-rich sulfonyl chlorides.

Sulfonyl Chloride	KSIE (MeOH)	KSIE (Water) ^a	Equation (3) Plot
<i>p</i> -methoxybenzene (2 , Z = OMe)	1.58 ^a	1.37 ^a	Figure 10
Dansyl (16)	1.88 ^b	(1.34) ^{b,c}	Figure 10
Mesityl (10 , Z = Me)	1.68 ^d	1.34 ^d	Figure 11
<i>p</i> -methoxy (10 , Z = OMe)	1.58 ^e	1.41 ^a	Figure 11

^a Data from Reference [13] at 25 °C; ^b Reference [79] at 35 °C; ^c Refers to 50/50 MeOH/water; ^d Reference [80] at 25 °C; ^e Reference [81] at 25 °C.

For solvolyses of **2**, Z = OMe, the changes in slopes based on Equation (3) are large, and the minimum value of log k_3 calculated from experimental data was lowered by log 2 = 0.3, before fitting the two correlation lines (Figure 10).

For solvolyses of **10**, Z = Me (Figure 11), and **16** (Figure 10), there are two minimum values of log k_3 between $Y = 0$ and $Y = 1$; each of these was lowered by 0.15 before fitting the two correlation lines; for solvolyses of **10**, Z = OMe, there is no clear minimum, so lines are drawn through the unadjusted values of log k_3 (coded in blue, Figure 11).




Figure 10. Plots of $\log k_3$ vs. Y (Equation (3)) for solvolyses in acetone-water of 4-methoxybenzene sulfonyl (**2**, $Z = \text{OMe}$) at 25 °C, and dansyl chlorides (**16**) at 35 °C; data from References [79,81,82].




Figure 11. Plots of $\log k_3$ vs. Y Equation (3) for solvolyses in acetone-water of 4-Z-2,6-di-methylbenzene sulfonyl chloride (**10**, $Z = \text{Me}, \text{OMe}$) at 25 °C; kinetic data from References [80,81].

2.4. Multi-Parameter Equations

Most of the above discussion has been concerned with solvolyses in water, acetone-water and methanol. Correlations of kinetic data for a wider range of solvents require multi-parameter correlations. In theory these are more comprehensive, but they are also more complex, and caution is recommended [83].

Retaining the assumption of third order rate laws and water as one of the solvents, the possibility of cosolvent acting as general base and/or as nucleophile leads to additional third order rate constants. Assuming that the cosolvent (e.g., acetone, acetonitrile, dioxan) acts as general base (or in some other

way), but not as nucleophile, two third order rate constants (k_{ww} and k_{wc}) contribute to k_{obs} , and Equation (5) is obtained [35,44].

$$k_{obs}/[\text{water}]^2 = k_{ww} + k_{wc} [\text{cosolvent}]/[\text{water}] \quad (5)$$

If the cosolvent is an alcohol, it could act as nucleophile and/or general base, and four third order rate constants are then possible. In such cases there are two products, and product ratios provide useful additional information [35,44].

A more general approach, applicable to a very wide range of solvents and solvent mixtures, is to extend the Grunwald-Winstein equation to include a term for solvent nucleophilicity. In this Equation (6), logarithms of observed first order rate constants in any solvent ($\log k$), relative to solvolyses in 80% ethanol-water (k_0), are correlated with Y and a term (N) for solvent nucleophilicity [5]; m is the response to changes in Y and l is the response to changes in N and c is a residual term.

$$\log (k/k_0) = lN + mY + c \quad (6)$$

Equation (6) was first applied to the S_N2 - S_N1 spectrum of solvolyses of primary and secondary alkyl tosylates [5]. Solvolyses closer to the S_N1 extreme have relatively high values of m and low values of l ; as the substrates become more susceptible to nucleophilic attack (e.g., 2-propyl, ethyl, methyl tosylates), values of m decrease and values of l increase in accordance with Equation (7). In terms of mechanism, a higher m implies more electrophilic “pull”, and a higher l implies greater nucleophilic “push”; a further implication of Equation (7) is that if there is more “pull” then less push is required (and *vice versa*).

$$l = (1 - m)/0.7 \quad (7)$$

For many of the studies of substrates discussed above, it has become standard practice to interpret l/m ratios. For an S_N1 reaction, $l = 0$, so $l/m = 0$. As l increases and m decreases, l/m ratios >2 can be obtained [40]. Consequently, l values and/or l/m ratios help to position substrates within a spectrum of mechanisms (e.g., for sulfonyl chlorides [40]). Values of m depend on how the N scale is defined [84], and published values are usually based on the N_T scale [85,86].

Deviations from a single correlation can be interpreted as evidence for a second reaction channel, and a correlation based on Equation (6) can often be obtained for each reaction channel (e.g., for solvolyses of a variety of ROCOCl , RSCl , ROCSCl , and RSCSCl substrates [87]). The two channels can be compared or characterized using l and m values or the l/m ratios.

2.5. Mechanistic Insights from Theoretical Calculations

Mechanistic conclusions from kinetic studies of hydrolyses of benzenesulfonyl chlorides (**2**) have recently been supported by theoretical calculations [7]. The calculations for water clusters in acetone indicate that reactions do not involve the formation of an intermediate and the S_N3 reaction is favoured by electron withdrawing substituents and by clusters having a small number of water molecules (*i.e.*, low Y).

The spectrum of mechanisms from nucleophilic attack by water (S_N2), often assisted by general base catalysis by a second water molecule (S_N3) can be illustrated (Figure 12) by the changes in bond lengths calculated [7] for a series of substrates (**2**, $Z = \text{OMe, Me, H, Cl and NO}_2$).




Figure 12. Calculated [7] changes in bond lengths (Angstroms) in transition states for hydrolyses of benzenesulfonyl chlorides (**2**) in a cluster of 17 water molecules in acetone.

The results, including numbering of atoms, are taken from Table 3 of Reference [7], and focus only on the substrate, the water molecule acting as nucleophile via O(16) and the water molecule acting as general base through O(22). Except for the approximately constant O(16)–H(17) distance, bond lengths extend significantly as the substituent changes from Z = NO₂ to OMe; as expected for the proposed S_N3 mechanism, the O(22)–H(17) distance in the second water molecule is considerably shorter than the value of 1.8 expected for a hydrogen bonded water molecule, and is shortest for Z = NO₂.

2.6. Comments on Mechanistic Details

The mechanism of Song and Jencks [9] is illustrated in Figure 13 for benzoyl fluorides (**3**). The key features are the same as those shown in Figure 1b. General base catalysis by a base (B, e.g., water) assists nucleophilic attack by activating the attacking water nucleophile and by preventing the reverse reaction. The transition state could lead to a tetrahedral intermediate or there may be an “uncoupled concerted reaction” [9,88] in which the C–F bond is cleaved before the intermediate can be formed.




Figure 13. General base catalysis mechanism [9] for hydrolysis of benzoyl fluoride (**3**).

In contrast, nucleophilic attack at the carbonyl group of chloro- and fluoroformates (**5,7**) is usually [29,77,87,89] described as shown in Figure 14. Two tetrahedral intermediates (**18,19**), not transition states are usually drawn. It is implied that there may be general base catalysis, but it not usually shown by these authors. However, general base catalysis is a key component of the proposed mechanism (Figures 1b and 13), so the fast deprotonation step is avoided, and the first tetrahedral intermediate (**18**) is by-passed.




Figure 14. Stepwise addition-elimination mechanism [29,77,87,89] for solvolyses of haloformates, illustrated for nucleophilic attack by solvent (SOH) on fluoroformates (**7**).

Also, it does not seem likely that the alkoxide intermediate (**19**) could be formed under typical reaction conditions of *ca.* 10^{-2} M substrate hydrolysed in unbuffered media, so the $\text{pH} < 3$; under these conditions, protonation of **19** would be expected, but the development of negative charge in the transition state is needed to explain the relatively high response to changes in Y for solvolyses of acid fluorides (Section 2.2.2 and Figure 3). As well as the proposed [29] developing charge on oxygen, there may be a partial charge on fluorine in a transition state similar to **17**. F/Cl rate ratios of *ca.* 1 [29] provide evidence for the addition character of either a transition state or a tetrahedral intermediate [88].

Previously, it has been proposed that tetrahedral intermediates such as **19** may be by-passed during nucleophilic substitution reactions of acid chlorides [68] and chloroformates [90]. A single transition state has also been proposed for acyl group transfer between phenolate nucleophiles [91] and also for $\text{S}_{\text{N}}(\text{Ar})$ reactions [92]. Therefore, although these reactions are often drawn to show stepwise mechanisms, there is substantial evidence for the concerted alternative.

The proposal that water as base assists water as nucleophile (Figure 1b) can be extended to hydroxide-catalysed hydrolyses of esters [93,94] and to $\text{S}_{\text{N}}(\text{Ar})$ reaction [95]. For base-catalysed hydrolysis of methyl formate, the kinetic isotope effect on the oxygen nucleophile was explained [93] by water acting as nucleophile and hydroxide acting as general base. This mechanism was later supported by proton inventory measurements on the base-catalysed hydrolysis of ethyl acetate [94]. Alkoxide-catalysed hydrolysis of a chlorotriazine indirectly indicates that hydroxide-catalysed hydrolysis also occurs in $\text{S}_{\text{N}}(\text{Ar})$ reactions [95].

3. Experimental Section

Kinetic data are from the references cited. Correlations were performed using Microsoft Excel.

4. Conclusions

Equation (3) provides a useful additional way to investigate mechanisms of hydrolyses, particularly for the $\text{S}_{\text{N}}2$ - $\text{S}_{\text{N}}3$ spectrum (Tables 1–3) and for changes in reaction channel (e.g., Figures 6, 7 and 9–11); it is more reliable than Equation (2) because solvent effects are included, and less complex than multi-parameter correlations (Section 2.4). KSIE data in water and methanol provide very useful additional information.

For sulfonyl chlorides, hydrolyses usually involve a second water molecule acting as a general base ($\text{S}_{\text{N}}2$ - $\text{S}_{\text{N}}3$ channel), but there is no reliable evidence for an addition-elimination process involving an intermediate. For the most electron-rich substrates, Equation (3) plots (Figures 10 and 11) support

previous evidence for a gradual change in transition structure to the S_N2-S_N1 (cationic) reaction channel in the more polar mixtures; in less polar mixtures (lower *Y*), the S_N2-S_N3 channel becomes significant.

The same trends may apply to hydrolyses of other acid derivatives (e.g., haloformates and carboxylic acid halides), except that cationic reactions are more favourable for carboxylic acid halides. When the alternative “addition-elimination” process occurs (e.g., for fluoroformates), tetrahedral intermediates may be by-passed.

Acknowledgments

Thanks are due to Professor Dennis N. Kevill (Northern Illinois University) for helpful comments on a preliminary draft of the manuscript.

Conflicts of Interest

The author declares no conflict of interest.

References

1. Streitwieser, A. Solvolytic displacement reactions at saturated carbon atoms. *Chem. Rev.* **1956**, *56*, 571–752.
2. Bentley, T.W.; Schleyer, P.v.R. Medium effects on the rates and mechanisms of solvolytic reactions. *Adv. Phys. Org. Chem.* **1976**, *8*, 1–67.
3. Williams, A. *Free Energy Relationships in Organic and Bioorganic Chemistry*; The Royal Society of Chemistry: Cambridge, UK, 2003.
4. Grunwald, E.; Winstein, S. Correlation of solvolysis rates. *J. Am. Chem. Soc.* **1948**, *70*, 846–854.
5. Schadt, F.L.; Bentley, T.W.; Schleyer, P.v.R. The S_N2-S_N1 spectrum 2. Quantitative treatments of nucleophilic solvent assistance. A scale of solvent nucleophilicities. *J. Am. Chem. Soc.* **1976**, *98*, 7667–7674.
6. Ruff, F.; Farkas, O. Concerted S_N2 mechanism for the hydrolysis of acid chlorides: comparisons of reactivities calculated by density functional theory with experimental data. *J. Phys. Org. Chem.* **2011**, *24*, 480–491.
7. Yamabe, S.; Zeng, G.; Guan, W.; Sakaki, S. S_N1-S_N2 and S_N2-S_N3 mechanistic changes revealed by transition states of the hydrolyses of benzyl and benzenesulfonyl chlorides. *J. Comput. Chem.* **2014**, *35*, 1140–1148.
8. Bentley, T.W.; Jones, R.O.; Kang, D.H.; Koo, I.S. The S_N3-S_N2 spectrum. Rate constants and product selectivities for solvolyses of benzenesulfonyl chlorides in aqueous alcohols. *J. Phys. Org. Chem.* **2009**, *22*, 799–806.
9. Song, B.D.; Jencks, W.P. Mechanisms of solvolysis of substituted benzoyl halides. *J. Am. Chem. Soc.* **1989**, *111*, 8470–8479.
10. Kevill, D.N.; D’Souza, M.J. Correlation of the rates of solvolysis of benzoyl fluoride and a consideration of leaving group effects. *J. Org. Chem.* **2004**, *69*, 7044–7050.
11. Kevill, D.N.; Ryu, Z.H. Reaction mechanism studies involving the correlation of the rates of solvolysis of benzoyl and *p*-nitrobenzoyl *p*-toluenesulfonates. *J. Chem. Res.* **2014**, *38*, 387–395.

12. Johnson, S.L. General base and nucleophilic catalysis of ester hydrolysis and related reactions. *Adv. Phys. Org. Chem.* **1967**, *5*, 237–330.
13. Koo, I.S.; Lee, I.; Oh, J.; Yang, K.; Bentley, T.W. Substituent effects on the kinetic solvent isotope effect in solvolyses of arenesulfonyl chlorides. *J. Phys. Org. Chem.* **1993**, *6*, 223–227.
14. Koo, I.S.; Yang, K.; Kang, K.; Lee, I. Transition state variation in solvolyses of para-substituted phenyl chloroformates in alcohol-water mixtures. *Bull. Korean Chem. Soc.* **1998**, *19*, 968–973.
15. Bentley, T.W.; Koo, I.S.; Norman, S.J. Distinguishing between solvation effects and mechanistic changes. Effects due to differences in solvation between aromatic rings and alkyl groups. *J. Org. Chem.* **1991**, *56*, 1604–1609.
16. Bentley, T.W.; Llewellyn, G.; Ryu, Z.H. Solvolytic reactions in fluorinated alcohols. Role of nucleophilic and other solvations effects. *J. Org. Chem.* **1998**, *63*, 4654–4659.
17. Fainberg, A.H.; Winstein, S. Correlation of solvolysis rates. III. *t*-Butyl chloride in a wide range of solvent mixtures. *J. Am. Chem. Soc.* **1956**, *78*, 2770–2777.
18. Kevill, D.N.; D'Souza, M.J. Sixty years of the Grunwald-Winstein equation: Development and recent applications. *J. Chem. Res.* **2008**, *2008*, 61–66.
19. Bentley, T.W.; Llewellyn, G. Y_X scales of solvent ionizing power. *Prog. Phys. Org. Chem.* **1990**, *17*, 121–158.
20. Bentley, T.W.; Koo, I.S.; Llewellyn, G.; Norman, S.J. Similarity models for solvent effects on reactivity. The dissociative channel for solvolyses of sulfonyl chlorides in binary aqueous mixtures. *Croat. Chem. Acta* **1992**, *65*, 575–583.
21. Bentley, T.W. Structural effects on the solvolytic reactivity of carboxylic and sulfonic acid chlorides. Comparisons with gas phase data for cation formation. *J. Org. Chem.* **2008**, *73*, 6251–6257.
22. Swain, C.G. Kinetic evidence for a termolecular reaction in displacement reactions of triphenyl methyl halides in benzene solution. *J. Am. Chem. Soc.* **1948**, *70*, 1119–1128.
23. Kevill, D.N.; D'Souza, M.J. Concerning the two reaction channels for solvolyses of ethyl chloroformate and ethyl chlorothioformate. *J. Org. Chem.* **1998**, *63*, 2120–2124.
24. Koh, H.J.; Kang, S.J. Correlation of the rates of solvolysis on 2,2,2-trichloroethyl chloroformate using the extended Grunwald-Winstein equation. *Bull. Korean Chem. Soc.* **2012**, *33*, 1729–1733.
25. Kevill, D.N.; Koyoshi, F.; D'Souza, M.J. Correlation of specific rates of solvolysis of solvolysis of aromatic carbamoyl chlorides, chloroformates, chlorothionoformates, and chlorodithionoformates revisited. *Int. J. Mol. Sci.* **2007**, *8*, 346–362.
26. Koh, H.J.; Kang, S.J.; Kevill, D.N. Kinetic studies of the solvolyses of 2,2,2-trichloroethyl-1,1-dimethyl chloroformate. *Bull. Korean Chem. Soc.* **2010**, *21*, 835–839.
27. Koo, I.S.; Yang, K.; Kang, K.; Lee, I.; Bentley, T.W. Stoichiometric solvation effects. Part 3. Product-rate correlations for solvolyses of *p*-nitrophenyl chloroformate in alcohol-water mixtures. *J. Chem. Soc. Perkin Trans. 2* **1998**, 1179–1183, doi:10.1039/A800179K.
28. D'Souza, M.J.; Kevill, D.N. Application of the Grunwald-Winstein equations to studies of solvolytic reactions of chloroformate and fluoroformate esters. *Recent Res. Dev. Org. Chem.* **2013**, *13*, 1–38.
29. Kevill, D.N.; D'Souza, M.J. Correlation of the rates of solvolysis of *n*-octyl fluoroformate and a comparison with *n*-octyl chloroformate. *J. Chem. Soc. Perkin Trans. 2* **2002**, 240–243, doi:10.1039/B109169G.

30. Lee, Y.W.; Seong, M.H.; Kyong, J.B.; Kevill, D.N. Correlation of the rates of solvolysis of *t*-butyl fluoroformate using the extended Grunwald-Winstein equation. *Bull. Korean Chem. Soc.* **2010**, *31*, 3366–3370.
31. Kevill, D.N.; Kyong, J.B. Multiple pathways in the solvolyses of 1-adamantyl fluoroformate. *J. Org. Chem.* **1992**, *57*, 258–265.
32. D'Souza, M.J.; McAneny, M.J.; Kevill, D.N.; Kyong, J.B.; Choi, S.H. Kinetic evaluation of the solvolysis of isobutyl chloro- and thiochloroformate esters. *Beilstein J. Org. Chem.* **2011**, *7*, 543–552.
33. Bentley, T.W.; Harris, H.C. Separation of mass law and solvent effects in the kinetics of solvolyses of *p*-nitrobenzoyl chloride in aqueous binary mixtures. *J. Org. Chem.* **1988**, *53*, 724–728.
34. Bentley, T.W.; Harris, H.C.; Ryu, Z.H.; Lim, G.T.; Sung, D.D.; Szajda, S.R. Mechanisms of solvolyses of acid chlorides and chloroformates. Chloroacetyl and phenylacetyl chloride as similarity models. *J. Org. Chem.* **2005**, *70*, 8963–8970.
35. Bentley, T.W.; Jones, R.O. Stoichiometric solvation effects. Part 1. New equations relating product selectivities to alcohol-water solvent compositions for hydrolyses of *p*-nitrobenzoyl chloride. *J. Chem. Soc. Perkin Trans. 2* **1993**, 2351–2357, doi:10.1039/P29930002351.
36. Koo, I.S.; Lee, S.I.; An, S.K.; Yang, K.; Lee, I. Nucleophilic substitution reaction of α -methoxy- α -(trifluoromethyl)phenylacetyl chloride in alcohol-water mixtures. *Bull. Korean Chem. Soc.* **1999**, *20*, 1451–1456.
37. Bunton, C.A.; Fendler, J.H. The hydrolysis of acetyl fluoride. *J. Org. Chem.* **1966**, *31*, 2307–2312.
38. Rogne, O. Solvolysis of dimethylsulphamoyl chloride in water and aqueous acetone. *J. Chem. Soc. B* **1969**, 663–665, doi:10.1039/J29690000663.
39. Ko, E.C.F.; Robertson, R.E. The hydrolysis of sulfamoyl chlorides. I. Hydrolysis of dimethylsulfamoyl chloride. Heat capacity of activation, the secondary γ -deuterium isotope effects, and solvent isotope effect. *J. Am. Chem. Soc.* **1972**, *94*, 573–575.
40. Ryu, Z.H.; Lee, S.W.; D'Souza, M.J.; Yaakoubd, L.; Feld, S.E.; Kevill, D.N. Correlation of the rates of solvolysis of two arenesulfonyl chlorides and of *trans*- β -styrenylsulfonyl chloride—Precursors in the development of new pharmaceuticals. *Int. J. Mol. Sci.* **2008**, *9*, 2639–2657.
41. Koo, I.S.; Yang, K.; Shin, H.B.; An, S.K.; Lee, J.P.; Lee, I. Stoichiometric solvation effects. Solvolysis of isopropylsulfonyl chloride. *Bull. Korean Chem. Soc.* **2004**, *25*, 699–703.
42. Koo, I.S.; Yang, K.; An, S.K.; Lee, C-K.; Lee, I. Stoichiometric solvation effects. Solvolysis of methanesulfonyl chloride. *Bull. Korean Chem. Soc.* **2000**, *21*, 1011–1014.
43. Koh, H.J.; Kang, S.J. Application of the extended Grunwald-Winstein equation to solvolyses of phenylmethanesulfonyl chloride in aqueous binary mixtures. *Bull. Korean Chem. Soc.* **2011**, *32*, 1897–1901.
44. Bentley, T.W.; Jones, R.O.; Koo, I.S. Stoichiometric solvation effects. Part 2. A new product-rate correlation for solvolyses of *p*-nitrobenzene sulfonyl chloride in alcohol-water mixtures. *J. Chem. Soc. Perkin Trans. 2* **1994**, 753–759, doi:10.1039/P29940000753.
45. Koo, I.S.; Yang, K.; Park, J.K.; Woo, M.Y.; Cho, J.M.; Lee, J.P.; Lee, I. Stoichiometric solvation effects. Solvolyses of trifluoromethane sulfonyl chloride. *Bull. Korean Chem. Soc.* **2001**, *26*, 1241–1245.

46. Kevill, D.N.; Park, B.-C.; Park, K.-H.; D'Souza, M.J.; Yaakoubd, L.; Mlynarski, S.L.; Kyong, J.B. Rate and product studies in the solvolyses of N,N-dimethylsulfamoyl chloride and 2-propanesulfonyl chloride. *Org. Biomol. Chem.* **2006**, *4*, 1580–1586.
47. D'Souza, M.J.; Yaakoubd, L.; Mlynarski, S.L.; Kevill, D.N. Concerted solvent processes for common sulfonyl chloride precursors used in the synthesis of sulphonamide-based drugs. *Int. J. Mol. Sci.* **2008**, *9*, 914–925.
48. Jenkins, F.E.; Hambly, A.N. Solvolyses of sulfonyl halides. I. The hydrolysis of aromatic sulfonyl chlorides in aqueous dioxan and aqueous acetone. *Aust. J. Chem.* **1961**, *14*, 190–204.
49. Kim, W.K.; Lee, I. Nucleophilic displacement at sulfur center (II). Solvolysis of benzenesulfonyl chloride in acetone-water mixtures. *J. Korean Chem. Soc.* **1973**, *17*, 163–168.
50. Kevill, D.N.; Ryu, Z.H. Correlation of the rates of solvolysis of acetyl *p*-toluenesulfonate (acetyl tosylate) and a comparison with acetyl halides. *J. Phys. Org. Chem.* **2013**, *26*, 983–988.
51. Kevill, D.N.; Ryu, Z.H.; Niedermeyer, M.A.; Koyoshi, F.; D'Souza, M.J. Rate and product studies in the solvolyses of methanesulfonic acid anhydride and a comparison with methane sulfonyl chloride solvolyses. *J. Phys. Org. Chem.* **2007**, *20*, 431–438.
52. Kevill, D.N.; Ryu, Z.H. Rate and product studied in solvolyses of two arenesulfonic acid anhydrides. *J. Chem. Res.* **2007**, 365–369.
53. Christensen, N.H. Kinetic solvent isotope effect in the hydrolyses of sulfonic anhydrides. *Acta Chem. Scand.* **1967**, *21*, 899–904.
54. Christensen, N.H. Kinetic studies of the hydrolysis of aromatic sulfonic anhydrides. *Acta Chem. Scand.* **1966**, *20*, 1955–1964.
55. Bentley, T.W.; Carter, G.E.; Roberts, K. Solvent ionizing power. Comparisons of solvolyses of 1-adamantyl chlorides, bromides, iodides and tosylates in protic solvents. *J. Org. Chem.* **1984**, *49*, 5183–5189.
56. Kevill, D.N.; Miller, B. Application of the N_T solvent nucleophilicity scale to attack at phosphorus. Solvolyses of *N,N,N,N*-tetramethyl diamidophosphorochloridate. *J. Org. Chem.* **2002**, *67*, 7399–7406.
57. Traylor, P.S.; Westheimer, F.H. Mechanism of hydrolysis of phosphorodiamidic chlorides. *J. Am. Chem. Soc.* **1965**, *87*, 553–559.
58. Choi, H.; Yang, K.; Koh, H.J.; Koo, I.S. Solvolysis reaction kinetics, rates and mechanisms for phenyl *N*-phenyl phosphoamidochloridate. *Bull. Korean Chem. Soc.* **2014**, *35*, 2465–2470.
59. Kevill, D.N.; Carver, J.S. Rate and product studies with dimethyl phosphorochloridate and phosphorochloridothionate. *Org. Biomol. Chem.* **2004**, *2*, 2040–2043.
60. Koh, H.J.; Kang, S.J.; Kevill, D.N. Rate and product studies in the solvolyses of two cyclic phosphorochloridate esters. *Phosphorus Sulfur Silicon* **2010**, *185*, 1404–1415.
61. Bentley, T.W.; Ebdon, D.; Llewellyn, G.; Abduljaber, M.H.; Miller, B.; Kevill, D.N. Chlorophosphate solvolyses. Evaluation of third order rate laws and rate-product correlations for solvolyses of diphenylphosphorochloridate in aqueous alcohols. *J. Chem. Soc. Dalton Trans.* **1997**, 3819–3825, doi:10.1039/A703409A.
62. Kevill, D.N.; Koh, H.J. Correlation of the rates of solvolysis of diphenylphosphinyl chloride using an extended form of the Grunwald-Winstein equation. *J. Phys. Org. Chem.* **2007**, *20*, 88–92.

63. Koh, H.J.; Kang, S.J.; Kevill, D.N. Correlation of the rates of solvolysis of diphenyl-thiophosphinyl chloride using an extended form of the Grunwald-Winstein equation. *Bull. Korean Chem. Soc.* **2008**, *29*, 1927–1931.
64. Dostrovsky, I.; Halman, M. Kinetic studies in the phosphinyl chloride and phosphochloridate series. Part 1. Solvolytic reactions. *J. Chem. Soc.* **1953**, 502–507, doi:10.1039/JR9530000508.
65. Lee, O.-S.; Yang, K.; Kang, K.D.; Koo, I.S.; Kim, C.-K.; Lee, I. *Ab initio* and DFT studies on hydrolyses of phosphorus halides. *J. Comput. Chem.* **2004**, *25*, 1740–1748.
66. Bentley, T.W.; Llewellyn, G.; McAlister, J.A. S_N2 mechanism for alcoholysis, aminolysis and hydrolysis of acetyl chloride. *J. Org. Chem.* **1996**, *61*, 7927–7932.
67. Bentley, T.W.; Carter, G.E.; Harris, H.C. S_N2 character of hydrolysis of benzoyl chloride. *J. Chem. Soc. Chem. Commun.* **1984**, 387–389, doi:10.1039/C39840000387.
68. Bentley, T.W.; Carter, G.E.; Harris, H.C. Competing S_N2 and carbonyl addition pathways for solvolyses of benzoyl chloride in aqueous media. *J. Chem. Soc. Perkin Trans. 2* **1985**, 983–990, doi:10.1039/P29850000983.
69. Bentley, T.W.; Koo, I.S. Trends in selectivity. Evidence from rates and products for simultaneous reaction channels in solvolyses of benzoyl chloride and substituted derivatives. *J. Chem. Soc. Perkin Trans. 2* **1989**, 1385–1392, doi:10.1039/P29890001385.
70. Bentley, T.W.; Shim, C.B. Dual reaction channels for solvolyses of acyl chlorides in alcohol-water mixtures. *J. Chem. Soc. Perkin Trans. 2* **1993**, 1659–1663, doi:10.1039/P29930001659.
71. Queen, A. Kinetics of the hydrolysis of acyl chlorides in water. *Can. J. Chem.* **1967**, *45*, 1619–1629.
72. D'Souza, M.J.; Reed, D.N.; Erdman, K.J.; Kyong, J.B.; Kevill, D.N. Grunwald-Winstein analysis—Isopropyl chloroformate solvolysis revisited. *Int. J. Mol. Sci.* **2009**, *10*, 862–879.
73. Lim, G.T.; Lee, Y.H.; Ryu, Z.H. Further kinetic studies of solvolytic reactions of isobutyl chloroformate in solvents of high ionizing power under conductometric conditions. *Bull. Korean Chem. Soc.* **2013**, *34*, 615–621.
74. Koh, H.J.; Kang, S.J. A kinetic study on solvolysis of 9-fluorenylmethyl chloroformate. *Bull. Korean Chem. Soc.* **2011**, *32*, 3799–3801.
75. Koo, I.S.; Yang, K.; Kang, D.H.; Park, H.J.; Kang, K.; Lee, I. Transition state variation in the solvolyses of phenylchlorothionoformate in alcohol-water mixtures. *Bull. Korean Chem. Soc.* **1999**, *20*, 577–580.
76. Ryu, Z.H.; Lee, Y.H.; Oh, Y. Stoichiometric effects. Correlation of the rates of solvolysis of isopropenyl chloroformate. *Bull. Korean Chem. Soc.* **2005**, *26*, 1761–1766.
77. Koh, H.J.; Kang, S.J. Kinetic studies of solvolyses of isopropenyl chloroformate. *Bull. Korean Chem. Soc.* **2010**, *31*, 1793–1796.
78. D'Souza, M.J.; Shuman, K.E.; Omondi, A.O.; Kevill, D.N. Detailed analysis for the solvolysis of isopropenyl chloroformate. *Eur. J. Chem.* **2011**, *2*, 130–135.
79. Koh, H.J.; Kang, S.J. Rate and product studies of 5-dimethylamino-naphthalene-1-sulfonyl chloride under solvolytic conditions. *Bull. Korean Chem. Soc.* **2014**, *35*, 2285–2289.
80. Koo, I.S.; Bentley, T.W.; Kang, D.H.; Lee, I. Limitations of the transition state variation model. Part 2. Dual reaction channels for solvolyses of 2,4,6-trimethylbenzenesulfonyl chloride. *J. Chem. Soc. Perkin Trans. 2* **1991**, 175–179, doi:10.1039/P29910000175.

81. Koo, I.S.; Bentley, T.W.; Llewellyn, G.; Yang, K. Limitations of the transition state variation model. Part 3. Solvolyses of electron-rich benzenesulfonyl chlorides. *J. Chem. Soc. Perkin Trans. 2* **1991**, 1175–1179, doi:10.1039/A701140G.
82. Koo, I.S.; Lee, O-K.; Lee, I. Limitations of the transition state variation model (5) dual reaction channels for solvolysis of dansyl chloride. *Bull. Korean Chem. Soc.* **1992**, *13*, 395–398.
83. Bentley, T.W.; Garley, M.S. Correlations and predictions of solvent effects on reactivity: some limitations of multi-parameter equations and comparisons with similarity models based on one parameter. *J. Phys. Org. Chem.* **2006**, *19*, 341–349.
84. Bentley, T.W. A comprehensive N_+ scale of nucleophilicity in an equation including a Swain-Scott selectivity parameter. *J. Phys. Org. Chem.* **2013**, *26*, 977–982.
85. Kevill, D.N.; Anderson, S.W. An improved scale of solvent nucleophilicity based on solvolyses of S-methyldibenzothiophenium ion. *J. Org. Chem.* **1991**, *56*, 1845–1850.
86. Kevill, D.N. Development and uses of scales of solvent nucleophilicity. In *Advances in Quantitative Structure-Property Relationships*; Charton, M., Ed.; Jai Press: Greenwich, CT, USA, 1996; Volume 1, pp. 81–115.
87. D'Souza, M.J.; Mahon, B.P.; Kevill, D.N. Analysis of the nucleophilic solvation effects in isopropyl chlorothioformate solvolyses. *Int. J. Mol. Sci.* **2010**, *11*, 2597–2611.
88. Kevill, D.N.; D'Souza, M.J. Correlation of the rates of solvolysis of phenyl chloroformate. *J. Chem. Soc. Perkin Trans. 2* **1997**, 1721–1724, doi:10.1039/A701140G.
89. D'Souza, M.J.; Deol, J.K.; Pavey, M.T.; Kevill, D.N. Statistical methods for the investigation of solvolysis mechanisms illustrated by the chlorides of the carbomethoxy protecting groups NVOc and FMOC. *J. Anal. Methods Chem.* **2015**, doi:10.1155/2015/941638.
90. Yew, K.H.; Koh, H.J.; Lee, H.W.; Lee, I. Nucleophilic substitution reactions of phenyl chloroformates. *J. Chem. Soc. Perkin Trans. 2* **1995**, 2263–2268, doi:10.1039/P29950002263.
91. Ba-Saif, S.; Luthra, A.K.; Williams, A. Concertedness in acyl group transfer in solution: A single transition state for acetyl group transfer between phenolate ion nucleophiles. *J. Am. Chem. Soc.* **1987**, *109*, 6362–6368.
92. Renfrew, A.H.M.; Rettura, D.; Taylor, J.A.; Whitmore, J.M.J.; Williams, A. Stepwise vs. concerted mechanisms at trigonal carbon. Transfer of the 1,3,5-triazinyl group between aryl oxide ions in aqueous solution. *J. Am. Chem. Soc.* **1995**, *109*, 5484–5491.
93. Marlier, J.F. Heavy atom isotope effects on the alkaline hydrolysis of methyl formate the role of hydroxide ion in ester hydrolysis. *J. Am. Chem. Soc.* **1993**, *115*, 5953–5956.
94. Mata-Segreda, J.F. Hydroxide as general base in the saponification of ethyl acetate. *J. Am. Chem. Soc.* **2002**, *124*, 2259–2262.
95. Bentley, T.W.; Morris, P.J.; Taylor, J.A. Alkoxide-catalysed hydrolysis of a chlorotriazine in alcohol-water mixtures: A new indirect probe for hydroxide-catalysed hydrolysis. *J. Chem. Soc. Perkin Trans. 2* **2000**, 2171–2176, doi:10.1039/B005949H.