

LETTER TO EDITOR

CALCULATION OF LD₅₀ VALUES FROM THE METHOD OF MILLER AND TAINTER, 1944

Muhammad Akram Randhawa

Department of Pharmacology, College of Medicine, University of Dammam, Dammam, Saudi Arabia

Dear Editor,

Acute toxicity of a drug can be determined by the calculation of LD₅₀, i.e., the dose that will kill 50% of animals of a particular species. Recently, we reported the LD₅₀ of thymoquinone, an active principle of *Nigella sativa*, in mice and rats by the method described by Miller and Tainter, 1944.^{1,2} Since then some post-graduate students and investigators have asked us to explain the details of calculation of LD₅₀ by this method, as the original article, being very old, is not easily available in the literature. We also faced problem in the calculations, however, we found some details for the calculation of LD₅₀ in the Fundamentals of Experimental Pharmacology by Ghosh³, which also is not easy to obtain in many parts of the world. Therefore, few lines are written, on behalf of the other authors, to elaborate the calculation of LD₅₀ by the method of Miller and Tainter for the benefit of young researchers.

Estimation of the dose range and percentage of mortality

An approximate LD₅₀ can be initially determined as a pilot study by a so called 'staircase method' using a small number of animals (2 each dose) and increasing the doses of the drug. Five doses can be chosen for determination of LD₅₀ starting from no death to 100% mortality. In our study for estimation of LD₅₀ of thymoquinone, 5 doses were given intraperitoneally to 5 groups of rats, 10 in each group (Table-1).

The animals were observed for first 2 hours and then at 6th and 24th hour for any toxic symptoms. After 24 hours, the number of deceased rats was counted in each group and percentage of mortality calculated.

Table-1: Results of the lethal doses of thymoquinone for the determination of LD₅₀ after intraperitoneal injection in rats (n=10)

Group	Dose (mg/kg)	Log dose	% Dead	*Corrected %	Probits
1	25	1.4	0	2.5	3.04
2	50	1.7	40	40	4.75
3	75	1.88	70	70	5.52
4	100	2	90	90	6.28
5	150	2.18	100	97.5	6.96

*Corrected % Formula for 0 and 100 is given in the text.

Conversion of percentage mortalities to probits and calculation of LD₅₀

The percentage of animals that had died at each dose level is then transformed to probit (Table-2).

Table-2: Transformation of percentage mortalities to probits

%	0	1	2	3	4	5	6	7	8	9
0	-	2.67	2.95	3.12	3.25	3.36	3.45	3.52	3.59	3.66
10	3.72	3.77	3.82	3.87	3.92	3.96	4.01	4.05	4.08	4.12
20	4.16	4.19	4.23	4.26	4.29	4.33	4.36	4.39	4.42	4.45
30	4.48	4.50	4.53	4.56	4.59	4.61	4.64	4.67	4.69	4.72
40	4.75	4.77	4.80	4.82	4.85	4.87	4.90	4.92	4.95	4.97
50	5.00	5.03	5.05	5.08	5.10	5.13	5.15	5.18	5.20	5.23
60	5.25	5.28	5.31	5.33	5.36	5.39	5.41	5.44	5.47	5.50
70	5.52	5.55	5.58	5.61	5.64	5.67	5.71	5.74	5.77	5.81
80	5.84	5.88	5.92	5.95	5.99	6.04	6.08	6.13	6.18	6.23
90	6.28	6.34	6.41	6.48	6.55	6.64	6.75	6.88	7.05	7.33

The percentage dead for 0 and 100 are corrected before the determination of probits as under:

Corrected % Formula for 0 and 100% mortality³:

For 0% dead: 100(0.25/n)

For 100% dead: 100(n-0.25/n)

The probit values are plotted against log-doses and then the dose corresponding to probit 5, i.e., 50%, is found out (**Figure-1**). In the present case the Log LD₅₀ is 1.76 and LD₅₀= 57.54 mg/kg.

Calculation of Standard Error (SE) of LD₅₀

The SE of LD₅₀ can be calculated from the following formula:³

$$\text{Approx. SE of LD}_{50} = \frac{(\text{Log LD}_{84} - \text{Log LD}_{16}) \dots (a)}{\sqrt{2N}}$$

where N is number of animals in each group.

The probits of 84 and 16 from Table-1 are 5.99 and 4.01 (Approx. 6 and 4), respectively. The log-LD values for the probits 6 and 4 are obtained from the line on the graph in Figure-1, which in the present case are 1.96 and 1.58 and their antilog are 91.2 and 38.02. Using these values in formula (a) the SE of LD₅₀ is 11.9. Therefore, LD₅₀ of thymoquinone when given intraperitoneally is 57.54±11.9, with 95% confidence interval of 45.64–69.44.




Figure-1: Plot of log-doses versus probits from Table-1 for calculation of LD₅₀ of thymoquinone administered intraperitoneally.

REFERENCES

1. Al-Ali A, Alkhawajah A, Randhawa MA, Shaikh NA. Oral and intraperitoneal LD₅₀ of thymoquinone, an active principle of *Nigella sativa*, in mice and rats. J Ayub Med Coll Abbottabad 2008;20(2):25-7.
2. Miller LC, Tainter ML. Estimation of LD₅₀ and its error by means of log-probit graph paper. Proc Soc Exp Bio Med 1944;57:261.
3. Ghosh MN. In Statistical Analysis, Fundamentals of Experimental Pharmacology, 2nd ed, Scientific Book Agency Calcutta, 1984. pp187-9.

Address for Correspondence:

Dr. Muhammad Akram Randhawa, Professor of Pharmacology, College of Medicine, University of Dammam, Dammam, Saudi Arabia.

Email: mrakramsa@yahoo.co.uk