

Radiocarbon

1982

CALIBRATION OF RADIOCARBON DATES:

Tables based on the consensus data of the
Workshop on Calibrating the Radiocarbon Time Scale

JEFFREY KLEIN*, J C LERMAN**, P E DAMON**,
and E K RALPH*

A calibration is presented for conventional radiocarbon ages ranging from 10 to 7240 years BP and thus covering a calendric range of 8000 years from 6050 BC to AD 1950. Distinctive features of this calibration include 1) an improved data set consisting of 1154 radiocarbon measurements on samples of known age, 2) an extended range over which radiocarbon ages may be calibrated (an additional 530 years), 3) separate 95% confidence intervals (in tabular form) for six different radiocarbon uncertainties (20, 50, 100, 150, 200, 300 years), and 4) an estimate of the non-Poisson errors related to radiocarbon determinations, including an estimate of the systematic errors between laboratories.

INTRODUCTION

It is now quite generally accepted that "conventional" radiocarbon dates need to be "calibrated" because of temporal variations in the radiocarbon content of atmospheric carbon dioxide. The discovery of this phenomenon was made largely by the pioneering work of de Vries (1958; 1959) and Willis, Tauber, and Münnich (1960), and subsequently has been carried on by more than a dozen radiocarbon laboratories worldwide (for a review see Damon, Lerman, and Long, 1978). The assessment of these variations relies on the measurement of ^{14}C activity in samples of known age. Dendrochronologically dated wood has proved to be an ideal material for such measurements, and currently all radiocarbon calibrations are based on measurements of ^{14}C activity in wood. The longest chronology extant is that of the bristlecone pine, resulting from the efforts of Schulman (1956) and Ferguson (1969; 1970; 1972). It reaches continuously to 8681 years ago, and to 8580 years ago with sufficient material to allow radiocarbon dating. This work includes measurements on wood as old as 8000 years.

Many calibrations have appeared during the past 13 years (Suess, 1979; 1970a; 1967; Clark, 1980; 1979; 1975; McKerrell, 1975; Damon *et al.*, 1974; Ralph, Michael, and Han, 1973; Switsur, 1973; Michael

* Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania 19104.

** Laboratory of Isotope Geochemistry, Department of Geosciences, University of Arizona, Tucson, Arizona 85721

and Ralph, 1972; Clark and Renfrew, 1972; Damon, Long, and Wallick, 1972; Wendland and Donley, 1971; Lerman, Mook, and Vogel, 1970; Ralph and Michael, 1970; Stuiver and Suess, 1966). Although all reflect similar long-term changes in atmospheric radiocarbon concentrations, they differ significantly in their treatments of shorter period variations. This diversity of available calibrations and the apparently conflicting results obtained when calibrating dates using one in preference to another has resulted in a suspicion on the part of many archaeologists regarding calibration, in particular, and radiocarbon dating, in general. Consequently, in 1978, it was suggested to the USA National Science Foundation that it was time to attempt a consensus among the divergent efforts of the many laboratories then involved in calibration research. With this as a goal, a workshop was held in Tucson, Arizona in early 1979, entitled, "Workshop on the Calibration of the Radiocarbon Dating Time Scale" (Damon *et al.*, 1980; Michael and Klein, 1979). This work is largely the implementation of the decisions reached at that meeting.

The Workshop participants decided to provide a calibration table suitable for the calibration of individual or "single" radiocarbon dates. A "single radiocarbon date" is defined as any radiocarbon date that is not associated with another radiocarbon date by a tight, independently determined relative chronology. Such a chronology is exemplified by tree rings, where the number of intervening rings determines the relative ages of samples, and by stratified samples, where the rate of stratification is known independently of the radiocarbon ages of samples contained therein. Included in the category of "single radiocarbon dates" are series of dates from samples thought to be coeval, or series in which the temporal sequence, or even the relative ages of its members is unknown.

A second decision of the participants of the Workshop was to provide the "user" with a realistic assessment of the precision of calibrated dates. A consideration of many factors is necessary in the estimation of this precision. These include the precision with which the sample's activity has been measured, involving not just the "counting" statistics quoted by the measurement laboratory, but also an estimate of the true reproducibility of the measurement, *i.e.*, the degree to which a particular result can be repeated by the same laboratory or any other laboratory on subsequent measurements. In addition, there is the precision to which the calibration function is known near a particular calendric date. This depends on the quantity and quality of data used in the construction of the calibration. Finally, there is the "shape" of the calibration "curve" in the region in which it is being employed. This factor is often the most influential in determining the magnitude of the uncertainty of a calibrated date, and although its importance has been recognized for some time (Renfrew and Clark, 1974; Grey and Damon, 1970) it is often ignored in the routine calibration of dates.

These objectives were implemented by providing a range of calibrated dates, representing the 95% confidence interval, for each radiocarbon age of specified precision. An advantage of specifying an interval,

rather than a midpoint and uncertainty, hinges on the fact that many confidence intervals are asymmetrically related to the value obtained from simply calibrating the ^{14}C date without consideration of uncertainties.

THE DATA

This calibration is based on the ^{14}C activity measurements performed by the radiocarbon laboratories at the Universities of Arizona, Groningen, California at La Jolla, Pennsylvania, and Yale, on 1154 samples of dendrochronologically dated wood, principally *Pinus longaeva* and *Sequoia gigantea* (bristlecone pine and giant sequoia). The data set consisting, for the most part, of an updated version of previously published data (current data sets in preparation by individual laboratories), was prepared for the "Workshop on the Calibration of the Radiocarbon Dating Time Scale." Only measurements on samples of wood containing 20 or fewer rings were used in this work so as not to attenuate significantly through averaging, variations occurring on the time scale of the order of 100 years. Beyond this consideration, no selection of the data was undertaken.

As one of the principal objectives of this analysis has been to understand more fully the nature and causes of the variability of radiocarbon dates, the data were examined carefully for signs of non-random errors. Much to our surprise and despite previous findings to the contrary (Damon, Lerman, and Long, 1978; Clark, 1975; Damon, 1970), there is significant evidence of systematic differences between the laboratories represented. Of the five laboratories, one shows an average systematic difference of approximately six per mil, roughly 50 radiocarbon years, significant at less than the 1% level. The other four laboratories agree within experimental uncertainties. Independent comparisons with a sixth laboratory have resulted in similar conclusions (Stuiver, pers commun, 1981). Systematic differences were determined by calculating residuals of each data set with respect to the calibration function calculated on the combined data set. If no systematic differences had existed, then the sum of residuals would have been consistent with zero for all laboratories; it was not. A table of these differences was reported earlier (Klein *et al*, 1980), and is included here with slight modifications (see Table 1). Since it is unlikely that the systematic errors between other radiocarbon laboratories are, in general, less than those encountered here (International Study Group, submitted for publication), we decided to leave the data as they were and to include the uncertainty related to interlaboratory standardization within the calibration uncertainty.

CONSTRUCTION OF TABLES

Though the method used to construct this calibration has been outlined elsewhere (Ralph and Klein, 1979; Klein *et al*, 1980) and will be described in more detail in a forthcoming article, it is briefly described here. The procedure may be divided into three steps: a "global" regression which describes the long period (of the order of a few thousand

years) secular changes in the atmospheric ^{14}C concentration; a series of short term intervals called "shingles" which describe variations of a few hundred years; and finally, the construction of the table itself from the combination of these functions.

First, paired dendrochronologic ages and radiocarbon ages are scaled logarithmically so that each ranges over the interval $[-1,1]$. This is done to avoid the pathology common with polynomial regressions, namely the dominance of measurements at large values of the independent variable in the determination of the coefficients of the function. Next, each measurement is weighted by an estimate of the inverse of its variance. But, as it is widely accepted that the uncertainties quoted by radiocarbon laboratories, based only on counting statistics, are underestimates of the "true" variability, the laboratory uncertainties were increased under the following assumptions: 1) the additional sources of variance are independent of the Poisson error of the activity measurement; 2) this added variance is of approximately the same magnitude for samples of similar age; 3) these "extra" components increase with the age of the sample, as demonstrated by the poorer reproducibility of radiocarbon dates for older samples (Currie and Polach, 1980; Pearson *et al.*, 1977; Clark, 1975; Currie, 1972). Consequently, the "counting" variance was increased by an additive term which was allowed to be a slowly increasing function of the age of the sample, hence:

$$w_i = \frac{1}{\sigma^2_i + \left(40 + \frac{x_i}{150} \right)^2}$$

This has the effect of increasing the smallest error to approximately 60 years for samples less than 1000 years old, and to approximately 115 years for samples with ages greater than 6000 years. These figures compare favorably with the error estimates of Otlet *et al.* (1980), *viz.*: 50 years for samples less than 5000 years and 100 years for samples less than 10,000 years old, and the estimates of Clark (1975), *viz.*: 50 years for samples less than 3000 years and 95 years for samples with ages greater than 3000 years.

Finally, the weighted, scaled radiocarbon ages are least squares regressed against their calendric (dendrochronologic) ages using a polynomial basis to obtain the long period trend curve. Polynomials were chosen since 1) a sample's radiocarbon age is, to first order, linearly related to its chronologic age, and 2) though the difference between a sample's uncalibrated age and its true age is bounded, and described reasonably well by a sine function (Damon, Long, and Wallick, 1972; Houtermans, 1971), a polynomial fit is better.

With Fisher's F-test as a criterion, the "best fit" was determined to be a polynomial of order six. Because of its low order, this function is insensitive to short-period variations in the ^{14}C inventory and, for the most part, reflects variations resulting from changes in the earth's magnet-

ic field. (See, eg, Sternberg and Damon, 1979; Lingenfelter and Ramaty, 1970; Damon, 1970; Bucha, 1970; Lal and Venkatavaradan, 1970; Suess, 1970b.) This function and the data are plotted in Figure 1.

The second step involves a piecewise Fourier analysis of the residuals around the polynominal regression. A piecewise regression, *ie*, one that divides the data into a number of similar intervals instead of considering the data set as a whole, was adopted because of several distinctive features observed in the variations of atmospheric ^{14}C . Such characteristic changes are represented by the variations in ^{14}C concentration occurring during the Spörer, Maunder, and Wolf minima (Stuiver and Quay, 1980a; 1980b; Damon, Long and Grey, 1966); by those occurring in the sixth millennium BP (de Jong, Mook, and Becker, 1979; de Jong and Mook, 1980), and by the peaks at 200 years, 150 years, etc, observed in the power spectra of Fourier analyses performed by various investigators (Neftel, Oeschger, and Suess, 1981; Suess, 1980; Lazear, Damon, and Sternberg, 1980; Siegenthaler, Heimann, and Oeschger, 1980; Houtermans, 1971). Damon (1977) has noted that although characteristic periods appear in the spectral analyses of atmospheric ^{14}C , their phase relationships are different depending upon the section of the 8000-year record analyzed. With this in mind, it seemed prudent to divide the entire time period into short segments and consider the fluctuations individually in each. Consequently, the calendric time scale was divided into 28 shingles, each 500 years long, and each overlapping the previous and next shingle by 250 years (50% overlap each end, 100% overlap for the entire shingle). Two Fourier analyses were carried out to a minimum period of 65 or 110 years, depending on the number of measurements in the shingle. The minimum periods were chosen with consideration of the attenuation factors predicted by various models for changes in the atmospheric ^{14}C activity resulting from changes of various durations in the production-forcing function (Oeschger *et al*, 1975; Houtermans, 1966). Such models predict attenuation factors on the order of 25 times for variations in production lasting less than 100 years. The result of these procedures is shown in Figure 2.

Two analyses were performed in order to assess the effects of outlying points on the calibration function. The first analysis used the unmodified data base as described in the section on data, whereas the second analysis used a "winsorized" data set in which the residuals used for winsorization were taken with respect to the function calculated on the unmodified data. "Winsorization" is a process which reduces the effect of a few aberrant measurements by limiting the effect on the mean of a single outlying point to less than $\sim 2.56\text{s}/n$, where s is the standard error estimated from the fourth quintile of the variance of the data, and n is the number of points in the interval. Winsorization, as employed here, is described elsewhere (Dixon, 1960). Winsorization was used instead of a simple rejection of "outlying" points for the following reasons: 1) the maximum rate of change of the ^{14}C concentration is not certain, and although it appears that changes of the order of a few per mil per year seem to be




Fig 1.

the rule (Stuiver and Quay, 1980b; Burchuladze *et al.*, 1980; Lerman, 1970a; 1970b; Lerman *et al.*, 1969; Lerman, Mook, and Vogel, 1967), it seemed preferable not to establish an arbitrary criterion for the rejection of suspect measurements, and 2) in the assessment of the "true" errors associated with radiocarbon dates, the rejection of measurements with large residuals furthers the practice of underestimating the scatter in the data.

Another problem is caused by unequal residuals at the ends of the regression intervals (endpoint effects) and this was eliminated by using a cosine weighted average of the overlapping functions. This weight is equal to one in the center of the interval and zero at the ends, producing a final calibration function that is both continuous and differentiable.

The combined uncertainty of the calibration and the "true" uncertainty of the data are estimated by averaging the residuals of the data around the final calibration function, using the following formula:

$$\sigma_{\text{calib}} = \sum_{\text{shingle}}^n \{(y_i - \hat{y}_i)^2 - \sigma_i^2\} / (n - a)$$

where the $y(i)$ are winsorized, but the $\sigma(i)$ are the unmodified laboratory estimates of the measurement uncertainty, and n is the number of measurements in the 500-year interval. The assumption is that

$$\text{Var}(y - \hat{y}) = \text{Var}(y) - \text{Var}(\hat{y}),$$

which is the natural decomposition, assuming the independence of y

Fig 1. The composite "workshop data set" is plotted against the 6th order polynomial regressed on the logarithmically scaled data. Calendric age minus conventional radiocarbon age is the ordinate; the calendric age is the abscissa. Positive values represent radiocarbon ages that are too young (too recent) and, consequently, atmospheric concentrations were greater than that of the standard atmosphere of 1890. Laboratories are identified by the following symbols: \triangle = Arizona; \circ = Pennsylvania; \square = La Jolla; \times = Groningen; \diamond = Yale; $+$ = Uppsala. Error bars are laboratory estimates of uncertainties calculated from counting statistics. The equation of the trend line in logarithmically compressed coordinates is:

$$\tilde{y}_i = \sum_{n=0}^6 a_n \tilde{x}_i^n$$

where $\tilde{x}_i = \alpha \log_{10}(x_i) + \beta$,

x_i is the dendrochronologic age in years before AD 1975, and the various coefficients are defined by:

$\alpha = 0.774607$	$a_3 = -1.249500$
$\beta = -2.024200$	$a_4 = 0.641460$
$a_0 = -0.023469$	$a_5 = 0.591000$
$a_1 = 1.205700$	$a_6 = -0.344350$
$a_2 = 0.143050$	

The predicted radiocarbon age (in years before AD 1975 and with $T_{1/2} = 5730$ years), y_i , is obtained from \tilde{y}_i , using the formula:

$$y_i = \exp \left(\frac{\tilde{y}_i - \beta}{\alpha} \right)$$

and \hat{y} . In fact, this is not the case for linear regression which always leaves residuals correlated with the original data, but this correlation has little effect on the value of this procedure in determining the magnitude of the combined uncertainty of the calibration and the true measurement variability.

Finally, the calibration tables were derived from the composite calibration function and the combined error of the calibration and the quoted error of the radiocarbon date being calibrated. This was done by adding together the variance of the calibration (which includes not only the error of the calibration proper, but also an estimate of the non-Poisson error associated with a typical radiocarbon date) and the variance




Fig 2. Graphic representation of the period covered by the calibration tables. The ordinate is the conventional radiocarbon age in years BP (1950 used as origin, ages calculated using the 5568-year half-life); the abscissa is the calendric date in years AD-BC. The same data set as in Figure 1 is plotted, but the data here have been winsorized as described in the text. The function includes both the trend analysis and the Fourier analysis of the residuals around the trend. If conventional radiocarbon years were equivalent to calendric years, all the data would fall on the diagonal line; that they do not is readily apparent. The maximum deviations between uncalibrated conventional radiocarbon dates and calendric dates occur ca 5200 bc.

of the particular date. The square root of this "total" variance was added to and subtracted from the composite calibration function, producing an uncertainty band in ^{14}C activity representative of the 95% confidence interval for a single determination of the ^{14}C activity in a sample of given age. This was converted to an uncertainty interval in calibrated age by determining the range of calendric dates for which the ^{14}C age was consistent (see Figure 3). With the exception of the post-industrial period, multiple calibration intervals were found to be statistically unjustifiable. Consequently, after combining the variances associated with the calibration and those associated with an individual date, the bound-




Fig 3A. CALIBRATED DATE (AD-BC)

Fig 3A-G. Calibration limits (monotonic) for radiocarbon uncertainties of 20, 100, 200, and 300 years. The data are the same as in Figure 2. The error bands include both the error of the calibration and an estimate of the possible systematic differences between laboratories.

The 90% confidence intervals plotted in these graphs are intended primarily for users with multiple dates and will provide calibration intervals shorter than those obtained from the tables. To calibrate a radiocarbon date, first locate the radiocarbon age (BP 1950) on the ordinate (vertical axis), then draw a horizontal line (parallel to the abscissa) through the calibration curves. The projection onto the x-axis of the intersections of this line with the "curves" of appropriate uncertainty gives the calibrated range of the date. Note that each graph spans 1400 radiocarbon years.

ing functions were made monotonic in calendric age before the calibration interval was determined. In the final table, separate intervals are provided for radiocarbon uncertainties of 20, 50, 100, 150, 200, and 300 years. The table represents the 95% confidence interval for the calibrated date and covers the range from 7240 to 10 BP (radiocarbon years). If we assume that the source of the non-counting error is independent of the counting error and similar for samples of similar age, then the procedure described above properly accounts for this error as well.

For samples less than 1000 years BP (radiocarbon) supplementary tables are provided following the main tables. Asterisks in the main table indicate dates for which multiple intervals exist (see Figure 4). The intervals in the main table represent the extremes in range of the multiple intervals in the supplementary tables.




Fig 3B.

INSTRUCTIONS FOR USING CALIBRATION TABLES

The tables on the following pages are to be employed in the calibration of single radiocarbon dates. One enters the tables with a radiocarbon age (years BP, 5568-year, "Libby," half-life) and uncertainty, and leaves with a 95% confidence interval containing the "true", calendric date. The radiocarbon age, rounded to the nearest 10 years and calculated using the Libby half-life, determines the row in which the calibrated age is to be found; the uncertainty determines the columns. All dates within the table have been rounded to the nearest five years. Each radiocarbon age is calibrated to a single calendric range for ages greater than 1000 years, though multiple dates are possible for younger samples. Radiocarbon samples with uncertainties between the tabulated values should have their uncertainties rounded to the nearest tabulated value (see table footnote). Hence, a sample with a date of 1960 BP \pm 30 would have a calibrated interval of 145 BC to AD 210, whereas 1960 BP \pm 40 would range from 155 BC to AD 215. It will normally not be necessary to




Fig 3C.

interpolate between tabulated ages, as rounding dates to the nearest five years does not significantly affect the calendric interval obtained. Negative values in the body of the table represent BC dates; positive, AD dates; and $-1/1$ represents the transition year between 1 BC and AD 1 (omitted in the widely-adopted chronology of Dionysius Exiguus (ca 525)).

Occasionally, there are large "jumps" in the length of the calibration intervals as read from the table, eg, between 1920 and 1930 BP \pm 20 or between 1770 and 1780 BP \pm 150 years. These are caused by "flat" regions in the calibration, *i.e.*, periods when the ^{14}C in the atmosphere has decreased at a rate greater than 1.2 per mil per 10 years, allowing multiple calendric ages for a single ^{14}C activity. In other calibrations, these periods have often been handled by assigning several calendric dates to a single radiocarbon age. However, as described previously, the ability to distinguish these as separate periods vanishes when the uncertainties of the calibration and radiocarbon activity measurement are considered. Reference to the calibration graphs should clarify this.




Fig 3D.

CALIBRATION INTERVAL FOR SAMPLES WITH UNCERTAINTIES
GREATER THAN 300 YEARS

The following procedure should be employed in calibrating ages of samples with radiocarbon uncertainties greater than 300 years. First, 60 years should be subtracted from the uncertainty of the date to be calibrated. This is to remove the uncertainty of the calibration, which is automatically added into the range in the tables. Then, the resultant uncertainty should be added to and subtracted from the radiocarbon age of the sample, producing two ages which are looked up in the calibration table, under the columns headed by sigma=20 years. The calibration interval is formed from the extremes of the intervals obtained from the table. That is, the lower limit of the interval [older limit] is equal to the lower limit of the calibration interval for the radiocarbon age plus the modified uncertainty. Similarly, the upper limit [younger limit] is the upper limit of the calibration range for the radiocarbon age minus the modified uncertainty. As an example, consider the calibration of 3200 ± 400 years. First, subtract 60 years from 400 to obtain 340 years, which,




Fig 3E.

alternately added to and subtracted from the sample's radiocarbon age produces 3540 and 2860, respectively. Looking up the appropriate limits for these two ages, the interval 2110 to 875 BC is obtained.

CALIBRATION OF DATES BEYOND TABULATED VALUES

At this time, the only data set of sufficient quality to provide retrospective assessment of atmospheric ^{14}C to a precision suitable for calibration consists of measurements on wood. This is largely because of the stringent requirements for a sample suitable for this purpose. The sample must 1) be independently datable, 2) contain carbon that is reliably associated with atmospheric ^{14}C at the date of the sample formation, and 3) contain sufficient quantities of carbon for an accurate activity measurement.

Beyond the existing range of dendrochronologically dated wood, we must rely either on samples of inferior quality (shorter or less certain chronology, or of smaller size, frequently containing too little carbon to




Fig 3F.

obtain an accurate date), or on "secondary" sources that estimate the constancy of cosmic rays from the measurements of other radionuclides, or from the inferred strength of the earth's magnetic field from archaeomagnetism. The consensus of these sources suggests that the cosmic ray flux reaching the earth and producing ^{14}C has probably remained constant to within $\pm 10\%$ over the past 50,000 years or more (Vogel, 1980; Barbetti, 1980; Forman and Shaeffer, 1980; Stuiver, 1971). A 10% uncertainty in a radiocarbon concentration represents an 800-year uncertainty in age, regardless of the age of the sample. Consequently, the current "best estimate" of the date of a sample older than 8000 years BP is obtained by assuming a constant atmospheric concentration of the ^{14}C , and using the 5730 half-life to calculate the date. An uncertainty of 1000 years, or the measurement uncertainty quoted by the laboratory, whichever is larger, would constitute a reasonable estimate of the uncertainty for the calendric age of the sample.




Fig 3G.

CONCLUDING REMARKS

It is the intent of the participants of the Workshop that this should be the first in a series of "consensus" calibrations, updated as warranted by improvements in the data base. At present, 1132 measurements of ^{14}C activity have been made on samples of bristlecone pine, the maximum age of which is 8000 years BP. There are 60 samples of wood currently being dated by the radiocarbon laboratories at the Universities of Arizona, California at La Jolla, Pennsylvania, and Washington which will extend the calibration another 550 years. An additional piece of wood, containing 500 rings, is still undatable dendrochronologically but from preliminary radiocarbon measurements appears to be approximately 9000 to 10,000 years old (Ferguson and Graybill, 1981). Another piece of wood, containing only 200 rings, also antedates the present master chronology.




Fig 4. The first 1400 radiocarbon years. Similar to the graph in Figure 3A, here, however, the calibration function is not monotonic, and corresponds to the supplementary tables for the most recent 1000 years. Note that for several ages, multiple calendric intervals are possible for a single radiocarbon age.

Perhaps within the next few years, these pieces will be linked with the present 8681-year chronology extending it to beyond 11,000 years ago.

Still other chronologies are being developed both in this country and in Europe. The University of Washington has made activity measurements on nearly 2000 years of Douglas fir (Stuiver and Quay, 1980a; Stuiver and Quay, 1981). A second bristlecone chronology, 3200 years long, has been established on wood found in Nevada (Graybill, pers commun, 1982). Several floating chronologies are being developed in Europe (Becker, 1979; 1980; Beer *et al.*, 1979; Lambert and Orcel, 1979; Pilcher *et al.*, 1977) and it is likely that within the next few years it will be possible to connect them with existing recent chronologies. When this is done, they will be valuable in checking and reinforcing the USA chronologies. Even now, they are of some value after their age has been fixed using "wiggle matching" (see eg, Clark and Sowray, 1973) because these data sets are of high quality and their combined use (although not done in this work) with the calibration data set strengthens and reduces the errors of the current calibration.

ACKNOWLEDGMENTS

Special thanks are due J W Tukey and R M Clark for their many suggestions that have resulted in significant improvements in the algorithms originally presented to the Workshop. Thanks are due as well to C W Ferguson and the directors and staffs of the radiocarbon laboratories responsible for the activity measurements on these known-age samples, without which this work would not have been possible. We would also like to acknowledge the assistance of the operations staff at the University of Arizona's Computer Center, especially Jackie Dombrowski and Barry Shaede for their unfailing help, particularly during the preparations of the graphs presented here. The patience of those who have waited for the final publication of this calibration, even delaying their own work in some cases, also should not be forgotten. And finally, we would like to thank the National Science Foundation for their support of this publication through their Grant BNS-8022250, and for their support, since 1956, of the bristlecone-pine project, under the direction of C W Ferguson, through various grants, most recently EAR 78-04436 and EAR-8018687 and for their support of the USA laboratories involved in calibration-related research. The US Department of Energy should also be acknowledged for their recent support of the bristlecone-pine project with Contracts EE-78-A-28-3274 and DE-AC02-81EV10680.

REFERENCES

- Barbetti, Mike, 1980, Geomagnetic strength over the last 50,000 years and changes in atmospheric ^{14}C concentration: emerging trends, *in* Stuiver, Minze and Kra, Renee, eds, Internat'l radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 192-199.
- Becker, Bernd, 1979, Holocene tree ring series from southern central Europe for archaeological dating, radiocarbon calibration, and stable isotope analysis, *in* Berger, Rainer and Suess, H F, eds, Radiocarbon dating, Internat'l radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 554-565.
- , 1980, Tree-ring dating and radiocarbon calibration in south-central Europe, *in* Stuiver, Minze and Kra, Renee, eds, Internat'l radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 219-226.

- Beer, J, Giertz, V, Möll, M, Oeschger, Hans, Riesen, T, and Strahm, C, 1979, The contribution of the Swiss lake-dwellings to the calibration of radiocarbon dates, in Berger, Rainer and Suess, H E, eds, Radiocarbon dating, Internat radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 566-584.
- Bucha, V, 1970, Influence of the earth's magnetic field on radiocarbon dating, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 501-511.
- Burchuladze, A A, Pagava, S V, Povinec, P, Togonidze, G I, and Usačev, S, 1980, Radiocarbon variations with the 41-year solar cycle during the last century: Nature, v 287, p 320-322.
- Clark, R M, 1975, A calibration curve for radiocarbon dates: Antiquity, v 49, p 251-266.
- _____, 1979, Calibration, cross-validation and carbon-14, I: Royal Statistical Soc Jour, ser A, v 142, no. 1, p 47-62.
- _____, 1980, Calibration, cross-validation and carbon-14, II: Royal Statistical Soc Jour, ser A, v 143, no. 2, p 177-194.
- Clark, R M and Renfrew, Colin, 1972, A statistical approach to the calibration of floating tree-ring chronologies using radiocarbon dates: Archæometry, v 14, p 5-19.
- Clark, R M and Sowray, A, 1973, Further statistical methods for the calibration of floating tree-ring chronologies: Archæometry, v 15, no. 2, p 255-266.
- Currie, L A, 1972, The evaluation of radiocarbon measurements and inherent statistical limitations in age resolution, in Rafter, T A and Grant-Taylor, T, eds, Internat radiocarbon dating conf, 8th, Proc: Wellington, Royal Soc New Zealand, p 598-611.
- Currie, L A and Polach, H A, 1980, Exploratory analysis of the international radiocarbon cross-calibration data: consensus values and interlaboratory error, in Stuiver, Minze and Kra, Renée, eds, Internat radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 3, p 933-935.
- Damon, P E, 1970, Climatic versus magnetic perturbation of the atmospheric C-14 reservoir, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 571-593.
- _____, 1977, Solar induced variations of energetic particles at one AU, in White, O R, ed, The solar output and its variation: Boulder, Colorado, Colorado Assoc Univ Press, p 429-448.
- Damon, P E, Ferguson, C W, Long, Austin, and Wallick, E I, 1974, Dendrochronologic calibration of the radiocarbon time scale: Am Antiquity, v 39, p 350-366.
- Damon, P E, Lerman, J C, and Long, Austin, 1978, Temporal fluctuations of atmospheric ^{14}C : causal factors and implications: Ann Rev Earth Planetary Sci, p 457-494.
- Damon, P E, Lerman, J C, Long, Austin, Bannister, B, Klein, Jeffrey, and Linick, T W, 1980, Report on the workshop on the calibration of the radiocarbon dating time scale, in Stuiver, Minze and Kra, Renée, eds, Internat radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 3, p 947-949.
- Damon, P E, Long, Austin, and Grey, D C, 1966, Fluctuation of atmospheric ^{14}C during the last six millennia: Jour Geophys Research, v 71, no. 4, p 1055-1063.
- Damon, P E, Long, Austin, and Wallick, E I, 1972, Dendrochronologic calibration of the carbon-14 time scale, in Rafter, T A and Grant-Taylor, T, eds, Internat radiocarbon dating conf, 8th, Proc: Wellington, Royal Soc New Zealand, p 44-59.
- Dionysius Exiguus, ca AD 525, see Bickerman, E J, 1968, Chronology of the ancient world: Ithaca, New York, Cornell Univ Press.
- Dixon, W J, 1960, Simplified estimation from censored normal samples: Ann Math Statistics, v 31, p 385-391.
- Ferguson, C W, 1969, A 7104-year annual tree-ring chronology for bristlecone pine, *Pinus aristata*, from the White Mountains, California: Tree Ring Bull, v 29, no. 3-4, p 1-29.
- _____, 1970, Dendrochronology of bristlecone pine, *Pinus aristata*: establishment of a 7484-year chronology in the White Mountains of eastern-central California, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 237-245.
- _____, 1972, Dendrochronology of bristlecone pine prior to 4000 BC, in Rafter, T A and Grant-Taylor, T, eds, Internat radiocarbon dating conf, 8th, Proc: Wellington, Royal Soc New Zealand, p A1-A10.
- Ferguson, C W and Graybill, D A, 1981, Dendrochronology of bristlecone pine: Terminal rept NSF Grant EAR-78-04436 & DOE no. EE-78-A-28-3274.
- Forman, M A and Schaeffer, O A, 1980, Cosmic ray intensity over long time scales: Preprint.
- Grey, D C and Damon, P E, 1970, Sunspots and radiocarbon dating in the Middle Ages, in Berger, Rainer, ed, Scientific methods in medieval archaeology: Berkeley, Univ California Press, p 167-182.

- Houtermans, J C, 1966, On the quantitative relationship between geophysical parameters and the natural radiocarbon inventory: *Zeitschr Physik*, v 193, p 1-12.
- 1971, Geophysical interpretations of the bristlecone radiocarbon measurements using the method of Fourier analysis of unequally spaced data: Doctoral dissert, Univ Bern.
- Jong, A F M de and Mook, W G, 1980, Medium-term atmospheric ^{14}C variations, in Stuiver, Minze and Kra, Renee, eds, Internatl radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 267-272.
- Jong, A F M de, Mook, W G, and Becker, Bernd, 1979, Confirmation of the Suess wiggles: 3200-3700 BC: *Nature*, v 280, p 48-49.
- Klein, Jeffrey, Lerman, J C, Damon, P E, and Linick, T W, 1980, Radiocarbon concentration in the atmosphere: 8000-year record of variations in tree rings: first results of a USA workshop, in Stuiver, Minze and Kra, Renee, eds, Internatl radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 3, p 950-961.
- Lal, Devendra and Venkatavaradan, V S, 1970, Analysis of the causes of C-14 variations in the atmosphere, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 549-569.
- Lambert, G and Orcel, C, 1979, Dendrochronology of Neolithic settlements in western Switzerland: new possibility for prehistoric calibration, in Berger, Rainer and Suess, H E, eds, Radiocarbon dating, Internatl radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 585-590.
- Lazcar, G, Damon, P E, and Sternberg, R S, 1980, The concept of dc gain in modeling secular variations in atmospheric ^{14}C , in Stuiver, Minze and Kra, Renee, eds, Internatl radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 318-327.
- Lerman, J C, 1970a, General discussion of magnitudes of $^{14}\text{C}/^{12}\text{C}$ variations (discussion), in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 121-126.
- 1970b, General discussion of the causes of $^{14}\text{C}/^{12}\text{C}$ variations (discussion), in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 327-333.
- Lerman, J C, Mook, W G, and Vogel, J C, 1967, Effect of the Tunguska meteor and sunspots on radiocarbon in tree rings: *Nature*, v 216, p 990-991.
- 1970, C-14 in tree rings from different localities, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 275-301.
- Lerman, J C, Mook, W G, Vogel, J C, and Waard, H de, 1969, Carbon-14 in Patagonian tree rings: *Science*, v 165, p 1123-1125.
- Lingenfelter, R E and Ramaty, R, 1970, Astrophysical and geophysical variations in ^{14}C production, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 513-537.
- McKerrell, H, 1975, Correction procedures for C-14 dates, in Watkins, T, ed, Radiocarbon: calibration and prehistory: Edinburgh, Edinburgh Univ Press, p 47-100.
- Michael, H N and Klein, Jeffrey, 1979, An international calibration for radiocarbon dates: *MASCA Jour*, v 1, no. 2, p 56-57.
- Michael, H N and Ralph, E K, 1972, Discussion of radiocarbon dates obtained from precisely dated sequoia and bristlecone pine samples, in Rafter, T A and Grant-Taylor, T, eds, Internatl radiocarbon dating conf, 8th, Proc: Wellington, Royal Soc New Zealand, p 28-43.
- NefTEL, Albrecht, Oeschger, Hans, and Suess, H E, 1981, Secular non-random variations of cosmogenic carbon-14 in the terrestrial atmosphere: *Earth and Planetary Sci Letters*, v 56, p 127-147.
- Oeschger, Hans, Siegenthaler, Ulrich, Schotterer, Ulrich and Gugelmann, A, 1975, A box diffusion model to study the carbon dioxide exchange in nature: *Tellus*, v 27, p 168-192.
- Otlet, R L, Walker, A J, Hewson, A D, and Burleigh, Richard, 1980, ^{14}C interlaboratory comparisons in the UK: experiment design, preparation and preliminary results, in Stuiver, Minze and Kra, Renee, eds, Internatl radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 3, p 936-946.
- Pearson, G W, Pilcher, J R, Baillie, M G L, and Hillam, J, 1977, Absolute radiocarbon dating using a low altitude European tree-ring calibration: *Nature*, v 270, p 25-28.
- Pilcher, J R, Hillam, J, Baillie, M G L, and Pearson, G W, 1977, A long subfossil oak tree-ring chronology from the north of Ireland: *New Phytol*, v 79, p 713-729.
- Ralph, E K and Klein, Jeffrey, 1979, Composite computer plots of ^{14}C dates for tree-ring dated bristlecone pines and sequoias, in Berger, Rainer and Suess, H E, eds, Radiocarbon dating, Internatl radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, 545-553.

- Ralph, E K and Michael, H N, 1970, MASCA radiocarbon dates for sequoia and bristlecone pine samples, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 619-624.
- Ralph, E K, Michael, H N, and Han, M C, 1973, Radiocarbon dates and reality: MASCA Newsletter, v 9, p 1-20.
- Renfrew, Colin and Clark, R M, 1974, Problems of the radiocarbon calendar and its calibration: Archaeometry, v 16, p 5-18.
- Schulman, E, 1956, Dendroclimatic changes in semiarid America: Tucson, Univ Arizona Press.
- Siegenthaler, Ulrich, Heimann, Martin, and Oeschger, Hans, 1980, ^{14}C variations caused by changes in the global carbon cycle, in Stuiver, Minze and Kra, Renee, eds, Internat radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 177-191.
- Sternberg, R S and Damon, P E, 1979, Sensitivity of radiocarbon fluctuations and inventory to geomagnetic and reservoir parameters, in Berger, Rainer and Suess, H E, eds, Radiocarbon dating, Internat radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 691-717.
- Stuiver, Minze, 1971, Evidence for the variation of atmospheric ^{14}C content in the Late Quaternary, in Turekian, K K, ed, The late Cenozoic glacial ages: New Haven, Yale Univ Press, p 57-70.
- Stuiver, Minze and Quay, P D, 1980a, Changes in atmospheric ^{14}C attributed to a variable sun: Science, v 207, p 11-19.
- 1980b, Patterns of atmospheric ^{14}C changes, in Stuiver, Minze and Kra, Renee, eds, Internat radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 166-176.
- 1981, Atmospheric ^{14}C changes resulting from fossil fuel CO_2 release and cosmic ray flux variability: Earth Planetary Sci Letters, v 53, p 349-362.
- Stuiver, Minze and Suess, H E, 1966, On the relationship between radiocarbon dates and true ages: Radiocarbon, v 8, p 534-540.
- Suess, H E, 1967, Bristlecone pine calibration of the radiocarbon time scale from 4100 BC to 1500 BC, in Radioactive dating and methods of low-level counting: Vienna, IAEA, p 143-151.
- 1970a, Bristlecone pine calibration of the radiocarbon time-scale 5200 BC to the present, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 303-311.
- 1970b, The three causes of the secular C-14 fluctuations, their amplitudes and time constants, in Olsson, I U, ed, Radiocarbon variations and absolute chronology, Nobel symposium, 12th, Proc: New York, John Wiley & Sons, p 595-604.
- 1979, A calibration table for conventional radiocarbon dates, in Berger, Rainer and Suess, H E, eds, Radiocarbon dating, Internat radiocarbon conf, 9th, Proc: Berkeley/Los Angeles, Univ California Press, p 777-785.
- 1980, The radiocarbon record in tree rings of the last 8000 years, in Stuiver, Minze and Kra, Renee, eds, Internat radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 200-209.
- Switsur, V R, 1973, The radiocarbon calendar recalibrated: Antiquity, v 47, p 131-137.
- Vries, Hessel de, 1958, Variation in the concentration of radiocarbon with time and location on earth: Koninkl Nederlandse Akad Wetensch Proc, ser B, v 61, p 94-102.
- 1959, Measurement and use of natural radiocarbon, in Abelson, P H, ed, Researches in geochemistry: New York, John Wiley & Sons, p 169-189.
- Vogel, J C, 1980, Accuracy of the radiocarbon time scale beyond 15,000 BP, in Stuiver, Minze and Kra, Renee, eds, Internat radiocarbon conf, 10th, Proc: Radiocarbon, v 22, no. 2, p 210-218.
- Wendland, W M and Donley, D L, 1971, Radiocarbon-calendar age relationship: Earth Planetary Sci Letters, v 2, p 135-139.
- Willis, E H, Tauber, Henrik and Münnich, K O, 1960, Variations in the atmospheric radiocarbon concentration over the past 1300 years: Radiocarbon, v 2, p 1-4.

TABLE 1
SYSTEMATIC DIFFERENCES OBSERVED BETWEEN LABORATORIES

Laboratory	Average deviation from mean ($\Delta\%_{\epsilon}$)
Arizona (A)	+3.0 \pm 1.7
Groningen (GrN)	+2.7 \pm 1.5
La Jolla (Lj)	-3.2 \pm 1.1
Pennsylvania (P)	+3.4 \pm 2.5
Yale (Y)	+3.2 \pm 2.0

TABLE 2
MAIN CALIBRATION TABLES (P 124)
(See instructions in text and in footnote below)

Look up under nearest tabulated value radiocarbon dates with uncertainties between tabulated values, hence:

for sigma =	look up under:
0 — 35	$\sigma = 20$
36 — 75	$\sigma = 50$
76 — 125	$\sigma = 100$
126 — 175	$\sigma = 150$
176 — 250	$\sigma = 200$
251 — 350	$\sigma = 300$
> 350	use the procedure described in the text

* in body of table indicates multiple calibrated ranges exist for these dates. See supplementary tables.

RADIOCARBON AGE (BP)	5568 HALF-LIFE	CALIBRATED RANGES (95% CONFIDENCE)			SIGMA=150 YRS.	SIGMA=200 YRS.	SIGMA=300 YRS.
		SIGMA= 20 YRS.	SIGMA= 50 YRS.	SIGMA=100 YRS.			
7240	-6545	-5625	-5615	-5595	-6630	-5565	-6825
7230	-6535	-5615	-5610	-5590	-6615	-5555	-6815
7220	-6520	-5605	-5630	-5580	-6605	-5550	-6820
7210	-6510	-5595	-6520	-5590	-6570	-5540	-6800
7200	-6495	-5590	-6505	-6535	-6580	-5565	-6790
7190	-6485	-5580	-6495	-6575	-6520	-5555	-6775
7180	-6470	-6470	-6480	-6565	-6510	-5545	-6750
7170	-6460	-5565	-6470	-5555	-6495	-5535	-6740
7160	-6445	-5555	-6455	-5550	-6485	-5530	-6725
7150	-6435	-5545	-6445	-5540	-6470	-5525	-6715
7140	-6420	-5540	-6430	-5530	-6460	-5515	-6700
7130	-6410	-5530	-6420	-5525	-6445	-5510	-6690
7120	-6395	-5525	-6405	-5520	-6435	-5505	-6685
7110	-6385	-5515	-6395	-5510	-6420	-5500	-6675
7100	-6370	-5510	-6380	-5505	-6410	-5490	-6665
7090	-6360	-5505	-6370	-5500	-6395	-5485	-6650
7080	-6345	-5500	-6355	-5495	-6385	-5480	-6640
7070	-6335	-5495	-6345	-5490	-6370	-5475	-6625
7060	-6320	-5490	-6330	-5485	-6360	-5470	-6615
7050	-6310	-5485	-6320	-5480	-6345	-5465	-6600
7040	-6295	-5480	-6305	-5475	-6335	-5460	-6590
7030	-6285	-5475	-6295	-5470	-6320	-5455	-6580
7020	-6270	-5470	-6280	-5465	-6310	-5450	-6565
7010	-6260	-5465	-6270	-5460	-6295	-5450	-6550
7000	-6245	-5460	-6255	-5455	-6285	-5445	-6540
6990	-6235	-5455	-6245	-5450	-6270	-5440	-6530
6980	-6220	-5450	-6230	-5445	-6260	-5435	-6520
6970	-6210	-5445	-6220	-5440	-6245	-5430	-6510
6960	-6195	-5440	-6205	-5435	-6235	-5425	-6500
6950	-6185	-5435	-6195	-5435	-6220	-5420	-6490
6940	-6170	-5430	-6180	-5430	-6210	-5415	-6485
6930	-6160	-5430	-6170	-5425	-6195	-5410	-6475
6920	-6145	-5425	-6155	-5420	-6185	-5410	-6465
6910	-6135	-5420	-6145	-5415	-6170	-5405	-6455
6900	-6120	-5415	-6130	-5410	-6160	-5400	-6440

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=300 YRS.							
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA=150 YRS.								
5568 HALF-LIFE												
6890	-6110	-5410	-6120	-5405	-6145	-5395	-6190	-5375	-6250	-5345	-6390	-5270
6880	-6095	-5405	-6105	-5405	-6135	-5390	-6180	-5370	-6235	-5340	-6375	-5265
6870	-6085	-5405	-6095	-5400	-6120	-5385	-6165	-5365	-6225	-5335	-6365	-5260
6860	-6070	-5400	-6080	-5395	-6110	-5380	-6155	-5360	-6210	-5320	-6350	-5255
6850	-6060	-5395	-6070	-5390	-6095	-5375	-6140	-5355	-6200	-5325	-6340	-5250
6840	-6045	-5390	-6055	-5385	-6085	-5370	-6130	-5350	-6185	-5320	-6325	-5245
6830	-6035	-5385	-6045	-5380	-6070	-5365	-6115	-5345	-6175	-5310	-6315	-5240
6820	-6025	-5380	-6030	-5375	-6060	-5365	-6105	-5340	-6160	-5305	-6300	-5235
6810	-6010	-5375	-6020	-5375	-6045	-5360	-6090	-5335	-6150	-5300	-6290	-5230
6800	-6000	-5375	-6005	-5370	-6035	-5355	-6080	-5330	-6135	-5295	-6275	-5225
6790	-5985	-5370	-5995	-5365	-6020	-5350	-6065	-5325	-6125	-5290	-6265	-5220
6780	-5975	-5365	-5985	-5360	-6010	-5345	-6055	-5320	-6110	-5285	-6250	-5215
6770	-5965	-5360	-5970	-5355	-6000	-5340	-6040	-5315	-6100	-5280	-6240	-5210
6760	-5950	-5355	-5960	-5350	-5985	-5335	-6030	-5305	-6085	-5275	-6225	-5210
6750	-5940	-5350	-5950	-5345	-5975	-5330	-6015	-5300	-6075	-5270	-6215	-5205
6740	-5930	-5345	-5935	-5340	-5965	-5325	-6005	-5295	-6060	-5265	-6200	-5200
6730	-5915	-5340	-5925	-5335	-5950	-5320	-5995	-5290	-6050	-5260	-6190	-5195
6720	-5905	-5335	-5915	-5330	-5940	-5315	-5980	-5285	-6035	-5255	-6175	-5190
6710	-5895	-5330	-5900	-5325	-5930	-5310	-5970	-5280	-6025	-5250	-6165	-5190
6700	-5885	-5325	-5890	-5320	-5915	-5305	-5960	-5275	-6010	-5245	-6150	-5185
6690	-5875	-5320	-5880	-5315	-5905	-5295	-5945	-5270	-6000	-5240	-6140	-5105
6680	-5860	-5315	-5870	-5310	-5895	-5290	-5935	-5265	-5990	-5235	-6125	-5095
6670	-5850	-5310	-5860	-5305	-5885	-5285	-5925	-5260	-5975	-5230	-6115	-5090
6660	-5840	-5305	-5845	-5300	-5870	-5280	-5910	-5255	-5965	-5225	-6100	-5080
6650	-5830	-5300	-5835	-5295	-5860	-5275	-5900	-5250	-5955	-5220	-6090	-5075
6640	-5820	-5295	-5825	-5290	-5850	-5270	-5890	-5245	-5940	-5215	-6075	-5070
6630	-5810	-5290	-5815	-5285	-5840	-5265	-5880	-5240	-5930	-5210	-6065	-5065
6620	-5795	-5285	-5805	-5280	-5830	-5260	-5865	-5235	-5920	-5210	-6050	-5055
6610	-5785	-5280	-5795	-5275	-5820	-5255	-5855	-5230	-5905	-5205	-6040	-5050
6600	-5775	-5275	-5785	-5270	-5805	-5250	-5845	-5225	-5895	-5200	-6025	-5055
6590	-5765	-5270	-5775	-5260	-5795	-5245	-5835	-5220	-5920	-5195	-6015	-5015
6580	-5755	-5265	-5760	-5255	-5785	-5240	-5825	-5215	-5875	-5190	-6005	-5005
6570	-5745	-5260	-5750	-5255	-5775	-5235	-5815	-5210	-5865	-5190	-5990	-4995
6560	-5735	-5255	-5740	-5250	-5765	-5230	-5805	-5210	-5850	-5185	-5980	-4985
6550	-5725	-5250	-5730	-5245	-5785	-5225	-5790	-5205	-5840	-5105	-5970	-4975

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=200 YRS.	SIGMA=300 YRS.
		SIGMA= 50 YRS.	SIGMA=100 YRS.	SIGMA=150 YRS.		
5568 HALF-LIFE						
6540	-5715	-5245	-5720	-5745	-5220	-5780
6530	-5705	-5235	-5710	-5735	-5220	-5770
6520	-5695	-5235	-5700	-5725	-5215	-5760
6510	-5685	-5230	-5690	-5725	-5210	-5750
6500	-5675	-5225	-5680	-5720	-5205	-5740
6490	-5665	-5220	-5670	-5715	-5200	-5730
6480	-5655	-5215	-5660	-5710	-5195	-5720
6470	-5645	-5215	-5650	-5675	-5195	-5710
6460	-5635	-5210	-5640	-5620	-5195	-5700
6450	-5625	-5205	-5630	-5620	-5185	-5655
6440	-5615	-5200	-5620	-5615	-5180	-5645
6430	-5605	-5195	-5610	-5610	-5165	-5610
6420	-5595	-5195	-5605	-5610	-5165	-5605
6410	-5590	-5190	-5595	-5615	-5160	-5605
6400	-5580	-5185	-5585	-5615	-5160	-5600
6390	-5570	-5180	-5575	-5610	-5155	-5620
6380	-5560	-5160	-5565	-5610	-5150	-5615
6370	-5555	-5155	-5560	-5605	-5145	-5610
6360	-5545	-5150	-5550	-5600	-5140	-5605
6350	-5535	-5085	-5540	-5605	-5075	-5605
6340	-5525	-5080	-5530	-5607	-5075	-5610
6330	-5515	-5070	-5520	-5605	-5070	-5605
6320	-5500	-5065	-5510	-5605	-5065	-5600
6310	-5485	-5060	-5500	-5600	-5055	-5595
6300	-5470	-5050	-5485	-5605	-5035	-5590
6290	-5445	-5040	-5465	-5602	-5020	-5595
6280	-5420	-5025	-5440	-5601	-5010	-5590
6270	-5400	-5010	-5420	-5600	-5000	-5585
6260	-5380	-5000	-5395	-5600	-4990	-5580
6250	-5365	-4995	-5380	-5600	-4985	-5575
6240	-5355	-4985	-5365	-5605	-4980	-5570
6230	-5345	-4980	-5355	-5607	-4975	-5565
6220	-5335	-4970	-5345	-5600	-4970	-5560
6210	-5325	-4965	-5335	-5600	-4965	-5555
6200	-5320	-4960	-5325	-5600	-4960	-5550

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE)			FOR MEASUREMENT UNCERTAINTIES OF		
		SIGMA= 50 YRS.	SIGMA=100 YRS.	SIGMA=150 YRS.	SIGMA=200 YRS.	SIGMA=300 YRS.	
5568 HALF-LIFE							
6190	-4950	-5320	-4945	-5350	-4920	-5405	-4890
6180	-4945	-5310	-4940	-5340	-4920	-5390	-4765
6170	-4940	-5305	-4935	-5330	-4915	-5380	-4755
6160	-4935	-5300	-4930	-5325	-4910	-5365	-4745
6150	-4930	-5290	-4925	-5315	-4905	-5355	-4740
6140	-4920	-5280	-4920	-5310	-4900	-5345	-4735
6130	-4925	-5275	-4915	-5305	-4895	-5340	-4735
6120	-4920	-5270	-4915	-5295	-4890	-5330	-4730
6110	-4915	-5265	-4910	-5290	-4765	-5325	-4725
6100	-4910	-5260	-4910	-5285	-4755	-5315	-4715
6090	-4905	-5255	-4900	-5280	-4750	-5310	-4705
6080	-4905	-5250	-4895	-5275	-4740	-5305	-4680
6070	-4900	-5245	-4900	-5250	-4720	-5300	-4675
6060	-4895	-5240	-4895	-5245	-4730	-5295	-4670
6050	-4890	-5235	-4890	-5240	-4770	-5260	-4670
6040	-4885	-5230	-4885	-5235	-4755	-5280	-4665
6030	-4880	-5225	-4880	-5230	-4750	-5250	-4650
6020	-4775	-5220	-4775	-5225	-4745	-5245	-4645
6010	-4750	-5215	-4750	-5220	-4740	-5240	-4640
6000	-4745	-5210	-4745	-5215	-4735	-5235	-4635
5990	-4740	-5205	-4740	-5205	-4730	-5230	-4630
5980	-4735	-5200	-4735	-5200	-4725	-5225	-4625
5970	-4730	-5195	-4730	-5195	-4720	-5210	-4620
5960	-4725	-5190	-4725	-5190	-4710	-5205	-4615
5950	-4700	-5185	-4700	-5185	-4690	-5195	-4610
5940	-4685	-5180	-4685	-5180	-4675	-5205	-4605
5930	-4680	-5180	-4680	-5180	-4675	-5200	-4600
5920	-4665	-5175	-4675	-5180	-4670	-5195	-4595
5910	-4660	-5075	-4675	-5175	-4665	-5190	-4590
5900	-4660	-5065	-4670	-5075	-4665	-5185	-4585
5890	-4570	-5195	-4575	-5205	-4560	-5230	-4440
5880	-4565	-5180	-4575	-5200	-4560	-5225	-4435
5870	-4565	-5175	-4575	-5190	-4555	-5220	-4435
5860	-4560	-5015	-4565	-5075	-4540	-5190	-4430
5850	-4560	-4990	-4560	-5020	-4555	-5070	-4430

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=300 YRS.					
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.						
5568 HALF-LIFE										
5840	-4965	-4555	-4995	-4445	-5185	-4425	-5210	-4405	-5280	-4120
5830	-4945	-4555	-4975	-4550	-5055	-4445	-5180	-4425	-5205	-4115
5820	-4935	-4550	-4950	-4550	-5040	-4440	-5175	-4420	-5200	-4110
5810	-4925	-4550	-4940	-4540	-5020	-4435	-5075	-4420	-5195	-3990
5800	-4915	-4545	-4930	-4460	-5000	-4435	-5070	-4415	-5195	-3980
5790	-4905	-4545	-4920	-4450	-4980	-4430	-5065	-4410	-5190	-3970
5780	-4895	-4460	-4910	-4445	-4960	-4430	-5055	-4410	-5185	-3965
5770	-4880	-4450	-4905	-4440	-4945	-4425	-5050	-4405	-5180	-3960
5760	-4855	-4445	-4895	-4440	-4935	-4420	-5035	-4405	-5175	-3950
5750	-4725	-4440	-4880	-4435	-4925	-4420	-5010	-4400	-5080	-4150
5740	-4720	-4440	-4855	-4430	-4920	-4415	-4900	-4395	-5075	-4140
5730	-4730	-4435	-4730	-4430	-4910	-4415	-4975	-4390	-5065	-4135
5720	-4715	-4430	-4725	-4425	-4905	-4410	-4960	-4380	-5060	-4130
5710	-4710	-4430	-4720	-4425	-4895	-4410	-4945	-4370	-5050	-4120
5700	-4700	-4425	-4715	-4420	-4885	-4405	-4935	-4355	-5040	-4120
5690	-4695	-4425	-4710	-4420	-4870	-4400	-4930	-4155	-5020	-4115
5680	-4690	-4420	-4705	-4415	-4850	-4400	-4920	-4145	-5000	-4110
5670	-4680	-4420	-4700	-4415	-4850	-4400	-4915	-4140	-4985	-4105
5660	-4565	-4415	-4690	-4410	-4725	-4395	-4905	-4135	-4970	-3985
5650	-4565	-4415	-4685	-4405	-4720	-4385	-4900	-4130	-4955	-3975
5640	-4560	-4410	-4670	-4405	-4715	-4365	-4890	-4125	-4945	-3970
5630	-4560	-4405	-4665	-4400	-4710	-4155	-4880	-4120	-4935	-3960
5620	-4555	-4405	-4560	-4395	-4705	-4150	-4865	-4115	-4920	-3950
5610	-4555	-4400	-4560	-4390	-4700	-4140	-4845	-4110	-4915	-3940
5600	-4550	-4395	-4555	-4380	-4690	-4135	-4730	-3995	-4915	-3930
5590	-4550	-4390	-4555	-4170	-4685	-4130	-4725	-3980	-4910	-3930
5580	-4545	-4380	-4550	-4155	-4670	-4125	-4720	-3975	-4900	-3925
5570	-4545	-4160	-4550	-4145	-4565	-4120	-4715	-3965	-4885	-3915
5560	-4540	-4140	-4545	-4130	-4560	-4110	-4705	-3955	-4870	-3905
5550	-4535	-4135	-4540	-4125	-4555	-3990	-4700	-3945	-4855	-3930
5540	-4535	-4130	-4540	-4120	-4550	-3980	-4690	-3935	-4855	-3920
5530	-4530	-4130	-4540	-4120	-4550	-3970	-4685	-3930	-4725	-3895
5520	-4525	-4125	-4535	-4115	-4550	-3965	-4670	-3920	-4720	-3890
5510	-4520	-4120	-4530	-4110	-4550	-3960	-4565	-3915	-4715	-3890
5500	-4440	-4115	-4530	-3995	-4545	-3960	-4565	-3915	-4785	-3875

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE)			FOR MEASUREMENT UNCERTAINTIES OF		
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.	SIGMA= 200 YRS.	SIGMA= 300 YRS.	
5568 HALF-LIFE							
5490	-4435	-4110	-4520	-3980	-4545	-3950	-4560
5480	-4435	-3985	-4410	-3975	-4540	-3945	-4560
5470	-4430	-3975	-4415	-3965	-4540	-3935	-4555
5460	-4430	-3970	-4415	-3960	-4535	-3930	-4555
5450	-4425	-3965	-4410	-3955	-4530	-3920	-4550
5440	-4425	-3955	-4410	-3945	-4525	-3915	-4550
5430	-4420	-3950	-4415	-3940	-4520	-3910	-4545
5420	-4420	-3940	-4425	-3930	-4440	-3905	-4545
5410	-4415	-3935	-4420	-3925	-4430	-3900	-4540
5400	-4415	-3925	-4420	-3915	-4435	-3895	-4540
5390	-4410	-3920	-4415	-3910	-4430	-3895	-4535
5380	-4410	-3915	-4415	-3905	-4430	-3890	-4530
5370	-4405	-3910	-4410	-3900	-4425	-3890	-4525
5360	-4405	-3905	-4410	-3900	-4420	-3885	-4520
5350	-4400	-3900	-4405	-3895	-4420	-3880	-4510
5340	-4395	-3895	-4405	-3890	-4420	-3880	-4515
5330	-4390	-3895	-4400	-3890	-4415	-3875	-4510
5320	-4385	-3890	-4395	-3885	-4415	-3875	-4505
5310	-4380	-3885	-4390	-3885	-4410	-3870	-4500
5300	-4370	-3885	-4385	-3880	-4405	-3870	-4500
5290	-4360	-3880	-4380	-3880	-4405	-3865	-4495
5280	-4350	-3880	-4370	-3875	-4400	-3860	-4490
5270	-4345	-3875	-4355	-3870	-4395	-3855	-4485
5260	-4335	-3875	-4350	-3870	-4395	-3800	-4480
5250	-4330	-3870	-4340	-3865	-4390	-3795	-4475
5240	-4325	-3865	-4335	-3860	-4380	-3790	-4470
5230	-4320	-3865	-4330	-3860	-4370	-3790	-4465
5220	-4305	-3860	-4325	-3805	-4360	-3785	-4460
5210	-4145	-3855	-4315	-3800	-4350	-3780	-4455
5200	-4140	-3800	-4300	-3795	-4345	-3780	-4450
5190	-4135	-3795	-4145	-3790	-4335	-3775	-4415
5180	-4130	-3790	-4140	-3785	-4330	-3690	-4415
5170	-4130	-3785	-4135	-3785	-4325	-3685	-4410
5160	-4125	-3785	-4130	-3780	-4315	-3680	-4410
5150	-4120	-3780	-4125	-3775	-4305	-3675	-4405

RADIOCARBON AGE (BP)	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA= 150 YRS.	SIGMA= 200 YRS.	SIGMA= 300 YRS.
	SIGMA= 20 YRS.	SIGMA= 50 YRS.	SIGMA= 100 YRS.			
5568 HALF-LIFE						
5140	-4115	-3780	-4120	-3775	-4140	-3675
5130	-4110	-3775	-4120	-4135	-4130	-3670
5120	-4110	-3695	-4115	-3685	-4130	-3670
5110	-4105	-3685	-4110	-3680	-4130	-3665
5100	-4100	-3680	-4105	-3675	-4125	-3665
5090	-4095	-3680	-4100	-3675	-4120	-3660
5080	-4090	-3675	-4095	-3670	-4115	-3660
5070	-4080	-3670	-4090	-3670	-4110	-3660
5060	-3955	-3670	-4085	-3665	-4105	-3655
5050	-3945	-3670	-4075	-3665	-4100	-3655
5040	-3935	-3665	-3950	-3660	-4100	-3650
5030	-3930	-3665	-3940	-3660	-4090	-3650
5020	-3920	-3660	-3930	-3660	-4085	-3650
5010	-3915	-3660	-3925	-3655	-4075	-3645
5000	-3910	-3655	-3915	-3655	-3950	-3640
4990	-3905	-3655	-3910	-3650	-3945	-3645
4980	-3900	-3655	-3905	-3650	-3935	-3645
4970	-3895	-3650	-3900	-3645	-3925	-3640
4960	-3895	-3650	-3895	-3645	-3920	-3640
4950	-3890	-3650	-3895	-3645	-3910	-3640
4940	-3885	-3640	-3890	-3635	-3905	-3635
4930	-3885	-3640	-3880	-3635	-3900	-3635
4910	-3880	-3640	-3885	-3635	-3900	-3635
4900	-3875	-3555	-3880	-3545	-3900	-3630
4890	-3870	-3545	-3875	-3535	-3905	-3625
4880	-3870	-3395	-3875	-3395	-3890	-3620
4870	-3395	-3395	-3875	-3395	-3885	-3620
4860	-3395	-3395	-3870	-3395	-3880	-3620
4850	-3385	-3385	-3865	-3385	-3875	-3615
4840	-3385	-3385	-3860	-3380	-3875	-3615
4830	-3385	-3385	-3855	-3385	-3870	-3615
4820	-3380	-3380	-3850	-3375	-3870	-3615
4810	-3375	-3375	-3795	-3365	-3865	-3615
4800	-3375	-3375	-3790	-3360	-3860	-3615

RADIOCARBON AGE (BP) 5568 HALF-LIFE	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT			SIGMA=200 YRS.	SIGMA=300 YRS.
		SIGMA= 50 YRS.	SIGMA=100 YRS.	SIGMA=150 YRS.		
4790	-3780	-3370	-3370	-3855	-3180	-3150
4780	-3780	-3370	-3370	-3850	-3175	-3145
4770	-3775	-3370	-3365	-3795	-3170	-3075
4760	-3775	-3365	-3365	-3790	-3170	-3055
4750	-3770	-3365	-3360	-3785	-3165	-3045
4740	-3765	-3365	-3360	-3785	-3160	-3040
4730	-3765	-3360	-3360	-3780	-3155	-3035
4720	-3760	-3360	-3355	-3780	-3150	-3025
4710	-3760	-3355	-3355	-3775	-3150	-3020
4700	-3755	-3355	-3350	-3775	-3170	-3075
4690	-3655	-3355	-3355	-3770	-3165	-3055
4680	-3655	-3350	-3350	-3765	-3160	-3050
4670	-3650	-3190	-3185	-3765	-3160	-2975
4660	-3650	-3190	-3185	-3760	-3155	-2955
4650	-3645	-3175	-3175	-3760	-3150	-2945
4640	-3640	-3170	-3165	-3770	-3165	-2955
4630	-3635	-3165	-3160	-3765	-3160	-2950
4620	-3625	-3165	-3155	-3765	-3155	-2940
4610	-3550	-3160	-3155	-3760	-3150	-2935
4600	-3520	-3155	-3155	-3755	-3150	-2930
4590	-3515	-3150	-3145	-3755	-3145	-2925
4580	-3510	-3145	-3145	-3645	-3030	-2920
4570	-3505	-3140	-3140	-3645	-3025	-2915
4560	-3500	-3160	-3155	-3645	-3025	-2910
4550	-3495	-3050	-3050	-3540	-3035	-2905
4540	-3490	-3045	-3030	-3530	-2970	-2945
4530	-3490	-3035	-3025	-3525	-2950	-2905
4520	-3485	-3030	-3015	-3515	-2940	-2905
4510	-3480	-3020	-3005	-3510	-2930	-2895
4500	-3475	-3015	-2995	-3505	-2925	-2890
4490	-3470	-3005	-3000	-3500	-2970	-2910
4480	-3465	-2990	-3495	-3500	-2950	-2890
4470	-3370	-2970	-3495	-3500	-2905	-2850
4460	-3370	-2955	-3490	-3490	-2900	-2875
4450	-3370	-2940	-3485	-3485	-2900	-2870

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=300 YRS.
		SIGMA= 50 YRS.	SIGMA=100 YRS.	SIGMA=150 YRS.	
5568 HALF-LIFE					
4440	-3365	-2930	-3370	-2920	-3480
4430	-3365	-2925	-3370	-2915	-3475
4420	-3365	-2920	-3365	-2910	-3470
4410	-3360	-2915	-3365	-2905	-3465
4400	-3360	-2910	-3365	-2900	-3370
4390	-3360	-2905	-3360	-2895	-3370
4380	-3355	-2900	-3360	-2895	-3365
4370	-3355	-2895	-3355	-2890	-3365
4360	-3350	-2895	-3355	-2885	-3365
4350	-3350	-2890	-3355	-2880	-3360
4340	-3345	-2885	-3350	-2880	-3360
4330	-3345	-2880	-3350	-2875	-3360
4320	-3340	-2880	-3345	-2870	-3355
4310	-3335	-2875	-3345	-2865	-3355
4300	-3330	-2870	-3340	-2860	-3350
4290	-3155	-2865	-3335	-2800	-3350
4280	-3150	-2860	-3160	-2860	-3350
4270	-3145	-2855	-3155	-2870	-3345
4260	-3140	-2860	-3150	-2665	-3340
4250	-3130	-2670	-3145	-2660	-3340
4240	-3040	-2665	-3135	-2660	-3330
4230	-3030	-2665	-3045	-2655	-3350
4220	-3025	-2660	-3035	-2650	-3345
4210	-3015	-2655	-3030	-2650	-3350
4200	-3005	-2650	-3020	-2645	-3345
4190	-2970	-2650	-3015	-2640	-3135
4180	-2940	-2645	-3000	-2640	-3045
4170	-2930	-2640	-2965	-2635	-3035
4160	-2920	-2640	-2940	-2560	-3030
4150	-2915	-2635	-2925	-2550	-3020
4140	-2905	-2560	-2920	-2545	-3015
4130	-2900	-2550	-2910	-2435	-3000
4120	-2895	-2550	-2905	-2430	-2970
4110	-2895	-2435	-2900	-2425	-2945
4100	-2890	-2430	-2895	-2420	-2930

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=300 YRS.					
		SIGMA= 50 YRS.	SIGMA=100 YRS.	SIGMA=150 YRS.						
5568 HALF-LIFE										
4090	-2885 *	-2425	-2895	-2400	-3025	-2310	-3155	-2180	-3365	-1965
4080	-2880	-2425	-2890	-2415	-2915	-2400	-3020	-2310	-3150	-1955
4070	-2875	-2420	-2885	-2415	-2910	-2395	-3010	-2305	-3145	-1945
4060	-2870	-2415	-2880	-2410	-2905	-2335	-2990	-2210	-3140	-1925
4050	-2865	-2415	-2875	-2405	-2900	-2325	-2960	-2200	-3130	-1905
4040	-2860	-2410	-2870	-2405	-2895	-2320	-2940	-2195	-3040	-1895
4030	-2795	-2410	-2865	-2400	-2890	-2315	-2930	-2185	-3035	-1890
4020	-2785	-2405	-2860	-2395	-2885	-2310	-2920	-2180	-3025	-1885
4010	-2780	-2400	-2795	-2395	-2885	-2310	-2915	-2175	-3020	-1875
4000	-2775	-2400	-2785	-2330	-2880	-2305	-2910	-2170	-3010	-1870
3990	-2770	-2395	-2780	-2325	-2875	-2210	-2905	-2160	-2965	-1865
3980	-2765	-2330	-2775	-2320	-2870	-2200	-2900	-2155	-2960	-1860
3970	-2755	-2325	-2770	-2315	-2865	-2190	-2895	-2150	-2945	-1855
3960	-2650	-2320	-2765	-2315	-2855	-2185	-2890	-2140	-2930	-1780
3950	-2645	-2315	-2760	-2305	-2795	-2180	-2885	-2130	-2920	-1770
3940	-2640	-2310	-2650	-2215	-2785	-2175	-2880	-2120	-2915	-1765
3930	-2640	-2305	-2645	-2205	-2780	-2165	-2880	-2110	-2910	-1760
3920	-2635	-2215	-2640	-2195	-2775	-2160	-2875	-2095	-2905	-1750
3910	-2630	-2205	-2640	-2190	-2770	-2155	-2870	-2010	-2900	-1740
3900	-2625	-2195	-2635	-2185	-2765	-2145	-2865	-1995	-2895	-1730
3890	-2620	-2190	-2630	-2180	-2755	-2140	-2855	-1985	-2890	-1730
3880	-2545	-2185	-2625	-2170	-2650	-2130	-2795	-1975	-2890	-1730
3870	-2540	-2180	-2620	-2165	-2645	-2120	-2785	-1970	-2885	-1725
3860	-2540	-2175	-2545	-2160	-2640	-2110	-2780	-1960	-2880	-1720
3850	-2535	-2165	-2540	-2150	-2640	-2090	-2775	-1950	-2875	-1715
3840	-2530	-2160	-2540	-2145	-2635	-2005	-2770	-1940	-2870	-1705
3830	-2530	-2150	-2535	-2135	-2630	-1990	-2765	-1915	-2865	-1700
3820	-2525	-2145	-2530	-2125	-2625	-1980	-2755	-1900	-2860	-1695
3810	-2525	-2135	-2530	-2115	-2620	-1975	-2650	-1895	-2845	-1690
3800	-2520	-2125	-2525	-2105	-2545	-1965	-2645	-1885	-2790	-1685
3790	-2515	-2115	-2520	-2025	-2540	-1960	-2640	-1880	-2785	-1680
3780	-2515	-2105	-2520	-2000	-2535	-1950	-2640	-1875	-2780	-1675
3770	-2510	-2025	-2515	-1990	-2535	-1935	-2635	-1870	-2770	-1670
3760	-2395	-2000	-2515	-1980	-2530	-1910	-2630	-1865	-2765	-1670
3750	-2395	-1990	-2510	-1970	-2530	-1900	-2625	-1790	-2760	-1670

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE)			FOR MEASUREMENT UNCERTAINTIES OF SIGMA=150 YRS.	SIGMA=200 YRS.	SIGMA=300 YRS.					
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.								
3740	-2385	-1980	-2500	-1965	-2525	-1890	-2620	-1775	-2650	-1710	-2895	-1570
3730	-2325	-1970	-2395	-1955	-2525	-1885	-2545	-1770	-2645	-1700	-2890	-1560
3720	-2320	-1965	-2390	-1945	-2520	-1880	-2540	-1760	-2645	-1695	-2885	-1555
3710	-2315	-1955	-2325	-1925	-2515	-1875	-2535	-1755	-2640	-1690	-2880	-1550
3700	-2310	-1945	-2320	-1905	-2515	-1870	-2535	-1745	-2635	-1685	-2875	-1540
3690	-2310	-1925	-2315	-1895	-2510	-1865	-2530	-1735	-2630	-1680	-2875	-1535
3680	-2305	-1905	-2310	-1890	-2395	-1785	-2530	-1725	-2625	-1675	-2870	-1525
3670	-2300	-1895	-2310	-1885	-2390	-1775	-2525	-1710	-2620	-1670	-2860	-1440
3660	-2295	-1890	-2305	-1875	-2385	-1770	-2525	-1705	-2610	-1670	-2855	-1430
3650	-2285	-1885	-2300	-1870	-2325	-1760	-2520	-1695	-2540	-1665	-2790	-1425
3640	-2185	-1875	-2295	-1865	-2320	-1755	-2515	-1690	-2540	-1660	-2785	-1420
3630	-2180	-1870	-2285	-1860	-2315	-1745	-2515	-1685	-2535	-1655	-2780	-1415
3620	-2175	-1865	-2185	-1855	-2310	-1735	-2510	-1680	-2530	-1645	-2775	-1405
3610	-2170	-1860	-2180	-1850	-2305	-1720	-2395	-1680	-2530	-1635	-2770	-1400
3600	-2160	-1780	-2175	-1765	-2300	-1710	-2390	-1675	-2525	-1580	-2765	-1395
3590	-2155	-1770	-2170	-1755	-2295	-1705	-2385	-1670	-2525	-1570	-2755	-1390
3580	-2145	-1765	-2160	-1750	-2290	-1695	-2325	-1665	-2520	-1565	-2650	-1385
3570	-2140	-1755	-2155	-1740	-2285	-1690	-2320	-1660	-2520	-1555	-2645	-1375
3560	-2130	-1750	-2150	-1730	-2185	-1685	-2315	-1655	-2515	-1550	-2640	-1370
3550	-2120	-1740	-2140	-1715	-2180	-1680	-2310	-1650	-2510	-1545	-2635	-1360
3540	-2110	-1730	-2130	-1705	-2170	-1675	-2305	-1640	-2505	-1535	-2635	-1355
3530	-2090	-1715	-2120	-1700	-2165	-1675	-2300	-1585	-2395	-1530	-2630	-1345
3520	-2005	-1705	-2110	-1695	-2160	-1670	-2295	-1575	-2390	-1450	-2625	-1335
3510	-1995	-1700	-2095	-1690	-2150	-1665	-2290	-1565	-2325	-1435	-2615	-1325
3500	-1985	-1695	-2005	-1685	-2145	-1660	-2280	-1560	-2320	-1430	-2545	-1270
3490	-1975	-1690	-1995	-1680	-2135	-1655	-2185	-1555	-2315	-1420	-2540	-1255
3480	-1970	-1685	-1985	-1675	-2125	-1650	-2175	-1545	-2310	-1415	-2535	-1250
3470	-1960	-1680	-1975	-1670	-2115	-1640	-2170	-1540	-2305	-1405	-2535	-1140
3460	-1950	-1675	-1970	-1665	-2105	-1635	-2165	-1535	-2305	-1400	-2530	-1130
3450	-1940	-1670	-1960	-1665	-2080	-1575	-2160	-1525	-2300	-1400	-2530	-1125
3440	-1920	-1670	-1955	-1660	-2060	-1565	-2150	-1440	-2295	-1390	-2525	-1120
3430	-1905	-1665	-1940	-1655	-1990	-1560	-2145	-1430	-2285	-1385	-2525	-1115
3420	-1885	-1660	-1920	-1645	-1980	-1555	-2135	-1425	-2185	-1380	-2515	-1115
3410	-1880	-1655	-1905	-1635	-1975	-1545	-2125	-1420	-2180	-1375	-2510	-1110
3400	-1885	-1645	-1895	-1585	-1965	-1545	-2115	-1415	-2175	-1365	-2510	-1110

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA= 300 YRS.
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.	
5568 HALF-LIFE					
3390	-1875	-1635	-1890	-1575	-1535
3380	-1870	-1585	-1885	-1565	-1950
3370	-1865	-1575	-1875	-1560	-1935
3360	-1860	-1565	-1870	-1550	-1910
3350	-1855	-1560	-1865	-1545	-1900
3340	-1845	-1555	-1860	-1540	-1890
3330	-1765	-1545	-1855	-1530	-1885
3320	-1760	-1540	-1850	-1520	-1880
3310	-1750	-1530	-1855	-1440	-1875
3300	-1740	-1525	-1860	-1430	-1870
3290	-1730	-1445	-1750	-1425	-1865
3280	-1715	-1435	-1740	-1420	-1860
3270	-1705	-1425	-1730	-1415	-1850
3260	-1695	-1420	-1720	-1410	-1770
3250	-1690	-1415	-1725	-1400	-1760
3240	-1685	-1410	-1700	-1395	-1755
3230	-1680	-1405	-1690	-1390	-1745
3220	-1675	-1400	-1685	-1385	-1735
3210	-1670	-1395	-1680	-1380	-1720
3200	-1670	-1385	-1675	-1370	-1710
3190	-1665	-1380	-1675	-1365	-1700
3180	-1660	-1375	-1670	-1355	-1695
3170	-1650	-1365	-1665	-1350	-1690
3160	-1645	-1360	-1660	-1340	-1685
3150	-1585	-1350	-1655	-1330	-1680
3140	-1575	-1345	-1645	-1310	-1675
3130	-1565	-1335	-1625	-1260	-1670
3120	-1560	-1320	-1575	-1255	-1665
3110	-1555	-1265	-1570	-1140	-1660
3100	-1545	-1255	-1560	-1135	-1655
3090	-1540	-1145	-1555	-1130	-1650
3080	-1530	-1135	-1545	-1125	-1640
3070	-1520	-1130	-1540	-1120	-1585
3060	-1430	-1125	-1535	-1115	-1575
3050	-1425	-1120	-1525	-1115	-1565

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA= 200 YRS.	SIGMA= 300 YRS.
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.		
5568 HALF-LIFE						
3040	-1420	-1120	-1505	-1110	-1660	-825
3030	-1410	-1115	-1455	-1105	-1655	-885
3020	-1405	-1110	-1420	-1105	-1650	-880
3010	-1400	-1110	-1415	-1100	-1640	-870
3000	-1395	-1105	-1405	-935	-1585	-860
2990	-1385	-1100	-1400	-925	-1520	-800
2980	-1380	-1095	-1395	-920	-1435	-835
2970	-1370	-1080	-1390	-915	-1425	-890
2960	-1365	-925	-1380	-910	-1400	-885
2950	-1355	-915	-1375	-905	-1415	-880
2940	-1345	-910	-1365	-905	-1405	-870
2930	-1335	-910	-1360	-900	-1400	-860
2920	-1325	-905	-1350	-895	-1395	-845
2910	-1270	-900	-1340	-890	-1390	-835
2900	-1260	-895	-1330	-885	-1380	-830
2890	-1250	-890	-1310	-880	-1375	-825
2880	-1245	-885	-1260	-870	-1370	-820
2870	-1240	-880	-1255	-860	-1360	-815
2860	-1235	-875	-1250	-845	-1350	-810
2850	-1235	-860	-1245	-835	-1340	-805
2840	-1230	-850	-1240	-830	-1330	-805
2830	-1225	-835	-1235	-825	-1315	-800
2820	-1220	-830	-1230	-820	-1265	-795
2810	-1215	-825	-1225	-815	-1255	-795
2800	-1210	-820	-1225	-810	-1250	-790
2790	-1105	-815	-1220	-805	-1245	-790
2780	-1100	-810	-1210	-805	-1240	-785
2770	-1095	-805	-1105	-800	-1235	-785
2760	-1090	-805	-1105	-795	-1230	-780
2750	-1075	-800	-1100	-795	-1225	-775
2740	-1045	-800	-1095	-790	-1225	-645
2730	-1035	-795	-1085	-785	-1220	-640
2720	-1030	-790	-1045	-785	-1215	-630
2710	-1025	-790	-1040	-785	-1210	-625
2700	-1020	-785	-1035	-780	-1205	-620

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF				SIGMA=300 YRS.					
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.	SIGMA= 200 YRS.						
5568 HALF-LIFE											
2690	-1020	-785	-1030	-775	-615	-1230	-440	-1330	-410	-1520	-200
2680	-1015	-780	-1025	-650	-615	-1230	-435	-1275	-410	-1435	-195
2670	-1005	-775	-1020	-640	-610	-1225	-430	-1260	-405	-1425	-190
2660	-985	-655	-1015	-635	-605	-1220	-425	-1255	-405	-1420	-185
2650	-880	-640	-1010	-625	-595	-1215	-420	-1245	-400	-1415	-180
2640	-875	-635	-1000	-625	-590	-1205	-420	-1245	-400	-1410	-180
2630	-860	-630	-880	-620	-580	-1105	-415	-1240	-395	-1400	-175
2620	-845	-625	-875	-615	-585	-1100	-410	-1235	-395	-1395	-170
2610	-835	-620	-865	-610	-585	-1095	-410	-1230	-395	-1390	-165
2600	-830	-615	-850	-605	-585	-1085	-410	-1225	-390	-1385	-165
2590	-825	-610	-840	-595	-585	-1010	-425	-1045	-405	-1220	-390
2580	-820	-605	-830	-590	-580	-1000	-420	-1040	-405	-1215	-385
2570	-815	-600	-825	-585	-580	-980	-420	-1035	-400	-1210	-380
2560	-810	-595	-820	-440	-875	-975	-415	-1030	-400	-1105	-265
2550	-805	-585	-815	-435	-865	-965	-415	-1025	-395	-1100	-245
2540	-805	-445	-810	-430	-850	-950	-410	-1020	-395	-1095	-205
2530	-600	-435	-810	-425	-840	-940	-410	-1015	-390	-1090	-195
2520	-800	-430	-805	-420	-830	-930	-405	-1010	-390	-1050	-190
2510	-795	-425	-800	-420	-825	-925	-405	-1000	-385	-1040	-185
2500	-790	-425	-800	-415	-820	-920	-400	-980	-385	-1035	-185
2490	-790	-420	-795	-415	-815	-905	-400	-875	-270	-1030	-180
2480	-785	-415	-790	-410	-810	-895	-395	-865	-260	-1025	-175
2470	-785	-415	-790	-410	-810	-895	-395	-850	-250	-1020	-170
2460	-780	-410	-785	-405	-805	-890	-390	-840	-200	-1015	-170
2450	-775	-410	-785	-405	-800	-890	-390	-830	-195	-1010	-165
2440	-775	-405	-780	-400	-800	-885	-385	-825	-190	-1005	-160
2430	-770	-405	-780	-400	-795	-885	-385	-820	-185	-885	-160
2420	-770	-400	-775	-395	-790	-875	-375	-815	-180	-880	-155
2410	-765	-400	-770	-395	-790	-860	-360	-810	-180	-870	-150
2400	-760	-400	-770	-395	-785	-855	-355	-815	-175	-855	-145
2390	-755	-395	-765	-390	-785	-850	-350	-805	-170	-845	-135
2380	-755	-395	-760	-390	-780	-845	-345	-800	-165	-835	-125
2370	-750	-390	-760	-385	-780	-840	-340	-795	-160	-830	-120
2360	-740	-390	-755	-275	-775	-835	-335	-775	-155	-820	-115
2350	-590	-385	-750	-265	-770	-820	-330	-770	-150	-820	-110

RADIOCARBON 5568 HALF-LIFE AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=200 YRS.	SIGMA=300 YRS.
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.		
2340	-580	-740	-245	-770	-180	-150
2330	-575	-755	-205	-765	-175	-45
2320	-560	-760	-195	-760	-170	-35
2310	-550	-755	-205	-760	-170	-30
2300	-535	-720	-565	-755	-165	-25
2290	-420	-190	-555	-750	-160	-775
2280	-420	-190	-540	-740	-155	-770
2270	-415	-185	-525	-745	-155	-770
2260	-415	-180	-420	-740	-170	-765
2250	-410	-175	-415	-740	-170	-760
2240	-410	-170	-415	-750	-165	-765
2230	-405	-170	-410	-755	-160	-755
2220	-405	-165	-410	-755	-155	-750
2210	-400	-160	-405	-755	-155	-745
2200	-400	-160	-405	-755	-155	-745
2190	-395	-155	-400	-765	-165	-760
2180	-395	-145	-400	-765	-160	-755
2170	-390	-135	-400	-765	-155	-750
2160	-390	-130	-395	-765	-150	-745
2150	-385	-125	-390	-765	-150	-740
2140	-385	-20	-390	-765	-15	-740
2130	-380	-15	-390	-765	-10	-740
2120	-380	-10	-385	-765	-10	-740
2110	-375	-11	-385	-765	-10	-740
2100	-375	-1/1	-380	-765	-15	-740
2090	-370	10	-375	-765	25	-740
2080	-365	15	-375	-765	30	-740
2070	-355	20	-370	-765	35	-740
2060	-385	25	-365	-765	40	-740
2050	-180	35	-360	-765	45	-740
2040	-175	40	-185	-780	210	-400
2030	-170	40	-180	-780	215	-395
2020	-170	45	-175	-780	200	-375
2010	-165	190	-175	-780	200	-370
2000	-160	195	-170	-780	205	-365

RADIOCARBON AGE (BP)	SIGMA = 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=300 YRS.							
		SIGMA = 50 YRS.	SIGMA = 100 YRS.	SIGMA = 150 YRS.								
5568 HALF-LIFE												
1990	-160	200	-165	210	-360	225	-385	240	-405	390	-750	575
1980	-155	205	-165	210	-185	225	-385	245	-400	405	-740	580
1970	-150	205	-160	215	-180	230	-380	250	-400	405	-595	585
1960	-145	210	-155	215	-175	235	-375	330	-395	420	-590	585
1950	-145	215	-155	220	-175	235	-375	330	-395	425	-580	590
1940	-135	215	-150	220	-170	235	-370	345	-390	430	-570	595
1930	-125	220	-145	225	-165	240	-370	360	-390	435	-560	595
1920	-120	220	-140	225	-165	240	-365	380	-390	440	-555	600
1910	-115	225	-130	230	-160	245	-365	400	-385	450	-530	605
1900	-110	225	-120	230	-155	250	-180	410	-385	455	-420	605
1890	-1/1	225	-15	235	-155	255	-180	415	-380	545	-420	610
1880	-1/1	230	-10	235	-150	330	-175	420	-375	555	-415	610
1870	10	230	-5	240	-145	340	-170	425	-375	560	-410	615
1860	15	235	-1/1	240	-140	355	-165	435	-370	565	-410	620
1850	25	240	5	245	-130	380	-165	440	-365	570	-405	620
1840	30	240	15	250	-20	395	-160	445	-360	575	-405	625
1830	35	245	20	255	-15	405	-155	450	-385	580	-405	630
1820	40	250	25	330	-10	415	-155	460	-400	580	-400	635
1810	45	255	35	340	-1/1	420	-150	550	-175	585	-400	640
1800	45	255	40	355	-1/1	425	-145	560	-170	590	-395	640
1790	50	335	40	380	10	430	-140	565	-170	590	-395	645
1780	55	350	45	395	15	440	-130	570	-165	595	-390	650
1770	60	370	50	405	20	445	-20	575	-160	600	-390	750
1760	65	395	55	415	25	450	-15	575	-160	600	-385	760
1750	70	405	60	420	35	455	-10	580	-155	605	-385	765
1740	70	410	60	425	40	550	-5	585	-150	605	-380	770
1730	75	420	65	430	40	560	-1/1	585	-145	610	-380	775
1720	80	425	70	440	45	565	5	590	-140	615	-375	785
1710	90	430	75	445	50	570	15	595	-135	615	-370	790
1700	220	435	80	450	55	570	20	595	-125	620	-370	850
1690	225	440	85	455	60	575	25	600	-20	625	-365	860
1680	225	450	100	550	65	580	30	600	-10	630	-185	865
1670	230	455	225	555	65	585	35	605	-5	630	-180	870
1660	230	545	225	565	70	585	40	610	-1/1	635	-180	875
1650	235	555	230	570	75	590	45	610	5	640	-175	880

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE)			FOR MEASUREMENT UNCERTAINTIES OF					
		SIGMA= 50 YRS.	SIGMA=100 YRS.	SIGMA=150 YRS.	SIGMA=200 YRS.	SIGMA=300 YRS.				
5568 HALF-LIFE										
1640	240	560	230	570	50	615	10	645	-170	885
1630	240	565	235	575	85	595	55	620	20	650
1620	245	570	235	580	100	600	60	620	25	670
1610	245	575	240	585	225	600	60	625	30	755
1600	250	580	245	585	225	605	65	630	35	760
1590	260	580	245	590	230	610	70	635	40	770
1580	335	585	250	590	230	610	75	635	45	775
1570	345	590	255	595	235	615	80	640	50	780
1560	365	590	270	600	235	620	85	645	50	785
1550	390	595	345	600	240	620	95	650	55	795
1540	400	595	360	605	245	625	225	750	60	855
1530	410	600	385	610	245	630	225	755	65	860
1520	415	605	400	610	250	630	230	765	70	865
1510	425	605	410	615	255	635	230	770	75	870
1500	430	610	415	615	265	640	235	775	80	875
1490	435	615	420	620	345	645	235	780	85	880
1480	440	615	425	625	360	650	240	790	90	885
1470	445	620	435	630	385	745	240	845	220	885
1460	455	625	440	630	400	755	245	855	225	890
1450	460	625	445	635	410	765	250	865	225	895
1440	470	630	450	640	415	770	255	870	230	900
1430	565	635	460	645	420	775	265	875	230	900
1420	570	640	470	650	425	780	340	880	235	905
1410	570	640	470	650	435	785	355	880	240	910
1400	575	645	475	655	440	795	380	885	240	915
1390	580	650	570	760	445	855	395	890	245	915
1380	585	750	575	765	450	860	405	890	245	915
1370	585	760	580	775	460	865	415	895	250	1015
1360	590	765	580	780	465	870	420	900	260	1020
1350	595	770	585	785	560	875	425	900	335	1025
1340	595	775	590	795	565	880	430	905	350	1030
1330	600	780	590	850	570	885	440	910	365	1030
1320	605	790	595	860	575	885	445	915	390	1035
1310	605	800	600	865	580	890	450	920	400	1040
1300	610	855	600	870	580	895	455	1005	410	1045

RADIOCARBON AGE (BP)	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE)			FOR MEASUREMENT UNCERTAINTIES OF		
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.	SIGMA= 200 YRS.	SIGMA= 300 YRS.	
5568 HALF-LIFE							
1290	610	860	605	875	585	900	465
1280	615	865	610	880	590	900	485
1270	620	870	615	885	590	905	565
1260	625	875	620	890	600	910	570
1250	630	880	625	895	605	1005	575
1240	635	885	630	900	605	1010	580
1230	640	890	635	905	610	1020	590
1220	645	895	640	905	615	1025	590
1210	650	900	645	905	620	1030	595
1200	655	905	650	910	625	1035	600
1190	660	905	655	915	630	1040	605
1180	665	910	660	920	635	1045	610
1170	670	915	665	925	640	1050	615
1160	675	915	675	930	645	1055	620
1150	680	915	685	935	650	1060	625
1140	685	900	670	905	655	1045	605
1130	685	900	670	905	640	1050	610
1120	690	900	675	905	645	1055	615
1110	695	900	680	905	650	1060	620
1100	695	900	685	905	655	1065	625
1090	700	900	690	905	660	1070	630
1080	705	905	695	910	670	1075	635
1070	710	905	700	915	675	1080	640
1060	715	905	705	915	680	1085	645
1050	720	905	710	915	685	1090	650
1040	725	905	715	915	690	1095	655
1030	730	905	720	915	695	1100	660
1020	735	905	725	915	700	1105	665
1010	740	905	730	915	705	1110	670
1000	745	905	735	915	710	1115	675
990	750	905	740	915	715	1120	680
980	755	905	745	915	720	1125	685
970	760	905	750	915	725	1130	690
960	765	905	755	915	730	1135*	695
950	770	905	760	915	735	1135*	700

RADIOCARBON AGE (BP)	SIGMA = 5568 HALF-LIFE	CALIBRATED RANGES (95% CONFIDENCE)			FOR MEASUREMENT	UNCERTAINTIES OF
		SIGMA = 20 YRS.	SIGMA = 50 YRS.	SIGMA = 100 YRS.		
940	1020	1215	920	895	1255	660
930	1025	1220	925	895	1255	660
920	1030	1225	930	900	1260	670
910	1030	1230	1020	1240	905	770
900	1035	1235	1025	1245	910	775
890	1040	1240	1035	1250	915	780
880	1045	1240	1035	1250	915	785
870	1045	1245	1035	1255	920	790
860	1050	1250	1040	1260	925	860
850	1055	1255	1045	1260	935	895
840	1060	1255*	1055	1265	1025	905
830	1115	1260	1055	1270	1025	905
820	1170	1265	1060	1275*	1030	925
810	1185	1265	1065	1285*	1035	930
800	1195	1270	1160	1295	1040	1335
790	1205	1275	1175	1310	1040	1335
780	1210	1285	1190	1315	1045	1340
770	1215	1300	1200	1320	1050	1345
760	1220	1300	1205	1325	1055	1350
750	1225	1315	1210	1330	1060	1360*
740	1230	1320	1215	1330	1070	1390*
730	1235	1325	1220	1335	1165	1395
720	1235	1330	1225	1340	1180	1400
710	1240	1335	1230	1340	1195	1400
700	1245	1335	1235	1345	1200	1405
690	1250	1340	1240	1350	1210	1405
680	1250	1345	1240	1385	1215	1410
670	1255	1350	1245	1395	1220	1410
660	1260	1350	1250	1395	1225	1415
650	1265	1390	1255	1400	1230	1415
640	1265	1395	1255	1405	1235	1415
630	1270	1400	1260	1405	1235	1420
620	1275	1400	1265	1405	1240	1420
610	1285	1405	1270	1410	1245	1425
600	1300	1405	1270	1410	1245	1425

SIGMA=300 YRS.
 SIGMA=200 YRS.
 SIGMA=150 YRS.
 SIGMA=100 YRS.
 SIGMA=50 YRS.

RADIOCARBON AGE (BP)	5568 HALF-LIFE	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF				SIGMA=200 YRS.								
		SIGMA= 20 YRS.	SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.									
590	1310	1410	1280	1415	1250	1430	1220	1505	1070	1645*	1645	1645*	910	1950*
580	1315	1410	1285	1415	1255	1435	1225	1510	1165	1645	1645	1645	915	1950*
570	1320	1410	1305	1420	1260	1475	1230	1515	1180	1650	1650	1650	920	1950*
560	1325	1415	1315	1425	1260	1485	1235	1595*	1195	1650	1650	1650	925	1950*
550	1330	1415	1320	1420	1265	1490	1240	1615	1200	1655	1655	1655	930	1950*
540	1335	1420	1325	1425	1270	1495	1240	1625	1210	1655	1655	1655	1020	1950
530	1335	1420	1325	1430	1275	1500	1245	1635	1215	1660	1660	1660	1025	1950
510	1340	1425	1330	1430	1280	1500	1250	1640	1220	1660	1660	1660	1030	1950
500	1350	1430	1335	1435	1295	1505	1255	1645	1225	1665	1665	1665	1030	1950
490	1355	1430	1340	1485	1310	1515	1270	1650	1230	1790*	1790*	1790*	1035	1950
480	1390	1435	1345	1490	1320	1605*	1265	1655	1235	1800*	1800*	1800*	1040	1950
470	1395	1480	1350	1495	1325	1620	1270	1655	1240	1950*	1950*	1950*	1045	1950
460	1400	1485	1385	1500	1330	1630	1270	1655	1245	1950*	1950*	1950*	1050	1950
450	1400	1490	1390	1505	1330	1635	1280	1660	1245	1950*	1950*	1950*	1055	1950
440	1405	1495	1395	1510	1335	1640	1285	1660	1250	1950*	1950*	1950*	1060	1950
430	1405	1500	1400	1515	1340	1645	1305	1665	1255	1950*	1950*	1950*	1155	1950
420	1410	1505	1405	1525	1340	1650	1315	1790*	1255	1950*	1950*	1950*	1170	1950
410	1410	1510	1405	1605*	1345	1650	1320	1795*	1260	1950*	1950*	1950*	1185	1950
400	1415	1515	1405	1620	1350	1655	1325	1800*	1265	1950*	1950*	1950*	1195	1950
390	1415	1525	1410	1630	1385	1655	1325	1950*	1270	1950	1950	1950	1205	1950
380	1415	1605	1410	1635	1395	1660	1330	1950*	1275	1950	1950	1950	1210	1950
370	1420	1620	1415	1640	1395	1660	1335	1950*	1280	1950	1950	1950	1215	1950
360	1420	1630	1415	1645	1400	1660	1335	1950*	1290	1950	1950	1950	1220	1950
350	1425	1635	1420	1650	1405	1665	1340	1950*	1305	1950	1950	1950	1225	1950
340	1425	1640	1420	1650	1405	1795*	1345	1950*	1315	1950	1950	1950	1230	1950
330	1430	1645	1420	1655	1410	1795*	1350	1950	1320	1950	1950	1950	1235	1950
320	1435	1650	1425	1655	1410	1800*	1355	1950	1325	1950	1950	1950	1240	1950
310	1440	1650	1425	1660	1410	1950*	1395	1950	1330	1950	1950	1950	1245	1950
300	1485	1655	1430	1660	1415	1950*	1395	1950	1330	1950	1950	1950	1250	1950
290	1490	1655	1435	1665	1415	1950*	1400	1950	1335	1950	1950	1950	1250	1950
280	1495	1660	1480	1665	1420	1950*	1405	1950	1340	1950	1950	1950	1255	1950
270	1500	1660	1485	1795*	1420	1950*	1405	1950	1345	1950	1950	1950	1260	1950
260	1505	1660	1490	1795*	1425	1950*	1405	1950	1350	1950	1950	1950	1260	1950
250	1510	1665	1495	1800*	1425	1950*	1410	1950	1350	1950	1950	1950	1260	1950

RADIOCARBON AGE (BP) 5568 HALF-LIFE	SIGMA= 20 YRS.	CALIBRATED RANGES (95% CONFIDENCE) FOR MEASUREMENT UNCERTAINTIES OF			SIGMA=300 YRS.
		SIGMA= 50 YRS.	SIGMA= 100 YRS.	SIGMA= 150 YRS.	
240	1515	1790*	1500	1430	1950
230	1525	1795*	1505	1430	1950
220	1620	1800*	1510	1435	1950
210	1630	1950*	1515	1435	1950
200	1640	1950*	1525	1485	1950
190	1645	1950*	1620	1490	1950
180	1645	1950*	1630	1495	1950
170	1650	1950*	1640	1500	1950
160	1650	1950*	1645	1505	1950
150	1655	1950	1645	1510	1950
140	1655	1950	1650	1520	1950
130	1660	1945	1650	1530	1950*
120	1660	1945	1655	1530	1950
110	1665	1940	1655	1535	1950
100	1665	1940*	1660	1640	1950
90	1670	1935*	1660	1945	1950
80	1670	1935*	1665	1940	1950
70	1675	1930*	1665	1940*	1950
60	1675	1930*	1670	1935*	1950
50	1680	1925*	1670	1935*	1950
40	1680	1920*	1670	1930*	1950
30	1815	1920*	1675	1930*	1945
20	1820	1915*	1680	1925*	1945
10	1680	1920*	1680	1920*	1940

TABLE 3
SUPPLEMENTARY CALIBRATION TABLES FOR THE MOST RECENT 1000 YEARS
This table lists calibration intervals only for the starred values in the main table, *i.e.*, only for ages consistent with
more than one calibration interval
Spaces between rows indicate steps of more than 10 years between tabulated radiocarbon ages.

SUPPLEMENTARY TABLES FOR SIGMA = 2.0		CALIBRATED RANGES (95% CONFIDENCE)		
RADIOCARBON AGE (BP)				
960	920	980	1005	1205
950	930	970	1015	1210
840	1060	1100	1140	1255
240	1515	1665	1760	1790
230	1525	1570	1605	1670
220	1620	1670	1720	1725
210	1630	1675	1715	1805
200	1640	1675	1710	1805
190	1645	1680	1705	1810
180	1645	1810	1845	1880
170	1650	1815	1840	1885
160	1650	1890	1915	1950
100	1665	1765	1790	1940
90	1670	1730	1795	1935
80	1670	1720	1800	1935
70	1675	1715	1800	1930
60	1675	1710	1805	1930
50	1680	1705	1810	1855
40	1680	1700	1810	1850
30	1815	1845	1885	1920
20	1820	1840	1885	1915

SUPPLEMENTARY TABLES FOR SIGMA = 50

RADIOCARBON AGE (BP)		CALIBRATED RANGES (95% CONFIDENCE)			
930	925	975	1010	1235	
920	930	965	1015	1235	
820	1060	1110	1130	1275	
810	1065	1095	1145	1285	
410	1405	1540	1560	1605	
270	1485	1665	1760	1795	
260	1490	1670	1725	1795	
250	1495	1670	1720	1800	
240	1500	1675	1715	1805	
230	1505	1675	1710	1805	
220	1510	1680	1705	1810	
210	1515	1810	1845	1880	
200	1525	1570	1605	1815	
190	1620	1890	1910	1950	
70	1665	1765	1790	1940	
60	1670	1730	1795	1935	
50	1670	1720	1800	1935	
40	1670	1715	1800	1930	
30	1675	1710	1805	1930	
20	1680	1705	1810	1855	
10	1680	1705	1810	1850	

SUPPLEMENTARY TABLES FOR SIGMA = 100

RADIOCARBON AGE (BP)	CALIBRATED RANGES (95% CONFIDENCE)				
870	920	980	1005	1290	
860	925	970	1010	1305	
850	935	955	1020	1315	
750	1060	1105	1135	1360	
740	1070	1090	1150	1390	
480	1320	1535	1565	1605	
340	1405	1665	1760	1795	
330	1410	1670	1725	1795	
320	1410	1670	1720	1800	
310	1410	1675	1715	1805	
300	1415	1675	1710	1805	
290	1415	1680	1705	1810	
280	1420	1810	1845	1880	
270	1420	1815	1840	1885	
260	1425	1830	1910	1950	
130	1530	1565	1610	1950	

SUPPLEMENTARY TABLES FOR SIGMA = 150

RADIOCARBON AGE (BP)	CALIBRATED RANGES (95% CONFIDENCE)			
790	925	975	1010	1400
780	935	960	1015	1405
680	1060	1110	1130	1425
670	1065	1095	1145	1430
560	1235	1525	1575	1595
420	1315	1665	1765	1790
410	1320	1670	1730	1795
400	1325	1670	1720	1800
390	1325	1675	1715	1800
380	1330	1675	1710	1805
370	1335	1680	1705	1810
360	1335	1680	1705	1810
350	1340	1815	1845	1855
340	1345	1820	1840	1885
60	1525	1575	1600	1950

SUPPLEMENTARY TABLES FOR SIGMA = 200

RADIOCARBON AGE (BP)	CALIBRATED RANGES (95% CONFIDENCE)			
950	655	725	750	1335
940	660	715	755	1340
930	670	705	765	1340
720	920	985	1005	1480
710	925	970	1010	1485
700	935	960	1020	1490
630	1045	1540	1560	1610
600	1060	1105	1135	1640
590	1070	1090	1150	1645
,				
500	1230	1665	1765	1790
490	1230	1670	1755	1795
480	1235	1670	1720	1800
470	1240	1670	1715	1800
460	1245	1675	1710	1805
450	1245	1675	1710	1805
440	1250	1680	1705	1810
430	1255	1815	1845	1880
420	1255	1820	1840	1885
410	1260	1890	1910	1950

SUPPLEMENTARY TABLES FOR SIGMA = 300

RADIOCARBON AGE (BP)	CALIBRATED RANGES (95% CONFIDENCE)				
790	660	720	750	1535	1605
780	665	710	760	1620	
770	675	695	765	1630	
660	885	1665	1770	1790	
650	890	1665	1760	1795	
640	890	1670	1725	1800	
630	895	1670	1720	1800	
620	900	1675	1715	1805	
610	900	1675	1710	1805	
600	905	1680	1705	1810	
590	910	1815	1845	1880	
580	915	1815	1840	1885	
570	920	1890	1910	1950	
560	925	980	1005	1950	
550	930	965	1015	1950	
440	1060	1100	1140	1950	