

Review

Calpain chronicle—an enzyme family under multidisciplinary characterization

By Hiroyuki SORIMACHI,^{*1,†} Shoji HATA^{*1} and Yasuko ONO^{*1}

(Communicated by Masanori OTSUKA, M.J.A.)

Abstract: Calpain is an intracellular Ca^{2+} -dependent cysteine protease (EC 3.4.22.17; Clan CA, family C02) discovered in 1964. It was also called CANP (Ca^{2+} -activated neutral protease) as well as CASF, CDP, KAF, *etc.* until 1990. Calpains are found in almost all eukaryotes and a few bacteria, but not in archaeobacteria. Calpains have a limited proteolytic activity, and function to transform or modulate their substrates' structures and activities; they are therefore called, "modulator proteases." In the human genome, 15 genes—*CAPN1*, *CAPN2*, *etc.*—encode a calpain-like protease domain. Their products are calpain homologs with divergent structures and various combinations of functional domains, including Ca^{2+} -binding and microtubule-interaction domains. Genetic studies have linked calpain deficiencies to a variety of defects in many different organisms, including lethality, muscular dystrophies, gastropathy, and diabetes. This review of the study of calpains focuses especially on recent findings about their structure–function relationships. These discoveries have been greatly aided by the development of 3D structural studies and genetic models.

Keywords: calpain, muscular dystrophy, calpainopathy, gastropathy, intracellular proteolysis, calcium ion

Introduction

An enzyme corresponding to calpain was first described in 1964 by Gordon Guroff (1933–1999).^{1),2)}

^{*1} Calpain Project, Department of Advanced Science for Biomolecules, Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan.

[†] Correspondence should be addressed: H. Sorimachi, Calpain Project, Department of Advanced Science for Biomolecules, Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa, Setagaya-ku, Tokyo 156-8506, Japan (e-mail: sorimachi-hr@igakuken.or.jp).

Abbreviations: aa: amino acid(s); aar: amino acid residue(s); C2 domain: PKC conserved domain 2; C2L domain: C2 domain-like domain; CANP: calcium-activated neutral protease; *CAPN*: calpain gene; *CAPNS*: calpain regulatory small subunit gene; CASF: Ca^{2+} -activated sarcoplasmic factor; CBS-1 and -2: Ca^{2+} -binding site in PC1 and PC2, respectively; CL: calpain catalytic large subunit; CysPc: calpain-like cysteine protease sequence motif defined in the conserved domain database of the National Center for Biotechnology Information (cd00044); GR domain: Gly-rich domain; KAF: kinase activating factor; MIT motif: microtubule interacting and transport motif; NIDDM: noninsulin-dependent diabetes mellitus (type 2 diabetes); PC1 and PC2 domains: calpain protease core domains 1 and 2, respectively; PEF: penta-EF-hand; PEF(L) and PEF(S) domains: PEF domains in calpain catalytic large subunit and regulatory small subunit, respectively; PKC: protein kinase C; QTL: quantitative trait locus.

Over the next several years, calpain was reported with various names and in various contexts by other researchers, including Edmond H. Fischer (1920–), Edwin G. Krebs (1918–2009),^{3),4)} Darrel E. Goll (1936–2008),⁵⁾ and Yasutomi Nishizuka (1932–2004).⁶⁾ In 1978, Kazutomo Imahori (1920–) and his colleagues purified chicken calpain to homogeneity and named it CANP (calcium-activated neutral protease).⁷⁾ This marked the beginning of serious, focused studies of calpain, and this protease was soon established as one of the most important and intriguing enzymes in the cell.^{8)–10)} That same year, an endogenous calpain-specific inhibitor protein was reported by Takashi Murachi (1926–1990) and his colleagues.¹¹⁾ They proposed the names calpain in place of CANP and calpastatin for its inhibitor protein in 1981.¹²⁾ In 1984, Koichi Suzuki (1939–2010) and his colleagues revealed the complete primary structure of the calpain catalytic subunit,¹³⁾ and calpain studies turned toward structure–function analyses. For more than ten years, both CANP and calpain were used in the literature. This complication was resolved by Suzuki in accordance with Imahori in 1990. He proposed the usage of calpain as an authorized name at the 8th International Conference




Fig. 1. Schematic structures of calpain superfamily members.

Calpain homologs have been identified in almost all eukaryotes and in some bacteria. Historically, there are several nomenclatures for domains. To avoid confusion in this review, the domains are called by the abbreviation corresponding to the structure, *i.e.*, PC1 (protease core domain 1), PC2 (protease core domain 2), C2L (C2 domain-like), PEF (penta-EF-hand) *etc.* Symbols: N, N-terminal region; PEF(L) and PEF(S), PEF domains of large catalytic and small regulatory subunits, respectively; GR, glycine-rich hydrophobic domain; AS, alternative splicing products; NS/IS1/IS2, CAPN3-characteristic sequences; C2, C2 domain; MIT, microtubule-interacting-and-transport motif; Zn, Zn-finger-containing domain; SOH, SOL-homology domain; DIS, CALPA-specific insertion sequence; TM, transmembrane domain; CSTN, calpastatin-like domain; IQ, a motif interactive with calmodulin.

on Proteolysis and Protein Turnover.¹⁴⁾ Since Suzuki's 1984 paper,¹³⁾ cDNA and genomic cloning studies have exploded, and thousands of calpains and related molecules were identified—among them 15 human calpain genes, which are now called *CAPNn* ($n = 1, 2, 3$, and $5-16$), which is short for *calpain*.¹⁵⁾ There are two genes for small regulatory subunits, *CAPNS1* and *CAPNS2*, short for *calpain* *small* subunit, and one for calpastatin, *CAST*.

Calpains (EC 3.4.22.17; Clan CA, family C02) constitute a distinct group of intracellular cysteine proteases found in almost all eukaryotes and a few bacteria (Fig. 1; for other reviews, see the two landmark reviews in 1991¹⁶⁾ and 2003¹⁷⁾ and later reviews¹⁸⁾⁻⁴⁹⁾). Calpains are first defined as cytosolic

proteases exhibiting Ca^{2+} -dependent proteolytic activity at a neutral pH. Most calpains are intracellular, and their activity is strictly regulated, as is the case with other intracellular proteolytic systems, such as the ubiquitin-proteasome system,⁵⁰⁾ the autophagy-lysosome system,⁵¹⁾ and caspases⁵²⁾ (Fig. 2). Caspases proteolyze substrates sequence-specifically, mostly for apoptosis, and they are activated by signal-dependent oligomerization. The former two systems act as "erasers" that degrade and eliminate substrate proteins. In contrast, calpains act in proteolytic processing, rather than degradation, *i.e.*, calpains proteolyze their substrates at a limited number of sites to transform or modulate their substrates' structures and activities. Therefore,




Fig. 2. Major intracellular proteolytic systems.

The ubiquitin–proteasome system degrades and eliminates specific substrate proteins in an ubiquitin-tagging system consisting of more than 1,000 ubiquitin ligases. The autophagy–lysosome system primarily degrades non-specific cell components, including proteins and micro-organisms, by compartmentalization by isolation membranes. The caspase system functions mainly for apoptosis. In contrast, calpains primarily use proteolytic processing, rather than degradation, to modulate or modify their substrates' activity, specificity, longevity, localization, and structure.

calpain is referred to as an intracellular “modulator” protease.^{53,54} Ubiquitin and autophagy systems are regulated in various steps mediated by many proteins, but calpain is distinctive in that a functional entity rests on a single molecule or two.

The importance of the calpains' physiological roles is reflected in the variety of defects caused by compromised calpain function in different tissues and organisms, including embryonic lethality (disruption of mouse *Capn2*),⁵⁵ muscular dystrophies (mutations in human *CAPN3*),⁵⁶ gastropathy (mutated *Capn8*),⁵⁷ lissencephaly⁵⁸ and tumorigenesis⁵⁹ (indirect defects in calpain functions), impaired neurogenesis in flies,⁶⁰ deficient sex determination in nematodes,⁶¹ defects in aleurone cell development in maize,⁶² and alkaline stress susceptibility in

fungi⁶³ and yeasts.^{64,65} These effects will be described in more detail in the following sections.

Mammalian gene products can be written as non-italic upper-case letters of their gene name, according to the international nomenclature system for a gene product; for example, mouse *Capn1* and *Capn2* produce CAPN1 and CAPN2, respectively. Although recently mammalian calpain gene products are often written as calpain-1, calpain-2, etc., this nomenclature is misleading as described below (“calpain” is also used for enzyme names), so we will use the above formal nomenclature (CAPN1, CAPN2, etc.) throughout this review. Some calpains were previously called by independent names, such as μ CL and p94. In this review, to clarify old and new mammalian calpain nomenclatures, we will note the

old after the current names: *e.g.*, CAPN1/ μ CL and CAPN3/p94.

1. A research history of calpain and its definition

Because of the scope of this review, it is important to define calpain, and the complexities of defining calpain are best explained in the context of its research history.^{(66)–(68)}

In 1964, Guroff partially purified a Ca^{2+} -dependent neutral protease from rat brain, which afterwards turned out to be calpain.¹⁾ In the same year, Meyer, Fischer and Krebs described the activation of rabbit skeletal muscle phosphorylase b kinase by Ca^{2+} and partially purified the responsible protein, which they called KAF (kinase activating factor).⁴⁾ Four years later, Huston and Krebs,³⁾ and Drummond and Duncan⁶⁹⁾ independently reported that this protein was a Ca^{2+} -dependent protease, although they concluded that the proteolytic activation of phosphorylase b kinase was unlikely to be physiological. Krebs and Fishers' main interest then moved to phosphorylation and its cascade, for which they were awarded a Nobel Prize in 1992.^{70),71)} Since then, Krebs has changed his mind about the physiological importance of proteolytic activation. Indeed, one of calpain's most important physiological functions is to modulate substrate protein functions by limited proteolysis, and this property must be considered in the definition—or at least, the classification—of calpain.

In 1972, Darrel E. Goll and his colleagues identified a factor that Ca^{2+} -dependently removes the Z-lines of skeletal muscle, and named it CASF (Ca²⁺-activated sarcoplasmic factor), which also turned out to be calpain.⁵⁾ This study is the beginning of research on calpain involved in post-mortem tenderization of meat.⁷²⁾ In 1977, Yasutomi Nishizuka and his colleagues identified calpain as a protein kinase C (PKC) activating factor,⁶⁾ and several calpain studies were done in line with PKC.^{73)–76)} In 1978, Imahori and his colleagues for the first time in the world purified chicken calpain to homogeneity and called it CANP. Following this remarkable study, Imahori's group established calpain research field by publishing important studies on enzymology of calpain such as substrate specificity,⁷⁷⁾ purification from human muscle,⁷⁸⁾ inhibitors,^{79),80)} autolysis,^{8),81)} and activation.^{82),83)}

Calpain purification studies (*ca.* 90% purity in 1976,⁸⁴⁾ and to homogeneity in 1978⁷⁾) revealed that functional native calpain is composed of large (*ca.* 80 kDa) and small (*ca.* 30 kDa) subunits, which is

another important property of calpain. The chicken μ /m-calpain catalytic subunit cDNA was cloned in 1984,¹³⁾ and since then many cDNAs and genomic DNAs corresponding to various calpain species have been cloned and sequenced. Little is known about the nature of calpain proteins as enzymes, however, and most calpains are known only by their sequence.

The most extensively studied calpains are the major ubiquitous mammalian μ - and m-calpains, and the major ubiquitous calpain in chicken, μ /m-calpain.¹⁴⁾ These are called “conventional” calpains in the wake of PKC;⁸⁵⁾ all others are termed “unconventional” calpains. The chicken μ /m-calpain has properties placing itself as an intermediate of μ - and m-calpains.^{7),77),86)} Its catalytic subunit was shown to be an ortholog of mammalian CAPN11 (see also 6.3.3).⁸⁷⁾

Among the amino acid (aa) sequences available, there is a group of peptidases that possess a protease domain that is significantly similar to one another but distinct from that of other peptidases.⁸⁸⁾ A search for “CysPC” in the conserved domain database of NCBI⁸⁹⁾ extracts the sequences of almost all the calpain homologs from various living organisms, including plants, fungi, yeast, and even bacteria. Calpains belong to the papain superfamily of cysteine proteases and have weak similarity to papains and cysteine cathepsins, although it is clearly less significant than the similarities between calpains. In this superfamily, calpain may evolutionarily be the oldest branch.⁸⁹⁾

Considering this situation, it is reasonable to define calpain mainly by the aa sequence in relation to the most-characterized mammalian μ - and m-calpains. Therefore, this review uses the following simple but clear definition: *calpains are defined as proteins that have aa sequences significantly similar to the protease domain of the human μ -calpain large subunit* (replacing μ -calpain with m-calpain in this definition would give the same result). Additional characteristics reflect calpain classifications: the conventional, classical, non-classical, ubiquitous, and/or tissue-specific calpains should be discussed separately when considering their physiological functions. In other words, sequence similarity does not necessarily reflect functional similarities between calpain species.

According to this definition, humans have 15 genes that encode calpains (Figs. 3 and 4). In other species, *Schistosoma mansoni* (Fig. 5), *Caenorhabditis elegans* (Fig. 6), *Anopheles gambiae* (Fig. 7), *Drosophila melanogaster* (Fig. 7), *Arabidopsis thaliana* (Fig. 1), *Emericella (Aspergillus) nidulans*




Fig. 3. Phylogenetic tree and schematic structures of human calpain-related molecules. A. Phylogenetic tree of human calpains. Method: The tree was drawn by the neighbor-joining/bootstrap method after aligning all the sequences using MAFFT v6.240 (at <http://align.genome.jp/mafft/>, strategy: E-INS-i). Atypical calpains are more divergent than classical calpains. The PalB subfamily consists of the strict PalB group, the TRA-3 group, and the CAPN10 group. B. Schematic structures: Black and green letters indicate ubiquitous and tissue/organ-specific calpains, respectively. See also Fig. 4. Symbols: L and XL, N-terminal and extended N-terminal regions of calpastatin. See Fig. 1 legend for other symbols.

| Gene | Chromosome location | Gene product | Aliases | Protease activity ^{a)} | Domains ^{b)} | | | Expression | Phenotype of gene deficiency in mice | Note |
|----------------------------|---------------------|----------------------|--|---------------------------------|-----------------------|----|-----|---|--------------------------------------|---|
| | | | | | C2L ^{c)} | C2 | PEF | | | |
| Catalytic subunits | | | | | | | | | | |
| <i>CAPN1</i> | 11q13 | CAPN1 | μ-calpain large subunit (μCL), μ80K, μCANP/calpain-I large subunit, calpain-1 | + | + | - | + | ubiquitous | platelet dysfunction | μ-calpain dimerizes with CAPNS1/30K |
| <i>CAPN2</i> | 1q41-q42 | CAPN2 | m-calpain large subunit (mCL), m80K, mCANP/calpain-II large subunit, calpain-2 | + | + | - | + | ubiquitous (except for erythrocytes) | embryonic lethal | m-calpain dimerizes with CAPNS1/30K |
| <i>CAPN3</i> | 15q15.1-q21.1 | CAPN3 | p94, calpain-3, calpain-3a, nCL-1 | + | + | - | + | skeletal muscle | muscular dystrophy | termination codon in human first exon |
| | | (CAPN3b, ...) | Lp82, Lp85, etc. | + | + | - | + | lens, retina | | Ups, Tps, and Mps are transcribed from an alternative promoter |
| | | CAPN3c, ... | p94:Δex15, Tp36, Mp18, Up84 etc. | + | + | - | + | ubiquitous | | |
| <i>CAPN5</i> | 11q14 | CAPN5 | hTRA-3, nCL-3, calpain-5 | + | + | + | - | ubiquitous (abundant in testis and brain) | sudden death ? | nematode TRA-3 homolog |
| <i>CAPN6</i> | Xq23 | CAPN6 | calpamodulin, CANPX, calpain-6 | - | + | + | - | embryonic muscles, placenta | n.d. | nematode TRA-3 homolog; the active site Cys->Lys in all mammals, and His->Tyr in mice and below; authentic C-H-N sequences in birds and below |
| <i>CAPN7</i> | 3p24 | CAPN7 | PalBH, calpain-7 | + | ++ | - | - | ubiquitous | n.d. | <i>Aspergillus</i> PalB and yeast Cpl1 homolog |
| <i>CAPN8</i> | 1q41 | CAPN8 | nCL-2, calpain-8, calpain-8a | + | + | - | + | gastrointestinal tracts | stress-induced gastric ulcer | G-calpain is a complex of CAPN8 and CAPN9 |
| | | CAPN8b | nCL-2' | + | - | - | - | gastrointestinal tracts | | |
| | | CAPN8c | nCL-2:Δex10,17-18 | + | + | - | +/- | gastrointestinal tracts | | |
| <i>CAPN9</i> | 1q42.11-q42.3 | CAPN9 | nCL-4, calpain-9, calpain-9a | + | + | - | + | gastrointestinal tracts | stress-induced gastric ulcer | G-calpain is a complex of CAPN8 and CAPN9 |
| | | CAPN9b | nCL-4:Δex8 | + | + | - | + | gastrointestinal tracts | | |
| <i>CAPN10</i> | 2q37.3 | CAPN10 | calpain-10, calpain-10a | + | ++ | - | - | ubiquitous | no significant phenotype | SNP is associated with NIDDM |
| | | CAPN10b, -10c, ... | | n.d. | +/- | - | - | | | |
| <i>CAPN11</i> | 6p12 | CAPN11 | calpain-11 | n.d. | + | - | - | testis | n.d. | |
| <i>CAPN12</i> | 19q13.2 | CAPN12 | calpain-12a | n.d. | + | - | + | hair follicle | n.d. | |
| | | CAPN12b, 12c, ... | | n.d. | + | - | +/- | | | |
| <i>CAPN13</i> | 2p22-p21 | CAPN13 | calpain-13 | n.d. | + | - | + | ubiquitous | n.d. | |
| <i>CAPN14</i> | 2p23.1-p21 | CAPN14 | calpain-14 | n.d. | + | - | + | ubiquitous | n.d. | |
| <i>CAPN15/SOLH</i> | 16p13.3 | CAPN15 | SOLH, calpain-15 | n.d. | - | - | - | ubiquitous | n.d. | <i>Drosophila</i> SOL homolog |
| <i>CAPN16/C6orf103</i> | 6q24.3 | CAPN16 | Demi-calpain, C6orf103, calpain-16 | - | - | - | - | ubiquitous | n.d. | contains only PC1 domain |
| Regulatory subunits | | | | | | | | | | |
| <i>CAPNS1</i> | 19q13.1 | CAPNS1 ^{d)} | CANP/calpain small subunit, 30K, css1, CAPNS1, CAPN4, calpain-s1 | - | - | - | + | ubiquitous | embryonic lethal | regulatory subunit for CAPN1/μCL and CAPN2/mCL |
| <i>CAPNS2</i> | 16q12.2 | CAPNS2 ^{d)} | calpain small subunit 2, 30K-2, css2, CAPNS2, calpain-s2 | - | - | - | + | ubiquitous | n.d. | intron-less gene; <i>CAPNS2</i> overlaps with <i>LPCAT2</i> (lysophosphatidylcholine acyltransferase 2) |
| Calpain inhibitor | | | | | | | | | | |
| <i>CAST</i> | 5q15 | calpastatin | CANP inhibitor | - | - | - | - | ubiquitous | excitotoxicity in neuronal cells | specific inhibitor for μ- and m- calpains; many splicing variants exist |

^{a)} + indicates that the molecule was experimentally shown to have protease activity, and - means that it does not have the active-site cysteine, asparagine, and/or histidine residues.

^{b)} + or - indicates that the molecule has or does not have a corresponding domain, respectively, and +/- indicates that it has one or more but fewer than five EF-hand motifs.

^{c)} Abbreviations: n.d., not yet determined/done; CANPX, calpain on chromosome X; nCL, novel calpain large subunit; LGMD2A, limb-girdle muscular dystrophy type 2A; TRA-3, transform genotypic hermaphrodites; PalB, phosphatase mutants: loss in activity at alkaline pH but normal or increased activity at acidic pH; PalBH, PalB homolog; SNP, single nucleotide polymorphism; NIDDM, non-insulin-dependent diabetes mellitus; SOL, small optic lobes; SOLH, SOL homolog.

^{d)} Note that these gene products are not "calpain" according to the definition of this review.

Fig. 4. Human calpain-related genes.
See also Fig. 3.




Fig. 5. Phylogenetic tree and schematic structures of *S. mansoni* calpains. A. Phylogenetic tree of *S. mansoni* and human calpains. See Fig. 3A for methods. B. Schematic structures. The human calpain most similar to each *S. mansoni* calpain is indicated in parentheses, with the aa sequence identity. The identity is given for each of the tandem classical calpain structures in XP_002577797. USP: ubiquitin-specific peptidase. See Fig. 1 legend for other symbols.




Fig. 6. Phylogenetic tree and schematic structures of *C. elegans* calpains. A. Phylogenetic tree of *C. elegans* and human calpains. See Fig. 3A for methods. B. Schematic structures. The human calpain most similar to each *C. elegans* calpain is indicated in parentheses, with the aa sequence identity. CI and CII stand for conserved regions I and II with unknown function, respectively. See Fig. 1 legend for other symbols.

(Fig. 1), and *Saccharomyces cerevisiae* (Fig. 1) have 7, 14, 7, 4, 1, 2, and 1 calpain genes, respectively. There is no calpain gene in *Encephalitozoon* or

Schizosaccharomyces pombe. In prokaryotes, 53 calpains from 42 bacteria were found among the 914 completely sequenced microbial genomes present in




Fig. 7. Phylogenetic tree and schematic structures of insect calpains. A. Phylogenetic tree of *A. gambiae* and human calpains. See Fig. 3A for methods. *D. melanogaster* (DM) homologs are shown in green letters. B. Schematic structures. The human calpain most similar to each insect calpain is indicated in parentheses, along with the aa sequence identity. The *D. melanogaster* calpain most similar to *A. gambiae* calpain is also indicated. P-rich: Pro-rich region. DIS: *Drosophila*-type insertion sequence. See Fig. 1 legend for other symbols.

A




Fig. 8. *Continued on next page.*

the database so far. Each of the bacterial species has 1–4 calpain genes (Fig. 8). However, most genome-sequenced bacteria, including *Escherichia coli* or any of the archaeobacteria, do not have calpain genes.

2. Conventional calpain—the basics

2.1. Domain structures of conventional calpains. The conventional calpains, μ - and m-calpains, are heterodimers consisting of a common small regulatory subunit (CAPNS1/30K; *ca.* 30 kDa) and a distinct, large (*ca.* 80 kDa) μ - or m-calpain catalytic subunit (CAPN1/ μ CL or CAPN2/mCL,

respectively) (Fig. 3). Calpain-I is an old alternative name for μ -calpain, named as an enzyme, and it is not the same as calpain-1, which is the name often used for a gene product, a subunit of an enzyme. Instead, calpain-I is a heterodimer of CAPN1/ μ CL and CAPNS1/30K. The distinction is similar for calpain-II and calpain-2.

CAPN1/ μ CL and CAPN2/mCL have approximately 60% aa identity. Accordingly, they have almost indistinguishable substrate and inhibitor specificities, and both are almost ubiquitously expressed. However, they differ greatly in their *in vitro*

Fig. 8. Phylogenetic tree and sequence alignment of bacterial calpains.

A. Phylogenetic tree of bacterial calpains and human CAPN1/ μ CL. See Fig. 3A for methods. B. Sequence alignment of the bacterial calpain protease domains and human CAPN1/ μ CL. Only the sequences of the protease domains are compared, with the start and end aa residue numbers indicated, as other sequences and lengths are somewhat divergent. Reversed fonts and gray shadow indicate residues conserved in all sequences or more than half, respectively. Triangles: active site residues. Asterisks: residues involved in Ca^{2+} -binding of the protease domain of human CAPN1/ μ CL. Sequences: 0: *H. sapiens* NP_001185798; 1: *Acaryochloris marina* MBIC11017 YP_001520592; 2: *Actinomyces odontolyticus* ATCC 17982 ZP_02043925; 3: *Actinomyces odontolyticus* F0309 ZP_06609922; 4: *Actinomyces* sp. oral taxon 848 str. F0332 ZP_06161744; 5: *Actinomyces* sp. oral taxon 848 str. F0332 ZP_06161892; 6: *Actinomyces* sp. oral taxon 848 str. F0332 ZP_06162803; 7: *Actinomyces* sp. oral taxon 848 str. F0332 ZP_06162809; 8: *Anabaena variabilis* ATCC 29413 YP_322403; 9: *Bacteroides ovatus* ATCC 8483 ZP_02064250; 10: *Bacteroides* sp. 1.1.14 ZP_06994459; 11: *Bacteroides* sp. 1.1.6 ZP_04847694; 12: *Bacteroides* sp. D2 ZP_05758284; 13: *Bacteroides thetaiotaomicron* VPI-5482 NP_812871; 14: *Brachybacterium faecium* DSM 4810 YP_003153779; 15: *Brachybacterium faecium* DSM 4810 YP_003155356; 16: *Bradyrhizobium* sp. BTAi1 YP_001241302; 17: *Bradyrhizobium* sp. ORS278 YP_001204811; 18: *Brevibacterium mcbrellneri* ATCC 49030 ZP_06806606; 19: *Cyanothecce* sp. PCC 7822 YP_003887732; 20: *Cyanothecce* sp. PCC 7822 YP_003887741; 21: *Frankia alni* ACN14a YP_716900; 22: *Frankia* sp. CcI3 YP_483525; 23: *Frankia* sp. EAN1pec YP_001504490; 24: *Frankia* sp. EuI1c YP_004014028; 25: *Frankia* sp. EUN1f ZP_06411021; 26: *Gemmata obscuriglobus* UQM 2246 ZP_02733073; 27: *Geobacter lovleyi* SZ YP_001952615; 28: *Gloeobacter violaceus* PCC 7421 NP_927086; 29: *Granulibacter thesedensis* CGDNIH1 YP_745367; 30: *Herpetosiphon aurantiacus* ATCC 23779 YP_001544076; 31: *Herpetosiphon aurantiacus* ATCC 23779 YP_001545619; 32: *Herpetosiphon aurantiacus* ATCC 23779 YP_001546212; 33: *Legionella longbeachae* D-4968 ZP_06188344; 34: *Legionella longbeachae* NSW150 YP_003455673; 35: *Microcystis aeruginosa* NIES-843 YP_001659697; 36: *NC10 bacterium* 'Dutch sediment' CBE68371; 37: *Nostoc* sp. PCC 7120 NP_484413; 38: *Nostoc* sp. PCC 7120 NP_485127; 39: *Nostoc* sp. PCC 7120 NP_487855; 40: *Photorhabdus luminescens* subsp. *laumondii* TTO1 NP_929692; 41: *Planctomyces limnophilus* DSM 3776 YP_003628665; 42: *Porphyromonas gingivalis* AAA25652; 43: *Porphyromonas gingivalis* W83 NP_905271; 44: *Rhodopseudomonas palustris* BisA53 YP_780017; 45: *Salinispora tropica* CNB-440 YP_001157346; 46: *Segniliparus rotundus* DSM 44985 YP_003659885; 47: *Streptomyces albus* J1074 ZP_06592718; 48: *Streptomyces viridochromogenes* DSM 40736 ZP_07305717; 49: *Synechococcus* sp. RS9916 ZP_01472825; 50: *Synechococcus* sp. RS9916 ZP_01472826; 51: *Synechococcus* sp. RS9916 ZP_01472827.

Ca^{2+} -requirement (μM versus mM) for proteolytic activity. Their functions are fundamental and essential, and are implicated in the regulation of signal transduction systems,^{90,91} cell motility,^{21,92} membrane repair,^{93,94} and apoptosis,^{95,96} although their precise roles in these events remain elusive.

The catalytic and regulatory subunits of conventional calpains can be divided into several domains (Fig. 3). The N-terminal anchor α -helix of the catalytic large subunit (also called anchor region or domain I) is autolyzed when activated by Ca^{2+} , resulting in its ability to function at a lower Ca^{2+} concentration,^{81,97–99} with a different substrate specificity,¹⁰⁰ and subunit dissociation in some cases^{100–103} but not others.^{104–106} Therefore, autolysis of this anchor helix is one of the important regulatory mechanisms for calpain activity.

In isolation, m-calpain's protease (CysPc) domain (also called domain II), has Ca^{2+} -dependent protease activity.¹⁰⁷ 3D structural analyses in the absence of Ca^{2+} showed that the CysPc domain is further divided into two core domains, PC1 (also called D-I or subdomain IIa) and PC2 (D-II or IIb) (Fig. 3). These two core domains fold into a single functional CysPc domain when bound to Ca^{2+} (Fig. 9).^{108–113} In other words, calpain remains structurally inactive in the absence of Ca^{2+} . This is reasonable, because calpain resides in the cytosol in

direct contact with many other proteins, and its activity must be strictly regulated.

Ca^{2+} also binds to C2 domain-like (C2L) domain (also called domain III)¹¹⁴ and penta-EF-hand (PEF) domain (domain IV) of the large catalytic subunit (PEF(L) domain), and that of the small regulatory subunit (PEF(S), also called domain VI).^{115,116} While C2L and the C2 domains have no sequence similarity, their 3D-structure is very similar (β -sandwich structure composed of eight anti-parallel β -strands). Two PEF domains (PEF(L) and PEF(S)) each contain five EF-hand motifs.^{48,117–119} The fifth EF-hand motif of each domain contributes to heterodimer formation.¹²⁰ It should be noted that the N-terminal anchor helix is in contact with the second EF-hand motif (EF-2) of PEF(S) of CAPNS1/30K, and the interaction is broken either by Ca^{2+} -binding to the EF-2 or by autolysis of the N-terminus upon activation.

Gly-rich (GR) domain (also called domain V) in the N-terminus of CAPNS1/30K contains hydrophobic Gly-clusters, most of which are autolyzed during activation. This domain was invisible in the 3D structure, indicating a very "soft" structure. In humans, the *CAPNS2* gene encodes a regulatory subunit homolog composed of GR and PEF(S) domains, whose physiological roles remain unclear.¹²¹




Fig. 9. Schematic of the 3D structure of inactive and active m-calpain. Surface-type schematic 3D structures of inactive (Ca^{2+} free) and active (Ca^{2+} - and calpastatin-bound) forms of m-calpain using PDB data, 1KFX¹⁰⁹) and 3DF0.¹¹¹) The oligopeptides represented by the yellow ribbon + ball-and-stick indicate portions of calpastatin bound to active m-calpain. The dotted lines indicate portions that were too mobile for the 3D structure to be determined. The active protease domain (CysPc) is formed by the fusion of core domains PC1 and PC2 upon the binding of one Ca^{2+} to each of the core domains. The active site is circled in black. Blue balls represent Ca^{2+} (not all are visible).




Fig. 10. A comparison of 3D structures of the Ca^{2+} -binding sites of the CAPN1/ μ CL protease domain IIa and SLO1 (Ca-bowl). Ca^{2+} -binding sites in (A) the CAPN1/ μ CL protease domain (2R9F) and (B) the high-conductance voltage- and Ca^{2+} -activated K^+ channel (BK or SLO1 channel) (3MT5). (C) The structures were superimposed for comparison.¹²⁶)

2.2. Activation and substrates. Ca^{2+} is required for the activation of conventional calpains, raising the question of which part of the calpain

molecule contains the Ca^{2+} -binding site that provokes calpain activation. It was previously believed that PEF domains were responsible for activating calpain

in response to Ca^{2+} , as they are the only Ca^{2+} -binding sites visible in the primary structure.¹³⁾ However, the 3D structure of calpain revealed that the PEF domains are somewhat distant from the protease domain (Fig. 9). CD spectrum measurements found no significant changes in calpain's secondary structure upon activation by Ca^{2+} binding.¹²²⁾ Biochemical analysis of recombinant m-calpain protease domains revealed that they respond directly to Ca^{2+} .¹⁰⁷⁾ Finally, the 3D structures of these domains showed that Ca^{2+} binds directly to PC1 and PC2, to activate conventional calpain (Fig. 9).^{112),113)}

A number of substrates have been reported for calpains, including protein kinases, phosphatases, phospholipases, cytoskeletal proteins, membrane proteins, cytokines, transcription factors, lens proteins, and calmodulin-binding proteins, but few of them have been shown to be physiological. Due to its complexity and "finesse", the exact rules governing calpain substrate specificity remain incomplete, although bioinformatics approaches revealed some cleavage-site preferences^{123),124)} (see also <http://calpain.org>). Substrate 3D structures also seem to be important factors for many cases.

2.3. Novelties deduced from the calpain 3D structure. The classical calpains CAPN1/ μ CL, CAPN2/mCL, and CAPN9/nCL-4 have highly conserved protease domains (59–71% aa identity); however, the Ca^{2+} -bound 3D structures of these domains reveal distinguishing features.^{112),113),125)}

First, comparing the m-calpain protease domains in the presence and absence of Ca^{2+} , all the α -helices and β -strands are conserved, consistent with previous CD spectrum results.¹²²⁾ Both PC1 structures can be aligned very closely ($\text{C}\alpha$ root-mean-squared deviation (r.m.s.d.) = 1.72 Å), whereas those of PC2 are moderately aligned (r.m.s.d. = 3.78 Å). As predicted from PC2's Ca^{2+} -unbound 3D structure, Ca^{2+} binding would cause it to be rotated and translated by 50° and 6 Å, if the alignment of PC2 is fixed. Surprisingly, a Ca^{2+} -induced Ca^{2+} -binding site (CBS-1 and CBS-2) is generated in each core domain, and Ca^{2+} binding to these sites causes the above-described PC2 rotation and translation for activation. These Ca^{2+} -binding sites have a novel and unique structure. Only recently, one structure was shown to have similarity with this site: the 3D structure of the " Ca^{2+} -bowl" of the high-conductance voltage- and Ca^{2+} -activated K^+ (BK or SLO1) channel has a similar Ca^{2+} -binding geometry to the μ -calpain CBS-1, although they share no similarity in their primary sequences (Fig. 10).¹²⁶⁾

Second, the active-site cleft is deeper and narrower than that of papain. Due to this constraint, the substrate must be in a fully extended conformation with its backbone stretched; this was verified by the 3D structures of the Ca^{2+} -bound active μ -calpain protease domain co-crystallized with leupeptin or E64.¹²⁷⁾ This finding explains calpain's preference for proteolyzing inter-domain unstructured regions.

Third, although the active protease domains of CAPN1, 2, and 9 are highly similar, a reversible intrinsic inhibitory/safety mechanism exists only for CAPN2/mCL. W106, which is adjacent to the active-site C105 of CAPN2/mCL, positions to interfere with substrates' access to the active site; in contrast, this mechanism does not occur in rat CAPN1/ μ CL.¹¹³⁾ Moreover, CAPN9/nCL-4 has a novel intrinsic inhibitory mechanism, *i.e.*, the misalignment of the catalytic residues.¹²⁵⁾ These findings together show that the activation/latency mechanisms of calpains, regardless of their highly conserved sequences, are somewhat specialized for each molecule, and are controlled by inter-domain interactions. In the whole 3D structure of m-calpain,^{110),111)} the above-mentioned W106 mechanism¹¹³⁾ was not found, indicating that inter-domain interactions impose extra controls over molecule-specific mechanisms.

Small-angle X-ray scattering analysis revealed direct semi-micro images of both active and inactive whole calpain molecules in the presence of E64,¹⁰⁵⁾ showing the following: in the presence of 100 μM Ca^{2+} and E64, human erythrocyte μ -calpain did not aggregate and had similar dimensions as in the absence of Ca^{2+} (maximal diameter = 120→130 Å, gyration radius = 36→39 Å). This was consistent with the previous CD spectrum results and with the 3D structure of CAPNS1/30K, whose conformation changes little in the presence or absence of Ca^{2+} .^{117),119)} On the other hand, a small but significant change was observed in the protease domain; it appeared to compact somewhat and partly detach from C2L domain.

In other words, upon activation, the protease domain dynamically changes its conformation without changing its secondary structure. This feature was finally confirmed when the whole m-calpain 3D structure was determined by co-crystallization with calpastatin repeat 1 or 4 (see 2.6) and Ca^{2+} ,^{110),111)} revealing one of the mechanisms used by calpastatin to inhibit conventional calpain activity. G174/G613 of rat calpastatin repeat 1/4, respectively, which is near the conserved TIPPLYR (179–185/618–624) motif in the reactive site of calpastatin, forces a




Fig. 11. Calpastatin region B binds and inhibits calpain.

Enlarged view of the interaction between calpastatin (yellow) and the catalytic cleft in the protease core domain PC1 (pink)–PC2 (red) of m-calpain (the 3D structure is from 3DF0). Calpastatin binds in the substrate orientation indicated by positions P3 to P3'. At P1, calpastatin distorts from the substrate path and projects residues 174–178 or 613–617, which form a kink between the P2 and P1' anchor sites.

kink (G174-IKE-G178/G613-ERD-D617) between its flanking residues, L173/L612 at the catalytic S2 site and T179/T618 at S1', to loop-out not to interact with the active site of calpain. A172/K611, L173/L612, T179/T618, I180/I619, and P181/P620 fit into the S3, S2, S1', S2', and S3' pocket of calpain's active site, respectively (Fig. 11).^{110,111} Therefore, calpastatin interacts tightly with calpain, but no proteolysis occurs. The importance of this “kink” in resisting proteolysis by calpain was confirmed by calpastatin mutagenesis studies. The deletion of K176, E177, or both, or the insertion of Phe between L612–G613, allows mutant calpastatins to be proteolyzed by calpain.^{110,111}

2.4. Genetic studies of conventional calpains.

The first calpain studies using genetically modified

mice were reported in 2000, when Arthur *et al.* and Zimmerman *et al.* independently demonstrated that disrupting the mouse gene (*Capns1*) for the conventional calpain regulatory subunit CAPNS1/30K causes embryonic lethality before E11.5.^{128,129} These reports showed that conventional calpains were necessary for mammalian life, which greatly encouraged researchers struggling to understand this enigmatic enzyme. The disruption of CAPNS1/30K causes the down-regulation of both the CAPN1/ μ CL and CAPN2/mCL proteins, indicating that CAPNS1/30K is required for the stable presence of both calpain catalytic subunits *in vivo*, and probably functions as an intramolecular chaperone. *In vitro*, CAPN2/mCL alone (without CAPNS1/30K) shows full proteolytic activity after being denatured and

then renatured by a long incubation with PEG or GroE.¹³⁰⁾ In cells, unfolded calpain large subunits are probably degraded by other proteases before they form active conformations.

Disrupting mouse *Capn1* or *Capn2* leads to contrasting results: *Capn2*^{-/-} mice die before the blastocyst stage, whereas *Capn1*^{-/-} mice appear normal and are fertile.^{55),131)} This suggests that μ - and m-calpain differ in their function and/or expression level, at least at specific developmental stage(s). *Capns1*^{-/-} embryonic stem (ES) cell growth and adhesion were not noticeably different from those of wild-type ES cells, while *Capn2*^{-/-} ES cells could not be obtained even after extensive screening.⁵⁵⁾

Cells from *Capns1*^{-/-} mice served as ideal tools to unequivocally demonstrate the role of calpain in specific cellular events: Dourdin *et al.* showed that calpain is required for cell migration;¹³²⁾ Mellgren *et al.* showed that calpain is required for the rapid, Ca²⁺-dependent repair of wounded plasma membrane;¹³³⁾ and Demarchi *et al.* showed that calpain is required for macroautophagy.¹³⁴⁾

2.5. Expression of conventional calpains.

Although *CAPN1* and *CAPN2* shows ubiquitous distribution, their gene expression patterns are slightly different in that *CAPN1* is rather constantly expressed whereas expression of *CAPN2* varies depending on tissues.^{135),136)} The promoter region of *CAPN2* lacks typical promoter elements such as TATA and CAAT boxes and has ca. 70% GC content in the -300 to -20 region with several potential GC boxes.¹³⁶⁾ The upstream region contains at least four tandemly reiterated regulatory sequences, each of which negatively regulates *CAPN2* expression.¹³⁶⁾ These features are common to several house-keeping genes such as adenosine deaminase¹³⁷⁾ and hypoxanthine phosphoribosyltransferase,¹³⁸⁾ and support that *CAPN2* gene is a typical house-keeping gene that shows ubiquitous and relatively low-level expression. On the other hand, *CAPN2* promoter region also has potential AP-1 binding site, which is responsible for phorbol ester-mediated induction of gene expression.^{139),140)} Indeed, proerythroblastic K562 cell line cells were reported to show upregulation of *CAPN2*/mCL expression by phorbol 12-myristate 13-acetate.¹⁴¹⁾

Although *CAPN1* gene has not been examined in literature, human genomic sequence showed that *CAPN1* has features similar to those of *CAPN2*, *i.e.*, no significant TATA and CAAT boxes with ca. 70% GC content in the -1000 to -1 region and with potential AP-1 binding site at the -80 position.

Analysis of *CAPNS1* gene again revealed similar features:¹⁴²⁾ the promoter region has G/C-rich sequences with several G-C boxes but without TATA or CAT box. Thus, three genes encoding subunits of the conventional calpains are all regarded to be regulated in the same manner as other house-keeping genes.

2.6. Structure and function of calpastatin.

Calpastatin is the only endogenous inhibitor protein for the conventional calpains. It is highly effective and specific, and does not inhibit any other enzymes. Calpastatin has an inhibitor unit that is repeated four times (Fig. 3B), and each unit inhibits one calpain molecule—although their inhibitory efficiencies vary.^{143)–145)} Calpastatin inhibits both μ - and m-calpains with similar efficiencies. Among other calpain homologs, CAPN8/nCL-2 and CAPN9/nCL-4, but not CAPN3/p94, are inhibited by calpastatin *in vitro*.^{53),146),147)}

Disruption of *Cast*, the mouse gene for calpastatin, does not produce a significant phenotype under normal, unstressed conditions.¹⁴⁸⁾ This suggests that conventional calpains are not normally activated dynamically, and that calpastatin is dispensable as a safety for calpain regulation. In accordance with this idea, intra-hippocampal injection of kainic acid (KA), which causes apoptotic neuronal cell death by excitotoxicity, resulted in significantly more DNA fragmentation in *Cast*^{-/-} mice than in wild-type mice.¹⁴⁸⁾ Moreover, the KA effect was reduced in transgenic (Tg) mice overexpressing calpastatin in neuronal cells, whereas Tg mice overexpressing the baculoviral caspase inhibitor p35 showed no change.¹⁴⁹⁾ These results indicated that kainic acid-induced apoptotic neuronal cell death is mediated by the conventional calpains, and that caspases are not involved in this process.

Spencer and colleagues developed Tg mice that overexpress calpastatin in muscles.^{150),151)} These mice appear healthy, without observable body or muscle mass changes or gross physiological, morphological, or behavioral defects. These mice were crossed with *mdx* mice, which have a nonsense *Dmd* mutation that causes mild muscular dystrophic phenotypes due to a lack of dystrophin; the dystrophic phenotype was significantly ameliorated in the resulting calpastatin-overexpressing *mdx* mice.¹⁵⁰⁾ Even in wild-type mice, calpastatin overexpression slows muscle atrophy during muscle unloading.¹⁵¹⁾

Studies using Tg mice also impact on food science, *e.g.*, the postmortem tenderization of muscles. Both calpastatin-overexpressing¹⁵²⁾ and *Capn1*^{-/-} mice¹⁵³⁾

showed reduced postmortem proteolysis of muscle proteins. Thus, calpain system has drawn attention to one of the targets for meat quality control.

3. Classification of calpain superfamily members

3.1. Classical and non-classical calpains.

Calpains have a wide variety of domains in addition to their protease domain, such as C2L and PEF domains (Fig. 1). The calpain superfamily is divided into several subfamilies according to the structures of these additional domains. Since the human conventional calpain catalytic subunits (CAPN1/ μ CL and CAPN2/mCL) are the reference point for calpain structure, calpains having a similar domain structure are called "classical" calpains (Fig. 3) in contrast to "non-classical" ones described below.^{46),154)} Classical calpains are defined as those having C2L and PEF domains in addition to the CysPc domain. Accordingly, non-classical calpains are those having not both of C2L and PEF domains. Among the 15 human calpains, nine are classical and six are non-classical (Fig. 3).

Human classical calpains include CAPN1/ μ CL, CAPN2/mCL, CAPN3/p94, CAPN8/nCL-2, CAPN9/nCL-4, and CAPN11–14, most of which are conserved among vertebrates; fishes have a duplicate set of most of them.⁸⁷⁾ In invertebrates, only a few classical calpains have been identified. The blood fluke *S. mansoni* has four (Fig. 5),¹⁵⁵⁾ the fruit fly *D. melanogaster* has three: CALPA/Dm-calpain, CALPB, and CALPC,³⁴⁾ and the African malaria mosquito *A. gambiae* has five (Fig. 7). No classical calpain homologs have been found in *C. elegans* (Fig. 6), trypanosomes,¹⁵⁶⁾ plants, fungi, or *S. cerevisiae*.

Non-classical calpain CysPc domains have aa identities with each other ranging from <30% to >75%, and they may have functional domains on the N- and/or C-terminal sides of CysPc domain (Figs. 1 and 3). These include transmembrane (TM), Zn-finger, and conserved domains with unknown functions such as SOH (SOL homology) domain. This cohort also includes several alternatively spliced products of classical calpain genes, such as *Capn8* and *CalpA* (Fig. 7B). Thus, non-classical calpains probably function differently from classical calpains, and not all are Ca²⁺-dependent. These features, together with the organization of mammalian calpain genes,¹⁵⁷⁾ strongly suggest that calpain molecules were generated evolutionarily by combining an ancestral calpain-type cysteine protease gene with genes encoding other functions.

3.2. Subfamilies. Non-classical calpains contain several subfamilies grouped according to their additional domain structures. The PalB subfamily is the most evolutionarily conserved, and is found in humans, yeasts, fungi, protists, all insects except *Drosophila*, and nematodes, but not plants (Fig. 12).^{63),64),158)–161)} PalB homologs commonly contain two C2L domains in tandem, each of which diverge greatly or moderately from that of conventional calpains, and have a conserved microtubule-interacting and transport (MIT) motif(s) at the N-terminus (Fig. 3).

Human CAPN10, the representative product of *CAPN10* from the longest transcript,¹⁶²⁾ also has both conserved and diverged C2L domains in succession at the C-terminus. The CAPN10 homologs are only found in vertebrates, and have no MIT domain, but they can be included in the PalB subfamily (Fig. 3).⁴⁶⁾

TRA-3 is found in nematodes and vertebrates, but not in insects or lower organisms (Fig. 13). Compared with classical calpains, TRA-3 homologs have a C2 domain at the C-terminus, instead of a PEF domain (Figs. 3 and 13). Therefore, TRA-3 homologs can also be included in the PalB subfamily, with its structural consensus of two tandem C2L/C2 domains at the C-terminus. TRA-3 homologs, like CAPN10 homologs, have no MIT domain.

The evolutionarily conserved SOL subfamily occurs in almost all animals, including humans, insects, and nematodes, and also in green algae. SOL homolog structures are characterized by having various numbers of Zn-finger motifs in the N-terminal domain. They also share an SOH domain (Figs. 1 and 3).

Plant calpains, called phytocalpains,¹⁶³⁾ were first described in 2002.⁶²⁾ The maize calpain homolog, DEK1 (defective kernel 1), is involved in aleurone cell development. DEK1 homologs are found in various other plants, including rice plants and *Arabidopsis*.^{164)–166)} They have one C2L domain at the C-terminus and a TM domain at the N-terminus. A calpain that is highly similar to DEK1 is also found in *Tetrahymena thermophila* (Fig. 14A). Some of the nematode calpains (CLP-3, -4, -6, and -7) have a domain structure (CysPc-C2L-COOH) similar to that of DEK1, but they do not have a TM domain (Fig. 6). These calpains are all grouped as the DEK1 subfamily.

The first member of the calpain bacterial subfamily was reported in 1992 from *Porphyromonas gingivalis*; it is called tpr (thiol protease).¹⁶⁷⁾ Sub-




Fig. 12. Continued on next page.

sequent bacterial genome projects revealed several bacterial calpain homologs. These are the most divergent calpain species, sharing similarity only in the protease domain (Fig. 8). Although *E. coli* was once reported to have a calpain-like protease,¹⁶⁸ its sequence rather belongs instead to the narrowly defined papain superfamily (clan CA-C1). In summary, all the calpain subfamily consensus structures are shown in Fig. 14B.

3.3. Ubiquitous and tissue-specific calpains.

In addition to their structural features, mammalian calpains are also independently categorized according to their tissue and organ distribution.¹⁶⁹ CAPN1/μCL, CAPN2/mCL, CAPN5/hTRA-3, CAPN7/

PalBH, CAPN10, CAPN13, and CAPN15/SOLH are ubiquitously expressed, whereas CAPN3/p94, CAPN8/nCL-2, CAPN9/nCL-4, CAPN6, CAPN11, and CAPN12 are expressed in specific tissues or organs (Figs. 3 and 4). Widely accepted assumption is that ubiquitous calpains play fundamental roles in all cells, whereas tissue-specific calpains have tissue-specific roles. In fact, defects in ubiquitous calpains may be lethal, as in *Capn2*^{-/-} mice,⁵⁵ whereas defects in tissue-specific calpains may cause tissue-specific diseases, like the muscular dystrophy caused by defective *CAPN3*.⁵⁶

At the same time, under various disease/damaged conditions such as muscular dystrophies,

Fig. 12. Phylogenetic tree and sequence alignment of PalB calpains.

A. Phylogenetic tree of strict PalB calpains and human CAPN1/ μ CL. See Fig. 3A for methods. B. Sequence alignment of PalBs and human CAPN1/ μ CL (the start and end aa residue numbers are indicated). Triangles: active site residues. Asterisks: residues involved in Ca^{2+} -binding of the protease domain of CAPN1/ μ CL. Arrows: domain boundaries. Deletion (-) and multiple (+) residue positions are indicated. Residue numbers of human CAPN1/ μ CL are given above the sequences. Residues conserved in all the sequences, all except human CAPN1/ μ CL, and in more than half the sequences are indicated by black reversed fonts, gray reversed fonts, and gray shadow, respectively. Sequences: 0: *H. sapiens* (human) CAPN1/ μ CL (NP_005177); 1: *H. sapiens* CAPN7/PalBH (NP_055111); 2: *X. tropicalis* (frog) (NP_998853); 3: *D. rerio* (zebrafish) (NP_001128580); 4: *Ciona intestinalis* (sea squirt) (XP_002121253); 5: *Tribolium castaneum* (red flour beetle) (XP_967682); 6: *Apis mellifera* (honey bee) (XP_001121978); 7: *Acyrtosiphon pisum* (pea aphid) (XP_001945029); 8: *A. gambiae* str. PEST (malaria mosquito) (XP_309874); 9: *S. mansoni* (XP_002579452); 10: *C. elegans* (NP_497964); 11: *T. brucei* TREU927 (XP_828540); 12: *L. major* (CAJ06528); 13: *Coprinopsis cinerea* okayama7#130 (common ink cap) (EAU91907); 14: *E. nidulans* FGSC-A4 (XP_657860); 15: *C. albicans* SC5314 (EAL03290); 16: *S. cerevisiae* (Q03792).




Fig. 13. Continued on next page.

cardiomyopathies, traumatic ischemia, and lissencephaly, conventional calpains tend to be overactivated—probably because the intracellular Ca^{2+} homeostasis is compromised by the disease—and that overactivity often exacerbates the disease. Conventional calpain inhibitors are currently used to prevent the progression of such diseases, a therapeutic technique first applied by Suzuki and colleagues.^{58),79),170),171)}

3.4. Expression of tissue-specific calpains.

Tissue-specificity of tissue-specific calpains should

be regulated by their promoter regions. Human, mouse and rat *CAPN3/Capn3* upstream regions have highly conserved structures containing TATA-box and several E-boxes, which probably are responsible for muscle-specific expression of *CAPN3/Capn3*.¹⁷²⁾ Intriguingly, *CAPN3/Capn3* has an alternative upstream promoter, which drives ubiquitous expression of *Up86 etc.* starting from an alternative initial exon with slight abundance in the brain and blood cells.^{173)–175)} Moreover, rodent *Capn3* has additional alternative promoter and initial exon between exons




Fig. 13. Continued on next page.

Fig. 13. Phylogenetic tree and sequence alignment of TRA-3 calpains.

A. Phylogenetic tree of TRA-3 calpains and human CAPN1/ μ CL. See Fig. 3A for methods. B. Sequence alignment of TRA-3 and human CAPN1/ μ CL (the start and end aa residue numbers are indicated). Sequences: 0: *H. sapiens* CAPN1/ μ CL NP005177; 1: *H. sapiens* CAPN5 NP004046; 2: *M. domestica* CAPN5 XP001377962; 3: *G. gallus* CAPN5 XP417278; 4: *X. tropicalis* CAPN5 NP001006736; 5: *D. rerio* CAPN5 NP001073476; 6: *D. rerio* CAPN5-2 XP001345114; 7: *H. sapiens* CAPN6 NP055104; 8: *M. domestica* CAPN6 XP001366895; 9: *G. gallus* CAPN6 XP420313; 10: *X. tropicalis* CAPN6 XP002938834; 11: *X. tropicalis* TRA-3 XP002941290; 12: *D. rerio* TRA-3 XP001339053; 13: *Strongylocentrotus purpuratus* CAPN5 XP001178522; 14: *S. mansoni* XP002578116; 15: *C. elegans* TRA-3/CLP-5 NP502751. For other explanation, see Fig. 12B.

1 and 2, and these are responsible for lens-specific expression of alternative splicing products of *Capn3* such as Lp82 and Rt88.^{176),177)} However, human *CAPN3* lost lens-specific products by in-frame termination in exon 1'.¹⁷⁸⁾ *Capn3* expression is also noted in testis.¹⁷³⁾

The promoter region of *Capn8* contains two TATA boxes with several GATA boxes, which are probably responsible for stomach expression of *Capn8*.¹⁵⁷⁾ Some cAMP responsive elements is also found in this region. Notably, *CAPN8* and *CAPN2* are close to each other on human chromosome 1q41 (mouse: chromosome 1), and their transcripts have overlapping complementary sequences, suggesting possible interaction with each other.¹⁵⁷⁾ In addition, *Capn8* has additional exon in the middle, which generates an alternative splicing product, CAPN8b/nCL-2'. Its physiological relevance, however, is elusive.^{157),179)} As for other tissue-specific calpain genes, their upstream sequences have not been analyzed in the detail.

4. Unconventional calpain functions in health and disease

4.1. Skeletal muscle-specific calpain. The first tissue-specific calpain, CAPN3/p94, was found in 1989.¹⁸⁰⁾ It is expressed predominantly in skeletal muscle. This classical calpain is approximately 50% identical to CAPN1/ μ CL and CAPN2/mCL (Fig. 3). However, it contains three additional characteristic regions NS, IS1, and IS2, located at the N-terminus, in PC2 domain, and between C2L and PEF domains, respectively. These regions give CAPN3/p94 some unique features.

The most characteristic property of CAPN3/p94 is its extremely rapid autodegradation—its half-life *in vitro* is less than 10 minutes—starting with autolysis in the NS and IS1 regions.^{181)–183)} This autodegradative activity depends on both IS1 and IS2, and conventional calpain inhibitors such as calpastatin and E64 have little effect on it.¹⁸¹⁾ Surprisingly, this autodegradation occurs Na⁺-dependently in the absence of Ca²⁺, establishing

CAPN3/p94 as the first example of an intracellular Na⁺-dependent enzyme.⁵⁴⁾ The physiological relevance of CAPN3/p94's Na⁺-dependency is still unclear, but its substrate specificities differ when it is activated by Ca²⁺ versus Na⁺. For example, LIM domain-binding protein 3, tropomyosin α -1 chain, and α -actinin-3 are candidate *in vivo* substrates for Na⁺-activated CAPN3/p94, whereas troponins T and I and Ras-specific guanine nucleotide-releasing factor 2 are for Ca²⁺ activation.⁵⁴⁾ Furthermore, CAPN3/p94 possesses a nuclear localization-signal-like sequence in the IS2 region, and is sometimes found in the nucleus as well as the cytosol.¹⁸¹⁾ CAPN3/p94 binds specifically to the N2A and M-line regions of the gigantic filamentous muscle protein connectin/titin (Mr > 3,000 kDa), and the binding site for N2A connectin/titin is the N-terminal vicinity of the IS2 region.¹⁸⁴⁾ Its autodegradative activity is almost completely suppressed *in vivo*, most probably by binding to N2A connectin/titin. This region also aligns important subcellular structures of muscle, such as the sarcoplasmic reticulum and T-tubules, and therefore represent the potential space for CAPN3/p94 to function.¹⁸³⁾

In 1995, *CAPN3* mutations were shown to be responsible for limb-girdle muscular dystrophy type 2A (LGMD2A), also called calpainopathy, by an analysis of families of La Réunion Island and by positional cloning.^{56),185)} *Capn3*^{-/-} mice recapitulate a human calpainopathy-like phenotype,^{186),187)} although milder, indicating that calpainopathy is indeed caused by *CAPN3* defects.¹⁸⁶⁾ Calpainopathy appears to be primarily caused by compromised CAPN3/p94 protease activity, rather than by damaged structural properties.¹⁸⁸⁾ This was confirmed by CAPN3/p94 knock-in (*Capn3*^{CS/CS}) mice, which express a structurally intact but inactive CAPN3/p94:C129S mutant, and also present a muscular dystrophy phenotype.¹⁸⁹⁾ Intriguingly, however, *Capn3*^{CS/CS} mice have a less severe phenotype than *Capn3*^{-/-} mice, indicating that the proteolytically inactive CAPN3/p94 retains some function, probably involving its structural properties.¹⁸⁹⁾




Fig. 14. Schematic structures of *T. thermophila* and *Z. mays* DEK1s.

A. The number of TM spanners is taken from prediction programs and may be corrected in the future. Domain names for *Z. mays* DEK1 are taken from ref. 62). B. Structural classification and consensus domain structures of calpain super family members. See Fig. 1 legend for symbols.

Many subsequent family analyses have been reported, identifying substantial numbers of both pathogenic and polymorphic mutations in *CAPN3*.^{190),191)} However, no mutational hot spot has been found, so a genetic diagnosis of calpainopathy is not easy. A total of 2,530 mutations, including 456 unique ones, have been reported in the *CAPN3* gene so far (available at the Leiden Muscular Dystrophy Pages <http://www.dmd.nl/>). Among the 80 gene entries with mutations responsible for muscular dystrophy, *CAPN3* has the third most, coming only after dystrophin and lamin A/C. One characteristic of *CAPN3* gene mutations is that more than half (62.7%) of them are nucleotide substitutions, of which missense mutations are dominant (60.3% among all substitutions, and 37.8% among all pathogenic mutations) (Fig. 15A).

Although some missense mutations found in LGMD2A patients are reported relatively frequently, such as R490Q/W, R769Q, and A236T, most of them are distributed all over the protein, between the N-terminus and C-terminus, including 215 unique mutations at 176 independent loci (21.4%) among the 821 aa residues (Fig. 15B). A comparison of vertebrate *CAPN3*/p94s (including the two gene products from fishes) revealed 241 residues that are conserved in all species (Fig. 16), including 83 of the above 176 missense loci, indicating that conserved aa residues are important for *CAPN3*/p94 functions. Conservation in the *CAPN3*/p94-characterizing regions NS, IS1, and IS2 is relatively low; however, it should be noted that some missense mutations, such as R49C/H, P319L, and S606L, which are in the NS, IS1, and IS2 regions respectively, occur relatively frequently. These observations suggest that sequence conservation of the *CAPN3*/p94-characterizing region is not necessarily required, but its length and a few important aa, usually near the borders of these regions, must be conserved for proper function.

LGMD2A and *CAPN3* mutations are the first—and so far, only—example of a clear cause–effect relationship between human disease and calpain gene mutations.

4.2. Gastrointestinal tract-specific calpains.

CAPN8/nCL-2 and *CAPN9*/nCL-4 are predominantly expressed in surface mucus cells, called pit cells, in the stomach. Smaller amounts are expressed in the goblet cells in the intestines.^{179),192),193)} Among mammalian calpains, *CAPN8*/nCL-2 is most similar to *CAPN2*/mCL (62% identical). Unlike m-calpain, however, *E. coli*-expressed recombinant *CAPN8*/nCL-2 exhibits Ca²⁺-dependent autolytic and casein-

olytic activity without *CAPNS1*/30K, and it forms a homo-di~oligomer *via* C2L domain *in vitro*.¹⁴⁶⁾ *Xenopus laevis* has a *CAPN8*/nCL-2 ortholog named xCL-2, which causes severe developmental defects when disrupted.¹⁹⁴⁾

CAPN9/nCL-4 is another classical calpain homolog specific for gastrointestinal tracts. It is equally similar to all other classical calpains at the aa level, suggesting that *CAPN9*/nCL-4 is the molecule closest to the ancestral calpain species.¹⁹³⁾ *CAPN9*/nCL-4 is reported to be involved in tumorigenesis¹⁹⁵⁾ and in lumen formation in breast epithelial cells induced by an adhesion molecule.¹⁹⁶⁾ Recombinant human *CAPN9*/nCL-4 requires *CAPNS1*/30K for its activity *in vitro*, and such a complex has Ca²⁺-dependent activity that is inhibited by calpastatin and other Cys protease inhibitors, similar to conventional calpains.¹⁴⁷⁾

In contrast, the *in vivo* characteristics of these calpains are rather different. Experiments using *Capn8*^{-/-} and *Capn9*^{-/-} mice showed that neither *CAPN8*/nCL-2 nor *CAPN9*/nCL-4 forms a stable complex with *CAPNS1*/30K; rather, they form a hybrid complex with each other, now designated as gastric calpain (G-calpain).⁵⁷⁾ As is the case for *Capns1*^{-/-}, disruption of either *Capn8* or *Capn9* caused down-regulation of another molecule, suggesting that each species contributes to the stability and the functionality of the other. Both *Capn8*^{-/-} and *Capn9*^{-/-} mice seem healthy under normal conditions, but they are significantly more susceptible to ethanol-induced gastric ulcers.⁵⁷⁾ *CAPN8*/nCL-2 knock-in (*Capn8*^{CS/CS}) mice that express the protease-inactive *CAPN8*/nCL-2:C105S mutant instead of wild-type *CAPN8*/nCL-2 also show stress-induced gastropathy. This indicates that *CAPN8*/nCL-2 and *CAPN9*/nCL-4, in the form of G-calpain, mediate key events for gastric mucosal defense.⁵⁷⁾

A single nucleotide polymorphism (SNP) database search revealed that human *CAPN8* and *CAPN9* have several aa-substituting SNPs.⁵⁷⁾ An *in vitro* expression study indicated that some of these SNPs compromised proteolytic activity of G-calpain. Thus, evaluation of the link between these SNPs and susceptibility to gastrointestinal disorders is on list of things to do.

4.3. The PalB subfamily. PalB was first identified in the ascomycete *E. nidulans* as a product of the gene responsible for the fungi's adaptation to alkaline conditions.⁶³⁾ Its orthologs were then widely identified, from *S. cerevisiae* to humans (Fig. 12).^{64),158)} The structural consensus for PalB




Fig. 15. Pathogenic CAPN3 mutations in calpainopathy patients.

A. Summary of calpainopathy mutation types. B. Calpainopathy missense mutation positions and frequencies (relative to the maximum value). Black vertical lines indicate aa residues that are conserved more than 80% of vertebrates as shown in Fig. 16B. α and β indicate α -helix and β -strand secondary structures of the corresponding m-calpain 3D structure. See Fig. 1 legend for other symbols.




Fig. 16. Phylogenetic tree and sequence alignment of CAPN3 proteins. A. Phylogenetic tree of vertebrate CAPN3. See Fig. 3A for methods. Explicit CAPN3 homologs are found only in vertebrates. B. Sequence alignment of CAPN3s. Fishes have two CAPN3 homologs (type-1 and -2); the latter does not have explicit IS1 or IS2 regions. Sequences: 1: *H. sapiens*, NP_000061; 2: *M. domestic*, ENSMODP00000022330; 3: *G. gallus*, NP_001004405; 4: *A. carolinensis*, ENSACAP00000015362; 5: *X. tropicalis*, ENSXETP00000026925; 6: *Takifugu rubripes* (type-1), ENSTRUP00000046251; 7: *Hippoglossus hippoglossus* (type-1), ACY78226; 8: *Salmo salar* (type-1), NP_001158880; 9: *D. rerio* (type-1), XP_001337065; 10: *T. rubripes* (type-2), ENSTRUP00000016935; 11: *D. rerio* (type-2), ENSDARP00000094244. For other explanation, see Fig. 12B.




Fig. 17. Pal-PacC pathway and corresponding molecules in yeast and human.

Schematics of signaling pathways involving PalB of fungi. In fungi, a membrane protein (PalH) senses ambient pH and transduces the alkaline signal to PalI. An arrestin homolog, PalF, further transduces the signal to PalA, which forms a complex with AnPef1. At the endosomal membrane, PalA forms the PalB active complex to proteolyze PacC at the C-terminus, thus activating this key transcription factor. Activated PacC regulates gene expression under alkaline conditions. This pathway is highly conserved in yeast and possibly conserved in human; corresponding molecules: PalH (fungi)-Rim21 (yeast)-? (human), PalI-Rim9-?, PalF-Rim8-arrestin, PalA-Rim20-Alix/AIP1, PalC-Ygr122w-?, AnPef1-Pef1/Ygr058w-ALG-2, AnSnf7-Snf7/Vps32-CHMP4s, PalB-Rim13/Cpl1-CAPN7/PalBH, and PacC-Rim101-C2H2 Zn-finger proteins? See text for details.

subfamily members is (MIT)₀₋₂-CysPc-C2L-C2L/C2 (Fig. 14B). Strict PalB homologs have (MIT)₂-CysPc-C2L-C2L (the upstream C2L was once called PBH (PalB-homology) because of its highly divergent sequence⁴⁸), whereas CAPN10 and TRA-3 homologs have CysPc-C2L-C2L and CysPc-C2L-C2, respectively.

4.3.1. Strict PalB homologs in fungi. E. nidulans, like many other microorganisms, grows over a wide pH range. The gene locus *palB* was identified from mutants that mimic the gene expression profiles for acid environments even under normal/neutral conditions, corresponding to *pal* genes (*pal* stands for “phosphatase mutants: loss in activity at alkaline pH but normal or increased activity at acidic pH”). PalB is involved in the proteolytic activation of a key transcription factor named PacC, which governs the entire regulation of pH-dependent changes in gene expression, such as the expression of alkaline phosphatase under alkaline conditions. This pH adaptation system is here called the Pal-PacC pathway (Fig. 17).

Mutations in six genes, *palA*, *palB*, *palC*, *palF*, *palH* and *palI*, that mimic the effects of growth at acid pH, and one, *pacC*, that mimics those at alkaline pH, were identified.¹⁹⁷⁻¹⁹⁹ These genes were characterized from 1995, starting with the identification of the *palB* gene product as a novel and divergent calpain homolog.^{63,200-203} Primary structures of these *pal* gene products showed that PalB was a calpain homolog, and PalH and PalI had TM domains. Protein interaction analyses showed that PalF, a distant homolog of mammalian arrestin, binds to the large cytosolic domain of PalH,²⁰⁴ and that PalF’s phosphorylation and ubiquitylation under alkaline conditions depend on PalH and PalI. These findings indicate that PalF–PalH interactions mediate pH signaling, similar to the arrestin–receptor interactions that regulate various processes in mammals, such as photo-signaling.²⁰⁵ These findings strongly imply the existence of an corresponding mammalian signaling pathway(s) involving arrestin and CAPN7/PalBH.

PacC (674 aa) usually forms an inactive, closed conformation, which protects it from activation by intramolecular interactions, and undergoes limited proteolysis to become active.^{206),207)} First, PacC is processed between aa 479–502 by a primary processing protease, PalB, in a pH-regulated manner. This changes PacC to an open form susceptible to proteolysis by the proteasome in a pH-independent constitutive manner, simultaneously creating the processed active form (252–254 aa).^{207),208)} It is thought that ubiquitylation is not involved in this process, which is a rare mode of action for proteasome.^{209)–212)}

4.3.2. Strict PalB homologs in yeasts and mammals. The above-described fungus Pal-PacC pathway is well conserved in budding yeasts. The *S. cerevisiae* PalB ortholog is Rim13/Cpl1. It was first identified in 1993 as one of the *RIM* genes, which reduce the expression of sporulation-specific genes when mutated.^{213),214)} Among the *RIM* gene products, Rim101 corresponds to fungus PacC, and plays a central role in this signaling system, as does PacC. The proteolytic regulation of Rim101 was first discovered in 1997,²¹⁵⁾ and the responsible protease was identified as Rim13/Cpl1 in 1999.⁶⁴⁾ Subsequently, *RIM8*, *9*, *20*, and *21*, which correspond to *palF*, *I*, *A*, and *H*, were identified.^{200),201)}

Saccharomyces' usual environment is rather acidic, and its alkaline adaptations function in near-neutral pH conditions. Therefore, this adaptive system, here called the Rim pathway, has practical importance for human health. Pathogenic and saprophytic yeasts like *Candida albicans* and *Yarrowia lipolytica* also have conserved Rim pathways. For example, *C. albicans* uses this pathway when it infects and invades mammalian skin in neutral pH environments, and disrupting the Rim pathway compromises its pathogenicity.^{216),217)} *Y. lipolytica*^{218),219)} and *Usitlago maydis*²²⁰⁾ are also reported to use an identical system.

The Pal-PacC and Rim pathways relate to membrane trafficking, and involve ESCRT (endosomal sorting complex required for transport) and Vps (vacuolar protein sorting) proteins.^{65),221)–229)} Based on a genomic study that showed Rim and Vps interactions, it is demonstrated that these two proteins had a functional relationship in *S. cerevisiae* and *C. albicans*.^{223),228),230)} Screening of suppressor mutants for *rim21* and *rim9* revealed functional relationships between the Rim and Vps proteins.⁶⁵⁾ These studies clarified components of the Rim signaling cascade and the hierarchy: *RIM8*, *9*, and *21* act upstream of *RIM13/CPL1* and *RIM20*, and

Rim13/Cpl1 proteolyzes Rim101 at the endosomal membranes with the aid of the ESCRT-III subcomplex as a scaffold (Fig. 17).

The Pal-PacC and Rim pathways are the first examples that genetics thoroughly uncovered the molecular components and mechanisms of calpain-mediated systems. These pathways also exemplify that calpains function as modulator proteases, affecting their substrates' functions by limited proteolysis. Theoretical extension of the knowledge of these pathways is anticipated to provide important keys for understanding other calpain systems, including those that involve CAPN7/PalBH.

There are few published reports on CAPN7/PalBH, a mammalian ortholog of PalB.^{158),231)–233)} Many studies have examined other homologous components; for instance, arrestin and Alix/AIP1, which correspond to PalA and F, respectively. It should be noted that interactions among Rim20, Fef1/Ygr058w, and Snf7/Vps32 are conserved in mammals as Alix1/AIP1, ALG-2, and CHMP4, respectively.^{234)–237)} Although multiple interactions of Alix/AIP1 have been reported,^{236),238)} the physiological roles of these proteins and CAPN7/PalBH have not been clarified. Another message from these studies is that it is essential to take ubiquitous unconventional calpains such as CAPN7/PalBH into account along with the conventional calpains when analyzing various biological phenomena, but very few reports achieve this.²³⁹⁾

4.3.3. TRA-3 homologs in the PalB subfamily. TRA-3/CLP-5 is classified into the PalB subfamily in the broad sense, but TRA-3/CLP-5 has a distal C2 instead of C2L domain, and no N-terminal MIT domain (Fig. 14B). The distal C2 domain was once called a T domain (T from TRA-3), because its primary sequence rather diverges from that of other C2 domains. TRA-3/CLP-5 was first identified by genetics as the product of the gene, *tra-3* (transformer: XX animals transformed into males), which is involved in the sex determination cascade of *C. elegans*.⁶¹⁾ Its action as an enzyme has not been characterized; however, its protease activity is necessary for female development in XX hermaphrodites, in the processing of the TRA-2A membrane protein.²⁴⁰⁾ In contrast, TRA-3/CLP-5 and another *C. elegans* calpain, CLP-1, are components of a neuronal necrotic death cascade, upstream of the aspartic proteases ASP-3 and ASP-4, which may correspond to mammalian cathepsins D and E.²⁴¹⁾ An SNP in *tra-3* is also reported to be involved in nematode body-size determination.²⁴²⁾

Mammals have two TRA-3 orthologs, CAPN5/hTRA-3 and CAPN6; both share more than 30% aa identity with TRA-3 (Fig. 13).^{243),244)} CAPN5/hTRA-3 has Ca²⁺-dependent autolytic activity and is sensitive to several calpain inhibitors.²⁴⁵⁾ CAPN5/hTRA-3 is expressed in almost all tissues in varying amounts. Analysis of *Capn5*^{-/-} mice showed that CAPN5/hTRA-3 is expressed in a subset of T cells, but is dispensable for development.²⁴⁶⁾ SNPs of human *CAPN5* were reported to be associated with polycystic ovary syndrome, diastolic blood pressure, and cholesterol levels,^{247),248)} and CAPN5/hTRA-3 and CAPN15/SOLH are upregulated in chronic intermittent, but not constant, heart hypoxia.²⁴⁹⁾

On the other hand, CAPN6 in eutherians (placental mammals) naturally has aa substitutions in the most important residue in the active site triad (C→K in humans, see Fig. 13B), strongly suggesting that eutherian CAPN6 has no proteolytic activity. Interestingly, CAPN6 in marsupialia (*Monodelphis domestica*; gray, short-tailed opossum) and in chicken (*Gallus gallus*) have the active-site residues Cys-His-Asn; moreover, frog (*X. tropicalis*) and fish (*Danio rerio*) have three TRA-3 homologs, which also have the active-site residues. This is in contrast to CAPN11, which is a classical calpain expressed specifically in the testis in eutherians, whereas the *CAPN11* gene is ubiquitously expressed in chicken but was lost in marsupialia. Mouse CAPN6 is predominantly expressed in embryonic muscles, placenta, and in several cultured cells. It is intriguing to consider the physiological function of eutherian CAPN6 and CAPN11, which are expressed in the placenta and testis, respectively. CAPN6 was shown to be involved in regulating microtubule dynamics,²⁵⁰⁾ and was subsequently shown to regulate cell motility in cultured cells,²⁵¹⁾ although the *in vivo* physiological functions of CAPN6 and CAPN11 are still unclear.

4.3.4. CAPN10 homologs in the PalB subfamily. CAPN10 also belongs to the PalB subfamily in the broad sense, but CAPN10 differs from PalB in that it has no N-terminal MIT domain (Fig. 14B). *CAPN10* was identified as a risk factor for non-insulin-dependent diabetes mellitus (NIDDM, type 2 diabetes) in a large-scale genetic association study.¹⁶²⁾ An association was found between G→A SNP (UCSNP-43, the G allele is the at-risk genotype) in intron 3 of *CAPN10* and susceptibility to NIDDM in Mexican Americans and in a Northern European population from the Botnia region of Finland. There is no molecular explanation why this

SNP causes susceptibility to NIDDM, although G/G homozygotes in UCSNP-43 were reported to have reduced *CAPN10* mRNA expression in skeletal muscle.^{252),253)}

The *Capn10* gene is also a candidate gene responsible for a phenotype of the Otsuka Long-Evans Tokushima Fatty rat, a NIDDM animal model.²⁵⁴⁾ Quantitative trait locus (QTL) analyses using *Capn10*^{-/-} mice and two strains, LG/J and SM/J, with low and high obesity phenotypes showed that *Capn10* is included in the obesity QTL, *Adip1*, indicating involvement of CAPN10 in obesity also in mice.²⁵⁵⁾

Studies using *Capn10*^{-/-} or CAPN10-overexpressing Tg mice suggest that CAPN10 is involved in type 2 ryanodine receptor-mediated apoptosis.²⁵⁶⁾ Although the phenotypes of these mice have not been described, they do not have embryonic lethality. Ubiquitous distribution and dynamic changes in CAPN10's cellular localization was shown by a CAPN10-specific antibody.²⁵⁷⁾ CAPN10 was also demonstrated to be involved in GLUT4 vesicle translocation during insulin-stimulated glucose uptake in adipocytes.²⁵⁸⁾ Arrington *et al.* showed that CAPN10, localized to mitochondria, mediates mitochondrial dysfunction by cleaving Complex I subunits and activating the mitochondrial permeability transition.²⁵⁹⁾

4.4. The SOL subfamily. The first genetic calpain study examined the *Drosophila* gene *small optic lobes (sol)*.⁶⁰⁾ A mutation in *sol*, such as *sol*¹⁰⁷⁸, results in the absence of certain classes of columnar neurons in the optic lobes, leading to specific alterations in flight and walking maneuvers. The longest transcripts from this gene encodes a 1,597-aa protein belonging to the calpain SOL subfamily. *Drosophila* SOL is composed of the N-terminal six-Zn-finger motif-containing domain and the C-terminal calpain-like protease domain (Fig. 7B). Most of the protease domain is deleted in the *sol*¹⁰⁷⁸ mutation, demonstrating that SOL as a protease is essential for forming proper optic lobes. Unfortunately, there have been no further reports dissecting the molecular mechanisms of neurodegeneration involving SOL.³⁴⁾ Mammals have one SOL ortholog, CAPN15/SOLH,^{260),261)} but its physiological role is not clear. In parallel with PalB subfamily, this evolutionarily interesting molecules including their mammalian homologs beg further study.

4.5. Phytocalpain. A plant calpain, now called phytocalpain, was first identified in the sugarcane EST database by Correa *et al.*, and was named SC

call.²⁶²⁾ A sequence similarity search using SC call as bait subsequently identified other phytocalpains: from the dicotyledons, *Vitis vinifera* (pinot noir grape), *A. thaliana*, and *Solanum melongena* (egg-plant); from the monocotyledons, *Zea mays* (maize), *Oryza sativa* (rice plant), and *Hordeum vulgare* (barley); and from gymnospermae, *Pinus taeda* (loblolly pine).¹⁶³⁾ A genetic study identified phytocalpain as the *defective kernel 1 (dek1)* gene product, DEK1, required for aleurone cell development in the maize endosperm.⁶²⁾ Since then, DEK1 homologs in *A. thaliana* and *Nicotiana benthamiana* (tobacco) have been found to have similar functions.^{166),263)}

A biochemical study using *E. coli*-expressed recombinant maize DEK1 CysPc-C2L domains showed that its Ca²⁺-activated caseinolytic activity depends on the predicted active site Cys residue, although this activity was much lower than that of the conventional calpains under the study conditions used.¹⁶⁴⁾ DEK1 is the only calpain homolog in the *Arabidopsis* genome,²⁶³⁾ and it is important for regulating growth in this plant. Although the full-length DEK1 protein localizes to membranes, intramolecular autolytic cleavage releases the calpain domain into the cytoplasm, and this domain is sufficient to fully complement *dek1* mutants.^{263),264)}

As shown in Fig. 14A, from the N-terminus of maize DEK1, a possible signal peptide is followed by the first TM region (A), eight-TM region (B1), large loop region (C), third TM domain with 14–15 spanners (B2), a domain of unknown function with no homology to other proteins (D), the calpain protease (CysPc) domain, and C2L domain.^{62),164)} The number of TM spanners of B2 region is predicted to be 15 or 14, respectively, by SOSUI or TMHMM Ver.2.0,^{265),266)} which makes the topology of C region and calpain domains on the same or opposite sides of cell membrane. This topology is very important for DEK1 function, and it should be examined biochemically.

Among the 15 human calpains, the protease domain of non-classical CAPN15/SOLH is the most similar to the DEK1 CysPc domain, with which it shows almost 40% aa sequence identity. Surprisingly, DEK1's C-terminal C2L domain is rather similar to that of classical mammalian calpains, such as CAPN1/ μ CL, CAPN3/p94, and CAPN8/nCL-2 (25–30% identity). Therefore, DEK1 is considered to consist of three evolutionarily different modules: a TM domain, a CysPc domain, and a C2L domain. In addition, it has been suggested that DEK1 homolog of the protista *T. thermophila* (Fig. 14A) resulted

from a lateral gene-transfer event from a green alga-type endosymbiont of ciliates as an explanation for this inter-kingdom similarity.⁴⁶⁾

4.6. Other calpain members. *Leishmania* and *Trypanosoma* have around 20 calpain homologs each, which are likely to contribute to cell morphogenesis, drug resistance, and stress-response mechanisms.^{267)–273)} Some trypanosome calpains have N-terminal domains with weak similarity to calpastatin. As described above, some of the calpain homologs have substitutions in one or more of the well-conserved active-site triad residues, Cys-His-Asn. This non-proteolytic family of calpain homologs includes eutherian CAPN6 (Fig. 3), some of the schistosome (Fig. 5) and nematode (Fig. 6) calpains, insect CALPC (Fig. 7B) and all of the *Trypanosoma* homologs.^{34),241),267)} The generation of these calpain species is interesting from an evolutionary viewpoint, and elucidating their physiological functions will reveal additional roles played by the calpain superfamily, e.g., possible non-proteolytic functions.

Concluding remarks

Gordon Guroff, Darrel E. Goll, Kazutomo Imahori, Koichi Suzuki, and Takashi Murachi established numerous milestones in the study of calpains, including its discovery,¹⁾ finding its relationship to skeletal muscle,⁵⁾ purifying it to homogeneity,⁷⁾ labeling calpain and calpastatin,¹²⁾ cloning calpain¹³⁾ and calpastatin cDNA,¹⁴³⁾ determining the 3D structure,¹⁰⁹⁾ and many more. Their highly comprehensive and educational reviews^{17),48),49),274),275)} have been cited many times. These heroes will always live in the memory of many calpain researchers. They made great strides in the study of calpains, but much work remains to elucidate calpain's physiological roles. The divergence of its physiological functions and the lack of a good activation marker are major obstacles. Recent progress in genetics and structural biology has led to new and invaluable information about calpains. Considering calpain's divergent physiological functions, breakthroughs in the field of calpain studies will affect an enormously wide range of life science fields.

Acknowledgments

We dedicate this paper to the memory of Prof. Koichi Suzuki. We thank Dr. Tamio Yamakawa, M.J.A., for giving us the opportunity to write this review, all the laboratory members for their invaluable support, and Drs. Leslie Miglietta and Grace

Gray for their excellent English editing. We also thank Drs. Peter L. Davies, Robert L. Campbell, and all calpain researchers who gave opinions at "Calpain Nomenclature Proposal and Comments" web site for valuable discussion about calpain structures and nomenclature. This work was supported in part by MEXT.KAKENHI 18076007 (H.S.), 20780106 (S.H.), 22770139 (Y.O.), JSPS.KAKENHI 20370055, and a Takeda Science Foundation research grant (H.S.).

References

- 1) Guroff, G. (1964) A neutral calcium-activated proteinase from the soluble fraction of rat brain. *J. Biol. Chem.* **239**, 149–155.
- 2) Lazarovici, P. (2000) Obituary: Dr. Gordon Guroff (1933–1999). *J. Mol. Neurosci.* **14**, 1–2.
- 3) Huston, R.B. and Krebs, E.G. (1968) Activation of skeletal muscle phosphorylase kinase by Ca^{2+} . II. Identification of the kinase activating factor as a proteolytic enzyme. *Biochemistry* **7**, 2116–2122.
- 4) Meyer, W.L., Fischer, E.H. and Krebs, E.G. (1964) Activation of skeletal muscle phosphorylase kinase by Ca^{2+} . *Biochemistry* **3**, 1033–1039.
- 5) Busch, W.A., Stromer, M.H., Goll, D.E. and Suzuki, A. (1972) Ca^{2+} -specific removal of Z lines from rabbit skeletal muscle. *J. Cell Biol.* **52**, 367–381.
- 6) Takai, Y., Yamamoto, M., Inoue, M., Kishimoto, A. and Nishizuka, Y. (1977) A proenzyme of cyclic nucleotide-independent protein kinase and its activation by calcium-dependent neutral protease from rat liver. *Biochem. Biophys. Res. Commun.* **77**, 542–550.
- 7) Ishiura, S., Murofushi, H., Suzuki, K. and Imahori, K. (1978) Studies of a calcium-activated neutral protease from chicken skeletal muscle. I. Purification and characterization. *J. Biochem.* **84**, 225–230.
- 8) Suzuki, K., Tsuji, S., Ishiura, S., Kimura, Y., Kubota, S. and Imahori, K. (1981) Autolysis of calcium-activated neutral protease of chicken skeletal muscle. *J. Biochem.* **90**, 1787–1793.
- 9) Suzuki, K. (1983) Reaction of calcium-activated neutral protease (CANP) with an epoxysuccinyl derivative (E64c) and iodoacetic acid. *J. Biochem.* **93**, 1305–1312.
- 10) Suzuki, K. and Ishiura, S. (1983) Effect of metal ions on the structure and activity of calcium-activated neutral protease (CANP). *J. Biochem.* **93**, 1463–1471.
- 11) Nishiura, I., Tanaka, K., Yamato, S. and Murachi, T. (1978) The occurrence of an inhibitor of Ca^{2+} -dependent neutral protease in rat liver. *J. Biochem.* **84**, 1657–1659.
- 12) Murachi, T., Tanaka, K., Hatanaka, M. and Murakami, T. (1981) Intracellular Ca^{2+} -dependent protease (calpain) and its high-molecular-weight endogenous inhibitor (calpastatin). *Adv. Enzyme Regul.* **19**, 407–424.
- 13) Ohno, S., Emori, Y., Imajoh, S., Kawasaki, H., Kisaragi, M. and Suzuki, K. (1984) Evolutionary origin of a calcium-dependent protease by fusion of genes for a thiol protease and a calcium-binding protein? *Nature* **312**, 566–570.
- 14) Suzuki, K. (1991) Nomenclature of calcium dependent proteinase. *Biomed. Biochim. Acta* **50**, 483–484.
- 15) Ohno, S., Minoshima, S., Kudoh, J., Fukuyama, R., Shimizu, Y., Ohmi-Imajoh, S., Shimizu, N. and Suzuki, K. (1990) Four genes for the calpain family locate on four distinct human chromosomes. *Cytogenet. Cell Genet.* **53**, 225–229.
- 16) Croall, D.E. and DeMartino, G.N. (1991) Calcium-activated neutral protease (calpain) system: structure, function, and regulation. *Physiol. Rev.* **71**, 813–847.
- 17) Goll, D.E., Thompson, V.F., Li, H., Wei, W. and Cong, J. (2003) The calpain system. *Physiol. Rev.* **83**, 731–801.
- 18) Zatz, M. and Starling, A. (2005) Mechanisms of disease: Calpains and disease. *N. Engl. J. Med.* **352**, 2413–2423.
- 19) Yamashima, T. (2004) Ca^{2+} -dependent proteases in ischemic neuronal death: a conserved 'calpain-cathepsin cascade' from nematodes to primates. *Cell Calcium* **36**, 285–293.
- 20) Wendt, A., Thompson, V.F. and Goll, D.E. (2004) Interaction of calpastatin with calpain: a review. *Biol. Chem.* **385**, 465–472.
- 21) Wells, A., Huttenlocher, A. and Lauffenburger, D.A. (2005) Calpain proteases in cell adhesion and motility. *Int. Rev. Cytol.* **245**, 1–16.
- 22) Turner, M.D., Cassell, P.G. and Hitman, G.A. (2005) Calpain-10: from genome search to function. *Diabetes Metab. Res. Rev.* **21**, 505–514.
- 23) Tian, W., Dewitt, S., Laffafian, I. and Hallett, M.B. (2004) Ca^{2+} , calpain and 3-phosphorylated phosphatidyl inositides; decision-making signals in neutrophils as potential targets for therapeutics. *J. Pharm. Pharmacol.* **56**, 565–571.
- 24) Sorimachi, H. and Kawabata, Y. (2003) Calpain and pathology in view of structure–function relationships. *Nippon Yakurigaku Zasshi* **122**, 21–29 (in Japanese).
- 25) Ray, S.K., Hogan, E.L. and Banik, N.L. (2003) Calpain in the pathophysiology of spinal cord injury: neuroprotection with calpain inhibitors. *Brain Res. Rev.* **42**, 169–185.
- 26) Ridderstrale, M., Parikh, H. and Groop, L. (2005) Calpain 10 and type 2 diabetes: are we getting closer to an explanation? *Curr. Opin. Clin. Nutr. Metab. Care* **8**, 361–366.
- 27) Ray, S.K. and Banik, N.L. (2003) Calpain and its involvement in the pathophysiology of CNS injuries and diseases: therapeutic potential of calpain inhibitors for prevention of neurodegeneration. *Curr. Drug Targets CNS Neurol. Disord.* **2**, 173–189.
- 28) Nixon, R.A. (2003) The calpains in aging and aging-related diseases. *Ageing Res. Rev.* **2**, 407–418.
- 29) Mehendale, H.M. and Limaye, P.B. (2005) Calpain:

- a death protein that mediates progression of liver injury. *Trends Pharmacol. Sci.* **26**, 232–236.
- 30) Liu, X., Van Vleet, T. and Schnellmann, R.G. (2004) The role of calpain in oncotic cell death. *Annu. Rev. Pharmacol. Toxicol.* **44**, 349–370.
 - 31) Laval, S.H. and Bushby, K.M.D. (2004) Limb-girdle muscular dystrophies—From genetics to molecular pathology. *Neuropathol. Appl. Neurobiol.* **30**, 91–105.
 - 32) Horikawa, Y. (2005) Type 2 diabetes and genetic variations of calpain 10 gene. *Nippon Rinsho* **63**, 155–159 (in Japanese).
 - 33) Harwood, S.M., Yaqoob, M.M. and Allen, D.A. (2005) Caspase and calpain function in cell death: bridging the gap between apoptosis and necrosis. *Ann. Clin. Biochem.* **42**, 415–431.
 - 34) Friedrich, P., Tompa, P. and Farkas, A. (2004) The calpain-system of *Drosophila melanogaster*: coming of age. *Bioessays* **26**, 1088–1096.
 - 35) Franco, S.J. and Huttenlocher, A. (2005) Regulating cell migration: calpains make the cut. *J. Cell Sci.* **118**, 3829–3838.
 - 36) Costelli, P., Reffo, P., Penna, F., Autelli, R., Bonelli, G. and Baccino, F.M. (2005) Ca²⁺-dependent proteolysis in muscle wasting. *Int. J. Biochem. Cell Biol.* **37**, 2134–2146.
 - 37) Carragher, N.O. (2006) Calpain inhibition: a therapeutic strategy targeting multiple disease states. *Curr. Pharm. Des.* **12**, 615–638.
 - 38) Branca, D. (2004) Calpain-related diseases. *Biochem. Biophys. Res. Commun.* **322**, 1098–1104.
 - 39) Biswas, S., Harris, F., Dennison, S., Singh, J.P. and Phoenix, D. (2005) Calpains: enzymes of vision? *Med. Sci. Monit.* **11**, RA301–RA310.
 - 40) Biswas, S., Harris, F., Dennison, S., Singh, J. and Phoenix, D.A. (2004) Calpains: targets of cataract prevention? *Trends Mol. Med.* **10**, 78–84.
 - 41) Bartoli, M. and Richard, I. (2005) Calpains in muscle wasting. *Int. J. Biochem. Cell Biol.* **37**, 2115–2133.
 - 42) Baud, L., Fouqueray, B., Bellocq, A. and Peltier, J. (2003) Calpains participate in inflammatory reaction development. *Med. Sci. (Paris)* **19**, 71–76 (in French).
 - 43) Kuchay, S.M. and Chishti, A.H. (2007) Calpain-mediated regulation of platelet signaling pathways. *Curr. Opin. Hematol.* **14**, 249–254.
 - 44) Liu, J., Liu, M.C. and Wang, K.K. (2008) Calpain in the CNS: from synaptic function to neurotoxicity. *Sci. Signal.* **1**, re1.
 - 45) Bertipaglia, I. and Carafoli, E. (2007) Calpains and human disease. *Subcell. Biochem.* **45**, 29–53.
 - 46) Croall, D.E. and Ersfeld, K. (2007) The calpains: modular designs and functional diversity. *Genome Biol.* **8**, 218.
 - 47) Suzuki, K., Hata, S., Kawabata, Y. and Sorimachi, H. (2004) Structure, activation, and biology of calpain. *Diabetes* **53**, S12–S18.
 - 48) Sorimachi, H., Ishiura, S. and Suzuki, K. (1997) Structure and physiological function of calpains. *Biochem. J.* **328**, 721–732.
 - 49) Murachi, T. (1983) Calpain and calpastatin. *Trends Biochem. Sci.* **8**, 167–169.
 - 50) Tanaka, K. (2009) The proteasome: overview of structure and functions. *Proc. Jpn. Acad., Ser. B, Phys. Biol. Sci.* **85**, 12–36.
 - 51) Mizushima, N. and Levine, B. (2010) Autophagy in mammalian development and differentiation. *Nat. Cell Biol.* **12**, 823–830.
 - 52) Tait, S.W. and Green, D.R. (2010) Mitochondria and cell death: outer membrane permeabilization and beyond. *Nat. Rev. Mol. Cell Biol.* **11**, 621–632.
 - 53) Ono, Y., Kakinuma, K., Torii, F., Irie, A., Nakagawa, K., Labeit, S., Abe, K., Suzuki, K. and Sorimachi, H. (2004) Possible regulation of the conventional calpain system by skeletal muscle-specific calpain, p94/calpain 3. *J. Biol. Chem.* **279**, 2761–2771.
 - 54) Ono, Y., Ojima, K., Torii, F., Takaya, E., Doi, N., Nakagawa, K., Hata, S., Abe, K. and Sorimachi, H. (2010) Skeletal muscle-specific calpain is an intracellular Na⁺-dependent protease. *J. Biol. Chem.* **285**, 22986–22998.
 - 55) Dutt, P., Croall, D.E., Arthur, S.C., De Veyra, T., Williams, K., Elce, J.S. and Greer, P.A. (2006) m-Calpain is required for preimplantation embryonic development in mice. *BMC Dev. Biol.* **6**, 3.
 - 56) Richard, I., Broux, O., Allamand, V., Fougerousse, F., Chiannilkulchai, N., Bourg, N., Brenguier, L., Devaud, C., Pasturaud, P., Roudaut, C., Hillaire, D., Passos-Bueno, M.-R., Zats, M., Tischfield, J.A., Fardeau, M., Jackson, C.E., Cohen, D. and Beckmann, J.S. (1995) Mutations in the proteolytic enzyme calpain 3 cause limb-girdle muscular dystrophy type 2A. *Cell* **81**, 27–40.
 - 57) Hata, S., Abe, M., Suzuki, H., Kitamura, F., Toyama-Sorimachi, N., Abe, K., Sakimura, K. and Sorimachi, H. (2010) Calpain 8/nCL-2 and Calpain 9/nCL-4 constitute an active protease complex, G-calpain, involved in gastric mucosal defense. *PLoS Genet.* **6**, e1001040.
 - 58) Yamada, M., Yoshida, Y., Mori, D., Takitoh, T., Kengaku, M., Umeshima, H., Takao, K., Miyakawa, T., Sato, M., Sorimachi, H., Wynshaw-Boris, A. and Hirotsune, S. (2009) Inhibition of calpain increases LIS1 expression and partially rescues *in vivo* phenotypes in a mouse model of lissencephaly. *Nat. Med.* **15**, 1202–1207.
 - 59) Kimura, Y., Koga, H., Araki, N., Mugita, N., Fujita, N., Takeshima, H., Nishi, T., Yamashima, T., Saido, T.C., Yamasaki, T., Moritake, K., Saya, H. and Nakao, M. (1998) The involvement of calpain-dependent proteolysis of the tumor suppressor NF2 (merlin) in schwannomas and meningiomas. *Nat. Med.* **4**, 915–922.
 - 60) Delaney, S.J., Hayward, D.C., Barleben, F., Fischbach, K.F. and Miklos, G.L. (1991) Molecular cloning and analysis of small optic lobes, a structural brain gene of *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* **88**, 7214–7218.
 - 61) Barnes, T.M. and Hodgkin, J. (1996) The *tra-3* sex determination gene of *Caenorhabditis elegans*

- encodes a member of the calpain regulatory protease family. *EMBO J.* **15**, 4477–4484.
- 62) Lid, S.E., Gruis, D., Jung, R., Lorentzen, J.A., Ananiev, E., Chamberlin, M., Niu, X., Meeley, R., Nichols, S. and Olsen, O.A. (2002) The defective kernel 1 (*dek1*) gene required for aleurone cell development in the endosperm of maize grains encodes a membrane protein of the calpain gene superfamily. *Proc. Natl. Acad. Sci. USA* **99**, 5460–5465.
- 63) Denison, S.H., Orejas, M. and Arst, H.N. Jr. (1995) Signaling of ambient pH in *Aspergillus* involves a cysteine protease. *J. Biol. Chem.* **270**, 28519–28522.
- 64) Futai, E., Maeda, T., Sorimachi, H., Kitamoto, K., Ishiura, S. and Suzuki, K. (1999) The protease activity of a calpain-like cysteine protease in *Saccharomyces cerevisiae* is required for alkaline adaptation and sporulation. *Mol. Gen. Genet.* **260**, 559–568.
- 65) Hayashi, M., Fukuzawa, T., Sorimachi, H. and Maeda, T. (2005) Constitutive activation of the pH-responsive Rim101 pathway in yeast mutants defective in late steps of the MVB/ESCRT pathway. *Mol. Cell. Biol.* **25**, 9478–9490.
- 66) Suzuki, K. (1991) Historical background, current status, and future prospect of research on calcium dependent protease. *Cell Technol.* **10**, 545–551 (in Japanese).
- 67) Suzuki, K. (1991) Determination of structure of calpain and East–West problem. *Cell Technol.* **10**, 719–727 (in Japanese).
- 68) Suzuki, K. (1991) In search of physiological function of calpain. *Cell Technol.* **10**, 878–885 (in Japanese).
- 69) Drummond, G.I. and Duncan, L. (1968) On the mechanism of activation of phosphorylase b kinase by calcium. *J. Biol. Chem.* **243**, 5532–5538.
- 70) Fischer, E.H. and Krebs, E.G. (1955) Conversion of phosphorylase b to phosphorylase a in muscle extracts. *J. Biol. Chem.* **216**, 121–132.
- 71) Krebs, E.G. and Fischer, E.H. (1955) Phosphorylase activity of skeletal muscle extracts. *J. Biol. Chem.* **216**, 113–120.
- 72) Koohmaraie, M. (1992) The role of Ca²⁺-dependent proteases (calpains) in post mortem proteolysis and meat tenderness. *Biochimie* **74**, 239–245.
- 73) Kishimoto, A., Kajikawa, N., Tabuchi, H., Shiota, M. and Nishizuka, Y. (1981) Calcium-dependent neural proteases, widespread occurrence of a species of protease active at lower concentrations of calcium. *J. Biochem.* **90**, 889–892.
- 74) Ogawa, Y., Takai, Y., Kawahara, Y., Kimura, S. and Nishizuka, Y. (1981) A new possible regulatory system for protein phosphorylation in human peripheral lymphocytes. I. Characterization of a calcium-activated, phospholipid-dependent protein kinase. *J. Immunol.* **127**, 1369–1374.
- 75) Takai, Y., Kaibuchi, K., Sano, K. and Nishizuka, Y. (1982) Counteraction of calcium-activated, phospholipid-dependent protein kinase activation by adenosine 3',5'-monophosphate and guanosine 3',5'-monophosphate in platelets. *J. Biochem.* **91**, 403–406.
- 76) Kishimoto, A., Kajikawa, N., Shiota, M. and Nishizuka, Y. (1983) Proteolytic activation of calcium-activated, phospholipid-dependent protein kinase by calcium-dependent neutral protease. *J. Biol. Chem.* **258**, 1156–1164.
- 77) Ishiura, S., Sugita, H., Suzuki, K. and Imahori, K. (1979) Studies of a calcium-activated neutral protease from chicken skeletal muscle. II. Substrate specificity. *J. Biochem.* **86**, 579–581.
- 78) Suzuki, K., Ishiura, S., Tsuji, S., Katamoto, T., Sugita, H. and Imahori, K. (1979) Calcium activated neutral protease from human skeletal muscle. *FEBS Lett.* **104**, 355–358.
- 79) Sugita, H., Ishiura, S., Suzuki, K. and Imahori, K. (1980) Ca-activated neutral protease and its inhibitors: in vitro effect on intact myofibrils. *Muscle Nerve* **3**, 335–339.
- 80) Sugita, H., Ishiura, S., Suzuki, K. and Imahori, K. (1980) Inhibition of epoxide derivatives on chicken calcium-activated neutral protease (CANP) *in vitro* and *in vivo*. *J. Biochem.* **87**, 339–341.
- 81) Suzuki, K., Tsuji, S., Kubota, S., Kimura, Y. and Imahori, K. (1981) Limited autolysis of Ca²⁺-activated neutral protease (CANP) changes its sensitivity to Ca²⁺ ions. *J. Biochem.* **90**, 275–278.
- 82) Suzuki, K. (1982) Activation mechanism of calcium-activated neutral protease. *Seikagaku* **54**, 1264–1270 (in Japanese).
- 83) Suzuki, K. and Tsuji, S. (1982) Synergistic activation of calcium-activated neutral protease by Mn²⁺ and Ca²⁺. *FEBS Lett.* **140**, 16–18.
- 84) Dayton, W.R., Goll, D.E., Zeece, M.G., Robson, R.M. and Reville, W.J. (1976) A Ca²⁺-activated protease possibly involved in myofibrillar protein turnover. Purification from porcine muscle. *Biochemistry* **15**, 2150–2158.
- 85) Ohno, S., Akita, Y., Hata, A., Osada, S., Kubo, K., Konno, Y., Akimoto, K., Mizuno, K., Saïdo, T., Kuroki, T. and Suzuki, K. (1991) Structural and functional diversities of a family of signal transducing protein kinases, protein kinase C family; two distinct classes of PKC, conventional cPKC and novel nPKC. *Adv. Enzyme Regul.* **31**, 287–303.
- 86) Sorimachi, H., Tsukahara, T., Okada-Ban, M., Sugita, H., Ishiura, S. and Suzuki, K. (1995) Identification of a third ubiquitous calpain species—chicken muscle expresses four distinct calpains. *Biochim. Biophys. Acta* **1261**, 381–393.
- 87) Macqueen, D.J., Delbridge, M.L., Manthri, S. and Johnston, I.A. (2010) A newly classified vertebrate calpain protease, directly ancestral to *CAPN1* and *2*, episodically evolved a restricted physiological function in placental mammals. *Mol. Biol. Evol.* **27**, 1886–1902.
- 88) Berti, P.J. and Storer, A.C. (1995) Alignment/phylogeny of the papain superfamily of cysteine proteases. *J. Mol. Biol.* **246**, 273–283.

- 89) Marchler-Bauer, A., Anderson, J.B., Chitsaz, F., Derbyshire, M.K., DeWeese-Scott, C., Fong, J.H., Geer, L.Y., Geer, R.C., Gonzales, N.R., Gwadz, M., He, S., Hurwitz, D.I., Jackson, J.D., Ke, Z., Lanczycki, C.J., Liebert, C.A., Liu, C., Lu, F., Lu, S., Marchler, G.H., Mullokandov, M., Song, J.S., Tasneem, A., Thanki, N., Yamashita, R.A., Zhang, D., Zhang, N. and Bryant, S.H. (2009) CDD: specific functional annotation with the Conserved Domain Database. *Nucleic Acids Res.* **37**, D205–D210.
- 90) Pontremoli, S. and Melloni, E. (1988) The role of calpain and protein kinase C in activation of human neutrophils. *Prog. Clin. Biol. Res.* **282**, 195–208.
- 91) Noguchi, M., Sarin, A., Aman, M.J., Nakajima, H., Shores, E.W., Henkart, P.A. and Leonard, W.J. (1997) Functional cleavage of the common cytokine receptor γ chain (γc) by calpain. *Proc. Natl. Acad. Sci. USA* **94**, 11534–11539.
- 92) Leloup, L., Shao, H., Bae, Y.H., Deasy, B., Stolz, D., Roy, P. and Wells, A. (2010) m-Calpain activation is regulated by its membrane localization and by its binding to phosphatidylinositol 4,5-bisphosphate. *J. Biol. Chem.* **285**, 33549–33566.
- 93) Mellgren, R.L., Zhang, W., Miyake, K. and McNeil, P.L. (2007) Calpain is required for the rapid, calcium-dependent repair of wounded plasma membrane. *J. Biol. Chem.* **282**, 2567–2575.
- 94) Mellgren, R.L. and Huang, X. (2007) Fetuin A stabilizes m-calpain and facilitates plasma membrane repair. *J. Biol. Chem.* **282**, 35868–35877.
- 95) Danial, N.N. and Korsmeyer, S.J. (2004) Cell death: critical control points. *Cell* **116**, 205–219.
- 96) Nakagawa, T. and Yuan, J. (2000) Cross-talk between two cysteine protease families. Activation of caspase-12 by calpain in apoptosis. *J. Cell Biol.* **150**, 887–894.
- 97) Hathaway, D.R., Werth, D.K. and Haerberle, J.R. (1982) Limited autolysis reduces the Ca^{2+} requirement of a smooth muscle Ca^{2+} -activated protease. *J. Biol. Chem.* **257**, 9072–9077.
- 98) Inomata, M., Kasai, Y., Nakamura, M. and Kawashima, S. (1988) Activation mechanism of calcium-activated neutral protease. Evidence for the existence of intramolecular and intermolecular autolyses. *J. Biol. Chem.* **263**, 19783–19787.
- 99) Cong, J., Goll, D.E., Peterson, A.M. and Kapprell, H.P. (1989) The role of autolysis in activity of the Ca^{2+} -dependent proteinases (μ -calpain and m-calpain). *J. Biol. Chem.* **264**, 10096–10103.
- 100) Nakagawa, K., Masumoto, H., Sorimachi, H. and Suzuki, K. (2001) Dissociation of m-calpain subunits occurs after autolysis of the N-terminus of the catalytic subunit, and is not required for activation. *J. Biochem.* **130**, 605–611.
- 101) Pal, G.P., Elce, J.S. and Jia, Z. (2001) Dissociation and aggregation of calpain in the presence of calcium. *J. Biol. Chem.* **276**, 47233–47238.
- 102) Suo, S., Koike, H., Sorimachi, H., Ishiura, S. and Suzuki, K. (1999) Association and dissociation of the calcium-binding domains of calpain by Ca^{2+} . *Biochem. Biophys. Res. Commun.* **257**, 63–66.
- 103) Li, H., Thompson, V.F. and Goll, D.E. (2004) Effects of autolysis on properties of μ - and m-calpain. *Biochim. Biophys. Acta* **1691**, 91–103.
- 104) Zhang, W. and Mellgren, R.L. (1996) Calpain subunits remain associated during catalysis. *Biochem. Biophys. Res. Commun.* **227**, 890–896.
- 105) Dainese, E., Minafra, R., Sabatucci, A., Vachette, P., Melloni, E. and Cozzani, I. (2002) Conformational changes of calpain from human erythrocytes in the presence of Ca^{2+} . *J. Biol. Chem.* **277**, 40296–40301.
- 106) Elce, J.S., Davies, P.L., Hegadorn, C., Maurice, D.H. and Arthur, J.S. (1997) The effects of truncations of the small subunit on m-calpain activity and heterodimer formation. *Biochem. J.* **326**, 31–38.
- 107) Hata, S., Sorimachi, H., Nakagawa, K., Maeda, T., Abe, K. and Suzuki, K. (2001) Domain II of m-calpain is a Ca^{2+} -dependent cysteine protease. *FEBS Lett.* **501**, 111–114.
- 108) Hosfield, C.M., Elce, J.S., Davies, P.L. and Jia, Z. (1999) Crystal structure of calpain reveals the structural basis for Ca^{2+} -dependent protease activity and a novel mode of enzyme activation. *EMBO J.* **18**, 6880–6889.
- 109) Strobl, S., Fernandez-Catalan, C., Braun, M., Huber, R., Masumoto, H., Nakagawa, K., Irie, A., Sorimachi, H., Bourenkow, G., Bartunik, H., Suzuki, K. and Bode, W. (2000) The crystal structure of calcium-free human m-calpain suggests an electrostatic switch mechanism for activation by calcium. *Proc. Natl. Acad. Sci. USA* **97**, 588–592.
- 110) Hanna, R.A., Campbell, R.L. and Davies, P.L. (2008) Calcium-bound structure of calpain and its mechanism of inhibition by calpastatin. *Nature* **456**, 409–412.
- 111) Moldoveanu, T., Gehring, K. and Green, D.R. (2008) Concerted multi-pronged attack by calpastatin to occlude the catalytic cleft of heterodimeric calpains. *Nature* **456**, 404–408.
- 112) Moldoveanu, T., Hosfield, C.M., Lim, D., Elce, J.S., Jia, Z. and Davies, P.L. (2002) A Ca^{2+} switch aligns the active site of calpain. *Cell* **108**, 649–660.
- 113) Moldoveanu, T., Hosfield, C.M., Lim, D., Jia, Z. and Davies, P.L. (2003) Calpain silencing by a reversible intrinsic mechanism. *Nat. Struct. Biol.* **10**, 371–378.
- 114) Tompa, P., Emori, Y., Sorimachi, H., Suzuki, K. and Friedrich, P. (2001) Domain III of calpain is a Ca^{2+} -regulated phospholipid-binding domain. *Biochem. Biophys. Res. Commun.* **280**, 1333–1339.
- 115) Minami, Y., Emori, Y., Imajoh-Ohmi, S., Kawasaki, H. and Suzuki, K. (1988) Carboxyl-terminal truncation and site-directed mutagenesis of the EF hand structure-domain of the small subunit of rabbit calcium-dependent protease. *J. Biochem.* **104**, 927–933.
- 116) Minami, Y., Emori, Y., Kawasaki, H. and Suzuki,

- K. (1987) E-F hand structure-domain of calcium-activated neutral protease (CANP) can bind Ca^{2+} ions. *J. Biochem.* **101**, 889–895.
- 117) Lin, G.D., Chattopadhyay, D., Maki, M., Wang, K.K., Carson, M., Jin, L., Yuen, P.W., Takano, E., Hatanaka, M., DeLucas, L.J. and Narayana, S.V. (1997) Crystal structure of calcium bound domain VI of calpain at 1.9 Å resolution and its role in enzyme assembly, regulation, and inhibitor binding. *Nat. Struct. Biol.* **4**, 539–547.
- 118) Maki, M., Narayana, S.V. and Hitomi, K. (1997) A growing family of the Ca^{2+} -binding proteins with five EF-hand motifs. *Biochem. J.* **328**, 718–720.
- 119) Blanchard, H., Grochulski, P., Li, Y., Arthur, J.S., Davies, P.L., Elce, J.S. and Cygler, M. (1997) Structure of a calpain Ca^{2+} -binding domain reveals a novel EF-hand and Ca^{2+} -induced conformational changes. *Nat. Struct. Biol.* **4**, 532–538.
- 120) Imajoh, S., Kawasaki, H. and Suzuki, K. (1987) The COOH-terminal E-F hand structure of calcium-activated neutral protease (CANP) is important for the association of subunits and resulting proteolytic activity. *J. Biochem.* **101**, 447–452.
- 121) Schad, E., Farkas, A., Jekely, G., Tompa, P. and Friedrich, P. (2002) A novel human small subunit of calpains. *Biochem. J.* **362**, 383–388.
- 122) Tsuji, S., Ishiura, S., Takahashi-Nakamura, M., Katamoto, T., Suzuki, K. and Imahori, K. (1981) Studies on a Ca^{2+} -activated neutral proteinase of rabbit skeletal muscle. II. Characterization of sulfhydryl groups and a role of Ca^{2+} ions in this enzyme. *J. Biochem.* **90**, 1405–1411.
- 123) Tompa, P., Buzder-Lantos, P., Tantos, A., Farkas, A., Szilagy, A., Banoczi, Z., Hudecz, F. and Friedrich, P. (2004) On the sequential determinants of calpain cleavage. *J. Biol. Chem.* **279**, 20775–20785.
- 124) duVerle, D., Takigawa, I., Ono, Y., Sorimachi, H. and Mamitsuka, H. (2010) CaMPDB: a resource for calpain and modulatory proteolysis. *Genome Inform.* **22**, 202–213.
- 125) Davis, T.L., Walker, J.R., Finerty, P.J. Jr., Mackenzie, F., Newman, E.M. and Dhe-Paganon, S. (2007) The crystal structures of human calpains 1 and 9 imply diverse mechanisms of action and auto-inhibition. *J. Mol. Biol.* **366**, 216–229.
- 126) Yuan, P., Leonetti, M.D., Pico, A.R., Hsiung, Y. and MacKinnon, R. (2010) Structure of the human BK channel Ca^{2+} -activation apparatus at 3.0 Å resolution. *Science* **329**, 182–186.
- 127) Moldoveanu, T., Campbell, R.L., Cuerrier, D. and Davies, P.L. (2004) Crystal structures of calpain-E64 and -leupeptin inhibitor complexes reveal mobile loops gating the active site. *J. Mol. Biol.* **343**, 1313–1326.
- 128) Arthur, J.S., Elce, J.S., Hegadorn, C., Williams, K. and Greer, P.A. (2000) Disruption of the murine calpain small subunit gene, *Capn4*: calpain is essential for embryonic development but not for cell growth and division. *Mol. Cell. Biol.* **20**, 4474–4481.
- 129) Zimmerman, U.J., Boring, L., Pak, J.H., Mukerjee, N. and Wang, K.K. (2000) The calpain small subunit gene is essential: its inactivation results in embryonic lethality. *IUBMB Life* **50**, 63–68.
- 130) Yoshizawa, T., Sorimachi, H., Tomioka, S., Ishiura, S. and Suzuki, K. (1995) A catalytic subunit of calpain possesses full proteolytic activity. *FEBS Lett.* **358**, 101–103.
- 131) Azam, M., Andrabi, S.S., Sahr, K.E., Kamath, L., Kuliopulos, A. and Chishti, A.H. (2001) Disruption of the mouse μ -calpain gene reveals an essential role in platelet function. *Mol. Cell. Biol.* **21**, 2213–2220.
- 132) Dourdin, N., Bhatt, A.K., Dutt, P., Greer, P.A., Arthur, J.S., Elce, J.S. and Huttenlocher, A. (2001) Reduced cell migration and disruption of the actin cytoskeleton in calpain-deficient embryonic fibroblasts. *J. Biol. Chem.* **276**, 48382–48388.
- 133) Mellgren, R.L., Zhang, W., Miyake, K. and McNeil, P.L. (2006) Calpain is required for the rapid, calcium-dependent repair of wounded plasma membrane. *J. Biol. Chem.* **282**, 2567–2575.
- 134) Demarchi, F., Bertoli, C., Copetti, T., Tanida, I., Brancolini, C., Eskelinen, E.L. and Schneider, C. (2006) Calpain is required for macroautophagy in mammalian cells. *J. Cell Biol.* **175**, 595–605.
- 135) Farkas, A., Tompa, P. and Friedrich, P. (2003) Revisiting ubiquity and tissue specificity of human calpains. *Biol. Chem.* **384**, 945–949.
- 136) Hata, A., Ohno, S., Akita, Y. and Suzuki, K. (1989) Tandemly reiterated negative enhancer-like elements regulate transcription of a human gene for the large subunit of calcium-dependent protease. *J. Biol. Chem.* **264**, 6404–6411.
- 137) Valerio, D., Duyvesteyn, M.G., Dekker, B.M., Weeda, G., Berkvens, T.M., van der Voorn, L., van Ormondt, H. and van der Eb, A.J. (1985) Adenosine deaminase: characterization and expression of a gene with a remarkable promoter. *EMBO J.* **4**, 437–443.
- 138) Melton, D.W., McEwan, C., McKie, A.B. and Reid, A.M. (1986) Expression of the mouse HPRT gene: deletional analysis of the promoter region of an X-chromosome linked housekeeping gene. *Cell* **44**, 319–328.
- 139) Angel, P., Imagawa, M., Chiu, R., Stein, B., Imbra, R.J., Rahmsdorf, H.J., Jonat, C., Herrlich, P. and Karin, M. (1987) Phorbol ester-inducible genes contain a common *cis* element recognized by a TPA-modulated *trans*-acting factor. *Cell* **49**, 729–739.
- 140) Lee, W., Mitchell, P. and Tjian, R. (1987) Purified transcription factor AP-1 interacts with TPA-inducible enhancer elements. *Cell* **49**, 741–752.
- 141) Nakamura, M., Mori, M., Morishita, Y., Mori, S. and Kawashima, S. (1992) Specific increase in calcium-activated neutral protease with low calcium sensitivity (m-calpain) in proerythroblastic K562 cell line cells induced to differentiation by phorbol 12-myristate 13-acetate. *Exp. Cell*

- Res. **200**, 513–522.
- 142) Miyake, S., Emori, Y. and Suzuki, K. (1986) Gene organization of the small subunit of human calcium-activated neutral protease. *Nucleic Acids Res.* **14**, 8805–8817.
- 143) Emori, Y., Kawasaki, H., Imajoh, S., Imahori, K. and Suzuki, K. (1987) Endogenous inhibitor for calcium-dependent cysteine protease contains four internal repeats that could be responsible for its multiple reactive sites. *Proc. Natl. Acad. Sci. USA* **84**, 3590–3594.
- 144) Maki, M., Takano, E., Mori, H., Kannagi, R., Murachi, T. and Hatanaka, M. (1987) Repetitive region of calpastatin is a functional unit of the proteinase inhibitor. *Biochem. Biophys. Res. Commun.* **143**, 300–308.
- 145) Maki, M., Takano, E., Mori, H., Sato, A., Murachi, T. and Hatanaka, M. (1987) All four internally repetitive domains of pig calpastatin possess inhibitory activities against calpains I and II. *FEBS Lett.* **223**, 174–180.
- 146) Hata, S., Doi, N., Kitamura, F. and Sorimachi, H. (2007) Stomach-specific calpain, nCL-2/calpain 8, is active without calpain regulatory subunit and oligomerizes through C2-like domains. *J. Biol. Chem.* **282**, 27847–27856.
- 147) Lee, H.J., Tomioka, S., Kinbara, K., Masumoto, H., Jeong, S.Y., Sorimachi, H., Ishiura, S. and Suzuki, K. (1999) Characterization of a human digestive tract-specific calpain, nCL-4, expressed in the baculovirus system. *Arch. Biochem. Biophys.* **362**, 22–31.
- 148) Takano, J., Tomioka, M., Tsubuki, S., Higuchi, M., Iwata, N., Itohara, S., Maki, M. and Saido, T.C. (2005) Calpain mediates excitotoxic DNA fragmentation via mitochondrial pathways in adult brains: evidence from calpastatin mutant mice. *J. Biol. Chem.* **280**, 16175–16184.
- 149) Higuchi, M., Tomioka, M., Takano, J., Shirotani, K., Iwata, N., Masumoto, H., Maki, M., Itohara, S. and Saido, T.C. (2005) Distinct mechanistic roles of calpain and caspase activation in neurodegeneration as revealed in mice overexpressing their specific inhibitors. *J. Biol. Chem.* **280**, 15229–15237.
- 150) Spencer, M.J. and Mellgren, R.L. (2002) Overexpression of a calpastatin transgene in mdx muscle reduces dystrophic pathology. *Hum. Mol. Genet.* **11**, 2645–2655.
- 151) Tidball, J.G. and Spencer, M.J. (2002) Expression of a calpastatin transgene slows muscle wasting and obviates changes in myosin isoform expression during murine muscle disuse. *J. Physiol.* **545**, 819–828.
- 152) Kent, M.P., Spencer, M.J. and Koohmaraie, M. (2004) Postmortem proteolysis is reduced in transgenic mice overexpressing calpastatin. *J. Anim. Sci.* **82**, 794–801.
- 153) Geesink, G.H., Kuchay, S., Chishti, A.H. and Koohmaraie, M. (2006) Micro-calpain is essential for postmortem proteolysis of muscle proteins. *J. Anim. Sci.* **84**, 2834–2840.
- 154) Dear, T.N. and Boehm, T. (2001) Identification and characterization of two novel calpain large subunit genes. *Gene* **274**, 245–252.
- 155) Andresen, K., Tom, T.D. and Strand, M. (1991) Characterization of cDNA clones encoding a novel calcium-activated neutral proteinase from *Schistosoma mansoni*. *J. Biol. Chem.* **266**, 15085–15090.
- 156) Ersfeld, K., Barraclough, H. and Gull, K. (2005) Evolutionary relationships and protein domain architecture in an expanded calpain superfamily in kinetoplastid parasites. *J. Mol. Evol.* **61**, 742–757.
- 157) Hata, S., Nishi, K., Kawamoto, T., Lee, H.J., Kawahara, H., Maeda, T., Shintani, Y., Sorimachi, H. and Suzuki, K. (2001) Both the conserved and the unique gene structure of stomach-specific calpains reveal processes of calpain gene evolution. *J. Mol. Evol.* **53**, 191–203.
- 158) Futai, E., Kubo, T., Sorimachi, H., Suzuki, K. and Maeda, T. (2001) Molecular cloning of PalBH, a mammalian homologue of the *Aspergillus* atypical calpain PalB. *Biochim. Biophys. Acta* **1517**, 316–319.
- 159) Li, M., Martin, S.J., Bruno, V.M., Mitchell, A.P. and Davis, D.A. (2004) *Candida albicans* Rim13p, a protease required for Rim101p processing at acidic and alkaline pHs. *Eukaryot. Cell* **3**, 741–751.
- 160) Futai, E., Sorimachi, H., Jeong, S.Y., Kitamoto, K., Ishiura, S. and Suzuki, K. (1999) *Aspergillus oryzae* palB^{ory} encodes a calpain-like protease: homology to *Emericella nidulans* PalB and conservation of functional regions. *J. Biosci. Bioeng.* **88**, 438–440.
- 161) Villalobo, E., Moch, C., Fryd-Versavel, G., Fleury-Aubusson, A. and Morin, L. (2003) Cysteine proteases and cell differentiation: excystation of the ciliated protist *Sterkiella histriomuscorum*. *Eukaryot. Cell* **2**, 1234–1245.
- 162) Horikawa, Y., Oda, N., Cox, N.J., Li, X., Orho-Melander, M., Hara, M., Hinokio, Y., Lindner, T.H., Mashima, H., Schwarz, P.E., del Bosque-Plata, L., Horikawa, Y., Oda, Y., Yoshiuchi, I., Colilla, S., Polonsky, K.S., Wei, S., Concannon, P., Iwasaki, N., Schulze, J., Baier, L.J., Bogardus, C., Groop, L., Boerwinkle, E., Hanis, C.L. and Bell, G.I. (2000) Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. *Nat. Genet.* **26**, 163–175.
- 163) Margis, R. and Margis-Pinheiro, M. (2003) Phyto-calpains: orthologous calcium-dependent cysteine proteinases. *Trends Plant Sci.* **8**, 58–62.
- 164) Wang, C., Barry, J.K., Min, Z., Tordsen, G., Rao, A.G. and Olsen, O.A. (2003) The calpain domain of the maize DEK1 protein contains the conserved catalytic triad and functions as a cysteine proteinase. *J. Biol. Chem.* **278**, 34467–34474.
- 165) Lid, S.E., Olsen, L., Nestestog, R., Aukerman, M., Brown, R.C., Lemmon, B., Mucha, M., Opsahl-Sorteberg, H.G. and Olsen, O.A. (2005)

- Mutation in the *Arabidopsis thaliana* *DEK1* calpain gene perturbs endosperm and embryo development while over-expression affects organ development globally. *Planta* **221**, 339–351.
- 166) Ahn, J.W., Kim, M., Lim, J.H., Kim, G.T. and Pai, H.S. (2004) Phytocalpain controls the proliferation and differentiation fates of cells in plant organ development. *Plant J.* **38**, 969–981.
- 167) Bourgeau, G., Lapointe, H., Peloquin, P. and Mayrand, D. (1992) Cloning, expression, and sequencing of a protease gene (*tpr*) from *Porphyr-omonas gingivalis* W83 in *Escherichia coli*. *Infect. Immun.* **60**, 3186–3192.
- 168) Wu, K.H. and Tai, P.C. (2004) Cys32 and His105 are the critical residues for the calcium-dependent cysteine proteolytic activity of CvaB, an ATP-binding cassette transporter. *J. Biol. Chem.* **279**, 901–909.
- 169) Sorimachi, H., Saido, T.C. and Suzuki, K. (1994) New era of calpain research. Discovery of tissue-specific calpains. *FEBS Lett.* **343**, 1–5.
- 170) Badalamente, M.A. and Stracher, A. (2000) Delay of muscle degeneration and necrosis in mdx mice by calpain inhibition. *Muscle Nerve* **23**, 106–111.
- 171) Bartus, R.T., Hayward, N.J., Elliott, P.J., Sawyer, S.D., Baker, K.L., Dean, R.L., Akiyama, A., Straub, J.A., Harbeson, S.L., Li, Z. and Powers, J. (1994) Calpain inhibitor AK295 protects neurons from focal brain ischemia. Effects of postocclusion intra-arterial administration. *Stroke* **25**, 2265–2270.
- 172) Sorimachi, H., Forsberg, N.E., Lee, H.J., Joeng, S.Y., Richard, I., Beckmann, J.S., Ishiura, S. and Suzuki, K. (1996) Highly conserved structure in the promoter region of the gene for muscle-specific calpain, p94. *Biol. Chem.* **377**, 859–864.
- 173) Kawabata, Y., Hata, S., Ono, Y., Ito, Y., Suzuki, K., Abe, K. and Sorimachi, H. (2003) Newly identified exons encoding novel variants of p94/calpain 3 are expressed ubiquitously and overlap the α -glucosidase C gene. *FEBS Lett.* **555**, 623–630.
- 174) De Tullio, R., Stifanese, R., Salamino, F., Pontremoli, S. and Melloni, E. (2003) Characterization of a new p94-like calpain form in human lymphocytes. *Biochem. J.* **375**, 689–696.
- 175) König, N., Raynaud, F., Feane, H., Durand, M., Mestre-Francès, N., Rossel, M., Ouali, A. and Benyamin, Y. (2003) Calpain 3 is expressed in astrocytes of rat and *Microcebus* brain. *J. Chem. Neuroanat.* **25**, 129–136.
- 176) Azuma, M., Fukiage, C., Higashine, M., Nakajima, T., Ma, H. and Shearer, T.R. (2000) Identification and characterization of a retina-specific calpain (Rt88) from rat. *Curr. Eye Res.* **21**, 710–720.
- 177) Ma, H., Fukiage, C., Azuma, M. and Shearer, T.R. (1998) Cloning and expression of mRNA for calpain Lp82 from rat lens: splice variant of p94. *Invest. Ophthalmol. Vis. Sci.* **39**, 454–461.
- 178) Fougerousse, F., Bullen, P., Herasse, M., Lindsay, S., Richard, I., Wilson, D., Suel, L., Durand, M., Robson, S., Abitbol, M., Beckmann, J.S. and Strachan, T. (2000) Human–mouse differences in the embryonic expression patterns of developmental control genes and disease genes. *Hum. Mol. Genet.* **9**, 165–173.
- 179) Sorimachi, H., Ishiura, S. and Suzuki, K. (1993) A novel tissue-specific calpain species expressed predominantly in the stomach comprises two alternative splicing products with and without Ca²⁺-binding domain. *J. Biol. Chem.* **268**, 19476–19482.
- 180) Sorimachi, H., Imajoh-Ohmi, S., Emori, Y., Kawasaki, H., Ohno, S., Minami, Y. and Suzuki, K. (1989) Molecular cloning of a novel mammalian calcium-dependent protease distinct from both m- and μ -types. Specific expression of the mRNA in skeletal muscle. *J. Biol. Chem.* **264**, 20106–20111.
- 181) Sorimachi, H., Toyama-Sorimachi, N., Saido, T.C., Kawasaki, H., Sugita, H., Miyasaka, M., Arahata, K., Ishiura, S. and Suzuki, K. (1993) Muscle-specific calpain, p94, is degraded by autolysis immediately after translation, resulting in disappearance from muscle. *J. Biol. Chem.* **268**, 10593–10605.
- 182) Ono, Y., Hayashi, C., Doi, N., Kitamura, F., Shindo, M., Kudo, K., Tsubata, T., Yanagida, M. and Sorimachi, H. (2007) Comprehensive survey of p94/calpain 3 substrates by comparative proteomics—Possible regulation of protein synthesis by p94. *Biotechnol. J.* **2**, 565–576.
- 183) Ono, Y., Torii, F., Ojima, K., Doi, N., Yoshioka, K., Kawabata, Y., Labeit, D., Labeit, S., Suzuki, K., Abe, K., Maeda, T. and Sorimachi, H. (2006) Suppressed disassembly of autolyzing p94/CAPN3 by N2A connectin/titin in a genetic reporter system. *J. Biol. Chem.* **281**, 18519–18531.
- 184) Sorimachi, H., Kinbara, K., Kimura, S., Takahashi, M., Ishiura, S., Sasagawa, N., Sorimachi, N., Shimada, H., Tagawa, K., Maruyama, K. and Suzuki, K. (1995) Muscle-specific calpain, p94, responsible for limb girdle muscular dystrophy type 2A, associates with connectin through IS2, a p94-specific sequence. *J. Biol. Chem.* **270**, 31158–31162.
- 185) Chiannilkulchai, N., Pasturaud, P., Richard, I., Auffray, C. and Beckmann, J.S. (1995) A primary expression map of the chromosome 15q15 region containing the recessive form of limb-girdle muscular dystrophy (LGMD2A) gene. *Hum. Mol. Genet.* **4**, 717–725.
- 186) Richard, I., Roudaut, C., Marchand, S., Baghdiguian, S., Herasse, M., Stockholm, D., Ono, Y., Suel, L., Bourg, N., Sorimachi, H., Lefranc, G., Fardeau, M., Sebille, A. and Beckmann, J.S. (2000) Loss of calpain 3 proteolytic activity leads to muscular dystrophy and to apoptosis-associated I κ B α /nuclear factor κ B pathway perturbation in mice. *J. Cell Biol.* **151**, 1583–1590.
- 187) Kramerova, I., Kudryashova, E., Tidball, J.G. and Spencer, M.J. (2004) Null mutation of calpain 3

- (p94) in mice causes abnormal sarcomere formation *in vivo* and *in vitro*. *Hum. Mol. Genet.* **13**, 1373–1388.
- 188) Ono, Y., Shimada, H., Sorimachi, H., Richard, I., Saido, T.C., Beckmann, J.S., Ishiura, S. and Suzuki, K. (1998) Functional defects of a muscle-specific calpain, p94, caused by mutations associated with limb-girdle muscular dystrophy type 2A. *J. Biol. Chem.* **273**, 17073–17078.
- 189) Ojima, K., Kawabata, Y., Nakao, H., Nakao, K., Doi, N., Kitamura, F., Ono, Y., Hata, S., Suzuki, H., Kawahara, H., Bogomolovas, J., Witt, C., Ottenheijm, C., Labeit, S., Granzier, H., Toyama-Sorimachi, N., Sorimachi, M., Suzuki, K., Maeda, T., Abe, K., Aiba, A. and Sorimachi, H. (2010) Role of dynamic distribution of muscle-specific calpain in physical-stress adaptation and muscular dystrophy in mice. *J. Clin. Invest.* **120**, 2672–2683.
- 190) Richard, I., Roudaut, C., Saenz, A., Pogue, R., Grimbergen, J.E., Anderson, L.V., Beley, C., Cobo, A.M., de Diego, C., Eymard, B., Gallano, P., Ginjaar, H.B., Lasa, A., Pollitt, C., Topaloglu, H., Urtizberea, J.A., de Visser, M., van der Kooij, A., Bushby, K., Bakker, E., Lopez de Munain, A., Fardeau, M. and Beckmann, J.S. (1999) Calpainopathy—a survey of mutations and polymorphisms. *Am. J. Hum. Genet.* **64**, 1524–1540.
- 191) Sáenz, A., Leturcq, F., Cobo, A.M., Poza, J.J., Ferrer, X., Otaegui, D., Camaño, P., Urtasun, M., Vilchez, J., Gutiérrez-Rivas, E., Empananza, J., Merlini, L., Paisán, C., Goicoechea, M., Blázquez, L., Eymard, B., Lochmuller, H., Walter, M., Bonnemann, C., Figarella-Branger, D., Kaplan, J.C., Urtizberea, J.A., Martí-Massó, J.F. and López de Munain, A. (2005) LGMD2A: genotype-phenotype correlations based on a large mutational survey on the calpain 3 gene. *Brain* **128**, 732–742.
- 192) Hata, S., Koyama, S., Kawahara, H., Doi, N., Maeda, T., Toyama-Sorimachi, N., Abe, K., Suzuki, K. and Sorimachi, H. (2006) Stomach-specific calpain, nCL-2, localizes in mucus cells and proteolyzes the β -subunit of coatamer complex, β -COP. *J. Biol. Chem.* **281**, 11214–11224.
- 193) Lee, H.J., Sorimachi, H., Jeong, S.Y., Ishiura, S. and Suzuki, K. (1998) Molecular cloning and characterization of a novel tissue-specific calpain predominantly expressed in the digestive tract. *Biol. Chem.* **379**, 175–183.
- 194) Cao, Y., Zhao, H. and Grunz, H. (2001) xCL-2 is a novel m-type calpain and disrupts morphogenetic movements during embryogenesis in *Xenopus laevis*. *Dev. Growth Differ.* **43**, 563–571.
- 195) Liu, K., Li, L. and Cohen, S.N. (2000) Antisense RNA-mediated deficiency of the calpain protease, nCL-4, in NIH3T3 cells is associated with neoplastic transformation and tumorigenesis. *J. Biol. Chem.* **275**, 31093–31098.
- 196) Chen, C.J., Nguyen, T. and Shively, J.E. (2010) Role of calpain-9 and PKC- δ in the apoptotic mechanism of lumen formation in CEACAM1 transfected breast epithelial cells. *Exp. Cell Res.* **316**, 638–648.
- 197) Caddick, M.X., Brownlee, A.G. and Arst, H.N. Jr. (1986) Regulation of gene expression by pH of the growth medium in *Aspergillus nidulans*. *Mol. Gen. Genet.* **203**, 346–353.
- 198) Arst, H.N. Jr., Bailey, C.R. and Penfold, H.A. (1980) A possible rôle for acid phosphatase in γ -amino-*n*-butyrate uptake in *Aspergillus nidulans*. *Arch. Microbiol.* **125**, 153–158.
- 199) Arst, H.N. Jr., Bignell, E. and Tilburn, J. (1994) Two new genes involved in signalling ambient pH in *Aspergillus nidulans*. *Mol. Gen. Genet.* **245**, 787–790.
- 200) Negrete-Urtasun, S., Denison, S.H. and Arst, H.N. Jr. (1997) Characterization of the pH signal transduction pathway gene *palA* of *Aspergillus nidulans* and identification of possible homologs. *J. Bacteriol.* **179**, 1832–1835.
- 201) Maccheroni, W. Jr., May, G.S., Martinez-Rossi, N.M. and Rossi, A. (1997) The sequence of *palF*, an environmental pH response gene in *Aspergillus nidulans*. *Gene* **194**, 163–167.
- 202) Denison, S.H., Negrete-Urtasun, S., Mingot, J.M., Tilburn, J., Mayer, W.A., Goel, A., Espeso, E.A., Peñalva, M.A. and Arst, H.N. Jr. (1998) Putative membrane components of signal transduction pathways for ambient pH regulation in *Aspergillus* and meiosis in *Saccharomyces* are homologous. *Mol. Microbiol.* **30**, 259–264.
- 203) Negrete-Urtasun, S., Reiter, W., Diez, E., Denison, S.H., Tilburn, J., Espeso, E.A., Peñalva, M.A. and Arst, H.N. Jr. (1999) Ambient pH signal transduction in *Aspergillus*: completion of gene characterization. *Mol. Microbiol.* **33**, 994–1003.
- 204) Herranz, S., Rodríguez, J.M., Bussink, H.J., Sanchez-Ferrero, J.C., Arst, H.N. Jr., Peñalva, M.A. and Vincent, O. (2005) Arrestin-related proteins mediate pH signaling in fungi. *Proc. Natl. Acad. Sci. USA* **102**, 12141–12146.
- 205) Ridge, K.D., Abdulaev, N.G., Sousa, M. and Palczewski, K. (2003) Phototransduction: crystal clear. *Trends Biochem. Sci.* **28**, 479–487.
- 206) Hervás-Aguilar, A., Rodríguez, J.M., Tilburn, J., Arst, H.N. Jr. and Peñalva, M.A. (2007) Evidence for the direct involvement of the proteasome in the proteolytic processing of the *Aspergillus nidulans* zinc finger transcription factor PacC. *J. Biol. Chem.* **282**, 34735–34747.
- 207) Diez, E., Alvaro, J., Espeso, E.A., Rainbow, L., Suarez, T., Tilburn, J., Arst, H.N. Jr. and Peñalva, M.A. (2002) Activation of the *Aspergillus* PacC zinc finger transcription factor requires two proteolytic steps. *EMBO J.* **21**, 1350–1359.
- 208) Mingot, J.M., Tilburn, J., Diez, E., Bignell, E., Orejas, M., Widdick, D.A., Sarkar, S., Brown, C.V., Caddick, M.X., Espeso, E.A., Arst, H.N. Jr. and Peñalva, M.A. (1999) Specificity determinants of proteolytic processing of *Aspergillus* PacC transcription factor are remote from the processing site, and processing occurs in yeast if pH signalling is bypassed. *Mol. Cell. Biol.* **19**,

- 1390–1400.
- 209) Attaix, D., Ventadour, S., Codran, A., Bechet, D., Taillandier, D. and Combaret, L. (2005) The ubiquitin–proteasome system and skeletal muscle wasting. *Essays Biochem.* **41**, 173–186.
- 210) Murakami, Y., Matsufuji, S., Kameji, T., Hayashi, S., Igarashi, K., Tamura, T., Tanaka, K. and Ichihara, A. (1992) Ornithine decarboxylase is degraded by the 26S proteasome without ubiquitination. *Nature* **360**, 597–599.
- 211) Orian, A., Whiteside, S., Israël, A., Stancovski, I., Schwartz, A.L. and Ciechanover, A. (1995) Ubiquitin-mediated processing of NF- κ B transcriptional activator precursor p105. Reconstitution of a cell-free system and identification of the ubiquitin-carrier protein, *E2*, and a novel ubiquitin-protein ligase, *E3*, involved in conjugation. *J. Biol. Chem.* **270**, 21707–21714.
- 212) Palombella, V.J., Rando, O.J., Goldberg, A.L. and Maniatis, T. (1994) The ubiquitin–proteasome pathway is required for processing the NF- κ B1 precursor protein and the activation of NF- κ B. *Cell* **78**, 773–785.
- 213) Su, S.S. and Mitchell, A.P. (1993) Molecular characterization of the yeast meiotic regulatory gene RIM1. *Nucleic Acids Res.* **21**, 3789–3797.
- 214) Su, S.S. and Mitchell, A.P. (1993) Identification of functionally related genes that stimulate early meiotic gene expression in yeast. *Genetics* **133**, 67–77.
- 215) Li, W. and Mitchell, A.P. (1997) Proteolytic activation of Rim1p, a positive regulator of yeast sporulation and invasive growth. *Genetics* **145**, 63–73.
- 216) Mitchell, B.M., Wu, T.G., Jackson, B.E. and Wilhelmus, K.R. (2007) *Candida albicans* strain-dependent virulence and Rim13p-mediated filamentation in experimental keratomycosis. *Invest. Ophthalmol. Vis. Sci.* **48**, 774–780.
- 217) Davis, D., Wilson, R.B. and Mitchell, A.P. (2000) RIM101-dependent and -independent pathways govern pH responses in *Candida albicans*. *Mol. Cell. Biol.* **20**, 971–978.
- 218) Lambert, M., Blanchin-Roland, S., Le Louedec, F., Lépingle, A. and Gaillardin, C. (1997) Genetic analysis of regulatory mutants affecting synthesis of extracellular proteinases in the yeast *Yarrowia lipolytica*: identification of a RIM101/pacC homolog. *Mol. Cell. Biol.* **17**, 3966–3976.
- 219) González-López, C.I., Ortiz-Castellanos, L. and Ruiz-Herrera, J. (2006) The ambient pH response Rim pathway in *Yarrowia lipolytica*: identification of YRIM9 and characterization of its role in dimorphism. *Curr. Microbiol.* **53**, 8–12.
- 220) Aréchiga-Carvajal, E.T. and Ruiz-Herrera, J. (2005) The RIM101/pacC homologue from the basidiomycete *Ustilago maydis* is functional in multiple pH-sensitive phenomena. *Eukaryot. Cell* **4**, 999–1008.
- 221) Rodríguez-Galán, O., Galindo, A., Hervás-Aguilar, A., Arst, H.N. Jr. and Peñalva, M.A. (2009) Physiological involvement in pH signaling of Vps24-mediated recruitment of *Aspergillus* PalB cysteine protease to ESCRT-III. *J. Biol. Chem.* **284**, 4404–4412.
- 222) Tilburn, J., Sánchez-Ferrero, J.C., Reoyo, E., Arst, H.N. Jr. and Peñalva, M.A. (2005) Mutational analysis of the pH signal transduction component PalC of *Aspergillus nidulans* supports distant similarity to BRO1 domain family members. *Genetics* **171**, 393–401.
- 223) Xu, W., Smith, F.J. Jr., Subaran, R. and Mitchell, A.P. (2004) Multivesicular body-ESCRT components function in pH response regulation in *Saccharomyces cerevisiae* and *Candida albicans*. *Mol. Biol. Cell* **15**, 5528–5537.
- 224) Cornet, M., Bidard, F., Schwarz, P., Da Costa, G., Blanchin-Roland, S., Dromer, F. and Gaillardin, C. (2005) Deletions of endocytic components VPS28 and VPS32 affect growth at alkaline pH and virulence through both RIM101-dependent and RIM101-independent pathways in *Candida albicans*. *Infect. Immun.* **73**, 7977–7987.
- 225) Barwell, K.J., Boysen, J.H., Xu, W. and Mitchell, A.P. (2005) Relationship of DFG16 to the Rim101p pH response pathway in *Saccharomyces cerevisiae* and *Candida albicans*. *Eukaryot. Cell* **4**, 890–899.
- 226) Blanchin-Roland, S., Da Costa, G. and Gaillardin, C. (2005) ESCRT-I components of the endocytic machinery are required for Rim101-dependent ambient pH regulation in the yeast *Yarrowia lipolytica*. *Microbiology* **151**, 3627–3637.
- 227) Boysen, J.H. and Mitchell, A.P. (2006) Control of Bro1-domain protein Rim20 localization by external pH, ESCRT machinery, and the *Saccharomyces cerevisiae* Rim101 pathway. *Mol. Biol. Cell* **17**, 1344–1353.
- 228) Kullas, A.L., Li, M. and Davis, D.A. (2004) Snf7p, a component of the ESCRT-III protein complex, is an upstream member of the RIM101 pathway in *Candida albicans*. *Eukaryot. Cell* **3**, 1609–1618.
- 229) Rothfels, K., Tanny, J.C., Molnar, E., Friesen, H., Comisso, C. and Segall, J. (2005) Components of the ESCRT pathway, DFG16, and YGR122w are required for Rim101 to act as a corepressor with Nrg1 at the negative regulatory element of the DIT1 gene of *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **25**, 6772–6788.
- 230) Ito, T., Chiba, T., Ozawa, R., Yoshida, M., Hattori, M. and Sakaki, Y. (2001) A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc. Natl. Acad. Sci. USA* **98**, 4569–4574.
- 231) Franz, T., Vingron, M., Boehm, T. and Dear, T.N. (1999) Capn7: a highly divergent vertebrate calpain with a novel C-terminal domain. *Mamm. Genome* **10**, 318–321.
- 232) Yorikawa, C., Takaya, E., Osako, Y., Tanaka, R., Terasawa, Y., Hamakubo, T., Mochizuki, Y., Iwanari, H., Kodama, T., Maeda, T., Hitomi, K., Shibata, H. and Maki, M. (2008) Human calpain 7/PalBH associates with a subset of ESCRT-III-related proteins in its N-terminal

- region and partly localizes to endocytic membrane compartments. *J. Biochem.* **143**, 731–745.
- 233) Osako, Y., Maemoto, Y., Tanaka, R., Suzuki, H., Shibata, H. and Maki, M. (2010) Autolytic activity of human calpain 7 is enhanced by ESCRT-III-related protein IST1 through MIT-MIM interaction. *FEBS J.* **277**, 4412–4426.
- 234) Katoh, K., Shibata, H., Suzuki, H., Nara, A., Ishidoh, K., Kominami, E., Yoshimori, T. and Maki, M. (2003) The ALG-2-interacting protein Alix associates with CHMP4b, a human homologue of yeast Snf7 that is involved in multi-vesicular body sorting. *J. Biol. Chem.* **278**, 39104–39113.
- 235) Missotten, M., Nichols, A., Rieger, K. and Sadoul, R. (1999) Alix, a novel mouse protein undergoing calcium-dependent interaction with the apoptosis-linked-gene 2 (ALG-2) protein. *Cell Death Differ.* **6**, 124–129.
- 236) Sadoul, R. (2006) Do Alix and ALG-2 really control endosomes for better or for worse? *Biol. Cell* **98**, 69–77.
- 237) Vito, P., Pellegrini, L., Guet, C. and D'Adamio, L. (1999) Cloning of AIP1, a novel protein that associates with the apoptosis-linked gene ALG-2 in a Ca²⁺-dependent reaction. *J. Biol. Chem.* **274**, 1533–1540.
- 238) Odorizzi, G. (2006) The multiple personalities of Alix. *J. Cell Sci.* **119**, 3025–3032.
- 239) Gafni, J., Hermel, E., Young, J.E., Wellington, C.L., Hayden, M.R. and Ellerby, L.M. (2004) Inhibition of calpain cleavage of huntingtin reduces toxicity: accumulation of calpain/caspase fragments in the nucleus. *J. Biol. Chem.* **279**, 20211–20220.
- 240) Sokol, S.B. and Kuwabara, P.E. (2000) Proteolysis in *Caenorhabditis elegans* sex determination: cleavage of TRA-2A by TRA-3. *Genes Dev.* **14**, 901–906.
- 241) Syntichaki, P., Xu, K., Driscoll, M. and Tavernarakis, N. (2002) Specific aspartyl and calpain proteases are required for neurodegeneration in *C. elegans*. *Nature* **419**, 939–944.
- 242) Kammenga, J.E., Doroszuk, A., Riksen, J.A., Hazendonk, E., Spiridon, L., Petrescu, A.J., Tijsterman, M., Plasterk, R.H. and Bakker, J. (2007) A *Caenorhabditis elegans* wild type defies the temperature-size rule owing to a single nucleotide polymorphism in *tra-3*. *PLoS Genet.* **3**, e34.
- 243) Dear, N., Matena, K., Vingron, M. and Boehm, T. (1997) A new subfamily of vertebrate calpains lacking a calmodulin-like domain: implications for calpain regulation and evolution. *Genomics* **45**, 175–184.
- 244) Mugita, N., Kimura, Y., Ogawa, M., Saya, H. and Nakao, M. (1997) Identification of a novel, tissue-specific calpain *htra-3*; a human homologue of the *Caenorhabditis elegans* sex determination gene. *Biochem. Biophys. Res. Commun.* **239**, 845–850.
- 245) Waghray, A., Wang, D.S., McKinsey, D., Hayes, R.L. and Wang, K.K. (2004) Molecular cloning and characterization of rat and human calpain-5. *Biochem. Biophys. Res. Commun.* **324**, 46–51.
- 246) Franz, T., Winckler, L., Boehm, T. and Dear, T.N. (2004) Capn5 is expressed in a subset of T cells and is dispensable for development. *Mol. Cell. Biol.* **24**, 1649–1654.
- 247) Gonzalez, A., Saez, M.E., Aragon, M.J., Galan, J.J., Vettori, P., Molina, L., Rubio, C., Real, L.M., Ruiz, A. and Ramirez-Lorca, R. (2006) Specific haplotypes of the CALPAIN-5 gene are associated with polycystic ovary syndrome. *Hum. Reprod.* **21**, 943–951.
- 248) Sáez, M.E., Martínez-Larrad, M.T., Ramirez-Lorca, R., Gonzalez-Sánchez, J.L., Zabena, C., Martínez-Calatrava, M.J., González, A., Morón, F.J., Ruiz, A. and Serrano-Ríos, M. (2007) Calpain-5 gene variants are associated with diastolic blood pressure and cholesterol levels. *BMC Med. Genet.* **8**, 1.
- 249) Fan, C., Iacobas, D.A., Zhou, D., Chen, Q., Lai, J.K., Gavrialov, O. and Haddad, G.G. (2005) Gene expression and phenotypic characterization of mouse heart after chronic constant or intermittent hypoxia. *Physiol. Genomics* **22**, 292–307.
- 250) Tonami, K., Kurihara, Y., Aburatani, H., Uchijima, Y., Asano, T. and Kurihara, H. (2007) Calpain 6 is involved in microtubule stabilization and cytoskeletal organization. *Mol. Cell. Biol.* **27**, 2548–2561.
- 251) Tonami, K., Kurihara, Y., Arima, S., Nishiyama, K., Uchijima, Y., Asano, T., Sorimachi, H. and Kurihara, H. (2011) Calpain 6, a microtubule-stabilizing protein, regulates Rac1 activity and cell motility through interaction with GEF-H1. *J. Cell Sci.* **124**, 1214–1223.
- 252) Baier, L.J., Permana, P.A., Yang, X., Pratley, R.E., Hanson, R.L., Shen, G.Q., Mott, D., Knowler, W.C., Cox, N.J., Horikawa, Y., Oda, N., Bell, G.I. and Bogardus, C. (2000) A calpain-10 gene polymorphism is associated with reduced muscle mRNA levels and insulin resistance. *J. Clin. Invest.* **106**, R69–R73.
- 253) Stumvoll, M., Fritsche, A., Madaus, A., Stefan, N., Weisser, M., Machicao, F. and Häring, H. (2001) Functional significance of the UCSNP-43 polymorphism in the CAPN10 gene for proinsulin processing and insulin secretion in nondiabetic Germans. *Diabetes* **50**, 2161–2163.
- 254) Muramatsu, Y., Taniguchi, Y., Kose, H., Matsumoto, K., Yamada, T. and Sasaki, Y. (2003) *Capn10*, a candidate gene responsible for type 2 diabetes mellitus in the OLETF rat. *IUBMB Life* **55**, 533–537.
- 255) Cheverud, J.M., Fawcett, G.L., Jarvis, J.P., Norgard, E.A., Pavlicev, M., Pletscher, L.S., Polonsky, K.S., Ye, H., Bell, G.I. and Semenkovich, C.F. (2010) Calpain-10 is a component of the obesity-related quantitative trait locus *Adip1*. *J. Lipid Res.* **51**, 907–913.
- 256) Johnson, J.D., Han, Z., Otani, K., Ye, H., Zhang, Y., Wu, H., Horikawa, Y., Misler, S., Bell, G.I. and Polonsky, K.S. (2004) RyR2 and calpain-10

- delineate a novel apoptosis pathway in pancreatic islets. *J. Biol. Chem.* **279**, 24794–24802.
- 257) Ma, H., Fukiage, C., Kim, Y.H., Duncan, M.K., Reed, N.A., Shih, M., Azuma, M. and Shearer, T.R. (2001) Characterization and expression of calpain 10. A novel ubiquitous calpain with nuclear localization. *J. Biol. Chem.* **276**, 28525–28531.
- 258) Paul, D.S., Harmon, A.W., Winston, C.P. and Patel, Y.M. (2003) Calpain facilitates GLUT4 vesicle translocation during insulin-stimulated glucose uptake in adipocytes. *Biochem. J.* **376**, 625–632.
- 259) Arrington, D.D., Van Vleet, T.R. and Schnellmann, R.G. (2006) Calpain 10: a mitochondrial calpain and its role in calcium-induced mitochondrial dysfunction. *Am. J. Physiol. Cell Physiol.* **291**, C1159–C1171.
- 260) Kamei, M., Webb, G.C., Young, I.G. and Campbell, H.D. (1998) *SOLH*, a human homologue of the *Drosophila melanogaster* small optic lobes gene is a member of the calpain and zinc-finger gene families and maps to human chromosome 16p13.3 near *CATM* (cataract with microphthalmia). *Genomics* **51**, 197–206.
- 261) Kamei, M., Webb, G.C., Heydon, K., Hendry, I.A., Young, I.G. and Campbell, H.D. (2000) *Solh*, the mouse homologue of the *Drosophila melanogaster* small optic lobes gene: organization, chromosomal mapping, and localization of gene product to the olfactory bulb. *Genomics* **64**, 82–89.
- 262) Correa, G.C., Margis-Pinheiro, M. and Margis, R. (2001) Identification, classification and expression pattern analysis of sugarcane cysteine proteinases. *Genet. Mol. Biol.* **24**, 275–283.
- 263) Johnson, K.L., Degan, K.A., Ross Walker, J. and Ingram, G.C. (2005) *AtDEK1* is essential for specification of embryonic epidermal cell fate. *Plant J.* **44**, 114–127.
- 264) Tian, Q., Olsen, L., Sun, B., Lid, S.E., Brown, R.C., Lemmon, B.E., Fosnes, K., Gruis, D.F., Opsahl-Sorteberg, H.G., Otegui, M.S. and Olsen, O.A. (2007) Subcellular localization and functional domain studies of DEFECTIVE KERNEL1 in maize and *Arabidopsis* suggest a model for aleurone cell fate specification involving CRINKLY4 and SUPERNUMERARY ALEURONE LAYER1. *Plant Cell* **19**, 3127–3145.
- 265) Hirokawa, T., Boon-Chieng, S. and Mitaku, S. (1998) SOSUI: classification and secondary structure prediction system for membrane proteins. *Bioinformatics* **14**, 378–379.
- 266) Krogh, A., Larsson, B., von Heijne, G. and Sonnhammer, E.L. (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580.
- 267) Ersfeld, K., Barraclough, H. and Gull, K. (2005) Evolutionary relationships and protein domain architecture in an expanded calpain superfamily in kinetoplastid parasites. *J. Mol. Evol.* **61**, 742–757.
- 268) Andrade, H.M., Murta, S.M., Chapeaurouge, A., Perales, J., Nirdé, P. and Romanha, A.J. (2008) Proteomic analysis of *Trypanosoma cruzi* resistance to Benzimidazole. *J. Proteome Res.* **7**, 2357–2367.
- 269) Giese, V., Dallagiovanna, B., Marchini, F.K., Pavoni, D.P., Krieger, M.A. and Goldenberg, S. (2008) *Trypanosoma cruzi*: a stage-specific calpain-like protein is induced after various kinds of stress. *Mem. Inst. Oswaldo Cruz* **103**, 598–601.
- 270) Hertz-Fowler, C., Ersfeld, K. and Gull, K. (2001) CAP5.5, a life-cycle-regulated, cytoskeleton-associated protein is a member of a novel family of calpain-related proteins in *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **116**, 25–34.
- 271) Olego-Fernandez, S., Vaughan, S., Shaw, M.K., Gull, K. and Ginger, M.L. (2009) Cell morphogenesis of *Trypanosoma brucei* requires the paralogous, differentially expressed calpain-related proteins CAP5.5 and CAP5.5V. *Protist* **160**, 576–590.
- 272) Pereira, F.M., Elias, C.G., d'Avila-Levy, C.M., Branquinha, M.H. and Santos, A.L. (2009) Cysteine peptidases in *Herpetomonas samuelpssoai* are modulated by temperature and dimethylsulfoxide-triggered differentiation. *Parasitology* **136**, 45–54.
- 273) Sangenito, L.S., Ennes-Vidal, V., Marinho, F.A., Da Mota, F.F., Santos, A.L., D'Avila-Levy, C.M. and Branquinha, M.H. (2009) Arrested growth of *Trypanosoma cruzi* by the calpain inhibitor MDL28170 and detection of calpain homologues in epimastigote forms. *Parasitology* **136**, 433–441.
- 274) Saïdo, T.C., Sorimachi, H. and Suzuki, K. (1994) Calpain: new perspectives in molecular diversity and physiological-pathological involvement. *FASEB J.* **8**, 814–822.
- 275) Imahori, K. (1987) Calcium-dependent proteases. *Tanpakushitsu Kakusan Koso* **32**, 93–95 (in Japanese).

(Received Jan. 14, 2011; accepted Mar. 4, 2011)

Profile

Hiroyuki Sorimachi was born in 1963 and started working on calpains in 1988 at The Tokyo Metropolitan Institute of Medical Science (Rinshoken) under the supervision of late Prof. Koichi Suzuki, after receiving his B.Sc. in Biochemistry from The University of Tokyo. In 1992, he moved to The University of Tokyo, and obtained his Ph.D. at The University of Tokyo. He continued to study calpains as a Prof. Suzuki's assistant professor, and in 1997 moved to the Laboratory of Prof. Keiko Abe at The University of Tokyo as an associate professor. He was awarded Young Investigator Award from Japan Biochemistry Society in 1999. In 2004, he returned to Rinshoken as a project leader of Calpain Project. His research interests include biochemistry and genetics of all kinds of calpains.



Profile

Shoji Hata was born in 1973 and started his research career as a post-doctoral researcher of Japan Society for the Promotion of Science (JSPS) in 2001 at The University of Tokyo, after receiving his Ph.D. from The University of Tokyo. In 2004, he moved to Rinshoken joining to Calpain Project. In 2010, he was promoted to a chief researcher at Rinshoken. He was awarded Outstanding Presentation Award and Young Investigator's Award at Rinshoken Research Workshop (RRW) in 2009 and at the Japanese Society of Proteases in Pathophysiology (JSP) in 2010. His research interests include biochemistry and genetics of tissue-specific calpains such as stomach-specific, testis-specific, and hair follicle-specific calpains.



Profile

Yasuko Ono was born in 1971 and started her research career as a post-doctoral researcher of JSPS in 1999 at The University of Tokyo, after receiving her Ph.D. from The University of Tokyo. From 2000, she studied on muscle regulatory proteins for three years at University of Arizona, USA as a post-doctoral research fellow abroad of JSPS. In 2003, she became a CREST postdoctoral researcher of Japan Science and Technology Agency at The University of Tokyo. In 2004, she moved to Rinshoken joining to Calpain Project. In 2009, she was promoted to a chief researcher at Rinshoken. She was awarded Murachi Award at FASEB Summer Research Conference in 2001, Young Investigator's Award of the JSPS in 2008, Outstanding Presentation Award at RRW in 2010, and The Young Scientists' Prize of the Commendation for Science and Technology by the Minister of Education, Culture, Sports, Science and Technology in 2011. She has been trying to enjoy tackling to challenge questions casted by the calpains, especially those by numbers 3 and 7.

