


RESEARCH

Open Access



Can a multicomponent multidisciplinary implementation package change physicians' and nurses' perceptions and practices regarding thrombolysis for acute ischemic stroke? An exploratory analysis of a cluster-randomized trial

Md Golam Hasnain¹ , Christopher R. Levi^{1,2}, Annika Ryan^{1,3}, Isobel J. Hubbard¹, Alix Hall³, Christopher Oldmeadow^{1,3}, Alice Grady^{1,3,4,5}, Amanda Jayakody¹, John R. Attia^{1,3,6} and Christine L. Paul^{1,3*}

Abstract

Background: The Thrombolysis ImPlementation in Stroke (TIPS) trial tested the effect of a multicomponent, multidisciplinary, collaborative intervention designed to increase the rates of intravenous thrombolysis via a cluster randomized controlled trial at 20 Australian hospitals (ten intervention, ten control). This sub-study investigated changes in self-reported perceptions and practices of physicians and nurses working in acute stroke care at the participating hospitals.

Methods: A survey with 74 statements was administered during the pre- and post-intervention periods to staff at 19 of the 20 hospitals. An exploratory factor analysis identified the structure of the survey items and linear mixed modeling was applied to the final survey domain scores to explore the differences between groups over time.

Result: The response rate was 45% for both the pre- (503 out of 1127 eligible staff from 19 hospitals) and post-intervention (414 out of 919 eligible staff from 18 hospitals) period. Four survey domains were identified: (1) hospital performance indicators, feedback, and training; (2) personal perceptions about thrombolysis evidence and implementation; (3) personal stroke skills and hospital stroke care policies; and (4) emergency and ambulance procedures. There was a significant pre- to post-intervention mean increase (0.21 95% CI 0.09; 0.34; $p < 0.01$) in scores relating to hospital performance indicators, feedback, and training; for the intervention hospitals compared to control hospitals. There was a corresponding increase in mean scores regarding perceptions about the thrombolysis evidence and implementation (0.21, 95% CI 0.06; 0.36; $p < 0.05$). Sub-group analysis indicated that the improvements were restricted to nurses' responses.

(Continued on next page)

* Correspondence: chris.paul@newcastle.edu.au

¹School of Medicine and Public Health (SMPH), University of Newcastle (UoN), Callaghan, NSW, Australia

³Hunter Medical Research Institute (HMRI), New Lambton Heights, NSW, Australia

Full list of author information is available at the end of the article



(Continued from previous page)

Conclusion: TIPS resulted in changes in some aspects of nurses' perceptions relating to the evidence for intravenous thrombolysis and its implementation and hospital performance indicators, feedback, and training. However, there is a need to explore further strategies for influencing the views of physicians given limited statistical power in the physician sample.

Trial registration: [ACTRN12613000939796](#), UTM: U1111-1145-6762.

Contributions to the literature

- This study demonstrated the ability of a multicomponent implementation intervention to influence the perceptions of clinical staff regarding hospital performance indicators, feedback, and training; and their individual perceptions about the evidence base supporting poststroke thrombolysis and its implementation.
- This change occurred in the context of a transient change in the proportion of patients receiving thrombolysis for acute stroke.
- This study illustrates the importance of exploring processes as well as outcomes in the context of implementation trials, in that a change in clinician perceptions was achieved alongside a transient change in clinical practice. This suggests the need to further explore the perceptions/practice nexus during implementation trials.

Background

Implementation of evidence-based recommendations is critical to delivering optimal clinical care to patients and achieving improvements in health outcomes [1]. However, the adoption of such recommendations into clinical practice often faces barriers [2] at the individual, organizational, and system levels [3].

Acute ischemic stroke (AIS) is one of the leading causes of mortality and morbidity globally [3] and in Australia [4]. Intravenous thrombolysis can improve clinical outcomes if administered within 4.5 h of symptom onset to eligible patients with AIS [5]. However, despite evidence of its efficacy and despite its inclusion in clinical guidelines [6], thrombolysis rates have remained persistently low over the last 10 years, at around 11% among all stroke cases [4]. Several potential barriers restricting the administration of intravenous thrombolysis in patients experiencing AIS [7] have been identified. At the individual level, Shiffman et al. demonstrated that physicians' knowledge of the desired behavior could lead to improvements across quality indicators in emergency care [8]. Similarly, improved skills and

opportunities to perform the desired behavior led to increased staff confidence in the care of patients with chronic cardiac failure [9]. Additionally, individual staff attitudes were associated with adherence to clinical guidelines [10]. In the case of AIS, the strong evidence base has been shown to increase physicians' level of certainty about the impact and administration of intravenous thrombolysis [11, 12], and their familiarity with, and motivation to adhere to, recommended guidelines [7, 13]. At the health systems level, resources and a supportive workplace environment have been shown to positively influence evidence-based practice [3]. Inefficient in-hospital processes for managing emergency stroke patients and a lack of appropriate infrastructure, staffing, and hospital capacity are known barriers to poor rates of intravenous thrombolysis [7, 13]. As both physicians and nurses have a central role in thrombolysis, their perceptions and practices potentially influence intravenous thrombolysis rates [14, 15].

The Thrombolysis ImPlementation in Stroke (TIPS) study investigated a multicomponent, multidisciplinary, collaborative intervention designed to increase rates of intravenous thrombolysis [16]. The TIPS intervention was developed using the Behaviour Change Wheel theoretical framework [16]. The study found significant differences in the rates of intravenous thrombolysis between intervention and control hospitals after 16 months of "active" intervention. However, after the end of the implementation package (assessed over a 12-month post-intervention period), the increased proportions of tPA in intervention hospitals were no longer significant [17]. Recognizing that the views of physicians and nurses are important to achieving practice change [18], the study included a pre- and post-intervention survey which aimed to assess attitudes toward thrombolysis and experiences of various barriers and enablers to thrombolysis implementation. Such data can provide important insights into the intermediate impacts of the intervention and help to shape future interventions aimed at increasing rates of intravenous thrombolysis. Physicians and nurses may face different tasks and challenges even in the same medical environment and therefore, their perceptions and attitudes toward patient care may vary [19]. Hence, it is critically important to explore perceptions and attitudes toward patient care

from both professions. As there are known differences between metropolitan and non-metropolitan hospitals such as staff experience, workload, infrastructure, etc. [20], the location of hospitals should be considered when assessing staff perceptions. Staff perceptions and practices are often explored using study-specific, non-validated measures. Therefore, wherever possible, it is important to identify the validity of the measure; for example, factor analysis can be used to assess how well a new measure captures the intended construct [21].

Therefore, the study aimed to (i) assess the validity of the staff survey measure through the exploratory factor analysis of the survey and (ii) investigate whether the perceptions and practices of the physicians and nurses involved in the TIPS study changed as a result of the TIPS intervention and whether any changes were specific to the practice group (physician or nurse) and hospital location (metropolitan or non-metropolitan).

Methods

Design and setting

The TIPS study recruited 20 hospitals across three states of Australia between 2011 and 2015. It evaluated the effectiveness of a multicomponent, multidisciplinary, collaborative intervention aimed at improving rates of intravenous thrombolysis [16, 17]. TIPS study hospitals were randomized to either receive the intervention activities (intervention hospitals) or continue with usual care (control hospitals) [17]. This study includes two cross-sectional surveys: one during the pre-intervention (2012–mid 2013) and another during the post-intervention (2015) period. We used STrengthening the Reporting of OBservational studies in Epidemiology (STROBE) guideline to report the study (Additional file 1). The survey was anonymous and administered to physicians and nurses employed in the participating hospitals [16].

TIPS intervention and its activities

Intervention components were developed in accordance with the behavior change wheel [16] and strategies with a preferred emphasis on the following components of behavior change wheel: education, persuasion, training, modeling, and enablement. Seven intervention components were delivered actively over 16 months which included pre-workshop meetings, collaborative communal workshops, site-based working groups, web-based training modules, regular telephone case monitoring, bi-monthly feedback of thrombolysis rate, and bi-monthly inter-site teleconferences. Intervention activities and the study timeline are described in Table 1.

Participants

All participating hospitals had a Stroke Unit or the staffing equivalent, and all were in the early stages of implementing intravenous thrombolysis. Participating hospitals included those that were publicly and privately funded and metropolitan and non-metropolitan based. Metropolitan and non-metropolitan hospitals were defined according to the Australian Standard Geographical Classification Remoteness Areas [22]. Eligible survey respondents were physicians and nurses who worked in the Stroke Units and/or the Emergency departments of the participating hospitals and had a role in assessing or managing acute stroke patients during the survey time period.

Procedure

Eligible survey participants were identified by a staff member within each hospital responsible for the organization of stroke care, and this was usually a nurse (survey coordinator). Eligible participants were invited to participate via a combination of email and personal communication. All surveys were completed in hard copy and these were deposited in collection boxes. The survey coordinator provided data on the estimated number of eligible staff, the number of surveys distributed, and the number returned.

Outcome measures

The survey was made up of 74 statements and in the first section, respondents were asked to rate their agreement using a five-point Likert Scale: Strongly disagree, disagree, agree, strongly agree, and not applicable [23] and the second section had some participants and hospital related information. A copy of this survey has been added to this manuscript (Additional file 2: Supplement 1). Survey items were selected from the previously published literature on behavior change and implementation of evidence-based practice. In addition, the National Stroke Foundation's Clinical Guidelines for the management of Stroke and its recommendation for hospital facilities and evidence for intravenous thrombolysis was also considered to finalize the survey items [24, 25]. The survey items were also piloted by a team of behavioral researchers, stroke clinicians, emergency physicians, and stroke nurses at the University of Newcastle and the Hunter New England Local Health District [26]. The survey was divided into two sections and they are as follows.

Section A: stroke care and intravenous thrombolysis

This section was titled "Your Views on tPA" and included 60 statements in total. Of these, 11 were directed at physicians only, and seven were directed at nurses only. Statements investigated issues related to knowledge and skills in assessing AIS and eligible

Table 1 Framework of interventional activity with the timeline

Aug–Sep 2013	↑	Oct 2013	↑	Jan–Dec 2014	↑	March–Nov 2014	↑	March–Nov 2014	↑	March–Dec 2014	↑	Meeting dates not on fixed period	↑	Oct 2014	↑
Executive buy-in (calls or visits)		Workshop 1		Online training modules		Bi-monthly teleconferences		Quarterly feedback on thrombolysis		Regular case monitoring		Site-based working group meetings		Workshop 2	
		1. Goals and target setting 2. Review of current thrombolysis rate 3. Situational analysis 4. Barriers and solutions 5. Forming a site working group		1. Competency-based assessments 2. Clinical decision-making skills 3. Eligibility criteria for stroke thrombolysis		1. Intrahospital collaboration 2. Barriers and solutions		1. Motivation 2. Tracking against targets 3. Feedback		1. Motivation 2. Case review 3. Training update 4. Site update		1. Multidisciplinary collaboration within the hospital 2. Case reviews Target settings 3. Barriers and solutions		1. Review of current and new goals 2. Sustainability of change achieved 3. Planning for the future	

patients for intravenous thrombolysis, hospital policy, and performance indicators for stroke care, impact, safety, and barriers to intravenous thrombolysis, the hospital's monitoring and feedback systems, its emergency service, the protocol used to identify, treat and follow-up patients with stroke, and staff skills levels and staff training facilities. An example of a Section A statement is: "This hospital has goals for improving performance in stroke care".

Section B: individual and hospital characteristics

This section was entitled "About You and Your Workplace" and included 14 questions. Of these, three were directed at physicians only, and one at nurses only. Questions investigated a respondent's age, gender, role within the hospital, and the number of years worked in a stroke unit and/or stroke care. They also investigated a hospital's intravenous thrombolysis rate, pre-arrival notification system from the ambulance, responsibilities around data entry, the proportion of stroke patients seen by Emergency physicians, number of patients referred to stroke care, proportion treated with intravenous thrombolysis, and the respondent's role and responsibilities around intravenous thrombolysis. An example of a Section B statement is "Does the hospital have arrangements in place for pre-arrival notification of stroke patients from the ambulance service?"

Statistical analysis

Identifying construct validity and internal consistency

The frequencies and percentages of the responses for each statement were examined. An exploratory factor analysis was conducted on all statements, except those with > 20% missing values or "not applicable" responses, as they were assessed as having limited relevance [27]. A principal factors method was undertaken to identify the underlying factor structure [28]. The number of factors to retain was determined by assessing the following criteria: (i) the Kaiser-Criteria (eigenvalue more than 1 rule); (ii) the break in the scree plot; and (iii) assessment of parallel analysis [29]. The number of factors suggested by these three criteria oblique rotation was done and the results of each assessed and compared to determine the most appropriate factor structure [29]. The final factor structure was determined as the structure that met the following criteria: items loading on only one factor ≥ 0.4 , minimal cross-loadings between items [30], and the structure that makes conceptual sense (Additional file 2: Supplement 3). Finally, to check and measure the reliability and internal consistency of the final factor structure, the Cronbach's alpha coefficient for each selected factor was calculated [31].

Calculation of scores

Following factor analysis, each selected factor was defined as a domain. Domain scores were calculated for each participant, by summing all statement responses in a domain and dividing them by the number of non-missing statements. The total score for each domain for each participant was four. Likert responses were allocated a score of strongly disagree = 1, disagree = 2, agree = 3, and strongly agree = 4. Only those who responded to at least 75% of the statements within each domain were calculated a domain score [32].

Measuring the effect of the intervention

Separate linear mixed models were conducted for each domain score to identify whether staff statements differed in response to the intervention. Between-group differences in the change in mean staff responses from pre-intervention to post-intervention were assessed. In each model, the main effect for the intervention group and time were included as fixed effects as well a group by time interaction term. Baseline thrombolysis rate was also included as a fixed effect in all models to control for this factor. A random intercept for the hospital was included to account for the clustered design of the trial. To assess our secondary aim, similar models were also conducted separately by profession (i.e., physician and nurse) and geographical location (i.e., metropolitan and non-metropolitan) to allow for assessment of these factors as potential effect modifiers. Due to violations in the assumption of homoscedasticity, robust errors were employed for models assessing domain 1 and 2 of staff barriers. For domain, three and four bootstrap estimation was employed due to violations in the assumption of normality.

Results

All 20 TIPS hospitals were invited to participate in the staff survey; 19 hospitals participated in the pre-intervention survey and 18 of those participated in the post-intervention survey. During the pre-intervention period, of the 1127 eligible staff, 503 returned completed surveys, whereas during the post-intervention period, of the 919 eligible staff, 414 returned completed surveys, equating to a 45% response rate for both surveys. Table 2 reports respondents' characteristics against the intervention and control hospitals and the pre- and post-intervention surveys.

Of the 48 statements included in the factor analysis, 33 statements were maintained representing four domains (Additional file 2: Supplement 2). Domain 1 included 14 statements related to individual and hospital-level performance indicators, feedback, and training. Domain 2 included nine statements related to individual-level perceptions about the evidence supporting thrombolysis

Table 2 Difference in participants' distribution and characteristics between intervention and control hospitals for both pre-intervention and post-intervention survey

	Pre-intervention 503 (55%)		Post-intervention 414 (45%)	
	Intervention 260 (52%)	Control 240 (48%)	Intervention 192 (49%)	Control 202 (51%)
Characteristics				
Age, n (%)				
≤ 25 years	24 (10)	17 (8)	17 (10)	13 (7)
> 25–45 years	149 (62)	117 (54)	101 (57)	109 (59)
> 45–60 years	60 (25)	76 (35)	50 (28)	58 (31)
> 60 years	9 (4)	5 (2)	8 (5)	6 (3)
Sex, n (%)				
Male	84 (34)	75 (32)	54 (29)	70 (36)
Female	164 (66)	160 (68)	130 (71)	125 (64)
Work experience in emergency/stroke, n (%)				
≤ 5 years	90 (35)	79 (34)	49 (27)	62 (32)
> 5–10 years	74 (29)	68 (30)	62 (34)	64 (33)
> 10–15 years	38 (15)	34 (15)	26 (14)	35 (18)
> 15 years	53 (21)	48 (21)	45 (25)	34 (17)
Staff type, n (%)				
Physician	74 (29)	69 (30)	47 (26)	67 (34)
Nurse	181 (71)	163 (70)	137 (74)	129 (66)
Distribution				
Location, n (%)				
Metropolitan	124 (48)	170 (71)	119 (62)	152 (75)
Regional	136 (52)	70 (29)	73 (38)	50 (25)
Baseline thrombolysis rate, n (%)				
Strata 1	106 (41)	134 (56)	82 (43)	123 (61)
Strata 2	127 (49)	76 (32)	83 (43)	63 (31)
Strata 3	27 (10)	30 (12)	27 (14)	16 (8)

and its implementation. Domain 3 included six statements related to staff stroke care skills and hospital stroke care policies. Domain 4 included four statements related to emergency and ambulance procedures. The Cronbach's alphas for domain 1, 2, 3, and 4 were 0.90, 0.79, 0.80, and 0.85 respectively.

When comparing results between the control and intervention hospitals, the domain 1 mean score in the intervention hospitals showed a significant mean increase of 0.21 (95% CI 0.09; 0.34; $p < 0.01$) from the pre- to the post-intervention surveys (Table 3), indicating a positive change in staff perceptions relating to hospital performance indicators, feedback, and training. Similarly, the domain 2 mean score showed a significant increase of 0.21 (95% CI 0.06; 0.36; $p < 0.05$) (Table 4), indicating a positive change in staff perceptions relating to the evidence supporting thrombolysis and its implementation.

The between-group differences for domains 3 and 4 were not significant indicating no change in staff perceptions relating to the individual's level of stroke care skills, the hospital's stroke care policies, and/or the emergency and ambulance procedures (Tables 5 and 6).

When comparing results between physicians and nurses, the domain 1, 2, and 4 mean scores in nurses showed a significant mean increase of 0.25 (95% CI 0.06; 0.44; $p < 0.05$), 0.18 (95% CI 0.01; 0.36; $p < 0.05$), and 0.18 (95% CI 0.02; 0.34) respectively. This indicated a change in nurses' perceptions toward hospital performance indicators, feedback and training, thrombolysis evidence base, and its implementation along with emergency and ambulance service procedure, which was not present in physicians. Moreover, the sub-group analysis based on emergency and stroke care physicians was also no longer significant.

When comparing results between metropolitan and non-metropolitan hospitals, the domain 1 mean score in the non-metropolitan hospitals showed a significant mean increase of 0.26 (95% CI 0.17; 0.35; $p < 0.001$) indicating a change in staff perception related to hospital performance indicators, feedback, and training. In domain 2, the mean score in the metropolitan hospitals showed a significant mean increase of 0.25 (95% CI 0.04; 0.46; $p < 0.05$), indicating a change in staff perception related to the evidence supporting thrombolysis and its implementation. There were no significant differences in domains 3 and 4.

Discussion

This is the first study to evaluate the effect of an implementation intervention aimed at increasing rates of intravenous thrombolysis on physicians' and nurses' perceptions utilizing a psychometrically tested tool for assessment on self-reported behavior. The assessment of the validity of the survey tool indicated that a four-factor structure with items loading on only one factor ≥ 0.4 , minimal cross-loadings between items, and a structure that makes conceptual sense. Thus, the content validity of structure was supported. Moreover, all the domains had satisfactory internal consistency reliability, measured through Cronbach's alpha; ranged 0.79–0.90. Therefore, the survey was considered an appropriate tool for assessing physicians' and nurses' perceptions. The TIPS intervention appeared to have some influence as would be expected according to the BCW framework, via strategies such as education and training. The intervention appeared to be effective in changing the perceptions of clinical staff in relation to their hospitals' performance indicators, feedback and training, and their individual perceptions about the evidence base supporting post-stroke thrombolysis and its implementation. However, this appeared to be the case only for nurses.

Table 3 Effect of intervention on domain 1 score = performance indicator, feedback and training

	Intervention group		Control group		Intervention vs. control group ^{a,b} ; <i>p</i> value
	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	
Overall	2.95 ± 0.48	3.17 ± 0.47	3.03 ± 0.50	3.02 ± 0.46	0.21 (0.09; 0.34)**; 0.001
Location					
Metropolitan	3.07 ± 0.52	3.21 ± 0.47	3.04 ± 0.50	3.03 ± 0.49	0.15 (− 0.07; 0.37); 0.175
Non-metropolitan	2.83 ± 0.40	3.11 ± 0.47	2.99 ± 0.51	2.99 ± 0.41	0.26 (0.17; 0.35)***; 0.000
Job role					
Physician	2.94 ± 0.41	3.05 ± 0.45	2.99 ± 0.51	2.96 ± 0.41	0.12 (− 0.08; 0.31); 0.243
Nurse	2.95 ± 0.50	3.19 ± 0.47	3.07 ± 0.48	3.05 ± 0.46	0.25 (0.06; 0.44)*; 0.010

**p* value < 0.05 considered as significant

***p* value < 0.01 considered as significant

****p* value < 0.001 considered as significant

^aChange from pre-intervention to post-intervention survey

^bLinear mixed model controlled for category based on baseline thrombolysis rate

Several studies have highlighted the use of constructive monitoring and feedback as a strategy for achieving positive changes in hospital-level performance [33, 34]. Moreover, identifying strategic goals is widely used as a means of enhancing organizational motivation, adherence, and autonomy, and in turn, improving processes of care [35, 36]. Hospital performance indicators such as those assessed in the survey can facilitate patient choice, can promote accountability, and finally can increase the quality of patient care [37]. In addition, stroke survivors are known to have complex needs and therefore, require the presence of a multidisciplinary team with specialized knowledge, skills, and experience in stroke [38]. The

development and delivery of stroke-specific education are therefore of vital importance to the provision of high-quality stroke care and to improve outcomes for people who have experienced stroke. TIPS intervention, utilized a “monitoring-evaluation-feedback” strategy that involved site-based leaders where the primary change agents provided regular feedback and conducted monitoring to site champions at each hospital via phone calls throughout the intervention phase, and a knowledge translation strategy that provided web-based training modules, case monitoring, and problem-solving activities [17]. Such knowledge translation strategies have been demonstrated to improve health professionals’ perceptions

Table 4 Effect of intervention on domain 2 score = perceptions about the evidence base for intravenous thrombolysis and its implementation

	Intervention group		Control group		Intervention vs. control group ^{a,b} ; <i>p</i> value
	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	
Overall	3.18 ± 0.47	3.29 ± 0.41	3.24 ± 0.43	3.14 ± 0.49	0.21 (0.06; 0.36)*; 0.007
Location					
Metropolitan	3.20 ± 0.49	3.33 ± 0.38	3.25 ± 0.43	3.12 ± 0.51	0.25 (0.04; 0.46)*; 0.021
Non-metropolitan	3.16 ± 0.46	3.22 ± 0.44	3.23 ± 0.43	3.19 ± 0.46	0.09 (− 0.20; 0.38); 0.523
Job role					
Physician	3.18 ± 0.48	3.10 ± 0.50	3.17 ± 0.40	2.90 ± 0.57	0.19 (− 0.10; 0.48); 0.206
Nurse	3.19 ± 0.47	3.36 ± 0.34	3.28 ± 0.44	3.27 ± 0.39	0.18 (0.01; 0.36)*; 0.039

**p* value < 0.05 considered as significant

***p* value < 0.01 considered as significant

****p* value < 0.001 considered as significant

^aChange from pre-intervention to post-intervention survey

^bLinear mixed model controlled for category based on baseline thrombolysis rate

Table 5 Effect of intervention on domain 3 score = personal stroke skills and hospital stroke care policies

	Intervention group		Control group		Intervention vs. control group ^{a,b} ; <i>p</i> value
	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	
Overall	3.48 ± 0.45	3.60 ± 0.39	3.48 ± 0.47	3.55 ± 0.43	0.04 (− 0.10; 0.18); 0.597
Location					
Metropolitan	3.52 ± 0.45	3.65 ± 0.36	3.48 ± 0.47	3.59 ± 0.39	0.01 (− 0.10; 0.13); 0.828
Non-metropolitan	3.44 ± 0.45	3.52 ± 0.43	3.46 ± 0.47	3.46 ± 0.51	0.09 (−v0.14; 0.22); 0.601
Job role					
Physician	3.49 ± 0.43	3.63 ± 0.40	3.48 ± 0.48	3.57 ± 0.40	0.04 (− 0.14; 0.22); 0.696
Nurse	3.49 ± 0.45	3.59 ± 0.39	3.48 ± 0.47	3.54 ± 0.46	0.04 (− 0.13; 0.21); 0.670

**p* value < 0.05 considered as significant

***p* value < 0.01 considered as significant

****p* value < 0.001 considered as significant

^aChange from pre-intervention to post-intervention survey

^bLinear mixed model controlled for category based on baseline thrombolysis rate

of evidence-based approaches and their implementation [39, 40]. However, why this was more effective in nurses, as opposed to physicians, in our study is uncertain.

Failure of the TIPS intervention to change the perceptions of physicians may be due to limited statistical power (*n* = 74 physicians). However, the evidence does indicate that it can be challenging to change physicians’ perceptions about their clinical practice because they are long-standing and widely held [41, 42]. It is possible that physicians are more cautious in these potentially high-risk circumstances, given their level of responsibility for decision-making. Finally, an additional explanation may be that more nurses than physicians were engaged with

the TIPS online learning modules made available to participants [17].

Hospital policies are often difficult to change as they involve complex systems [37, 38]. Given this complexity and a very limited focus on policy in the TIPS intervention, it is perhaps unsurprising that no change was found for the domain scores related to hospital policy and individual skills. While skills may be more amenable to change than policies, a ceiling effect may have been evident in relation to skills: According to results from the pre-intervention survey [26], most participants in the TIPS were skilled in post-stroke thrombolysis. For example, 98% could correctly assess stroke and 83% could correctly identify patients eligible for intravenous

Table 6 Effect of intervention on domain 4 score = perceptions toward emergency service

	Intervention group		Control group		Intervention vs. control group ^{a,b} ; <i>p</i> value
	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	Pre-intervention (Mean ± SD)	Post-intervention (Mean ± SD)	
Overall	3.08 ± 0.61	3.40 ± 0.49	3.15 ± 0.58	3.36 ± 0.56	0.10 (− 0.07; 0.27); 0.178
Location					
Metropolitan	3.13 ± 0.58	3.44 ± 0.48	3.18 ± 0.54	3.34 ± 0.53	0.14 (− 0.09; 0.38); 0.219
Non-metropolitan	3.03 ± 0.63	3.34 ± 0.52	3.09 ± 0.66	3.41 ± 0.62	− 0.02 (− 0.36; 0.30); 0.860
Job role					
Physician	3.09 ± 0.67	3.13 ± 0.50	3.13 ± 0.55	3.26 ± 0.55	− 0.11 (− 0.38; 0.16); 0.398
Nurse	3.07 ± 0.58	3.50 ± 0.45	3.17 ± 0.57	3.41 ± 0.55	0.18 (0.02; 0.34)*; 0.041

**p* value < 0.05 considered as significant

***p* value < 0.01 considered as significant

****p* value < 0.001 considered as significant

^aChange from pre-intervention to post-intervention survey

^bLinear mixed model controlled for category based on baseline thrombolysis rate

thrombolysis. The result from Grady et al. [25] was very similar, i.e., 98% of emergency physicians reported they could identify a stroke patient clearly, and 76% could identify patients eligible for intravenous thrombolysis. Finally, perceptions related to emergency services were not part of the TIPS intervention which accords with the null finding in relation to the domain score regarding perception toward emergency care.

Unfortunately, the study struggled with several limitations. As mentioned before, the number of participating physicians was low, which may have limited the study's statistical power in relation to physician data, and the data were self-reported. The study was also not powered to evaluate the effect of the changes in perception at an individual site level. Therefore, the study suggests a need to explore in more depth (e.g., via more comprehensive mixed methods) the experience of the staff involved in implementation interventions and intervention uptake/adherence to developing better and sustainable intervention for the future.

Conclusion

The TIPS intervention appeared to have more impact on changing the perception of nurses than physicians, particularly in the domains of hospitals' performance, and feedback and training and perceptions about the thrombolysis evidence and its implementation. This further reinforces evidence about how challenging it is to change clinical practice and to effectively and efficiently bridge the evidence-practice gap.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13012-019-0940-0>.

Additional file 1. STROBE Statement.

Additional file 2. Supplement 1. Domains selected by factor analysis. Supplement 2. Domains selected by factor analysis. Supplement 3. Conceptual sense for the Factor structure.

Abbreviations

AIS: Acute ischemic stroke; ASGC: Australian Standard Geographical Classification; NHMRC: National Health and Medical Research Council; NSW: New South Wales; TIPS: Thrombolysis Implementation in Stroke

Acknowledgments

We are thankful to all staffs and patients from the participating hospitals for their kind participation and support during the study period.

Authors' contributions

MGH, CLP, and CRL designed and prepared the study concept. MGH and AH analyzed the data with advice from CLP and CO. MGH drafted the manuscript with advice from CLP, JRA, AR, AG, AJ, AH, CO, IH, and CRL on the method of data analysis and presentation of results. CLP, JRA, AR, AG, AJ, AH, CO, IH, and CRL were involved in critical revision of the manuscript. All authors read and approved the final manuscript.

Funding

This study was funded by the National Health and Medical Research Council (NHMRC) partnership grant (569328), part-funded by an NHMRC Practitioner Fellowship (1043913) and NHMRC Translating Research Into Practice fellowship, and included partnership grant contribution funding from Boehringer Ingelheim, in-kind support from Agency for Clinical Innovation Stroke Care Network/Stroke Services New South Wales (NSW), Stroke Foundation and NSW Cardiovascular Research Network-National Heart Foundation, with a cash contribution from the Victorian Stroke Clinical Network and infrastructure funding from the Hunter Medical Research Institute and The University of Newcastle.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Ethical approval was obtained from the Hunter New England Human Research Ethics Committee and from the University of Newcastle, Human Research Ethics Committee. Written informed consent was taken from all the participants.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Medicine and Public Health (SMPH), University of Newcastle (UoN), Callaghan, NSW, Australia. ²The Sydney Partnership for Health, Education, Research & Enterprise (SPHERE), Liverpool, NSW, Australia. ³Hunter Medical Research Institute (HMRI), New Lambton Heights, NSW, Australia. ⁴Hunter New England Local Health District, Population Health, Wallsend, NSW, Australia. ⁵Priority Research Centre for Health Behaviour, University of Newcastle, Callaghan, NSW, Australia. ⁶John Hunter Hospital, New Lambton Heights, NSW, Australia.

Received: 23 May 2019 Accepted: 23 September 2019

Published online: 27 November 2019

References

- Grol R. Successes and failures in the implementation of evidence-based guidelines for clinical practice. *Med Care*. 2001;39(8 suppl 2):146–54.
- Carey M, Buchan H, Sanson-Fisher R. The cycle of change: implementing best-evidence clinical practice. *Int J Qual Health Care*. 2009;21:37–43.
- Kuklina EV, Tong X, George MG, Bansil P. Epidemiology and prevention of stroke: a worldwide perspective. *Expert Rev Neurother*. 2012;12(2):199–208.
- Stroke Foundation. Interim response to ACEM thrombolysis review. Published on: 24 2016 Published on: Aug 7, 2014 cited on: may 10, 2017. Available on: <https://strokefoundation.org.au/News/2016/03/24/Interim-Response-to-ACEM-review-of-tPA>
- Lees KR, Bluhmki E, von Kummer R, et al. ECASS, ATLANTIS, NINDS and EPITHET rt-PA study group. Time to treatment with intravenous alteplase and outcome in stroke: an updated pooled analysis of ECASS, ATLANTIS, NINDS, and EPITHET trials. *Lancet* 2010;375(9727):1695–1703.
- Stroke Foundation. Clinical guidelines for stroke management 2017. Available on: <https://informme.org.au/Guidelines/Clinical-Guidelines-for-Stroke-Management-2017>. Accessed on: 4 January, 2019.
- Eissa A, Krass I, Bajorek BV. Barriers to the utilization of thrombolysis in acute ischemic stroke. *J Clin Pharm Ther*. 2012, 37, 399–409.
- Shiffman RN, Dixon J, Brandt C, Essaihi A, Hsiao A, Michel G, et al. The guideline implementability appraisal (GLIA): development of an instrument to identify obstacles to guideline implementation. *BMC Med Inform Decis Mak* 2005; 5:23.
- Asch SM, Baker DW, Keesey JW, Broder M, Schonlau M, et al. Does the collaborative model improve care for chronic heart failure? *Med Care* 2005; 43(7):667–675.
- Grol R, Dalhuijsen J, Thomas S, Veld C. Attributes of clinical guidelines that influence use of guidelines in general practice: observational study. *BMJ* 1998 Sep; 317:858–861.

11. Meurer WJ, Majersik JJ, Frederiksen SM, Kade AM, Sandretto AM, Scott PA. Provider perceptions of barriers to the emergency use of tPA for acute ischemic stroke: a qualitative study. *BMC emerg*. 2011;11:5.
12. Kwan J, Hand P, Sandercock P. Improving the efficiency of delivery of thrombolysis for acute stroke: a systematic review. *QJM*. 2004;97:273–9.
13. Ehlers L, Groth L, Bech M, Andersen G, Kjølby M. Organisational barriers to thrombolysis treatment of acute ischaemic stroke. *Curr Med Res Opin*. 2002; 23:2833–9.
14. Morgenstern LB, Bartholomew LK, Grotta JC, Staub L, King M, Chan W. Sustained benefit of a community and professional intervention to increase acute stroke therapy. *Arch Intern Med*. 2003 Oct 13;163(18):2198–202.
15. Scott PA, Meurer WJ, Frederiksen SM, et al. A multilevel intervention to increase community hospital use of alteplase for acute stroke (INSTINCT): a cluster-randomised controlled trial. *Lancet Neurol* 2012; 12(2):139–148.
16. Paul CL, Levi CR, D'Este CA, Parsons MW, Bladin CF, Lindley RI, et al. Thrombolysis Implementation in stroke (TIPS): evaluating the effectiveness of a strategy to increase the adoption of best evidence practice—protocol for a cluster randomised controlled trial in acute stroke care. *Implement Sci*. 2014;9:38.
17. Levi CR, Attia JR, D'Este C, et al. A cluster randomised trial of thrombolysis implementation support in metropolitan and regional Australian stroke centres; lessons for individual and systems behaviour change (Unpublished).
18. Baatiema L, Otim ME, Mnatzaganian G, de-Graft Aikins A, Coombes J, Somerset S. Health professionals' views on the barriers and enablers to evidence-based practice for acute stroke care: a systematic review. *Implement Sci*. 2017; 12(1):74.
19. Huang CH, Wu HH, Lee YC. The perceptions of patient safety culture: a difference between physicians and nurses in Taiwan. *Appl Nurs Res*. 2018; 40:39–44.
20. Australian Bureau of Statistics (ABS) 2017. Health service usage and health related actions, Australia 2014–15. ABS cat. No. 4364.0.55.002. Canberra: ABS.
21. Gorsuch RL. Exploratory factor analysis: its role in item analysis. *J Pers Assess*. 1997;68(3):532–60.
22. Australian Government Department of Health. Remoteness classification systems. Available on: <http://www.health.gov.au/internet/publications/publishing.nsf/Content/work-res-ruraid-toc~work-res-ruraid-lis~work-res-ruraid-lis-e> Accessed on: 7 January, 2019.
23. Sullivan GM, Artino AR. Analyzing and interpreting data from likert-type scales. *J Grad Med Educ* 2013;5(4):541–542.
24. Grady A, Bryant J, Carey M, Paul C, Sanson-Fisher R. Enablers of the implementation of tissue plasminogen activator in acute stroke care: a cross-sectional survey. 2014; *PLoS ONE* 9(12): e114778.
25. Grady AM, Bryant J, Carey ML, Paul CL, Sanson-Fisher RW, Levi CR. Agreement with evidence for tissue plasminogen activator use among emergency physicians: a cross-sectional survey. *BMC Res Notes*. 2015;8:267.
26. Paul CL, D'Este CA, Ryan A, Attia JR, Oldmeadow C, Kerr E, Henskens F, Grady A, Levi CR. Staff perspectives from Australian hospitals seeking to improve implementation of thrombolysis Care for Acute Stroke. *SAGE Open Med* 2019; 7: 2050312119865656.
27. Schmitt N. Uses and abuses of coefficient alpha. *Psychol Assess*. 1996;8(4): 350–3.
28. Brunius Enlund K, Brunius C, Hanson J, Hagman R, Höglund OV, Gustås P, et al. Development and validation of two questionnaires: dental home care and dental health in Swedish dogs. *PLoS One*. 2019;14(1):e0204581.
29. Asún RA, Rdz-Navarro K, Alvarado JM. Developing multidimensional Likert scales using item factor analysis: the case of four-point items. *Sociol Methods Res*. 2016;45(1):109–33.
30. Costello AB, Osborne JW. Best practices in exploratory factor analysis: four recommendations for getting the most from your analysis. *Practical Assessment Research & Evaluation*. 2005;10(7)
31. Floyd FJ, Widaman KF. Factor analysis in the development and refinement of clinical assessment instruments. *Psychol Assess*. 1995;7(3):286–99.
32. Trockel M, Bohman B, Lesure E, Hamidi MS, Walle D, Roberts L, et al. A brief instrument to assess both burnout and professional fulfillment in physicians: reliability and validity, including correlation with self-reported medical errors, in a sample of resident and practicing physicians. *Acad Psychiatry* 2017;42(1):11–24.
33. Majeed A, Lester H, Bindman A. Improving the quality of care with performance indicators. *BMJ*. 2007;335:916–8.
34. Marshall M, Campbell S, Hacker J, Roland M. Quality indicators for general practice. London: Royal Society of Medicine; 2002.
35. Wade DT. Goal planning in stroke rehabilitation: why? *Top Stroke Rehabil* 1999; 6:1–7.
36. Levack WM, Dean SG, Siegert RJ, McPherson KM. Purposes and mechanisms of goal planning in rehabilitation: the need for a critical distinction. *Disabil Rehabil*. 2006;28:741–9.
37. Botje D, Ten Asbroek G, Plochg T, Anema H, Kringos DS, Fischer C, Wagner C, Klazinga NS. Are performance indicators used for hospital quality management: a qualitative interview study amongst health professionals and quality managers in the Netherlands. *BMC Health Serv Res*. 2016;16(1):574.
38. Jones SP, Miller C, Gibson JME, Cook J, Price C, Watkins CL. The impact of education and training interventions for nurses and other health care staff involved in the delivery of stroke care: an integrative review. *Nurse Educ Today*. 2018;61:249–57.
39. Brownson RC, Fielding JE, Maylahn CM. Evidence-based public health: a fundamental concept for public health practice. *Annu Rev Public Health*. 2009;30:175–201.
40. Nutley S, Walter I, Davies H. From knowing to doing: a framework for understanding the evidence into practice agenda. *Eval Program Plann*. 2003;9:125–48.
41. Mostofian F, Ruban C, Simunovic N, Bhandari M. Changing physician behavior: what works? *Am J Manag Care*. 2015;21(1):75–84.
42. Alvanzo AAH, Cohen GM, Nettleman M. Changing physician behavior: half-empty or half-full? *Clinical Governance*. 2003;8(1):69–78.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

