

Cancer-associated bone disease

RIZZOLI, René, *et al.*

Reference

RIZZOLI, René, *et al.* Cancer-associated bone disease. *Osteoporosis international*, 2013, vol. 24, no. 12, p. 2929-2953

DOI : 10.1007/s00198-013-2530-3

PMID : 24146095

Available at:

<http://archive-ouverte.unige.ch/unige:33723>

Disclaimer: layout of this document may differ from the published version.



**UNIVERSITÉ
DE GENÈVE**

Cancer-associated bone disease

R. Rizzoli · J.-J. Body · M.-L. Brandi · J. Cannata-Andia · D. Chappard · A. El Maghraoui · C. C. Glüer · D. Kendler · N. Napoli · A. Papaioannou · D. D. Pierroz · M. Rahme · C. H. Van Poznak · T. J. de Villiers · G. El Hajj Fuleihan · for the International Osteoporosis Foundation Committee of Scientific Advisors Working Group on Cancer-Induced Bone Disease

Received: 5 August 2013 / Accepted: 25 September 2013 / Published online: 22 October 2013

© International Osteoporosis Foundation and National Osteoporosis Foundation 2013

Abstract Bone is commonly affected in cancer. Cancer-induced bone disease results from the primary disease, or from therapies against the primary condition, causing bone fragility. Bone-modifying agents, such as bisphosphonates and denosumab, are efficacious in preventing and delaying cancer-related bone disease. With evidence-based care

pathways, guidelines assist physicians in clinical decision-making. Of the 57 million deaths in 2008 worldwide, almost two thirds were due to non-communicable diseases, led by cardiovascular diseases and cancers. Bone is a commonly affected organ in cancer, and although the incidence of metastatic bone disease is not well defined, it is

This position paper has been endorsed by the Committee of Scientific Advisors of IOF

R. Rizzoli (✉)
 Division of Bone Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland
 e-mail: Rene.Rizzoli@unige.ch

J.-J. Body
 Department of Medicine, CHU Brugmann, Université Libre de Bruxelles, Brussels, Belgium

M.-L. Brandi
 Department of Internal Medicine, University of Florence, Florence, Italy

J. Cannata-Andia
 Bone and Mineral Research Unit, Instituto Reina Sofia, REDinREN, ISCIII, Hospital Universitario Central de Asturias, Universidad de Oviedo, Oviedo, Spain

D. Chappard
 GEROM-Research Group on Bone Remodeling and bioMaterials, LUNAM University, Angers, France

A. El Maghraoui
 Rheumatology Department, Military Hospital Mohammed V, Rabat, Morocco

C. C. Glüer
 Sektion Biomedizinische Bildgebung, Klinik für Diagnostische Radiologie, Universitätsklinikum Schleswig-Holstein, Kiel, Germany

D. Kendler
 Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

N. Napoli
 Division of Endocrinology, University Campus Bio-Medico, Rome, Italy

A. Papaioannou
 Department of Medicine, McMaster University, Hamilton, Ontario, Canada

D. D. Pierroz
 International Osteoporosis Foundation (IOF), Nyon, Switzerland

M. Rahme · G. El Hajj Fuleihan
 Calcium Metabolism and Osteoporosis Program, WHO Collaborating Center for Osteoporosis and Metabolic Bone Disorders, American University of Beirut Medical Center, Beirut, Lebanon

C. H. Van Poznak
 University of Michigan, Ann Arbor, MI, USA

T. J. de Villiers
 Panorama MediClinic and Department of Gynaecology, Faculty of Health Sciences, Stellenbosch University, Cape Town, South Africa

estimated that around half of patients who die from cancer in the USA each year have bone involvement. Furthermore, cancer-induced bone disease can result from the primary disease itself, either due to circulating bone resorbing substances or metastatic bone disease, such as commonly occurs with breast, lung and prostate cancer, or from therapies administered to treat the primary condition thus causing bone loss and fractures. Treatment-induced osteoporosis may occur in the setting of glucocorticoid therapy or oestrogen deprivation therapy, chemotherapy-induced ovarian failure and androgen deprivation therapy. Tumour skeletal-related events include pathologic fractures, spinal cord compression, surgery and radiotherapy to bone and may or may not include hypercalcaemia of malignancy while skeletal complication refers to pain and other symptoms. Some evidence demonstrates the efficacy of various interventions including bone-modifying agents, such as bisphosphonates and denosumab, in preventing or delaying cancer-related bone disease. The latter includes treatment of patients with metastatic skeletal lesions in general, adjuvant treatment of breast and prostate cancer in particular, and the prevention of cancer-associated bone disease. This has led to the development of guidelines by several societies and working groups to assist physicians in clinical decision making, providing them with evidence-based care pathways to prevent skeletal-related events and bone loss. The goal of this paper is to put forth an IOF position paper addressing bone diseases and cancer and summarizing the position papers of other organizations.

Keywords Bone · Cancer · IOF · Skeletal-related events

Abbreviations

ABCSG	Austrian Breast & Colorectal Cancer Study Group
ADT	Androgen deprivation therapy
AI	Aromatase inhibitor
ALP	Alkaline phosphatase
ASCO	American Society of Clinical Oncology
ATAC	Arimidex, Tamoxifen Alone or in Combination
AZURE	Adjuvant Zoledronic Acid to Reduce Recurrence
BTM	Bone turnover marker
BMD	Bone mineral density
CI	Confidence interval
CMF	Cyclophosphamide–methotrexate–5 fluorouracil
CT	Computed tomography
CTX	Cross-linked terminal telopeptide
DKK1	Dickkopf-related protein
DXA	Dual-energy X-ray absorptiometry
ER	Estrogen receptor

ET	Endothelin
ESCEO	European Society for Clinical and Economical Aspects of Osteoporosis and Osteoarthritis
FAC	5-Fluorouracil–doxorubicin–cyclophosphamide
FDG	Fluorodeoxyglucose
FGF	Fibroblast growth factor
FRAX	WHO fracture risk assessment tool
GnRH	Gonadotropin-releasing hormone
HR	Hazard ratio
IGF	Insulin-like growth factor
IES	Intergroup Exemestane Study
IL	Interleukin
IV	Intravenous
MRI	Magnetic resonance imaging
NCCN	National Comprehensive Cancer Network
NTX	N-terminal telopeptide
ONJ	Osteonecrosis of the jaw
OPG	Osteoprotegerin
PDGF-BB	Platelet-derived growth factor-BB
PET	Positron emission tomography
PSA	Prostate-specific antigen
PTHrP	Parathyroid hormone-related protein
RANK	Receptor activator of nuclear factor kappa-B
RANKL	Receptor activator of nuclear factor kappa-B ligand
RCT	Randomized controlled trial
RR	Relative risk
SABRE	Study of Anastrozole with the Bisphosphonate Risedronate
SERM	Selective oestrogen receptor modulator
sFRP	Secreted frizzled-related protein
SRE	Skeletal-related event
TGF- β	Transforming growth factor-beta
TNF- α	Tumour necrosis factor-alpha
TRAcP	Tartrate-resistant acid phosphatase
TSH	Thyroid-stimulating hormone
VEGF	Vascular endothelial growth factor
VFA	Vertebral fracture assessment
WHO	World Health Organization
Z-Fast and ZO-Fast	Zometa-Femara Adjuvant Synergy Trials
α CTX	Alpha cross-linked telopeptides type

Epidemiology of cancer-associated bone disease

Bone metastasis

Cancer affects nearly 12.7 million people and is associated with over 7 million deaths in 2008 [1]. Cancer is a rising global

health burden and it is estimated that in 2030, cancer deaths will be tallied at over 13 million (World Health Organization (WHO) 2010 Global Status Report [2]). Addressing the morbidity and mortality of cancer is an important public health concern. On purpose, we are limiting our analysis to cancer bone involvement in adults and do not discuss cancer in children.

Metastases from cancer are associated with 90 % of cancer deaths [3]. It was estimated that one half of people who die from cancer in the USA have bone involvement [4–6]. Different tumour types may have preferential sites of metastases; however, the vast majority of tumours metastasize to bone, albeit at varying frequencies. The term metastatic bone disease reflects the spread of a tumour to the bone. This term may be applied to solid tumours, as well as to multiple myeloma, where the tumour is intrinsic to the bone marrow. In multiple myeloma, 70–95 % of patients have tumour bone disease [7]. In breast cancer, bone is often the first site of distant metastases [5] with approximately one half of patients experiencing bone metastases as the site of first relapse [8]. In advanced breast or prostate cancer, metastatic bone disease is present in the vast majority of patients. Bone metastases may also be seen in 15–30 % of cancers of the lung, gastrointestinal tract (colon and stomach) and the genitourinary (bladder, kidney and uterus) [9] (Table 1). In advanced thyroid cancer and melanoma, bone metastases are also frequently present [7]. Skeletal-related events (SREs) occur relatively commonly and include pathologic fractures (20.7 %), spinal cord compression (0.9 %), surgery (1.2 %) and radiotherapy (8.0 %) to bone and may or may not include hypercalcaemia of malignancy while skeletal complication also refers to pain and other symptoms, thus impairing quality of life, and also decreasing survival [7] and may encompass SREs as well. New anticancer and supportive care treatment modalities are alleviating symptoms, including bone-related ones, and maintaining or improving quality of life, at the expense in some instances of accelerated bone loss and fractures.

Cancer-related bone loss and fracture in patients who do not have bone metastases

Patients with cancer may be at increased risk of bone loss secondary to cancer disease treatment [10]. With improved

Table 1 Incidence of metastatic bone disease [42]

	Incidence (%)
Myeloma	70–95
Breast	65–75
Prostate	65–75 (NCCN, Grawlow)
Renal	20–25
Melanoma	14–45
Thyroid	60
Lung	30–40
Bladder	40

survival rates in many types of cancer, early identification and treatment of osteoporosis among cancer patients could prevent unnecessary fractures, morbidity and reductions in quality of life. A recent prospective study in Germany of 1,041 cancer patients, mean age of 57 years, 78 % female, found elevated rates of osteoporosis compared to the general population [10]. The prevalence of osteoporosis in both men and women with cancer in complete remission was 16 % (95 % confidence interval (CI) 13.8–18.2) and osteopenia 44 % (defined using WHO criteria). Rates of osteoporosis were not statistically different among various cancer types, which included breast, gynaecological, prostate, colorectal and haematological cancers, although sample size was too small to detect potential differences in rates between subtypes [10].

Women who have been treated medically for breast cancer may be at increased risk for bone loss and fractures [11, 12]. In a case-controlled study of over 1,200 women with newly diagnosed breast cancer and no metastases, the annual incidence of vertebral fractures was 2.72 % compared to 0.53 % in the control arm, i.e. a fivefold increase, with rates adjusted for age, prevalent fractures and duration of follow-up [12]. Similarly, in the prospective observational arm of the Women's Health Initiative, fracture rates in breast cancer survivors were increased by 68.6 fractures per 10,000 person-years, which compared to rates in women without breast cancer is a 15 % increase, after adjustment for age, ethnicity, weight and geographic location [11]. Kanis et al. [12] found a fivefold higher prevalence of vertebral fractures in women with breast cancer, but without bone metastases, than in women of the same age (odds ratio 4.7, 95 % CI 2.3–9.9).

Women with oestrogen receptor positive (ER+) breast cancer treated with aromatase inhibitors (AIs) as adjuvant endocrine therapy are at increased risk of rapid bone loss, and of fractures [13–15]. For example in the landmark Arimidex, Tamoxifen Alone or in Combination (ATAC) trial, average bone loss rates in women assigned to the AI anastrozole were 1–2 %/year, well above those recorded in women in the tamoxifen arm [16].

Men with prostate cancer are at particularly high risk of osteoporosis [17] and of fracture [18] in part due to treatment with androgen deprivation therapy (ADT). Rates of bone mineral density (BMD) decrease could be as high as 3.0 to 5.6 % within the first year of ADT, depending on the measured site, with annual decreases of 1.1 to 2.3 % thereafter [19].

Pathophysiology

Fractures in cancer-associated bone disease can result from the direct or systemic effect of the tumour itself or from therapies used to treat the primary disease. In the former case, they are related to local effects of metastatic deposit in bone and/or to generalized bone loss from tumour-produced systemically

circulating bone resorbing hormones or cytokines. These comprise parathyroid hormone-related protein (PTHrP), like in lung and breast cancer, or tumour stimulated secretion by the osteoblast of local bone resorbing factors such as receptor activator of nuclear factor kappa-B ligand (RANKL), interleukin (IL)-6 or IL-3, like in multiple myeloma. Alternatively, bone loss may result from gonadal ablation by chemotherapy or endocrine ablative therapy to treat the primary disease. In some tumours, more than one mechanism may be operating [20].

Bone metastasis

The bone microenvironment represents a fertile soil capable of favouring the growth of malignant cells coming from a distant tumour (metastasis) or of haematological origin (myeloma or lymphoma). Relationships between bone remodelling and metastatic cells are summarized by the “seed and the soil” theory proposed in 1879 by Stephan Paget, who noticed that some cancer cells had an increased propensity to migrate and expand in bone (cited in [21]). The preferential targeting of the skeleton by some tumours is in part explained by tumour-specific factors, and by relevant modulators in the bone microenvironment, that enhance tumour growth. Indeed, some particular cancer cells (breast, prostate, malignant lymphocytes and plasma cells) possess characteristics that favour their anchorage in the bone marrow. The pathophysiology of bone metastasis has been reviewed extensively elsewhere [22–24]. Here, we will summarize the main aspects of the interactions between malignant cells, bone cells and bone remodelling.

The metastatic cascade

From a primary tumour, malignant cells can acquire the capacity to metastasize due to an increased motility and invasiveness and a special tropism to bone or bone marrow. They may produce or express various adhesive molecules for, e.g. integrins, which can bind to ligand molecules–receptors expressed by the stromal cells of the bone marrow or to non-collagenic proteins present in the bone matrix, such as osteopontin. When circulating within the blood stream, cancer cells can reach the sinusoid capillaries of the bone marrow, which also contain large pores [25]. Malignant cells can adhere to these endotheliums and extravase into the bone marrow environment.

Bone metastases are usually classified as osteolytic, osteoblastic/osteosclerotic, or mixed (osteolytic and osteoblastic), based on their appearance on X-ray images. Many patients may have both osteolytic and osteosclerotic metastases, and individual bone lesions can harbour both features. Predominantly osteolytic metastases are typical of multiple myeloma, renal cell cancer, thyroid cancer, non-small lung cancer, some gastro-intestinal tumours, and melanomas.

Predominantly osteosclerotic lesions are most often observed in prostate cancer, carcinoid, gastrinoma and medulloblastoma. Mixed lesions occur most commonly in breast cancer, gastrointestinal tumours and most squamous cell cancers at their primary site. In osteolytic metastases, bone resorption is due to a stimulation of osteoclastogenesis (and not by a direct action of the tumour cells), and areas of metaplastic bone at the margin of the lytic lesions are observed [26] such as in non-small lung carcinoma, non-Hodgkin’s lymphoma and thyroid cancer. In osteosclerotic tumours, there is an increase in osteoclastogenesis but the stimulation of bone formation is more pronounced.

Development of osteolytic metastasis

Malignant cells release a number of molecules that favour osteoclastogenesis via the receptor activator of nuclear factor kappa-B (RANK)/RANKL/osteoprotegerin (OPG) system. PTHrP may activate this bone resorption pathway and is detected by immunohistochemistry in about 90 % of bone metastases from breast cancer, i.e. more often than in soft tissue metastases [27]. RANKL has also been shown to trigger the migration to bone of melanoma and of some epithelial cancer cells that express the RANK receptor, such as breast cancer cells. In a mouse model, by blocking RANKL, OPG resulted in a reduction in tumour burden in bone, but not in other organs [28]. Other bone resorbing factors such IL-1, IL-6, IL-8, IL-11 and TNF- α have also been identified. Osteoclastogenesis in both cortical and trabecular bones is increased in the vicinity of the tumour by a paracrine mechanism. In advanced metastatic bone disease, hypercalcaemia reflects the release of large amounts of calcium mobilized due to breakdown of the calcified matrix. During resorption, large amounts of deeply entrapped growth factors in bone are released and activated in the microenvironment. Transforming growth factor-beta (TGF- β), insulin-like growth factor (IGF)-I, and IGF-II promote the growth of the tumour cells locally through their receptors, e.g. breast cancer cells express receptors for TGF- β [24]. α CTX fragments of degraded collagen are also released from the eroded areas and represent a strong chemoattractant for recruiting locally new malignant cells [29].

Development of osteosclerotic metastasis

Prostatic adenocarcinoma is predominantly associated with osteosclerosis metastases. On histological bone sections, a large number of osteoblasts building new trabeculae are observed in the vicinity of the tumour cells. The neo-trabeculae made of woven bone are anchored at the surface of pre-existing trabeculae and fill the marrow cavity. Prostate adenocarcinoma cells are able to release a number of cytokines that induce osteoclastogenesis including PTHrP. However, malignant cells also express a variety of proteases, e.g. prostate-

specific antigen (PSA) which partially degrades PTHrP or IGF-binding proteins, and limit osteoclastogenesis. One characteristic of prostate cancer cells is their production of cytokines that favour osteoblastogenesis: ET-1 [30, 31], IGF-I and IGF-II, FGF-1 and FGF-2, VEGF. In turn, activated osteoblasts can release large amounts of IL-6, TGF- β and PDGF-BB which are potent growth factors for the tumour cells [32–34]. In both osteoblastic and osteoclastic metastases, a vicious circle is established since the malignant cells stimulate osteoblast or osteoclastic activity, which in turn stimulates tumour growth and progression.

Bone loss in myelomas and lymphomas

Lytic bone lesions are observed on X-ray in about 95 % of patients with advanced myeloma, in contrast to what is observed in lymphomas, although both entities are B cells and B- and T cells malignancies, respectively. Bone involvement is related to an excessive bone resorption through increased osteoclast number encountered in the close vicinity of myeloma cells [35] together with a decreased osteoblast activity [36–38]. In myeloma, the cytokines produced by plasma cells and bone cell progenitors (IL-6, macrophage inflammatory protein-1 α /CCL-3) induce the genesis of mature multinucleated osteoclasts. In lymphoma, when B cells invade the bone [39], a mixed population of multi- and mononuclear osteoclasts is observed, mononuclear TRAcP⁺ cells being only capable of microresorption [40]. Osteolytic lesions are usually rare in B cell lymphoma (8–10 % of cases) and occur only when the number of these cells is high [41]. In myelomas and lymphomas, a marked reduction in osteoblast activity has been identified by histomorphometric analysis [36]. Plasma cells can release several factors such as DKK1, sFRP 2, which act on the Wnt pathway and reduce the osteoblast number and activity [38]. The lesions observed in myeloma are predominantly osteolytic and can produce removal of whole trabeculae and perforation of cortical bone.

Cancer-related bone loss and fracture

A variety of mechanisms are responsible for bone loss in patients with cancer treatment-associated bone loss. The mechanisms may vary according to patient profile and chemotherapeutic regimen used. These include hypogonadism induced by chemotherapy, hormone ablative therapy, glucocorticoid, surgical castration, irradiation [42–44] or any combination of the above. Indeed, bone loss has been observed in lymphoma survivors who received therapy regimens including corticosteroids, alkylating agents and radiation therapy, all of which can cause hypogonadism [45, 46]. The highest rates for bone loss observed are in premenopausal women who experience cytotoxic or endocrine-induced acute ovarian ablation, reaching 8 % at the spine and 4 % at the hip, within the

first year [47, 48], compared to half those rates in postmenopausal women receiving AIs [49–51].

Chemotherapy-induced hypogonadism

Predictors of premature ovarian failure in women with breast cancer receiving cytotoxic chemotherapy include patient's age and type of regimen used. It was reported to occur in 60–85 % of women receiving adjuvant therapy with CMF, and in 50 % of women receiving the FAC regimens, with age-specific rates of 33 % in women aged 30–39 years, 96 % for women between 40 and 49 years and 100 % in women above age 50 [52] (Table 2). Permanent ovarian failure was observed following individual therapeutic doses of various chemotherapy agents including cyclophosphamide, chlorambucil and mitomycin-C.

Cyclophosphamide appears to be the most common agent implicated in chemotherapy-associated amenorrhea. Premature menopause is dependent on its cumulative dose [52]. Chlorambucil and cyclophosphamide can cause prolonged azoospermia in male patients [53]. In addition, gonadal toxicity was evident in patients, especially those with testicular cancer, receiving a cumulative dose of cisplatin greater than 400 mg/m² [54]. Bone loss was also observed in patients made hypogonadic by cytotoxic drugs used in haematopoietic stem cell transplantations [55]. In premenopausal women, tamoxifen treatment is associated with bone loss, through its antiestrogen effects, whereas it is rather bone protective after the menopause being a partial oestrogen agonist [56, 57].

Table 2 Cancer treatment regimens directly and indirectly associated with bone loss [43]

Direct effects	Indirect effects
Androgen deprivation therapy	Hypogonadism
Oestrogen suppression	Hyperparathyroidism
Glucocorticoids/corticosteroids	Vitamin D deficiency
Methotrexate	Gastrectomy
Megestrol acetate	Hyperprolactinemia
Platinum compounds	
Cyclophosphamide	
Doxorubicin	
Interferon-alpha	
Valproic acid	
Cyclosporine	
Vitamin A	
NSAIDS	
Estramustine	
Ifosfamide	
Radiotherapy	
Combination chemotherapy regimens	

Hormone deprivation therapy

Osteoporosis stemming from hypogonadism is frequently seen in survivors of breast and prostate cancer, as therapeutic hypogonadism is an important strategy for controlling these hormone-dependent tumours [58].

Aromatase inhibitors

In women with ER+ breast cancers (about 70 % of tumours), AIs aim to reduce oestrogen levels by inhibiting the aromatization of androgens and their conversion to oestrogens in peripheral tissues [15]. Third generation non-steroidal (anastrozole and letrozole) and steroidal (exemestane which is similar to androstenedione) AI drugs inhibit the aromatase enzyme by 96–99 % [51], with substantial reduction in oestrogen concentrations (Table 2). AI-induced bone loss is generally more rapid and severe than bone loss in normal postmenopausal women [15], and should be taken into account especially when treated women with low BMD and/or fracture [59]. The skeletal effects observed are inversely correlated with baseline BMD and serum estradiol concentrations. Osteoporosis is more prevalent in women starting AI early after menopause, and there is only a partial recovery of BMD following the withdrawal of AI. Bone loss is accompanied by higher fracture risk [12, 15, 60–63]. This differs greatly from the effects of tamoxifen or raloxifene given in the prevention of breast cancer recurrence where increases in bone turnover may be partially averted and bone loss prevented. AIs are superior to tamoxifen as adjuvant therapy of ER+ breast cancer, with longer disease-free survival and without the risk of endometrial hyperplasia and cancer, cerebrovascular and venous thromboembolic events, but exhibit other toxicities such as arthralgias [15, 64]. Recently, AIs have been shown to further reduce the risk of recurrence after the diagnosis of ER+ breast cancer, either when given instead of tamoxifen or when administered sequentially after few years of tamoxifen therapy, and are thus now recommended in the adjuvant setting [65]. There is evidence that even the low levels of oestrogen in postmenopausal women are important for bone health. In the MORE clinical trial, comparing raloxifene to placebo, there was an inverse correlation between oestrogen levels and prospectively recorded fracture risk in those women assigned to the placebo group [66]. Letrozole, anastrozole and exemestane increase bone turnover [64] and decrease BMD, and letrozole and anastrozole increase the relative risk of vertebral and non-vertebral fractures by 40 %, when compared with tamoxifen. After a few years of AI use, women have a 20–35 % increased fracture risk [67]. For instance, fracture risk has been reported to increase by 55–115 % with anastrozole in the ATAC [68] and Austrian Breast & Colorectal Cancer Study Group (ABCSCG) [69] trials, by 15–50 % with letrozole in the BIG-198 [70] and MA17 [71]

trials, and by 41 % with exemestane in the IES trial [72], as compared to tamoxifen or placebo. However, recent data show that bone loss associated to treatment with aromatase inhibitors in breast cancer patients is influenced by CYP19 polymorphisms [73].

Sex hormone deprivation therapy with GnRH agonists and antagonists

Long-acting gonadotropin-releasing hormone (GnRH) agonists with increased receptor affinity or prolonged half-life lead to persistent activation of gonadotropin-releasing hormone receptors, causing an initial release of pituitary gonadotropins followed by a down-regulation of GnRH receptor and suppression of gonadotropin secretion. Consequently, ovarian sex steroids production is suppressed. GnRH agonists are effective in the management of endometriosis and of breast cancer in premenopausal women by suppressing oestrogen levels but they are inducing bone loss (Table 2). A BMD decrease of about 6 %/year is observed in patients on GnRH agonists with a recovery of bone mass after discontinuation. GnRH agonists may not increase the risk of fragility fractures in women with normal BMD [43].

In men with hormone-sensitive prostate cancer, castration or ADT can be induced surgically or medically with GnRH agonists or antagonists and a combination of GnRH agonist with androgen biosynthetic blockade. ADT is effective in reducing tumour extension, growth and improving survival [74]. ADT can be enhanced by the addition of androgen biosynthetic blockade such as cyproterone. The latter two strategies have a different effect on the skeleton. ADT by either bilateral orchiectomy or GnRH agonist increases the risk of fractures [18, 75–77]. In contrast, anti-androgens alone increase rather than decrease BMD in two randomized trials [78, 79]. Androgen deficiency-mediated decrease in lean body mass, increase in fat mass and impaired muscular strength may contribute to increased fracture risk [80–84] (Table 2). Abiraterone is an inhibitor of CYP17A1, an enzyme required in androgen synthesis. It reduces androgen and precursors steroids, and is associated with a better survival. Abiraterone is administered with prednisone 5 mg twice daily, or use of another steroid. At this time, there is insufficient bone data on this treatment regimen [85].

In men with prostate carcinoma, BMD of hip, ultradistal radius and lumbar spine decreases by 2–5 % after 12 months of ADT, and the risk of vertebral and hip fractures increases by 40–50 % [18, 77]. Fracture risk increases linearly with the number of GnRH injections. It has been estimated that fracture risk in men on ADT is as high as 20 % by 5 years of treatment [18, 86]. Older men and those with osteoporosis risk factors in addition to age are at the highest risk [87].

Radiation-induced hypogonadism and direct toxic effects on bone

Both female and male cancer survivors who received irradiation to the cranium, ovaries or testes can display hypogonadism [88]. The association between male hypogonadism and osteoporosis is further supported by reports of fractures occurring after external beam radiation therapy to the prostate bed for prostate cancer [89–92] (Table 2). In addition, pelvic or rib fracture can occur in relation with bone local irradiation [93, 94].

Glucocorticoids

The majority of therapeutic regimens for many haematopoietic malignancies involve high-dose glucocorticoids, usually administered over extended periods of time. Glucocorticoids, which are often used as a pain adjuvant, palliative agent, antiemetic or as part of the treatment, initially increase bone resorption, then later suppress osteoblast activity reducing thereby bone formation [88] (Table 2). Prolonged exposure to corticosteroids is the third leading cause of osteoporosis, after hypogonadism and advancing age [95, 96]. The risk of fracture increases by 50–100 % in recipients of oral corticosteroids [97].

Other indirect effects of cancer therapies

There are other indirect effects on bone health that result from cancer therapies [43]. These effects include hypovitaminosis D, cachexia (sarcopenia) and decreased mobility (Table 2).

Fracture risk assessment in patients with cancer and adjuvant therapies

Patients with cancer-associated osteoporosis are generally younger than patients with postmenopausal osteoporosis. As people with cancer survive longer, cancer-related skeletal event and their treatments are becoming increasingly recognized as important co-morbidities [98]. Early evaluation of risk factors for osteoporosis, including family history of fracture and assessment of peripheral neuropathy that may have occurred secondary to cancer therapy, medication review, physical examination, fall risk assessment, diet and exercise assessment and counselling, as well as changes in BMD over time, are of prime importance. Fracture risk assessment in this unique population can be challenging. It should include evaluation of BMD and clinical risk factors as detailed below.

Bone mineral density

BMD measurement by dual X-ray absorptiometry (DXA) is the most common clinical tool to directly measure bone mineral

mass and indirectly evaluate bone strength [99]. BMD should be measured by DXA at spine and hip with measurements at the 1/3 radius considered if either one of these sites is not available. Malignancies in bone may either be lytic (e.g. myeloma) or blastic (e.g. breast or prostate), and if present in the region of interest, artifact is introduced. Infrequently, DXA images may give an indication of skeletal metastases requiring other imaging follow-up. Although radius BMD changes little with osteoporosis therapy, hypogonadism (e.g. from ADT) and hyperparathyroidism often lead to loss of forearm BMD [100]. Several guidelines recommend that postmenopausal women with breast cancer on AI, premenopausal women with ovarian failure secondary to cancer treatment and prostate cancer patients on ADT should have their BMD measured [42, 101]. For instance, the American Society of Clinical Oncology (ASCO) recommends DXA BMD testing in postmenopausal women taking AIs and premenopausal women who develop treatment-related premature menopause [102]. The Belgian Bone Club recommends measuring DXA BMD in all women starting AIs or medical castration therapy [103]. The UK Expert Group recommends measuring DXA BMD within 3 to 6 months of commencing AIs in all women, except for those ≥ 75 years of age, in whom treatment decisions are based upon age and clinical risk factor assessment, independently of BMD [101]. Similarly, an International Expert Panel recommends BMD measurement in patients with breast cancer initiating or receiving AI therapy [104] as does also the European Society for Clinical and Economical Aspects of Osteoporosis and Osteoarthritis (ESCEO) [15]. Patients with bone metastases receiving monthly bone resorption inhibitors do not require BMD assessment.

Clinical risk factors: 10-year absolute fracture risk

WHO fracture risk assessment tool (FRAX) provides an algorithm applicable for men older than 40 years and for postmenopausal women: 10-year fracture risk of the hip and of major osteoporotic fracture (wrist, proximal humerus, hip and clinical spine) with and without BMD. Including BMD improves the predictive performance of the score, that is, improves sensitivity without decreasing specificity [105, 106]. Although the FRAX calculator has provided a major advance in our assessment of fracture risk, it does not take into consideration some other risk factors, due to the nature of the clinical information available in the cohorts the model was developed from [107].

There is no way of estimating the impact of malignancy or its treatment from the FRAX algorithm [108], though cancer treatment-induced bone loss could be considered as secondary osteoporosis, the role of which in FRAX fracture assessment is entirely captured by BMD. Clinicians should thus use their clinical judgement to quantify their patient's individual fracture risk.

There are two studies that used the FRAX calculation in men on ADT. In a cross-sectional study [109], FRAX

identified more men at risk for fracture than BMD alone. Age was a very important risk factor. Adler reported [87] that FRAX derived estimates using femoral neck BMD or calculated without BMD defined different populations at risk.

Other risk factors

Fall risk, bone turnover markers and many other risk factors for fracture and bone loss are not included in FRAX. A variety of measures of prospective fall risk have been shown to be useful in predicting fractures. They include questions or questionnaire-based tools, simple physical performance tests, measurements of muscle mass and devices that measure some aspect of strength, balance or integrated function [110, 111]. Bone turnover markers hold promise in fracture risk prediction and for monitoring treatment. However, there are still uncertainties about their use in clinical practice [112].

Medications

Numerous medications other than glucocorticoids, such as proton pump inhibitors, can predispose to bone loss (Table 2). They have not been included in FRAX modelling due to the infrequency of their use in the general population, their lack of evaluation in epidemiologic studies and their presumed lesser effects on bone metabolism.

Prevalent fracture

Prevalent fracture is a powerful predictor for further fractures, e.g. hip fracture risk is increased by more than twofold by a prevalent hip or spine fracture, independently of BMD, some of them being pathologic fractures. Indeed, a spine fracture increases the risk of hip fracture (relative risk (RR)=2.5), subsequent spine fracture (RR=4.4), forearm fracture (RR=1.7) or proximal humerus fracture (RR=1.9) [113, 114]. A greater number or higher severity of vertebral fracture increases the risk of vertebral and non-vertebral fracture. However, number, severity, location of fractures and glucocorticoid dose and duration are not captured in the current FRAX algorithm [107, 115]. Pathologic fracture versus osteoporotic fracture may be difficult to distinguish from plain radiographs and consideration should be given to other imaging such as computed tomography (CT), magnetic resonance imaging (MRI) or isotope bone scanning. Asymptomatic vertebral fractures are very common when appropriate imaging is applied [116]. Concerns for metastatic osseous lesions may be confirmed by biopsy.

Detection of metastatic cancer to bone

Among the many techniques used for imaging metastases, each has advantages and limitations [117]. The two “classical”

methods, radiography and scintigraphy, both have limited sensitivity and specificity. CT and MRI are preferable because they provide more accurate information [118, 119]. Modern devices permit scans that cover a large part of the body with acceptable radiations exposure (for CT) and in reasonable time (for MRI). While MRI is more suited to detect early infiltration when the tumour is still restricted to bone marrow and offers better insight into soft tissue involvement, CT provides a better characterization of bone integrity. More sophisticated approaches like positron emission tomography (PET), today usually used in conjunction with CT (PET/CT), may also be considered, but the choice of the radiotracer is important. FDG PET is most commonly used, but the magnitude of the improvement over scintigraphy is controversial [120, 121]. ^{18}F -Fluoride PET or ^{18}F -choline PET [122] may be better choices but are more expensive and rarely available. The choice of bone imaging is impacted by the primary tumour type, the patient's symptoms and location of area(s) in question.

Compromised bone integrity may endanger the spinal cord or other critical structures. Urgent surgical intervention to stabilize the skeleton and relieve pressure on nerves may be required. For long bones, lesions that involve 50 % of the diameter of the cortex or that are larger than 2.5–3 cm are considered at risk for pathological fracture. Clinical judgment is required to evaluate all skeletal complications of metastatic disease. Particularly at the spine, the assessment of mechanical stability is complex. New approaches for evaluation of stability based on CT may be more accurate, allowing both a better quantification of regional bone density and specific imaging of bone at high risk of pathologic fracture. In addition, finite element analysis methods developed successfully for evaluation of bone strength in osteoporosis [123] may be adapted in the future to metastatic bone [124, 125].

Prevention and treatment

Prevention of skeletal-related events

Bisphosphonates

Several studies have shown that bisphosphonates have anti-tumour potentials with direct and indirect effects that result not only in less bone loss but less tumour burden as well. They interrupt the vicious cycle in the bone microenvironment between tumour cells and osteoclasts described in section on pathophysiology above. Bisphosphonates, in particular zoledronic acid, also enhance $\gamma\delta$ T cell differentiation (a T lymphocyte subpopulation that plays a main role against tumour cells) and have also potent anti-angiogenic activity in the adjuvant setting [23]. There is no clear impact of osteoclast inhibiting therapy on tumour burden. However, two recent phase III studies suggest that potent osteoclast inhibition may impact overall survival [126, 127].

Breast cancer in the adjuvant setting The efficacy of bisphosphonates in reducing SREs in women with breast cancer and bone metastatic disease is unequivocal, as demonstrated in a recent Cochrane meta-analysis [128]. While they reduce the occurrence of SREs by 15 to 40 % [129], they do not impact SREs in subjects without bone metastases [130]. In the frame of adjuvant therapies, evidence from a secondary endpoint in the Zometa-Femara Adjuvant Therapy Synergy Trial (ZO-Fast) supports the hypothesis that bisphosphonates have an anti-metastatic role in breast cancer patients. In more than 1,000 recruited patients, those in the upfront group had lower disease recurrence or death than patients in the delayed group (1.1 vs 2.3 %). No local disease recurrences occurred in upfront patients while it occurred in 0.6 % of the delayed group. Efficacy of zoledronic acid was confirmed also in premenopausal women with a significant improvement in disease-free survival shown in the ABCSG-12 trial [131]. Both loco-regional and extraskelatal metastases risk were reduced, suggesting a systemic anti-tumour effect exerted by zoledronic acid. The results of these two adjuvant bisphosphonate studies must be viewed in conjunction with the outcomes of three other phase III adjuvant bisphosphonates trials. The Adjuvant Zoledronic Acid to Reduce Recurrence (AZURE) trial randomized 3,360 women with stage II or III early stage breast cancer to receive standard adjuvant systemic therapy with or without zoledronic acid. The bisphosphonate dosing was incrementally decelerated over the course of 5 years; however, total dose of zoledronic acid administered was much greater than either ZoFast or ABCSG-12. The primary endpoint in AZURE was disease-free survival. At a median follow-up of 59 months, AZURE demonstrated disease recurrence or death in 377 women in the zoledronic acid group and 375 in the control group [130]. In subset analysis, postmenopausal women appeared to have gained benefit from the addition of zoledronic acid. Note that to reach a neutral overall study outcome, premenopausal women had to have displayed a worse outcome with the addition of zoledronic acid. There is much speculation as to account for these findings. In patients with bone metastasis, a retrospective analysis based on 578 patients showed that zoledronic acid treatment normalized N-terminal telopeptide (NTX) levels in 81 % of the treated patients within the third month. Moreover normalization of NTX was associated with reduced risk of SREs and improved overall survival [132]. The National Surgical Adjuvant Breast and Bowel Project study B-34 [133] randomized 3,323 women to adjuvant clodronate or placebo with disease-free survival in women with stage I, II or III breast cancer as primary endpoint. At a median follow-up of 90.7 months, there were 286 events in the clodronate group and 312 events in the controls, which was a non-significant difference (95 % CI 0.78, 1.07; $p=0.27$). Similarly, in the study GAIN, 3,023 women with lymph node positive breast cancer were randomized to either oral ibandronate or placebo. There was equal disease free and overall survival in the

two groups [134]. In another phase III study, patients were randomized to clodronate, ibandronate or zoledronic acid (without control or placebo arm). The results are due soon [135].

Multiple meta-analyses have been performed using published data and came to different conclusions, depending on the studies selected. It is anticipated that the Early Breast Cancer Trialists' Collaborative Group will perform a meta-analysis based upon raw data and provide thus greater insight into the situation.

Prostate cancer Earlier studies using pamidronate and clodronate failed to demonstrate a reduction in SREs in patients with prostate cancer and bone metastases. In a randomized controlled trial (RCT) in patients with castration-resistant prostate cancer with bone metastases, zoledronic acid (4 mg, 3-weekly) or placebo reduced SREs ($p=0.009$) and prolonged the median time to first SRE from 321 to 488 days ($p=0.009$) [136]. Bisphosphonates have not been shown to prevent bone metastases due to prostate cancer in any study. An ongoing study in men with metastatic prostate cancer compares the early use of zoledronic acid (within 3 months of initiation of ADT) to standard zoledronic acid (on diagnosis of castration resistance). The primary end point is the proportion of subjects experiencing SREs.

Denosumab

RANKL is a key mediator of metastatic bone resorption. Denosumab is a human monoclonal antibody that binds and neutralizes human RANKL. It prevents RANKL from activating RANK on osteoclasts, inhibiting osteoclast formation, function and survival, and hence reducing bone resorption. Therefore, RANKL inhibition through denosumab is a therapeutic target for preventing and treating bone metastases.

Breast cancer In a study evaluating the efficacy of denosumab in 2,046 breast cancer patients with bone metastases in a double-blind double-dummy trial, denosumab (120 mg monthly) was superior to zoledronic acid (4 mg monthly with adaptation of the dose to the renal function) in suppressing bone turnover and delaying or preventing SREs. Denosumab increased the time to first on-study SRE by 18 % compared with zoledronic acid (hazard ratio (HR), 0.82; $p<0.001$ for non-inferiority and $p=0.01$ for superiority). The median time to first on-study SRE was 26.4 months for the zoledronic acid group and had not been reached for the denosumab treatment group. Denosumab also delayed the time to first and subsequent (multiple) on-study SREs by 23 % compared with zoledronic acid (multiple event analysis; $p=0.001$). The mean skeletal morbidity rate (defined as the ratio of the number of SREs per patient divided by the patient's time at risk) was also lower with denosumab than with zoledronic acid (0.45 vs 0.58 events per patient per year; $p=0.004$), which represents a reduction of 22 % with denosumab. Overall survival and

disease progression were similar in the two treatment groups. Safety profile, including onset of osteonecrosis of the jaw, was similar between both groups [137].

Prostate cancer In a study comparing denosumab (120 mg monthly) to zoledronic acid (4 mg monthly) for the prevention of SREs in 1,904 men with castration-resistant prostate cancer metastatic to bone, the primary end point was the time to first SRE. Denosumab proved to be non-inferior ($p=0.0002$) and superior ($p=0.008$) compared to zoledronic acid. Denosumab delayed the time to first on-study SRE by 18 % compared with zoledronic acid (HR, 0.82) and the median time to first on-study SRE was 3.6 months longer with denosumab compared with zoledronic acid. Denosumab also significantly delayed the time to first and subsequent on-study SREs by 18 % compared with zoledronic acid ($p=0.008$). Overall survival and disease progression evaluated by changes in PSA levels did not differ between both groups [138].

In the adjuvant setting, when denosumab 120 mg every 4 weeks was compared to placebo in preventing bone metastases in 1,432 men with castration-resistant prostate cancer and at high risk of developing bone metastases, bone metastasis-free survival was higher ($p=0.028$) and time to first bone metastasis increased ($p=0.032$) in the denosumab group. Denosumab improved bone metastasis-free survival by a median of 4.2 months compared with placebo (29.5 vs 25.2 months; HR, 0.85; $p=0.028$). Time to symptomatic bone metastasis was also prolonged. Overall survival was similar in both groups, but study medications were stopped when the first bone metastasis was diagnosed. Denosumab was associated with higher incidence of hypocalcaemia (1.7 vs 0.3 %) and osteonecrosis of the jaw (4.6 vs 0 % at the end of year 3) [139].

Other solid tumours and multiple myeloma In a third study with an identical design, 1,776 patients with multiple myeloma or bone metastases and solid tumours (except those of prostate or breast) were randomized to receive denosumab or zoledronic acid. Approximately 40 % of patients in the study had non-small cell lung cancer and 10 % multiple myeloma. Denosumab was non-inferior to zoledronic acid in delaying the time to first on-study SRE (HR, 0.84; $p=0.0007$), but, after adjustment for multiple comparisons, the difference between the groups was not statistically significant for treatment superiority ($p=0.06$). The median time to first on-study SRE was 20.6 months for denosumab and 16.3 months for zoledronic acid. Overall survival and disease progression were similar in both treatment groups [140].

Integrated analysis of the three phase III trials in metastatic patients

The identical design of the three phase III studies conducted in patients with metastatic bone disease allowed for a pre-

planned integrated analysis of the data involving a total of more than 5,700 patients. Denosumab increased the time to first on-study SRE by 17 % over zoledronic acid (HR, 0.83; 95 % CI, 0.76–0.90; $p<0.001$ for superiority). The median time to first SRE was 27.7 months with denosumab and 19.4 months with zoledronic acid. This integrated analysis showed that denosumab delayed the time to first and subsequent on-study SREs by 18 % compared with zoledronic acid (HR, 0.82; 95 % CI, 0.75–0.89; $p<0.001$ for superiority). Overall survival and disease progression were similar with both treatments [132]. The integrated analysis showed that the cumulative incidence of confirmed osteonecrosis of the jaw (ONJ) after 3 years was not negligible ($n=89$ from total of 5,723 patients; 1.6 %) with no significant difference between denosumab and zoledronic acid treatments (1.8 vs 1.3 %; $p=0.13$) [141]. The role of denosumab in the management of hypercalcaemia of malignancy has not yet been evaluated and clinical trials should be conducted to more clearly document this effect.

Conclusion for the prevention of skeletal-related events in breast cancer

In a recent meta-analysis of 34 RCTs conducted in women with various stages of breast cancer, Wong et al. have demonstrated the unequivocal efficacy of bone resorption inhibitors in reducing the incidence of SREs in women with bone metastatic disease [128]. Bisphosphonates reduced the risk of SREs by 15 % (RR=0.85, 95 % CI, 0.77–0.94), in nine studies of 2,806 women with bone metastases. This benefit was observed with intravenous (IV) zoledronic acid (4 mg/3–4 weeks) with RR 0.59 (95 % CI, 0.42–0.82), IV pamidronate (90 mg/3–4 weeks) (RR=0.77; 95 % CI, 0.69–0.87) and IV ibandronate (RR=0.80; 95 % CI, 0.67–0.96), with one direct comparison in a large study confirming the equivalent efficacy of zoledronate and pamidronate. In three studies comprising 3,405 patients with skeletal metastases, denosumab was superior to a bisphosphonate in reducing the risk of SREs (RR=0.78; 95 % CI, 0.72–0.85) [128]. Such therapies did not influence the outcomes of women without bone metastases at study entry. The authors also concluded that there is insufficient evidence to make a conclusion about the role of adjuvant bisphosphonates in reducing visceral metastases, loco-regional recurrence and total recurrence or improving survival. Toxicity was generally mild kidney toxicity and ONJ were reported at similar rates for patients on denosumab compared to zoledronic acid.

Prevention of bone loss and fractures

As in idiopathic osteoporosis, preventive measures to maintain bone health include lifestyle changes, adequate calcium and

vitamin D, regular physical exercise and pharmacotherapy such as bisphosphonates, denosumab or selective oestrogen receptor modulators (SERMs).

Lifestyle, calcium and vitamin D

Lifestyle interventions are crucial in order to improve quality of life and maximize any pharmacological treatment in cancer patients. The American Cancer Society, the World Cancer Research Fund/American Institute for Cancer Research and the American College of Sports Medicine have released guidelines for cancer survivors [142–144], recommending at least 30 min daily of physical activity for at least 5 days per week. Exercise interventions are also beneficial for muscle strength and bone mass, reducing also the risk of falls and, in turn, potentially of hip fractures.

Patients with malignancies are exposed to risk factors that may predispose to vitamin D deficiency. Among them, inadequate sunlight exposure, poor dietary intakes and treatment with medications that reduce vitamin D absorption are the most common [145, 146]. Indeed, 76–88 % of breast cancer survivors have low vitamin D levels, i.e. <20 ng/mL [147–149]. A recent study has shown that low vitamin D levels are highly prevalent among newly diagnosed breast cancer patients with nearly 44 % with vitamin D insufficiency [150]. As expected, African-Americans are more affected than Caucasians [150]. Clinical trials using vitamin D in cancer patients are lacking. In a recent double-blind placebo-controlled randomized phase II trial, breast cancer patients on AIs, treated with vitamin D₂, had less aromatase inhibitor-induced musculoskeletal symptoms than those on placebo. Vitamin D-treated patients had also a stable BMD at the femoral neck (0.45 %±0.72), compared to a 1.39 % decrease seen in the placebo group [151]. The ASCO recommends supplementation with calcium and vitamin D in breast cancer patients, mostly if treated with AIs [102]. However, routine 25-hydroxyvitamin D measurement is not part of the current guidelines in breast cancer patients. Given the lack of clinical trials, no guidelines on specific levels of supplementation for vitamin D or calcium are available. Vitamin D supplementation should be given in order to bring serum 25-hydroxyvitamin D levels to 20 ng/mL (50 nmol/L) or higher [152]. According to data obtained in postmenopausal women, a daily intake of at least 800 IU should be advised together with a daily calcium intake (from food and supplements) of 1,000 mg. The intake of calcium from the diet should be favoured because it has been suggested that high doses of calcium supplements were associated with increased cardiovascular risks [153]. These recommendations have been extended from National Comprehensive Cancer Network (NCCN) Bone Health in Cancer Care Task Force to premenopausal women at risk of cancer treatment-associated bone loss.

As advised by the NCCN Prostate Cancer Guidelines (available at www.nccn.org), similar preventive measures,

with calcium and vitamin D supplementation, should be applied to patients with prostate cancer, particularly if on ADT. A sufficient intake of calcium and vitamin D should be emphasized in those on intravenous bisphosphonates or denosumab to avoid the risk of hypocalcaemia.

Bisphosphonates

Bisphosphonates have been successfully used in patients with metastatic bone diseases to reduce SREs [154]. However, their use has been extended to prevent bone loss in patients with both breast or prostate cancer in the adjuvant setting, who are treated with chemotherapy or hormonal therapy [131]. This approach is in accordance with the wide use of bisphosphonates in idiopathic osteoporosis [155].

Breast cancer In premenopausal women with breast cancer, adjuvant chemotherapy induces early menopause, a strong factor of accelerated bone loss. Few trials have evaluated the efficacy of bisphosphonates to attenuate chemotherapy-induced bone loss in young women [156–164]. The administration of oral clodronate at daily doses of 1,600 mg, oral risedronate at a dose of 30 mg/day, cyclical intravenous pamidronate at doses of 60 mg every 3 months and intravenous zoledronic acid at a dose of 4 mg every 3 months prevented chemotherapy-induced bone loss in these studies. Numerous other studies have confirmed such findings and are summarized in Table 3.

Bisphosphonates have also been shown to prevent AI-induced bone loss. The SABRE trial has tested in an open label approach the effect of risedronate in breast cancer patients on anastrozole. At 24 months follow-up, medium-risk patients on risedronate had a significant increase in lumbar spine and total hip BMD compared with anastrozole and placebo (2.2 vs -1.8 and 1.8 vs -1.1 %, respectively). In the high-risk group, lumbar spine and total hip BMD increased (3.0 and 2.0 %) while those on anastrozole had a 2.1 % decrease [165]. In the ARIBON study, early stage breast cancer patients receiving anastrozole were treated with ibandronate (high risk), ibandronate or placebo (medium risk), or anastrozole only (low risk). At 24 months, ibandronate-treated patients (150 mg orally every month), compared to placebo, had a higher BMD and suppressed markers of bone turnover [166]. Overall, these data confirm that oral bisphosphonates improve BMD and may normalize bone turnover in AI-treated patients.

Prevention of AI-induced bone loss has been tested also using intravenous infusions of zoledronic acid. The Z-Fast and ZO-Fast were designed to compare effects of zoledronic acid (4 mg intravenously) initiated either with letrozole and every 6 months thereafter (upfront group) or given only when bone loss became significant or a fragility fracture occurred (delayed group) [147]. In over 1,667 patients, the upfront

Table 3 Guidelines for bone management in breast cancer

Resources	Target population	Indication	Type of treatment, concentration, duration	Monitoring	Other recommendations
American Society of Clinical Oncology (ASCO) [102, 154]	Patients with metastatic breast cancer	Patients with metastatic breast cancer with evidence of bone metastases	sc denosumab 120 mg every 4 weeks IV pamidronate 90 mg over 2 h every 3–4 weeks IV zoledronic acid 4 mg over 15 min every 3–4 weeks	Serum creatinine monitored before each dose of BP If serum creatinine clearance >60 mL/min, no change Annual DXA test	To prevent ONJ, dental examination and appropriate treatment before BP Start the current standards of care for cancer bone pain management at the onset of pain, in concert with the initiation of the bone-modifying agent therapy Adequate lifestyle modifications (smoking, alcohol, exercise) Calcium and vitamin D supplements
International panel [189]	Postmenopausal patients treated with AI Patients with metastatic breast-cancer	Prevention of bone loss Patients with metastatic breast-cancer Patients who cannot attend regular hospital care	BP therapy with AI IV pamidronate over 2 h every 3–4 weeks IV ibandronate over 15 min every 3–4 weeks IV zoledronic acid over 15 min every 3–4 weeks Oral ibandronate or clodronate	Serum creatinine monitored before each dose of BP Dose of BP adapted to renal function Subsequent monitoring for zoledronic acid and pamidronate Treatment for 2 years even in case of bone event. Beyond 2 years, individual risk assessment BMT test for people with risk factors: T-score < -1.5, age >65 years, corticosteroids for more than 6 months, family history of hip fracture, or personal history of fragility fracture after 50 BMD test	To prevent ONJ, dental examination and appropriate treatment before BP Pain control, analgesic therapy and BP by itself is a major factor of quality of life To avoid renal toxicity with IV BP, hydration before treatment Calcium and vitamin D supplements In case of oral administration, instruction to avoid GI problems
International Society of Geriatric Oncology (SIOG) [190]	Adjuvant setting Elderly patients with bone metastases	Patients receiving AI Prevention of SREs	Preventive BP, and especially IV zoledronic acid IV pamidronate 60 mg/h at max IV ibandronate, for creatinine clearance <30 mL/min, 2 m of ibandronate every 3–4 weeks (1-h infusion) IV zoledronic acid 4 mg, with dose adjustments for patients with mild to severe renal impairment and not	Creatinine clearance instead of serum creatinine should be monitored before each dose of BP Less renally toxic agent should be used	To prevent ONJ, dental examination and appropriate treatment before BP Zoledronic acid, ibandronate and pamidronate useful in the management of pain Optimization of the hydration status

Table 3 (continued)

Resources	Target population	Indication	Type of treatment, concentration, duration	Monitoring	Other recommendations
UK guidance document [101, 191]	Patients with advanced breast cancer	<p>Patients newly diagnosed with bone metastases</p> <p>Patients with pain</p> <p>Patients with long bone involvement and significant cortical destruction</p> <p>Patients with or at risk of spinal cord compression</p>	<p>recommended in patients with severe renal impairment (<30 mL/min)</p> <p>Oral clodronate 800 mg: creatinine clearance between 10 and 30 mL/min</p> <p>Oral ibandronate 50 mg weekly: creatinine clearance < 30 mL/min</p> <p>BPs</p> <p>Single fraction radiotherapy (8 Gy)</p> <p>Assessment by an orthopaedic surgeon, surgery intervention to prevent fracture</p> <p>Assessment by a spinal specialized team, surgery, corticosteroids, radiotherapy</p>	<p>BMD (DXA) test at spine/hip within 3 months of start of the treatment and then DXA after 1 year</p>	<p>In case of oral administration, instruction to avoid GI problems</p> <p>Supportive care</p>
	Post-menopausal patients treated with AI for breast cancer	<p>Treatment depends on baseline BMD (T-score < -2.0) or occurrence of a vertebral fragility fracture or previous low trauma hip fracture or an annual bone loss $\geq 4\%$ or age >75 years and ≥ 1 clinical risk factors</p>	<p>Oral alendronate 70 mg weekly or risedronate 35 mg, oral ibandronate 150 mg monthly, IV ibandronate 3 mg 3-monthly or IV zoledronic acid 4 mg 6-monthly</p>	<p>BMD (DXA) test at spine/hip within 3 months of start of the treatment and then DXA after 1 year</p>	<p>Calcium and vitamin D supplements</p> <p>Lifestyle advice</p>
	Patients who experience premature menopause due to chemotherapy, or ovarian suppression, ablation or removal	<p>Patients with DXA T-score < -1.0 or vertebral fragility fracture or previous low trauma hip fracture</p>	<p>Oral alendronate 70 mg weekly or risedronate 35 mg, oral ibandronate 150 mg monthly, IV ibandronate 3 mg 3-monthly or IV zoledronic acid 4 mg 6-monthly</p>	<p>BMD (DXA) test at spine/hip within 3 months of start of the treatment and then DXA after 1 year</p> <p>Monitoring thereafter depends on baseline BMD</p>	<p>Calcium and vitamin D supplements</p> <p>Lifestyle advice</p>
Belgian Bone Club [103]	Patients treated with AI for breast cancer	<p>Patients with a T-score < -2.5 or history fragility fracture or a T-score between -1.0 and -2.5 with additional risk factors</p>	<p>IV zoledronic acid 4 mg every 6 months</p> <p>Other bisphosphonates may be considered</p>	<p>At start of treatment, BMD test by DXA and assessment of risk factors for osteoporotic fracture</p> <p>BMD monitoring every 1–2 years of osteopenic, osteoporotic patients</p>	<p>Adequate lifestyle modifications (smoking, alcohol, exercise)</p> <p>Calcium and vitamin D supplements</p>
Practical guidance [104, 187]	Patients treated with AI for breast cancer	<p>For patients with 2 risk factors (T-score < -1.5, age >65 years, low BMI, family</p>	<p>IV zoledronic acid 4 mg every 6 months for at least 2 years and possibly as long as AI treatment</p>	<p>Monitoring of BMD every 2 years</p> <p>For patients with a T-score ≥ -2.0 and no additional</p>	<p>Yo prevent ONJ, dental examination and appropriate treatment before BP</p>

Table 3 (continued)

Resources	Target population	Indication	Type of treatment, concentration, duration	Monitoring	Other recommendations
ESCEO [15]	Women treated with AI for breast cancer	history of hip fracture, corticosteroids, smoking) Osteoporotic women (T-score hip/spine < -2.5, or ≥1 prevalent fracture) Women aged ≥75 irrespective of BMD Patients with T-score < -1.5 + ≥1 clinical risk or T-score < -1.0 + ≥2 clinical risk factors	Oral risedronate 35 mg/week (limited efficacy data) sc denosumab 60 mg every 6 months (limited efficacy data) IV zoledronic acid 4 mg every 6 months sc denosumab Oral BPs For the entire period of AI: BP	risk factors, BMD monitoring every 1–2 years PTH, calcium, 25-hydroxyvitamin D to be tested before start of AI treatment Baseline risk of osteoporotic fracture by DXA and evaluation of clinical risk factors	Adequate lifestyle modifications (smoking, alcohol, exercise) Calcium and vitamin D supplements Adequate lifestyle modifications (smoking, alcohol, exercise) Calcium and vitamin D supplements

AI aromatase inhibitor, BMD bone mineral density, BP bisphosphonate, DXA dual X-ray absorptiometry, ER+ oestrogen receptor positive, GI gastrointestinal, IV intravenous, ONJ osteonecrosis of the jaw, sc subcutaneous, SRE skeletal-related event

regimen significantly improved lumbar spine BMD (+5.2 %) and total hip BMD (+3.5 %) versus the delayed group. N-telopeptide and ALP dropped by 21.3 and 12.8 % in the upfront group and increased by 21.7 and 24.9 % in the delayed group. No differences were found between the two groups in terms of fractures, although the study was not designed to test this endpoint. Disease recurrence was less frequent in the upfront group and no case of osteonecrosis of the jaw was reported. Encouraging data were also obtained in premenopausal women in the ABCSG where zoledronic acid prevented bone loss in a subgroup of premenopausal cancer patients [50]. These and other studies are presented in Table 3.

Prostate cancer Alendronate, risedronate, zoledronic acid, denosumab and teriparatide are approved for the treatment/prevention of osteoporotic fractures in men with osteoporosis with some limitations according to regional policies. The use of teriparatide is not recommended in cancer-treated patients. Available RCTs in the setting of ADT and prostate cancer have shown efficacy in the prevention of bone loss, a significant increase in BMD and normalization of bone turnover in the dosages normally recommended for idiopathic osteoporosis [167, 168]. Pamidronate (60 mg IV 12-weekly) in a small prospective study in men with non-metastatic prostate cancer significantly increased BMD at the spine and hip [83]. Zoledronic acid (4 mg IV as single dose) in a RCT of ADT and non-metastatic prostate cancer significantly increased BMD at spine and hip, when compared to placebo [169]. There is presently no published trial of the effect of a bisphosphonate on fracture risk in the ADT/prostate cancer setting. These and other studies are presented in Table 5.

Denosumab

Breast cancer The efficacy of denosumab was evaluated in 252 patients receiving adjuvant AI in the HALT-BC trial. Patients randomly received either placebo or denosumab (60 mg) every 6 months. In the denosumab group, BMD in all examined skeletal sites was increased already after 1 month of treatment, and at 12 and 24 months was significantly higher than placebo [170, 171]. At 24 months, the difference between both treatment groups was 7.6 % at the lumbar spine and 4.7 % at the total hip. At the lumbar spine, BMD increased by more than 3 % in 80 % of denosumab-treated patients compared with 10 % in the placebo group. An increase in BMD with denosumab was observed at lumbar spine after 1 year (+5.5 %) and at radius after 2 years (+6.1 %). No serious adverse events were attributed to denosumab in the trial. An ongoing clinical trial is investigating the anti-fracture efficacy of denosumab in women with breast cancer receiving AI in an adjuvant setting [172].

Prostate cancer In the HALT Prostate trial, 1,468 men receiving ADT for prostate cancer and being at high risk for fracture (history of osteoporotic fracture, age ≥ 70 years or low BMD) were randomized to either denosumab (60 mg subcutaneously 6-monthly) or placebo for 3 years. At 2 years, BMD of the lumbar spine (the primary endpoint) increased by 5.6 % in the denosumab group as compared with a decrease of 1.0 % in the placebo group ($p < 0.001$). Denosumab increased BMD at various skeletal sites (femoral neck, total hip, distal radius) and reduced the 3-year incidence of new vertebral fractures by 62 % (1.5 vs 3.9 % in the placebo group; HR, 0.38; 95 % CI, 0.19–0.78; $p = 0.006$), fractures at any site by 72 % ($p = 0.10$) and multiple fractures at any site by 72 % ($p = 0.006$). In a further post hoc analysis, a trend was found toward a positive effect on non-vertebral fractures [173].

Selective oestrogen receptor modulators

In a RCT of 646 men with prostate cancer and ADT, subjects randomly received the SERM toremifene (80 mg oral daily) or placebo for 2 years. Toremifene significantly reduced the risk of new vertebral fractures by 50 % ($p = 0.05$), increased BMD at the spine and hip, and decreased markers of bone turnover [174]. Raloxifene improved BMD at the hip and tended to improve BMD at the spine in another small open-label study, but has not been studied with fracture endpoints [175].

Official recommendations and comparison of guidelines

Metastatic bone disease

Tables 3 (breast), 4 (multiple myeloma) and 5 (prostate) summarize the various national and international recommendations for the management of bone disease in cancer patients and the suggested approaches to bone assessment and follow-up. The 2003 recommendations from the ASCO for the prevention and treatment SREs in patients with metastatic breast cancer [102] were updated in 2011 [154] (Table 3). The 2011 differ from the 2003 ones in the addition of denosumab to intravenous bisphosphonates. Bone-modifying agent therapy is only recommended for patients with breast cancer with evidence of bone metastases; denosumab 120 mg subcutaneously every 4 weeks, intravenous pamidronate 90 mg over no less than 2 h or zoledronic acid 4 mg over no less than 15 min every 3 to 4 weeks is recommended. ASCO also issued clinical practice guidelines to manage lytic bone disease or compression fractures in patients with multiple myeloma with intravenous bisphosphonates [176, 177]. IV zoledronic acid or pamidronate is recommended for preventing SREs in patients with multiple myeloma. Zoledronic acid is preferred over oral clodronate in newly diagnosed patients with multiple myeloma because of its potential anti-myeloma effects and survival benefits. Bisphosphonates should be

administered every 3 to 4 weeks IV during initial therapy. Zoledronic acid or pamidronate should be continued in patients with active disease and should be resumed after disease relapse, if discontinued in patients achieving complete or very good partial response (Table 4).

Although several studies suggest the potential usefulness of bone-modifying agents in preventing extension in other parenchymas, including skeletal metastases, in patients without metastatic bone disease at study entry, the evidence is not conclusive. The Cancer Care Ontario group has also issued recommendations to use zoledronic acid in adult patients with renal carcinoma and bone metastases [133].

Treatment-induced bone loss

The ASCO Guidelines Update committee on adjuvant endocrine therapy for women with ER+ breast cancer recommends that postmenopausal women with hormone receptor-positive breast cancer consider incorporating an aromatase inhibitor therapy at some point during adjuvant treatment, either as upfront therapy or as sequential treatment after tamoxifen. The optimal timing and duration of aromatase inhibition remain unresolved. The ASCO panel supports careful consideration of side effect profiles and patient preferences in deciding whether and when to incorporate AI therapy (Table 3). Risk stratification based on BMD T-score and clinical risk factors has been the recommended approach by several organizations to identify patients who most benefit from inhibitors of bone resorption. The issue of use of bone-modifying agents in the management of adjuvant-associated bone loss in patients with breast cancer is to be covered by ASCO in a separate guideline update that is eagerly awaited [154, 178].

General recommendations

Cancer without known skeletal metastases and not requiring therapy to lower sex steroid

Risk assessment should be applied as in non-cancer patients [179–182]. A detailed history and a focused physical examination are recommended to identify risk factors for low BMD, falls and fractures, as well as undiagnosed vertebral fractures. Hip and spine BMD should be measured with DXA according to the local guidelines for DXA use in non-cancer patients. FRAX should be calculated using femoral neck BMD and pharmacotherapy introduced according to guidelines for non-cancer patients.

For postmenopausal women or men over age 50 with densitometric osteoporosis (T-score ≤ -2.5 at the total hip, femoral neck or lumbar spine) or prevalent fragility fracture, laboratory investigations to rule out secondary causes of bone loss are recommended (blood cell count, calcium, phosphate, alkaline phosphatase, TSH, 25-hydroxyvitamin D, creatinine,

Table 4 Guidelines for bone management in multiple myeloma

Resources	Target population	Indication	Type of treatments, concentration, duration	Monitoring	Other recommendations
American Society of Clinical Oncology (ASCO) [176, 177]	Multiple myeloma (MM) patients	For MM patients with lytic destruction of bone or spine compression fracture from osteopenia on plain radiographs or imaging and for MM patients with osteopenia, but no radiographic evidence of lytic bone disease	IV pamidronate 90 mg over 2 h every 3–4 weeks for 2 years IV zoledronic acid 4 mg over 15 min every 3–4 weeks for 2 years Oral clodronate	Serum creatinine before each dose of pamidronate or Zoledronic acid Regular check of serum calcium, electrolytes, phosphate magnesium, haematocrit/haemoglobin Intermittent evaluation (every 3–6 months) of albuminuria in patients who develop renal deterioration with no apparent cause during BP therapy should stop it Renal function should be carefully monitored and doses reduced according to manufacturers' guidance	For patients with pre-existing renal impairment, reduced dosage of zoledronic acid and initial pamidronate dose Zoledronic acid is not recommended in patients with severe renal impairment IV pamidronate or zoledronic acid for patients with pain caused by osteolytic disease as an adjuvant therapy To prevent ONJ, dental examination and appropriate treatment before BP To prevent ONJ, dental examination and appropriate treatment before BP Oral calcium and vitamin D supplements with zoledronic acid
Haematology-oncology Task Force of the British Committee for Standards In Haematology and UK Myeloma Forum [192]	MM patients	For symptomatic MM patients with evident bone lesions or not For MM patients with fracture	IV pamidronate IV zoledronic acid Oral clodronate Stabilization of long bone fractures and subsequent radiotherapy (8 Gy single fraction). Local radiation of 8 Gy single fraction for pain control IV (or oral) pamidronate over 2–4 h for 2 years IV zoledronic acid Oral clodronate	creatinine clearance	To prevent ONJ, dental examination and appropriate treatment before BP Patients with compromised renal function should have creatinine clearance rates, serum electrolytes, albuminuria BP useful in the management of pain To prevent ONJ, dental examination and appropriate treatment before BP
European Myeloma Network [193]	MM patients	For MM patients with lytic bone disease or severe osteoporosis	IV pamidronate monthly for 2 years IV zoledronic acid for monthly for 2 years IV pamidronate IV zoledronic acid Radiotherapy (10–30 Gy) as a palliative treatment for uncontrolled pain Orthopaedic consultation Vertebroplasty or kyphoplasty IV or oral long-term administration of bisphosphonates	Pamidronate is favoured over zoledronic acid Renal function	To prevent ONJ, dental examination and appropriate treatment before BP To prevent ONJ, dental examination and appropriate treatment before BP
Mayo Clinic Consensus Statement [194]	MM patients	For MM patients with lytic disease evident on plain radiographs or osteopenia/osteoporosis	IV pamidronate monthly for 2 years IV zoledronic acid for monthly for 2 years	creatinine clearance	To prevent ONJ, dental examination and appropriate treatment before BP Patients with compromised renal function should have creatinine clearance rates, serum electrolytes, albuminuria BP useful in the management of pain To prevent ONJ, dental examination and appropriate treatment before BP
National Comprehensive Cancer Network (NCCN) [195]	MM patients	For MM patients receiving primary myeloma therapy For MM patients with long-bone fractures or bony compression of spinal cord For MM patients with symptomatic vertebral compression fractures Patients with stage III or relapsed disease receiving conventional dose therapy	IV pamidronate IV zoledronic acid Radiotherapy (10–30 Gy) as a palliative treatment for uncontrolled pain Orthopaedic consultation Vertebroplasty or kyphoplasty IV or oral long-term administration of bisphosphonates	Renal function	To prevent ONJ, dental examination and appropriate treatment before BP To prevent ONJ, dental examination and appropriate treatment before BP
ESMO [196]	MM patients	Prevention of SREs	IV pamidronate 60 mg/h at max. IV zoledronic acid 4 mg, with dose adjustments for patients with mild to severe renal impairment and not recommended in patients with severe renal impairment (<30 mL/min) IV ibandronate oral clodronate 800 mg: creatinine clearance between 10 and 30 mL/min	Creatinine clearance instead of serum creatinine should be monitored in each patient before each dose Less renally toxic agent should be used	To prevent ONJ, dental examination and appropriate treatment before BP BP useful in the management of pain Optimization of the hydration status In case of oral administration, instruction to avoid GI problems
International Society of Geriatric Oncology (SIOG) [190]	Elderly patients with MM	Prevention of SREs	IV pamidronate 60 mg/h at max. IV zoledronic acid 4 mg, with dose adjustments for patients with mild to severe renal impairment and not recommended in patients with severe renal impairment (<30 mL/min) IV ibandronate oral clodronate 800 mg: creatinine clearance between 10 and 30 mL/min	Creatinine clearance instead of serum creatinine should be monitored in each patient before each dose Less renally toxic agent should be used	To prevent ONJ, dental examination and appropriate treatment before BP BP useful in the management of pain Optimization of the hydration status In case of oral administration, instruction to avoid GI problems

BMD bone mineral density, *BP* bisphosphonate, *DXA* dual X-ray absorptiometry, *GI* gastrointestinal, *IV* intravenous, *MM* multiple myeloma, *ONJ* osteonecrosis of the jaw, *sc* subcutaneous, *SRE* skeletal-related event

Table 5 Guidelines for bone management in prostate cancer

Resources	Target population	Indication	Type of treatments, concentration, duration	Monitoring	Other recommendations
International panel [189]	Patients with prostate cancer	Patients with hormone refractory prostate cancer metastatic to bone	IV zoledronic acid 4 mg every 3 weeks for 15 min	Serum creatinine before each dose Dose of BP adapted to renal function Subsequent monitoring for zoledronic acid Treatment for 2 years even in case of bone event; beyond 2 years, individual risk assessment	To prevent ONJ, dental examination and appropriate treatment before BP Pain control, analgesic therapy and BP by itself is a major factor of quality of life To avoid renal toxicity with iv BP, hydration before treatment Calcium and vitamin D In case of oral administration, instruction to avoid GI problems
International Society of Geriatric Oncology (SIOG) [190]	Elderly patients with prostate cancer	Prevention of SREs	IV zoledronic acid 4 mg with dose adjustments for patients with mild to severe renal impairment and not recommended in patients with severe renal impairment (<30 mL/min)	Creatinine clearance instead of serum creatinine monitored in each patient before each dose Less renally toxic agent should be used	To prevent ONJ, dental examination and appropriate treatment before BP BP useful in the management of pain Optimization of the hydration status
Belgian Bone Club [103]	Patients with prostate cancer	Patients treated with ADT and T-score < -2.5 or history fragility fracture or a T-score between -1.0 and -2.5 with also risk factors	IV zoledronic acid 4 mg once a year Oral alendronate, weekly	At start of treatment, BMD test by DXA and assessment of risk factors for osteoporotic fracture BMD monitoring every 1–2 years of osteopenic, osteoporotic patients	Adequate lifestyle modifications (smoking, alcohol, exercise) Calcium and vitamin D supplements
ESA, ANZBMS and USANZ [197]	Patients with non-metastatic prostate cancer	Patients receiving ADT	BPs (pamidronate, zoledronic acid, alendronate, risedronate)	DXA and thoracolumbar spine DXA in case of osteopenia Monitoring of metabolic health	Adequate lifestyle modifications (smoking, alcohol, exercise) Calcium and vitamin D supplements
Expert panel [198]	Patients with prostate cancer	Patients beginning or receiving ADT with T-score < -2.5	IV zoledronic acid 4 mg 15-min infusion every 3–4 weeks	DXA test at baseline and repeat every 12 months	Adequate lifestyle modifications (smoking, alcohol, exercise) Calcium and vitamin D supplements

ADT androgen deprivation therapy, BMD bone mineral density, BP bisphosphonate, DXA dual X-ray absorptiometry, GI gastrointestinal, IV intravenous, ONJ osteonecrosis of the jaw, sc subcutaneous, SRE skeletal-related event

serum protein electrophoresis) as in idiopathic osteoporosis. The use of bone turnover markers to improve fracture risk assessment or to monitor therapy remains controversial. However, in patients with skeletal metastatic disease, elevated markers of bone resorption appear to be associated with poorer prognosis and increased mortality [183].

Lateral thoracic and lumbar spine radiography, or VFA by DXA, if clinical evidence is suggestive of a vertebral fracture, should be performed. Vertebral fractures are defined as deformities of vertebrae with reduction greater than 20 % of vertebral dimension. Thirty percent of vertebral fractures are asymptomatic and therefore imaging in patients at risk should be encouraged. Such fractures are highly predictive of future fracture and have a marked influence on FRAX outcome.

Non-metastatic cancer treated with endocrine therapy

All of the above can be applied to this group of patients. Women who are taking AI and men who are undergoing ADT should be assessed for fracture risk, and osteoporosis therapy to prevent fractures should be considered (Figs. 1 and 2).

Follow-up

Baseline status, osteoporosis and bone fracture risk factors, and the type of cancer therapy used influence the frequency, profile and duration of examinations included in the follow-up of these patients [76, 101, 104].

Most of the cancer treatment-induced bone loss is explained by the hypogonadal state induced by the therapies.

Thus, the follow-up currently focusses on the measurement of BMD and perhaps of bone turnover markers (BTMs) [101]. The goal of therapy is the maintenance (with some agents) or the increase (with other agents) in BMD. BTMs may respond with increases, with some therapies and decreases with other therapies. Measuring BMD or BTMs may increase adherence to therapy in individual patients.

A general approach and practical algorithm is presented in Figs. 1 and 2, which also emphasize the important role played by osteoporosis risk factors in the follow-up and management of these patients. Other organizations such as ESCEO, ASCO and the UK Expert Group have also issued other algorithms that are also anchored on BMD T-scores cut-offs and risk factors [15, 101, 102]. Indications and limitations can be region-specific according to the approval or insurance reimbursement policies.

Breast cancer

BTMs may be used to monitor response to antiresorptive therapy. The most commonly used specific markers are serum cross-linked terminal telopeptide (CTX) on a serum sample collected between 7 and 10 h in the morning in fasting state and urinary NTX expressed as a ratio to creatinine and measured on a second morning void urine sample. With the use of bisphosphonates as antiresorptive therapy, a 50–70 % reduction of CTX and NTX is expected in the first 3 months of treatment, a plateau is observed thereafter. It is generally accepted that the goal is to reduce bone resorption by more than the least significant change, keeping the bone resorption markers into the lower half of the reference range for healthy young women [101, 184]. Currently, a reduction in

Fig. 1 Management of patients with breast cancer treated with aromatase inhibitors (AIs). Adapted from [15]





Fig. 2 Management of patients with non-metastatic prostate cancer on androgen deprivation therapy (ADT)



bone resorption greater than 50 % indicates that the least significant change has been achieved. It has been shown that concomitant diseases and recent fractures can influence BTM levels, thus caution needs to be taken when interpreting the results [185].

BMD value and the number of risk factors of each individual patient should be known before the initiation of any kind of bone-sparing therapy in all cancer treatment-associated bone loss (Figs. 1 and 2). Conversely, as the increases in BMD on treatment are small, BMD should be measured preferably at lumbar spine where the least significant change is around 3 %. BMD should be measured every 18–24 months [101, 186] in patients treated with bone resorption inhibitors.

Elderly women (>65 years) and patients with T-score < -2.0 and at least one more additional risk factor should receive bone protection with bisphosphonates irrespective of BMD if they are receiving AI therapy (Fig. 1).

Different management [101] is proposed in cancer therapy-induced premature menopause compared with biologically occurring menopause. In treatment-induced menopause or in programmed ovarian suppression before the age of 45 years, the evaluation of BMD should be indicated in two sites (spine and hip) and BMD should be repeated 12 months [187] after post-chemotherapy amenorrhea occurrence. The frequency of further monitoring would depend on the baseline BMD and the type of treatment. When AIs are concomitantly used, monitoring (every 18–24 months) and therapeutic intervention with calcium, vitamin D supplements and bisphosphonates are recommended [101].

Patients with a T-score between -1 and -2 should be considered “medium risk”. However, patients with medium risk with a decline of BMD >4 % per year (either site) should initiate bone protection therapy. Finally if BMD is ≥ 1 , the 10-year risk for osteoporosis is very low and only lifestyle measures should be prescribed with re-evaluation in 24 months.

Prostate cancer

Bisphosphonates have been shown to attenuate bone loss, but optimal regimen (IV vs oral route of administration, every 6 months vs every 12 months) and the long-term effect on fracture risk remain to be established. Therefore, there is a need for bone monitoring measuring BMD with DXA. Denosumab has also been shown to increase bone mass in ADT patients who are at high risk of fracture and decrease BTMs. BMD and BTMs should be followed during treatment [84].

The frequency of monitoring will depend on the baseline DXA, the presence of osteoporosis risk factors and the use of bisphosphonates. Differently to breast cancer, high, medium and low-risk groups have been clearly established in prostate cancer, but a similar approach to breast cancer for monitoring is currently adopted [188].

If T-score is ≤ -2.5 patients should be monitored and managed as high-risk patients. If T-score is between -1 and -2.5, a more frequent monitoring (12 months) is advised. Patients with a T-score ≥ -1 can be considered at lower risk and BMD studies

repeated every 18–24 months [101, 186]. If bisphosphonates are used in the frame of ADT, BMD should be measured every 18 to 24 months (Fig. 2).

Conclusions

Bone is often affected in the course of cancer either as bone metastases or as bone loss and fragility resulting from anti-cancer therapies. These various disease complications are associated with an important morbidity and with a largely compromised quality of life. To preserve bone and reduce this morbidity, efficacious therapies are available, with a highly favourable risk–benefits ratio. Expert reports from various societies are providing guidance for the identification of patients at risk of bone disease, their management and follow-up.

Acknowledgments The authors thank Ms Aida Farha, Medical Information Specialist, Saab Medical Library, American University of Beirut, for her assistance in designing comprehensive and complex searches of the various medical literature resources.

Conflicts of interest René Rizzoli has received speaker or advisory board fees from Amgen, MSD, GSK, Servier, Danone and Takeda. Jean-Jacques Body has received speaker and consulting fees from Amgen and Novartis. Jorge B. Cannata-Andía has received research grants and speaker or advisory board fees from Amgen, Abbott, Servier and Shire. David Kendler has received honoraria for speaking, consulting and/or research grants from Amgen, Novartis, GSK, Eli Lilly, Merck, Johnson&Johnson, Pfizer and Roche. Alexandra Papaioannou has been a consultant, or on a speaker's bureau, or received unrestricted grants from Amgen, Eli Lilly, Merck Canada Inc., Novartis, Pfizer and Warner Chilcott. Tobias J de Villiers has received speaker or advisory board fees from Amgen, Merck, Adcock Ingram and Pfizer. Catherine Van Poznak has received research support from Amgen and Novartis. Ghada El-Hajj Fuleihan has received grants to support courses on osteoporosis from Novartis Pharmaceuticals and Les Laboratoires Servier. Abdellah El Maghraoui, Nicola Napoli, Dominique D Pierroz and Maria Rahme have no conflict of interest.

References

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D (2011) Global cancer statistics. *CA Cancer J Clin* 61:69–90
- WHO (2011) Global status report on noncommunicable diseases 2010. Description of the global burden of NCDs, their risk factors and determinants. WHO, Geneva
- Weigelt B, Peterse JL, van 't Veer LJ (2005) Breast cancer metastasis: markers and models. *Nat Rev Cancer* 5:591–602
- Coleman RE (2006) Clinical features of metastatic bone disease and risk of skeletal morbidity. *Clin Cancer Res* 12:6243s–6249s
- Coleman RE, Rubens RD (1987) The clinical course of bone metastases from breast cancer. *Br J Cancer* 55:61–66
- Mundy GR (2002) Metastasis to bone: causes, consequences and therapeutic opportunities. *Nat Rev Cancer* 2:584–593
- Coleman RE (2001) Metastatic bone disease: clinical features, pathophysiology and treatment strategies. *Cancer Treat Rev* 27:165–176
- Cazzaniga M, Pronzato P, Leto di Priolo SL, De Matteis A, Di Costanzo F, Passalacqua R, Rosso R, Torri V (2004) Patterns of relapse and modalities of treatment of breast cancer: the 'IRIS' Project, a multicenter observational study. *Oncology* 66:260–268
- Roodman GD (2004) Mechanisms of bone metastasis. *N Engl J Med* 350:1655–1664
- Reuss-Borst M, Hartmann U, Scheede C, Weiss J (2012) Prevalence of osteoporosis among cancer patients in Germany: prospective data from an oncological rehabilitation clinic. *Osteoporos Int* 23:1437–1444
- Chen Z, Maricic M, Bassford TL, Pettinger M, Ritenbaugh C, Lopez AM, Barad DH, Gass M, Leboff MS (2005) Fracture risk among breast cancer survivors: results from the Women's Health Initiative Observational Study. *Arch Intern Med* 165:552–558
- Kanis JA, McCloskey EV, Powles T, Paterson AH, Ashley S, Spector T (1999) A high incidence of vertebral fracture in women with breast cancer. *Br J Cancer* 79:1179–1181
- Eastell R (2007) Aromatase inhibitors and bone. *J Steroid Biochem Mol Biol* 106:157–161
- McCloskey E (2006) Effects of third-generation aromatase inhibitors on bone. *Eur J Cancer* 42:1044–1051
- Rizzoli R, Body JJ, De Censi A, Reginster JY, Piscitelli P, Brandi ML, European Society for Clinical Economical Aspects of Osteoporosis and Osteoarthritis (ESCEO) (2012) Guidance for the prevention of bone loss and fractures in postmenopausal women treated with aromatase inhibitors for breast cancer: an ESCEO position paper. *Osteoporos Int* 23:2567–2576
- Eastell R, Adams JE, Coleman RE, Howell A, Hannon RA, Cuzick J, Mackey JR, Beckmann MW, Clack G (2008) Effect of anastrozole on bone mineral density: 5-year results from the anastrozole, tamoxifen, alone or in combination trial 18233230. *J Clin Oncol* 26:1051–1057
- Morote J, Morin JP, Orsola A, Abascal JM, Salvador C, Trilla E, Raventos CX, Cecchini L, Encabo G, Reventos J (2007) Prevalence of osteoporosis during long-term androgen deprivation therapy in patients with prostate cancer. *Urology* 69:500–504
- Shahinian VB, Kuo YF, Freeman JL, Goodwin JS (2005) Risk of fracture after androgen deprivation for prostate cancer. *N Engl J Med* 352:154–164
- Morote J, Orsola A, Abascal JM, Planas J, Trilla E, Raventos CX, Cecchini L, Encabo G, Reventos J (2006) Bone mineral density changes in patients with prostate cancer during the first 2 years of androgen suppression. *J Urol* 175:1679–1683, discussion 1683
- Roodman GD (2009) Pathogenesis of myeloma bone disease. *Leukemia* 23:435–441
- Guise TA, Mundy GR (1998) Cancer and bone. *Endocr Rev* 19:18–54
- Chappard D, Bouvard B, Basle MF, Legrand E, Audran M (2011) Bone metastasis: histological changes and pathophysiological mechanisms in osteolytic or osteosclerotic localizations. A review. *Morphologie* 95:65–75
- Clézardin P, Teti A (2007) Bone metastasis: pathogenesis and therapeutic implications. *Clin Exp Metastasis* 24:599–608
- Guise TA (2000) Molecular mechanisms of osteolytic bone metastases. *Cancer* 88:2892–2898
- Pugsley MK, Tabrizchi R (2000) The vascular system. An overview of structure and function. *J Pharmacol Toxicol Methods* 44:333–340
- Chappard D, Libouban H, Legrand E, Ifrah N, Masson C, Basle MF, Audran M (2010) Computed microtomography of bone specimens for rapid analysis of bone changes associated with malignancy. *Anat Rec (Hoboken)* 293:1125–1133
- Powell GJ, Southby J, Danks JA, Stillwell RG, Hayman JA, Henderson MA, Bennett RC, Martin TJ (1991) Localization of parathyroid hormone-related protein in breast cancer metastases: increased incidence in bone compared with other sites. *Cancer Res* 51:3059–3061
- Jones DH, Nakashima T, Sanchez OH et al (2006) Regulation of cancer cell migration and bone metastasis by RANKL. *Nat Geosci* 440:692–696

29. Leeming DJ, Dellings G, Koizumi M, Henriksen K, Karsdal MA, Li B, Qvist P, Tanko LB, Byrjalsen I (2006) Alpha CTX as a biomarker of skeletal invasion of breast cancer: immunolocalization and the load dependency of urinary excretion. *Cancer Epidemiol Biomarkers Prev* 15:1392–1395
30. Granchi S, Brocchi S, Bonaccorsi L, Baldi E, Vinci MC, Forti G, Serio M, Maggi M (2001) Endothelin-1 production by prostate cancer cell lines is up-regulated by factors involved in cancer progression and down-regulated by androgens. *Prostate* 49:267–277
31. Takuwa Y, Masaki T, Yamashita K (1990) The effects of the endothelin family peptides on cultured osteoblastic cells from rat calvariae. *Biochem Biophys Res Commun* 170:998–1005
32. Lu Y, Zhang J, Dai J, Dehne LA, Mizokami A, Yao Z, Keller ET (2004) Osteoblasts induce prostate cancer proliferation and PSA expression through interleukin-6-mediated activation of the androgen receptor. *Clin Exp Metastasis* 21:399–408
33. Nakashima J, Tachibana M, Horiguchi Y, Oya M, Ohigashi T, Asakura H, Murai M (2000) Serum interleukin 6 as a prognostic factor in patients with prostate cancer. *Clin Cancer Res* 6:2702–2706
34. Shariat SF, Andrews B, Kattan MW, Kim J, Wheeler TM, Slawin KM (2001) Plasma levels of interleukin-6 and its soluble receptor are associated with prostate cancer progression and metastasis. *Urology* 58:1008–1015
35. Bataille R, Chappard D, Basle M (1995) Excessive bone resorption in human plasmacytomas: direct induction by tumour cells in vivo. *Br J Haematol* 90:721–724
36. Bataille R, Chappard D, Klein B (1992) Mechanisms of bone lesions in multiple myeloma. *Hematol Oncol Clin North Am* 6:285–295
37. Colla S, Zhan F, Xiong W, Wu X, Xu H, Stephens O, Yaccoby S, Epstein J, Barlogie B, Shaughnessy JD Jr (2007) The oxidative stress response regulates DKK1 expression through the JNK signaling cascade in multiple myeloma plasma cells. *Blood* 109:4470–4477
38. Tian E, Zhan F, Walker R, Rasmussen E, Ma Y, Barlogie B, Shaughnessy JD Jr (2003) The role of the Wnt-signaling antagonist DKK1 in the development of osteolytic lesions in multiple myeloma. *N Engl J Med* 349:2483–2494
39. Barretina J, Junca J, Llano A, Gutierrez A, Flores A, Blanco J, Clotet B, Este JA (2003) CXCR4 and SDF-1 expression in B-cell chronic lymphocytic leukemia and stage of the disease. *Ann Hematol* 82:500–505
40. Josselin N, Libouban H, Dib M, Ifrah N, Legrand E, Basle MF, Audran M, Chappard D (2009) Quantification of dendritic cells and osteoclasts in the bone marrow of patients with monoclonal gammopathy. *Pathol Oncol Res* 15:65–72
41. Marcelli C, Chappard D, Rossi JF, Jaubert J, Alexandre C, Dessauw P, Baldet P, Bataille R (1988) Histologic evidence of an abnormal bone remodeling in B-cell malignancies other than multiple myeloma. *Cancer* 62:1163–1170
42. Coleman RE, Lipton A, Roodman GD et al (2010) Metastasis and bone loss: advancing treatment and prevention. *Cancer Treat Rev* 36:615–620
43. Stava CJ, Jimenez C, Hu MI, Vassilopoulou-Sellin R (2009) Skeletal sequelae of cancer and cancer treatment. *J Cancer Surviv* 3:75–88
44. VanderWalde A, Hurria A (2011) Aging and osteoporosis in breast and prostate cancer. *CA Cancer J Clin* 61:139–156
45. Michaud LB (2010) Managing cancer treatment-induced bone loss and osteoporosis in patients with breast or prostate cancer. *Am J Health Syst Pharm* 67:S20–30, quiz S31–23
46. Wadhwa VK, Parr NJ (2009) Peripheral or axial bone density measurements to identify osteoporosis in prostate cancer patients undergoing androgen deprivation therapy? *Urology* 73:1347–1351
47. Hadji P (2010) Guidelines for osteoprotection in breast cancer patients on an aromatase inhibitor. *Breast Care (Basel)* 5:290–296
48. Shapiro CL, Manola J, Leboff M (2001) Ovarian failure after adjuvant chemotherapy is associated with rapid bone loss in women with early-stage breast cancer. *J Clin Oncol* 19:3306–3311
49. Eastell R, Hannon RA, Cuzick J, Dowsett M, Clack G, Adams JE (2006) Effect of an aromatase inhibitor on bmd and bone turnover markers: 2-year results of the Anastrozole, Tamoxifen, Alone or in Combination (ATAC) trial (18233230). *J Bone Miner Res* 21:1215–1223
50. Gnant MF, Mlineritsch B, Luschin-Ebengreuth G et al (2007) Zoledronic acid prevents cancer treatment-induced bone loss in premenopausal women receiving adjuvant endocrine therapy for hormone-responsive breast cancer: a report from the Austrian Breast and Colorectal Cancer Study Group. *J Clin Oncol* 25:820–828
51. Hadji P (2009) Aromatase inhibitor-associated bone loss in breast cancer patients is distinct from postmenopausal osteoporosis. *Crit Rev Oncol Hematol* 69:73–82
52. Bines J, Oleske DM, Cobleigh MA (1996) Ovarian function in premenopausal women treated with adjuvant chemotherapy for breast cancer. *J Clin Oncol* 14:1718–1729
53. Meistrich ML (2009) Male gonadal toxicity. *Pediatr Blood Cancer* 53:261–266
54. Bokemeyer C, Berger CC, Kuczyk MA, Schmoll HJ (1996) Evaluation of long-term toxicity after chemotherapy for testicular cancer. *J Clin Oncol* 14:2923–2932
55. Weilbaecher KN (2000) Mechanisms of osteoporosis after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 6:165–174
56. Love RR, Barden HS, Mazess RB, Epstein S, Chappell RJ (1994) Effect of tamoxifen on lumbar spine bone mineral density in postmenopausal women after 5 years. *Arch Intern Med* 154:2585–2588
57. Love RR, Mazess RB, Barden HS, Epstein S, Newcomb PA, Jordan VC, Carbone PP, DeMets DL (1992) Effects of tamoxifen on bone mineral density in postmenopausal women with breast cancer. *N Engl J Med* 326:852–856
58. Hu MI, Gagel RF, Jimenez C (2007) Bone loss in patients with breast or prostate cancer. *Curr Osteoporos Rep* 5:170–178
59. Bouvard B, Hoppe E, Soulie P et al (2012) High prevalence of vertebral fractures in women with breast cancer starting aromatase inhibitor therapy. *Ann Oncol* 23:1151–1156
60. Ghazi M, Roux C (2009) Hormonal deprivation therapy-induced osteoporosis in postmenopausal women with breast cancer. *Best Pract Res Clin Rheumatol* 23:805–811
61. Reid DM (2009) Prevention of osteoporosis after breast cancer. *Maturitas* 64:4–8
62. Rozenberg S, Carly B, Liebens F, Antoine C (2009) Risks of osteoporosis associated with breast cancer treatment: the need to access to preventive treatment. *Maturitas* 64:1–3
63. Yamamoto DS, Viale PH (2009) Update on identifying and managing osteoporosis in women with breast cancer. *Clin J Oncol Nurs* 13:E18–29
64. McCloskey EV, Hannon RA, Lakner G, Fraser WD, Clack G, Miyamoto A, Finkelman RD, Eastell R (2007) Effects of third generation aromatase inhibitors on bone health and other safety parameters: results of an open, randomised, multi-centre study of letrozole, exemestane and anastrozole in healthy postmenopausal women. *Eur J Cancer* 43:2523–2531
65. Winer EP, Hudis C, Burstein HJ et al (2005) American Society of Clinical Oncology technology assessment on the use of aromatase inhibitors as adjuvant therapy for postmenopausal women with hormone receptor-positive breast cancer: status report 2004. *J Clin Oncol* 23:619–629
66. Cummings SR, Duong T, Kenyon E, Cauley JA, Whitehead M, Krueger KA, Multiple Outcomes of Raloxifene Evaluation T (2002) Serum estradiol level and risk of breast cancer during treatment with raloxifene. *JAMA* 287:216–220
67. Gibson K, O'Bryan CL (2008) Screening and management of osteoporosis in breast cancer patients on aromatase inhibitors. *J Oncol Pharm Pract* 14:139–145
68. Arimidex T, Alone or in Combination (ATAC) Trialists' Group, Forbes JF, Cuzick J, Buzzdar A, Howell A, Tobias JS, Baum M

- (2008) Effect of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: 100-month analysis of the ATAC trial. *Lancet Oncol* 9:45–53
69. Jakesz R, Jonat W, Gnani M et al (2005) Switching of postmenopausal women with endocrine-responsive early breast cancer to anastrozole after 2 years' adjuvant tamoxifen: combined results of ABCSG trial 8 and ARNO 95 trial. *Lancet* 366:455–462
 70. Coates AS, Keshaviah A, Thurlimann B et al (2007) Five years of letrozole compared with tamoxifen as initial adjuvant therapy for postmenopausal women with endocrine-responsive early breast cancer: update of study BIG 1-98. *J Clin Oncol* 25:486–492
 71. Perez EA, Josse RG, Pritchard KI et al (2006) Effect of letrozole versus placebo on bone mineral density in women with primary breast cancer completing 5 or more years of adjuvant tamoxifen: a companion study to NCIC CTG MA.17. *J Clin Oncol* 24:3629–3635
 72. Coombes RC, Kilburn LS, Snowdon CF et al (2007) Survival and safety of exemestane versus tamoxifen after 2–3 years' tamoxifen treatment (Intergroup Exemestane Study): a randomised controlled trial. *Lancet* 369:559–570
 73. Napoli N, Rastelli A, Ma C et al (2013) Genetic polymorphism at Val(80) (rs700518) of the CYP19A1 gene is associated with aromatase inhibitor associated bone loss in women with ER (+) breast cancer. *Bone* 55:309–314
 74. Malcolm JB, Derweesh IH, Kincade MC, DiBlasio CJ, Lamar KD, Wake RW, Patterson AL (2007) Osteoporosis and fractures after androgen deprivation initiation for prostate cancer. *Can J Urol* 14:3551–3559
 75. Alibhai SM, Breunis H, Timilshina N et al (2010) Impact of androgen-deprivation therapy on physical function and quality of life in men with nonmetastatic prostate cancer. *J Clin Oncol* 28:5038–5045
 76. Saad F, Adachi JD, Brown JP, Canning LA, Gelmon KA, Josse RG, Pritchard KI (2008) Cancer treatment-induced bone loss in breast and prostate cancer. *J Clin Oncol* 26:5465–5476
 77. Shahinian VB, Kuo YF, Freeman JL, Orihuela E, Goodwin JS (2005) Increasing use of gonadotropin-releasing hormone agonists for the treatment of localized prostate carcinoma. *Cancer* 103:1615–1624
 78. Sieber PR, Keiller DL, Kahnoski RJ, Gallo J, McFadden S (2004) Bicalutamide 150 mg maintains bone mineral density during monotherapy for localized or locally advanced prostate cancer. *J Urol* 171:2272–2276, quiz 2435
 79. Smith MR, Goode M, Zietman AL, McGovern FJ, Lee H, Finkelstein JS (2004) Bicalutamide monotherapy versus leuprolide monotherapy for prostate cancer: effects on bone mineral density and body composition. *J Clin Oncol* 22:2546–2553
 80. Allain TJ (2006) Prostate cancer, osteoporosis and fracture risk. *Gerontology* 52:107–110
 81. McLeod N, Huynh CC, Rashid P (2006) Osteoporosis from androgen deprivation therapy in prostate cancer treatment. *Aust Fam Physician* 35:243–245
 82. Moyad MA (2003) Osteoporosis and prostate cancer. *Urol Oncol* 21:374
 83. Smith MR (2003) Management of treatment-related osteoporosis in men with prostate cancer. *Cancer Treat Rev* 29:211–218
 84. Smith MR, Egerdie B, Hernandez Toriz N et al (2009) Denosumab in men receiving androgen-deprivation therapy for prostate cancer. *N Engl J Med* 361:745–755
 85. Ryan CJ, Molina A, Griffin T (2013) Abiraterone in metastatic prostate cancer. *N Engl J Med* 368:1458–1459
 86. Townsend MF, Sanders WH, Northway RO, Graham SD Jr (1997) Bone fractures associated with luteinizing hormone-releasing hormone agonists used in the treatment of prostate carcinoma. *Cancer* 79:545–550
 87. Adler RA (2011) Management of osteoporosis in men on androgen deprivation therapy. *Maturitas* 68:143–147
 88. Aksnes LH, Bruland OS (2007) Some musculo-skeletal sequelae in cancer survivors. *Acta Oncol* 46:490–496
 89. Chaganti RK, Parimi N, Lang T, Orwoll E, Stefanick ML, Nevitt M, Lane NE (2010) Bone mineral density and prevalent osteoarthritis of the hip in older men for the Osteoporotic Fractures in Men (MrOS) Study Group. *Osteoporos Int* 21:1307–1316
 90. Daniell HW (1997) Osteoporosis after orchiectomy for prostate cancer. *J Urol* 157:439–444
 91. Daniell HW, Clark JC, Pereira SE, Niazi ZA, Ferguson DW, Dunn SR, Figueroa ML, Stratte PT (2001) Hypogonadism following prostate-bed radiation therapy for prostate carcinoma. *Cancer* 91:1889–1895
 92. Grigsby PW, Perez CA (1986) The effects of external beam radiotherapy on endocrine function in patients with carcinoma of the prostate. *J Urol* 135:726–727
 93. Igdem S, Alco G, Ercan T, Barlan M, Ganiyusufoglu K, Unalan B, Turkan S, Okkan S (2010) Insufficiency fractures after pelvic radiotherapy in patients with prostate cancer. *Int J Radiat Oncol Biol Phys* 77:818–823
 94. Overgaard M (1988) Spontaneous radiation-induced rib fractures in breast cancer patients treated with postmastectomy irradiation. A clinical radiobiological analysis of the influence of fraction size and dose–response relationships on late bone damage. *Acta Oncol* 27:117–122
 95. El Maghraoui A (2004) Corticosteroid-induced osteoporosis. *Presse Med* 33:1213–1217
 96. Sambrook PN (2005) How to prevent steroid induced osteoporosis. *Ann Rheum Dis* 64:176–178
 97. Sambrook PN, Diamond T, Ferris L, Fiatarone-Singh M, Flicker L, MacLennan A, Nowson C, O'Neill S, Greville H (2001) Corticosteroid induced osteoporosis. Guidelines for treatment. *Aust Fam Physician* 30:793–796
 98. Lustberg MB, Reinbolt RE, Shapiro CL (2012) Bone health in adult cancer survivorship. *J Clin Oncol* 30:3665–3674
 99. Kanis JA, McCloskey EV, Johansson H, Cooper C, Rizzoli R, Reginster JY, Scientific A, Board of the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO) and the Committee of Scientific Advisors of the International Osteoporosis Foundation (IOF) (2013) European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int* 24:23–57
 100. Bruder JM, Ma JZ, Basler JW, Welch MD (2006) Prevalence of osteopenia and osteoporosis by central and peripheral bone mineral density in men with prostate cancer during androgen-deprivation therapy. *Urology* 67:152–155
 101. Reid DM, Doughty J, Eastell R, Heys SD, Howell A, McCloskey EV, Powles T, Selby P, Coleman RE (2008) Guidance for the management of breast cancer treatment-induced bone loss: a consensus position statement from a UK Expert Group. *Cancer Treat Rev* 34(Suppl 1):S3–18
 102. Hillner BE, Ingle JN, Chlebowski RT et al (2003) American Society of Clinical Oncology 2003 update on the role of bisphosphonates and bone health issues in women with breast cancer. *J Clin Oncol* 21:4042–4057
 103. Body JJ, Bergmann P, Boonen S, Boutsens Y, Devogelaer JP, Goemaere S, Reginster JY, Rozenberg S, Kaufman JM (2007) Management of cancer treatment-induced bone loss in early breast and prostate cancer—a consensus paper of the Belgian Bone Club. *Osteoporos Int* 18:1439–1450
 104. Hadji P, Body JJ, Aapro MS, Brufsky A, Coleman RE, Guise T, Lipton A, Tubiana-Hulin M (2008) Practical guidance for the management of aromatase inhibitor-associated bone loss. *Ann Oncol* 19:1407–1416
 105. van den Bergh JP, van Geel TA, Lems WF, Geusens PP (2010) Assessment of individual fracture risk: FRAX and beyond. *Curr Osteoporos Rep* 8:131–137

106. Watts NB (2011) The Fracture Risk Assessment Tool (FRAX(R)): applications in clinical practice. *J Womens Health (Larchmt)* 20: 525–531
107. Kanis JA, Hans D, Cooper C et al (2011) Interpretation and use of FRAX in clinical practice. *Osteoporos Int* 22:2395–2411
108. Kanis JA, McCloskey EV, Johansson H, Oden A, Strom O, Borgstrom F (2010) Development and use of FRAX in osteoporosis. *Osteoporos Int* 21(Suppl 2):S407–413
109. Saylor PJ, Kaufman DS, Michaelson MD, Lee RJ, Smith MR (2010) Application of a fracture risk algorithm to men treated with androgen deprivation therapy for prostate cancer. *J Urol* 183:2200–2205
110. Cameron ID, Gillespie LD, Robertson MC, Murray GR, Hill KD, Cumming RG, Kerse N (2012) Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst Rev* 12, CD005465
111. Karlsson MK, Magnusson H, von Schewelow T, Rosengren BE (2013) Prevention of falls in the elderly—a review. *Osteoporos Int* 24:747–762
112. Vasikaran S, Eastell R, Bruyere O et al (2011) Markers of bone turnover for the prediction of fracture risk and monitoring of osteoporosis treatment: a need for international reference standards. *Osteoporos Int* 22:391–420
113. Kanis JA, Johansson H, Oden A et al (2004) A family history of fracture and fracture risk: a meta-analysis. *Bone* 35:1029–1037
114. Klotzbuecher CM, Ross PD, Landsman PB, Abbott TA 3rd, Berger M (2000) Patients with prior fractures have an increased risk of future fractures: a summary of the literature and statistical synthesis. *J Bone Miner Res* 15:721–739
115. McCloskey EV, Vasireddy S, Threlkeld J, Eastaugh J, Parry A, Bonnet N, Beneton M, Kanis JA, Charlesworth D (2008) Vertebral fracture assessment (VFA) with a densitometer predicts future fractures in elderly women unselected for osteoporosis. *J Bone Miner Res* 23:1561–1568
116. Diamond TH, Bucci J, Kersley JH, Aslan P, Lynch WB, Bryant C (2004) Osteoporosis and spinal fractures in men with prostate cancer: risk factors and effects of androgen deprivation therapy. *J Urol* 172:529–532
117. Costelloe CM, Rohren EM, Madewell JE et al (2009) Imaging bone metastases in breast cancer: techniques and recommendations for diagnosis. *Lancet Oncol* 10:606–614
118. Bristow RG (2008) Biomarkers of clinical trials using molecular inhibitors and radiotherapy: state-of-the-art. Preface. *Cancer Metastasis Rev* 27:337–338
119. Steinborn MM, Heuck AF, Tiling R, Bruegel M, Gauger L, Reiser MF (1999) Whole-body bone marrow MRI in patients with metastatic disease to the skeletal system. *J Comput Assist Tomogr* 23: 123–129
120. Shie P, Cardarelli R, Brandon D, Erdman W, Abdulrahim N (2008) Meta-analysis: comparison of F-18 fluorodeoxyglucose-positron emission tomography and bone scintigraphy in the detection of bone metastases in patients with breast cancer. *Clin Nucl Med* 33: 97–101
121. Yang HL, Liu T, Wang XM, Xu Y, Deng SM (2011) Diagnosis of bone metastases: a meta-analysis comparing (1)(8)FDG PET, CT, MRI and bone scintigraphy. *Eur Radiol* 21:2604–2617
122. Cook GJ (2010) PET and PET/CT imaging of skeletal metastases. *Cancer Imaging* 10:1–8
123. Keaveny TM, Kopperdahl DL, Melton LJ 3rd, Hoffmann PF, Amin S, Riggs BL, Khosla S (2010) Age-dependence of femoral strength in white women and men. *J Bone Miner Res* 25:994–1001
124. Keyak JH, Kaneko TS, Rossi SA, Pejic MR, Tehranzadeh J, Skinner HB (2005) Predicting the strength of femoral shafts with and without metastatic lesions. *Clin Orthop Relat Res* 439:161–170
125. Tanck E, van Aken JB, van der Linden YM, Schreuder HW, Binkowski M, Huizenga H, Verdonchot N (2009) Pathological fracture prediction in patients with metastatic lesions can be improved with quantitative computed tomography based computer models. *Bone* 45:777–783
126. Dearnaley DP, Mason MD, Parmar MK, Sanders K, Sydes MR (2009) Adjuvant therapy with oral sodium clodronate in locally advanced and metastatic prostate cancer: long-term overall survival results from the MRC PR04 and PR05 randomised controlled trials. *Lancet Oncol* 10:872–876
127. Morgan GJ, Davies FE, Gregory WM et al (2010) First-line treatment with zoledronic acid as compared with clodronic acid in multiple myeloma (MRC Myeloma IX): a randomised controlled trial. *Lancet* 376:1989–1999
128. Wong MH, Stockler MR, Pavlakis N (2012) Bisphosphonates and other bone agents for breast cancer. *Cochrane Database Syst Rev* 2, CD003474
129. Kohno N, Aogi K, Minami H, Nakamura S, Asaga T, Iino Y, Watanabe T, Goessl C, Ohashi Y, Takashima S (2005) Zoledronic acid significantly reduces skeletal complications compared with placebo in Japanese women with bone metastases from breast cancer: a randomized, placebo-controlled trial. *J Clin Oncol* 23: 3314–3321
130. Coleman RE, Marshall H, Cameron D et al (2011) Breast-cancer adjuvant therapy with zoledronic acid. *N Engl J Med* 365:1396–1405
131. Gnant M, Mlineritsch B, Schippinger W et al (2009) Endocrine therapy plus zoledronic acid in premenopausal breast cancer. *N Engl J Med* 360:679–691
132. Lipton A, Siena S, Rader M, Bilynsky B, Viniegra M, Richardson G, Beuzebec P, Clemens MR, Ke C, Jun S (2010) Comparison of denosumab versus zoledronic acid (za) for treatment of bone diseases in advanced cancer patients: an integrated analysis of 3 pivotal trials. *Ann Oncol* 21(viii):379
133. MacKenzie M, Quinton-Gladstone C, Haynes AE, Hotte S (2010) Zoledronic acid for the treatment of bone metastases secondary to renal cell carcinoma. Toronto (ON): Cancer Care Ontario program in Evidence-based Care CED-CCO Special Advice Report No 18
134. Möbus V, Diel IJ, Harbeck N, et al. (2011) GAIN Study: a phase III trial to compare ETC. vs. EC-TX and Ibandronate vs. observation in patients with node-positive primary breast cancer- 1st interim efficacy analysis. 34th annual CTRC-AACR San Antonio Breast Cancer Symposium
135. S0307 Zoledronate, Clodronate, or Ibandronate in Treating Women Who Have Undergone Surgery for Stage I, Stage II, or Stage III Breast Cancer. <http://clinicaltrials.gov/ct2/show/NCT00127205?term=s0307&rank=2>. Accessed June 2013
136. Saad F, Gleason DM, Murray R, Tchekmedyian S, Venner P, Lacombe L, Chin JL, Vinholes JJ, Goas JA, Chen B (2002) A randomized, placebo-controlled trial of zoledronic acid in patients with hormone-refractory metastatic prostate carcinoma. *J Natl Cancer Inst* 94:1458–1468
137. Stopeck AT, Lipton A, Body JJ et al (2010) Denosumab compared with zoledronic acid for the treatment of bone metastases in patients with advanced breast cancer: a randomized, double-blind study. *J Clin Oncol* 28:5132–5139
138. Fizazi K, Carducci M, Smith M et al (2011) Denosumab versus zoledronic acid for treatment of bone metastases in men with castration-resistant prostate cancer: a randomised, double-blind study. *Lancet* 377:813–822
139. Smith MR (2011) Denosumab to prevent bone metastasis-free survival in men with castrate-resistant prostate cancer: results of a global Phase 3, randomised, double-blind trial. AUA meeting Washington DC: Oral presentation
140. Henry DH, Costa L, Goldwasser F et al (2011) Randomized, double-blind study of denosumab versus zoledronic acid in the treatment of bone metastases in patients with advanced cancer (excluding breast and prostate cancer) or multiple myeloma. *J Clin Oncol* 29:1125–1132

141. Brown JE, Barrios CH, Diel IJ et al (2010) Incidence and outcomes of osteonecrosis of the jaw from an integrated analysis of three pivotal randomized double-blind, double-dummy phase 3 trials comparing denosumab and zoledronic acid for treatment of bone metastases in advanced cancer patients or myeloma. *Bone Suppl* 1:S18–S19
142. Doyle C, Kushi LH, Byers T et al (2006) Nutrition and physical activity during and after cancer treatment: an American Cancer Society guide for informed choices. *CA Cancer J Clin* 56:323–353
143. Schmitz KH, Courneya KS, Matthews C et al (2010) American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc* 42:1409–1426
144. WCRF/AICR (2007) Food, nutrition and the prevention of cancer: a global perspective expert report. WCRF/AICR, London
145. Holick MF (2007) Vitamin D deficiency. *N Engl J Med* 357:266–281
146. Sato Y, Iwamoto J, Kanoko T, Satoh K (2005) Amelioration of osteoporosis and hypovitaminosis D by sunlight exposure in hospitalized, elderly women with Alzheimer's disease: a randomized controlled trial. *J Bone Miner Res* 20:1327–1333
147. Brufsky A, Bundred N, Coleman R, Lambert-Falls R, Mena R, Hadji P, Jin L, Schenk N, Ericson S, Perez EA (2008) Integrated analysis of zoledronic acid for prevention of aromatase inhibitor-associated bone loss in postmenopausal women with early breast cancer receiving adjuvant letrozole. *Oncologist* 13:503–514
148. Geisler J, Lonning PE, Krag LE et al (2006) Changes in bone and lipid metabolism in postmenopausal women with early breast cancer after terminating 2-year treatment with exemestane: a randomised, placebo-controlled study. *Eur J Cancer* 42:2968–2975
149. Neuhaus ML, Sorensen B, Hollis BW et al (2008) Vitamin D insufficiency in a multiethnic cohort of breast cancer survivors. *Am J Clin Nutr* 88:133–139
150. Napoli N, Vattikuti S, Ma C, Rastelli A, Rayani A, Donepudi R, Asadfar M, Yarramaneni J, Ellis M, Armamento-Villareal R (2010) High prevalence of low vitamin D and musculoskeletal complaints in women with breast cancer. *Breast J* 16:609–616
151. Rastelli AL, Taylor ME, Gao F, Armamento-Villareal R, Jamalabadi-Majidi S, Napoli N, Ellis MJ (2011) Vitamin D and aromatase inhibitor-induced musculoskeletal symptoms (AIMSS): a phase II, double-blind, placebo-controlled, randomized trial. *Breast Cancer Res Treat* 129:107–116
152. Rizzoli R, Boonen S, Brandi ML, Bruyere O, Cooper C, Kanis JA, Kaufman JM, Ringe JD, Weryha G, Reginster JY (2013) Vitamin D supplementation in elderly or postmenopausal women: a 2013 update of the 2008 recommendations from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Curr Med Res Opin* 29(4):305–313
153. Bolland MJ, Grey A, Reid IR (2013) Calcium supplements and cardiovascular risk in the Women's Health Initiative. *Osteoporos Int* 24(8): 2371–2372
154. Van Poznak CH, Temin S, Yee GC et al (2011) American Society of Clinical Oncology executive summary of the clinical practice guideline update on the role of bone-modifying agents in metastatic breast cancer. *J Clin Oncol* 29:1221–1227
155. Rizzoli R (2011) Bisphosphonates for post-menopausal osteoporosis: are they all the same? *QJM* 104:281–300
156. Delmas PD, Balena R, Confravreux E, Hardouin C, Hardy P, Bremond A (1997) Bisphosphonate risidronate prevents bone loss in women with artificial menopause due to chemotherapy of breast cancer: a double-blind, placebo-controlled study. *J Clin Oncol* 15: 955–962
157. Fuleihan Gel H, Salamoun M, Mourad YA, Chehal A, Salem Z, Mahfoud Z, Shamseddine A (2005) Pamidronate in the prevention of chemotherapy-induced bone loss in premenopausal women with breast cancer: a randomized controlled trial. *J Clin Endocrinol Metab* 90:3209–3214
158. Hershman DL, McMahon DJ, Crew KD, Cremers S, Irani D, Cucchiara G, Brafman L, Shane E (2008) Zoledronic acid prevents bone loss in premenopausal women undergoing adjuvant chemotherapy for early-stage breast cancer. *J Clin Oncol* 26:4739–4745
159. Hershman DL, McMahon DJ, Crew KD, Shao T, Cremers S, Brafman L, Awad D, Shane E (2010) Prevention of bone loss by zoledronic acid in premenopausal women undergoing adjuvant chemotherapy persist up to one year following discontinuing treatment. *J Clin Endocrinol Metab* 95:559–566
160. Kim JE, Ahn JH, Jung KH et al (2011) Zoledronic acid prevents bone loss in premenopausal women with early breast cancer undergoing adjuvant chemotherapy: a phase III trial of the Korean Cancer Study Group (KCSG-BR06-01). *Breast Cancer Res Treat* 125:99–106
161. Powles TJ, McCloskey E, Paterson AH, Ashley S, Tidy VA, Nevantaus A, Rosenqvist K, Kanis J (1998) Oral clodronate and reduction in loss of bone mineral density in women with operable primary breast cancer. *J Natl Cancer Inst* 90:704–708
162. Saarto T, Blomqvist C, Valimaki M, Makela P, Sama S, Elomaa I (1997) Chemical castration induced by adjuvant cyclophosphamide, methotrexate, and fluorouracil chemotherapy causes rapid bone loss that is reduced by clodronate: a randomized study in premenopausal breast cancer patients. *J Clin Oncol* 15:1341–1347
163. Shapiro CL, Halabi S, Hars V et al (2011) Zoledronic acid preserves bone mineral density in premenopausal women who develop ovarian failure due to adjuvant chemotherapy: final results from CALGB trial 79809. *Eur J Cancer* 47:683–689
164. Vehmanen L, Saarto T, Elomaa I, Makela P, Valimaki M, Blomqvist C (2001) Long-term impact of chemotherapy-induced ovarian failure on bone mineral density (BMD) in premenopausal breast cancer patients. The effect of adjuvant clodronate treatment. *Eur J Cancer* 37:2373–2378
165. Van Poznak C, Hannon RA, Mackey JR, Campone M, Apffelstaedt JP, Clack G, Barlow D, Makris A, Eastell R (2010) Prevention of aromatase inhibitor-induced bone loss using risidronate: the SABRE trial. *J Clin Oncol* 28:967–975
166. Lester JE, Dodwell D, Purohit OP, Gutcher SA, Ellis SP, Thorpe R, Horsman JM, Brown JE, Hannon RA, Coleman RE (2008) Prevention of anastrozole-induced bone loss with monthly oral ibandronate during adjuvant aromatase inhibitor therapy for breast cancer. *Clin Cancer Res* 14:6336–6342
167. Greenspan SL, Resnick NM, Nelson JB (2006) Once weekly oral alendronate prevents bone loss in men on androgen deprivation therapy for prostate cancer. *J Urol* 175(suppl 4):975, abstract
168. Taxel P, Dowsett R, Richter L, Fall P, Klepinger A, Albertsen P (2010) Risidronate prevents early bone loss and increased bone turnover in the first 6 months of luteinizing hormone-releasing hormone-agonist therapy for prostate cancer. *BJU Int* 106:1473–1476
169. Michaelson MD, Kaufman DS, Lee H, McGovern FJ, Kantoff PW, Fallon MA, Finkelstein JS, Smith MR (2007) Randomized controlled trial of annual zoledronic acid to prevent gonadotropin-releasing hormone agonist-induced bone loss in men with prostate cancer. *J Clin Oncol* 25:1038–1042
170. Ellis GK, Bone HG, Chlebowski R, Paul D, Spadafora S, Fan M, Kim D (2009) Effect of denosumab on bone mineral density in women receiving adjuvant aromatase inhibitors for non-metastatic breast cancer: subgroup analyses of a phase 3 study. *Breast Cancer Res Treat* 118:81–87
171. Ellis GK, Bone HG, Chlebowski R, Paul D, Spadafora S, Smith J, Fan M, Jun S (2008) Randomized trial of denosumab in patients receiving adjuvant aromatase inhibitors for nonmetastatic breast cancer. *J Clin Oncol* 26:4875–4882
172. Study to determine treatment effects of denosumab in patients with breast cancer receiving aromatase inhibitor therapy. <http://www.clinicaltrials.gov/ct2/show/NCT00556374?term=denosumab+and+cancer&rank=10>. Accessed June 2013
173. Saad F, Smith MR, Egerdie B, Tammela TL, Feldman RA, Heracek J, Szwedowski M, Ke C, Leder B, Goessl C (2009) Denosumab for

- prevention of fractures in men receiving androgen deprivation therapy (ADT) for prostate cancer. *J Clin Oncol* 27:15s, abstract 5056
174. Smith MR, Morton RA, Barnette KG, Sieber PR, Malkowicz SB, Rodriguez D, Hancock ML, Steiner MS (2010) Toremifene to reduce fracture risk in men receiving androgen deprivation therapy for prostate cancer. *J Urol* 184:1316–1321
 175. Smith MR, Fallon MA, Lee H, Finkelstein JS (2004) Raloxifene to prevent gonadotropin-releasing hormone agonist-induced bone loss in men with prostate cancer: a randomized controlled trial. *J Clin Endocrinol Metab* 89:3841–3846
 176. Kyle RA, Yee GC, Somerfield MR et al (2007) American Society of Clinical Oncology 2007 clinical practice guideline update on the role of bisphosphonates in multiple myeloma. *J Clin Oncol* 25:2464–2472
 177. Terpos E, Morgan G, Dimopoulos MA et al (2013) International myeloma working group recommendations for the treatment of multiple myeloma-related bone disease. *J Clin Oncol* 31:2347–2357
 178. Van Poznak CH, Von Roenn JH, Temin S (2011) American society of clinical oncology clinical practice guideline update: recommendations on the role of bone-modifying agents in metastatic breast cancer. *J Oncol Pract* 7:117–121
 179. Lentle B, Cheung AM, Hanley DA, et al. (2011) Osteoporosis Canada 2010 Guidelines for the Assessment of Fracture Risk. *Can Assoc Radiol J* 62(4):243–250
 180. Leslie WD, Schousboe JT (2011) A review of osteoporosis diagnosis and treatment options in new and recently updated guidelines on case finding around the world. *Curr Osteoporos Rep* 9:129–140
 181. Papaioannou A, Morin S, Cheung AM et al (2010) 2010 clinical practice guidelines for the diagnosis and management of osteoporosis in Canada: summary. *CMAJ* 182:1864–1873
 182. Watts NB, Bilezikian JP, Camacho PM et al (2010) American Association of Clinical Endocrinologists Medical Guidelines for Clinical Practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract* 16(Suppl 3):1–37
 183. Coleman R, Costa L, Saad F et al (2011) Consensus on the utility of bone markers in the malignant bone disease setting. *Crit Rev Oncol Hematol* 80:411–432
 184. Eastell R, Barton I, Hannon RA, Chines A, Gamero P, Delmas PD (2003) Relationship of early changes in bone resorption to the reduction in fracture risk with risedronate. *J Bone Miner Res* 18:1051–1056
 185. Hannon R, Eastell R (2000) Preanalytical variability of biochemical markers of bone turnover. *Osteoporos Int* 11(Suppl 6):S30–44
 186. Watts NB, Lewiecki EM, Bonnick SL et al (2009) Clinical value of monitoring BMD in patients treated with bisphosphonates for osteoporosis. *J Bone Miner Res* 24:1643–1646
 187. Hadji P, Aapro MS, Body JJ, Bundred NJ, Brufsky A, Coleman RE, Gnant M, Guise T, Lipton A (2011) Management of aromatase inhibitor-associated bone loss in postmenopausal women with breast cancer: practical guidance for prevention and treatment. *Ann Oncol* 22:2546–2555
 188. Mottet N, Bellmunt J, Bolla M et al (2011) EAU guidelines on prostate cancer. Part II: treatment of advanced, relapsing, and castration-resistant prostate cancer. *Eur Urol* 59:572–583
 189. Aapro M, Abrahamsson PA, Body JJ et al (2008) Guidance on the use of bisphosphonates in solid tumours: recommendations of an international expert panel. *Ann Oncol* 19:420–432
 190. Body JJ, Coleman R, Clezardin P, Ripamonti C, Rizzoli R, Aapro M, International Society of Geriatric Oncology (2007) International Society of Geriatric Oncology (SIOG) clinical practice recommendations for the use of bisphosphonates in elderly patients. *Eur J Cancer* 43:852–858
 191. Coleman RE, Bertelli G, Beaumont T, Kunkler I, Miles D, Simmonds PD, Jones AL, Smith IE (2012) UK guidance document: treatment of metastatic breast cancer. *Clin Oncol (R Coll Radiol)* 24:169–176
 192. Bird JM, Owen RG, D'Sa S et al (2011) Guidelines for the diagnosis and management of multiple myeloma 2011. *Br J Haematol* 154:32–75
 193. Terpos E, Sezer O, Croucher PI et al (2009) The use of bisphosphonates in multiple myeloma: recommendations of an expert panel on behalf of the European Myeloma Network. *Ann Oncol* 20:1303–1317
 194. Lacy MQ, Dispenzieri A, Gertz MA et al (2006) Mayo clinic consensus statement for the use of bisphosphonates in multiple myeloma. *Mayo Clin Proc* 81:1047–1053
 195. National Comprehensive Cancer Network (2012) NCCN Guidelines version 1.2012 Multiple Myeloma.
 196. Harrouseau JL, Greil R, Kloke O, Force EGT (2005) ESMO Minimum Clinical Recommendations for diagnosis, treatment and follow-up of multiple myeloma. *Ann Oncol* 16(Suppl 1):i45–47
 197. Grossmann M, Hamilton EJ, Gilfillan C, Bolton D, Joon DL, Zajac JD (2011) Bone and metabolic health in patients with non-metastatic prostate cancer who are receiving androgen deprivation therapy. *Med J Aust* 194:301–306
 198. Saad F, Higano CS, Sartor O, Colombel M, Murray R, Mason MD, Tubaro A, Schulman C (2006) The role of bisphosphonates in the treatment of prostate cancer: recommendations from an expert panel. *Clin Genitourin Cancer* 4:257–262