

# Pacific Journal of Mathematics

## **CARATHEODORY THEOREMS IN CONVEX PRODUCT STRUCTURES**

JOHN ROBERT REAY

## CARATHEODORY THEOREMS IN CONVEX PRODUCT STRUCTURES

JOHN R. REAY

Various attempts have been made to place convexity in an axiomatic setting. Recently J. Eckhoff has considered the classic theorem of Radon in several different settings. Most of his work is done in what we call an Eckhoff space, i.e., in a finite product of euclidean spaces where convex sets are defined as the cartesian products of usual convex sets in each component space. The purpose of this paper is to investigate the closely related theorem of Caratheodory and its generalizations in this setting.

The papers of F.W. Levi [5] and Dauzer, Grunbaum, Klee [3] have various approaches to axiomatic settings of convexity, and a good bibliography for before 1961. See the papers of Eckhoff [4] and Bonnice-Reay [2] for more recent results and references.

1. Eckhoff spaces. The pair  $(E, \mathcal{E})$  denotes an *Eckhoff space* provided (1)  $E$  is a direct cartesian product  $E = \prod_{i=1}^n E_i$  where each  $E_i$  is a  $d_i$ -dimensional euclidean space with  $\mathcal{E}_i$  the family of all convex sets of  $E_i$ , and (2)  $\mathcal{E} = \{\prod_{i=1}^n A_i : A_i \in \mathcal{E}_i\}$  is the family of all *product-convex* sets in  $E$ . For any set  $X \subset E$ , the set  $E(X) = \bigcap \{A : X \subset A \in \mathcal{E}\}$  is called the *product-convex hull* of  $X$ . Let  $\pi_i : E \rightarrow E_i$  denote the usual projection. Then we can consider  $E$  as a linear space of dimension  $d = \sum_{i=1}^n d_i$ , and  $E(X) = \prod_{i=1}^n (\text{conv } \pi_i X)$  where  $\text{conv } B$  denotes the usual convex hull of  $B$  in each euclidean space  $E_i$ . The cardinality of  $B$  will be denoted by  $|B|$ . Using the notation of Bonnice-Klee [1] and others, we say that  $\text{int}_r B$  is the set of all points  $p$  for which there exists an  $r$ -dimensional simplex contained in  $B$  and containing  $p$  in its relative interior.

2. Caratheodory-type theorems. By a Caratheodory-type theorem we mean a result which asserts that if a point is embedded in the (axiomatically defined) hull of a set  $X$ , then it is similarly embedded in the hull of a sufficiently small subset of  $X$ . Note that the case  $n = 1$  of Theorem 1 below is the result usually called Caratheodory's theorem.

**THEOREM 1.** *If  $X$  is any subset of an Eckhoff space  $E = \prod_{i=1}^n E_i$  of dimension  $d = \sum d_i$  and if  $p \in E(X)$ , then  $p \in E(Y)$  for some  $Y \subset X$  with  $|Y| \leq d + \delta$ , where  $\delta = 1$  if  $n = 1$  and  $\delta = 0$  if  $n > 1$ . Fur-*

thermore, if  $m = |Y|$  is the cardinality of a smallest subset  $Y$  of  $X$  for which  $p \in E(Y)$ , then  $p \in \text{int}_r E(Y)$  where

$$\max(0, m - n) \leq r \leq (m - 1)|\{d_i : d_i \geq m\}| + \sum_{d_i < m} d_i.$$

*Proof.* If  $n = 1$  the upper and lower bounds on  $r$  reduce to  $r = m - 1$ , that is,  $p$  lies interior to the  $(m - 1)$ -simplex determined by the  $m$  points of  $Y$ .

Assume  $n = 2$ . It suffices to show that there is a set  $Y \subset X$  for which  $|Y| \leq d_1 + d_2 = d$  and  $\pi_i p \in \text{conv } \pi_i Y$  for  $i = 1, 2$ . Applying Caratheodory's theorem to  $E_1$ , there is a subset  $Y_1 \subset X$  with  $|Y_1| \leq d_1 + 1$  for which  $\pi_1 p \in \text{conv } \pi_1 Y_1$ . Now if  $\pi_2 p \in \text{conv } \pi_2 Y_1$  as well, then  $p \in E(Y_1)$  and we are done. Otherwise choose a set  $Y_2 \subset X$  of minimal cardinality such that  $\pi_2 p \in \text{conv } \pi_2 (Y_1 \cup Y_2)$ . Since we may choose one of the  $d_2 + 1$  points of Caratheodory's theorem arbitrarily (see [6], Lemmas 4.1-4.4) it follows that  $|Y_2| \leq d_2$ . Thus letting  $Y = Y_1 \cup Y_2$  it follows that  $p \in E(Y)$  and  $|Y| \leq |Y_1| + |Y_2| \leq d_1 + d_2 + 1$ . We are therefore done unless both  $|Y_1| = d_1 + 1$  and  $|Y_2| = d_2$  and  $Y_1 \cap Y_2 = \emptyset$ . In this case we will show that one correctly chosen point may be removed from  $Y_1 \cup Y_2$ .

*Case 1.*  $\pi_2 p \in \text{conv } \pi_2 Y_2$ . In this case we reverse the roles of  $E_1$  and  $E_2$  in the above argument, i.e., let  $Y_2$  be as above and redefine  $Y_1$  to be a set of minimal cardinality so that  $\pi_1 p \in \text{conv } (Y_2 \cup Y_1)$ . Then  $|Y_1| \leq d_1$  and  $|Y_2| = d_2$  and  $p \in E(Y_2 \cup Y_1)$ .

*Case 2.*  $\pi_2 p \notin \text{conv } \pi_2 Y_2$ . In this case  $|Y_2| = d_2$  so for each point  $y \in Y_1$  it is true that  $\pi_2 p \in \text{conv } \pi_2 (\{y\} \cup Y_2)$ . Thus some point of  $\pi_1 Y_2$  in the space  $E_1$  may be used to replace a particular point of  $Y_1$ , say  $y_1$ . Then  $\pi_1 p \in \text{conv } \pi_1 (Y_2 \cup (Y_1 - \{y_1\}))$  and it is still true that  $\pi_2 p \in \text{conv } \pi_2 ((Y_1 - \{y_1\}) \cup Y_2)$ . This establishes Case 2, and hence proves the first statement of the theorem if  $n = 2$ .

The case when  $n \geq 3$  now follows easily. As in the case  $n = 2$  there exists a set  $Y_1 \cup Y_2 \subset X$  for which  $\pi_i p \in \text{conv } \pi_i (Y_1 \cup Y_2)$  for  $i = 1, 2$  and  $|Y_1 \cup Y_2| \leq d_1 + d_2$ . For each  $i \geq 3$  there is a set  $Y_i \subset X$ , by Caratheodory's theorem in  $E_i$ , such that  $|Y_i| \leq d_i$  and  $\pi_i p \in \text{conv } \pi_i (Y_1 \cup Y_2 \cup Y_i)$ . The set  $Y = \bigcup_{i=1}^n Y_i$  then has the desired properties;  $p \in E(Y)$  and  $|Y| \leq \sum_{i=1}^n d_i = d$ . This establishes the first half of the theorem.

To prove the last statement, let  $Y \subset X$  be a smallest subset for which  $p \in E(Y)$  and suppose  $|Y| = m$ . Then for each  $i$ ,  $\pi_i p \in \text{int}_{r(i)} \text{conv } \pi_i Y$  for some largest nonnegative integer  $r(i) \leq d_i$ , and  $p \in \text{int}_r E(Y)$  where  $r = \sum_{i=1}^n r(i)$ . It follows that  $r$  assumes a minimal value whenever each  $r(i)$  is as small as possible, within the constraint  $|Y| = m$ .

This is achieved when the points of  $Y$  are used as “inefficiently as possible”, specifically, when for some partition  $Y = Y_1 \cup \dots \cup Y_n$  we have  $\pi_i p \in \text{conv } \pi_i Y_i$  and the points of  $\pi_i(Y \sim Y_i)$  are not used in  $E_i$ . For example, if  $\pi_i(Y \sim Y_i)$  is a single point in  $E_i$  then the points of  $\pi_i Y$  are the vertices of a simplex in  $E_i$  of dimension  $|Y_i|$ , and  $\pi_i p$  is interior to the subsimplex  $\text{conv } \pi_i Y_i$ . In any case, Caratheodory’s theorem (case  $n = 1$ ) implies that  $r(i) = |Y_i| - 1$ . Thus  $r = \Sigma_i r(i) = (\Sigma_i |Y_i|) - n = m - n$ , and in general  $r \geq m - n$ . The other inequality on  $r$  follows from the fact that the  $m$  points of  $Y$  projected onto each space  $E_i$  can have a convex hull of dimension at most  $\min(m - 1, d_i)$  in  $E_i$ . This proves Theorem 1.

EXAMPLES. The following examples show that the bounds in Theorem 1 cannot, in general, be improved.

(1) For each  $i = 1, 2, \dots, n$ , let  $X_i$  be a subset of  $E_i$  for which  $X_i \cup \{0\}$  form the vertices of a nondegenerate  $d_i$ -simplex, and let  $p_i$  be in the relative interior of the simplex  $\text{conv } X_i$ . Define  $p \in E$  by the relations  $\pi_i p = p_i$ . For each point  $x_i$  in each set  $X_i$  define the point  $\bar{x}_i \in E$  by the relations  $\pi_i \bar{x}_i = x_i$  and  $\pi_j \bar{x}_i = 0$  if  $j \neq i$ . Let  $X \subset E$  be the set of all such points  $\bar{x}_i$ . Then clearly  $p \in E(X)$ , but  $p \notin E(Y)$  for any proper subset of  $Y$  of  $X$ , and  $|X| = \Sigma |X_i| = \Sigma d_i = d$ . Furthermore  $p \in \text{int}_r E(X)$  where

$$r = \Sigma(|X_i| - 1) = |X| - n = m - n .$$

(2) As a second example, let  $m$  be any integer for which  $1 \leq m \leq \max \{d_i + 1 : i = 1, \dots, n\}$ . For each subspace  $E_i$  if  $m \leq d_i + 1$  let  $\{x_{ij}\}_{j=1}^m$  be the vertices of a nondegenerate simplex in  $E_i$ . If  $m > d_i + 1$  let  $\{x_{ij} : j = 1, \dots, d_i + 1\}$  be the vertices of a nondegenerate simplex, and let  $x_{i1} = x_{ij}$  for  $j = (d_i + 2), \dots, m$ . In either case choose a point  $p_i$  in the relative interior of this simplex. Now define  $p \in E$  by the relations  $\pi_i p = p_i$  and let  $X = \{x_j\}_{j=1}^m \subset E$  where each  $x_j$  is defined by  $\pi_i x_j = x_{ij} \in E_i$ . Then  $|X| = m$  and  $p \in E(X)$  but  $p \notin E(Y)$  for any proper subset  $Y$  of  $X$ . Also  $p \in \text{int}_r E(X)$  where

$$r = (m - 1)|\{d_i : d_i \geq m\}| + \Sigma_{d_i < m} d_i .$$

The case where  $n = 1$  and  $r = d$  in Theorem 2 below is commonly called Steinitz’s theorem.

**THEOREM 2.** *If  $X$  is any subset of an Eckhoff space  $E = \prod_{i=1}^n E_i$  and if  $r \geq 0$  is the largest integer for which  $p \in \text{int}_r E(X)$ , then  $p \in \text{int}_r E(Y)$  for some subset  $Y \subset X$  with  $|Y| \leq 2r + \delta$  where  $\delta$  is the number of spaces  $E_i$  for which  $\pi_i p \notin \text{int}_1 \text{conv } \pi_i X$ .*

*Proof.* Let  $r(i)$  be the largest integer for which  $\pi_i p \in \text{int}_{r(i)} \text{conv} \pi_i X$ . Thus  $r = \Sigma r(i)$ . By the Bonnice-Klee theorem (see [1], Th. 2.5) and the maximality of  $r(i)$  there is a subset  $Y_i$  of  $X$  for which  $\pi_i p \in \text{int}_{r(i)} \text{conv} \pi_i Y_i$  and  $|Y_i| \leq 2r(i)$  if  $r(i) > 0$ , and  $|Y_i| = 1$  if  $r(i) = 0$ . Thus letting  $Y = \bigcup_{i=1}^n Y_i$  we have  $p \in \text{int}_r E(Y)$  and  $|Y| = |\bigcup Y_i| \leq \Sigma |Y_i| \leq 2\Sigma_{r(i)>0} r(i) + \Sigma_{\{i|r(i)=0\}} 1 = 2r + \delta$ . This proves Theorem 2.

Using the techniques from the examples given above, it is easy to construct sets  $X$  in Eckhoff spaces which show that the bounds of Theorem 2 cannot, in general, be improved. A further generalization may be obtained by considering  $p \in \text{int}_s E(X)$  where  $0 < s < r$ , and ask the cardinality of the smallest  $Y \subset X$  for which  $p \in \text{int}_s E(Y)$ . This is the spirit of the Bonnice-Klee Theorem (see [1] and [6]). Another approach is to add further information about the set  $X$ , and ask how the bound on  $|Y|$  may be improved. For example, if it is known that  $k_i$  is the dimension of the highest-dimensional simplex with vertices in  $\pi_i X \subset E_i$  and having  $\pi_i p$  in its relative interior, then the bound on  $|Y|$  can, in general, be improved. See Bonnice-Reay [2] for a bibliography and results of this type. Also connectedness or symmetry conditions on  $X$  may lead to an improvement of the bound on  $|Y|$ . See [6] for a bibliography and results of this type.

These theorems and others which depend even more upon the structure of  $X$  are similar to the above theorems, but are much more complicated and are therefore omitted.

## REFERENCES

1. W. Bonnice and V. Klee, *The generation of convex hulls*, Math. Ann., **152** (1963), 1-29.
2. W. Bonnice and J. Reay, *Relative interiors of convex hulls*, Proc. Amer. Math. Soc., **20** (1969), 246-250.
3. L. Danzer, B. Grunbaum, and V. Klee, *Helly's theorem and its relatives*, Proc. of Symp. in Pure Math., vol. 7, "Convexity", Amer. Math. Soc., 1963, 101-180.
4. J. Eckhoff, *Der Satz von Radon in konvexen Productstrukturen I-II*, Monatsh. Math., **72** (1968), 303-314; **73** (1969), 7-30.
5. F. W. Levi, *On Helly's theorem and the axioms of convexity*, J. Indian Math. Soc., (N.S.) **15** (1951), 65-76.
6. J. R. Reay, *Generalizations of a theorem of Caratheodory*, Amer. Math. Soc. Memoir No. 54, 1965.

Received October 16, 1969.

WESTERN WASHINGTON STATE COLLEGE

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

B. D. Arendt and C. J. Stuth, <i>On the structure of commutative periodic semigroups</i> .....	1
B. D. Arendt and C. J. Stuth, <i>On partial homomorphisms of semigroups</i> ....	7
Leonard Asimov, <i>Extensions of continuous affine functions</i> .....	11
Claude Elias Billigheimer, <i>Regular boundary problems for a five-term recurrence relation</i> .....	23
Edwin Ogilvie Buchman and F. A. Valentine, <i>A characterization of the parallelepiped in <math>E^n</math></i> .....	53
Victor P. Camillo, <i>A note on commutative injective rings</i> .....	59
Larry Jean Cummings, <i>Decomposable symmetric tensors</i> .....	65
J. E. H. Elliott, <i>On matrices with a restricted number of diagonal values</i> ...	79
Garth Ian Gaudry, <i>Bad behavior and inclusion results for multipliers of type <math>(p, q)</math></i> .....	83
Frances F. Gulick, <i>Derivations and actions</i> .....	95
Langdon Frank Harris, <i>On subgroups of prime power index</i> .....	117
Jutta Hausen, <i>The hypo residuum of the automorphism group of an abelian <math>p</math>-group</i> .....	127
R. Hrycay, <i>Noncontinuous multifunctions</i> .....	141
A. Jeanne LaDuke, <i>On a certain generalization of <math>p</math> spaces</i> .....	155
Marion-Josephine Lim, <i>Rank preservers of skew-symmetric matrices</i> .....	169
John Hathway Lindsey, II, <i>On a six dimensional projective representation of the Hall-Janko group</i> .....	175
Roger McCann, <i>Transversally perturbed planar dynamical systems</i> .....	187
Theodore Windle Palmer, <i>Real <math>C^*</math>-algebras</i> .....	195
Don David Porter, <i>Symplectic bordism, Stiefel-Whitney numbers, and a Novikov resolution</i> .....	205
Tilak Raj Prabhakar, <i>On a set of polynomials suggested by Laguerre polynomials</i> .....	213
B. L. S. Prakasa Rao, <i>Infinitely divisible characteristic functionals on locally convex topological vector spaces</i> .....	221
John Robert Reay, <i>Caratheodory theorems in convex product structures</i> ....	227
Allan M. Sinclair, <i>Eigenvalues in the boundary of the numerical range</i> .....	231
David R. Stone, <i>Torsion-free and divisible modules over matrix rings</i> .....	235
William Jennings Wickless, <i>A characterization of the nil radical of a ring</i> .....	255