

Cardiac dysfunction in cancer patients: beyond direct cardiomyocyte damage of anticancer drugs. Novel cardio-oncology insights from the joint 2019 meeting of the ESC Working Groups of Myocardial Function and Cellular Biology of the Heart

Short title: Novel cardio-oncology insights

Carlo G Tocchetti, MD, PhD, FISC, FHFA^{1,2*}, Pietro Ameri, MD, PhD, FHFA^{3*}, Rudolf A. de Boer MD, PhD, FHFA⁴, Yuri D'Alessandra, PhD⁵, Michele Russo, PhD^{1,6}, Daniela Sorriento, PhD⁷, Michele Ciccarelli, MD, PhD, FESC⁸, Bernadett Kiss⁹, Luc Bertrand, PhD¹⁰, Dana Dawson, MD, PhD, FESC¹¹, Ines Falcao-Pires, PhD¹², Mauro Giacca, MD, PhD^{13,14,15}, Nazha Hamdani, PhD¹⁶, Wolfgang A. Linke, PhD¹⁷, Manuel Mayr, MD, PhD¹⁵, Jolanda van der Velden, PhD¹⁸, Serena Zacchigna, MD, PhD^{13,14}, Alessandra Ghigo, PhD⁶, Emilio Hirsch, PhD⁶, Alexander R Lyon, MD, PhD, FHFA¹⁹, Anikó Görbe, MD, PhD^{9,20}, Péter Ferdinandy, MD, PhD, MBA, FESC^{9,20}, Rosalinda Madonna, MD, PhD^{21,22}, Stephane Heymans, MD, PhD, FESC^{23, 24}, Thomas Thum, MD, PhD, FAHA, FESC, FHFA²⁵

*CGT and PA share first authorship

¹ Department of Translational Medical Sciences, Federico II University, Naples, Italy

² Interdepartmental Center of Clinical and Translational Sciences (CIRCET), Federico II University, Naples, Italy

³ Cardiovascular Disease Unit, IRCCS Ospedale Policlinico San Martino, Genova, Italy – IRCCS Italian Cardiovascular Network & Department of Internal Medicine; University of Genova, Italy

⁴ University of Groningen, University Medical Center Groningen, Department of Cardiology, AB31, PO Box 30.001, 9700 RB Groningen, the Netherlands

⁵ Immunology and Functional Genomics Unit, Centro Cardiologico Monzino IRCCS, Milan, Italy

⁶ Department of Molecular Biotechnology and Health Sciences, Molecular Biotechnology Center, University of Torino, Torino, Italy

⁷ Department of Advanced Biomedical Sciences, Federico II University, Naples, Italy

⁸ University of Salerno, Italy

⁹ Cardiometabolic Research Group and MTA-SE System Pharmacology Research Group, Department of Pharmacology and Pharmacotherapy, Semmelweis University, Budapest, Hungary

¹⁰ IREC Institute, Pole of Cardiovascular Research, Université catholique de Louvain, Brussels, Belgium

¹¹ School of Medicine and Dentistry, University of Aberdeen, United Kingdom

¹² Unidade de Investigação e Desenvolvimento Cardiovascular, Departamento de Cirurgia e Fisiologia, Faculdade de Medicina, Universidade do Porto, Portugal

¹³ Department of Medicine, Surgery and Health Sciences and Cardiovascular Department, Centre for Translational Cardiology, Azienda Sanitaria Universitaria Integrata Trieste, Italy

¹⁴ International Center for Genetic Engineering and Biotechnology (ICGEB), Trieste, Italy

¹⁵ King's British Heart Foundation Centre, King's College London, London, United Kingdom

¹⁶ Department of Molecular and Experimental Cardiology, Ruhr Universität Bochum, Bochum Germany and Department of Cardiology, St. Joseph Hospital, Ruhr University Bochum, Witten, Germany

¹⁷ Institute of Physiology 2, University of Muenster, Germany

¹⁸ Department of Physiology, Amsterdam UMC, Vrije Universiteit, Amsterdam Cardiovascular Sciences Institute, Amsterdam, The Netherlands

¹⁹ Cardio-Oncology Service, Royal Brompton Hospital and Imperial College London, UK

²⁰ Pharmahungary Group, Szeged, Hungary

²¹ Institute of Cardiology, University of Pisa, Pisa, Italy

²² Center for Cardiovascular Biology and Atherosclerosis Research, McGovern School of Medicine, University of Texas Health Science Center at Houston, Houston, TX, USA

²³ Cardiovascular Research Institute Maastricht (CARIM), Department of Cardiology, Maastricht University Medical Centre, Maastricht University, The Netherlands

²⁴ Department of Cardiovascular Sciences, University of Leuven, Belgium

²⁵ Institute for Molecular and Translational Therapeutic Strategies (IMTTS), Hannover Medical School, Hannover, Germany

Address for correspondence:

Carlo Gabriele Tocchetti, MD, PhD, FISC, FHFA

UOS Gestione del paziente oncologico in Medicina Interna

Dipartimento di Scienze Mediche Traslazionali

Centro Interdipartimentale di Ricerca Clinica e Traslazionale (CIRCET)

Universita' degli Studi di Napoli Federico II

Via Pansini 5

80131 Napoli, ITALY

Phone +39-081-746-2242

Fax +39-081-746-2246

carlo.gabriele.tocchetti@unina.it

OR

Thomas Thum, MD, PhD, FAHA, FESC, FHFA

Institute of Molecular and Translational Therapeutic Strategies (IMTTS)

Hannover Medical School

Carl-Neuberg-Str. 1

30627 Hannover, GERMANY

Phone +49-511-532-9174

thum.thomas@mh-hannover.de

ABSTRACT

In the Western countries cardiovascular disease and cancer are the leading causes of death in the ageing population. Recent epidemiological data suggest that cancer is more frequent in patients with prevalent or incident cardiovascular disease, in particular heart failure. Indeed, there is a tight link in terms of shared risk factors and mechanisms between heart failure and cancer. Heart failure induced by anticancer therapies has been extensively studied, primarily focusing on the toxic effects that antitumor treatments exert on cardiomyocytes. In this Cardio-Oncology update, members of the ESC WGs of Myocardial Function and of Cellular Biology of the Heart discuss novel evidence interconnecting cardiac dysfunction and cancer via pathways in which cardiomyocytes may be involved, but are not central. In particular, the multiple roles of cardiac stromal cells (endothelial cells, fibroblasts) and inflammatory cells are highlighted. Also, the gut microbiota is depicted as a new player at the crossroads between heart failure and cancer. Finally, the role of non-coding RNAs in Cardio-Oncology is also addressed. All these insights are expected to fuel additional research efforts in the field of Cardio-Oncology.

1. Introduction

In the industrialized world, cardiovascular (CV) disease and cancer are the leading causes of death in the ageing population ¹. Left ventricular dysfunction (LVD) and heart failure (HF) are not rare across the broad population of cancer patients. In cancer patients, CV disease is the most frequent non-cancer cause of death ². HF and cancer share the same risk factors (e.g. ageing, smoking, obesity, diabetes, dyslipidemia, alcohol intake, inflammation) ^{3, 4}. Furthermore cancer and HF may have ancillary factors linking the two together ⁵. Registries have observed that HF patients have a higher cumulative incidence of cancer, with a worse prognosis when both co-exist ⁶, suggesting that cancer surveillance may be useful in the management of HF patients ⁷. Finally, an increased cumulative incidence of cancer among HF patients 30 days after MI has been reported, compared to HF-free patients 30 days after MI ⁸.

When considering these observations, it should be taken into account that there may be a surveillance bias, due to the fact that these study patients usually undergo an intense follow-up program that may lead to anticipate cancer diagnosis, sometimes discovering malignancies that would have gone undiscovered. Moreover, some of the most common therapies used to treat HF patients may play a role in revealing tumors otherwise asymptomatic (e.g. a latent intestinal neoplasm can bleed due to anti-thrombotic therapy)³. Clinical presentations can also be difficult to distinguish between HF and cancer, since the 2 conditions can share some common symptoms (fatigue, dyspnea, weight loss, muscle wasting, oedema) ^{1,3}. This may delay the diagnosis of new-onset cancer in HF patients due to the overlap in clinical manifestation. Furthermore, CV function and predictors of exercise capacity have been shown to be impaired in patients with cancer *per se*⁹. Hence, symptoms due to a tumor may overlap with those of HF and be attributed to heart disease. This may even delay cancer diagnosis, as symptoms might be thought of as due to advancing disease rather than new cancer ³. Although the relationship

between cancer and HF is not well-defined in clinical studies, there are increasing data to suggest mechanistic links between the two conditions that we discuss in our manuscript.

Beside these reciprocal relations, cancer and HF carry an independent risk of mortality and also limit optimal treatment of the other condition when they co-exist, contributing to higher mortality. In addition, the cardiotoxicity risk related to treatment with anticancer drugs may unmask or deteriorate pre-existing HF³. The mechanisms driving HF triggered by anticancer therapies have been extensively investigated over the last 20 years and important insights have been uncovered¹⁰⁻¹². Nonetheless, major questions are still open, and the answers to these questions may lay the foundations for new strategies to detect, monitor and treat cancer-therapy induced cardiotoxicity. On the other hand, research into the common pathways linking cancer and HF regardless of anticancer drugs has just begun¹³.

The latest insights in translational Cardio-Oncology were discussed during the joint meeting of the Working Groups of Myocardial Function and the WG of Cellular Biology of the Heart of the European Society of Cardiology, held in Naples, Italy, in May 2019. In particular, given the systemic involvement of both HF and cancer, the Cardio-Oncology session focused on the contribution of organs, systems and cells other than cardiomyocytes to the pathogenesis of cardiac dysfunction in cancer patients, and to the interconnection between cancer and HF, primarily via inflammation. Opportunities and the current limitations in the use of microRNAs (miRNA) in cardio-oncology were also discussed. These topics are reviewed here, to provide the reader with updated information and further stimulate research in the field.

2. Role of non-cardiomyocytes in cancer treatment-related cardiotoxicity

The heart is a multicellular organ composed by cardiomyocytes, fibroblasts, neurons, endothelial and hematopoietic-derived cells. In fact, cardiomyocytes are not the most abundant cell type¹⁴. The different cardiac cell populations have diverse functions, but also interact

through complex intercellular communications¹⁵. Most studies performed so far have focused on the effects of anticancer drugs on cardiomyocytes, in both *in vitro* systems and *in vivo* models¹⁶ (see table 1). Briefly, among the many forms of cardiotoxicity caused by several anticancer drugs (table 2), cardiac dysfunction due anthracyclines such as doxorubicin (DOXO) has historically been the most relevant¹⁷. From a pathophysiological point of view, anthracyclines induce cardiomyocyte death, mainly apoptosis and necrosis, via different molecular mechanisms, including but not limited to induction of oxidative stress, activation of DNA damage responses and impairment of mitochondrial biogenesis and metabolism¹⁸⁻²⁰. Among other mechanisms involved in anthracycline-induced cardiotoxicity, abnormalities in myocardial energetics have also been studied^{21, 22}. Also biological drugs, designed to target specific oncologic pathways may be cardiotoxic, since these pathways play a major role in the maintenance of cardiac homeostasis, especially during stressful conditions, such as hypertension or hypertrophy²³. For instance, *human epidermal growth factor receptor 2 (HER/ErbB2)* and *angiogenesis inhibitors* profoundly affect cardiomyocytes metabolism and contractile proteins, as discussed in comprehensive reviews^{16, 24-26}.

In addition, antitumor therapies likely also affect non-cardiomyocytes in the heart. For instance, DOXO has been shown to exert toxic effects on cultured cardiac endothelial cells²⁷ and fibroblasts²⁸⁻³⁰. This direct activity on non-cardiomyocytes may partly account for the cardiotoxicity of the drug, e.g. endothelial cells lose their barrier function with increased permeability and myocardial injury.

The impact of the toxicity of DOXO and any other antitumor treatment on non-cardiomyocytes can be better understood when it is placed into the context of the intercellular cross-talks in the heart. This concept is exemplified by the current knowledge about the cardiotoxicity of anti-HER2 drugs³¹. Besides being expressed in breast cancer cells, HER2/ErbB2 is also physiologically present in cardiomyocytes together with another receptor

tyrosine kinases (RTK) of the same family, HER4/ErbB4³². Upon binding of HER4/ErbB4 by neuregulin-1 (NRG) and other ligands secreted by cardiac microvascular endothelial cells, HER2/ErbB2 and HER4/ErbB4 form heterodimers and initiate protective signaling cascades. Therefore, drugs targeting HER/ErbB2 are postulated to disrupt the NRG1-HER2/ErbB2-mediated endothelial cell-cardiomyocyte crosstalk and make cardiomyocytes more vulnerable to other stressors (Figure 1). It is notable that trastuzumab, used in the treatment of human epidermal growth factor receptor (HER)-2+ breast cancer, also directly damages cardiomyocytes and endothelial cells^{33, 34}.

The inhibitors of the RTK for vascular endothelial growth factor (VEGF) and platelet-derived growth factor (PDGF) cause cardiac microvascular dysfunction secondary to depletion of coronary microvascular pericytes³⁵. The resulting myocardial hypoxia leads to sustained expression of hypoxia-inducible factor alpha (HIF- α), which was demonstrated to be sufficient to cause cardiomyopathy^{36,37}. Indeed, enhanced vascular permeability and reversible microvascular vasoconstriction have been reported in patients receiving therapies targeting VEGF and PDGF receptor (VEGFR and PDGFR, respectively)³⁸. Moreover, this mechanism of toxicity well explains the clinical observation that cardiomyopathy associated with anti-VEGFR/PDGFR agents is reversible³⁹.

However, evidence obtained over the last years suggests that blockade of VEGF signaling also interrupt endothelial cell-cardiomyocyte communication (Figure 1). VEGF binds VEGFR on endothelial cells to stimulate angiogenesis, but also to induce the release of angiocrines (including ErbB4 and ErbB1 ligands) that modulate the function and homeostasis of adjacent cardiomyocytes⁴⁰. Thus, drugs that inhibit VEGFR may alter cardiac function by interfering with the VEGF-VEGFR signaling axis, as well as by promoting endothelial cell dysfunction and death^{41, 42}. High-throughput screening of RTK inhibitors pinpointed those

targeting VEGFR2 and PDGFR as the most toxic in human induced pluripotent stem cell (hiPSC)–derived endothelial cells ⁴³.

Experimental models and analyses of human biopsies indicate that some features of HF with preserved ejection fraction (HFpEF) are at least in part driven by cardiac endothelial cell dysfunction. This latter elicits inflammatory infiltration of the myocardium, fibroblasts activation to deposit collagen excessively and increased stiffness triggered by a reduction of nitric oxide-dependent signaling ⁴⁴⁻⁴⁶. Consistently with the epidemiological finding that the risk of HFpEF is correlated with prior radiotherapy for breast cancer ⁴⁷, similar features were demonstrated in rats receiving cardiac radiation ⁴⁸. Since non-proliferating cardiomyocytes are considered resistant to ionizing radiation, other cell types, and in particular endothelial cells, are predicted to be the main target of radiation therapy leading leads to HF ⁴⁹.

Fibroblasts also regulate cardiomyocytes and inflammatory cells through their secretome ^{50, 51}. In a recent study, DOXO caused both apoptosis of cardiac fibroblasts and secretion of Fas ligand, which in turn promoted cardiomyocyte death in a paracrine manner ⁵². Conditional deletion of ataxia telangiectasia mutated kinase (ATM) in cardiac fibroblasts attenuated cardiac cell apoptosis, LVD and mortality in response to DOXO, suggesting that fibroblast are central in the pathogenesis of DOXO cardiotoxicity through ATM. The interactions between fibroblasts and other cardiac cell types, and the mechanisms in the cardiotoxicity of anticancer therapies, are an important area for future research ⁵². Senescence of fibroblasts and possibly other cardiac stromal cells is especially worth being investigated, since it has been proposed that it plays a major role in the pathogenesis of heart disease ⁵³.

In conclusion, oncological drugs and radiotherapy induce abnormalities in non-cardiomyocytes, which secondarily derange the networks with cardiomyocytes and may lead to LVD and HF. Additional studies are needed, ^{54, 55} also considering that cardiotoxicity may be evident in an already damaged myocardium, but may remain latent or hidden in the healthy

heart⁵⁶. Since cardiac diseases and their comorbidities significantly change the global cardiac transcriptome, proteome and metabolome, it is not surprising that several drugs may act differently on the diseased versus healthy hearts^{57,58}. Novel cardiac safety testing platforms involving combined experimental models of cardiac diseases in the presence and absence of major cardiovascular co-morbidities and/or co-treatments are needed⁵⁹. In this regard, cardiac organoids may allow modelling the complexity of the interactions between the different cardiac cell populations and, thereby, comprehensively evaluate the effects of anticancer therapies⁵⁵.

3. Interconnections between cancer and heart failure

Recently, attention has been drawn to the fact that cancer and heart disease have a reciprocal relationship: while the presence of cancer may cause LVD, the presence of HF associates with excess incident cancer^{3-5,60}. The communication between these two threatening diseases is complex, intriguing and involves many components.

First, during life and aging, several risk factors accumulate, which lead to chronic inflammation, oxidative stress, and protein and DNA instability. Classical CV risk factors, including obesity, diabetes, dyslipidemia and inflammation, are also associated with the development of cancer. Many of these risk factors lead to accumulation of fat mass, which is an active endocrine organ, secreting inflammatory factors and adipokines, which in turn have been associated with new onset CV disease (CVD) and new onset cancer^{3-5,60}.

Second, genetic mutations that accumulate throughout life, such as clonal hematopoiesis of indeterminate potential (CHIP), defined as the presence of clonal leukocytes with impaired immune properties derived by acquired mutation in hematopoietic stem cells, have been associated with both cancer and CVD, including HF⁶¹⁻⁶⁵. These mutations usually occur in a few genes, including DNMT3A, TET2, ASXL1, PPM1D, JAK2, TP53, SF3B1, and SRSF2⁶⁶. The risk of developing CHIP increases with aging and, although it rarely results in development

of hematologic malignancies, it seems to be tightly linked to increased CV events and worse HF prognosis^{61-65, 67}.

Also, genetic mutations in sarcomeric proteins predispose to HF in patients undergoing chemotherapy. Unrecognized rare variants in cardiomyopathy-associated genes, particularly *Titin* truncating variants, have been shown to increase the risk for systolic dysfunction and cardiac events in a relatively small population of both children and adults undergoing chemotherapy. In specific populations, genotype variant testing, along with cumulative chemotherapy dosage and traditional cardiovascular risk factors, may be useful to improve the identification of cancer patients with a higher risk for developing HF upon chemotherapy⁶⁸.

Other CV risk factors, such as hypertension and trace albuminuria, have been related to cancer development. Therefore, systemic risk factors likely exert effects on several damage pathways, and it is hypothesized that individual additional risk factors, such as genetic predisposition or pre-existing conditions, will also contribute to the risk of one or both conditions.

Third, cancer and CVD are both associated with profound changes in tissue structure, either growth of entirely new tissue or tissue deformation, remodeling, and scarring of pre-existing tissues, such as heart, endothelial cells and matrix. Neoplasms are characterized by stroma, which is matrix tissue supporting the tumor, providing a scaffold, structure, and connections to adjacent organs. Further, most cancers, and especially metastases, rely on strong neovascularization requiring mitogenic endothelial cells and pericytes, where multiple growth factors play a role. In comparison, damaged cardiac tissue leads to dysfunctional cardiomyocytes, and also may develop extracellular matrix remodeling, fibrosis and scar. Matrix is produced by activated fibroblasts and multiple cell types homing in, including monocytes, macrophages and neutrophils. The cardiac scar is not a static structure, but rather is a dynamic and secreting structure⁶⁹.

4. Psychological convergence of HF and cancer

There is a well established psychological impact on patients suffering from chronic conditions, notably heart failure. This is one of the main aims of cardiac rehabilitation programmes in these patients. Unfortunately, rehabilitation programmes have only recently been implemented in cancer patients, in a generic “one fits all” umbrella rather than bespoke guidelines for specific cancers. It is notable however, the recognition that both cardiac dysfunction syndromes and cancers have a significant impact in regards to neuronal changes. Whilst these have only just been thought of, the molecular and cellular mechanisms of neurobiology change remain relatively unknown. It is likely that both neuronal changes per se as well as modifications in signaling and transmission underlie the clinical states of depression or cognitive changes in these patients. The most likely culprit remains the chronic systemic inflammatory state present in both, probably responsible for an enhanced level of oxidative stress, DNA damage, mitochondrial dysfunction as well as synaptic modifications^{70; 71}.

Whilst there is available evidence to support a link between certain chemotherapies and peripheral neuropathy (for example cisplatin), the issue of clinical states of depression/cognitive changes and them per se being a basis for autonomic dysfunction seen in these patients is far more complex and yet undemonstrated. At this current time it does not have the level of evidence and merits further exploration.

5. Inflammation at the crossroad between cancer, cardiotoxicity of anticancer therapies and heart failure

Abnormal inflammation is increasingly recognized as a common driver of CVD and cancer^{72, 73}. HF is characterized by a state of mild chronic systemic inflammation, with increased circulating concentrations of pro-inflammatory cytokines, such as tumor necrosis

factor-alpha (TNF- α), interleukin-1 (IL-1), and interleukin-6 (IL-6). Myocardial injury itself triggers the recruitment and the activation of immune cells, which in turn produce pro-inflammatory cytokines and contribute to a self-perpetuating inflammatory state that underlies adverse tissue remodeling, primarily associated with capillary dysfunction and fibrosis ⁷⁴. Doxorubicin-induced damage also involves inflammation (Figure 2), with upregulation of pro-inflammatory toll-like receptor 4 (TLR4) in macrophages ⁷⁵, higher levels as TNF- α and IL-6 and reduced levels of the anti-inflammatory cytokine IL-10 ⁷⁶. Cardiac function was preserved and survival improved in TLR2 knock-out mice after DOXO exposure compared to wild-types ⁷⁷. DOXO also induces local modulators of inflammation and fibrosis, produced by both macrophages and fibroblasts. Increased production of the matricellular protein thrombospondin-2 (TSP2) is protective in mice treated with DOXO. Enhanced myocyte damage in the absence of TSP-2 was associated with impaired activation of the Akt signaling pathway. Inhibition of Akt phosphorylation in cardiomyocytes significantly reduced TSP-2 expression, unveiling a unique feedback loop between Akt and TSP-2 ⁷⁸. Importantly, CCL2/CCR2-dependent recruitment of functional antigen-presenting cells into tumors is a desired therapeutic effect of anthracyclines ⁷⁹.

Indeed, for decades oncologists have been developing strategies to modulate inflammation in order to achieve therapeutic anticancer immune responses ⁸⁰. The first attempts were not really successful, since cancer escapes T-cell-mediated cancer-specific immunity via inhibitory pathways mediated by cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), programmed cell death protein 1 (PD-1), and programmed cell death ligand 1 (PD-L1) (all depressing the antineoplastic activity of T lymphocytes) ⁸¹. On the opposite, in the last years, Immune Checkpoint Inhibitors (ICIs), such as monoclonal antibodies (mAbs) targeting CTLA-4, PD-1 and PD-L1, have dramatically improved the outcome of many malignancies, but serious immune related cardiovascular adverse events have been observed ⁸²⁻⁸⁴ (Figure 2). Interfering

with the CTLA-4 and PD-1 axes can bring to autoimmune myocarditis and dilated cardiomyopathy⁸⁵, suggesting that these molecules play an important role in preventing autoimmunity⁸⁶. Hence, immunosuppressive therapies may be necessary to halt immune related adverse events (IRAEs) and major adverse cardiovascular events (MACE)⁸⁷⁻⁸⁹.

More recently, engineered T cells with chimeric antigen receptors (CAR-T cells) have been approved by the U.S. Food and Drug Administration (FDA) as the first genetically modified autologous T-cell immunotherapeutic agents that target CD-19. CD-19 is broadly expressed on most B-cell malignancies and has limited expression beyond B-cell lineage^{90,91}. Unfortunately, CAR-T cells are burdened by cytokine release syndrome (CRS) that is due to elevated levels of inflammatory cytokines released by activated CAR-T cells and other immune cells such as macrophages, with fever and tachycardia that may be associated with hypotension and hypoxia. Also, cardiac dysfunction and extremely serious complications such as vascular leak syndrome with circulatory collapse and multiorgan failure can be dreadful side effects of these therapies^{92, 93} (Figure 2). Beside CAR-T cells, bispecific antibodies such as blinatumumab (that targets CD19 and CD3 and is increasingly used in the treatment of Philadelphia chromosome negative B cell acute lymphoblastic leukemia (ALL)) can also lead to CRS and cardiomyopathy⁹⁴.

Interestingly, inflammation in cancer plays a dual role. On the one hand it is essential to recognize and destroy cancer cells; on the other hand it provides a fertile milieu for tumorigenesis and plays key roles in different steps of tumor development, from initiation and promotion to invasion and metastasis. Tumor-associated inflammation favors proliferation and survival of malignant cells, promotes angiogenesis and metastasis, undermines adaptive immune responses, and potentially interferes with responses to hormones and chemotherapeutic agents^{95,96}. The finding that anti-inflammatory agents are effective in the prevention of cancer and CVD further advocates inflammation as a common contributor to both diseases. A 2019

study concluded that chronic systemic low-grade inflammation, measured by CRP levels <10 mg/L, is a risk factor for incident cancer, in particular lung cancer, in patients with stable CVD. The relation between inflammation and incident cancer is seen in former and current smokers and is uncertain in never smokers⁹⁷. Blockade of the pro-inflammatory cytokine IL-1 β with canakinumab was shown to significantly reduce the rate of recurrent CV events in patients with previous myocardial infarction (CANTOS trial). At the same time, blocking IL-1 β appeared to protect from lung cancer mortality^{98,99}. Mice exposed to DOXO showed an increase in serum IL-1 β along with other inflammatory factors¹⁰⁰. Moreover, the IL-1 β receptor antagonism protects against DOXO cardiotoxicity¹⁰¹. Similarly, the IL-6 inhibitor tocilizumab can protect against MACE in CAR-T patients⁹³.

The experience of IL-1 β blockade highlights that the identification of key players of the inflammatory response is important to tackle both cancer and heart disease. Among intriguing candidates are PI3Ks, and more specifically the PI3K γ isoform that is enriched in both cardiomyocytes and leukocytes (Figure 2). This implies a key role for this isoform not only in the control of cardiomyocyte pathobiology, but also in the orchestration of the inflammatory response associated to different types of cardiovascular injury¹⁰². PI3K γ is upregulated in patients as well as in mouse models of atherosclerosis, and directs leukocyte infiltration of the arterial wall, which is a key pathogenic event in atherosclerosis¹⁰³. PI3K γ -mediated inflammation is also pivotal to the cardiac response to pressure overload¹⁰⁴.

Besides directing the cardiac response to stress, macrophage PI3K γ expression critically contributes to tumor growth and progression. Intriguingly, macrophages play opposite roles in non-oncological inflammatory conditions and cancer. In response to pathogens or injury, macrophages express cytokines that stimulate cytotoxic T cells to clear infected or damaged cells. Conversely, in cancer macrophages express anti-inflammatory cytokines that induce immune suppression, inhibit T cell-mediated tumor killing and promote resistance to

immunotherapies (i.e. T cell checkpoint inhibitors). PI3K γ has been recently proposed as the molecular switch controlling immune stimulation and suppression in cancer ¹⁰⁵. The unique feature of macrophage PI3K γ , playing a maladaptive role both in heart disease and in cancer, makes this enzyme the ideal pharmacological target to “kill two birds with one stone”, i.e. to halt the tumor and at the same time treat the heart ¹⁰⁶. This is particularly relevant for cancer patients treated with chemotherapy and suffering from iatrogenic cardiotoxicity ^{15, 105 107}. Results from clinical trials assessing the combined anticancer effect of such compounds in a context of cardiac protection are awaited.

6. The gut microbiome in Cardio-Oncology

HF has long been recognized to be associated with altered gut function ^{108, 109}. Low cardiac output in HF results in intestinal ischaemia, with congestion of the splanchnic circulation, bowel wall oedema and impaired intestinal barrier function (Figure 3). This condition increases the overall inflammatory state as well as oxidative stress as a consequence of HF-induced ischaemia and congestion within the gut via enhanced bacterial translocation and the presence of bacterial products in the blood circulation. Increased leakiness modifies the gut environment and affects its resident microbial population ¹¹⁰.

Among the conditions that can influence the gut composition, including individual genetic variability, lifestyle, colonization and delivery at birth ¹¹¹⁻¹¹³, also changes in diet, presence of diseases and relative treatments have to be considered ¹¹⁴. Interestingly, genetic composition of gut microbiota, defined as microbiome, also influences cancer development and progression in different ways ¹¹⁵. Several types of cancers (head and neck, lung, colorectal and cervical carcinomas) promote a shift in microbiome composition ¹¹⁶⁻¹¹⁸. In addition, chemotherapy directly impacts the gut microbiota and its efficacy is strongly influenced by microbiome composition (Figure 3) ^{119, 120}.

Metabolites generated by the gut microbiota derive from the fermentation of indigestible fibres to short-chain fatty acids, that have protective properties (reducing inflammation, oxidative stress^{121, 122} and improving vascular tone). Dietary sources of choline, phosphatidylcholine, l-carnitine, and other methylamine-containing nutrients provide substrates for microbiota-mediated generation of trimethylamine (TMA) that accesses the portal circulation and is converted by the hepatic flavin-containing monooxygenase (FMO) family of enzymes into trimethylamine *N*-oxide (TMAO, Table 3). TMAO can favor the development of atherosclerosis, thrombosis, kidney disease, and HF (Figure 3). High plasma levels of TMAO have been suggested to be predictive of cardiovascular events of mortality, independently from renal function and cardiovascular comorbidities¹¹⁰.

Additionally, the bacterial transformation of bile acids can result in altered bile acid profiles, that in turn can impact systemic inflammatory and fibrotic processes¹¹⁰. Importantly, microbiota-derived peptide mimics may also drive HF, by inducing a lethal inflammatory cardiomyopathy. Cardiac myosin-specific TH17 cells are being imprinted in the intestine by a commensal *Bacteroides* species peptide mimic. These cells promote cardiac inflammation and dysfunction in genetically susceptible individuals¹²³.

Several studies reported SCFAs-producing bacteria perturbation in patients with CVDs¹²⁴. Among these SCFA generated by the gut microbiota, butyrate (BUT) has multiple beneficial effects for our cardiovascular system through different mechanisms^{125-131, 120} (Table 3). BUT exerts major epigenetic effects, acting as a potent inhibitor of histone deacetylase (HDACs) activity. Inhibition of HDACs is well-known to protect the heart from pathologic hypertrophy and ischaemia¹³²⁻¹³⁵. Among HDAC inhibitors, BUT has been shown to exert anti-neoplastic properties *in vitro*^{136, 120}; while its derivatives can enhance the anticancer cytotoxic effects of DOXO while protecting against cardiotoxicity¹³⁷ and can decrease cardiac apoptosis and myocardial dysfunction induced by DOXO, by lowering endoplasmic reticulum stress-initiated

apoptotic signalling and HDAC-inhibition mechanisms^{138, 139}. The cardioprotective effect of BUT and analogues is associated with the production of anti-inflammatory molecules, cytoprotection, modulation of angiogenesis, limiting the occurrence of cardiotoxic manifestations caused by DOXO treatments, with reduction of nitrosative and oxidative stress, counteracting mitochondrial dysfunction¹²¹. In turn, DOXO is reported to induce GUT-microbiota dysbiosis in mice, while the administration of BUT attenuates the inflammation state induced by DOXO¹⁴⁰, fuelling nutraceutical as a new promising area of research to cardio-oncology

7. Opportunities and limitations in the use of noncoding RNAs in Cardio-Oncology

Multiple evidence seems to suggest an involvement of circulating microRNAs (miRNAs) in anthracyclines-induced cardiotoxicity both *in vivo* and in the clinical setting, evidencing a very heterogeneous situation. In particular, when focusing on DOXO, miR-1^{141, 142} and miR-34a¹⁴³⁻¹⁴⁶, showed a drug-induced regulation in tissues and plasma samples, both in patients and animal models. miR-1 is one of the most investigated and most highly expressed miRNAs in cardiac and skeletal muscle, both in physiological^{146, 147} and pathologic^{148, 149} condition. While many groups have indicated miR-1 as a specific circulating marker of heart disease, there is no clear indication about its unambiguous cardiac origin, particularly in anthracyclines-induced toxicity, which is a systemic phenomenon. Similarly, miR-34a was demonstrated to be modulated by anthracyclines both in experimental models^{144, 145, 150} and in breast cancer patients¹⁵¹. Piegari and co-authors, showed that tissue regulation of miR-34a by DOXO was not restricted only to the heart¹⁴⁴, hinting at a multi-tissue contribution to the circulating levels of this miRNA. Indeed, besides cardiomyocytes, smooth muscle cells, fibroblasts, cardiac progenitor cells and endothelial cells may also play a role in DOXO-induced cardiomyopathy¹⁵². Acute DOXO treatment in mice was shown to reduce microvessel density

and VEGF-A expression with a parallel increase in miR-320a¹⁵³. Inhibition of miR320a improved cardiac function, decreased apoptosis, and increased microvessel density in DOXO-treated mice, while overexpression of miR-320a worsened DOXO-induced LV dysfunction¹⁵⁴. Conversely, overexpression of the miR-320a target VEGF-A prevented detrimental effects of miR-320a in DOXO-cardiotoxicity experimental model confirming VEGF as a direct downstream target molecule¹⁵³. Mechanistically, the overexpression of the pro-hypertrophic miR-212/132 cluster in primary rodent and human iPSC-derived cardiomyocytes as well as in in vivo models has been shown to inhibit doxorubicin-induced toxicity¹⁵⁵. Also, another class of noncoding RNAs, circular RNAs, may play a crucial role in mediating cardiotoxicity of doxorubicin; indeed, overexpression of the RNA binding protein Quaking 5 (*Qki5*) strongly attenuated the toxic effect of doxorubicin in a mouse model by regulating a set of circular RNAs including those derived from titin (*Ttn*;¹⁵⁶).

The role of miRNAs as markers of cardiotoxicity has also been investigated. Ruggeri and coworkers¹⁵⁰ showed that after one month from DOXO administration, only a part of the drug-treated mice presented cardiac dysfunction, similarly to the clinical context. miR-1 was again among the circulating miRNAs regulated after cardiotoxicity onset, together with miR-499-5p. In an acute DOXO cardiotoxicity model, the same authors showed that miR-34a-5p and miR-451a were dysregulated in all cardiac chambers, with miR34a-5p showing opposite trends of regulation between the atria and the ventricles of treated mice. In another study using DOXO both in vivo and in vitro acutely and chronically treated cardiomyocytes, DOXO-dependent downregulation of miR-30 led to increased cardiomyocyte apoptosis and abnormalities of cardiomyocyte β -adrenergic receptor signaling¹⁵⁷.

Importantly, only part of circulating miRNAs overlapped with their cardiac counterparts, suggesting only a partial contribution of the heart to the variations in circulating levels of miRNAs upon drug administration. Limitations of the studies are the number of

animals, the number of screened miRNAs (often only selected cardiovascular miRNAs), the acute phase observed, the absence of tumor in the experimental models, and the lack of additional cancer treatments.

Besides the few miRNAs showing a “reproducible sensitivity” to anthracycline treatment, there is a highly heterogeneous picture composed by past and present investigations. While the discrepancies in terms of results could be in part explained by the different experimental models and by the different malignancies and therapies adopted in patients-based investigations, there are at least two fundamental issues that should be addressed in future works. A striking feature of many, if not all, published papers is that no study described a decline of LVEF below the “normal” threshold of 50%, possibly because of lack of a long-term follow-up. Moreover, the vast majority of human-based research studies concentrated on the acute phase of cardiotoxicity ¹⁵⁸, and the same limitations often apply also to *experimental* researches, which rarely go beyond a few days’ time span from treatment to sacrifice.

Additional data on the main non-coding RNAs are summarized in Table 4.

8. Concluding remarks

We discussed several of the novel exciting insights that are emerging in the ever-expanding field of cardio-oncology. More research is required to identify and investigate the pathways and mechanisms underpinning the intimate relationship between CVD and cancer. Current studies focus on shared risk factors, both acquired/modifiable and genetic. The substantial structural changes in diseased organs prompt further studies in an effort to learn how disease in one organ may communicate with another organ. Learning from each disease mechanisms may help to combat both CVD disease and cancer.

Funding

CGT was supported by a Ricerca di Ateneo/Federico II University grant, PA is supported by the Italian Ministry of Health (GR-2018-12365661, CHANGE Study), PF was supported by the National Research, Development and Innovation Office of Hungary (OTKA KH_17 125570, the National Heart Program NVKP 16-1-2016-0017) and by the Higher Education Institutional Excellence Program of the Ministry of Human Capacities in Hungary, within the framework of the Therapeutic Development thematic program of the Semmelweis University.

TT received funding from the EU-ERANET grant EXPERT. LB is Research director of FRS-FNRS, Belgium., RM received funding from Incyte Biosciences.

RAdB is supported by the European Research Council (ERC CoG 818715, SECRETE-HF) and the CVON grants from Netherlands Heart Foundation (DOSIS, 2014-40; SHE-PREDICTS-HF, 2017-21; RED-CVD, 2017-11; and PREDICT2, 2018-30).

This manuscript has been possible thanks to the support of the ERA-Net-CVD project MacroERA, 01KL1706, and IMI2-CARDIATEAM (N° 821508) to SH. We acknowledge the support from the Netherlands Cardiovascular Research Initiative, an initiative with support of the Dutch Heart Foundation, CVON2016-Early HFPEF, 2015-10, CVON She-PREDICTS, grant 2017-21, CVON Arena-PRIME, 2017-18. Furthermore, we acknowledge the support of FWO G091018N and FWO G0B5930N to SH.

IFP is supported by Fundo Europeu de Desenvolvimento Regional (FEDER) through Compete 2020 – Programa Operacional Competitividade E Internacionalização (POCI), the project DOCNET (norte-01-0145-feder-000003) is supported by Norte Portugal regional operational programme (norte 2020), under the Portugal 2020 partnership agreement, through the European Regional Development Fund (ERDF).

Conflict of interest

PF is the founder and CEO of Pharmahungary Group, a group of R&D companies. TT has filed and licensed patents in the field of noncoding RNAs. TT is founder and shareholder of Cardior Pharmaceuticals GmbH. The UMCG, which employs RAdB has received research grants and/or fees from AstraZeneca, Abbott, Bristol-Myers Squibb, Novartis, Novo Nordisk, and Roche. RAdB received speaker fees from Abbott, AstraZeneca, Novartis, and Roche.

REFERENCES

1. Anker MS, von Haehling S, Landmesser U, Coats AJS, Anker SD. Cancer and heart failure-more than meets the eye: Common risk factors and co-morbidities. *European journal of heart failure*. 2018;20:1382-1384
2. Strongman H, Gadd S, Matthews A, Mansfield KE, Stanway S, Lyon AR, Dos-Santos-Silva I, Smeeth L, Bhaskaran K. Medium and long-term risks of specific cardiovascular diseases in survivors of 20 adult cancers: A population-based cohort study using multiple linked uk electronic health records databases. *Lancet*. 2019;394:1041-1054
3. Ameri P, Canepa M, Anker MS, Belenkov Y, Bergler-Klein J, Cohen-Solal A, Farmakis D, Lopez-Fernandez T, Lainscak M, Pudil R, Ruschitska F, Seferovic P, Filippatos G, Coats A, Suter T, Von Haehling S, Ciardiello F, de Boer RA, Lyon AR, Tocchetti CG, Heart Failure Association Cardio-Oncology Study Group of the European Society of C. Cancer diagnosis in patients with heart failure: Epidemiology, clinical implications and gaps in knowledge. *European journal of heart failure*. 2018;20:879-887
4. Meijers WC, de Boer RA. Common risk factors for heart failure and cancer. *Cardiovascular research*. 2019;115:844-853
5. Meijers WC, Maglione M, Bakker SJL, Oberhuber R, Kieneker LM, de Jong S, Haubner BJ, Nagengast WB, Lyon AR, van der Vegt B, van Veldhuisen DJ, Westenbrink BD, van der Meer P, Sillje HHW, de Boer RA. Heart failure stimulates tumor growth by circulating factors. *Circulation*. 2018;138:678-691
6. Banke A, Schou M, Videbaek L, Moller JE, Torp-Pedersen C, Gustafsson F, Dahl JS, Kober L, Hildebrandt PR, Gislason GH. Incidence of cancer in patients with chronic heart failure: A long-term follow-up study. *European journal of heart failure*. 2016;18:260-266
7. Hasin T, Gerber Y, McNallan SM, Weston SA, Kushwaha SS, Nelson TJ, Cerhan JR, Roger VL. Patients with heart failure have an increased risk of incident cancer. *Journal of the American College of Cardiology*. 2013;62:881-886
8. Hasin T, Gerber Y, Weston SA, Jiang R, Killian JM, Manemann SM, Cerhan JR, Roger VL. Heart failure after myocardial infarction is associated with increased risk of cancer. *Journal of the American College of Cardiology*. 2016;68:265-271
9. Cramer L, Hildebrandt B, Kung T, Wichmann K, Springer J, Doehner W, Sandek A, Valentova M, Stojakovic T, Scharnagl H, Riess H, Anker SD, von Haehling S. Cardiovascular function and predictors of exercise capacity in patients with colorectal cancer. *Journal of the American College of Cardiology*. 2014;64:1310-1319
10. Antoniadou C, Small HY, Guzik T. The evolution of cardiovascular research online: Online and on demand. *Cardiovascular research*. 2018;114:e9
11. Moslehi J, Fujiwara K, Guzik T. Cardio-oncology: A novel platform for basic and translational cardiovascular investigation driven by clinical need. *Cardiovascular research*. 2019;115:819-823
12. Molinaro M, Ameri P, Marone G, Petretta M, Abete P, Di Lisa F, De Placido S, Bonaduce D, Tocchetti CG. Recent advances on pathophysiology, diagnostic and therapeutic insights in cardiac dysfunction induced by antineoplastic drugs. *BioMed research international*. 2015;2015:138148
13. Bertero E, Ameri P, Maack C. Bidirectional relationship between cancer and heart failure: Old and new issues in cardio-oncology. *Cardiac failure review*. 2019;5:106-111

14. Pinto AR, Ilinykh A, Ivey MJ, Kuwabara JT, D'Antoni ML, Debuque R, Chandran A, Wang L, Arora K, Rosenthal NA, Tallquist MD. Revisiting cardiac cellular composition. *Circulation research*. 2016;118:400-409
15. Skelly DA, Squiers GT, McLellan MA, Bolisetty MT, Robson P, Rosenthal NA, Pinto AR. Single-cell transcriptional profiling reveals cellular diversity and intercommunication in the mouse heart. *Cell reports*. 2018;22:600-610
16. Ky B, Vejpongsa P, Yeh ET, Force T, Moslehi JJ. Emerging paradigms in cardiomyopathies associated with cancer therapies. *Circulation research*. 2013;113:754-764
17. Ewer MS, Von Hoff DD, Benjamin RS. A historical perspective of anthracycline cardiotoxicity. *Heart failure clinics*. 2011;7:363-372
18. Varricchi G, Ameri P, Cadeddu C, Ghigo A, Madonna R, Marone G, Mercurio V, Monte I, Novo G, Parrella P, Pirozzi F, Pecoraro A, Spallarossa P, Zito C, Mercurio G, Pagliaro P, Tocchetti CG. Antineoplastic drug-induced cardiotoxicity: A redox perspective. *Frontiers in physiology*. 2018;9:167
19. Zhang S, Liu X, Bawa-Khalfe T, Lu LS, Lyu YL, Liu LF, Yeh ET. Identification of the molecular basis of doxorubicin-induced cardiotoxicity. *Nature medicine*. 2012;18:1639-1642
20. Amgalan D, Garner TP, Pekson R, Jia XF, Yanamandala M, Paulino V, Liang FG, Corbalan JJ, Lee J, Chen Y, Karagiannis GS, Sanchez LR, Liang H, Narayanagari S-R, Mitchell K, Lopez A, Margulets V, Scarlata M, Santulli G, Asnani A, Peterson RT, Hazan RB, Condeelis JS, Oktay MH, Steidl U, Kirshenbaum LA, Gavathiotis E, Kitsis RN. A small-molecule allosteric inhibitor of bax protects against doxorubicin-induced cardiomyopathy. *Nature Cancer*. 2020;1:315-328
21. Maslov MY, Chacko VP, Hirsch GA, Akki A, Leppo MK, Steenbergen C, Weiss RG. Reduced in vivo high-energy phosphates precede adriamycin-induced cardiac dysfunction. *American journal of physiology. Heart and circulatory physiology*. 2010;299:H332-337
22. Gupta A, Rohlfen C, Leppo MK, Chacko VP, Wang Y, Steenbergen C, Weiss RG. Creatine kinase-overexpression improves myocardial energetics, contractile dysfunction and survival in murine doxorubicin cardiotoxicity. *PloS one*. 2013;8:e74675
23. Suter TM, Ewer MS. Cancer drugs and the heart: Importance and management. *European heart journal*. 2013;34:1102-1111
24. De Keulenaer GW, Doggen K, Lemmens K. The vulnerability of the heart as a pluricellular paracrine organ: Lessons from unexpected triggers of heart failure in targeted erbb2 anticancer therapy. *Circulation research*. 2010;106:35-46
25. Eschenhagen T, Force T, Ewer MS, de Keulenaer GW, Suter TM, Anker SD, Avkiran M, de Azambuja E, Balligand JL, Brutsaert DL, Condorelli G, Hansen A, Heymans S, Hill JA, Hirsch E, Hilfiker-Kleiner D, Janssens S, de Jong S, Neubauer G, Pieske B, Ponikowski P, Pirmohamed M, Rauchhaus M, Sawyer D, Sugden PH, Wojta J, Zannad F, Shah AM. Cardiovascular side effects of cancer therapies: A position statement from the heart failure association of the european society of cardiology. *European journal of heart failure*. 2011;13:1-10
26. Moslehi JJ. Cardiovascular toxic effects of targeted cancer therapies. *The New England journal of medicine*. 2016;375:1457-1467
27. Luu AZ, Chowdhury B, Al-Omran M, Teoh H, Hess DA, Verma S. Role of endothelium in doxorubicin-induced cardiomyopathy. *JACC. Basic to translational science*. 2018;3:861-870
28. Zhang W, St Clair D, Butterfield A, Vore M. Loss of mrp1 potentiates doxorubicin-induced cytotoxicity in neonatal mouse cardiomyocytes and cardiac fibroblasts. *Toxicological sciences : an official journal of the Society of Toxicology*. 2016;151:44-56
29. Narikawa M, Umemura M, Tanaka R, Hikichi M, Nagasako A, Fujita T, Yokoyama U, Ishigami T, Kimura K, Tamura K, Ishikawa Y. Doxorubicin induces trans-differentiation and mmp1 expression in cardiac fibroblasts via cell death-independent pathways. *PloS one*. 2019;14:e0221940
30. Ghosh AK, Rai R, Park KE, Eren M, Miyata T, Wilsbacher LD, Vaughan DE. A small molecule inhibitor of pai-1 protects against doxorubicin-induced cellular senescence. *Oncotarget*. 2016;7:72443-72457

31. Mercurio V, Pirozzi F, Lazzarini E, Marone G, Rizzo P, Agnetti G, Tocchetti CG, Ghigo A, Ameri P. Models of heart failure based on the cardiotoxicity of anticancer drugs. *Journal of cardiac failure*. 2016;22:449-458
32. Vermeulen Z, Segers VF, De Keulenaer GW. ErbB2 signaling at the crossing between heart failure and cancer. *Basic research in cardiology*. 2016;111:60
33. Wilkinson EL, Sidaway JE, Cross MJ. Cardiotoxic drugs herceptin and doxorubicin inhibit cardiac microvascular endothelial cell barrier formation resulting in increased drug permeability. *Biology open*. 2016;5:1362-1370
34. Kitani T, Ong SG, Lam CK, Rhee JW, Zhang JZ, Oikonomopoulos A, Ma N, Tian L, Lee J, Telli ML, Witteles RM, Sharma A, Sayed N, Wu JC. Human-induced pluripotent stem cell model of trastuzumab-induced cardiac dysfunction in patients with breast cancer. *Circulation*. 2019;139:2451-2465
35. Chintalgattu V, Rees ML, Culver JC, Goel A, Jiffar T, Zhang J, Dunner K, Jr., Pati S, Bankson JA, Pasqualini R, Arap W, Bryan NS, Taegtmeier H, Langley RR, Yao H, Kupferman ME, Entman ML, Dickinson ME, Khakoo AY. Coronary microvascular pericytes are the cellular target of sunitinib malate-induced cardiotoxicity. *Science translational medicine*. 2013;5:187ra169
36. Moslehi J, Minamishima YA, Shi J, Neuberg D, Charytan DM, Padera RF, Signoretti S, Liao R, Kaelin WG, Jr. Loss of hypoxia-inducible factor prolyl hydroxylase activity in cardiomyocytes phenocopies ischemic cardiomyopathy. *Circulation*. 2010;122:1004-1016
37. Bekeredjian R, Walton CB, MacCannell KA, Ecker J, Kruse F, Outten JT, Sutcliffe D, Gerard RD, Bruick RK, Shohet RV. Conditional hif-1alpha expression produces a reversible cardiomyopathy. *PloS one*. 2010;5:e11693
38. Dobbin SJH, Mangion K, Berry C, Roditi G, Basak S, Sourbron S, White J, Venugopal B, Touyz RM, Jones RJ, Petrie MC, Lang NN. Cardiotoxicity and myocardial hypoperfusion associated with anti-vascular endothelial growth factor therapies: Prospective cardiac magnetic resonance imaging in patients with cancer. *European journal of heart failure*. 2020
39. Uraizee I, Cheng S, Moslehi J. Reversible cardiomyopathy associated with sunitinib and sorafenib. *The New England journal of medicine*. 2011;365:1649-1650
40. Kivela R, Hemanthakumar KA, Vaparanta K, Robciuc M, Izumiya Y, Kidoya H, Takakura N, Peng X, Sawyer DB, Elenius K, Walsh K, Alitalo K. Endothelial cells regulate physiological cardiomyocyte growth via vegfr2-mediated paracrine signaling. *Circulation*. 2019;139:2570-2584
41. Chiusa M, Hool SL, Truetsch P, Djafarzadeh S, Jakob SM, Seifriz F, Scherer SJ, Suter TM, Zuppinger C, Zbinden S. Cancer therapy modulates vegf signaling and viability in adult rat cardiac microvascular endothelial cells and cardiomyocytes. *Journal of molecular and cellular cardiology*. 2012;52:1164-1175
42. Tocchetti CG, Gallucci G, Coppola C, Piscopo G, Cipresso C, Maurea C, Giudice A, Iaffaioli RV, Arra C, Maurea N. The emerging issue of cardiac dysfunction induced by antineoplastic angiogenesis inhibitors. *European journal of heart failure*. 2013;15:482-489
43. Sharma A, BurrIDGE PW, McKeithan WL, Serrano R, Shukla P, Sayed N, Churko JM, Kitani T, Wu H, Holmstrom A, Matsa E, Zhang Y, Kumar A, Fan AC, Del Alamo JC, Wu SM, Moslehi JJ, Mercola M, Wu JC. High-throughput screening of tyrosine kinase inhibitor cardiotoxicity with human induced pluripotent stem cells. *Science translational medicine*. 2017;9
44. Paulus WJ, Tschope C. A novel paradigm for heart failure with preserved ejection fraction: Comorbidities drive myocardial dysfunction and remodeling through coronary microvascular endothelial inflammation. *Journal of the American College of Cardiology*. 2013;62:263-271
45. Triposkiadis F, Butler J, Abboud FM, Armstrong PW, Adamopoulos S, Atherton JJ, Backs J, Bauersachs J, Burkhoff D, Bonow RO, Chopra VK, de Boer RA, de Windt L, Hamdani N, Hasenfuss G, Heymans S, Hulot JS, Konstam M, Lee RT, Linke WA, Lunde IG, Lyon AR, Maack C, Mann DL, Mebazaa A, Mentz RJ, Nihoyannopoulos P, Papp Z, Parissis J, Pedrazzini T, Rosano G, Rouleau J, Seferovic PM, Shah AM, Starling RC, Tocchetti CG, Trochu JN, Thum T, Zannad F,

- Brutsaert DL, Segers VF, De Keulenaer GW. The continuous heart failure spectrum: Moving beyond an ejection fraction classification. *European heart journal*. 2019;40:2155-2163
46. Lourenco AP, Leite-Moreira AF, Balligand JL, Bauersachs J, Dawson D, de Boer RA, de Windt LJ, Falcao-Pires I, Fontes-Carvalho R, Franz S, Giacca M, Hilfiker-Kleiner D, Hirsch E, Maack C, Mayr M, Pieske B, Thum T, Tocchetti CG, Brutsaert DL, Heymans S. An integrative translational approach to study heart failure with preserved ejection fraction: A position paper from the working group on myocardial function of the european society of cardiology. *European journal of heart failure*. 2018;20:216-227
 47. Saiki H, Petersen IA, Scott CG, Bailey KR, Dunlay SM, Finley RR, Ruddy KJ, Yan E, Redfield MM. Risk of heart failure with preserved ejection fraction in older women after contemporary radiotherapy for breast cancer. *Circulation*. 2017;135:1388-1396
 48. Saiki H, Moulay G, Guenzel AJ, Liu W, Decklever TD, Classic KL, Pham L, Chen HH, Burnett JC, Russell SJ, Redfield MM. Experimental cardiac radiation exposure induces ventricular diastolic dysfunction with preserved ejection fraction. *American journal of physiology. Heart and circulatory physiology*. 2017;313:H392-H407
 49. Mercurio V, Cuomo A, Della Pepa R, Ciervo D, Cella L, Pirozzi F, Parrella P, Campi G, Franco R, Varricchi G, Abete P, Marone G, Petretta M, Bonaduce D, Pacelli R, Picardi M, Tocchetti CG. What is the cardiac impact of chemotherapy and subsequent radiotherapy in lymphoma patients? *Antioxidants & redox signaling*. 2019;31:1166-1174
 50. Hirsch E, Nagai R, Thum T. Heterocellular signalling and crosstalk in the heart in ischaemia and heart failure. *Cardiovascular research*. 2014;102:191-193
 51. Bang C, Batkai S, Dangwal S, Gupta SK, Foinquinos A, Holzmann A, Just A, Remke J, Zimmer K, Zeug A, Ponimaskin E, Schmiedel A, Yin X, Mayr M, Halder R, Fischer A, Engelhardt S, Wei Y, Schober A, Fiedler J, Thum T. Cardiac fibroblast-derived microRNA passenger strand-enriched exosomes mediate cardiomyocyte hypertrophy. *The Journal of clinical investigation*. 2014;124:2136-2146
 52. Zhan H, Aizawa K, Sun J, Tomida S, Otsu K, Conway SJ, McKinnon PJ, Manabe I, Komuro I, Miyagawa K, Nagai R, Suzuki T. Ataxia telangiectasia mutated in cardiac fibroblasts regulates doxorubicin-induced cardiotoxicity. *Cardiovascular research*. 2016;110:85-95
 53. Meyer K, Hodwin B, Ramanujam D, Engelhardt S, Sarikas A. Essential role for premature senescence of myofibroblasts in myocardial fibrosis. *Journal of the American College of Cardiology*. 2016;67:2018-2028
 54. Sayed N, Ameen M, Wu JC. Personalized medicine in cardio-oncology: The role of induced pluripotent stem cell. *Cardiovascular research*. 2019;115:949-959
 55. Archer CR, Sargeant R, Basak J, Pilling J, Barnes JR, Pointon A. Characterization and validation of a human 3d cardiac microtissue for the assessment of changes in cardiac pathology. *Scientific reports*. 2018;8:10160
 56. Ferdinandy P, Baczkó I, Bencsik P, Giricz Z, Gorbe A, Pacher P, Varga ZV, Varro A, Schulz R. Definition of hidden drug cardiotoxicity: Paradigm change in cardiac safety testing and its clinical implications. *European heart journal*. 2019;40:1771-1777
 57. Ferdinandy P, Hausenloy DJ, Heusch G, Baxter GF, Schulz R. Interaction of risk factors, comorbidities, and comedications with ischemia/reperfusion injury and cardioprotection by preconditioning, postconditioning, and remote conditioning. *Pharmacological reviews*. 2014;66:1142-1174
 58. Perrino C, Barabasi AL, Condorelli G, Davidson SM, De Windt L, Dimmeler S, Engel FB, Hausenloy DJ, Hill JA, Van Laake LW, Lecour S, Leor J, Madonna R, Mayr M, Prunier F, Sluijter JPG, Schulz R, Thum T, Ytrehus K, Ferdinandy P. Epigenomic and transcriptomic approaches in the post-genomic era: Path to novel targets for diagnosis and therapy of the ischaemic heart? Position paper of the european society of cardiology working group on cellular biology of the heart. *Cardiovascular research*. 2017;113:725-736

59. Makkos A, Szantai A, Paloczi J, Pipis J, Kiss B, Poggi P, Ferdinandy P, Chatgialiloglu A, Gorbe A. A comorbidity model of myocardial ischemia/reperfusion injury and hypercholesterolemia in rat cardiac myocyte cultures. *Frontiers in physiology*. 2019;10:1564
60. de Boer RA, Meijers WC, van der Meer P, van Veldhuisen DJ. Cancer and heart disease: Associations and relations. *European journal of heart failure*. 2019;21:1515-1525
61. Genovese G, Kahler AK, Handsaker RE, Lindberg J, Rose SA, Bakhoum SF, Chambert K, Mick E, Neale BM, Fromer M, Purcell SM, Svantesson O, Landen M, Hoglund M, Lehmann S, Gabriel SB, Moran JL, Lander ES, Sullivan PF, Sklar P, Gronberg H, Hultman CM, McCarroll SA. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *The New England journal of medicine*. 2014;371:2477-2487
62. Dorsheimer L, Assmus B, Rasper T, Ortmann CA, Ecke A, Abou-El-Ardat K, Schmid T, Brune B, Wagner S, Serve H, Hoffmann J, Seeger F, Dimmeler S, Zeiher AM, Rieger MA. Association of mutations contributing to clonal hematopoiesis with prognosis in chronic ischemic heart failure. *JAMA cardiology*. 2019;4:25-33
63. Fuster JJ, Walsh K. Somatic mutations and clonal hematopoiesis: Unexpected potential new drivers of age-related cardiovascular disease. *Circulation research*. 2018;122:523-532
64. Jaiswal S, Natarajan P, Silver AJ, Gibson CJ, Bick AG, Shvartz E, McConkey M, Gupta N, Gabriel S, Ardissino D, Baber U, Mehran R, Fuster V, Danesh J, Frossard P, Saleheen D, Melander O, Sukhova GK, Neuberg D, Libby P, Kathiresan S, Ebert BL. Clonal hematopoiesis and risk of atherosclerotic cardiovascular disease. *The New England journal of medicine*. 2017;377:111-121
65. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, Lindsley RC, Mermel CH, Burt N, Chavez A, Higgins JM, Moltchanov V, Kuo FC, Kluk MJ, Henderson B, Kinnunen L, Koistinen HA, Ladenvall C, Getz G, Correa A, Banahan BF, Gabriel S, Kathiresan S, Stringham HM, McCarthy MI, Boehnke M, Tuomilehto J, Haiman C, Groop L, Atzmon G, Wilson JG, Neuberg D, Altshuler D, Ebert BL. Age-related clonal hematopoiesis associated with adverse outcomes. *The New England journal of medicine*. 2014;371:2488-2498
66. Libby P, Sidlow R, Lin AE, Gupta D, Jones LW, Moslehi J, Zeiher A, Jaiswal S, Schulz C, Blankstein R, Bolton KL, Steensma D, Levine RL, Ebert BL. Clonal hematopoiesis: Crossroads of aging, cardiovascular disease, and cancer: Jacc review topic of the week. *Journal of the American College of Cardiology*. 2019;74:567-577
67. Yura Y, Sano S, Walsh K. Clonal hematopoiesis: A new step linking inflammation to heart failure. *JACC. Basic to translational science*. 2020;5:196-207
68. Garcia-Pavia P, Kim Y, Restrepo-Cordoba MA, Lunde IG, Wakimoto H, Smith AM, Toepfer CN, Getz K, Gorham J, Patel P, Ito K, Willcox JA, Arany Z, Li J, Owens AT, Govind R, Nunez B, Mazaika E, Bayes-Genis A, Walsh R, Finkelman B, Lupon J, Whiffin N, Serrano I, Midwinter W, Wilk A, Bardaji A, Ingold N, Buchan R, Tayal U, Pascual-Figal DA, de Marvao A, Ahmad M, Garcia-Pinilla JM, Pantazis A, Dominguez F, John Baksi A, O'Regan DP, Rosen SD, Prasad SK, Lara-Pezzi E, Provencio M, Lyon AR, Alonso-Pulpon L, Cook SA, DePalma SR, Barton PJR, Aplenc R, Seidman JG, Ky B, Ware JS, Seidman CE. Genetic variants associated with cancer therapy-induced cardiomyopathy. *Circulation*. 2019;140:31-41
69. de Boer RA, De Keulenaer G, Bauersachs J, Brutsaert D, Cleland JG, Diez J, Du XJ, Ford P, Heinzl FR, Lipson KE, McDonagh T, Lopez-Andres N, Lunde IG, Lyon AR, Pollesello P, Prasad SK, Tocchetti CG, Mayr M, Sluijter JPG, Thum T, Tschope C, Zannad F, Zimmermann WH, Ruschitzka F, Filippatos G, Lindsey ML, Maack C, Heymans S. Towards better definition, quantification and treatment of fibrosis in heart failure. A scientific roadmap by the committee of translational research of the heart failure association (hfa) of the european society of cardiology. *European journal of heart failure*. 2019;21:272-285
70. Li M, Caeyenberghs K. Longitudinal assessment of chemotherapy-induced changes in brain and cognitive functioning: A systematic review. *Neuroscience and biobehavioral reviews*. 2018;92:304-317

71. Ogren JA, Fonarow GC, Woo MA. Cerebral impairment in heart failure. *Current heart failure reports*. 2014;11:321-329
72. Bertero E, Canepa M, Maack C, Ameri P. Linking heart failure to cancer. *Circulation*. 2018;138:735-742
73. Libby P, Kobold S. Inflammation: A common contributor to cancer, aging, and cardiovascular diseases-expanding the concept of cardio-oncology. *Cardiovascular research*. 2019;115:824-829
74. Prabhu SD, Frangogiannis NG. The biological basis for cardiac repair after myocardial infarction: From inflammation to fibrosis. *Circ Res*. 2016;119:91-112
75. Wang L, Chen Q, Qi H, Wang C, Wang C, Zhang J, Dong L. Doxorubicin-induced systemic inflammation is driven by upregulation of toll-like receptor tlr4 and endotoxin leakage. *Cancer research*. 2016;76:6631-6642
76. Pecoraro M, Del Pizzo M, Marzocco S, Sorrentino R, Ciccarelli M, Iaccarino G, Pinto A, Popolo A. Inflammatory mediators in a short-time mouse model of doxorubicin-induced cardiotoxicity. *Toxicology and applied pharmacology*. 2016;293:44-52
77. Nozaki N, Shishido T, Takeishi Y, Kubota I. Modulation of doxorubicin-induced cardiac dysfunction in toll-like receptor-2-knockout mice. *Circulation*. 2004;110:2869-2874
78. van Almen GC, Swinnen M, Carai P, Verhesen W, Cleutjens JP, D'Hooge J, Verheyen FK, Pinto YM, Schroen B, Carmeliet P, Heymans S. Absence of thrombospondin-2 increases cardiomyocyte damage and matrix disruption in doxorubicin-induced cardiomyopathy. *Journal of molecular and cellular cardiology*. 2011;51:318-328
79. Ma Y, Mattarollo SR, Adjemian S, Yang H, Aymeric L, Hannani D, Portela Catani JP, Duret H, Teng MW, Kepp O, Wang Y, Sistigu A, Schultze JL, Stoll G, Galluzzi L, Zitvogel L, Smyth MJ, Kroemer G. Ccl2/ccr2-dependent recruitment of functional antigen-presenting cells into tumors upon chemotherapy. *Cancer research*. 2014;74:436-445
80. Lesterhuis WJ, Haanen JB, Punt CJ. Cancer immunotherapy--revisited. *Nature reviews. Drug discovery*. 2011;10:591-600
81. Varricchi G, Galdiero MR, Tocchetti CG. Cardiac toxicity of immune checkpoint inhibitors: Cardio-oncology meets immunology. *Circulation*. 2017;136:1989-1992
82. Hu JR, Florido R, Lipson EJ, Naidoo J, Ardehali R, Tocchetti CG, Lyon AR, Padera RF, Johnson DB, Moslehi J. Cardiovascular toxicities associated with immune checkpoint inhibitors. *Cardiovascular research*. 2019;115:854-868
83. Lyon AR, Yousaf N, Battisti NML, Moslehi J, Larkin J. Immune checkpoint inhibitors and cardiovascular toxicity. *The Lancet. Oncology*. 2018;19:e447-e458
84. Escudier M, Cautela J, Malissen N, Ancedy Y, Orabona M, Pinto J, Monestier S, Grob JJ, Scemama U, Jacquier A, Lalevee N, Barraud J, Peyrol M, Laine M, Bonello L, Paganelli F, Cohen A, Barlesi F, Ederhy S, Thuny F. Clinical features, management, and outcomes of immune checkpoint inhibitor-related cardiotoxicity. *Circulation*. 2017;136:2085-2087
85. Nishimura H, Okazaki T, Tanaka Y, Nakatani K, Hara M, Matsumori A, Sasayama S, Mizoguchi A, Hiai H, Minato N, Honjo T. Autoimmune dilated cardiomyopathy in pd-1 receptor-deficient mice. *Science*. 2001;291:319-322
86. Varricchi G, Marone G, Mercurio V, Galdiero MR, Bonaduce D, Tocchetti CG. Immune checkpoint inhibitors and cardiac toxicity: An emerging issue. *Current medicinal chemistry*. 2018;25:1327-1339
87. Awadalla M, Mahmood SS, Groarke JD, Hassan MZO, Nohria A, Rokicki A, Murphy SP, Mercaldo ND, Zhang L, Zlotoff DA, Reynolds KL, Alvi RM, Banerji D, Liu S, Heinzerling LM, Jones-O'Connor M, Bakar RB, Cohen JV, Kirchberger MC, Sullivan RJ, Gupta D, Mulligan CP, Shah SP, Ganatra S, Rizvi MA, Sahni G, Tocchetti CG, Lawrence DP, Mahmoudi M, Devereux RB, Forrester BJ, Mandawat A, Lyon AR, Chen CL, Barac A, Hung J, Thavendiranathan P, Picard MH, Thuny F, Ederhy S, Fradley MG, Neilan TG. Global longitudinal strain and cardiac events in patients with immune checkpoint inhibitor-related myocarditis. *Journal of the American College of Cardiology*. 2020;75:467-478

88. Salem JE, Allenbach Y, Vozy A, Brechot N, Johnson DB, Moslehi JJ, Kerneis M. Abatacept for severe immune checkpoint inhibitor-associated myocarditis. *The New England journal of medicine*. 2019;380:2377-2379
89. Zhang L, Zlotoff DA, Awadalla M, Mahmood SS, Nohria A, Hassan MZO, Thuny F, Zubiri L, Chen CL, Sullivan RJ, Alvi RM, Rokicki A, Murphy SP, Jones-O'Connor M, Heinzerling LM, Barac A, Forrester BJ, Yang EH, Gupta D, Kirchberger MC, Shah SP, Rizvi MA, Sahni G, Mandawat A, Mahmoudi M, Ganatra S, Ederhy S, Zatarain-Nicolas E, Groarke JD, Tocchetti CG, Lyon AR, Thavendiranathan P, Cohen JV, Reynolds KL, Fradley MG, Neilan TG. Major adverse cardiovascular events and the timing and dose of corticosteroids in immune checkpoint inhibitor-associated myocarditis. *Circulation*. 2020;141:2031-2034
90. Ganatra S, Carver JR, Hayek SS, Ky B, Leja MJ, Lenihan DJ, Lenneman C, Mousavi N, Park JH, Perales MA, Ryan TD, Scherrer-Crosbie M, Steingart RM, Yang EH, Zaha V, Barac A, Liu JE. Chimeric antigen receptor t-cell therapy for cancer and heart: Jacc council perspectives. *Journal of the American College of Cardiology*. 2019;74:3153-3163
91. June CH, Sadelain M. Chimeric antigen receptor therapy. *The New England journal of medicine*. 2018;379:64-73
92. Riegler LL, Jones GP, Lee DW. Current approaches in the grading and management of cytokine release syndrome after chimeric antigen receptor t-cell therapy. *Therapeutics and clinical risk management*. 2019;15:323-335
93. Alvi RM, Frigault MJ, Fradley MG, Jain MD, Mahmood SS, Awadalla M, Lee DH, Zlotoff DA, Zhang L, Drobni ZD, Hassan MZO, Bassily E, Rhea I, Ismail-Khan R, Mulligan CP, Banerji D, Lazaryan A, Shah BD, Rokicki A, Raje N, Chavez JC, Abramson J, Locke FL, Neilan TG. Cardiovascular events among adults treated with chimeric antigen receptor t-cells (car-t). *Journal of the American College of Cardiology*. 2019;74:3099-3108
94. Aldoss I, Khaled SK, Budde E, Stein AS. Cytokine release syndrome with the novel treatments of acute lymphoblastic leukemia: Pathophysiology, prevention, and treatment. *Current oncology reports*. 2019;21:4
95. Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature*. 2008;454:436-444
96. Galdiero MR, Garlanda C, Jaillon S, Marone G, Mantovani A. Tumor associated macrophages and neutrophils in tumor progression. *Journal of cellular physiology*. 2013;228:1404-1412
97. Van't Klooster CC, Ridker PM, Hjortnaes J, van der Graaf Y, Asselbergs FW, Westerink J, Aerts J, Visseren FLJ. The relation between systemic inflammation and incident cancer in patients with stable cardiovascular disease: A cohort study. *European heart journal*. 2019;40:3901-3909
98. Ridker PM, MacFadyen JG, Thuren T, Everett BM, Libby P, Glynn RJ, Group CT. Effect of interleukin-1beta inhibition with canakinumab on incident lung cancer in patients with atherosclerosis: Exploratory results from a randomised, double-blind, placebo-controlled trial. *Lancet*. 2017;390:1833-1842
99. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, Fonseca F, Nicolau J, Koenig W, Anker SD, Kastelein JJP, Cornel JH, Pais P, Pella D, Genest J, Cifkova R, Lorenzatti A, Forster T, Kobalava Z, Vida-Simiti L, Flather M, Shimokawa H, Ogawa H, Dellborg M, Rossi PRF, Troquay RPT, Libby P, Glynn RJ, Group CT. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med*. 2017;377:1119-1131
100. Sauter KA, Wood LJ, Wong J, Iordanov M, Magun BE. Doxorubicin and daunorubicin induce processing and release of interleukin-1beta through activation of the nlrp3 inflammasome. *Cancer biology & therapy*. 2011;11:1008-1016
101. Zhu J, Zhang J, Xiang D, Zhang Z, Zhang L, Wu M, Zhu S, Zhang R, Han W. Recombinant human interleukin-1 receptor antagonist protects mice against acute doxorubicin-induced cardiotoxicity. *European journal of pharmacology*. 2010;643:247-253
102. Ghigo A, Damilano F, Braccini L, Hirsch E. Pi3k inhibition in inflammation: Toward tailored therapies for specific diseases. *Bioessays*. 2010;32:185-196

103. Fougerat A, Gayral S, Gourdy P, Schambourg A, Ruckle T, Schwarz MK, Rommel C, Hirsch E, Arnal JF, Salles JP, Perret B, Breton-Douillon M, Wymann MP, Laffargue M. Genetic and pharmacological targeting of phosphoinositide 3-kinase-gamma reduces atherosclerosis and favors plaque stability by modulating inflammatory processes. *Circulation*. 2008;117:1310-1317
104. Damilano F, Franco I, Perrino C, Schaefer K, Azzolino O, Carnevale D, Cifelli G, Carullo P, Ragona R, Ghigo A, Perino A, Lembo G, Hirsch E. Distinct effects of leukocyte and cardiac phosphoinositide 3-kinase gamma activity in pressure overload-induced cardiac failure. *Circulation*. 2011;123:391-399
105. Kaneda MM, Messer KS, Ralainirina N, Li H, Leem CJ, Gorjestani S, Woo G, Nguyen AV, Figueiredo CC, Foubert P, Schmid MC, Pink M, Winkler DG, Rausch M, Palombella VJ, Kutok J, McGovern K, Frazer KA, Wu X, Karin M, Sasik R, Cohen EE, Varner JA. PI3kgamma is a molecular switch that controls immune suppression. *Nature*. 2016;539:437-442
106. Li M, Sala V, De Santis MC, Cimino J, Cappello P, Pianca N, Di Bona A, Margaria JP, Martini M, Lazzarini E, Pirozzi F, Rossi L, Franco I, Bornbaum J, Heger J, Rohrbach S, Perino A, Tocchetti CG, Lima BHF, Teixeira MM, Porporato PE, Schulz R, Angelini A, Sandri M, Ameri P, Sciarretta S, Lima-Junior RCP, Mongillo M, Zaglia T, Morello F, Novelli F, Hirsch E, Ghigo A. Phosphoinositide 3-kinase gamma inhibition protects from anthracycline cardiotoxicity and reduces tumor growth. *Circulation*. 2018;138:696-711
107. Gangadhara G, Dahl G, Bohnacker T, Rae R, Gunnarsson J, Blaho S, Oster L, Lindmark H, Karabelas K, Pemberton N, Tyrchan C, Mogemark M, Wymann MP, Williams RL, Perry MWD, Papavoine T, Petersen J. A class of highly selective inhibitors bind to an active state of pi3kgamma. *Nat Chem Biol*. 2019;15:348-357
108. Anker SD, Egerer KR, Volk HD, Kox WJ, Poole-Wilson PA, Coats AJ. Elevated soluble cd14 receptors and altered cytokines in chronic heart failure. *The American journal of cardiology*. 1997;79:1426-1430
109. Krack A, Sharma R, Figulla HR, Anker SD. The importance of the gastrointestinal system in the pathogenesis of heart failure. *European heart journal*. 2005;26:2368-2374
110. Tang WHW, Li DY, Hazen SL. Dietary metabolism, the gut microbiome, and heart failure. *Nature reviews. Cardiology*. 2019;16:137-154
111. Backhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, Li Y, Xia Y, Xie H, Zhong H, Khan MT, Zhang J, Li J, Xiao L, Al-Aama J, Zhang D, Lee YS, Kotowska D, Colding C, Tremaroli V, Yin Y, Bergman S, Xu X, Madsen L, Kristiansen K, Dahlgren J, Wang J. Dynamics and stabilization of the human gut microbiome during the first year of life. *Cell host & microbe*. 2015;17:690-703
112. Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer N, Knight R. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proceedings of the National Academy of Sciences of the United States of America*. 2010;107:11971-11975
113. Goodrich JK, Waters JL, Poole AC, Sutter JL, Koren O, Blekhman R, Beaumont M, Van Treuren W, Knight R, Bell JT, Spector TD, Clark AG, Ley RE. Human genetics shape the gut microbiome. *Cell*. 2014;159:789-799
114. Kinross JM, Darzi AW, Nicholson JK. Gut microbiome-host interactions in health and disease. *Genome medicine*. 2011;3:14
115. Zitvogel L, Daillere R, Roberti MP, Routy B, Kroemer G. Anticancer effects of the microbiome and its products. *Nature reviews. Microbiology*. 2017;15:465-478
116. Nakatsu G, Li X, Zhou H, Sheng J, Wong SH, Wu WK, Ng SC, Tsoi H, Dong Y, Zhang N, He Y, Kang Q, Cao L, Wang K, Zhang J, Liang Q, Yu J, Sung JJ. Gut mucosal microbiome across stages of colorectal carcinogenesis. *Nature communications*. 2015;6:8727
117. Mitra A, MacIntyre DA, Lee YS, Smith A, Marchesi JR, Lehne B, Bhatia R, Lyons D, Paraskevaidis E, Li JV, Holmes E, Nicholson JK, Bennett PR, Kyrgiou M. Cervical intraepithelial neoplasia

- disease progression is associated with increased vaginal microbiome diversity. *Scientific reports*. 2015;5:16865
118. Yu G, Gail MH, Consonni D, Carugno M, Humphrys M, Pesatori AC, Caporaso NE, Goedert JJ, Ravel J, Landi MT. Characterizing human lung tissue microbiota and its relationship to epidemiological and clinical features. *Genome biology*. 2016;17:163
 119. Karin M, Jobin C, Balkwill F. Chemotherapy, immunity and microbiota--a new triumvirate? *Nature medicine*. 2014;20:126-127
 120. Roy S, Trinchieri G. Microbiota: A key orchestrator of cancer therapy. *Nature reviews. Cancer*. 2017;17:271-285
 121. Russo M, Guida F, Paparo L, Trinchese G, Aitoro R, Avagliano C, Fiordelisi A, Napolitano F, Mercurio V, Sala V, Li M, Sorriento D, Ciccarelli M, Ghigo A, Hirsch E, Bianco R, Iaccarino G, Abete P, Bonaduce D, Calignano A, Berni Canani R, Tocchetti CG. The novel butyrate derivative phenylalanine-butyramide protects from doxorubicin-induced cardiotoxicity. *European journal of heart failure*. 2019;21:519-528
 122. Paparo L, Calignano A, Tocchetti CG, Di Scala C, Russo R, Bonaduce D, Canani RB. The influence of fiber on gut microbiota: Butyrate as molecular player involved in the beneficial interplay between dietary fiber and cardiovascular health. 2017:61-71
 123. Gil-Cruz C, Perez-Shibayama C, De Martin A, Ronchi F, van der Borght K, Niederer R, Onder L, Lutge M, Novkovic M, Nindl V, Ramos G, Arnoldini M, Slack EMC, Boivin-Jahns V, Jahns R, Wyss M, Mooser C, Lambrecht BN, Maeder MT, Rickli H, Flatz L, Eriksson U, Geuking MB, McCoy KD, Ludewig B. Microbiota-derived peptide mimics drive lethal inflammatory cardiomyopathy. *Science*. 2019;366:881-886
 124. Tang WH, Kitai T, Hazen SL. Gut microbiota in cardiovascular health and disease. *Circulation research*. 2017;120:1183-1196
 125. Kim MH, Kang SG, Park JH, Yanagisawa M, Kim CH. Short-chain fatty acids activate gpr41 and gpr43 on intestinal epithelial cells to promote inflammatory responses in mice. *Gastroenterology*. 2013;145:396-406 e391-310
 126. Maslowski KM, Vieira AT, Ng A, Kranich J, Sierro F, Yu D, Schilter HC, Rolph MS, Mackay F, Artis D, Xavier RJ, Teixeira MM, Mackay CR. Regulation of inflammatory responses by gut microbiota and chemoattractant receptor gpr43. *Nature*. 2009;461:1282-1286
 127. Brown AJ, Goldsworthy SM, Barnes AA, Eilert MM, Tcheang L, Daniels D, Muir AI, Wigglesworth MJ, Kinghorn I, Fraser NJ, Pike NB, Strum JC, Steplewski KM, Murdock PR, Holder JC, Marshall FH, Szekeres PG, Wilson S, Ignar DM, Foord SM, Wise A, Dowell SJ. The orphan g protein-coupled receptors gpr41 and gpr43 are activated by propionate and other short chain carboxylic acids. *The Journal of biological chemistry*. 2003;278:11312-11319
 128. Le Poul E, Loison C, Struyf S, Springael JY, Lannoy V, Decobecq ME, Brezillon S, Dupriez V, Vassart G, Van Damme J, Parmentier M, Detheux M. Functional characterization of human receptors for short chain fatty acids and their role in polymorphonuclear cell activation. *The Journal of biological chemistry*. 2003;278:25481-25489
 129. Li L, Hua Y, Ren J. Short-chain fatty acid propionate alleviates akt2 knockout-induced myocardial contractile dysfunction. *Experimental diabetes research*. 2012;2012:851717
 130. Mujumdar VS, Tummalapalli CM, Aru GM, Tyagi SC. Mechanism of constrictive vascular remodeling by homocysteine: Role of ppar. *American journal of physiology. Cell physiology*. 2002;282:C1009-1015
 131. Ali FY, Armstrong PC, Dhanji AR, Tucker AT, Paul-Clark MJ, Mitchell JA, Warner TD. Antiplatelet actions of statins and fibrates are mediated by ppars. *Arteriosclerosis, thrombosis, and vascular biology*. 2009;29:706-711
 132. Antos CL, McKinsey TA, Dreitz M, Hollingsworth LM, Zhang CL, Schreiber K, Rindt H, Gorczynski RJ, Olson EN. Dose-dependent blockade to cardiomyocyte hypertrophy by histone deacetylase inhibitors. *The Journal of biological chemistry*. 2003;278:28930-28937
 133. Gallo P, Latronico MV, Gallo P, Grimaldi S, Borgia F, Todaro M, Jones P, Gallinari P, De Francesco R, Ciliberto G, Steinkuhler C, Esposito G, Condorelli G. Inhibition of class i histone deacetylase

- with an apicidin derivative prevents cardiac hypertrophy and failure. *Cardiovascular research*. 2008;80:416-424
134. Granger A, Abdullah I, Huebner F, Stout A, Wang T, Huebner T, Epstein JA, Gruber PJ. Histone deacetylase inhibition reduces myocardial ischemia-reperfusion injury in mice. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2008;22:3549-3560
 135. Kong Y, Tannous P, Lu G, Berenji K, Rothermel BA, Olson EN, Hill JA. Suppression of class i and ii histone deacetylases blunts pressure-overload cardiac hypertrophy. *Circulation*. 2006;113:2579-2588
 136. Rephaeli A, Rabizadeh E, Aviram A, Shaklai M, Ruse M, Nudelman A. Derivatives of butyric acid as potential anti-neoplastic agents. *International journal of cancer*. 1991;49:66-72
 137. Rephaeli A, Waks-Yona S, Nudelman A, Tarasenko I, Tarasenko N, Phillips DR, Cutts SM, Kessler-Ickson G. Anticancer prodrugs of butyric acid and formaldehyde protect against doxorubicin-induced cardiotoxicity. *British journal of cancer*. 2007;96:1667-1674
 138. Fu HY, Sanada S, Matsuzaki T, Liao Y, Okuda K, Yamato M, Tsuchida S, Araki R, Asano Y, Asanuma H, Asakura M, French BA, Sakata Y, Kitakaze M, Minamino T. Chemical endoplasmic reticulum chaperone alleviates doxorubicin-induced cardiac dysfunction. *Circulation research*. 2016;118:798-809
 139. Daosukho C, Chen Y, Noel T, Sompol P, Nithipongvanitch R, Velez JM, Oberley TD, St Clair DK. Phenylbutyrate, a histone deacetylase inhibitor, protects against adriamycin-induced cardiac injury. *Free radical biology & medicine*. 2007;42:1818-1825
 140. Huang K, Liu Y, Tang H, Qiu M, Li C, Duan C, Wang C, Yang J, Zhou X. Glabridin prevents doxorubicin-induced cardiotoxicity through gut microbiota modulation and colonic macrophage polarization in mice. *Frontiers in pharmacology*. 2019;10:107
 141. Nishimura Y, Kondo C, Morikawa Y, Tonomura Y, Torii M, Yamate J, Uehara T. Plasma mir-208 as a useful biomarker for drug-induced cardiotoxicity in rats. *Journal of applied toxicology : JAT*. 2015;35:173-180
 142. Rigaud VO, Ferreira LR, Ayub-Ferreira SM, Avila MS, Brandao SM, Cruz FD, Santos MH, Cruz CB, Alves MS, Issa VS, Guimaraes GV, Cunha-Neto E, Bocchi EA. Circulating mir-1 as a potential biomarker of doxorubicin-induced cardiotoxicity in breast cancer patients. *Oncotarget*. 2017;8:6994-7002
 143. Gioffre S, Ricci V, Vavassori C, Ruggeri C, Chiesa M, Alfieri I, Zorzan S, Buzzetti M, Milano G, Scopece A, Castiglioni L, Sironi L, Pompilio G, Colombo GI, D'Alessandra Y. Plasmatic and chamber-specific modulation of cardiac micrnas in an acute model of dox-induced cardiotoxicity. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*. 2019;110:1-8
 144. Piegari E, Russo R, Cappetta D, Esposito G, Urbanek K, Dell'Aversana C, Altucci L, Berrino L, Rossi F, De Angelis A. MicroRNA-34a regulates doxorubicin-induced cardiotoxicity in rat. *Oncotarget*. 2016;7:62312-62326
 145. Desai VG, J CK, Vijay V, Moland CL, Herman EH, Lee T, Han T, Lewis SM, Davis KJ, Muskhelishvili L, Kerr S, Fuscoe JC. Early biomarkers of doxorubicin-induced heart injury in a mouse model. *Toxicology and applied pharmacology*. 2014;281:221-229
 146. Horak M, Novak J, Bienertova-Vasku J. Muscle-specific micrnas in skeletal muscle development. *Developmental biology*. 2016;410:1-13
 147. Gomes CP, Oliveira GP, Jr., Madrid B, Almeida JA, Franco OL, Pereira RW. Circulating mir-1, mir-133a, and mir-206 levels are increased after a half-marathon run. *Biomarkers : biochemical indicators of exposure, response, and susceptibility to chemicals*. 2014;19:585-589
 148. Navickas R, Gal D, Laucevicus A, Taparauskaite A, Zdanyte M, Holvoet P. Identifying circulating micrnas as biomarkers of cardiovascular disease: A systematic review. *Cardiovascular research*. 2016;111:322-337
 149. Coenen-Stass AML, Wood MJA, Roberts TC. Biomarker potential of extracellular mirnas in duchenne muscular dystrophy. *Trends in molecular medicine*. 2017;23:989-1001

150. Ruggeri C, Gioffre S, Chiesa M, Buzzetti M, Milano G, Scopece A, Castiglioni L, Pontremoli M, Sironi L, Pompilio G, Colombo GI, D'Alessandra Y. A specific circulating microRNA cluster is associated to late differential cardiac response to doxorubicin-induced cardiotoxicity in vivo. *Disease markers*. 2018;2018:8395651
151. Freres P, Bouznad N, Servais L, Josse C, Wenric S, Poncin A, Thiry J, Moonen M, Oury C, Lancellotti P, Bours V, Jerusalem G. Variations of circulating cardiac biomarkers during and after anthracycline-containing chemotherapy in breast cancer patients. *BMC cancer*. 2018;18:102
152. Fiedler J, Gupta SK, Thum T. Identification of cardiovascular microRNA targetomes. *Journal of molecular and cellular cardiology*. 2011;51:674-681
153. Yin Z, Zhao Y, Li H, Yan M, Zhou L, Chen C, Wang DW. Mir-320a mediates doxorubicin-induced cardiotoxicity by targeting vegf signal pathway. *Aging*. 2016;8:192-207
154. Chatterjee S, Gupta SK, Bar C, Thum T. Noncoding rnas: Potential regulators in cardioncology. *American journal of physiology. Heart and circulatory physiology*. 2019;316:H160-H168
155. Gupta SK, Garg A, Avramopoulos P, Engelhardt S, Streckfuss-Bomeke K, Batkai S, Thum T. Mir-212/132 cluster modulation prevents doxorubicin-mediated atrophy and cardiotoxicity. *Molecular therapy : the journal of the American Society of Gene Therapy*. 2019;27:17-28
156. Gupta SK, Garg A, Bar C, Chatterjee S, Foinquinos A, Milting H, Streckfuss-Bomeke K, Fiedler J, Thum T. Quaking inhibits doxorubicin-mediated cardiotoxicity through regulation of cardiac circular rna expression. *Circulation research*. 2018;122:246-254
157. Roca-Alonso L, Castellano L, Mills A, Dabrowska AF, Sikkel MB, Pellegrino L, Jacob J, Frampton AE, Krell J, Coombes RC, Harding SE, Lyon AR, Stebbing J. Myocardial mir-30 downregulation triggered by doxorubicin drives alterations in beta-adrenergic signaling and enhances apoptosis. *Cell death & disease*. 2015;6:e1754
158. Oatmen KE, Toro-Salazar OH, Hauser K, Zellars KN, Mason KC, Hor K, Gillan E, Zeiss CJ, Gatti DM, Spinale FG. Identification of a novel microRNA profile in pediatric patients with cancer treated with anthracycline chemotherapy. *American journal of physiology. Heart and circulatory physiology*. 2018;315:H1443-H1452
159. Bae S, Ulrich CM, Neuhaus ML, Malysheva O, Bailey LB, Xiao L, Brown EC, Cushing-Haugen KL, Zheng Y, Cheng TY, Miller JW, Green R, Lane DS, Beresford SA, Caudill MA. Plasma choline metabolites and colorectal cancer risk in the women's health initiative observational study. *Cancer research*. 2014;74:7442-7452
160. Mondul AM, Moore SC, Weinstein SJ, Karoly ED, Sampson JN, Albanes D. Metabolomic analysis of prostate cancer risk in a prospective cohort: The alpha-tocolpherol, beta-carotene cancer prevention (atbc) study. *International journal of cancer*. 2015;137:2124-2132
161. Wang C, Jing Q. Non-coding rnas as biomarkers for acute myocardial infarction. *Acta pharmacologica Sinica*. 2018;39:1110-1119
162. Chistiakov DA, Orekhov AN, Bobryshev YV. Cardiac-specific mirna in cardiogenesis, heart function, and cardiac pathology (with focus on myocardial infarction). *Journal of molecular and cellular cardiology*. 2016;94:107-121
163. Kura B, Kalocayova B, Devaux Y, Bartekova M. Potential clinical implications of mir-1 and mir-21 in heart disease and cardioprotection. *International journal of molecular sciences*. 2020;21
164. Kashyap D, Kaur H. Cell-free mirnas as non-invasive biomarkers in breast cancer: Significance in early diagnosis and metastasis prediction. *Life sciences*. 2020;246:117417
165. Sheervalilou R, Lotfi H, Shirvaliloo M, Sharifi A, Nazemiyeh M, Zarghami N. Circulating mir-10b, mir-1 and mir-30a expression profiles in lung cancer: Possible correlation with clinicopathologic characteristics and lung cancer detection. *International journal of molecular and cellular medicine*. 2019;8:118-129
166. Peng J, Yuan C, Wu Z, Wang Y, Yin W, Lin Y, Zhou L, Lu J. Upregulation of microRNA1 inhibits proliferation and metastasis of breast cancer. *Molecular medicine reports*. 2020;22:454-464

167. Han C, Shen JK, Hornicek FJ, Kan Q, Duan Z. Regulation of microRNA-1 (mir-1) expression in human cancer. *Biochimica et biophysica acta. Gene regulatory mechanisms*. 2017;1860:227-232
168. Fulzele S, Mendhe B, Khayrullin A, Johnson M, Kaiser H, Liu Y, Isaacs CM, Hamrick MW. Muscle-derived mir-34a increases with age in circulating extracellular vesicles and induces senescence of bone marrow stem cells. *Aging*. 2019;11:1791-1803
169. Gioffre S, Chiesa M, Cardinale DM, Ricci V, Vavassori C, Cipolla CM, Masson S, Sandri MT, Salvatici M, Ciceri F, Latini R, Staszewsky LI, Pompilio G, Colombo GI, D'Alessandra Y. Circulating microRNAs as potential predictors of anthracycline-induced troponin elevation in breast cancer patients: Diverging effects of doxorubicin and epirubicin. *Journal of clinical medicine*. 2020;9
170. Boon RA, Iekushi K, Lechner S, Seeger T, Fischer A, Heydt S, Kaluza D, Treguer K, Carmona G, Bonauer A, Horrevoets AJ, Didier N, Girmatsion Z, Biliczki P, Ehrlich JR, Katus HA, Muller OJ, Potente M, Zeiher AM, Hermeking H, Dimmeler S. MicroRNA-34a regulates cardiac ageing and function. *Nature*. 2013;495:107-110
171. Zhu JN, Fu YH, Hu ZQ, Li WY, Tang CM, Fei HW, Yang H, Lin QX, Gou DM, Wu SL, Shan ZX. Activation of mir-34a-5p/sirt1/p66shc pathway contributes to doxorubicin-induced cardiotoxicity. *Scientific reports*. 2017;7:11879
172. Lin Y, Lin Z, Fang Z, Li H, Zhi X, Zhang Z. Plasma microRNA-34a as a potential biomarker for early diagnosis of esophageal cancer. *Clinical laboratory*. 2019;65
173. Zhang L, Liao Y, Tang L. MicroRNA-34 family: A potential tumor suppressor and therapeutic candidate in cancer. *Journal of experimental & clinical cancer research : CR*. 2019;38:53
174. Tokumaru Y, Katsuta E, Oshi M, Sporn JC, Yan L, Le L, Matsuhashi N, Futamura M, Akao Y, Yoshida K, Takabe K. High expression of mir-34a associated with less aggressive cancer biology but not with survival in breast cancer. *International journal of molecular sciences*. 2020;21
175. Sommariva E, D'Alessandra Y, Farina FM, Casella M, Cattaneo F, Catto V, Chiesa M, Stadiotti I, Brambilla S, Dello Russo A, Carbuicchio C, Vettor G, Riggio D, Sandri MT, Barbuti A, Vernillo G, Muratori M, Dal Ferro M, Sinagra G, Moimas S, Giacca M, Colombo GI, Pompilio G, Tondo C. Mir-320a as a potential novel circulating biomarker of arrhythmogenic cardiomyopathy. *Scientific reports*. 2017;7:4802
176. Galeano-Otero I, Del Toro R, Guisado A, Diaz I, Mayoral-Gonzalez I, Guerrero-Marquez F, Gutierrez-Carretero E, Casquero-Dominguez S, Diaz-de la Llera L, Baron-Esquivias G, Jimenez-Navarro M, Smani T, Ordonez-Fernandez A. Circulating mir-320a as a predictive biomarker for left ventricular remodelling in stemi patients undergoing primary percutaneous coronary intervention. *Journal of clinical medicine*. 2020;9
177. Bostjancic E, Zidar N, Glavac D. MicroRNAs and cardiac sarcoplasmic reticulum calcium atpase-2 in human myocardial infarction: Expression and bioinformatic analysis. *BMC genomics*. 2012;13:552
178. Vila-Navarro E, Duran-Sanchon S, Vila-Casadesus M, Moreira L, Gines A, Cuatrecasas M, Lozano JJ, Bujanda L, Castells A, Gironella M. Novel circulating miRNA signatures for early detection of pancreatic neoplasia. *Clinical and translational gastroenterology*. 2019;10:e00029
179. Sole C, Tramonti D, Schramm M, Goicoechea I, Armesto M, Hernandez LI, Manterola L, Fernandez-Mercado M, Mujika K, Tuneu A, Jaka A, Tellaetxe M, Friedlander MR, Estivill X, Piazza P, Ortiz-Romero PL, Middleton MR, Lawrie CH. The circulating transcriptome as a source of biomarkers for melanoma. *Cancers*. 2019;11
180. Zhao W, Sun Q, Yu Z, Mao S, Jin Y, Li J, Jiang Z, Zhang Y, Chen M, Chen P, Chen D, Xu H, Ding S, Yu Z. Mir-320a-3p/elf3 axis regulates cell metastasis and invasion in non-small cell lung cancer via pi3k/akt pathway. *Gene*. 2018;670:31-37
181. Ge X, Cui H, Zhou Y, Yin D, Feng Y, Xin Q, Xu X, Liu W, Liu S, Zhang Q. Mir-320a modulates cell growth and chemosensitivity via regulating adam10 in gastric cancer. *Molecular medicine reports*. 2017;16:9664-9670

182. Jeong HS, Kim JY, Lee SH, Hwang J, Shin JW, Song KS, Lee S, Kim J. Synergy of circulating mir-212 with markers for cardiovascular risks to enhance estimation of atherosclerosis presence. *PloS one*. 2017;12:e0177809
183. Masson S, Batkai S, Beermann J, Bar C, Pfanne A, Thum S, Magnoli M, Balconi G, Nicolosi GL, Tavazzi L, Latini R, Thum T. Circulating microRNA-132 levels improve risk prediction for heart failure hospitalization in patients with chronic heart failure. *European journal of heart failure*. 2018;20:78-85
184. Ucar A, Gupta SK, Fiedler J, Erikci E, Kardasinski M, Batkai S, Dangwal S, Kumarswamy R, Bang C, Holzmann A, Remke J, Caprio M, Jentsch C, Engelhardt S, Geisendorf S, Glas C, Hofmann TG, Nessling M, Richter K, Schiffer M, Carrier L, Napp LC, Bauersachs J, Chowdhury K, Thum T. The mirna-212/132 family regulates both cardiac hypertrophy and cardiomyocyte autophagy. *Nature communications*. 2012;3:1078
185. Duell EJ, Lujan-Barroso L, Sala N, Deitz McElyea S, Overvad K, Tjonneland A, Olsen A, Weiderpass E, Busund LT, Moi L, Muller D, Vineis P, Aune D, Matullo G, Naccarati A, Panico S, Tagliabue G, Tumino R, Palli D, Kaaks R, Katzke VA, Boeing H, Bueno-de-Mesquita HBA, Peeters PH, Trichopoulou A, Lagiou P, Kotanidou A, Travis RC, Wareham N, Khaw KT, Ramon Quiros J, Rodriguez-Barranco M, Dorronsoro M, Chirlaque MD, Ardanaz E, Severi G, Boutron-Ruault MC, Rebours V, Brennan P, Gunter M, Scelo G, Cote G, Sherman S, Korc M. Plasma microRNAs as biomarkers of pancreatic cancer risk in a prospective cohort study. *International journal of cancer*. 2017;141:905-915
186. Weber DG, Gawrych K, Casjens S, Brik A, Lehnert M, Taeger D, Pesch B, Kollmeier J, Bauer TT, Johnen G, Bruning T. Circulating mir-132-3p as a candidate diagnostic biomarker for malignant mesothelioma. *Disease markers*. 2017;2017:9280170
187. Jiang X, Chen X, Chen L, Ma Y, Zhou L, Qi Q, Liu Y, Zhang S, Luo J, Zhou X. Upregulation of the mir-212/132 cluster suppresses proliferation of human lung cancer cells. *Oncology reports*. 2015;33:705-712
188. Lin L, Wang Z, Jin H, Shi H, Lu Z, Qi Z. Mir-212/132 is epigenetically downregulated by sox4/ezh2-h3k27me3 feedback loop in ovarian cancer cells. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*. 2016
189. Oliosio D, Dauriz M, Bacchi E, Negri C, Santi L, Bonora E, Moghetti P. Effects of aerobic and resistance training on circulating micro-rna expression profile in subjects with type 2 diabetes. *The Journal of clinical endocrinology and metabolism*. 2019;104:1119-1130
190. Li J, Wan W, Chen T, Tong S, Jiang X, Liu W. Mir-451 silencing inhibited doxorubicin exposure-induced cardiotoxicity in mice. *BioMed research international*. 2019;2019:1528278
191. Song L, Su M, Wang S, Zou Y, Wang X, Wang Y, Cui H, Zhao P, Hui R, Wang J. Mir-451 is decreased in hypertrophic cardiomyopathy and regulates autophagy by targeting tsc1. *Journal of cellular and molecular medicine*. 2014;18:2266-2274
192. Bai H, Wu S. Mir-451: A novel biomarker and potential therapeutic target for cancer. *OncoTargets and therapy*. 2019;12:11069-11082
193. Shen Y, Ding Y, Ma Q, Zhao L, Guo X, Shao Y, Niu C, He Y, Zhang F, Zheng D, Wei W, Liu F. Identification of novel circulating mirna biomarkers for the diagnosis of esophageal squamous cell carcinoma and squamous dysplasia. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2019;28:1212-1220
194. Wang Q, Shang J, Zhang Y, Zhou Y, Tang L. Mir-451a restrains the growth and metastatic phenotypes of papillary thyroid carcinoma cells via inhibiting zeb1. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*. 2020;127:109901
195. Dong Y, Wang G. Knockdown of lncrna snhg12 suppresses cell proliferation, migration and invasion in breast cancer by sponging mir-451a. *International journal of clinical and experimental pathology*. 2020;13:393-402

196. Kaur A, Mackin ST, Schlosser K, Wong FL, Elharram M, Delles C, Stewart DJ, Dayan N, Landry T, Pilote L. Systematic review of microRNA biomarkers in acute coronary syndrome and stable coronary artery disease. *Cardiovascular research*. 2020;116:1113-1124
197. Pereira-da-Silva T, Coutinho Cruz M, Carrusca C, Cruz Ferreira R, Napoleao P, Mota Carmo M. Circulating microRNA profiles in different arterial territories of stable atherosclerotic disease: A systematic review. *American journal of cardiovascular disease*. 2018;8:1-13
198. Lai L, Chen J, Wang N, Zhu G, Duan X, Ling F. Mirna-30e mediated cardioprotection of ace2 in rats with doxorubicin-induced heart failure through inhibiting cardiomyocytes autophagy. *Life sciences*. 2017;169:69-75
199. Zhang X, Dong S, Jia Q, Zhang A, Li Y, Zhu Y, Lv S, Zhang J. The microRNA in ventricular remodeling: The mir-30 family. *Bioscience reports*. 2019;39
200. Shi M, Mu Y, Zhang H, Liu M, Wan J, Qin X, Li C. MicroRNA-200 and microRNA-30 family as prognostic molecular signatures in ovarian cancer: A meta-analysis. *Medicine*. 2018;97:e11505
201. Wang C, Chen L, Yang Y, Zhang M, Wong G. Identification of bladder cancer prognostic biomarkers using an ageing gene-related competitive endogenous rna network. *Oncotarget*. 2017;8:111742-111753
202. Han W, Cui H, Liang J, Su X. Role of microRNA-30c in cancer progression. *Journal of Cancer*. 2020;11:2593-2601
203. Croset M, Pantano F, Kan CWS, Bonnelye E, Descotes F, Alix-Panabieres C, Lecellier CH, Bachelier R, Allioli N, Hong SS, Bartkowiak K, Pantel K, Clezardin P. Mirna-30 family members inhibit breast cancer invasion, osteomimicry, and bone destruction by directly targeting multiple bone metastasis-associated genes. *Cancer research*. 2018;78:5259-5273

Figure legend.

Figure 1

Besides directly affecting cardiomyocytes and the other cardiac cell populations, cancer treatments may disrupt the intercellular communications between cardiomyocytes and non-cardiomyocytes. The inset in the lower part of the Figure shows key endothelial cell (green)-cardiomyocyte (red) paracrine signaling axes that may be impaired by antitumor therapies.

NRG-1: neuregulin-1; VEGF: vascular endothelial growth factor; VEGFR2: vascular endothelial growth factor receptor type 2; NO: nitric oxide; cGMP: cyclic guanosine monophosphate; PKG: protein kinase G

Figure 2

Inflammation at the intersection of the anti-cancer action and cardiac side effects of major oncological treatments.

Besides directly killing tumor cells, doxorubicin triggers cardiac inflammation via activation of macrophages and fibroblasts and the ensuing release of local modulators of inflammation and fibrosis, such as TNF- α , IL-1 β and IL-6. Major players of the inflammatory response induced by doxorubicin include macrophage TLR-4, the matricellular protein thrombospondin-2 (TSP-2) and leukocyte PI3K γ . On the other hand, immune check point inhibitors (ICIs) inhibit molecules such as cytotoxic-T-lymphocyte-associated antigen 4 (CTLA-4), programmed cell death 1 (PD-1) and its ligand PD-L1. As a consequence, anti-tumor immune cell responses are reactivated and lead to tumor cell death, but concomitantly drives myocarditis. Although these new immunotherapies have notable anti-cancer effects, multiple mechanisms of immune

resistance exist, and these might be overcome by using PI3K γ inhibitors that re-shape the tumor immune microenvironment. Finally, engineered T cells with chimeric antigen receptors (CAR-T cells) boosts T cell-mediated tumor killing, but are burdened by cytokine release syndrome (CRS) leading to extremely serious complications, including cardiac and vascular dysfunction, and ultimately to multiorgan failure.

Figure 3

GUT Microbiome Dysbiosis can be influenced by both HF and cancer. HF has long been associated with congestion of splanchnic circulation, leading to bowel wall edema, impaired intestinal barrier function and increased systemic inflammation, that drastically affect GUT microbiome composition and response to HF treatments. At the same time, cancer-mediated disruption of metabolism and the production of cancer-derived metabolites modifies the microbiome. Such altered gut microbiome generates cardiotoxic metabolites such as TMAO and Bile Acids, eventually leading to HF worsening.

Table 1. Main direct toxic effects of cancer therapies on cardiomyocytes

Cellular toxicity	Treatment(s) most commonly involved
Type II topoisomerase poisoning *	Anthracyclines
Mitochondrial dysfunction	Anthracyclines, VEGFR / multitargeted RTK inhibitors
Oxidative stress	Anthracyclines
Impaired autophagy	Anthracyclines, proteasome inhibitors
Altered protein handling	Proteasome inhibitors
Induction of HIF pathways	VEGFR / multitargeted RTK inhibitors

* This toxicity is peculiar of anthracyclines.

VEGFR: vascular endothelial growth factor receptor; RTK: receptor tyrosine kinase; HIF: hypoxia-inducible factor.

Table 2. Cardiovascular toxicities of cancer therapies

Type of toxicity	Treatment(s) most commonly involved
LVD, HF	Anthracyclines, HER2-targeting drugs, VEGFR / multitargeted RTK inhibitors, proteasome inhibitors, radiation therapy (HFpEF)
Myocardial ischemia	Fluoropyrimidines, VEGFR inhibitors, radiation therapy
Myocarditis	ICIs, cyclophosphamide (rarely)
Atrial fibrillation	Ibrutinib
QT prolongation	Arsenic trioxide, vandetanib, androgen deprivation therapy (enzalutamide)
Valvular heart disease	radiation therapy
Pericarditis	ICIs, cyclophosphamide
Hypertension	VEGFR inhibitors
Peripheral artery disease	Nilotinib, ponatinib
Vascular thrombosis *	Cisplatin, nilotinib, ponatinib, thalidomide and lenalidomide, VEGFR inhibitors, proteasome inhibitors, aromatase inhibitors
Pulmonary arterial hypertension	Dasatinib, cyclophosphamide

* Acute myocardial ischemia will ensue if thrombosis occurs at coronary artery atherosclerotic plaques.

LVD: left ventricular dysfunction; HF: heart failure; VEGFR: vascular endothelial growth factor receptor; RTK: receptor tyrosine kinase; HFpEF: heart failure with preserved ejection fraction; ICIs: immune checkpoint inhibitors

Table 3**Role of TMAO and Butyrate in Cardio-Oncology**

Cardiovascular field		Cancer field	
TMAO	Butyrate	TMAO	Butyrate
Prognostic 5 years follow-up marker in patients with heart failure.			
Prognostic biomarker in chronic systolic HF.	Inhibition of maladaptive hypertrophy and heart failure ^{132, 133, 135}		
Predictive biomarker in patients with Acute Heart Failure	Reduction of myocardial ischemia-reperfusion injury ¹³⁴⁾ Cardioprotective action against DOXO toxicity ¹³⁷	Predictive biomarker of colorectal cancer ¹⁵⁹⁾	Induction of cytodifferentiation and inhibition leukemic cells proliferation; inhibition of Lewis lung carcinoma cells growth ¹³⁶
Prognostic biomarker in patients associated with NYHA III and IV ischaemic aetiology and adverse outcomes	Alleviation of DOXO-induced ER stress ^{138,139)} Prevention of DOXO-induced mitochondrial dysfunction and ROS/RNS production ¹²¹	Predictive biomarker of aggressive prostate cancer ¹⁶⁰	Increase of the antineoplastic effect of DOXO ¹³⁷
Predictive biomarker for mortality and CV mortality in HFrEF but not HFpEF patients ^{110*}			

*High levels of TMAO in patients have been suggested to be predictive of cardiovascular events of mortality, independently from renal function and cardiovascular comorbidities¹¹⁰.

		Cardiovascular field		Cancer field	
		Circulating forms (biomarkers)	Intracellular forms (epigenetic activity)	Circulating forms (biomarkers)	Intracellular forms (epigenetic activity)
miR-1	<p>Marker Of Myocardial Infarction.¹⁶¹;</p> <p>Biomarker Of Atherosclerosis, Coronary Artery Disease, Acute Coronary Syndrome.¹⁴⁸</p>	<p>Involved In Cardiogenesis, Heart Function, Cardiac Pathology¹⁶²</p> <p>Involved In Heart Disease And Cardioprotection.¹⁶³</p>	<p>Non-Invasive Biomarkers In Breast Cancer: Early Diagnosis And Metastasis Prediction¹⁶⁴</p> <p>Correlation With Clinico-Pathologic Characteristics And Lung Cancer Detection.¹⁶⁵</p>	<p>Inhibition Of Proliferation And Metastasis Of Breast Cancer.¹⁶⁶;</p> <p>Differential Expression In Different Human Cancers.¹⁶⁷</p>	
miR-34a	<p>Biomarker Of Aging¹⁶⁸</p> <p>Marker Of Anthracycline Treatment¹⁶⁹</p>	<p>Regulation Of Cardiac Ageing And Function.¹⁷⁰</p> <p>Contribution To Doxorubicin-</p>	<p>Non-Invasive Biomarkers In Breast Cancer: Early Diagnosis And Metastasis Prediction¹⁶⁴</p>	<p>Potential Tumor Suppressor And Therapeutic Candidate In Cancer¹⁷³</p>	

		Induced Cardiotoxicity. ¹⁷¹	Potential Biomarker For Early Diagnosis Of Esophageal Cancer ¹⁷²	Associated With Aggressive Breast Cancer ¹⁷⁴
miR-320a	Biomarker Of Arrhythmogenic Cardiomyopathy ¹⁷⁵ Predictive Biomarker For Left Ventricular Remodelling ¹⁷⁶	Mediator Of Doxorubicin- Induced Cardiotoxicity ¹⁵³ Involved In Human Myocardial Infarction ¹⁷⁷	Early Detection Of Pancreatic Neoplasia ¹⁷⁸ Circulating Biomarker Of Melanoma ¹⁷⁹	Regulation Of Cell Metastasis And Invasion In Non-Small Cell Lung Cancer ¹⁸⁰ Modulation Of Cell Growth And Chemosensitivity In Gastric Cancer ¹⁸¹
miR-212/132 cluster	Estimation Of Atherosclerosis Presence ¹⁸²	Prevention Of Doxorubicin- Mediated Atrophy And Cardiotoxicity ¹⁵⁵	Biomarker Of Pancreatic Cancer Risk ¹⁸⁵	Suppression Of Proliferation Of Human Lung Cancer Cells ¹⁸⁷

	Risk Prediction For Heart Failure ¹⁸³	Regulation Of Cardiac Hypertrophy And Cardiomyocyte Autophagy ¹⁸⁴	Diagnostic Biomarker Of Malignant Mesothelioma ¹⁸⁶	Regulated In Ovarian Cancer Cells ¹⁸⁸
miR-451a	Modulated In Acute Model Of Doxorubicin-Induced Cardiotoxicity ¹⁴³ Marker Of Type 2 Diabetes ¹⁸⁹	Regulation Of Doxorubicin-Induced Cardiotoxicity ¹⁹⁰ Decreased In Hypertrophic Cardiomyopathy And Regulates Autophagy ¹⁹¹	Biomarker And Potential Therapeutic Target For Cancer ¹⁹² Biomarker For The Diagnosis Of Esophageal Squamous Cell Carcinoma And Squamous Dysplasia ¹⁹³	Regulation Of Papillary Thyroid Carcinoma Cells ¹⁹⁴ Involved In Breast Cancer ¹⁹⁵
miR-30	Biomarkers Of Acute Coronary Syndrome And Stable Coronary Artery Disease ¹⁹⁶	Cardioprotection In Doxorubicin-Induced Heart Failure And Inhibition Of Cardiomyocytes Autophagy ¹⁹⁸	Prognosis Of Ovarian Cancer ²⁰⁰ Prognosis Of Bladder Cancer ²⁰¹	Involved In Cancer Progression ²⁰² Involved In Breast Cancer

	<p>Marker Of Stable Atherosclerotic Disease¹⁹⁷</p>	<p>Involvement In Ventricular Remodeling: The Mir-30 Family¹⁹⁹</p>		<p>Invasion, Osteomimicry, And Bone Destruction²⁰³</p>
--	---	---	--	---

1311





