

# Cardiovascular health, traffic-related air pollution and noise: are associations mutually confounded? A systematic review

Louis-François Tétreault · Stéphane Perron ·  
Audrey Smargiassi

Received: 31 March 2013 / Revised: 25 June 2013 / Accepted: 27 June 2013 / Published online: 26 July 2013  
© The Author(s) 2013. This article is published with open access at Springerlink.com

## Abstract

**Objectives** This review assessed the confounding effect of one traffic-related exposure (noise or air pollutants) on the association between the other exposure and cardiovascular outcomes.

**Methods** A systematic review was conducted with the databases Medline and Embase. The confounding effects in studies were assessed by using change in the estimate with a 10 % cutoff point. The influence on the change in the estimate of the quality of the studies, the exposure assessment methods and the correlation between road noise and air pollutions were also assessed.

**Results** Nine publications were identified. For most studies, the specified confounders produced changes in estimates <10 %. The correlation between noise and pollutants, the quality of the study and of the exposure assessment do not seem to influence the confounding effects.

**Conclusions** Results from this review suggest that confounding of cardiovascular effects by noise or air pollutants is low, though with further improvements in exposure assessment, the situation may change. More studies using pollution indicators specific to road traffic are needed to properly assess if noise and air pollution are subjected to confounding.

**Keywords** Noise · Air pollution · Confounding · Cardiovascular

## Introduction

Studies have shown that exposure to road traffic noise and air pollutant emissions can lead to adverse health effects such as annoyance (World Health Organisation (2011), sleep disturbance (Pirrera et al. 2010), respiratory problems (Health Effects Institute Panel on the Health Effects of Traffic-Related Air Pollution 2010) and cancer (Beelen et al. 2008). A few studies (Finkelstein et al. 2004; Gan et al. 2010; Hoffmann et al. 2006, 2009; Maheswaran and Elliott 2003) have also reported associations with cardiovascular (CV) outcomes. CV health effects could be explained either by noise or air pollution associated with traffic.

On the one hand, there is evidence linking traffic noise to ischemic heart diseases (World Health Organisation 2011; Babisch and Kamp 2009; Babisch 2006) and hypertension (World Health Organisation Regional Office for Europe 2011; van Kempen and Babisch 2012). On the other hand, there is also evidence linking traffic-generated air pollution to CV diseases (Hoek et al. 2002; Brook et al. 2010). However, in a recent report of the Health Effects Institute Panel on the Health Effects of Traffic-Related Air Pollution (2010), the evidence of the effects of traffic-

---

This article is part of the special issue: "Environment and Health Reviews".

---

L.-F. Tétreault · A. Smargiassi (✉)  
Département de santé environnementale et santé au travail,  
University of Montreal, 2375 Chemin de la Côte-Sainte-  
Catherine, Montréal, QC H3T 1A8, Canada  
e-mail: asmargia@santepub-mtl.qc.ca

L.-F. Tétreault  
e-mail: ltetreau@santepub-mtl.qc.ca

S. Perron  
Département de médecine sociale et préventive,  
University of Montreal, 1301 Sherbrooke est,  
Montreal, QC H2L 1M3, Canada  
e-mail: sperron@santepub-mtl.qc.ca

S. Perron  
Direction de santé publique de l'Agence de la santé et des  
services sociaux de Montréal, Québec, Canada

related air pollution on cardiovascular mortality was considered suggestive, but not sufficient. Nonetheless, there is increasing evidence connecting air pollution to overall cardiovascular mortality (Chen et al. 2013), myocardial infarction (Nuvolone et al. 2011; Rosenlund et al. 2006), atherosclerosis (Adar et al. 2013) and atrial fibrillation (Liao et al. 2010). Although noise or air pollution can confound the effect of each other, the underlying physiological mechanisms are likely different.

A few mechanisms are postulated to explain the effect of noise on CV diseases. Noise can act as a general stressor disturbing the body homeostasis through the “stress syndrome” (Babisch and Kamp 2009; Westman and Walters 1981; World Health Organisation Regional Office for Europe 2011; Amato et al. 2010; Gan et al. 2012). Noise can induce stress by two different pathways. The stress response generated by the direct pathway consists of a neural activation by the noise. In the indirect pathway, the activation of the stress response is created by a cognitive interpretation of the noise (Westman and Walters 1981; Gan et al. 2012). The activation of this pathway can be influenced by the perception of the noise, the perceived control over the sound and noise sensitivity of the subjects (European Environment Agency (EEA) 2010). There are also pathways by which noise could increase the risk of cardiovascular diseases. One of those is the disruption of sleeping patterns. Studies have associated the lack of reduction of blood pressure (BP) during the night (“BP dipping”) caused by noise with an increased risk of cardiovascular outcomes (Vardeny et al. 2011; Haralabidis et al. 2011). It is also suggested that short sleep durations may result in higher ghrelin and lower leptin concentrations (Taheri et al. 2004). The deregulation of those hormones linked with appetite regulation could potentially lead to obesity and higher risk of cardiovascular diseases.

For air pollution, several pathways are hypothesized to explain its impact on CV diseases. Firstly, air pollutants could generate an increase in lung oxidative stress and inflammation. Some pollutants could also migrate through the pulmonary epithelium into blood. Those pathways could lead to a systemic and vascular inflammation, increasing the risk of hypertension and thrombogenesis. Another hypothesized mechanism is the activation of the pulmonary reflex by particulate matter leading to the activation of the sympathetic system. A chronic activation of this system could lead to hypertension, plaque instability and cardiac arrhythmias (Brook et al. 2004; Burgan et al. 2010; Mills et al. 2007).

As motor vehicles are the predominant source of both air pollution and noise in many cities (Allen and Adar 2011), the reported associations between road traffic exposure and CV diseases could be influenced by a confounding or interaction between those two pollutants. Yet, few studies have aimed to untangle the possible effects of road traffic

noise and air pollution. This is fundamental to better steer public health interventions and policies aimed at reducing CV effects of road traffic. For example, if road traffic induces CV effects through noise levels, then regulations to reduce car air pollutant emissions may not tackle the problem and zoning by-laws could be more appropriate. In this article, we reviewed epidemiological evidences that looked at the confounding effect of one traffic-related exposure (noise or air pollution) on the association between the exposure to traffic-related noise or air pollution and CV outcomes.

## Methods

### Data sources

The bibliographic databases used were Medline and Elsevier Embase on the Ovid SP portal. Only studies published until November 2012 were considered. No other temporal limitation was applied.

### Extraction strategy

The strategy used to conduct this review consisted of a combination of keywords representing three distinct categories: (1) exposure to traffic air pollution, (2) traffic noise exposure and (3) cardiovascular outcomes (see “Appendix” for specific keywords). Only peer review articles written in English or French on road traffic were reviewed. The studies were then selected manually according to the following exclusion criteria:

- Commentaries, editorials, review articles
- Studies not related to road traffic
- Studies not performed on humans
- Studies with no simultaneous exposure to noise and air pollutants
- Studies with no assessment of cardiovascular effects
- Studies not reporting the impact of confounding variables

Studies using the same cohort but assessing different CV outcomes were included. The references of each selected article were consulted to ensure that all pertinent information was gathered in our review. Experts in the field were also consulted to see if further articles could be added. Finally, gray literature (OAIster database, WHO and the New York Academy of Medicine) was also consulted. Studies presenting the associations before and after the authors controlled for the co-variables of interest (e.g., traffic noise or air pollution) were reviewed.

The articles selected were then separated into two categories: studies of CV outcomes where the noise exposure effect was adjusted for air pollutants and studies evaluating

the effect of air pollutants, adjusted for the noise effect. In this review, cardiovascular findings of the original studies are reported before and after the control of either road traffic noise or air pollution effects. The effect of the supposed confounders was assessed using the percentage of variation in the estimate as follows: the confounding effect (C) of noise or air pollution was evaluated by a change in the point estimate (CIE) with a cutoff point of 10 % (Eq. 1) (Bliss et al. 2012; Vittinghoff et al. 2012).

$$\frac{\text{Adjusted point estimate} - \text{Unadjusted point estimate}}{\text{Unadjusted point estimate}} \times 100 = C \quad (1)$$

To assess the quality of the study and identify possible biases within the studies reviewed, two authors (LFT and SP) reviewed independently each study. The methodology applied has been described previously (Perron et al. 2012). Briefly, we considered the study's design selection and classification bias relevant to our review. For each type of bias, the distinction between major and minor biases was assessed qualitatively by both authors. We defined major biases as those that could invalidate the results presented in the study. Minor biases are expected to affect the association studied, but unlikely to invalidate the conclusion.

## Results

Figure 1 presents a flowchart describing the stages of selection for the studies used in this review. Our keyword search yielded 221 starting articles ("Appendix"), which were reduced to 16 articles by applying the various inclusion and exclusions criteria. A large number of irrelevant articles had to be sifted by hand because of the double meaning of the terms sound and noise. We added four articles identified by snowballing and expert consultation. Of these 20 articles, a final 11 studies were discarded because it was impossible to assess the impact of the confounding variables of interest (Fig. 1). We did not find any document in the gray literature that could benefit this review.


Nine studies evaluating the relationship between CV outcomes, noise and traffic-related air pollution met our criterion (Beelen et al. 2009; Clark et al. 2012; de Kluizenaar et al. 2007; Dratva et al. 2012; Gan et al. 2012; Selander et al. 2009; Sorensen et al. 2011, 2012a, b). Of those articles, one assessed CV disease mortalities in general (Beelen et al. 2009), four assessed ischemic heart diseases (Beelen et al. 2009; Gan et al. 2012; Selander et al. 2009; Sorensen et al. 2012a), two assessed cerebrovascular diseases (Beelen et al. 2009; Sorensen et al. 2011), four

looked at blood pressure or hypertension (Clark et al. 2012; Dratva et al. 2012; Sorensen et al. 2012b; de Kluizenaar et al. 2007), and one article was identified for both heart failure and cardiac dysrhythmia (Beelen et al. 2009).

Tables 1 and 2 present the associations between cardiovascular outcomes and either road traffic air pollution or noise. The tables also summarize the population characteristics of each study, the health outcomes monitored, as well as the noise and air pollutants indicators used. When available, the background exposure levels to pollution and noise of the subjects were reported.

Six studies used day–evening–night noise levels ( $L_{den}$ ), as indicators of exposure to noise levels from road traffic (Beelen et al. 2009; de Kluizenaar et al. 2007; Gan et al. 2012; Selander et al. 2009; Sorensen et al. 2011, 2012a, b). Others studies used equivalent noise levels over 24 h ( $L_{eq,24h}$ ) (Selander et al. 2009), night noise levels ( $L_{night}$ ) (Dratva et al. 2012) or daily noise levels ( $L_{day}$ ) (Clark et al. 2012; Dratva et al. 2012). All studies reported used dispersion noise models, but only four reported validation information on the model used (Beelen et al. 2009; Sorensen et al. 2011, 2012a, b). In those studies, the average difference between the estimated noise levels and measured noise levels was 3.1 dB or less. Regarding exposure to air pollutants, seven studies used nitrogen oxide ( $NO_x$  or  $NO_2$ ) as an indicator (Clark et al. 2012; Dratva et al. 2012; Gan et al. 2012; Selander et al. 2009; Sorensen et al. 2011, 2012a, b). Two articles reported results based on particulate matter ( $PM_{10}$  or  $PM_{2.5}$ ) (de Kluizenaar et al. 2007; Gan et al. 2012). Two studies used measurements of black smoke or black carbon to account for road traffic pollutants (Beelen et al. 2009; Gan et al. 2012). Of those studies, two applied land use regression models (Beelen et al. 2009; Gan et al. 2012) to assess the air pollution exposure, while the other studies employed dispersion models (Clark et al. 2012; de Kluizenaar et al. 2007; Dratva et al. 2012; Selander et al. 2009; Sorensen et al. 2011, 2012a, b). As noted in Tables 1 and 2, seven out of the nine studies reviewed reported validation information on their air pollution model (Beelen et al. 2009; Clark et al. 2012; Dratva et al. 2012; Gan et al. 2012; Sorensen et al. 2011, 2012a, b). The Clark et al. study reported an average difference between the estimated and the measured pollutant levels of 2.4 ppb. The remaining studies presented  $R^2$  that ranged from 0.52 to 0.75. All articles presented average yearly outdoor exposure for both noise and air pollutant levels as a proxy of individual exposure. The correlations between traffic noise and air pollutants in the studies reviewed range from 0.16 to 0.72. It is surprising that studies where both noise and air pollutants were modeled, relying on some of the same variables (e.g., distance to traffic source), could generate low and disparate correlations.

**Fig. 1** Stages of the selection of studies for analysis



### Quality of studies

Table 3 presents the assessment of the quality of the studies by LFT and SP. All studies, with the exception of Selander et al. (2009) and Dratva et al. (2012), only considered the exposure at home or at work/school, which we considered as a minor classification bias likely to be non-differential. Two studies also used medical administrative databases to identify the cause of death, potentially leading to another minor classification bias. Four studies reported response rates between 30 and 60 % which could lead to a selection bias and one did not report the response rate. Finally, four studies used cross-sectional designs to assess associations with cardiovascular outcomes reported in this review. Most studies contained less than four minor biases. We did not identify any major bias in the studies assessed.

### Noise effects controlled for air pollution effects

As illustrated in Table 1, four studies assessed the association between noise levels and ischemic heart diseases

while controlling for air pollutants. Gan et al. (2012) reported an association between death from coronary heart disease and a 10 dB(A) elevation of the  $L_{den}$  level of 1.13 (95 % CI 1.06, 1.21) before and 1.09 (95 % CI 1.01, 1.18) after controlling for  $NO_2$ ,  $PM_{2.5}$  and black carbon. Models adjusted for  $PM_{2.5}$  and  $NO_2$  only in Gan et al. (2012), produced small modifications of the point estimate (Gan et al. 2012). The IRR estimated by Sorensen et al. (2012a) for the association between the incidence of myocardial infarction and  $L_{den}$  was 1.10 (95 % CI 1.03, 1.19) per increase of 10 dB(A); this IRR increased to 1.12 (95 % CI 1.02, 1.22) after adjustment for  $NO_x$ . Selander et al. (2009) only reported an association between myocardial infarction and  $L_{eq,24h}$  after controlling for  $NO_2$  (1.12 (95 % CI 0.95, 1.33) with the reference category <50 dB[A]). While Selander et al. (2009) did not present point estimates before and after adjusting for noise, the authors reported a 7 % change of the crude coefficient when compared with the model adjusted for air pollution. The percentage changes in the point estimates in the studies mentioned above were all below 10 % (ranging between 0 and 7 %). Beelen et al.

**Table 1** Estimation of the effects of exposure to noise levels on cardiovascular mortality and morbidity while controlling for air pollution (Sweden [1992–1994], Denmark [1993–2006], Canada [1994–2002], Switzerland [2002–2003], Netherlands [1997–1998] and, [1987–1996])

Studies	Characteristics of the study <sup>a</sup>	Traffic noise exposure (noise indicator) [validation of the model]	Air pollution exposure (air pollution indicator) [validation of the model]	Exposure levels	Health outcome (definition)	Main findings: (95 % confidence interval)	Percentage of change in the estimate
de Kluizenaar et al. (2007)	Groningen sample: N: 40,856 28–75 years Groningen, Netherlands 1997–1998	Dispersion model : The Standaard, Kartering Method 2 implemented in Urbis ( $L_{den}$ ) [no information on validation]	Dispersion models: (i) Local traffic contribution from the model CAR II (ii) The Gaussian dispersion model “Pluim” ( $PM_{10}$ ) [no information on validation]	Average (SD) $L_{den}$ in the Groningen sample: No AHT: 53.3 (6.9) dB(A) AHT: 54.6 (7.0) dB(A) Median (5–95 percentile) $PM_{10}$ level in the Groningen sample: No AHT: 33.5 (32.8–37.5) $\mu g/m^3$ AHT: 33.6 (32.9–37.6) $\mu g/m^3$	Self-reported antihypertensive medication intake	OR per 10 dB(A) increase Full sample OR <sub>A</sub> 1.01 (0.96–1.06) OR <sub>P</sub> 1.03 (0.96–1.11) Subgroup 45–55 years OR <sub>A</sub> 1.08 (0.97–1.20) OR <sub>P</sub> 1.19 (1.02–1.40)	Full sample 1.98 % Subgroup 45–55 years 10.19 %
				PREVENT cohort sub sample N: 8,592 28–75 years Groningen, Netherlands 1997–1998	Hypertension: use of antihypertensive medication (pharmacy record) or systolic blood pressure $\geq 140$ and diastolic blood pressure $\geq 90$ (mean of the last 2 measurements from the 2 visits)	OR per 10 dB(A) increase Full sample OR <sub>A</sub> 1.07 (0.98–1.18) OR <sub>P</sub> 1.08 (0.95–1.23) Subgroup 45–55 years OR <sub>A</sub> 1.27 (1.08–1.49) OR <sub>P</sub> 1.39 (1.08–1.77)	Full sample 0.93 % Subgroup 45–55 years 9.45 %
Beelen et al. (2009)	Cohort N: 117,528 55–69 years Netherlands 1987–1996	Dispersion model :Empara ( $L_{den}$ ) [Measured vs. Estimated: on average <2–3 dB(A)]	Land use regression (black smoke) [ $R^2 = 0.59$ ]	Average $L_{den}$ level (SD) : 52 (7) dB(A) Black smoke average level: 13.9 (2.2) $\mu g/m^3$ NO <sub>2</sub> average level: 30 $mg/m^3$	Mortality from: ischemic heart disease, cardiovascular disease, cerebrovascular disease, heart failure and cardiac dysrhythmia (ICD 9 for 1986–1995 and ICD 10 for 1996)	RR compared to a reference category of $\leq 50$ dB(A) Overall cardiovascular mortality RR <sub>A</sub> 1.25 (1.01–1.53) RR <sub>P</sub> 1.17 (0.94–1.45) Ischemic heart disease mortality RR <sub>A</sub> 1.15 (0.86–1.53) RR <sub>P</sub> 1.01 (0.74–1.36) Cerebrovascular mortality RR <sub>A</sub> 0.88 (0.52–1.50) RR <sub>P</sub> 0.95 (0.55–1.66) Heart failure mortality RR <sub>A</sub> 1.99 (1.05–3.79) RR <sub>P</sub> 1.90 (0.96–3.78) Cardiac dysrhythmia mortality RR <sub>A</sub> 1.23 (0.50–3.01) RR <sub>P</sub> 1.23 (0.48–3.13)	Overall cardiovascular mortality 6.40 % Ischemic heart disease mortality 12.17 % Cerebrovascular mortality 7.95 % Heart failure mortality 4.52 % Cardiac dysrhythmia mortality 0.00 %

Table 1 continued

Studies	Characteristics of the study <sup>a</sup>	Traffic noise exposure (noise indicator) [validation of the model]	Air pollution exposure (air pollution indicator) [validation of the model]	Exposure levels	Health outcome (definition)	Main findings: (95 % confidence interval)	Percentage of change in the estimate
Selander et al. (2009)	Case-control N: 2,095 (controls) + 1,571 (cases) 45–70 years Stockholm country, Sweden 1992–1994	Dispersion model: simplified Nordic prediction method ( $L_{Aeq,24h}$ ) [No information on validation]	Dispersion model: Gaussian Air Quality Dispersion model ( $NO_2$ ) [No information on validation]	$L_{Aeq,24h}$ : NA $NO_2$ : median level 12.9 $\mu g/m^3$ for controls Median level for cases: NA	Myocardial infarction (coronary records, hospital discharge register and the National Cause of Death at statistic Sweden)	OR compared to a reference category of $\leq 50$ dB Full sample OR <sub>p</sub> 1.12 (0.95–1.33)	7 %
Sorensen et al. (2011)	Cohort N: 51,485 55–64 years Copenhagen or Aarhus, Denmark 1993–2006	Dispersion model: Sound plan with the Nordic prediction method, DANSIM and INM 3 ( $L_{den}$ ) [Measured vs. Estimated: on average 0.2 dB]	Dispersion model: Danish AirGis ( $NO_x$ ) [ $R^2 = 0.75$ ]	Median $L_{den}$ (5–95 percentile): $< 64.5$ years 57.8 (NA) dB(A) and $\geq 64.5$ years 58.2 (NA) dB(A) $NO_x$ 5–95 percentile) median levels at $L_{den} \leq 60$ dB 18.5 (14.1–28.3) $\mu g/m^3$ and $L_{den} > 60$ dB 34.3 (16.9–137) $\mu g/m^3$	Stroke (hospital discharge register ICD 8 and 10)	IRR per 10 dB(A) increase Full sample IRR <sub>A</sub> 1.10 (1.03–1.18) IRR <sub>p</sub> 1.14 (1.03–1.25)	3.64 %
Sorensen et al. (2012a)	Cohort N: 50,614 55–64 years Copenhagen or Aarhus, Denmark 1993–2006	Dispersion model: Sound plan with the Nordic prediction method, DANSIM and INM 3 ( $L_{den}$ ) [Measured vs. Estimated: on average 0.2 dB]	Dispersion model: Danish AirGis ( $NO_x$ ) [ $R^2 = 0.75$ ]	Median $L_{den}$ (5–95 percentile) 56.4 (48.5–70.1) dB(A) Median $NO_x$ (5–95 percentile) 20.8 (14.4–88.0) $\mu g/m^3$	Myocardial infarction (ICD 10)	IRR per 10 dB(A) increase Full sample (per 10 dB(A)) IRR <sub>crude</sub> 1.10 (1.03–1.19) IRR <sub>Adj</sub> 1.12 (1.02–1.22)	1.82 %
Gan et al. (2012)	Cohort N: 445,868 45–85 years Vancouver Canada 1994–2002	Dispersion model: CadnaA base model using the EMME/2 for traffic volume ( $L_{den}$ ) [No information on validation]	Land use regression ( $NO_2$ , $PM_{2.5}$ and black carbon) [ $NO_2$ : $R^2 = 0.56$ $PM_{2.5}$ $R^2 = 0.52$ Black carbon NA]	Average $L_{den}$ (SD): 63.4 (5.0) dB(A) $PM_{2.5}$ average level (SD): 4.10 (1.64) $\mu g/m^3$ $NO_2$ average level (SD): 32.3 (8.1) $\mu g/m^3$ $NO_x$ average level (SD): 32.2 (12.0) $\mu g/m^3$ Black carbon average level (SD): 1.50 (1.1) $10^{-5}/m$	Ischemic heart diseases mortalities (ICD-9 and ICD-10)	RR per increase of 10 dB(A) $PM_{2.5}$ RR <sub>A</sub> 1.13 (1.06–1.21) RR <sub>p</sub> 1.13 (1.06–1.21) $NO_2$ + $PM_{2.5}$ RR <sub>A</sub> 1.13 (1.06–1.21) RR <sub>p</sub> 1.12 (1.05–1.21) Black carbon + $NO_2$ + $PM_{2.5}$ RR <sub>A</sub> 1.13 (1.06–1.21) RR <sub>p</sub> 1.09 (1.01–1.18)	0.00 % $NO_2$ + $PM_{2.5}$ 0.88 % Black carbon + $NO_2$ + $PM_{2.5}$ 3.54 %

**Table 1** continued

Studies	Characteristics of the study <sup>a</sup>	Traffic noise exposure (noise indicator) [validation of the model]	Air pollution exposure (air pollution indicator) [validation of the model]	Exposure levels	Health outcome (definition)	Main findings: (95 % confidence interval)	Percentage of change in the estimate
Dratva et al. (2012)	Cross-sectional N: 6,450 28–72 years Switzerland 2002–2003	Dispersion model: SONABASE ( $L_{day}$ and $L_{night}$ ) [Measured vs. Estimated: on average $\pm 2.6$ dB(A) (day) $\pm 3.1$ dB(A) (night)]	Dispersion model: PolluMap Gaussian dispersion model (NO <sub>2</sub> ) [R <sup>2</sup> = 0.72]	Average (SD) $L_{day}$ : 50.5 (7.2) dB(A) Average (SD) $L_{night}$ : 38.7 (7.8) dB(A) Average (SD) levels for: NO <sub>2</sub> : 23.0 (9.9) $\mu\text{g}/\text{m}^3$ PM <sub>10</sub> : 21.3 (7.1) $\mu\text{g}/\text{m}^3$	Blood pressure (measured by the Riva-Rocci method by trained field workers)	Increase in BP per 10 dB (A) Night time systolic BP $\beta_A$ : -0.01 (-0.6 to 0.59) $\beta_{AP}$ : 0.15 (-0.48 to 0.77) Nighttime diastolic BP $\beta_A$ : -0.05 (-0.41 to 0.30) $\beta_{AP}$ : -0.15 (-0.36 to 0.39) Daytime systolic BP $\beta_A$ : -0.11 (-0.68 to 0.47) $\beta_{AP}$ : 0.05 (-0.56 to 0.07) Daytime diastolic BP $\beta_A$ : -0.10 (-0.44 to 0.24) $\beta_{AP}$ : -0.04 (-0.40 to 0.33)	Nighttime systolic BP 1,600 % Nighttime diastolic BP 200 % Daytime systolic BP 145.45 % Daytime diastolic BP 60 %

<sup>a</sup> Final sample sizes used for analysis

OR odds ratio, NA not available, IRR incident rate ratio, A adjusted for some of the following potential confounding factors: age, education, employment, marital status, study area, mean pulse, hearing impairment, noise at work, crowding, home ownership, mother’s educational level, language spoken at home, parental support for schoolwork, classroom window glazing, body mass index, cuff size, room temperature, birth weight, parental high blood pressure, prematurity, smoking status, family history of CVD, physical inactivity smoking intensity, intake of fruits, intake of vegetables, intake of coffee, alcohol intake diabetes, antihypertensive medication, high blood pressure, long-standing illness and other comorbidity (see original article for details), P adjusted for some potential confounding factors and air pollution levels, AHT antihypertensive treatment, SD standard deviation,  $\beta$  regression coefficient,  $PM_{2.5}$  particles with a diameter of 2.5  $\mu\text{m}$  or smaller,  $PM_{10}$  particles with a diameter of larger than 2.5  $\mu\text{m}$ , but smaller than 10  $\mu\text{m}$ ,  $NO_2$  Nitrogen dioxide,  $NO_x$  Nitrogen oxide,  $dB(A)$  A-weighted decibels,  $L_{day}$  integrated A-weighted sound level over 16 h (0600–2200),  $L_{night}$  integrated A-weighted sound level over 8 h (2200–0600),  $L_{A,eq,24h}$  integrated A-weighted sound level over 24 h,  $L_{den}$  integrated A-weighted sound level over 24 h (day, evening and night) in which sound levels during the evening (1900–2300 hours) are increased by 5 dB(A) and those during the night (2300–0700 hours) by 10 dB(A)

**Table 2** Estimation of effects of exposure to ambient air pollutants on cardiovascular mortality and morbidity while controlling for noise (Netherlands [1987–1996], Denmark [2000–2002] and UK [2001–2003])

Studies	Characteristics of the study <sup>a</sup>	Traffic noise exposure (noise indicator) [validation of the model]	Air pollution exposure (air pollution indicator) [validation of the model]	Exposure levels	Health outcome (definition)	Main findings : air pollutants effects adjusted for noise effects estimate	Percentage of change in the estimate
Beelen et al. (2009)	Cohort N 117,528 55–69 years Netherlands 1987–1996	Dispersion model :Empara ( $L_{den}$ ) [Measured vs. Estimated: on average <2–3 dB(A)]	Land use regression (black smoke) [ $R^2 = 0.59$ ]	Average $L_{den}$ level (SD) : 52 (7) dB(A) Black smoke average level: 13.9 (2.2) $\mu\text{g}/\text{m}^3$ NO <sub>2</sub> average level : 30 $\text{mg}/\text{m}^3$	Mortality from: ischemic heart disease, cardiovascular disease, cerebrovascular disease, heart failure and cardiac dysrhythmia (ICD 9 for 1986–1995 and ICD 10 for 1996)	RR for an increase of 10 $\mu\text{g}/\text{m}^3$ of black smoke and adjusted for traffic intensity Overall cardiovascular mortality RR <sub>A</sub> 1.11 (0.96–1.28) RR <sub>PT</sub> 1.11 (0.95–1.28) [RR <sub>A</sub> 1.01 (1.00–1.02) RR <sub>PT</sub> 1.01 (0.99–1.02)] <sup>†</sup> Ischemic heart disease mortality RR <sub>A</sub> 1.01 (0.83–1.22) RR <sub>PT</sub> 1.01 (0.83–1.22) [RR <sub>A</sub> 1.00 (0.98–1.02) RR <sub>PT</sub> 1.00 (0.98–1.02)] <sup>†</sup> Cerebrovascular mortality RR <sub>A</sub> 1.39 (0.99–1.94) RR <sub>PT</sub> 1.41 (1.01–1.97) [RR <sub>A</sub> 1.03 (1.00–1.07) RR <sub>PT</sub> 1.03 (1.00–1.07)] <sup>†</sup> Heart failure mortality RR <sub>A</sub> 1.75 (1.00–3.05) RR <sub>PT</sub> 1.76 (1.01–3.08) [RR <sub>A</sub> 1.06 (1.00–1.12) RR <sub>PT</sub> 1.06 (1.00–1.12)] <sup>†</sup> Cardiac dysrhythmia mortality RR <sub>A</sub> 0.96 (0.51–1.79) RR <sub>PT</sub> 0.94 (0.50–1.76) [RR <sub>A</sub> 1.00 (0.99–1.06) RR <sub>PT</sub> 0.99 (0.93–1.06)] <sup>†</sup>	Overall cardiovascular mortality 0.00 % [0.00 %] <sup>†</sup> Ischemic heart disease mortality 0.00 % [0.00 %] <sup>†</sup> Cerebrovascular mortality 1.44 % [0.14 %] <sup>†</sup> Heart failure mortality 0.57 % [0.06 %] <sup>†</sup> Cardiac dysrhythmia mortality 2.08 % [0.21 %] <sup>†</sup> Systolic BP 35.90 %
Sorensen et al. (2012b)	Cohort (cross-sectional to assess the BP results) N: 44,436 55–64 years Copenhagen or Aarthus, Denmark 2000–2002	Dispersion model: Sound plan with the Nordic prediction method, DANSIM and INM 3 ( $L_{den}$ ) [Measured vs. Estimated: on average 0.2 dB]	Dispersion model: Danish AirGis (NO <sub>x</sub> ) [ $R^2 = 0.75$ ]	Median baseline $L_{den}$ 5–95 percentile) 56.3 (48.4–70.0) dB(A) Median baseline NO <sub>x</sub> (5–95 percentile) 20.2 (14.3–86.8) $\mu\text{g}/\text{m}^3$	Difference in BP (mmHg)	Regression coefficient for a doubling in NO <sub>x</sub> level Systolic BP $\beta_A$ : -0.39 (-0.64; -0.13) $\beta_P$ : -0.53 (-0.88; -0.19)	



Table 2 continued

Studies	Characteristics of the study <sup>a</sup>	Traffic noise exposure (noise indicator) [validation of the model]	Air pollution exposure (air pollution indicator) [validation of the model]	Exposure levels	Health outcome (definition)	Main findings : air pollutions effects adjusted for noise effects estimate	Percentage of change in the estimate
Clark et al. (2012)	Cross-sectional N: 719 9–10 years UK 2001–2003	Dispersion model: simplified form of the UK standard calculation of road traffic noise ( $L_{Day}$ ) [No information on validation]	Dispersion model: King's College London Emissions Toolkit ( $NO_2$ ) [Measured vs. Estimated: on average 2.4 ppb]	Average $L_{day}$ : 50 dB(A) $NO_2$ average level 42.73 $\mu g/m^3$	Blood pressure measured using automatic blood pressure meters (OMORON 711)	Regression coefficient for an increase of 1 $\mu g/m^3$ $NO_2$ increase Systolic BP $\beta_A$ : 0.058 (–0.092 to 0.210) $\beta_P$ : 0.070 (–0.120 to 0.259) Diastolic BP $\beta_A$ : 0.033 (–0.084 to 0.151) $\beta_P$ : 0.088 (–0.059 to 0.236)	Systolic BP 20.69 % Diastolic BP 166.67 %

RR Risk ratio, SD standard deviation,  $\beta$  regression coefficient,  $NO_2$  Nitrogen dioxide,  $NO_x$  Nitrogen oxide,  $dB(A)$  A-weighted decibels,  $L_{day}$  Integrated A-weighted sound level over 16 h (0600–2200),  $L_{den}$  Integrated A-weighted sound level over 24 h (day, evening and night) in which sound levels during the evening (1900–2300 hours) are increased by 5 dB(A) and those during the night (2300–0700 hours) by 10 dB(A), A adjusted for some of the following potential confounding factors: age, education, employment, marital status, study area, mean pulse, hearing impairment, noise at work, crowding, home ownership, mother's educational level, language spoken at home, parental support for schoolwork, classroom window glazing, body mass index, cuff size, room temperature, birth weight, parental high blood pressure, prematurity, smoking status, family history of CVD, physical inactivity smoking intensity, intake of fruits, intake of vegetables, intake of coffee, alcohol intake diabetes, antihypertensive medication, high blood pressure, long-standing illness and other comorbidity, P adjusted for potential confounding factors including air pollution levels

<sup>a</sup> Final sample sizes used for analysis

<sup>†</sup> Results for an increase of 1  $\mu g/m^3$

**Table 3** Quality assessment of the studies reviewed (UK [2001–2003], Sweden [1992–1994], Denmark [1993–2006], Canada [1994–2002], Switzerland [2002–2003], Netherlands [1997–1998] and, [1987–1996])

Author(s)	Selection biases		Classification biases		Study design
	Major	Minor	Major	Minor	
Beelen et al. (2009)	None	Approximately, 85 % of the population at baseline had no paid job.	None	Input data from 2000 for the noise model paired with the 1986 home address Cause of death based on non-validated medico administrative databases Exposure assessed with the residential address only	Case cohort
Sorensen et al. (2011)	None	Response rate between 30 and 60 %	None	Exposure assessed with the residential address only	Cohort
de Kluizenaar et al. (2007)	None	Response rate between 30 and 60 %	None	The air pollution indicator was not specific to road traffic Exposure assessed with the residential address only	Cross-sectional
Sorensen et al. (2012a)	None	Response rate between 30 and 60 %	None	Exposure assessed with the residential address only	Cohort
Sorensen et al. (2012b)	None	Response rate between 30 and 60 %	None	Exposure assessed with the residential address only	Cross-sectional <sup>a</sup>
Gan et al. (2012)	None	None	None	Cause of death based on non-validated medico administrative databases Exposure assessed with the residential address only	Cohort
Selander et al. (2009)	None	None	None	None	Case control
Dratva et al. (2012)	None	No direct information on response rate	None	None	Cross-sectional <sup>a</sup>
Clark et al. (2012)	None	Exclusion of 7 of the 9 school because of missing air pollution exposure	None	Exposure assessed at school only	Cross-sectional <sup>a</sup>

<sup>a</sup> Design used for the CV outcome of interest

(2009) also reported a reduction of the RR between ischemic heart disease mortality and annual  $L_{den}$  level after adjusting for black smoke and traffic intensity (from 1.15 (95 % CI 0.86, 1.53) to 1.01 (95 % CI 0.74, 1.36) with the reference category <50 dB[A]). However, this study was the only one that presented a variation of >10 % in the point estimate (12.17 %) following adjustment for air pollution.

For cerebrovascular diseases, Sorensen et al. (2011) published a positive association between hospitalization for stroke and  $L_{den}$  before (1.18 (95 % CI 1.11, 1.26) per 10 dB[A]) and after (1.14 (95 % CI 1.03, 1.25) per 10 dB[A]) controlling for  $NO_x$ . The crude association between  $L_{den}$  level and cerebrovascular mortalities found in Beelen et al. (2009) was 0.88 (95 % CI 0.52, 1.50) and moved toward unity 0.95 (95 % CI 0.55, 1.66) after controlling for black smoke. Both studies reported a percentage change in their estimate of <10 % (respectively 3.64 and 7.95 %).

Two studies assessed the effect of noise on blood pressure. In the first (de Kluizenaar et al. 2007), reported associations between  $L_{den}$  and self-reported antihypertensive medication intake or hypertension were, respectively, 1.01 (95 % CI 0.96, 1.06) and 1.07 (0.98; 1.18) before controlling for  $PM_{10}$ . Controlling for air pollutants for both outcomes resulted in a small change in the odds ratios, respectively, of 1.03 (95 % CI 0.96, 1.11) and 1.08 (95 % CI 0.95, 1.23). So adjusting for  $PM_{10}$  produced a CIE of 0.93 % for hypertension and 1.96 % for self-reported antihypertensive medication intake. Once stratified by age in both samples (the Groningen sample and the prevent cohort subsample), the only subgroup presenting significant associations was composed of individuals between 45 and 55 years old. In this age group, the percentage CIE was near our cutoff point for confounding effects (9.45 % in the prevent cohort and 10.19 % in the Groningen sample). The second study assessing blood pressure (Dratva et al. 2012) presented no significant association between road traffic noise ( $L_{night}$  and  $L_{day}$ ) and either systolic or diastolic blood pressure before and after adjustment for  $NO_2$ . The regression coefficient did, however, vary extensively before and after adjustment for air pollutants resulting in CIE ranging from 60 to 1,600 %. The effects of noise on overall cardiovascular diseases, heart failure and cardiac dysrhythmia were reported only in Beelen et al. (2009). All associations in the final model were reduced or were identical after an adjustment for black smoke. The percentage CIE was <10 %.

The correlation between road traffic noise and air pollution reported in the studies on noise effects described above does not seem to influence the CIE produced by adjusting for air pollution levels. Studies that presented weak and high correlations (see “Appendix”) were both

subject to large CIE. The quality of the approach used to estimate the confounder exposure levels (i.e., air pollution) does not appear to impact the CIE either. As presented in Table 1, the largest CIE were observed neither in studies with the small  $R^2$  nor in those with the large  $R^2$ . Though CIE does not seem to be linked to the quality of the study (quantity of biases), cohort studies appear to generally report smaller CIE than studies using case–control or cross-sectional designs.

#### Air pollution controlled for noise

As shown in Table 2, three studies assessed the association between air pollutant levels and cardiovascular diseases while controlling for noise levels. Beelen et al. (2009) reported associations between black smoke and mortality from overall cardiovascular diseases, heart failure ischemic diseases, cardiac dysrhythmia and cerebrovascular diseases. The percentage changes in the estimates for the associations reported in Beelen et al. (2009) ranged from 0.00 to 2.08 %, well below our predefined cutoff point for confounding effects. The two remaining studies reported associations with blood pressure. Sorensen et al. (2012b) presented a negative association between  $\text{NO}_x$  levels and systolic BP ( $-0.39$  (95 % CI  $-0.64, -0.13$ ) for doubling the 1 year concentrations). This association was stronger after adjustment for  $L_{\text{den}}$  ( $-0.53$  (95 % CI  $-0.88, -0.19$ ) (CI obtained from a personal communication with Mette Sorensen 01-06-2012). Adjusting for noise levels led to a 35.90 % change in the regression coefficient. In Clark et al. (2012), the regression coefficient representing the association between diastolic BP and  $\text{NO}_2$  varied from 0.033 (95 % CI  $-0.084, 0.151$ ) to 0.088 (95 % CI  $-0.059, 0.236$ ) per one point increase of  $\text{NO}_2$  ( $\mu\text{g}/\text{m}^3$ ); before and after adjusting for noise levels ( $L_{\text{day}}$ ). The regression coefficient for the association between systolic BP and  $\text{NO}_2$  increased from 0.058 (95 % CI  $-0.092, 0.210$ ) to 0.070 (95 % CI  $-0.120, 0.259$ ) per one point increase of  $\text{NO}_2$  ( $\mu\text{g}/\text{m}^3$ ), before and after adjusting for noise levels ( $L_{\text{day}}$ ). The modification of the regression coefficient for diastolic and systolic BP was, respectively, 166.67 and 20.69 %.

The correlation between road traffic noise and air pollution reported in the studies on traffic-related pollutants described above does not seem to influence the CIE produced by controlling for air pollution levels. The CIE also appears to be independent of the number of biases. On the other hand, studies with cross-sectional design presented higher CIE than the case–control study. Since none of the studies had validation information on noise exposure estimates, we could not assess the impact of the quality of the approach used to estimate the confounder exposure levels (i.e., noise) in these studies.

#### Interaction

Regarding studies that assessed the interaction between air pollutant levels and noise on cardiovascular outcomes, only two were identified. Selander et al. (2009) did not report a significant interaction between annual  $\text{NO}_2$  levels and  $L_{\text{eq},24\text{h}}$ . Gan et al. (2012) did not find a statistically significant interaction between black carbon and noise levels ( $L_{\text{den}}$ ) for ischemic heart diseases.

#### Discussion

This review aimed to assess the confounding effects of one traffic-related exposure (either noise or air pollutants) on the association between its counterpart and cardiovascular outcomes. In general, the results of the nine studies reviewed here showed that when associations between noise and CV diseases were adjusted for air pollutants, modifications of the point estimates for cardiovascular diseases were  $<10$  %, after controlling for the air pollutants, with the exception of the studies by de Kluizenaar et al. (2007), Beelen et al. (2009) and Dratva et al. (2012) where the CIE was higher than our cutoff point for confounding effects. The Beelen et al. (2009) study reported a marked decrease of the strength of the association after adjustment for air pollution and traffic intensity. Yet, the simultaneous adjustment for traffic intensity and black smoke makes the evaluation of confounding by black smoke difficult in this study. Nonetheless, no association between road traffic and CV outcomes (before and after adjustment) was found in both the Dratva et al. (2012) and the Beelen et al. (2009) studies, rendering the CIE meaningless. By its definition, a confounder must modify the association between the exposure and the outcome. To confound, such associations must be present at least before or after adjustment. de Kluizenaar et al. (2007) reported CIE of  $<2$  % in both the Groningen sample and the prevent cohort. However, CIEs nearing 10 % were observed in the 45–55 years subgroup, which could indicate the presence of a small confounding between the two exposures in this particular subgroup. Similar findings were found for associations between air pollutant levels and CV diseases, although the number of studies was limited ( $N = 3$ ): controlling for noise levels either changed the point estimates for CV diseases by  $<10$  % or the study did not present an association between the exposure and the outcome (before and after adjustment). Only the Sorensen et al. (2012b) study presented an indication of confounding by traffic noise in the association between  $\text{NO}_x$  levels and blood pressure. Nonetheless, overall these findings suggest an independent effect of noise and air pollution on CV diseases, particularly ischemic disease for which there were a

greater number of studies. The review also points to the absence of comparability between studies. Most studies were difficult to compare because different noise or air pollution indicators were used, the pollution levels were assessed using different techniques and very few studies assessed comparable health outcomes.

In this review, we also tried to verify if the impacts of noise and air pollutants on CV were subject to the same interactive effects. Though both studies (Gan et al. 2012; Selander et al. 2009) that assessed interaction effects did not find any effect, one cannot conclude that there was an absence of interaction effect between noise and air pollutant levels with so few studies. This is particularly true in the light of the point raised by Selander et al. (2009) that the interaction analysis might have lacked power. It should also be noted that both studies used different noise indicators, the air pollutant or the cardiovascular outcome, to identify possible interaction effects. Those results do suggest, though, that if a multiplicative interaction exists, it is likely to be small.

Due to limitations of the literature, we cannot conclusively ascertain the independence of the effects of the two risks on any CV health outcome. Nonetheless, the results reported tend to indicate that the impacts of traffic noise and air pollution on cardiovascular outcome are distinct, or at least that they are not completely dependent on one another. Furthermore, the correlation between noise and air pollutant levels does not seem to influence the confounding effects. A wide range of correlations between noise and air pollutants were reported in the studies reviewed and this could be partially explained by differences in the urban structure at each location (building height, distance of buildings to sidewalks, street width, traffic intensity and distance to major road). This would suggest that confounding between traffic-related noise and air pollution is a study-dependant issue. However, high correlations between noise and pollutant levels were not associated with greater confounding effects in the studies reviewed. Additionally, the quality of the exposure assessment of the confounding variables and the quality of the studies (number of biases) do not seem to influence the confounding effects. However, the reported CIE seemed higher in studies that used a cross-sectional design. This may be because these studies mainly used linear regressions and presented regression coefficients. In fact, the approach that we used to assess confounding effects was developed for risk ratios and may not be applicable for linear regression. Nonetheless, this approach has been suggested by some authors in linear regression text books (Vittinghoff et al. 2012).

This review is subjected to a few limitations. First, only a small number of studies were available to be reviewed, which reduced the strength of our findings. Secondly, with any systematic review, the possibility of publication bias is present. We tried to minimize this bias by searching the

gray literature and reviewing some non-English publications, but this bias cannot be excluded. We also tried to minimize the selective reporting bias, which could be a major one in this review. Indeed, the principal objective of most of the studies reviewed was not to assess the confounding effect between noise and air pollutants from road traffic. As shown in Fig. 1, more than half of the studies presenting the basic characteristics to be included did not report the impact of the co-variables of interest or the regression coefficients. It is also possible that some authors assessing the effect of either noise or air pollutant levels on CV diseases found that the corresponding co-variable was not significant and did not report it. This omission would, however, strengthen our results, as it would seem to indicate the independence of the effects of both exposures. Finally, the strategy used to assess the confounding effect in studies could also be criticized. The 10 % variation in the estimate is an arbitrary cutoff point that does not necessarily rule out confounding or inform on the statistical variability. Nonetheless, this cutoff point is widely used in the literature and was identified as the least biased in a simulation presented in Maldonado and Greenland (1993), in the absence of prior knowledge of confounders. Another potential limitation in our assessment of confounding is that the 10 % cutoff implies that effects are linear. In the case of nonlinear effect estimates, the absence of CIE does not preclude confounding (Janes et al. 2010). This might be the case if noise had a threshold effect, and future studies should address this.

To assess clearly the presence of confounding, future studies should use coherent noise and air pollution indicators. Those indicators should be chosen according to the effects examined. It would be better to use maximum noise levels ( $L_{\max}$ ) or equivalent noise over 1 h ( $L_{\text{eq},1\text{h}}$ ) to assess acute effects, and  $L_{\text{eq},24\text{h}}$  to assess chronic effects. The use of  $L_{\text{den}}$  or  $L_{\text{DN}}$  as the noise metrics in the studies included in this review could be contested. Those noise metrics created to assess annoyance increase the weight noise levels occurring in evening or at night and are therefore not representative of the actual sound exposure. The source of the exposure is also an important factor to take into account while choosing an indicator. For a surrogate of all pollutants emitted by on-road traffic, one should use individual pollutants such as black carbon,  $\text{NO}_x$  and ultrafine particulate that are more source specific. More studies on the relationship between noise outdoor exposure levels and personal exposure levels should be conducted. This is important given that this relationship could differ between noise and air pollution. Furthermore, future studies should present adjusted results for both noise and air pollution modeled separately, so that the impact of the pollutant that was controlled for can be assessed. As shown in Tables 1 and 2, only few studies provided validation information for the exposure models used in their study. The absence of such information prevents the reader from judging the

quality of the exposure assessment and thereby precluding the reader to judge from assessing the quality of the adjustments made. Finally, more studies are needed to find out if the confounding effect is specific to subcategories of CV outcomes. Ideally, those studies would need an epidemiologic design enabling them to assess the chronic effects of both traffic-related pollutants. We also recommend that those studies focus their research on ischemic heart diseases, hypertension or the fluctuation in blood pressure, for which mechanisms should yield more conclusive results.

### Conclusion

Results from this review suggest that confounding of cardiovascular effects by noise or air pollutants is low, on average, though heterogeneity across studies and areas within studies has been reported. The quality of exposure assessment of the confounding variable, the quality of studies as well as the correlation between noise and pollutant levels do not seem to influence the confounding effect, though with further improvements in exposure assessment, the situation may change. More studies using air pollution indicators specific to road traffic are needed to properly assess if road noise and pollutant effects on CV outcomes are subjected to the confounding effect of one another.

**Acknowledgments** This study was made possible through a financial contribution from the Public Health Agency of Canada through the National Collaborating Centre for Environmental Health. We would like to thank Hugh Davies, Helen Ward and Allan Brand for their helpful comments on the writing of this manuscript.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## Appendix

### Search strategy

1. cardiovascular.mp. (906078)
2. hypertension.mp. (887805)
3. arterial tension.mp. (560)
4. blood pressure.mp. (753655)
5. arrhythmia.mp. (783)
6. myocardial infraction.mp. (811)
7. stroke.mp. (382573)
8. vasoconstriction.mp. (79992)
9. ischemia.mp. (461795)
10. heart.mp. (2590525)
11. coagulation.mp. (227773)
12. arteries.mp. (364558)
13. blood flow.mp. (467842)
14. electrocardiogram.mp. (112029)
15. cardio<sup>a</sup>.mp. (1557392)
16. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 (5132805)
17. noise.mp. (184815)
18. sound.mp. (142837)
19. 17 or 18 (309938)
20. 16 and 19 (31528)
21. air pollution.mp. (104657)
22. particles.mp. (301041)
23. particulate.mp. (84570)
24. nitrogen oxide.mp. (9147)
25. ozone.mp. (39435)
26. diesel.mp. (12621)
27. motor vehicles.mp. (6278)
28. 21 or 22 or 23 or 24 or 25 or 26 or 27 (493806)
29. 20 and 28 (328)
30. remove duplicates from 29 (221)

See Table 4.

**Table 4** Correlations between sound levels and air pollution levels in urban areas (UK [2001–2003], Sweden [1992–1994] and [2004–2005], Denmark [1993–2006], Canada [1994–2002] and [2001], Switzerland [2002–2003], Spain [1995–2000] and [2008], Belgium [2009], Germany [2009], USA [2006–2007], Netherlands [1997–1998], [1987–1996] and [2006])

Studies	Characteristics		Noise		Air pollution		Correlation between noise and air pollution levels									
			Method	Method	Air pollution exposure (air pollutants)	Method		NO <sub>2</sub>	NO <sub>x</sub>	PM <sub>10</sub>	PM <sub>2.5</sub>	UFP	Black smoke	O <sub>3</sub>	Black carbon	
Dratva et al. (2012)	Cross-sectional N: 6,450 2002–2003 Switzerland	$L_{\text{Day}}$ and $L_{\text{night}}$	Dispersion model: SONABASE <sup>M</sup>		NO <sub>2</sub> PM <sub>10</sub>	PolluMap Gaussian dispersion <sup>M</sup>		0.28–0.29		0.16–0.17						
Clark et al. (2012)	Cross-sectional N: 719 2001–2003 UK	$L_{\text{Day}}$	Dispersion model: simplified form of the UK standard calculation of road traffic noise <sup>M</sup>		NO <sub>2</sub>	King's college London Emissions Toolkit <sup>M</sup>		0.46								
Gan et al. (2012)	N: 445,868 1994–2002 Vancouver Canada	$L_{\text{den}}$	Annual average of equivalent SPL for 24 h (CadmA model using the EMME/2 for traffic volume) <sup>M</sup>		NO <sub>2</sub> NO <sub>x</sub> PM <sub>2.5</sub>	Annual average levels (land-use regression)		0.33	0.39	–	0.14	–	–	–	–	0.44
Sorensen et al. (2011)	N: 51,485 1995, 2000 and 2005 Copenhagen or Aarhus, Denmark	$L_{\text{den}}$	Annual average of equivalent SPL for 24 h (sound plan model with the nordic prediction method, DANSIM and INM) <sup>M</sup>		NO <sub>x</sub>	Annual average levels (Danish AirGIS modelling system) <sup>M</sup>		–	0.62	–	–	–	–	–	–	–
Forster et al. (2011)	77 sites 2008 Girona, Spain	$L_{A,eq,24h}$	Average of equivalent SPL for 24 h (Girona traffic noise model) <sup>M</sup>		NO <sub>2</sub>	Annual average estimated using NO <sub>2</sub> monthly measurements <sup>M</sup>		0.62	–	–	–	–	–	–	–	–
Gan et al. (2010)	1 site 2009 Wolffstraat, Belgium	$L_{A,eq,15min}$	15 min average of equivalent SPL		NO <sub>2</sub> NO <sub>x</sub> UFP	15 min average concentration (integrated multi gas measurement platform [Airpointer®, Recordum Austria])		0.29	0.46	–	–	0.38	–	–	–	–
Boogaard et al. (2009)	264 sites (132 routes) 2006 Netherlands	$L_{A,eq,1min}$	1 min average SPL (measurement from 10:00 to 16:00)		PM <sub>2.5</sub>	One minute average concentration (condensation particle counter)		–	–	–	0.009	–	–	–	–	–

**Table 4** continued

Studies	Characteristics		Noise		Air pollution		Correlation between noise and air pollution levels						
	Noise exposure (noise indicator)	Method	Air pollution exposure (air pollutants)	Method	NO <sub>2</sub>	NO <sub>x</sub>	PM <sub>10</sub>	PM <sub>2.5</sub>	UFP	Black smoke	O <sub>3</sub>	Black carbon	
Weber (2009)	50 sites 2008 Essen, Germany	Equivalent SPL for 20 s	PM <sub>2.5</sub>	Average of air pollution level at the beginning and the end of the noise monitoring period values (condensational particulate counter 1 s)	-	-	-	0.41–0.81	-	-	-	-	
Selander et al. (2009)	N: 3,666 1970 to 1992–1994 Stockholm county, Sweden	Annual average of equivalent SPL for 24 h (simplified version of the nordic prediction method) <sup>M</sup>	NO <sub>2</sub>	Annual average levels (Stockholm county dispersion model) <sup>M</sup>	0.6	-	-	-	-	-	-	-	
Beelen et al. (2009)	N: 117,528 1986 home address with 2000–2001 noise input data and 1987–1996 air pollutants input data Netherlands	Annual average of equivalent SPL for 24 h (empirical noise model) <sup>M</sup>	black smoke	Annual levels (land use and regression model (sum of regional, urban and local traffic contributions)) <sup>M</sup>	-	-	-	-	-	0.24	-	-	
Allen et al. (2009)	105 sites 2006–2007 Chicago and Riverside county USA	Equivalent SPL for 5 min (measurement from 10:00 to 16:00)	NO <sub>2</sub> , NO <sub>x</sub> and UFP	14 days average (Ogawa sampler)	0.16–0.62 <sup>a</sup>	0.49–0.62 <sup>a</sup>	-	-	0.22–0.41 <sup>a</sup>	-	-	-	
Davies et al. (2009)	103 sites 2001 Vancouver, Canada	Equivalent SPL for 5 min (measurement at 103 sites from 8:00 to 18:00)	NO <sub>2</sub> and NO <sub>x</sub>	14 days average (passive sampler)	0.53	0.64	-	-	-	-	-	-	
de Kluizenaar et al. (2007)	N: 40,856 1997–1998 Sweden Groningen, Netherlands	Annual average of equivalent SPL for 24 h (Urbis model) [standard Karterings Method 2]) <sup>M</sup>	PM <sub>10</sub>	Annual average levels (Netherlands' standard Dutch models for local air pollution calculations: the street model CAR I and dispersion model "Pluim") <sup>M</sup>	-	-	0.72	-	-	-	-	-	

Table 4 continued

Studies	Characteristics		Noise	Air pollution		Correlation between noise and air pollution levels						
	Method	Method		Air pollution exposure (air pollutants)	Method	NO <sub>x</sub>	PM <sub>10</sub>	PM <sub>2.5</sub>	UFP	Black smoke	O <sub>3</sub>	Black carbon
Persson et al. (2007)	$L_{A,eq,24h}$	Annual average of equivalent SPL for 24 h (simplified version of the nordic prediction method) <sup>M</sup>	NO <sub>x</sub>	Modified Gaussian dispersion model (ENVIMAN) <sup>M</sup>	0.50							
ALNAP- (2006) <sup>†</sup>	$L_{den}$	Annual average of equivalent SPL for 24 h (harmonoise model) <sup>M</sup>	NO <sub>2</sub> , and PM <sub>10</sub>	Annual average (network emission model (NEMO)) <sup>M</sup>	0.12–0.48 <sup>a</sup>	0.09–0.39 <sup>a</sup>	–	–	–	–	–	–
Linares et al. (2006)	$L_{A,eq,24h}$	Equivalent SPL for 24 h	NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub> , PM <sub>10</sub> and NO <sub>x</sub>	Daily average (from the Madrid city air pollution network)	0.138	0.089	–	–	–	–	–	–0.275
Ising et al. (2004)	$L_{night}$	Equivalent SPL from 22:00 to 6:00	NO <sub>2</sub>	Exposure from 58–93 h (palms tube)	0.836	–	–	–	–	–	–	–
Tobias et al. (2001)	$L_{eq,24h}$	Equivalent SPL for 24 h	NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub> and NO <sub>x</sub>	Daily average (from the Madrid city air pollution network) except for O <sub>3</sub> (1 h maximum value)	0.32	0.35	–	–	–	–	–	–0.42
Kiaebø et al. (2000)	$L_{A,eq,24h}$	Annual average of equivalent SPL for 24 h (nordic prediction method) <sup>M</sup>	NO <sub>2</sub> , PM <sub>10</sub> , PM <sub>2.5</sub>	3 month average of hourly estimations (dispersion model EPISODE) <sup>M</sup>	0.48	0.34	0.39	–	–	–	–	–
BEG-(1998) <sup>b</sup>	$L_{dn}$	Annual average of equivalent SPL for 24 h (sound plan software) <sup>M</sup>	NO <sub>2</sub> and PM <sub>10</sub>	NA (Gaussian propagation model) <sup>M</sup>	0.63	0.61	–	–	–	–	–	–

<sup>a</sup> Lowest and highest correlation of all the sampling sessions

<sup>M</sup> Exposure asses using models

<sup>b</sup> The description of those studies were available only in Lercher et al. 2011 (those studies are either not publish in English or French or are not publish at all)

*N* final sample size used for analysis

PM<sub>2.5</sub> particles with a diameter of 2.5 μm or smaller, PM<sub>10</sub> particles with a diameter of larger than 2.5 μm but smaller than 10 μm, NO<sub>2</sub> nitrogen dioxide, NO<sub>x</sub> nitrogen oxide, O<sub>3</sub> Ozone, dB(A) a-weighted decibels,  $L_{A,eq,24h}$  integrated A-weighted sound level over 24 h,  $L_{day}$  integrated sound level over 16 h (0600–2200),  $L_{night}$  integrated A-weighted sound level over 8 h (2200–0600),  $L_{A,eq,20s}$  integrated A-weighted sound level over 20 s,  $L_{A,eq,1min}$  integrated A-weighted sound level over 1 min,  $L_{A,eq,5min}$  integrated A-weighted sound level over 5 min,  $L_{A,eq,15min}$  integrated A-weighted sound level over 15 min,  $L_{dn}$  integrated A-weighted sound level during the night (22h00–07h00) are increased by 10 dB(A),  $L_{den}$  integrated A-weighted sound level over 24 h (day, evening and night) in which sound levels during the evening (1900–2300 hours) are increased by 5 dB(A) and those during the night (2300–0700 hours) by 10 dB(A), Last validation 12th April 2012



## References

- Adar SD, Sheppard L, Vedal S, Polak JF, Sampson PD, Diez Roux AV, Budoff M, Jacobs DR Jr, Barr RG, Watson K, Kaufman JD (2013) Fine particulate air pollution and the progression of carotid intima-medial thickness: a prospective cohort study from the multi-ethnic study of atherosclerosis and air pollution. *PLoS Med* 10(4):e1001430
- Allen RW, Adar SD (2011) Are both air pollution and noise driving adverse cardiovascular health effects from motor vehicles? *Environ Res* 111(1):184–185
- Allen RW, Davies H, Cohen MA, Mallach G, Kaufman JD, Adar SD (2009) The spatial relationship between traffic-generated air pollution and noise in 2 US cities. *Environ Res* 109(3):334–342
- Amato MC, Giordano C, Galia M, Criscimanna A, Vitabile S, Midiri M, Galluzzo A (2010) Visceral Adiposity Index: a reliable indicator of visceral fat function associated with cardiometabolic risk. *Diabetes Care* 33(4):920–922. doi:10.2337/dc09-1825
- Babisch W (2006) Transportation noise and cardiovascular risk: updated review and synthesis of epidemiological studies indicate that the evidence has increased. *Noise Health* 8(30):1–29
- Babisch W, Kamp I (2009) Exposure–response relationship of the association between aircraft noise and the risk of hypertension. *Noise Health* 11(44):161–168. doi:10.4103/1463-1741.53363
- Beelen R, Hoek G, van den Brandt PA, Goldbohm RA, Fischer P, Schouten LJ, Armstrong B, Brunekreef B (2008) Long-term exposure to traffic-related air pollution and lung cancer risk. *Epidemiology* 19(5):702–710. doi:10.1097/EDE.0b013e318181b3ca
- Beelen R, Hoek G, Houthuijs D, Van Den Brandt PA, Goldbohm RA, Fischer P, Schouten LJ, Armstrong B, Brunekreef B (2009) The joint association of air pollution and noise from road traffic with cardiovascular mortality in a cohort study. *Occup Environ Med* 66(4):243–250
- Bliss R, Weinberg J, Webster T, Vieira V (2012) Determining the probability distribution and evaluating sensitivity and false positive rate of a confounder detection method applied to logistic regression. *J Biom Biostat* 3(4):142. doi:10.4172/2155-6180.1000142
- Boogaard H, Borgman F, Kamminga J, Hoek G (2009) Exposure to ultrafine and fine particles and noise during cycling and driving in 11 Dutch cities. *Atmos Environ* 43(27):4234–4242
- Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, Luepker R, Mittleman M, Samet J, Smith SC Jr, Tager I (2004) Air pollution and cardiovascular disease: a statement for healthcare professionals from the expert panel on population and prevention science of the American Heart Association. *Circulation* 109(21):2655–2671. doi:10.1161/01.CIR.0000128587.30041.C8
- Brook RD, Rajagopalan S, Pope CA, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC, Whitsel L, Kaufman JD, American Heart Association Council on Epidemiology and Prevention, Council on the Kidney in Cardiovascular Disease, and Council on Nutrition, Physical Activity and Metabolism (2010) Particulate matter air pollution and cardiovascular disease. *Circulation* 121(21):2331–2378. doi:10.1161/CIR.0b013e318181d8ce1
- Burgan O, Smargiassi A, Perron S, Kosatsky T (2010) Cardiovascular effects of sub-daily levels of ambient fine particles: a systematic review. *Environ Health* 9:26. doi:10.1186/1476-069X-9-26
- Chen H, Goldberg MS, Burnett RT, Jerrett M, Wheeler AJ, Villeneuve PJ (2013) Long-term exposure to traffic-related air pollution and cardiovascular mortality. *Epidemiology* 24(1):35–43. doi:10.1097/EDE.0b013e318276c005
- Clark C, Crombie R, Head J, van Kamp I, van Kempen E, Stansfeld SA (2012) Does traffic-related air pollution explain associations of aircraft and road traffic noise exposure on children's health and cognition? A secondary analysis of the United Kingdom sample from the RANCH project. *Am J Epidemiol* 176(4):327–337. doi:10.1093/aje/kws012
- Davies HW, Vlaanderen JJ, Henderson SB, Brauer M (2009) Correlation between co-exposures to noise and air pollution from traffic sources. *Occup Environ Med* 66(5):347–350
- de Kluijzenaar Y, Gansevoort RT, Miedema HM, de Jong PE (2007) Hypertension and road traffic noise exposure. *J Occup Environ Med* 49(5):484–492. doi:10.1097/JOM.0b013e318058a9ff
- Dratva J, Phuleria HC, Foraster M, Gaspoz JM, Keidel D, Kunzli N, Liu LJ, Pons M, Zemp E, Gerbase MW, Schindler C (2012) Transportation noise and blood pressure in a population-based sample of adults. *Environ Health Perspect* 120(1):50–55. doi:10.1289/ehp.1103448
- European Environment Agency (EEA) (2010) Good practice guide on noise exposure and potential health effects. European Environment Agency (EEA), Copenhagen
- Finkelstein MM, Jerrett M, Sears MR (2004) Traffic air pollution and mortality rate advancement periods. *Am J Epidemiol* 160(2):173–177. doi:10.1093/aje/kwh181
- Foraster M, Deltell A, Basagaña X, Medina-Ramón M, Aguilera I, Bouso L, Grau M, Phuleria HC, Rivera M, Slama R, Sunyer J, Targa J, Kunzli N (2011) Local determinants of road traffic noise levels versus determinants of air pollution levels in a Mediterranean city. *Environ Res* 111(1):177–183. ISSN 0013-9351. doi:10.1016/j.envres.2010.10.013. <http://www.sciencedirect.com/science/article/pii/S0013935110001878>
- Gan WQ, Tamburic L, Davies HW, Demers PA, Koehoorn M, Brauer M (2010) Changes in residential proximity to road traffic and the risk of death from coronary heart disease. *Epidemiology* 21(5):642–649
- Gan WQ, Davies HW, Koehoorn M, Brauer M (2012) Association of long-term exposure to community noise and traffic-related air pollution with coronary heart disease mortality. *Am J Epidemiol* 175(9):898–906. doi:10.1093/aje/kwr424
- Haralabidis AS, Dimakopoulou K, Velonaki V, Barbaglia G, Mussin M, Giampaolo M, Selander J, Pershagen G, Dudley M-L, Babisch W, Swart W, Katsouyanni K, Jarup L, HYENA Consortium (2011) Can exposure to noise affect the 24 h blood pressure profile? Results from the HYENA study. *J Epidemiol Community Health* 65(6):535–541. doi:10.1136/jech.2009.102954
- HEI Panel on the Health Effects of Traffic-Related Air Pollution (2010) Traffic-related air pollution: a critical review of the literature on emissions, exposure, and health effects vol special report 17. Health Effects Institute, Boston, MA
- Hoek G, Brunekreef B, Goldbohm S, Fischer P, van den Brandt PA (2002) Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet* 360(9341):1203–1209
- Hoffmann B, Moebus S, Stang A, Beck EM, Dragano N, Möhlenkamp S, Schmermund A, Memmesheimer M, Mann K, Erbel R, Jöckel KH, Heinz Nixdorf RECALL Study Investigative Group (2006) Residence close to high traffic and prevalence of coronary heart disease. *Eur Heart J* 27(22):2696–2702. doi:10.1093/eurheartj/ehl278
- Hoffmann B, Moebus S, Dragano N, Möhlenkamp S, Memmesheimer M, Erbel R, Jöckel KH, Heinz Nixdorf Recall Investigative Group (2009) Residential traffic exposure and coronary heart disease: results from the Heinz Nixdorf Recall Study. *Biomarkers* 14(s1):74–78. doi:10.1080/13547500902965096
- Ising H, Lange-Asschenfeldt H, Moriske HJ, Born J, Eilts M (2004) Low frequency noise and stress: bronchitis and cortisol in

- children exposed chronically to traffic noise and exhaust fumes. *Noise Health* 6(23):21–28
- Janes H, Dominici F, Zeger S (2010) On quantifying the magnitude of confounding. *Biostatistics* 11(3):572–582. doi:[10.1093/biostatistics/kxq007](https://doi.org/10.1093/biostatistics/kxq007)
- Klæboe R, Kolbenstvedt M, Clench-Aas J, Bartonova A (2000) Oslo traffic study: part 1: an integrated approach to assess the combined effects of noise and air pollution on annoyance. *Atmos Environ* 34(27):4727–4736
- Lercher P, Botteldooren D, Widmann U, Uhrner U, Kammeringer E (2011) Cardiovascular effects of environmental noise: research in Austria. *Noise Health* 13(52):234–250
- Liao D, Shaffer ML, He F, Rodriguez-Colon S, Wu R, Whitsel EA, Bixler EO, Cascio WE (2010) Fine particulate air pollution is associated with higher vulnerability to atrial fibrillation—the APACR study. *J Toxicol Environ Health, Part A* 74(11):693–705. doi:[10.1080/15287394.2011.556056](https://doi.org/10.1080/15287394.2011.556056)
- Linares C, Diaz J, Tobias A, De Miguel JM, Otero A (2006) Impact of urban air pollutants and noise levels over daily hospital admissions in children in Madrid: a time series analysis. *Int Arch Occup Environ Health* 79(2):143–152. doi:[10.1007/s00420-005-0032-0](https://doi.org/10.1007/s00420-005-0032-0)
- Maheswaran R, Elliott P (2003) Stroke mortality associated with living near main roads in England and Wales. *Stroke* 34(12):2776–2780. doi:[10.1161/01.str.0000101750.77547.11](https://doi.org/10.1161/01.str.0000101750.77547.11)
- Maldonado G, Greenland S (1993) Simulation study of confounder-selection strategies. *Am J Epidemiol* 138(11):923–936
- Mills NL, Tornqvist H, Robinson SD, Gonzalez MC, Soderberg S, Sandstrom T, Blomberg A, Newby DE, Donaldson K (2007) Air pollution and atherothrombosis. *Inhal Toxicol* 19(Suppl 1):81–89. doi:[10.1080/08958370701495170](https://doi.org/10.1080/08958370701495170)
- Nuvolone D, Balzi D, Chini M, Scala D, Giovannini F, Barchielli A (2011) Short-term association between ambient air pollution and risk of hospitalization for acute myocardial infarction: results of the cardiovascular risk and air pollution in tuscan (RISCAT) study. *Am J Epidemiol* 174(1):63–71. doi:[10.1093/aje/kwr046](https://doi.org/10.1093/aje/kwr046)
- Perron S, Tétéreault LF, King N, Plante C, Smargiassi A (2012) Review of the effect of aircraft noise on sleep disturbance in adults. *Noise Health* 14(57):58–67. doi:[10.4103/1463-1741.95133](https://doi.org/10.4103/1463-1741.95133)
- Persson R, Bjork J, Ardo J, Albin M, Jakobsson K (2007) Trait anxiety and modeled exposure as determinants of self-reported annoyance to sound, air pollution and other environmental factors in the home. *Int Arch Occup Environ Health* 81(2):179–191. doi:[10.1007/s00420-007-0204-1](https://doi.org/10.1007/s00420-007-0204-1)
- Pirrerá S, De Valck E, Cluydts R (2010) Nocturnal road traffic noise: a review on its assessment and consequences on sleep and health. *Environ Int* 36(5):492–498
- Rosenlund M, Berglind N, Pershagen G, Hallqvist J, Jonson T, Bellander T (2006) Long-term exposure to urban air pollution and myocardial infarction. *Epidemiology* 17(4):383–390
- Selander J, Nilsson ME, Bluhm G, Rosenlund M, Lindqvist M, Nise G, Pershagen G (2009) Long-term exposure to road traffic noise and myocardial infarction. *Epidemiology* 20(2):272–279
- Sorensen M, Hvidberg M, Andersen ZJ, Nordsborg RB, Lillelund KG, Jakobsen J, Tjønneland A, Overvad K, Raaschou-Nielsen O (2011) Road traffic noise and stroke: a prospective cohort study. *Eur Heart J*. 32(6):737–744
- Sorensen M, Andersen ZJ, Nordsborg RB, Jensen SS, Lillelund KG, Beelen R, Schmidt EB, Tjønneland A, Overvad K, Raaschou-Nielsen O (2012a) Road traffic noise and incident myocardial infarction: a prospective cohort study. *PLoS One* 7(6):e39283. doi:[10.1371/journal.pone.0039283](https://doi.org/10.1371/journal.pone.0039283)
- Sorensen M, Hoffmann B, Hvidberg M, Ketzel M, Jensen SS, Andersen ZJ, Tjønneland A, Overvad K, Raaschou-Nielsen O (2012b) Long-term exposure to traffic-related air pollution associated with blood pressure and self-reported hypertension in a Danish cohort. *Environ Health Perspect* 120(3):418–424. doi:[10.1289/ehp.1103631](https://doi.org/10.1289/ehp.1103631)
- Taheri S, Lin L, Austin D, Young T, Mignot E (2004) Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 1(3):e62
- Tobias A, Diaz J, Saez M, Carlos Alberdi J (2001) Use of poisson regression and Box-Jenkins models to evaluate the short-term effects of environmental noise levels on daily emergency admissions in Madrid, Spain. *Eur J Epidemiol* 17(8):765–771
- van Kempen E, Babisch W (2012) The quantitative relationship between road traffic noise and hypertension: a meta-analysis. *J Hypertens* 30(6):1075–1086. doi:[10.1097/HJH.0b013e3283252ac54](https://doi.org/10.1097/HJH.0b013e3283252ac54)
- Vardeny O, Peppard PE, Finn LA, Faraco JH, Mignot E, Hla KM (2011) [beta]2 adrenergic receptor polymorphisms and nocturnal blood pressure dipping status in the Wisconsin Sleep Cohort Study. *J Am Soc Hypertens* 5(2):114–122
- Vittinghoff E, Glidden DV, Shiboski SC, McCulloch CE (2012) *Regression methods in biostatistics. Linear, logistic, survival, and repeated measures models*, 2nd edn. Springer Science+Business Media, Boston
- Weber S (2009) Spatio-temporal covariation of urban particle number concentration and ambient noise. *Atmos Environ* 43(34):5518–5525
- Westman JC, Walters JR (1981) Noise and stress: a comprehensive approach. *Environ Health Perspect* 41:291–309
- World Health Organisation (2011) Burden of disease from environmental noise. WHO Regional Office for Europe, Copenhagen
- World Health Organisation Regional Office for Europe (2011) Burden of disease from environmental noise. World Health Organisation, Copenhagen