

# Catalase-Dependent Peroxidative Metabolism in the Alveolar Macrophage during Phagocytosis

J. BERNARD L. GEE, CHARLES L. VASSALLO, PAUL BELL, JAMES KASKIN, R. E. BASFORD, and JAMES B. FIELD

*From the Departments of Medicine and Biochemistry, University of Pittsburgh School of Medicine and Veterans Administration Hospital, Pittsburgh, Pennsylvania 15213*

**ABSTRACT** Evidence for the presence of peroxidative metabolism in rabbit alveolar macrophages (AM) has been obtained from the following observations: (a) catalase is present in high concentrations; (b) peroxidase activity could not be detected employing guaiacol as substrate; (c) the irreversible inhibition of AM catalase by aminotriazole served as a detection system for  $H_2O_2$  and demonstrated increased intracellular  $H_2O_2$  after phagocytosis; (d) formate oxidation, a marker of catalase-dependent peroxidations, occurs in resting AM and is increased by phagocytosis; (e) measurements of  $H_2O_2$  accumulation in a dialysate of AM demonstrated two-fold increase during phagocytosis; and (f) aminotriazole diminishes  $O_2$  utilization and  $^{14}CO_2$  production from labeled glucose and pyruvate. It is concluded that, while catalase-dependent  $H_2O_2$  metabolism is not essential for particle entry, this pathway represents one of the metabolic pathways stimulated by particle entry in the AM.

## INTRODUCTION

The alveolar macrophage (AM) plays an important role in pulmonary bacterial clearance by functioning as a phagocyte. While phagocytosis by the AM requires oxygen and stimulates both  $O_2$  consumption ( $Q_{O_2}$ ) and  $^{14}CO_2$  production from labeled glucose (2), the relation of these metabolic pathways to  $H_2O_2$  has not been defined in the AM. The classical work of Iyer, Islam, and Quastel (3) first indicated that hydrogen peroxide plays an important role in these pathways in another phago-

cytic cell, namely the polymorphonuclear leucocyte (PMN). Their evidence indicated that the oxidation of reduced pyridine nucleotides is accompanied by the generation of  $H_2O_2$ . The generation of  $H_2O_2$  was detected indirectly employing formate oxidation as a marker. Further evidence for  $H_2O_2$ -dependent metabolism in the PMN has been provided by a number of workers notably Karnovsky (4), Roberts and Quastel (5), and Paul and Sbarra (6). This peroxidative metabolism has recently been incupated in intracellular bactericidal mechanisms in the PMN (7, 8).

This study was designed to provide evidence concerning  $H_2O_2$  metabolism in resting and phagocytosing AM. Several approaches have been used. First, inferences have been drawn from the presence of enzymes utilizing  $H_2O_2$  as a substrate. Second, the catalase inhibitor aminotriazole (AT) may be used to detect intracellular  $H_2O_2$  (9) since it causes an irreversible inhibition of catalase:  $H_2O_2$  compound 1 as opposed to a reversible inhibition of free catalase (10). Third, formate oxidation may be used as a marker of catalase-dependent  $H_2O_2$  metabolism (3) and changes in  $H_2O_2$  concentration in intact cells may be detected by fluorimetric measurements of  $H_2O_2$  concentration by the method of Paul and Sbarra (6).

## METHODS

*Alveolar macrophages.* The AM were harvested by a modification of the technique of Myrvik, Leake, and Fariss (11) in which AM are obtained by pulmonary lavage from New Zealand white rabbits. The lavage fluid was Krebs Ringer phosphate solution (KRPS) (12) at pH 7.4 to which 5.5 mM glucose was added. Average cell yield was  $1 \times 10^6$  cells/rabbit. Red cell contamination was less than 5%. Polymorphonuclear leucocytes were less than 2% of the nucleated cells.

*Leucocytes and red blood cells.* Human leucocytes were separated by sedimentation of five parts of heparinized blood with one part 6% dextran in normal saline. The plasma,

This work was presented in part before the American Society of Clinical Investigation May 1968 (1).

Dr. Vassallo's present address is the Veterans Administration Hospital, Cincinnati, Ohio 45220.

Received for publication 12 May 1969 and in revised form 9 February 1970.

which contained the leucocytes, was pipetted off and the cells separated by centrifugation at 3000 rpm at room temperature. The WBC were resuspended in KRPS before freeze-thawing for peroxidase determination. Red cell contamination of this WBC preparation varied between 5 and 15% of WBC. Rabbit red cells were prepared from heparinized blood by centrifugation. The buffy coat was removed and cells resuspended in KRPS.

**Phagocytosis studies.** Heat-killed (75°C for 30 min) coagulase-negative *Staphylococcus albus* were employed as test particles. Bacteria to cell ratios were approximately 20/1 unless otherwise specified. Phagocytosis was assessed by light microscopy of Gram's stained preparations obtained after 30 min incubation at 37°C in KRPS containing 5.5 mM glucose and 15% fresh rabbit serum. A semiquantitative expression of particle entry was employed, namely the number of cells containing 0-5, 6-10, and > 11 bacteria were counted. At least 50 cells were examined.

**Catalase activity.** Catalase activity was determined by the method of Feinstein (13), employing 0.1 M sodium perborate. Perborate utilization in 5 min was measured by titration with a 0.1 N solution of potassium permanganate after the reaction was stopped with 1 N H<sub>2</sub>SO<sub>4</sub>. The catalase activities of AM, red cells, and purified beef liver catalase (Worthington Biochemical Corp., Freehold, N. J.) were determined after incubation for 30 min in KRPS containing 15% homologous serum and 5.5 mM glucose at pH 7.4. Measurements were performed on two preparations of AM, namely hand homogenized preparations and after three cycles of freeze-thawing employing acetone-dry ice. Both total extract and the supernatant obtained by centrifuging both types of preparation at 8000 rpm in a Servall centrifuge RC2 (7710 g) for 10 min were assayed. Activity was expressed as milliequivalents of perborate utilized in 5 min; 1 U representing the utilization of 1 mEq of perborate. The method showed good reproducibility, a coefficient of variation  $\pm 2\%$ , and linear concentration response when perborate utilization during 5 min was employed. Prolongation of incubation time of intact AM up to 3 hr did not increase the catalase activity. It should be noted that cell counts were measured in a limited number of experiments since all experiments to determine the effects of preparative procedures, AT, or particle exposure were internally controlled by comparison with control aliquots of the same AM preparations. In experiments employing aminotriazole, this was added at the beginning of the 30 min incubation period.

Dialysis experiments to assess the binding of aminotriazole to catalase were performed on 6-ml aliquots of a freeze-thaw extract of the AM in KRPS contained in a cellophane membrane. Dialysis was carried out for 3 hr against 100 ml of KRPS which changed hourly.

Additional experiments were performed to measure the effects of AT on cells previously exposed to bacteria. The procedure employed was the following; 4-ml aliquots of AM suspended in KRPS, containing 5.5 mM glucose and 15% fresh rabbit serum, were incubated for 30 min in the presence or absence of heat-killed *S. albus*. The cells were partially separated from bacteria by centrifugation at 1000 rpm for 15 min, washed in 10 ml of KRPS, recentrifuged, and resuspended in KRPS:serum:glucose with or without 20 mM AT. In some experiments aminotriazole-<sup>14</sup>C was employed. After 30 min incubation at 37°C, the cells were separated, washed, and resuspended in 1 ml of KRPS for freeze-thawing. Aliquots were taken from freeze-thaw extracts for catalase measurements and for counting where

aminotriazole-<sup>14</sup>C was employed. Counting was performed in Bray's solution (14) containing 2% v/v of 10% hyamine.

**Peroxidase activity.** This was determined by a modification of the guaiacol method of Chance and Maehly (15). Whole extracts of freeze-thawed cells were employed. The assay medium contained 0.1 M phosphate buffer at pH 7.4, 0.5 ml of 100 mM guaiacol, 0.2 ml of extract sample, and 0.02 ml of ice-cold 10 mM H<sub>2</sub>O<sub>2</sub>. Absorbancy changes due to tetraguaiacol formation were measured at 750 m $\mu$  in a Zeiss spectrophotometer and the time required to produce an 0.05 U increase in absorbancy recorded. Results were expressed in reciprocal seconds per 10<sup>9</sup> cells. The method showed a coefficient of variation of  $\pm 7\%$  and a linear response with time and number of cells. The effect of excess catalase on the estimation of peroxidase was examined in view of the presence of catalase in the AM and in view of the contamination of the white blood cell (WBC) preparation by catalase containing red cells. No effects on the peroxidase estimation performed on WBC preparations were observed under two conditions: (a) the addition of sufficient red cells to double the red cell contamination; and (b) the addition of purified beef liver catalase to increase threefold the initial catalase activity of the WBC preparation.

**Hydrogen peroxide determination.** Hydrogen peroxide was determined spectrophotometrically on dialysates of AM as described by Paul and Sbara (6). The nonfluorescent dye, diacetyl-2,7-dichlorofluorescein (LDADCF), was synthesized by the method of Brandt and Keston (16) and the fluorescence of the oxidized product of alkali-activated LDADCF was measured with an Aminco-Bowman spectrofluorimeter. The excitation wave length was 340 m $\mu$  and the emission wave length 525 m $\mu$ .

Aliquots of AM (20-60  $\times 10^6$  cells) in the presence and absence of heat-killed *S. albus* (bacteria to cell ratio, 150:1) were incubated at 37°C in KRPS containing 15% homologous serum, final volume 5 ml, in dialysis tubing immersed in 60 ml of KRPS. Samples were removed from the dialysate at appropriate times for determination of the H<sub>2</sub>O<sub>2</sub> produced. Control experiments were also performed in which dialysate obtained from serum KRPS and, where appropriate, bacteria were assayed. The difference in fluorescence between experimental and the relevant control dialysate was taken as the measure of H<sub>2</sub>O<sub>2</sub> produced by the AM. This was reduced by over 90% when commercial catalase (1 perborate U) was added to the cell suspension. Data were expressed as the quantity of H<sub>2</sub>O<sub>2</sub> in the total dialysate per 1  $\times 10^6$  cells in dialysis bag.

**Metabolic studies.** All studies were performed in KRPS containing 5.5 mM glucose and 15% homologous serum at pH 7.4. Approximately 25  $\times 10^6$  cells were present in each flask in a final volume of 4 ml. Bacteria cell ratio was 150:1. O<sub>2</sub> consumption was measured for 1 hr in a modified Gilson respirometer (Gilson Medical Electronics, Inc., Middleton, Wis.) after respirometer gassing with 100% O<sub>2</sub>. Other studies were performed for 1 hr in a Dubnoff metabolic shaker with air as the gas phase.

Approximately 0.3  $\mu$ Ci of the following labeled substrates were employed without the addition of unlabeled substrate: sodium formate-<sup>14</sup>C<sup>1</sup> (3 mCi/mmmole), 1-sodium acetate-<sup>14</sup>C<sup>2</sup> (40 mCi/mmmole), and 1-sodium pyruvate-<sup>14</sup>C<sup>2</sup> (4.0 mCi/mmmole). The studies with 1-glucose-<sup>14</sup>C<sup>2</sup> (4.0 mCi/mmmole) and 6-glucose-<sup>14</sup>C<sup>2</sup> (3.8 mCi/mmmole) employed the addition of approximately 0.3  $\mu$ Ci of tracer glucose to each flask in which the total glucose concentration was 5.5 mM. The

<sup>1</sup> New England Nuclear Corp., Boston, Mass.

<sup>2</sup> Nuclear-Chicago Corporation, Des Plaines, Ill.

TABLE I  
The Effect of Extraction Procedures on  
Catalase Activity of AM

Freeze-thawed extract		Homogenate	
Total	Supernatant	Total	Supernatant
0.55 ± 0.05	0.52 ± 0.05	0.22 ± 0.04	0.02 ± 0.01

Catalase activity expressed as milliequivalents of perborate utilized in 5 min. Mean ± 1 SD of three duplicate experiments shown.

counts obtained with 6-glucose-<sup>14</sup>C were adjusted to the same specific activity as 1-glucose-<sup>14</sup>C by the ratio 1.05. <sup>14</sup>CO<sub>2</sub> was released with 6 N H<sub>2</sub>SO<sub>4</sub> in the closed flask and entrapped in 10% hyamine hydroxide in the center cup. The hyamine was transferred quantitatively into the counting vials containing 2,5-diphenyloxazole (PPO) and 1-4-bis-2-4-methyl-(5-phenyloxazolyl)benzene in toluene and counting was performed with a liquid scintillation counter (Tri-Carb, Packard Instruments Co., Downers Grove, Ill.). All experiments were performed in triplicate.

*Aminotriazole.* 3-amino-1,2,4-triazole was obtained from Nutritional Biochemicals Corp., Cleveland, Ohio and recrystallized from ethanol. Labeled 5-aminotriazole-<sup>14</sup>C (1.5 mCi/mmmole) was obtained from Tracerlab Div. of Waltham, Mass.

## RESULTS

*Catalase and peroxidase activity.* The catalase activity in two types of AM preparations is indicated in Table I, which compares duplicate measurements of the activity present in total extract and supernatant obtained from aliquots of the same AM preparations after either homogenization or freeze-thawing. Two points emerge. First, freeze-thaw preparations demonstrate 2.5 times more catalase activity than homogenized preparations. Second, 95% of the activity was found in the supernatant after freeze-thawing while most of the activity remains in the particulate fraction after homogenization. The freeze-thaw supernatant was therefore employed in the ensuing measurements of the catalase activity. Further observations, not shown here, indicated that three cycles of freeze-thawing produced maximal catalase activity. No loss of catalase activity

TABLE II  
The Effects of Aminotriazole on AM Catalase Activity

Experiment	Control	Aminotriazole	
		100 mM	20 mM
1	0.29	0.06	0.16
2	0.24	0.02	0.08

Means of two duplicate experiments are indicated; catalase activity expressed as milliequivalents of perborate utilized in 5 min.

from the freeze-thaw supernatant obtained by centrifugation at 7710 g occurred after high speed centrifugation (up to 100,000 g) indicating that three cycles of freeze-thawing produced maximal solubilization of catalase.

The catalase activity, measured in 11 separate preparations of the AM freeze-thaw supernatant, was  $37.3 \times 10^{-2}$  U/10<sup>6</sup> cells with 1 SD of  $\pm 3.1 \times 10^{-2}$  U. This is approximately 6 times more than rabbit red cells catalase whose activity was  $6.3 \times 10^{-2}$  U/10<sup>6</sup> cells with 1 SD of  $0.7 \times 10^{-2}$  U. Peroxidase measurements in the AM showed remarkably little activity,  $6.8 \pm 0.9$  reciprocal seconds/10<sup>6</sup> cells. This may be contrasted with the measured peroxidase activity of human polymorphonuclear leucocytes, namely  $483 \pm 31$  reciprocal seconds/10<sup>6</sup> cells. The minimal peroxidase activity of the AM preparation may reflect PMN contamination of these preparations.

*The action of aminotriazole on catalase activity.* The effects of AT on the catalase activity of the AM are indicated in Table II. Concentrations of 100 and 20 mM produce approximately 85 and 66% inhibition of the catalase activity respectively.

Evidence that the inhibition of catalase by AT is dependent on the presence of H<sub>2</sub>O<sub>2</sub> (see Discussion) is provided by the data in Table III. The inhibition is shown to be of the irreversible type since it cannot be reversed by dialysis. Further, ethanol, which alone has no action on catalase, prevents the inhibitory action of subsequently added AT by decomposing catalase H<sub>2</sub>O<sub>2</sub> compound 1 to free catalase (9). Ethanol did not diminish the inhibition of catalase pretreated with AT providing further evidence that the AT inhibition is of the irreversible type.

*Particle entry in the presence of aminotriazole.* The results of the semiquantitative intracellular bacterial counts in the presence or absence of 100 mM AT are presented in Table IV. No differences were observed.

TABLE III  
Factors Influencing AT Inhibition of Catalase Activity

Experiment	Control	AT		Ethanol		
		No dialysis	Dialysis	Ethanol alone	AT after ethanol	AT before ethanol
1	0.75	0.04	0.04	0.75	0.73	0.04
2	0.75	0.05	0.06	0.74	0.72	0.05

Six aliquots of AM freeze-thaw extract were employed for each experiment. Duplicate measurements of catalase activity (milliequivalents of perborate used in 5 min) were obtained under the indicated conditions. AT and/or ethanol in final concentrations of 100 mM and 0.45 M respectively were added to the appropriate aliquots in the sequences indicated. The aliquots were allowed to stand at room temperatures for 30 min after each addition before assay or dialysis.

*Particle entry and catalase activity.* The possibility that particle entry increases  $H_2O_2$  levels in the AM was next examined. Table V indicates a comparison of the catalase activity measured on aliquots of the same AM preparations at "rest" and during phagocytosis in the presence and absence of AT.

The data demonstrate the marked increase in the inhibition of catalase by AT in the AM which have phagocytosed the bacteria. The increased inhibition by AT after phagocytosis suggests an increase in  $H_2O_2$  levels after phagocytosis. The possibility that this increased AT inhibition of catalase might be due to an increased intracellular AT concentration after phagocytosis is excluded by experiments 3 and 4 employing aminotriazole- $^{14}C$  (Table V). The counts due to labelled aminotriazole recovered from aliquots of the AM extract employed for measurement of catalase activity were similar in the extracts from both resting and phagocytosing AM.

In all four duplicate experiments there was a small increase (12%) in the control catalase activity after phagocytosis. This change in catalase activity after phagocytosis was not due to loss of catalase during the washing procedures since, while the pooled washing fluid contained 8% of the final AM catalase activity, the amount of catalase detected was similar in the washings from "resting" and phagocytosing AM. Further, since no catalase activity could be detected in heat-killed *S. Albus*, it seems unlikely that this increased catalase is derived from the phagocytosed bacteria. The significance of this increase in intracellular catalase activity after phagocytosis is unclear.

*Hydrogen peroxide.* The quantities of  $H_2O_2$  present in the dialysis fluid at varying incubation times in resting and phagocytosing AM are indicated in Fig. 1.

TABLE IV  
*The Effects of Aminotriazole on Bacterial Entry*

Experiment	Bacteria per cell	Control			Aminotriazole		
		1-5	6-10	>11	1-5	6-10	>11
1	No. of AM	9	32	9	7	28	15
2		12	28	10	14	23	13
3		11	33	6	9	28	13
Mean		11	31	8	10	26	14

The number of AM containing the indicated number of particles are presented. 50 cells were examined in each of three paired experiments. 20 mM aminotriazole.

There was a progressive rise in the quantity of  $H_2O_2$  recovered by dialysis as incubation is prolonged. Further the quantity recovered at any given time from phagocytosing AM was approximately double that observed in resting AM. In studies employing intact cells, up to 0.34  $\mu$ moles of  $H_2O_2$  were recovered from  $10^6$  AM during phagocytosis.

*The effects of aminotriazole on  $O_2$  utilization.* The effect of AT on  $Q_{O_2}$  in "resting" and phagocytosing AM are presented in Table VI. Each experiment was performed on aliquots obtained from the same AM preparations and thus each experiment was internally controlled. AT diminishes the  $Q_{O_2}$  in the resting AM. Though these changes are small, they are consistent in all experiments and the mean values are significantly different by Student's *t* test for paired samples at  $P < 0.01$  level. The maximum effect at concentrations of 0.1 M AT, was an approximately 14% reduction of  $Q_{O_2}$ . Particle entry consistently stimulated  $Q_{O_2}$  by 55%. ( $P < 0.01$ ). The effects of AT on the  $Q_{O_2}$  increase

TABLE V  
*The Effects of Phagocytosis on Catalase Activity and on Aminotriazole Inhibition of Catalase*

Experiment	Rest				Phagocytosis			
	Control	+AT	Inhibition %	AT- $^{14}C$ cpm	Control	+AT	Inhibition %	AT- $^{14}C$ cpm
1	1.00	0.48	52	—	1.20	0.10	75	—
	0.98	0.48			1.22	0.08		
2	1.00	0.54	48	—	1.12	0.29	60	—
	1.00	0.54			1.14	0.35		
3	1.48	1.23	16	191	1.67	0.33	80	209
				190				191
4	1.80	1.11	37	187	1.94	0.65	65	184
	1.76	1.10		192	1.84	0.62		208
Average	1.31	0.82	39	190	1.48	0.45	75	198

Aminotriazole (AT) 20 mM, AT- $^{14}C$  = 5-aminotriazole- $^{14}C$ , 0.01  $\mu$ Ci of AT- $^{14}C$  added. Duplicate measurements of catalase activity (milliequivalents of perborate utilized in 5 min) are indicated except for experiment 3. For procedures, see Methods.




FIGURE 1  $H_2O_2$  recovery from resting and phagocytosing AM.  $H_2O_2$  expressed as  $m\mu\text{moles}/10^6$  cells in dialysis bag. Mean  $\pm 1$  SE are indicated; numbers in parentheses are number of observations.

specifically associated with phagocytosis are evident from a comparison of the right-hand two columns of Table V. The  $Q_{O_2}$  due to phagocytosis is greater in the absence of AT than in the presence of AT. The statistical significance of the differences in  $Q_{O_2}$  due to phagocytosis in the presence and absence of AT was examined by an analysis of variance (17). This analysis demonstrated that the "interactions" relating  $Q_{O_2}$  in resting and phagocytosing AM in presence and absence of AT were statistically different at concentrations of 0.1 M AT ( $P < 0.01$ ) and at 0.04 M AT ( $P < 0.05$ ) but not at 0.02 M AT ( $P > 0.05$ ).

**Formate oxidation.**  $^{14}CO_2$  production from sodium formate- $^{14}C$  was employed to assess the activity of catalase/ $H_2O_2$  compounds in the intact AM. The stimulation of  $^{14}CO_2$  production from labeled formate by phagocytosis is demonstrated by the data in Table VII. The effects of AT on formate oxidation in the resting

and phagocytosing AM are also shown in Table VII. AT causes an inhibition of formate oxidation both in resting and phagocytosing AM at both 100 mM and 20 mM concentration ( $P < 0.01$  for both concentrations).

**Oxidation of glucose, pyruvate, and acetate.** The results of measurements of  $^{14}CO_2$  production from 1-glucose- $^{14}C$ , 6-glucose- $^{14}C$ , 1-pyruvate- $^{14}C$ , 1-acetate- $^{14}C$  in resting and phagocytosing AM are presented in Table VIII. Data obtained in the absence of AT indicated the following: (a) resting AM generate 3-6 times more  $^{14}CO_2$  from 1-glucose- $^{14}C$  than from 6-glucose- $^{14}C$ ; (b) phagocytosis stimulates  $^{14}CO_2$  production from both labeled forms of glucose, by a factor of varying between 2 and 4 times; (c)  $^{14}CO_2$  production from pyruvate is approximately doubled by phagocytosis; and (d)  $^{14}CO_2$  production from acetate is not stimulated by phagocytosis.

The effects of AT on  $^{14}CO_2$  production from these labeled substrates are also indicated in Table VIII. Concentrations of 100 mM AT diminished  $^{14}CO_2$  production from pyruvate and both  $^1C$  and  $^6C$ -labeled glucose. Concentrations of 20 mM AT diminished  $^{14}CO_2$  production from pyruvate alone. Neither concentration of AT affected acetate conversion to  $CO_2$ .

## DISCUSSION

**Evidence for the presence of  $H_2O_2$  metabolism.** The demonstration that catalase is present in high concentrations in the AM provides prima facie evidence for the existence of peroxidative metabolism. This was supported by using the catalase inhibitor, AT, and formate oxidation. The validity of these approaches is indicated by the following considerations. AT is known to produce two types of inhibition of catalase (10), a reversible inhibition of free catalase and an irreversible inhibition in which, according to Margoliash and Novogrodsky, the AT is bound to the protein moiety of catalase: $H_2O_2$  compound 1 (10). The distinction between these two types of inhibition is important, since the second type of inhibition has been used to detect the presence of  $H_2O_2$

TABLE VI  
The Effects of AT on  $Q_{O_2}$  in Resting and Phagocytosing AM

Concentration of AT	No. of Experiments	Resting		$\Delta Q_{O_2}$ due to AT in resting AM (A-B)	Phagocytosing		$\Delta Q_{O_2}$ due to phagocytosis AT absent (C-A)	$\Delta Q_{O_2}$ due to phagocytosis AT present (D-B)
		A Control	B AT		C Control	D AT		
<i>M</i>								
0.1	6	4.3 $\pm$ 1.3	3.7 $\pm$ 1.2	-0.6 $\pm$ 0.08	6.7 $\pm$ 1.4	5.1 $\pm$ 1.3	+2.4 $\pm$ 0.12	+1.4 $\pm$ 0.12
0.04	5	3.9 $\pm$ 0.61	3.6 $\pm$ 0.71	-0.3 $\pm$ 0.05	6.2 $\pm$ 0.56	5.6 $\pm$ 0.42	+2.3 $\pm$ 0.10	+2.0 $\pm$ 0.10
0.02	3	3.9 $\pm$ 0.41	3.7 $\pm$ 0.20	-0.2 $\pm$ 0.07	6.1 $\pm$ 0.42	5.7 $\pm$ 0.36	+2.2 $\pm$ 0.14	+2.0 $\pm$ 0.10

$Q_{O_2}$  expressed as  $\mu\text{l/hr}$  per  $10^6$  cells at  $37^\circ\text{C}$  and ambient pressure. Mean  $\pm 1$  SE are indicated for A thru D; mean  $\pm 1$  SE of differences between paired samples are indicated for  $\Delta Q_{O_2}$  values.

in biologic systems, notably the red cell (9). The evidence that the inhibition is of the irreversible type in the AM is twofold. First, dialysis of the AT-treated extract failed to modify the catalase inhibition. Thus, the AT was bound to some component of the extract, presumably catalase protein (10). Second, pretreatment of the AM with ethanol completely prevented the AT inhibition and ethanol is known to decompose catalase:  $H_2O_2$  compounds to free catalase and acetaldehyde (9, 18). These features of the AT inhibition of AM catalase, therefore, suggest the presence of catalase:  $H_2O_2$  compound I which only exists in the presence of low concentrations of  $H_2O_2$  (10). Evidence compatible with increased  $H_2O_2$  levels after phagocytosis has also been obtained from the observation that AM catalase is more sensitive to AT inhibition in the presence of intracellular bacteria (Table IV). While the use of AT to detect  $H_2O_2$  is indirect, it has the merit of demonstrating that  $H_2O_2$  exists intracellularly. Thus in those experiments (Table V) in which washed AM were employed for the preparation of freeze-thaw extracts, it was demonstrated that the AT was irreversibly bound to intracellular catalase. Comparable observations of increased catalase sensitivity to AT after phagocytosis in the PMN have been obtained by Reed (19).

Formate oxidation has been used as a marker of catalase-dependent  $H_2O_2$  metabolism in a number of tissues (20, 21) including the PMN (3, 5, 7) since Keilin and Hartree showed that catalase as opposed to a number of other hemoproteins was required for the coupled peroxidation of formate (18). The foregoing data indicate that formate oxidation by AM is almost completely inhibited by 100 mM AT, and is therefore presumably largely catalase-dependent. Phagocytosis has been shown to increase formate oxidation (Table VII) indicating increased  $H_2O_2$  formation during phagocytosis.

TABLE VII  
The Effects of Aminotriazole of  $^{14}CO_2$  Production from Formate- $^{14}C$

Concentration of AT	Resting		Phagocytosing	
	Control	AT	Control	AT
<i>M</i>				
0.1	36.8 ± 12.1	7.9 ± 0.5	108.8 ± 25.0	3.9 ± 0.4
0.02	50.8 ± 5.1	23.0 ± 2.7	96.9 ± 21.0	32.5 ± 3.6

Data expressed as cpm/10<sup>6</sup> cells per 200,000 cpm added to medium. Mean ± 1 SE of two triplicate experiments are indicated.

This increase is quantitatively similar to that observed in the PMN (3).

The assay of  $H_2O_2$  (Fig. 1) confirms that  $H_2O_2$  is produced by resting AM. The increased quantity of  $H_2O_2$  obtained from AM during phagocytosis suggests increased intracellular  $H_2O_2$  concentrations. While changes in cell permeability to  $H_2O_2$  during phagocytosis could explain the increased  $H_2O_2$  observed during phagocytosis, the preceding lines of evidence suggest that increased intracellular  $H_2O_2$  levels are the correct explanation. Our data for the AM are similar to those of Paul and Sbarra (6) who obtained values of 0.054 nmoles/10<sup>6</sup> cells from resting PMN and of 0.092 nmoles/10<sup>6</sup> cells from PMN phagocytosing *S. albus* after 30-min incubation periods.

It is noteworthy that the three indices of  $H_2O_2$  formation, namely AT inhibition of catalase, formate oxidation, and  $H_2O_2$  present in dialysates are all increased approximately twofold during phagocytosis.

*Relation between peroxidative and glucose metabolism.* The classical studies of Iyer and colleagues (3) on PMN first suggested the relationship between increased  $Q_{O_2}$ , increased  $H_2O_2$  formation and increased glucose oxida-

TABLE VIII  
The Effects of AT on  $^{14}CO_2$  Production from Substrates- $^{14}C$

Substrate	No. of experiments	Concentration of AT	Resting			Phagocytosing		
			Control	AT	P value	Control	AT	P value
<i>M</i>								
1-Glucose- $^{14}C$	4	0.1	22 ± 7.1	8 ± 3.1	<0.05	90 ± 20.1	51 ± 16.1	<0.01
	2	0.02	23 ± 1.2	23 ± 2.1	>0.1	76 ± 3.1	70 ± 4.2	>0.1
6-Glucose- $^{14}C$	3	0.1	2.3 ± 0.3	0.5 ± 0.1	<0.01	9.1 ± 0.7	1.7 ± 0.7	<0.01
	2	0.02	4.7 ± 0.8	4.0 ± 0.6	>0.1	10.5 ± 1.3	10.1 ± 1.3	>0.1
1-Pyruvate- $^{14}C$	3	0.1	162 ± 14	101 ± 6	<0.01	381 ± 21	130 ± 14	<0.01
	3	0.02	104 ± 10	83 ± 6	<0.05	179 ± 19	116 ± 12	<0.01
1-Acetate- $^{14}C$	3	0.1	35 ± 6	32 ± 5	>0.1	38 ± 8	30 ± 6	>0.1
	2	0.02	25 ± 8	25 ± 7	>0.1	23 ± 5	27 ± 5	>0.1

Data expressed as cpm/10<sup>6</sup> cells per hr of 200,000 initial counts; mean ± 1 SE are indicated.

tion during phagocytosis. Our studies indicate a similar relationship in the AM by providing evidence of increased  $H_2O_2$  metabolism during phagocytosis and by confirming the studies of Ouchi, Selvaraj, and Sbarra (2) on  $Q_{O_2}$  and glucose metabolism. The latter also demonstrated that phagocytosis by rabbit AM stimulates  $Q_{O_2}$  and  $^{14}CO_2$  production from labeled glucose and pyruvate. Their observation that 1-4-succinate- $^{14}C$  oxidation is unaffected by phagocytosis is in accord with our observations on acetate oxidation since both produce  $CO_2$  specifically via the Krebs cycle. Both sets of observations are different from the results of Oren, Farnham, Saito, Milofsky, and Karnovsky (22), who observed smaller increases in  $Q_{O_2}$  and glucose conversion to  $CO_2$  during phagocytosis by guinea pig AM. These differences may reflect either species differences or a difference in cell harvesting technique in that Oren employed minced guinea pig lung. Thus, at least in rabbit AM, there is an increase in  $Q_{O_2}$ , glucose oxidation, and  $H_2O_2$  levels during phagocytosis and analogy with the PMN suggests these to be inter-related.

The definition of the role of catalase in these interrelations must await knowledge of the  $H_2O_2$  generating pathway. Further communications<sup>3</sup> will provide evidence that, in the AM,  $H_2O_2$  is in part generated by a D-amino acid oxidase, an enzyme recently described in PMN (23), and also that the  $H_2O_2$  is coupled to triphosphopyridine nucleotide, reduced form of, (NADPH) in the AM by a glutathione system (24), similar to that described in the rat leucocyte (19). However, two comments may be made on the basis of the current studies. First, the virtual absence of myeloperoxidase activity precludes it serving as an  $H_2O_2$ -generating NADPH oxidase in the AM as has been proposed for the PMN (25). Second, while caution is required in interpreting evidence based on the actions of AT, it is possible that the actions of AT are relatively specific for catalase. At least AT has no general toxic actions on the AM since particle entry, cell viability (eosin uptake technique), acetate oxidation (Table VIII), and ATP concentrations (1) are uninfluenced by 100 mM AT. On the assumption of the catalase specificity two alternative explanations of the AT inhibition of  $Q_{O_2}$  and  $^{14}CO_2$  production from glucose and pyruvate appear reasonable. First, catalase activity may determine the  $H_2O_2$  production by an oxidase. The importance of the association of  $H_2O_2$ -generating oxidases and catalase has recently been stressed by deDuve and Baudhuin (21). Second, catalase may serve in coupled peroxidations linked to diphosphopyridine nucleotide, reduced form of, (NADH), as demonstrated by several workers (18, 20, 21) and NADH

<sup>3</sup> Vogt, M., C. Thomas, J. B. L. Gee, and R. E. Basford. Unpublished observations.

is a known regulator of glycolysis and pyruvate metabolism.

*Significance of peroxidative metabolism.* It is evident that inhibition of catalase activity does not seriously affect particle entry. Increased  $H_2O_2$  metabolism, therefore, represents a secondary event after particle entry. Peroxidative metabolism represents a mechanism capable of increasing glucose metabolism in a manner similar to that observed in the liver, kidney, and red cells (20, 21, 26, 27). Finally, the role of  $H_2O_2$  as an intracellular bactericidal agent in PMN has been recently emphasized (6, 7, 8). The tempting idea that  $H_2O_2$  serves a similar role in the AM requires further study particularly since the AM lack white cell peroxidase activity. This enzyme represents an important component of the bactericidal peroxidative systems (8, 28, 29). Deficiency of this peroxidase in PMN is associated with impairment of Candidacidal activity and subnormal killing rates of certain bacteria (30-32).

#### ACKNOWLEDGMENTS

We gratefully acknowledge the helpful discussions with Dr. E. D. Robin and the skilled technical help of Miss Joan Yogodnik.

This work was supported by Grants IR01 U1-00850, Health Service and Research Foundation Grant, and Training Grant HE 05706.

#### REFERENCES

1. Gee, J. B. L., E. D. Robin, J. B. Field, J. D. Smith, A. R. Tanser, and J. Kaskin. 1968. Roles of ATP and of peroxidative metabolism during phagocytosis in alveolar macrophages *J. Clin. Invest.* **47**: 38a.
2. Ouchi, E., R. J. Selvaraj, and A. J. Sbarra. 1965. The biochemical activities of rabbit alveolar macrophages during phagocytosis. *Exp. Cell Res.* **40**: 456.
3. Iyer, G. Y. N., M. F. Islam, and J. H. Quastel. 1961. Biochemical aspects of phagocytosis. *Nature (London)*. **192**: 535.
4. Karnovsky, M. L. 1962. Metabolic basis of phagocytic activity. *Physiol. Rev.* **42**: 143.
5. Roberts, J., and J. H. Quastel. 1964. Oxidation of reduced triphosphopyridine nucleotide by guinea pig polymorphonuclear leucocytes. *Nature (London)*. **202**: 85.
6. Paul, B., and A. J. Sbarra. 1968. The role of the phagocyte in host-parasite interactions. XIII. The direct quantitative estimation of  $H_2O_2$  in phagocytising cells. *Biochem. Biophys. Acta.* **156**: 168.
7. Holmes, B., A. R. Page, and R. A. Good. 1967. Studies of the metabolic activity of leucocytes from patients with a genetic abnormality of phagocytic function. *J. Clin. Invest.* **46**: 1422.
8. McRipley, R. J., and A. J. Sbarra. 1967. Role of the phagocyte in host-parasite interactions. XII. Hydrogen peroxide-myeloperoxidase bactericidal system in the phagocyte. *J. Bacteriol.* **94**: 1425.
9. Cohen, G., and P. Hochstein. 1964. Generation of hydrogen peroxide in erythrocytes by hemolytic agents. *Biochemistry*. **3**: 895.

10. Margoliash, E., and A. Novogrodsky. 1958. A study of the inhibition of catalase by 3-amino-1,2,4-triazole. *Biochem. J.* **68**: 468.
11. Myrvik, Q. N., E. S. Leake, and B. Fariss. 1961. Studies on pulmonary alveolar macrophages from the normal rabbit: a technique to procure them in a high state of purity. *J. Immunol.* **86**: 128.
12. Umbreit, J. W., R. H. Burris, and J. F. Stauffer. 1964. Preparation of tissues. In *Manometric Techniques*. J. W. Umbreit, R. H. Burris, J. F. Stauffer, editors. Burgess Publishing Co., Minneapolis, Minn. 149.
13. Feinstein, R. N. 1949. Perborate as substrate in a new assay of catalase. *J. Biol. Chem.* **180**: 1197.
14. Bray, G. A. 1960. A simple efficient liquid scintillation for counting aqueous solutions in a liquid scintillation counter. *Anal. Biochem.* **1**: 279.
15. Chance, B., and A. C. Maehly. 1955. Assay of catalases and peroxidase. In *Methods in Enzymology*. S. Colowick, editor. Academic Press, Inc., New York. **2**: 764.
16. Brandt, R., and A. S. Keston. 1965. Synthesis of diacetyl-dichlorofluorescein: a stable reagent for fluorometric analysis. *Anal. Biochem.* **11**: 6.
17. Edwards, A. L. 1964. Experimental design in psychological research. Holt, Rinehart and Winston, New York. Chap. 9.
18. Keilin, D., and E. F. Hartree. 1955. Catalase, peroxidase and metmyoglobin as catalysts of coupled peroxidatic reactions. *Biochem. J.* **60**: 310.
19. Reed, P. W. 1969. Glutathione and the hexose monophosphate shunt in phagocytising and hydrogen peroxide treated rat leukocytes. *J. Biol. Chem.* **244**: 2459.
20. Portwich, F., and H. Aebi. 1960. Erfassung der Peroxybildung tierischer Gewebe mittels peroxydatischer Umsetzung. *Helv. Physiol. Pharmacol. Acta.* **18**: 312.
21. deDuve, C., and P. Baudhuin. 1966. Peroxisomes (microbodies and related particles). *Physiol. Rev.* **46**: 323.
22. Oren, R., A. E. Farnham, K. Saito, E. Milofsky, and M. L. Karnovsky. 1963. Metabolic patterns in three types of phagocytising cells. *J. Cell Biol.* **17**: 487.
23. Cline, M. J., and R. I. Lehrer. 1969. D-amino acid oxidase in leucocytes: a possible D-amino-acid-linked antimicrobial system. *Proc. Nat. Acad. Sci. U. S. A.* **62**: 756.
24. Gee, J. B. L., and R. E. Basford. 1969. Effects of sulfhydryl inhibitors on phagocytosis by alveolar macrophages (AM). *Fed. Proc.* **28**: 445.
25. Roberts, J., and Z. Camacho. 1967. Oxidation of NADPH by polymorphonuclear leucocytes during phagocytosis. *Nature (London)*. **216**: 606.
26. Hochstein, P., and H. Utley. 1968. Hydrogen peroxide detoxication by glutathione peroxidase and catalase in rat liver homogenates. *Mol. Pharmacol.* **4**: 574.
27. Jacob, H. S., and J. H. Jandl. 1966. Effects of sulfhydryl inhibition on red blood cells. III. Glutathione in the regulation of the hexose monophosphate pathway. *J. Biol. Chem.* **241**: 4243.
28. Klebanoff, S. J. 1967. Iodination of bacteria: a bactericidal mechanism. *J. Exp. Med.* **126**: 1063.
29. Klebanoff, S. J., and L. R. White. 1969. Iodination defect in the leucocytes of a patient with chronic granulomatous disease of childhood. *N. Engl. J. Med.* **280**: 460.
30. Lehrer, R. I., and M. J. Cline. 1969. Leukocyte myeloperoxidase deficiency and disseminated candidiasis: the role of myeloperoxidase in resistance to *Candida* infection. *J. Clin. Invest.* **48**: 1478.
31. Lehrer, R. I., J. Hanifin, and M. J. Cline. 1969. Defective bactericidal activity in myeloperoxidase-deficient human neutrophils. *Nature (London)*. **223**: 78.
32. Salmon, S. E., M. J. Cline, J. Schultz, and R. I. Lehrer. 1970. Myeloperoxidase deficiency: immunologic study of a genetic leukocyte defect. *N. Engl. J. Med.* **282**: 250.