



REVIEW ARTICLE

OPEN ACCESS

Cataract: A major secondary diabetic complication

*Piyush Patel^{1,2}, Nurudin Jivani³, Shailesh Malaviya², Tushar Gohil², Yagnik Bhalodia³

¹School of Pharmacy, RK University, Bhavnagar Highway, Kasturbadham, Rajkot-360 020, Gujarat, India

²Rajkot-Bhavanagr Highway, Smt. R.B. Patel Mahila Pharmacy College, Atkot-360 040, Gujarat, India

³Surendranagar-Ahmedabad Highway, C.U. Shah College of Pharmacy and Research, Wadhawan-363 030, Gujarat, India

ABSTRACT

Cataract is a visual impairment occurs due to the opacification of crystalline lens. It affects around 17 billion peoples worldwide, although incidence of cataracts is increasing day by day among the elderly persons. Still today except surgery no other effective treatment have been successfully developed so far, thus this present review is focused to highlights the etiological aspects, risk factors along with possible pharmacological prevention and animal models which are widely used for screening anticataract activity. The present review includes the list of plants and their phytoconstituents which have been evaluated pharmacologically for the treatment of cataract. From the review it can be concluded that antioxidant properties of plants phytochemicals are responsible for the anticataract activity.

Key Words: Cataract, aldose reductase, galactose, selenite, antioxidant.

INTRODUCTION

A cataract is opacity of the lens that interferes with vision, and is the most frequent cause of visual impairment worldwide, especially for the elderly because the incidence of cataracts increases with increasing age. From a public health perspective, it is important to identify the risk factors that affect the development and progression of cataract. Cataract is a multifactorial disease process and is induced by various toxic factors, environmental, stressors, and gene mutations. Cataract can be classified into four types; nuclear, cortical, posterior subcapsular (PSC), and mixed. Although the etiology of each cataract type remains elusive, cataracts are known from studies on many animal models and humans to be associated with damage or death of lens epithelial cells (LECs) (Li *et al.*, 1995). Cataract is the leading cause of blindness worldwide and covers around 42% of overall visual impairment. Cataract is mainly responsible for almost 80% blindness in India (Gupta *et al.*, 2009). Cataract is also produces by the advanced glycation

end products (AGE) is cause of blindness worldwide (Thiagarajan *et al.*, 2003). AGE related cataract is leading cause of blindness and visual impairment worldwide. Chronic hyperglycemia can cause various types of secondary diabetic complications including diabetic cataract. Form literature it can be found that hyperglycemia increase the risk of development of cataract. Even though aging is another factor for development of cataract, other risk factor such as nutritional deficiencies, trace metals, exposure to sunlight, smoking are also responsible for development of cataract.

CAUSES OF CATARACT

Various factor such as diabetes, oxidation of lens, dehydration, daylight, diet and lipid peroxidation attributes to the generation of lens opacification in elderly persons (Kothadia *et al.*, 2011). Other risk factors such as smoking, environmental factor, lack of consumption of antioxidants, nutritional deficiency and diabetes can also increase the risk of development of cataract. During hyperglycemia extracellular glucose diffuse into the lens, which can leads to the post-translational modification. Cataract progress forms the synthesis and accumulation of excessive sorbitol in the lens fibre and consequent osmotic stress. Sorbitol is synthesis from aldose

*Corresponding Author:

Piyush M. Patel, Assistant Professor
Smt. R.B. Patel Mahila Pharmacy College
Kailash Nagar, Bhavnagar-High Way
Atkot- 360 040, Rajkot, India
E-mail: pyush_28@yahoo.co.in
Contact No.: (02821)288349

reductase utilising the NADPH and does not cross the cell membranes; it can accumulate in the cells and can cause cell damage due to disturbing osmotic homeostasis (Gupta *et al.*, 2009). Deficient glutathione is another mechanism behind the formation of cataract. In cataract reduced glutathione level were found to be significantly reduced when compared to normal (Thiagarajan *et al.*, 2003). Oxidative stress is another mechanism involve in development of cataract, which cause the oxidation of lens protein. Reduction in concentration of glutathione, antioxidant enzymes such as catalase, superoxide dismutase (SOD), glutathione Reductase, glutathione peroxidase with increase in age where the main factor involving the generation of cataract (Graw, 2009). Diabetes causes the increased level of oxidised DNA, proteins and lipids, which are also limiting factor in various diabetic complications (Suryanarayana *et al.*, 2005). Glucose autoxidation, formation of AGE, and activation of polyol pathway cause the intracellular accumulation of sorbitol, which cause the various types of ocular lesions, alteration of membrane permeability, loss of glutathione and diminuation of protein synthesis. Also diabetic patient has polymorphism in the promoter region of aldose reductase (ARL2). gene, causing the cataract, retinopathy, and neuropathy. Production of hydrogen peroxide through glucose auto-oxidation is also associated with cataract formation (Pastene *et al.*, 2007). Aldose Reductase key enzyme of polyol pathway catalyses the reduction of glucose into sugar alcohol sorbitol, which is subsequently metabolised to fructose by sorbitol dehydrogenase. Sorbitol as an osmolyte that leads to osmotic swelling, changes in the membrane permeability, leakage of glutathione, myo-inositol, the generation of free radicals and hydrogen peroxide which primarily causing the diabetic complication such as cataract, retinopathy and neuropathy (Jung *et al.*, 2011). Hydrogen peroxide at higher concentration can cause lens opacification and tissue damage similar to that found in human cataract (Javadzadeh *et al.*, 2009)

PHARMACOLOGICAL PREVENTION

There is sufficient evidence that oxidative stress play a pilot role in mechanism of cataractogenesis, hence there is increasing interest in developing suitable antioxidant nutrient from natural or

synthetic origin that could be effective in delaying or prevention the progression of cataract (Awasthi *et al.*, 1996). Presently few medicines, eye drops, exercise and glass are available to prevent the cataract. The symptom of early cataract can be improved by new eyeglasses, anti-glare sunglasses or magnifying lens, and if these measures do not help surgery is last effective treatment (Gupta *et al.*, 2009). From earlier studies, it was found that person consuming diet with high content of antioxidants can reduce the development of cataract (Jacques *et al.*, 1991). The plants and plant-derived products have been used as therapeutics in traditional medicine due to their less adverse effects and economical benefit. Currently the only available treatment for disease is surgical removal of cataractous lens and followed by replacement with synthetic implants. Efforts have been taken to explore the traditional medicine to delay and retard the progression of cataract. Several numbers of plants and synthetic compounds has been reported to possess anti-cataract activity (Rooban *et al.*, 2010).

ANIMAL MODEL TO ASSESS ANTI-CATARACT ACTIVITY

Various animal models have been used for the screening the plants against cataract. Out of that galactose induced cataract is commonly used, it produces the large amount of reduced form, galactitol and finally into glucose. Furthermore galactitol is not subsequently metabolised as compared to the sorbitol. In this model it is supposed that factor initiating galactose cataracts in young rats are similar to those involve in the human galactose cataract model. Three mechanism involve in the formation of cataract are oxidation, polyol pathway and non enzymatic glycation (Kinoshita, 1990). Naphthalene induced cataract is similiary used to test herbal drugs as anticataract due to its similarities to that of age related cataract in humans. Naphthalene is converted into stable compound naphthalene-1,2-dihydrodiol and its metabolism to 1,2-napthoquinone by an enzyme dihydrodiol dehydrogenase. Epithelial mitochondrion is the target of 1,2-nathoquinone toxicity. Formation of the 1,2- napthoquinone is considered as underlying mechanism of cataract development in naphthalene induced animal model. Aldose reductase is the key enzyme in the metabolism of naphthalene

Table 1: List of plants with anticataract potential.

Plant name with family	Parts of plant and extract/fraction	Mechanism of action	References
<i>Adhatoda vasica</i> (Acanthaceae)	Aqueous, ethanol and chloroform extracts	Aldose reductase activity	(Cacche and Dhole, 2011)
<i>Allium cepa</i> (Liliaceae)	Juice	Reduces the oxidative stress	(Javadzadeh <i>et al.</i> , 2009)
<i>Cassia fistula</i> (Acanthaceae)	Aqueous, ethanolic and chloroform extracts	Aldose reductase activity	(Cacche and Dhole, 2011)
<i>Angelica dahurica</i> (Umbellifereae)	byakangeleisin and furanocoumarin from the roots	Inhibits the galactitol and sorbitol accumulation	(Shin <i>et al.</i> , 1998)
<i>Citrus aurantium</i> (Rutaceae)	Hydromethanol peel extracts	Prevent the peroxidative damage	(Umamaheshwari <i>et al.</i> , 2011)
<i>Cochlospermum religiosum</i> (Cochlospermaceae)	Isorhmnetin -3-glucoside, bioactive flavonoids form leaves	Inhibits the oxidative stress	(Devi <i>et al.</i> , 2010)
<i>Emilia sonchifolia</i> (Asteraceae)	Flavonoids	Inhibits the oxidative damage and lens opacification	(Lija <i>et al.</i> , 2006)
<i>Erigeron annuus</i> (Asteraceae)	Caffeoylquinic acids and four flavonoids from ethyl acetate soluble fractions of stems and leaves	Prevent the lens opacity	(Jang <i>et al.</i> , 2010)
<i>Ginko biloba</i> (Ginkgoaceae)	Extract of plant	Inhibit the oxidative stress	(Ertekin <i>et al.</i> , 2004)
<i>Momordica charantia</i> (Cucurbitaceae)	lyophilized extracts	Inhibit the oxidative damage	(Rathi <i>et al.</i> , 2002)
<i>Moringa oliefera</i> (Moringaceae)	flavonoids fractions of leaves	Inhibit the oxidative damage and reduces the lipid peroxidation.	(Sasikala <i>et al.</i> , 2010)
<i>Ocimum sanctum</i> (Lamiaceae)	Extract of plant	inhibition of polyol accumulation	(Halder <i>et al.</i> , 2003)
<i>Origanum vulgare</i>	Extract of plant	Inhibit the oxidative damage and reduces the lipid peroxidation.	(Dailami <i>et al.</i> , 2010)
<i>Pterocarpus marsupium</i> (Fabaceae)	aqueous extract	Inhibits the diabetes induced cataract	(Vats <i>et al.</i> , 2004)
<i>Silybum marianum</i> (Compositae)	seeds extract	Inhibit the oxidative damage	(Huseini <i>et al.</i> , 2004)
<i>Trigonella foenum graecum</i> (Fabaceae)	Seed extract	Inhibit the generated oxidative stress	(Lija <i>et al.</i> , 2006)
<i>Vitex negundo</i> (verbanaceae)	flavonoids fraction	Inhibits the oxidative stress	(Rooban <i>et al.</i> , 2011)
<i>Withania somnifera</i> (solanaceae)	Extract of plant	Inhibits the oxidative stress	(Thiagarajan <i>et al.</i> , 2003)

(Germansky and Jamall, 1988). Selenite induced cataract is model for oxidative stress induced cataract. Selenite cataract is occurs due to the acute exposure of the selenite to the lens at particular age which cause the nuclear cataract (Huang *et al.*, 1982). Furthermore selenite induced oxidative stress causes the nuclear opacity through the calpain proteolysis of lens protein. It is strong sulfhydryl oxidant and it is resembles cataract caused by oxidative stress. Similar to human senile cataract, this type of cataract is accompanied by decrease in activities of the antioxidants enzymes such as

superoxide dismutase, glutathione peroxidase (Shearer *et al.*, 1997).

PLANT PRODUCTS AGAINST CATARACTS

Based on literature assessment from earlier studies it was found that disturbance in the oxidative state in lens can be corrected by giving antioxidant such as ascorbic acid and besides also in body there are enzymatic and non enzymatic antioxidant system to protect cellular organelles from free radicals (Bhuyan *et al.*, 1981). Hyperglycemia generates the

reactive oxygen species, which causes the lipid peroxidations and thus play major role in development of diabetic secondary complications in kidney, blood vessels, eye and nerve (Patel *et al.*, 2011). The natural antioxidant such as ascorbic acid, vitamin E, and carotinoids have been screened and it was found that antioxidant has protective action against cataract. Flavonoids found in various food and beverages have strong antioxidant and anticataract activity (Durukan *et al.*, 2006; Patel *et al.*, 2011). one isoflavonoid glycoside isolated from ethyl acetate soluble fraction of *Viola hondoensis* was screened for aldose reductase inhibitory activity and it was found that it possess potent aldose reductase inhibitory potential (Chung *et al.*, 2008). Tea a major source of dietary quercetin and other flavonoids had been reported to have anticataract activity; and ability of isoflavone genistein in delaying the progression of cataract induced by the dietary galactose has been reported (Vinson and Zhang, 2005). Therefore development of pharmacologically and biochemically tested compound would be of great importance for the management of diabetic complication.

PLANTS WITH ANTICATARACT ACTIVITY

A list of plants with anticataract potential is given in table 1.

DISCUSSION

Cataract is disorder associated with several risk factors such as diabetes, aging etc. For the treatment of cataract there is no medicine available except surgery. However it has also some complication such as posterior capsular opacification, endophthalmitis and uncontrolled residual refractive errors (Javadzadeh *et al.*, 2009). So this type of therapeutic approach can reduces health burden. In last few years researchers are trying to find out the phytoconstituents that could beneficial against the cataractogenesis. From earlier studies it was found that *Ocimum sanctum* possess significant antioxidant potential with anticataract activity (Devi *et al.*, 2010). Reduction in glutathione level is another mechanism for retardation of cataract. Several phytoconstituents from herbal origin may indirectly inhibits the consumption of glutathione or it can stimulate the glutathione synthesis which may be

due to modulating effect on enzymes that affects glutathione level in lens architecture. In another study, it was found that *Trigonella foenum-graceum* can enhance the glutathione level due to inhibition of consumption of glutathione (Gupta *et al.*, 2010). Several factors are involved in generation of cataract, but exact mechanism of cataract formation is still unknown. Several researchers are trying to determine the mechanism of cataract formation by use of different *In vitro* and *In vivo* models. Amongst all the experimental models, the galactose induced cataract model is widely used due to similarity of this model to that of galactose induced human cataractogenesis (Gupta *et al.*, 2009; Suryanarayana *et al.*, 2005). Human cataract is not single disease but it is group of disease involve many risk factor. Oxidative stress is main mechanism behind cataract formation and in which antioxidant level is found to be less as compared to normal person. According to previous studies, it was found that significant increased level of free radicals and low concentration of glutathione content were observed in aqueous humor. Many study shows that flavonoids, a group of phytoconstituents with antioxidant properties can prevent the oxidative damage and cataract progression. Onion a flavonoids rich staple food is example in which major flavonoids are identified for antioxidant properties including quercetin, quercetin-4-glucoside and quercetin-3,4'-diglucoside (Javadzadeh *et al.*, 2009). According to world health organisation (WHO) reports it seems that about 80% of world population are uses herbal medicine for health needs (Kim *et al.*, 2010). Many studies have done to identify the natural or synthetic compound that inhibit aldose reductase and flavonoids are found to most potent aldose reductase inhibitors (Constantino *et al.*, 1999). Plant phytochemicals including polyphenols, which are currently regarded as natural antioxidant they are important for human health against various types of enzymes, which reduces the blood glucose level in diabetes, and are capable of reducing the oxidative stress by scavenging the free radicals and prevention cell damage (Vishvanath *et al.*, 2010; Girija *et al.*, 2011).

CONCLUSION

This review provides the relevant update information about the pathophysiological aspects of cataract, risk factor involved and suitable treatment available for cataract. It also provides the database of plants and their phytoconstituents responsible for anticataract activity. Since eye is unique organ, it is constantly exposed to oxidative stress and protection of lens from these is critical for essential. So far, no single compound has found wide spread acceptance for these indication, although many compound have been screened. Thus, there is need for further research in this area. Therefore, this review enables as an important segment for development of effective medicine for treatment of cataract.

REFERENCES

- Awasthi S, Srivatava SK, Piper J T, singhal SS, chaubey M, Awasthi YC, (1996). Curcumin protects against 4-Hydroxy-2-transnonenal induced cataract formation in rat lens. *American Journal of Clinical Nutrition*, 64: 761-766. PMID:8901798
- Bhuyan, K.C., Bhuyan, D.K., Podos, S.M., (1981). Cataract induced by selenium in rat. II. Increased lipid peroxidation and impairment of enzymatic defenses against oxidative damage. *IRCS Medical Science*, 9, 195-196.
- Cacche RN, Dhole NA, (2011). Aldose reductase inhibitory, anticataract and antioxidant potential of selected medicinal plants from marathawada region, india. *Natural Product Research*, 25:760-763. [DOI](#) PMID:21462076
- Chung IM, Kim MY, Park WH, Moon HI (2008). Aldose reductase inhibitor from *viola hondoensis* *American Journal of Chinese Medicine*, 36:799-803. [DOI](#) PMID:18711775
- Constantino, L., Rastelli, G., Vianello, P., Cignarella, G., & Barlocco, D (1999). Diabetes complications and their potential prevention: aldose reductase inhibition and other approaches. *Medical Research Reviews*, 19, 3- 23. [DOI](#)
- Dailami K N, Azadbakht M, Pharm Z R, Lasgari M. (2010). Prevention of selenite induced cataractogenesis by *origanum vulgare* extract. *Pakistan Journal of Biological Science*. 13:743-747. [DOI](#) PMID:21850936
- Devi VG, Rooban BN, Sasikala V, Sahasranaman AT, Abraham A (2010). Isorhamnetin-3-glucoside alleviates oxidative stress and opacification in senile cataract in vitro. *Toxicology In vitro*, 24:1662-1669.
- Durukan AH, Evreklioglu C, hurmeric V, Kerimoglu H, Erdurman C, Bayraktar MZ (2006). Ingestion of IH636 grape seeds proanthocyanidin extract to prevent selenite induced oxidative stress in experimental cataract. *Journal of Cataract and Refractive Surgery*, 32: 1041-1045. [DOI](#)
- Ertekin M V, Kocer I, Karslioglu I, Taysi S, gepdiremen A, Sezen O (2004). Effects of oral Ginkgo biloba supplementation on cataract formation and oxidative stress occurring in lens of rats exposed to cranium radiotherapy. *Japanese Journal of Ophthalmology*, 48:499-502. [DOI](#) PMID:15486777
- Germansky, M.; Jamall, S (1988.). Organ-specific effects of naphthalene on tissue peroxidation, glutathione peroxidases, and superoxide dismutase in the rat. *Archives of Toxicology*, 61:480-483. [DOI](#) PMID:3190445
- Girija K, Lakshman K, chandrika U, Ghosh SS, divya T (2011). Antidiabetic and anticholesterolemic activity of methanol extracts of three species of *Amaranthus* Asian Pacific *Journal Tropical Biomedicine*, 11:133-138.
- Graw J (2009). Mouse model of cataract. *Journal of genetics*, 88:469-486. [DOI](#) PMID:20090208
- Gupta S K, Kalaiselvan A, Srivastava S, Saxena R, Agrawal S S, (2010). *Trigonella Foenum graecum* (fenugreek seed). protects against selenite induced oxidative stress in experimental cataractogenesis *Biological Trace Element Research*, 136:258-268. [DOI](#) PMID:19823776
- Gupta SK, Kalaiselvan V, Saxena R, Agrawal SS (2009). Advances in pharmacological strategies for the prevention of cataract development. *Indian Journal of Ophthalmology*, 57:175-83. [DOI](#) PMID:19384010 PMID:2683438
- Gupta SK, Kalaiselvan V, Sharma A, Srivastava S (2009). Anticataract potential of *phyllanthus niruri* in galactose induced cataractogenesis of rat. *International Journal of Ophthalmology*, 2; 286-289.
- Gupta SK, Kalaiselvan V, srivastava S, Saxena R, Agrawal SS, (2009). Inhibitory effect of *trigonella foenum-graecum* on galactose induced cataract in rat. In vitro and in vivo studies. *Journal of Ophthalmic and Vision Research*, 4; 213-219.
- Halder N, Joshi S, Gupta S K, (2003). Lens aldose reductase inhibiting potential of some indigenous plants. *Journal of Ethnopharmacology*, 86:113-116. [DOI](#)
- Hsieh P, Huang G, Ho Y, Lin Y, Huang S, Chiang Y (2010). Activities of antioxidants α -glucosidase inhibitors and aldose reductase inhibitors on aqueous extracts of four *Flemingia* species in Taiwan. *Botanical Studies*, 51:293-302.
- Huang, L.L., Zhang, C.Y., Hess, C.Y., (1982). Biochemical changes and cataract formation in lenses in rats receiving multiple low doses of sodium selenite. *Experimental Eye Research*, 55, 671-678. [DOI](#)
- Huseini H F, Zaree A B, Zarch AB, Heshmat R, (2004). The effect of herbal medicine *silybum marianum* (L.). Gaerth. Seed extract on galactose induced cataract formation in rats. *Journal of Medicinal Plants Research*, 3: 58-62.

- Jacques, P.F., & Taylor, A. (1991). Micronutrients and age-related cataracts, In: A. Bendich, & C. E. Butterworth (Eds.), *Micronutrients in health and in disease prevention*, New York: Dekker. Pp. 359-379.
- Jang, D.S., Yoo, N.H., Kim, N.H., Lee, Y.M., Kim, C.S., Kim, J. (2010). 3,5-di-o-caffeyol-epi-quinic acid from the leaves and stems of *Erigeron annuus* inhibits protein glycation, aldose reductase, and cataractogenesis. *Biological and Pharmaceutical Bulletin*, 33: 329-333. [\[DOI\]](#)
- Javadzadeh A, Ghorbanihaghjo A, Bonyadi S, Rashidi M R, Mesgari M, Rashtchizadeh N, et al. (2009). Preventive effect of onion juice on selenite induced cataract. *Indian Journal of Ophthalmology*, 57:185-189. [\[DOI\]](#) PMID:19384011 PMCID:2683439
- Jung HA, Isla MD, Kwon YS, Jin SE, son YK, Park JJ (2011). Extraction and identification of three major aldose Reductase inhibitors from *Artemisia Montana* food chem. *Toxicology*, 49; 376-384.
- Kim H M, Mok S, Lee JM, Che EJ, Choi K, Ku J (2010). Inhibition of aldose reductase from rat lenses by methanol extracts from Korean folk plants. *Journal of natural products*, 16: 285-290.
- Kinoshita, J. H (1990). A thirty year journey in the polyol pathway. *Experimental Eye Research*, 50, 567-573. [\[DOI\]](#)
- Kothadia AD, shenoy AM, Shabaraya AR, Rajan MS, Viradia UM, Patel NH, (2011) Evaluation of cataract prevention action of phycocyanin. *International Journal of pharmaceutical Science and Drug Research*, 3; 42-44.
- Li WC, Kuszak JR, Dunn K, et al (1995). Lens epithelial cell apoptosis appears to be a common cellular basis for non-congenital cataract development in humans and animals, *Journal of Cell Biology*, 130:169-81. [\[DOI\]](#) PMID:7790371
- Lija Y, Biju PG, Reeni A, Cibin TR, Sahasranamam V, Abraham A (2006). Modulation of selenite cataract by the flavonoid fraction of *Emilia sonchifolia* in experimental animal model. *Phytotherapy Research*, 20:1091-1095. [\[DOI\]](#) PMID:17009203
- Pastene E, Avello M, letelier ME, sanzana E, vega M, Gonzalez M, (2007). Preliminary studies on antioxidant and anticataract activity of *cheilanthes glauca* (cav.) Mett. Through various in vitro models. *Electronic Journal of Food and Plants Chemistry*, 2: 1-8.
- Patel DK, Kumar R, Prasad SK, Hemalata S (2011). *Pedaliium murex* Linn (pedaliaceae). fruits; a comparative antioxidant activity of it's different fractions. *Asian Pacific Journal of Tropical Biomedicine*, 1: 395-400. [http://dx.doi.org/10.1016/S2221-1691\(11\)60087-7](http://dx.doi.org/10.1016/S2221-1691(11)60087-7)
- Patel DK, kumar R, Prasad SK, Sairam K, Hemalatha S, (2011). Antidiabetic and in vitro antioxidant potential of *Hybanthus enneaspermum* (Linn.). F. Muell in streptozotocin-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine*, 1: 316-322 [\[DOI\]](#)
- Rathi S S, Grover J K, Vikrant V, Biswas N R, (2002). Prevention of experimental diabetic cataract by Indian ayurvedic plant extracts. *Phytotherapy Research*, 16:774-777. [\[DOI\]](#) PMID:12458487
- Rooban BN, Sasikala V, Sahasranamam V, Abraham A (2010). *Vitex negundo* modulates selenite induced opacification and cataractogenesis in rat pups. *Biological Trace Element Research*, 138:282-292. [\[DOI\]](#) PMID:20174976
- Rooban BN, Sasikala V, Sahasranaman V, Abraham A (2011). Amelioration of selenite toxicity and cataractogenesis in cultured rat lens by *vitex negundo*. *Graefes Archive of Clinical and Experimental Ophthalmology*, 249:685-692. [\[DOI\]](#) PMID:21240520
- Sasikala V, Rooban B N, Priya S C, Sahasranamam V, Abraham A (2010). *Moringa oleifera* prevents selenite induced cataractogenesis in rat pups. *Journal of Ocular Pharmacology and Therapeutics*, 26: 441-447. [\[DOI\]](#) PMID:20879807
- Shearer TR, Ma H, Fukiage C, Azuma M (1997). Selenite nuclear cataract: Review of the model. *Molecular Vision*, 3: 8. PMID:9238097
- Shin KH, Lim SS, Kim DK (1998). Effect of byakangelicin an aldose reductase inhibitor, on galactosamine cataract, the polyol contents and Na+ K+ ATPase activity in sciatic nerve of streptozotocin induced diabetic rats. *Phytomedicine* 5: 121-127. [\[DOI\]](#)
- Suryanarayana P, saraswat M, Mrudula T, Krishna TP, Krishnaswamy K, Reddy GB, (2005). Curcumin and turmeric delay streptozotocin induced diabetic cataract in rats. *Investigative ophthalmology and visual science*, 46:2092-2099. [\[DOI\]](#)
- Thiagarajan G., Venu T., Balasubramanian D. (2003). Approche to relieve the burden of cataract blindness through natural antioxidants: use of *ashwagandha* (withania somnifera). *Current science*, 85(7): 1065-1071.
- Umamaheshwari M, Ashokkumar K, Lalitha V, Sivashanmugam AT, subhadradevi V (2011). Anticataract and antioxidant activities of citrus aurantium L. Peel extract against naphthalene induced cataractogenesis in rats. *Journal of Pharmacy Research*, 4:680-682.
- Vats V S, Yadav S P, Biswas N R, Grover J K, (2004). Anticataract activity of *pterocarpus marsupium* bark and *trigonella foenum graceum* seeds extracts in alloxan diabetic rats. *Journal of Ethnopharmacology*, 93:289-294. [\[DOI\]](#) PMID:15234767
- Vinson, J.A., Zhang, J. (2005). Black and green teas equally inhibit diabetic cataracts in a streptozotocin-induced rat model of diabetes, *Journal of Agricultural and Food Chemistry*, 53:3710-3713. [\[DOI\]](#) PMID:15853424
- Vishvanath GLS, Vaidya S.K., Ramesh C., Krishnadas N., (2010). Antioxidant and antimutagenic activity of bark extract of *Terminalia arjuna*, *Asia Pacific Journal of Tropical Medicine*, 3:695-670.