



HAL
open science

Categorical models of Linear Logic with fixed points of formulas

Thomas Ehrhard, Farzad Jafarrahmani

► **To cite this version:**

Thomas Ehrhard, Farzad Jafarrahmani. Categorical models of Linear Logic with fixed points of formulas. 2021. hal-03013356v2

HAL Id: hal-03013356

<https://hal.archives-ouvertes.fr/hal-03013356v2>

Preprint submitted on 5 Feb 2021 (v2), last revised 18 May 2021 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Categorical models of Linear Logic with fixed points of formulas

Thomas Ehrhard and Farzad Jafarrahmani
IRIF, CNRS and Université de Paris

Abstract—We develop a categorical semantics of μLL , a version of propositional Linear Logic with least and greatest fixed points extending David Baelde’s propositional μMALL with exponentials. Our general categorical setting is based on Seely categories and on strong functors acting on them. We exhibit two simple instances of this setting. In the first one, which is based on the category of sets and relations, least and greatest fixed points are interpreted in the same way. In the second one, based on a category of sets equipped with a notion of totality (non-uniform totality spaces) and relations preserving it, least and greatest fixed points have distinct interpretations. This latter model shows that μLL enjoys a denotational form of normalization of proofs.

I. INTRODUCTION

Propositional Linear Logic is a well-established logical system introduced by Girard in [1]. It provides a fine-grain analysis of proofs in intuitionistic and classical logic, and more specifically of their cut-elimination. LL features a logical account of the structural rules (weakening, contraction) which are handled implicitly in intuitionistic and classical logic. For this reason, LL has many useful outcomes in the Curry-Howard based approach to the theory of programming: logical understanding of evaluation strategies, new syntax of proofs/programs (proof-nets), connections with other branches of mathematics (linear algebra, functional analysis, differential calculus), new operational semantics (geometry of interaction) etc.

However propositional LL is not a reasonable programming language, by lack of data-types and iteration or recursion principles. This is usually remedied by extending propositional LL to the 2nd order, thus defining a logical system in which Girard’s System F [2] can be embedded. Another option to turn propositional LL into a programming language – closer to usual programming – is to extend it with least and greatest fixed points of formulas. Such an extension was early suggested by Girard in an unpublished note [3], though the first comprehensive proof-theoretic investigation of such an extension of LL is recent: in [4] Baelde considers an extension μMALL of Multiplicative Additive LL sequent calculus with least and greatest fixed points. His motivations arose from a proof-search and system verification perspective and therefore his μMALL logical system is a predicate calculus. Our purpose is to develop a more Curry-Howard oriented point of view on LL with fixed points and therefore we stick to the proposition calculus setting of [2]. But, unlike [4] we

include the exponentials in our system from the beginning¹, so we call it μLL rather than propositional μMALL and we consider it as an alternative to the “system F” approach to representing programs in LL. The ν -introduction rule of μLL (Park’s rule, that is rule ($\nu\text{-rec}$) of Section II-F1) leads to subtle cut-elimination rewrite rules for which Baelde could prove cut-elimination in μMALL , showing for instance that a proof of the type of integers $\mu\zeta(1 \oplus \zeta)$ necessarily reduces to an integer (in contrast with LL, μMALL enjoys only a restricted form of sub-formula property). There are alternative proof-systems for the same logic, involving infinite or cyclic proofs, see [5], whose connections with the aforementioned finitary proof-system are not completely clear yet.

Since the proof-theory (and hence the “operational semantics”) of μLL is still under development, it is important to investigate its categorical semantics, whose definition does not rely on the precise choice of inference and rewrite rules we equip μLL with, see the *Outcome* § below. We develop here a categorical semantics of μLL extending the standard notion of Seely category² of classical LL, see [6]. Such a model of μLL consists of a Seely category \mathcal{L} and of a class of functors $\mathcal{L}^n \rightarrow \mathcal{L}$ for all possible arities n which will be used for interpreting μLL formulas with free variables. These functors have to be equipped with a strength to deal properly with contexts in the rule ($\nu\text{-rec}$), see Section II-F2 for a discussion on these contexts in particular.

Then we develop a simple instance of this setting which consists in taking for \mathcal{L} the category of sets and relations, a well-known Seely model of LL. The *variable sets* are the strong functors we consider on this category. They are the pairs $\mathbb{F} = (\overline{\mathbb{F}}, \widehat{\mathbb{F}})$ where $\widehat{\mathbb{F}}$ is the strength and $\overline{\mathbb{F}} : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ is a functor which is Scott-continuous in the sense that it commutes with directed unions of morphisms. This property implies that $\overline{\mathbb{F}}$ maps injections to injections and is cocontinuous on the category of sets and injections. There is no special requirement about the strength $\widehat{\mathbb{F}}$ beyond naturality, monoidality and compatibility with the comultiplication of the comonad $!_-$. Variable sets form a Seely model of μLL where linear negation is the identity on objects. The formulas

¹Exponentials are not considered in μMALL because some form of exponential can be encoded using inductive/coinductive types, however these exponentials are not fully satisfactory from our point of view because their denotational interpretation does not satisfy all required isomorphisms; specifically, the *Seely isos* are lacking.

²Sometimes called new-Seely category: it is a cartesian symmetric monoidal closed category with a $*$ -autonomous structure and a comonad $!_-$ with a strong symmetric monoidal structure from the cartesian product to the tensor product.

$\mu\zeta F$ and $\nu\zeta F$ are interpreted as the same variable set, exactly as \otimes and \wp are interpreted in the same way (and similarly for additives and exponentials). This denotational “degeneracy” at the level of types is a well known feature of **Rel** which does not mean at all that the model is trivial. For instance normal multiplicative exponential LL proofs which have distinct relational interpretations have distinct associated proof-nets [7], [8].

Last we enrich this model **Rel** by considering sets equipped with an additional structure of *totality*: a *non-uniform totality space* (NUTS) is a pair $X = (|X|, \mathcal{T}(X))$ where $|X|$ is a set and $\mathcal{T}(X)$ is a set of subsets of $|X|$ which intuitively represent the total, that is, terminating computations of type X . This set $\mathcal{T}(X)$ is required to coincide with its bidual for a duality expressed in terms of non-empty intersections. This kind of definition by duality is ubiquitous in LL since [1] and has been categorically formalized as *double glueing* in [9]. We don’t use this categorical formalization here however as it would not simplify the presentation. One nice feature of this specific duality is that the bidual of a set of subsets of $|X|$ is simply its upwards-closure (wrt. inclusion)³, see Lemma 17. Given two NUTS X and Y there is a natural notion of *total relation* $t \subseteq |X| \times |Y|$ giving rise to a category **Nuts** which is easily seen to be a Seely model of LL. To turn it into a categorical model of μ LL, we need a notion of strong functors $\mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$. Rather than considering them directly as functors, we define *variable non-uniform totality spaces* (VNUTS) as pairs $\mathbb{X} = (|\mathbb{X}|, \mathcal{T}(\mathbb{X}))$ where $|\mathbb{X}| : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ is a variable set and, for each tuple $\vec{X} = (X_1, \dots, X_n)$ of VNUTS’s, $\mathcal{T}(\mathbb{X})(\vec{X})$ is a totality structure on the set $|\mathbb{X}|(|\vec{X}|)$. It is also required that the action of the functor $|\mathbb{X}|$ on **Nuts** morphisms and the strength $\widehat{\mathbb{X}}$ respect this totality structures. Then it is easy to derive from such a VNUTS \mathbb{X} a strong functor $\mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$ and we prove that, equipped with these strong functors, **Nuts** is a model of μ LL. Most proofs are given in an Appendix.

a) *Outcome*: One major benefit of this construction is that it gives a value to all proofs of μ LL, invariant by cut-elimination. Moreover, the fact that this value is total shows in a *syntax independent way* that when π is for instance a μ LL proof of $1 \oplus 1$ (the type of booleans), the value associated with π is non-empty, that is, π has a defined boolean value true or false⁴. We could also obtain this by a normalization theorem: π reduces to one of the two normal proofs of $1 \oplus 1$ (and if we prove for instance a Church-Rosser theorem we will know that this proof is unique). Such proofs would depend of course on the actual presentation of the syntax whereas our denotational argument does not.

b) *Related work*: There is a vast literature on extending logic with fixed point that we cannot reasonably

³This is a major simplification wrt. notions of totality on coherence spaces [10] or Loader’s totality spaces [11] where biduality is much harder to deal with.

⁴Or both because our **Nuts** model accepts non-determinism. By adding a *non-uniform* coherence relation as defined in [12], [13] to the model one can show that this value is actually a uniquely defined boolean. See also Section IV-C1.

summarize, see the discussions in [14], [5]. Cut-elimination of such systems has been extensively investigated, see for instance [15], [16], [17], [18]. Closer to ours is the work of Santocanale [19] and its categorical interpretation in μ -bicomplete categories [20] which, unlike most contributions in this field, considers also categorical interpretations of proofs. Santocanale *et al.* consider circular proofs whereas we use Park’s rule. A deeper difference lies in the logic itself: from an LL point of view the logic considered by Santocanale *et al.* is *purely additive linear logic with least and greatest fixed points* μ ALL which seems too weak in our Curry-Howard perspective. And indeed μ -bicomplete categories do not provide the monoidal and exponential structures required for interpreting μ LL.

In [21], that we became aware of only recently (and seems related to the earlier report [22]), Loader extends the simply typed λ -calculus with inductive types and develops its denotational semantics. His models are cartesian closed categories \mathcal{C} equipped with a class of strong functors and seem very close to ours (Section II-F): one might think that any of our models yields a Loader model as its Kleisli category. This is not the case because in a Loader model the category \mathcal{C} is cocartesian⁵ whereas the Kleisli category of a Seely category is not cocartesian in general: this would require to have an iso between $!(X \oplus Y)$ and $!X \oplus !Y$ which is usually absent. Loader studies two concrete instances of his models: one is based on recursion theory (partial equivalence relations) and the other on a notion of domains with totality described as a model of LL and might give rise to one of our Seely models, this point requires further studies. Our NUTS are quite different from Loader totality domains which feature a notion of “consistency” enforcing some kind of determinism and, combined with totality, allow the Kleisli category to be cocartesian as well. Our model is based on **Rel** and therefore is compatible with non-determinism [23] and PCF recursion. This is important for us because we would like to consider rules beyond Park’s rule for inductive and coinductive types, based on PCF fixed points with further guardedness conditions⁶ in the spirit of [24], [25], [26] or even on infinite terms in the spirit of [5].

We mention also the work of Clairambault [27], [28] who investigates the game with totality semantics of an extension of intuitionistic logic with least and greatest fixed points (independently of [21], [22]). A Kleisli-like connection with his work should be sought too.

c) *Notations*: We use the following conventions: \vec{a} stands for a list (a_1, \dots, a_n) . A unary operation f is extended to lists of arguments in the obvious way: $f(\vec{a}) = (f(a_1), \dots, f(a_n))$. When we write natural transformations, we very often omit the objects where they are taken keeping them implicit for the sake of readability, when they can easily be retrieved from the context. If \mathcal{A} is a category then $\text{Obj}(\mathcal{A})$ is its class of objects and if $A, B \in \text{Obj}(\mathcal{A})$ then $\mathcal{A}(A, B)$ is

⁵To account for the disjunction of his logical system which is crucial for defining interesting data-types such as the integers.

⁶For guaranteeing termination.

the set of morphisms from A to B in \mathcal{A} (all the categories we consider are locally small). If $\mathcal{F} : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ is a functor and $A \in \text{Obj}(\mathcal{A})$ then $\mathcal{F}_A : \mathcal{B} \rightarrow \mathcal{C}$ is the functor defined by $\mathcal{F}_A(B) = \mathcal{F}(A, B)$ and $\mathcal{F}_A(f) = \mathcal{F}(A, f)$ (we often write A instead of Id_A).

II. CATEGORICAL MODELS OF LL

A. Seely categories.

We recall the basic notion of categorical model of LL. Our main reference is the notion of a *Seely category* as presented in [6]. We refer to that survey for all the technical material that we do not recall here.

A Seely category is a symmetric monoidal closed category (SMCC) $(\mathcal{L}, \otimes, 1, \lambda, \rho, \alpha, \gamma)$ where $\lambda_X \in \mathcal{L}(1 \otimes X, X)$, $\rho_X \in \mathcal{L}(X \otimes 1, X)$, $\alpha_{X,Y,Z} \in \mathcal{L}((X \otimes Y) \otimes Z, X \otimes (Y \otimes Z))$ and $\gamma_{X,Y} \in \mathcal{L}(X \otimes Y, Y \otimes X)$ are natural isomorphisms satisfying coherence diagrams that we do not record here. We use $X \multimap Y$ for the object of linear morphisms from X to Y , $\text{ev} \in \mathcal{L}((X \multimap Y) \otimes X, Y)$ for the evaluation morphism and cur for the linear curryfication map $\mathcal{L}(Z \otimes X, Y) \rightarrow \mathcal{L}(Z, X \multimap Y)$. We assume \mathcal{L} to be $*$ -autonomous with dualizing object \perp (this object is part of the structure of a Seely category). We use X^\perp for the object $X \multimap \perp$ of \mathcal{L} (the dual, or linear negation, of X). It is also assumed that \mathcal{L} is cartesian with final object \top , product $X_1 \& X_2$ with projections π_1, π_2 . By $*$ -autonomy \mathcal{L} is cocartesian with initial object 0 , coproduct \oplus and injections $\bar{\pi}_i$.

We also assume to be given a comonad $!_ : \mathcal{L} \rightarrow \mathcal{L}$ with counit $\text{der}_X \in \mathcal{L}(!_X, X)$ (*dereliction*) and comultiplication $\text{dig}_X \in \mathcal{L}(!_X, !!X)$ (*digging*) together with a strong symmetric monoidal structure (Seely natural isos $m^0 : 1 \rightarrow !\top$ and m^2 with $m^2_{X_1, X_2} : !X_1 \otimes !X_2 \rightarrow !(X_1 \& X_2)$ for the functor $!_$, from the symmetric monoidal category $(\mathcal{L}, \&)$ to the symmetric monoidal category (\mathcal{L}, \otimes) satisfying an additional coherence condition wrt. dig). This strong monoidal structure allows to define a lax monoidal structure (μ^0, μ^2) of $!_$ from (\mathcal{L}, \otimes) to itself. More precisely $\mu^0 \in \mathcal{L}(1, !1)$ and $\mu^2_{X_1, X_2} \in \mathcal{L}(!_X \otimes !_Y, !(X_1 \& X_2))$ are defined using m^0 and m^2 (and are not isos in most cases). Also, for each object $X \in \text{Obj}(\mathcal{L})$, there is a canonical structure of commutative \otimes -comonoid on $!X$ given by $w_X \in \mathcal{L}(!_X, 1)$ and $\text{contr}_X \in \mathcal{L}(!_X, !_X \otimes !_X)$. The definition of these morphisms involves all the structure of $!_$ explained above, and in particular the Seely isos. We use $?_$ for the ‘‘De Morgan dual’’ of $!_$: $?X = !(X^\perp)^\perp$ and similarly for morphisms.

B. Oplax monoidal comonads

Let \mathcal{M} be a symmetric monoidal category (with the same notations as above for the tensor product) and $(T, \varepsilon, \mu) : \mathcal{M} \rightarrow \mathcal{M}$ be a comonad (ε is the unit and μ the multiplication). An *oplax monoidal structure* on T consists of a morphism $\theta^0 \in \mathcal{M}(T1, 1)$ and a natural transformation $\theta^2_{X_1, X_2} \in \mathcal{M}(T(X_1 \otimes X_2), T(X_1) \otimes T(X_2))$ subject to standard symmetric monoidality and compatibility with ε and μ , this latter reading $(\varepsilon_{X_1} \otimes \varepsilon_{X_2}) \theta_{X_1, X_2} = \varepsilon_{X_1 \otimes X_2}$ and:

$$\begin{array}{ccc} T(X_1 \otimes X_2) & \xrightarrow{\theta_{X_1, X_2}^2} & TX_1 \otimes TX_2 & \xrightarrow{\mu_{X_1} \otimes \mu_{X_2}} & T^2 X_1 \otimes T^2 X_2 \\ & & \downarrow \mu_{X_1 \otimes X_2} & & \uparrow \theta_{TX_1, TX_2} \\ T^2(X_1 \otimes X_2) & \xrightarrow{T(\theta_{X_1, X_2}^2)} & T(TX_1 \otimes TX_2) & & \end{array}$$

Then the Kleisli category \mathcal{M}_T has a canonical symmetric monoidal structure, with unit 1 and tensor product $X_1 \otimes X_2$ defined as in \mathcal{M} for objects. Given $f_i \in \mathcal{M}_T(X_i, Y_i)$, $f_1 \otimes_T f_2 \in \mathcal{M}_T(X_1 \otimes X_2, Y_1 \otimes Y_2)$ is defined as

$$T(X_1 \otimes X_2) \xrightarrow{\theta_{X_1, X_2}^2} TX_1 \otimes TX_2 \xrightarrow{f_1 \otimes f_2} Y_1 \otimes Y_2 .$$

Let $F_T : \mathcal{M} \rightarrow \mathcal{M}_T$ be the canonical functor which acts as the identity on objects and maps $f \in \mathcal{M}(X, Y)$ to $f \varepsilon_X \in \mathcal{M}_T(X, Y)$.

C. Eilenberg-Moore category and free comodules

Let \mathcal{L} be a Seely category. Since $!_$ is a comonad we can define the category $\mathcal{L}^!$ of $!_$ -coalgebras (Eilenberg-Moore category of $!_$). An object of this category is a pair $P = (\underline{P}, h_P)$ where $\underline{P} \in \text{Obj}(\mathcal{L})$ and $h_P \in \mathcal{L}(\underline{P}, !_\underline{P})$ is such that $\text{der}_{\underline{P}} h_P = \text{Id}$ and $\text{dig}_{\underline{P}} h_P = !h_P h_P$. Then $f \in \mathcal{L}^!(P, Q)$ if $f \in \mathcal{L}(\underline{P}, \underline{Q})$ and $h_Q f = !f h_P$. The functor $!_$ can be seen as a functor from \mathcal{L} to $\mathcal{L}^!$ mapping X to $(!X, \text{dig}_X)$ and $f \in \mathcal{L}(X, Y)$ to $!f$. It is right adjoint to the forgetful functor $\mathcal{L}^! \rightarrow \mathcal{L}$. Given $f \in \mathcal{L}(\underline{P}, X)$, we use $f^! \in \mathcal{L}^!(P, !X)$ for the morphism associated with f by this adjunction, one has $f^! = !f h_P$. If $g \in \mathcal{L}^!(Q, P)$, we have $f^! g = (fg)^!$. Then $\mathcal{L}^!$ is cartesian with final object $(1, h_1 = \mu^0)$ still denoted as 1 and product $P_1 \otimes P_2 = (\underline{P}_1 \otimes \underline{P}_2, h_{P_1 \otimes P_2})$ with

$$h_{P_1 \otimes P_2} : \underline{P}_1 \otimes \underline{P}_2 \xrightarrow{h_{P_1} \otimes h_{P_2}} !_\underline{P}_1 \otimes !_\underline{P}_2 \xrightarrow{\mu_{!P_1, !P_2}^2} !(P_1 \otimes P_2) .$$

This category is also cocartesian with initial object $(0, h_0)$ still denoted as 0 and coproduct $P_1 \oplus P_2 = (\underline{P}_1 \oplus \underline{P}_2, h_{P_1 \oplus P_2})$ with $h_{P_1 \oplus P_2}$ defined as follows. For $i = 1, 2$ one defines $h^i : \underline{P}_i \rightarrow !(P_1 \oplus P_2)$ as $\underline{P}_i \xrightarrow{h_{P_i}} !_\underline{P}_i \xrightarrow{! \bar{\pi}_i} !(P_1 \oplus P_2)$ and then $h_{P_1 \oplus P_2}$ is the unique morphism $\underline{P}_1 \oplus \underline{P}_2 \rightarrow !(P_1 \oplus P_2)$ such that $h_{P_1 \oplus P_2} \bar{\pi}_i = h^i$ for $i = 1, 2$. More details can be found in [6]. We use $\text{contr}_P \in \mathcal{L}^!(P, P \otimes P)$ (*contraction*) for the diagonal and $w_P \in \mathcal{L}^!(P, 1)$ (*weakening*) for the unique morphism to the final object.

1) *The LL model of free comodules on a given coalgebra:* Given an object P of $\mathcal{L}^!$, we can define a functor $\text{fc}_P : \mathcal{L} \rightarrow \mathcal{L}$ which maps an object X to $\underline{P} \otimes X$ and a morphism f to $\underline{P} \otimes f$. This functor is clearly an oplax monoidal comonad (with structure maps defined using w_P , contr_P and the monoidal structure of \mathcal{L})⁸. A coalgebra for this comonad is a *P-comodule*. By Section II-B the Kleisli category $\mathcal{L}[P] = \mathcal{L}_{\text{fc}_P}$ of this comonad (that is, the category of free P -comodules) has a canonical structure of symmetric monoidal category (SMC).

⁷In this paper we could restrict to the case where P is a tensor of ‘‘free coalgebras’’ $(!X_i, \text{dig}_{X_i})$ but it is more natural to deal with the general case, which will be quite useful in further work, see Section V.

⁸The definition of this comonad uses only the comonoid structure of \underline{P} . The $!_$ -structure will be used later.

We set $F_P = F_{fc_P} : \mathcal{L} \rightarrow \mathcal{L}[P]$. Girard showed in [29] that $\mathcal{L}[P]$ is a Seely model of LL with operations on objects defined in the same way as in \mathcal{L} , and using the coalgebra structure of P for the operations on morphisms. Intuitively, P should be considered as a given context and $\mathcal{L}[P]$ as a model in this context. This idea appears at various places in the literature, see for instance [30], [31]. Let us summarize this construction. If $f_i \in \mathcal{L}[P](X_i, Y_i)$ for $i = 1, 2$ then $f_1 \otimes_P f_2 = f_1 \otimes_{fc_P} f_2 \in \mathcal{L}[P](X_1 \otimes X_2, Y_1 \otimes Y_2)$ is given by

$$\begin{array}{ccc} \underline{P} \otimes X_1 \otimes X_2 & \xrightarrow{\text{contr}_P \otimes \text{Id}} & \underline{P} \otimes \underline{P} \otimes X_1 \otimes X_2 \\ & & \parallel \\ Y_1 \otimes Y_2 & \xleftarrow{f_1 \otimes f_2} & \underline{P} \otimes X_1 \otimes \underline{P} \otimes X_2 \end{array}$$

The object of linear morphisms from X to Y in $\mathcal{L}[P]$ is $X \multimap Y$, and the evaluation morphism $\text{ev}_P \in \mathcal{L}[P]((X \multimap Y) \otimes X, Y)$ is simply $F_P(\text{ev})$. Then it is easy to check that if $f \in \mathcal{L}[P](Z \otimes X, Y)$, that is $f \in \mathcal{L}(\underline{P} \otimes Z \otimes X, Y)$, the morphism $\text{cur}(f) \in \mathcal{L}[P](Z, X \multimap Y)$ satisfies the required monoidal closeness equations. With these definitions, the category $\mathcal{L}[P]$ is $*$ -autonomous, with \perp as dualizing object. Specifically, given $f \in \mathcal{L}[P](X, Y)$, then $f^{\perp[P]}$ is the following composition of morphisms:

$$\underline{P} \otimes Y^{\perp} \xrightarrow{\underline{P} \otimes f^{\perp}} \underline{P} \otimes (\underline{P} \multimap X^{\perp}) \xrightarrow{\text{ev}} X^{\perp}$$

using implicitly the iso between $(Z \otimes X)^{\perp}$ and $Z \multimap X^{\perp}$, and the $*$ -autonomy of \mathcal{L} allows to prove that indeed $f^{\perp[P]^{\perp[P]}} = f$. The category $\mathcal{L}[P]$ is easily seen to be cartesian with \top as final object, $X_1 \& X_2$ as cartesian product (and projections defined in the obvious way, applying F_P to the projections of \mathcal{L}). Last we define a functor $!_P : \mathcal{L}[P] \rightarrow \mathcal{L}[P]$ by $!_P X = !X$ and, given $f \in \mathcal{L}[P](X, Y)$, we define $!_P f \in \mathcal{L}[P](!X, !Y)$ as $\underline{P} \otimes !X \xrightarrow{h_P \otimes !X} \underline{P} \otimes !X \xrightarrow{\mu^2} !(\underline{P} \otimes X) \xrightarrow{!f} !Y$ and this functor has a comonad structure $(\text{der}[P], \text{dig}[P])$ defined by $\text{der}[P] = F_P(\text{der})$ and $\text{dig}[P] = F_P(\text{dig})^9$.

Remark 1. Any $p \in \mathcal{L}^!(P, Q)$ induces a functor $\mathcal{L}[p] : \mathcal{L}[Q] \rightarrow \mathcal{L}[P]$ which acts as the identity on objects and maps $f \in \mathcal{L}[Q](X, Y)$ to $\mathcal{L}[p](f) = f(p \otimes X) \in \mathcal{L}[P](X, Y)$. This functor is strict monoidal symmetric and preserves all the constructions of LL, for instance $\mathcal{L}[p](\text{dig}[Q]) = \text{dig}[P]$ (simply because $\mathcal{L}[p] \circ F_Q = F_P$) and also $\mathcal{L}[p](!_Q f) = !_P(\mathcal{L}[p](f))$. We can actually consider $\mathcal{L}[_]$ as a functor from \mathcal{L}^{op} to the category of Seely categories and functors which preserve their structure on the nose. This functor could probably more suitably be considered as a fibration in the line of [32], Section 7.

D. Strong functors on \mathcal{L}

Given $n \in \mathbb{N}$, an n -ary strong functor on \mathcal{L} is a pair $\mathbb{F} = (\overline{\mathbb{F}}, \widehat{\mathbb{F}})$ where $\overline{\mathbb{F}} : \mathcal{L}^n \rightarrow \mathcal{L}$ is a functor and $\widehat{\mathbb{F}}_{X, \vec{Y}} \in \mathcal{L}(!X \otimes \overline{\mathbb{F}}(\vec{Y}), \overline{\mathbb{F}}(!X \otimes \vec{Y}))$ is a natural transformation, called the *strength* of \mathbb{F} . We use the notation

⁹The definition of $!_P f$ requires P to be a $!$ -coalgebra and not simply a commutative \otimes -comonoid. Of course if $!$ is the free exponential as in [29] the latter condition implies the former.

$Z \otimes (Y_1, \dots, Y_n) = (Z \otimes Y_1, \dots, Z \otimes Y_n)$. It is assumed moreover that the diagrams of Figure 1 commute, expressing the monoidality of this strength as well as its compatibility with the comultiplication of $!$. The main purpose of this definition is that for any object P of $\mathcal{L}^!$ one can lift \mathbb{F} to a functor $\mathbb{F}[P] : \mathcal{L}[P]^n \rightarrow \mathcal{L}[P]$ as follows. First one sets $\mathbb{F}[P](\vec{X}) = \overline{\mathbb{F}}(\vec{X})$. Then, given $\vec{f} \in \mathcal{L}[P]^n(\vec{X}, \vec{Y})$ we define $\mathbb{F}[P](\vec{f}) \in \mathcal{L}[P](\mathbb{F}(\vec{X}), \mathbb{F}(\vec{Y}))$ as

$$\begin{array}{ccc} \underline{P} \otimes \mathbb{F}(\vec{X}) & \xrightarrow{h_P \otimes \text{Id}} & !\underline{P} \otimes \mathbb{F}(\vec{X}) \xrightarrow{\widehat{\mathbb{F}}} \mathbb{F}(!\underline{P} \otimes \vec{X}) \xrightarrow{\overline{\mathbb{F}}(\text{der}_P \otimes \vec{X})} \mathbb{F}(\underline{P} \otimes \vec{X}) \\ & & \mathbb{F}(\vec{f}) \downarrow \\ & & \mathbb{F}(\vec{Y}) \end{array}$$

The fact that we have defined a functor results from the three diagrams of Figure 1 and from the definition of w_P and contr_P based on the Seely isomorphisms.

Remark 2. Since the seminal work of Moggi [33] strong functors play a central role in semantics for representing effects. Hasegawa [34] showed that functors on a model of LL which have a strength wrt. \otimes could be lifted to the Klesli category of $!$. Our strong functors differ from Hasegawa's in two respects: first the strength is required only for $!$ -ed objects, and second it has to satisfy a compatibility condition wrt. dig .

1) *Operations on strong functors:* There is an obvious unary identity strong functor \mathcal{I} and for each object Y of \mathcal{L} there is an n -ary Y -valued constant strong functor \mathcal{K}^Y ; in the first case the strength natural transformation is the identity morphism and in the second case, it is defined using $w_{!X}$. Let \mathbb{F} be an n -ary strong functor and $\mathbb{G}_1, \dots, \mathbb{G}_n$ be k -ary strong functors. Then one defines a k -ary strong functor $\mathbb{H} = \mathbb{F} \circ (\mathbb{G}_1, \dots, \mathbb{G}_n)$: the functorial component $\overline{\mathbb{H}}$ is defined in the obvious compositional way. The strength is

$$!X \otimes \overline{\mathbb{H}}(\vec{Y}) \xrightarrow{\widehat{\mathbb{H}}} \overline{\mathbb{F}}(!X \otimes \overline{\mathbb{G}}_i(\vec{Y}))_{i=1}^n \xrightarrow{\overline{\mathbb{F}}(\widehat{\mathbb{G}}_i^k)} \overline{\mathbb{F}}(\overline{\mathbb{G}}_i(!X \otimes \vec{Y}))_{i=1}^n$$

and is easily seen to satisfy the commutations of Figure 1. Given an n -ary strong functor, we can define its *De Morgan dual* \mathbb{F}^{\perp} which is also an n -ary strong functor. On objects, we set $\overline{\mathbb{F}^{\perp}}(\vec{Y}) = \overline{\mathbb{F}}(\vec{Y}^{\perp})^{\perp}$ and similarly for morphisms. The strength of \mathbb{F}^{\perp} is defined as the Curry transpose of the following morphism (remember that $!X \multimap \vec{Y}^{\perp} = (!X \otimes \vec{Y})^{\perp}$ up to canonical iso):

$$\begin{array}{ccc} !X \otimes \overline{\mathbb{F}}(\vec{Y}^{\perp})^{\perp} \otimes \overline{\mathbb{F}}(!X \multimap \vec{Y}^{\perp}) & \cong & !X \otimes \overline{\mathbb{F}}(!X \multimap \vec{Y}^{\perp}) \otimes \overline{\mathbb{F}}(\vec{Y}^{\perp})^{\perp} \\ & & \downarrow \widehat{\mathbb{F} \otimes \text{Id}} \\ \overline{\mathbb{F}}(\vec{Y}^{\perp}) \otimes \overline{\mathbb{F}}(\vec{Y}^{\perp})^{\perp} & \xleftarrow{\overline{\mathbb{F}}(\text{ev}) \otimes \text{Id}} & \overline{\mathbb{F}}(!X \otimes (!X \multimap \vec{Y}^{\perp})) \otimes \overline{\mathbb{F}}(\vec{Y}^{\perp})^{\perp} \\ & & \downarrow \text{ev } \gamma \\ & & \perp \end{array}$$

Then it is possible to prove, using the $*$ -autonomy of \mathcal{L} , that $\mathbb{F}^{\perp \perp}$ and \mathbb{F} are canonically isomorphic (as strong functors)¹⁰.

¹⁰In the concrete settings considered in this paper, these canonical isos are actuality identity maps.

$$\begin{array}{ccc}
(!X_1 \otimes !X_2) \otimes \overline{\mathbb{F}}(\overrightarrow{Y}) & \xrightarrow{m^2 \otimes \overline{\mathbb{F}}(\overrightarrow{Y})} & !(X_1 \& X_2) \otimes \overline{\mathbb{F}}(\overrightarrow{Y}) \\
!X_1 \otimes \overline{\mathbb{F}}_{X_2, \overrightarrow{Y}} \downarrow & & \downarrow \overline{\mathbb{F}}_{X_1 \& X_2, \overrightarrow{Y}} \\
!X_1 \otimes \overline{\mathbb{F}}(!X_2 \otimes \overrightarrow{Y}) & & \\
\overline{\mathbb{F}}_{X_1, !X_2 \otimes \overrightarrow{Y}} \downarrow & & \\
\overline{\mathbb{F}}(!X_1 \otimes !X_2 \otimes \overline{\mathbb{F}}(\overrightarrow{Y})) & \xrightarrow{\overline{\mathbb{F}}(m^2 \otimes \overrightarrow{Y})} & \overline{\mathbb{F}}(!(X_1 \& X_2) \otimes \overrightarrow{Y})
\end{array}
\quad
\begin{array}{ccc}
1 \otimes \overline{\mathbb{F}}(\overrightarrow{Y}) & \xrightarrow{m^0 \otimes \overline{\mathbb{F}}(\overrightarrow{Y})} & !\top \otimes \overline{\mathbb{F}}(\overrightarrow{Y}) & \quad & !X \otimes \overline{\mathbb{F}}(\overrightarrow{Y}) & \xrightarrow{\text{dig}_X \otimes \overline{\mathbb{F}}(\overrightarrow{Y})} & !!X \otimes \overline{\mathbb{F}}(\overrightarrow{Y}) \\
\Downarrow & & \downarrow \overline{\mathbb{F}}_{\top, \overrightarrow{Y}} & & \overline{\mathbb{F}}_{X, \overrightarrow{Y}} \downarrow & & \downarrow \overline{\mathbb{F}}_{!X, \overrightarrow{Y}} \\
\overline{\mathbb{F}}(1 \otimes \overrightarrow{Y}) & \xrightarrow{\overline{\mathbb{F}}(m^0 \otimes \overrightarrow{Y})} & \overline{\mathbb{F}}(!\top \otimes \overrightarrow{Y}) & & \overline{\mathbb{F}}(!X \otimes \overrightarrow{Y}) & \xrightarrow{\overline{\mathbb{F}}(\text{dig}_X \otimes \overrightarrow{Y})} & \overline{\mathbb{F}} (!!X \otimes \overrightarrow{Y})
\end{array}$$

Fig. 1. Monoidality and dig diagrams for strong functors

As a direct consequence of the definition of \mathbb{F}^\perp and of the canonical iso between $\mathbb{F}^{\perp\perp}$ and \mathbb{F} we get:

Lemma 1. $(\mathbb{F} \circ (\mathbb{G}_1, \dots, \mathbb{G}_n))^\perp = \mathbb{F}^\perp \circ (\mathbb{G}_1^\perp, \dots, \mathbb{G}_n^\perp)$ up to canonical iso.

The bifunctor \otimes can be turned into a strong functor: one defines the strength as¹¹

$$!X \otimes Y_1 \otimes Y_2 \xrightarrow{\text{contr}_{!X} \otimes \text{Id}} !X \otimes !X \otimes Y_1 \otimes Y_2 \cong !X \otimes Y_1 \otimes !X \otimes Y_2$$

By De Morgan duality, this endows \mathfrak{A} with a strength as well. The bifunctor \oplus is also endowed with a strength, simply using the distributivity of \otimes over \oplus (which results from the monoidal closeness of \mathcal{L}). By duality again, $\&$ inherits a strength. The functor $!_-$ is equipped with the strength

$$!X \otimes !Y \xrightarrow{\text{dig}_X \otimes !Y} !!X \otimes !Y \xrightarrow{\mu^2} !(X \otimes Y) .$$

E. Fixed Points of strong functors

The following facts are standard in the literature on fixed points of functors.

Definition 2. Let \mathcal{A} be a category and $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{A}$ be a functor. A *coalgebra*¹² of \mathcal{F} is a pair (A, f) where A is an object of \mathcal{A} and $f \in \mathcal{A}(A, \mathcal{F}(A))$. Given two coalgebras (A, f) and (A', f') of \mathcal{F} , a coalgebra morphism from (A, f) to (A', f') is an $h \in \mathcal{A}(A, A')$ such that $f' h = \mathcal{F}(h) f$. The category of coalgebras of the functor \mathcal{F} will be denoted as $\mathbf{Coalg}_{\mathcal{A}}(\mathcal{F})$. The notion of algebra of an endofunctor is defined dually (reverse the directions of the arrows f and f') and the corresponding category is denoted as $\mathbf{Alg}_{\mathcal{A}}(\mathcal{F})$.

By Lambek's Lemma, if (A, f) with $f \in \mathcal{A}(A, \mathcal{F}(A))$ is a final object in $\mathbf{Coalg}_{\mathcal{A}}(\mathcal{F})$ then f is an iso. We assume that this iso is always the identity¹³ as this holds in our concrete models so that this final object $(\nu\mathcal{F}, \text{Id})$ satisfies $\mathcal{F}(\nu\mathcal{F}) = \nu\mathcal{F}$. We focus on coalgebras rather than algebras for reasons which will become clear when we deal with fixed points of strong functors. This universal property of $\nu\mathcal{F}$ gives us a powerful tool for proving equalities of morphisms.

¹¹This definition, as well as the following one, shows that our assumption that the strength is available for "context object" of shape $!X$ only cannot be disposed of.

¹²Not to be confused with the coalgebras of Section II-C which must satisfy additional properties of compatibility with the comonad structure of $!_-$.

¹³This assumption is highly debatable from the view point of category theory where the notion of equality of objects is not really meaningful. It will be dropped in a longer version of this paper.

Lemma 3. Let $A \in \text{Obj}(\mathcal{A})$ and let $f_1, f_2 \in \mathcal{A}(A, \nu\mathcal{F})$. If there exists $l \in \mathcal{A}(A, \mathcal{F}(A))$ such that $\mathcal{F}(f_i) l = f_i$ for $i = 1, 2$, then $f_1 = f_2$.

Lemma 4. Let $\mathcal{F} : \mathcal{B} \times \mathcal{A} \rightarrow \mathcal{A}$ be a functor such that, for all $B \in \text{Obj}(\mathcal{B})$, the category $\mathbf{Coalg}_{\mathcal{A}}(\mathcal{F}_B)$ has a final object. Then there is a functor $\nu\mathcal{F} : \mathcal{B} \rightarrow \mathcal{A}$ such that $(\nu\mathcal{F}(B), \text{Id})$ is the final object of $\mathbf{Coalg}_{\mathcal{A}}(\mathcal{F}_B)$ (so that $\mathcal{F}(B, \nu\mathcal{F}(B)) = \nu\mathcal{F}(B)$) for each $B \in \text{Obj}(\mathcal{B})$, and, for each $g \in \mathcal{B}(B, B')$, $\nu\mathcal{F}(g)$ is uniquely characterized by $\mathcal{F}(g, \nu\mathcal{F}(g)) = \nu\mathcal{F}(g)$.

We consider now the same $\nu\mathcal{F}$ operation applied to strong functors on a model \mathcal{L} of LL. Let \mathbb{F} be an $n + 1$ -ary strong functor on \mathcal{L} (so that $\overline{\mathbb{F}}$ is a functor $\mathcal{L}^n \times \mathcal{L} \rightarrow \mathcal{L}$). Assume that for each $\overrightarrow{X} \in \text{Obj}(\mathcal{L}^n)$ the category $\mathbf{Coalg}_{\mathcal{L}}(\overline{\mathbb{F}}_{\overrightarrow{X}})$ has a final object. We have defined $\nu\overline{\mathbb{F}} : \mathcal{L}^n \rightarrow \mathcal{L}$ characterized by $\overline{\mathbb{F}}(\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X})) = \nu\overline{\mathbb{F}}(\overrightarrow{X})$ and $\overline{\mathbb{F}}(\overrightarrow{f}, \nu\overline{\mathbb{F}}(\overrightarrow{f})) = \nu\overline{\mathbb{F}}(\overrightarrow{f})$ for all $\overrightarrow{f} \in \mathcal{L}^n(\overrightarrow{X}, \overrightarrow{X}')$ (Lemma 4). For each $Y, \overrightarrow{X} \in \mathcal{L}$, we define $\widehat{\nu\overline{\mathbb{F}}}_{Y, \overrightarrow{X}} \in \mathcal{L}(!Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X}), \nu\overline{\mathbb{F}}(!Y \otimes \overrightarrow{X}))$. We have

$$!Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X}) = !Y \otimes \overline{\mathbb{F}}(\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X})) \xrightarrow{\overline{\mathbb{F}}_{Y, (\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X}))}} \overline{\mathbb{F}}(!Y \otimes \overrightarrow{X}, !Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X}))$$

exhibiting a $\overline{\mathbb{F}}_{!Y \otimes \overrightarrow{X}}$ -coalgebra structure on $!Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X})$. Since $\nu\overline{\mathbb{F}}(!Y \otimes \overrightarrow{X})$ is the final coalgebra of the functor $\overline{\mathbb{F}}_{!Y \otimes \overrightarrow{X}}$, we define $\widehat{\nu\overline{\mathbb{F}}}_{Y, \overrightarrow{X}}$ as the unique morphism $!Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X}) \rightarrow \nu\overline{\mathbb{F}}(!Y \otimes \overrightarrow{X})$ such that the following diagram commutes

$$\begin{array}{ccc}
!Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X}) & \xrightarrow{\overline{\mathbb{F}}_{Y, (\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X}))}} & \overline{\mathbb{F}}(!Y \otimes \overrightarrow{X}, !Y \otimes \nu\overline{\mathbb{F}}(\overrightarrow{X})) \\
\searrow \widehat{\nu\overline{\mathbb{F}}}_{Y, \overrightarrow{X}} & & \downarrow \overline{\mathbb{F}}_{!Y \otimes \overrightarrow{X}, \widehat{\nu\overline{\mathbb{F}}}_{Y, \overrightarrow{X}}} \\
& & \overline{\mathbb{F}}(!Y \otimes \overrightarrow{X}, \nu\overline{\mathbb{F}}(!Y \otimes \overrightarrow{X})) = \nu\overline{\mathbb{F}}(!Y \otimes \overrightarrow{X})
\end{array}$$

Lemma 5. Let \mathbb{F} be an $n + 1$ -ary strong functor on \mathcal{L} such that for each $\overrightarrow{X} \in \text{Obj}(\mathcal{L}^n)$, the category $\mathbf{Coalg}_{\mathcal{L}}(\overline{\mathbb{F}}_{\overrightarrow{X}})$ has a final object $\nu\overline{\mathbb{F}}_{\overrightarrow{X}}$. Then there is a unique n -ary strong functor $\nu\mathbb{F}$ such that $\overline{\mathbb{F}}(\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X})) = \nu\overline{\mathbb{F}}_{\overrightarrow{X}}$ (and hence $\overline{\mathbb{F}}(\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X})) = \nu\overline{\mathbb{F}}(\overrightarrow{X})$), $\overline{\mathbb{F}}(\overrightarrow{f}, \nu\overline{\mathbb{F}}(\overrightarrow{f})) = \nu\overline{\mathbb{F}}(\overrightarrow{f})$ for all $\overrightarrow{f} \in \mathcal{L}^n(\overrightarrow{X}, \overrightarrow{X}')$ and $\overline{\mathbb{F}}(!Y \otimes \overrightarrow{X}, \widehat{\nu\overline{\mathbb{F}}}_{Y, \overrightarrow{X}}) \xrightarrow{\overline{\mathbb{F}}_{Y, (\overrightarrow{X}, \nu\overline{\mathbb{F}}(\overrightarrow{X}))}} \widehat{\nu\overline{\mathbb{F}}}_{Y, \overrightarrow{X}}$.

Lemma 6. Let \mathbb{F} be an $n + 1$ -ary strong functor on \mathcal{L} such that for each $\overrightarrow{X} \in \text{Obj}(\mathcal{L}^n)$, the category $\mathbf{Alg}_{\mathcal{L}}(\overline{\mathbb{F}}_{\overrightarrow{X}})$ has an initial object $\mu\overline{\mathbb{F}}_{\overrightarrow{X}}$. Then there is a unique n -ary strong functor $\mu\mathbb{F}$ such that $\overline{\mathbb{F}}(\overrightarrow{X}, \mu\overline{\mathbb{F}}(\overrightarrow{X})) = \mu\overline{\mathbb{F}}_{\overrightarrow{X}}$ (and hence $\overline{\mathbb{F}}(\overrightarrow{X}, \mu\overline{\mathbb{F}}(\overrightarrow{X})) = \mu\overline{\mathbb{F}}(\overrightarrow{X})$), $\overline{\mathbb{F}}(\overrightarrow{f}, \mu\overline{\mathbb{F}}(\overrightarrow{f})) = \mu\overline{\mathbb{F}}(\overrightarrow{f})$ for all $\overrightarrow{f} \in \mathcal{L}^n(\overrightarrow{X}, \overrightarrow{X}')$

and $\widehat{\mathbb{F}}(!Y \otimes \overrightarrow{X}, \widehat{\mu\mathbb{F}}_{Y, \overrightarrow{X}}) \widehat{\mathbb{F}}_{Y, (\overrightarrow{X}, \overline{\mu\mathbb{F}}(\overrightarrow{X}))} = \widehat{\mu\mathbb{F}}_{Y, \overrightarrow{X}}$. Moreover $(\mu\mathbb{F})^\perp = \nu(\mathbb{F}^\perp)$

Proof: Apply Lemma 5 to the strong functor \mathbb{F}^\perp . ■

F. A categorical axiomatization of models of μLL

Our general definition of Seely categorical model of μLL is based on the notions and results above. We refer in particular to Section II-D for the basic definitions of operations on strong functors in our LL categorical setting.

Definition 7. A *categorical model* or *Seely model* of μLL is a pair $(\mathcal{L}, \overrightarrow{\mathcal{L}})$ where

- 1 \mathcal{L} is a Seely category
- 2 $\overrightarrow{\mathcal{L}} = (\mathcal{L}_n)_{n \in \mathbb{N}}$ where \mathcal{L}_n is a class of strong functors $\mathcal{L}^n \rightarrow \mathcal{L}$, and $\mathcal{L}_0 = \text{Obj}(\mathcal{L})$
- 3 if $\mathbb{X} \in \mathcal{L}_n$ and $\mathbb{X}_i \in \mathcal{L}_k$ (for $i = 1, \dots, n$) then $\mathbb{X} \circ \overrightarrow{\mathbb{X}} \in \mathcal{L}_k$ and all k projection strong functors $\mathcal{L}^k \rightarrow \mathcal{L}$ belong to \mathcal{L}_k
- 4 the strong functors \otimes and $\&$ belong to \mathcal{L}_2 , the strong functor $!_-$ belongs to \mathcal{L}_1 and, if $\mathbb{X} \in \mathcal{L}_n$, then $\mathbb{X}^\perp \in \mathcal{L}_n$
- 5 and last, for all $\mathbb{X} \in \mathcal{L}_1$ the category $\text{Coalg}_{\mathcal{L}}(\overline{\mathbb{X}})$ (see Section II-E) has a final object. So for any $\mathbb{X} \in \mathcal{L}_{k+1}$ there is a strong functor $\nu\mathbb{X} : \mathcal{L}^k \rightarrow \mathcal{L}$ (see Lemma 5). It is required that $\nu\mathbb{X} \in \mathcal{L}_k$.

Our goal is now to outline the interpretation of μLL formulas and proofs in such a model. This requires first to describe the syntax of formulas and proofs.

Remark 3. One can certainly also define a notion of categorical model of μLL in a linear-non-linear adjunction setting as presented in [6]. This is postponed to further work.

1) *Syntax of μLL :* We assume to be given an infinite set of propositional variables \mathcal{V} (ranged over by Greek letters ζ, ξ, \dots). We introduce a language of propositional LL formulas with least and greatest fixed points.

$$A, B, \dots := 1 \mid \perp \mid A \otimes B \mid A \wp B \\ \mid 0 \mid \top \mid A \oplus B \mid A \& B \mid !A \mid ?A \mid \zeta \mid \mu\zeta A \mid \nu\zeta A.$$

The notion of closed formula is defined as usual, the two last constructions being the only binders.

Remark 4. In contrast with second-order LL or dependent type systems where open formulas play a crucial role, in the case of fixed points, all formulas appearing in sequents and other syntactical devices allowing to give types to programs will be closed. In our setting, open types/formulas appear only locally, for allowing the expression of (least and greatest) fixed points.

We can define two basic operations on formulas.

- *Substitution:* $A[B/\zeta]$, taking care of not binding free variables (uses α -conversion).
- *Negation* or *dualization:* defined by induction on formulas $1^\perp = \perp, \perp^\perp = 1, (A \wp B)^\perp = A^\perp \otimes B^\perp, (A \otimes B)^\perp = A^\perp \wp B^\perp, 0^\perp = \top, \top^\perp = 0, (A \& B)^\perp = A^\perp \oplus B^\perp, (A \oplus B)^\perp = A^\perp \& B^\perp, (!A)^\perp = ?A^\perp, (?A)^\perp = !A^\perp, \zeta^\perp = \zeta, (\mu\zeta A)^\perp = \nu\zeta A^\perp$ and $(\nu\zeta A)^\perp = \mu\zeta A^\perp$. Obviously $A^{\perp\perp} = A$ for any formula A .

Remark 5. The only subtle point of this definition is negation of propositional variables: $\zeta^\perp = \zeta$. This entails $(B[A/\zeta])^\perp = B^\perp[A^\perp/\zeta]$ by an easy induction on B . If we consider B as a compound logical connective with placeholders labeled by variables then B^\perp is its De Morgan dual. This definition of ζ^\perp is also a natural way of preventing the introduction of fixed points wrt. variables with negative occurrences. As an illustration, if we define as usual $A \multimap B$ as $A^\perp \wp B$ then we can define $E = \mu\zeta(1 \& (!\zeta \multimap \zeta))$ which *looks like* the definition of a model of the pure λ -calculus as a recursive type. But this is only an illusion since we actually have $E = \mu\zeta(1 \& (? \zeta \wp \zeta))$ so that $!E \multimap E$ is not a retract of E . And indeed if it were possible to define a type D such that $!D \multimap D$ is isomorphic to (or is a retract of) D then we would be able to type all pure λ -terms in our system and this would contradict the fact that μLL enjoys strong normalization and has a denotational semantics based on totality as shown below.

Our logical system extends the usual unilateral sequent calculus of classical propositional LL [1], see also [6] Section 3.1 and 3.13. In this setting we deal with sequents $\vdash A_1, \dots, A_n$ where the A_i 's are formulas. It is important to notice that the order of formulas in this list is not relevant, which means that we keep the exchange rule implicit as it is usual in sequent calculus. To the standard rules¹⁴ of [6] Fig. 1, we add the two next introduction rules for fixed point formulas which are essentially borrowed to [4] (see Section II-F2)

$$\frac{\vdash \Gamma, F[\mu\zeta F/\zeta]}{\vdash \Gamma, \mu\zeta F} (\mu\text{-fold}) \quad \frac{\vdash \Delta, A \quad \vdash ?\Gamma, A^\perp, F[A/\zeta]}{\vdash \Delta, ?\Gamma, \nu\zeta F} (\nu\text{-rec}).$$

By taking, in the last rule, $\Delta = A^\perp$ and proving the left premise by an axiom, we obtain the following derived rule

$$\frac{\vdash ?\Gamma, A^\perp, F[A/\zeta]}{\vdash ?\Gamma, A^\perp, \nu\zeta F} (\nu\text{-rec}').$$

The corresponding cut-elimination rule is described in Section II-F4. For the other connectives (which are the standard connectives of LL), the cut-elimination rules are the usual ones as described in [1], [6].

2) *Comments:* Let us summarize and comment the differences between our system and Baelde's μMALL .

- Baelde's logical system is a *predicate calculus* whereas our system is a *propositional calculus*. Indeed, Baelde is mainly interested in applying μMALL to program verification where the predicate calculus is essential for expressing properties of programs. We have a Curry-Howard perspective where formulas are seen at types and proofs as programs and where a propositional logical system is sufficient.
- Our system has exponentials whereas Baelde's system has not because they can be encoded in μMALL to some extent. However the exponentials encoded in that way do not satisfy all required isos (in particular the "Seely morphisms" are not isos with Baelde's exponentials) and this is a serious issue

¹⁴Notice that the *promotion rule* of LL has a condition on contexts similar to that of the rule $(\nu\text{-rec})$ below: to deduce $\vdash \Delta, !A$ from $\vdash \Delta, A$ it is required that all formulas in the context Δ are of shape $?B$, that is $\Delta = ?\Gamma$.

if we want to encode some form of λ -calculus in the system and consider it as a programming language.

- Our $(\nu\text{-rec})$ rule differs from Baelde's by the fact that we admit a context in the right premise. Notice that all formulas of this context must bear a $?_$ modality: this restriction is absolutely crucial for allowing to express the cut-elimination rule in Section II-F4 which uses an operation of substitution of proofs in formulas and this operation uses structural rules on the context. The semantic counterpart of this operation is described in Section II-D1 where it appears clearly that it uses the fact that P is an object of $\mathcal{L}^!$. Such a version of $(\nu\text{-rec})$ with a context would be problematic in Baelde's system by lack of built-in exponentials.

3) *Syntactic functoriality of formulas*: The reduction rule for the $(\mu\text{-fold})/(\nu\text{-rec})$ cut requires the possibility of substituting a proof for a propositional variable in a formula. More precisely, let $(\zeta, \xi_1, \dots, \xi_k)$ be a list of pairwise distinct propositional variables containing all the free variables of a formula F and let $\vec{C} = (C_1, \dots, C_k)$ be a sequence of closed formulas. Let π be a proof of $\vdash ?\Gamma, A^\perp, B$, then one defines¹⁵ a proof $F \left[\pi/\zeta, \vec{C}/\vec{\xi} \right]$ of

$$\vdash ?\Gamma, (F \left[A/\zeta, \vec{C}/\vec{\xi} \right])^\perp, F \left[B/\zeta, \vec{C}/\vec{\xi} \right]$$

by induction on F , adapting the corresponding definition in [4]. We illustrate this definition by two inductive steps. Observe, in these examples, how the exchange rule is used implicitly.

Assume first that $F = \mu\xi G$ (so that $(\zeta, \xi, \xi_1, \dots, \xi_k)$ is a list of pairwise distinct variables containing all free variables of G). Let $G' = G \left[\vec{C}/\vec{\xi} \right]$ whose only possible free variables are ζ and ξ . The proof $F \left[\pi/\zeta, \vec{C}/\vec{\xi} \right]$ is defined by

$$\frac{\frac{\frac{\vdots G \left[\pi/\zeta, (\mu\xi G') [B/\zeta]/\xi, \vec{C}/\vec{\xi} \right]}{\vdash ?\Gamma, (G' [A/\zeta, (\mu\xi G') [B/\zeta]/\xi)]^\perp, G' [B/\zeta, (\mu\xi G') [B/\zeta]/\xi]}{\vdash ?\Gamma, (G' [A/\zeta, (\mu\xi G') [B/\zeta]/\xi)]^\perp, (\mu\xi G') [B/\zeta]} (\mu\text{-fold})}{\vdash ?\Gamma, ((\mu\xi G') [A/\zeta])^\perp, (\mu\xi G') [B/\zeta]} (\nu\text{-rec}')$$

Notice that this case uses the additional parameters \vec{C} in the definition of this substitution with $k + 1$ parameters in the inductive hypothesis. To see that the last inference in this deduction is an instance of $(\nu\text{-rec}')$, set $H = G'^\perp [A^\perp/\zeta]$ and notice that $(G' [A/\zeta, (\mu\xi G') [B/\zeta]/\xi])^\perp = H \left[(\mu\xi G') [B/\zeta]^\perp/\xi \right]$ and $((\mu\xi G') [A/\zeta])^\perp = \nu\xi H$. Another example is $F = G_1 \otimes G_2$: $F \left[\pi/\zeta, \vec{C}/\vec{\xi} \right]$ is defined as

$$\frac{\frac{\frac{\frac{\vdots G_1 \left[\pi/\zeta, \vec{C}/\vec{\xi} \right]}{\vdash ?\Gamma, (G'_1 [A/\zeta])^\perp, G'_1 [B/\zeta]} (\otimes)}{\vdash ?\Gamma, ?\Gamma, (G'_1 [A/\zeta])^\perp, (G'_2 [A/\zeta])^\perp, G'_1 [B/\zeta] \otimes G'_2 [B/\zeta]} (\otimes)}{\vdash ?\Gamma, (G'_1 [A/\zeta])^\perp, (G'_2 [A/\zeta])^\perp, G'_1 [B/\zeta] \otimes G'_2 [B/\zeta]} (\otimes)}{\vdash ?\Gamma, (G'_1 [A/\zeta])^\perp \wp (G'_2 [A/\zeta])^\perp, G'_1 [B/\zeta] \otimes G'_2 [B/\zeta]} (\wp)} (\otimes)$$

Observe that we use in an essential way the fact that all formulas of the context are of shape $?H$ (even if F is exponential-free) when we apply contraction rules on this context.

4) *Cut elimination*: The only reduction that we will mention here is $(\mu\text{-fold})/(\nu\text{-rec})$. Let θ be

$$\frac{\frac{\frac{\vdots \pi}{\vdash \Delta, F [\mu\zeta F/\zeta]} (\mu\text{-fold})}{\vdash \Delta, \mu\zeta F} \quad \frac{\frac{\frac{\vdots \lambda}{\vdash \Delta, A^\perp} \quad \frac{\vdots \rho}{\vdash ?\Gamma, A, (F [A/\zeta])^\perp}}{\vdash \Delta, ?\Gamma, (\mu\zeta F)^\perp} (\nu\text{-rec})}{\vdash \Delta, \Delta, ?\Gamma} (\text{cut})$$

and let ρ' be the proof

$$\frac{\vdots \rho}{\vdash ?\Gamma, A, (F [A/\zeta])^\perp} (\nu\text{-rec}') \quad \frac{}{\vdash ?\Gamma, A, (\mu\zeta F)^\perp}$$

Then θ reduces to the proof shown in Figure 2. This reduction rule uses the functoriality of formulas as well as the $?_$ -contexts in the $(\nu\text{-rec})$ rule.

Remark 6. In [4] it is shown that μMALL enjoys cut-elimination. We will show in a further paper how this method based on reducibility can be adapted to our μLL . Notice that a cut-free proof has not the sub-formula property in general because of rule $(\nu\text{-rec})$. Though, the normalization theorem makes sure that a proof of a sequent *which does not contain any ν -formula* reduces to a cut-free proof enjoying the sub-formula property.

5) *Interpreting formulas and proofs (outline)*: We assume to be given a μLL Seely model $(\mathcal{L}, \vec{\mathcal{L}})$, see Section II-F. With any formula A and any repetition-free sequence $\vec{\zeta} = (\zeta_1, \dots, \zeta_k)$ of type variables containing all the free variables of A , we associate $\llbracket A \rrbracket_{\vec{\zeta}} \in \mathcal{L}_k$ in the obvious way, for instance $\llbracket A \otimes B \rrbracket_{\vec{\zeta}} = \otimes \circ (\llbracket A \rrbracket_{\vec{\zeta}}, \llbracket B \rrbracket_{\vec{\zeta}}) \in \mathcal{L}_k$ by conditions (4) and (3) in Definition 7 and $\llbracket \nu\zeta A \rrbracket_{\vec{\zeta}} = \nu(\llbracket A \rrbracket_{\vec{\zeta}, \zeta})$ using condition (5). Then $\llbracket A^\perp \rrbracket_{\vec{\zeta}} = \llbracket A \rrbracket_{\vec{\zeta}}^\perp$ up to a natural isomorphism. In this outline, we keep symmetric monoidality isomorphisms of \mathcal{L} and of $\!_!$ implicit (see for instance [35] how *monoidal trees* allow to take them into account). With any $\Gamma = (A_1, \dots, A_n)$ we associate an object $\llbracket \Gamma \rrbracket$ of \mathcal{L} and with any proof π of $\vdash \Gamma$ we associate a morphism $\llbracket \pi \rrbracket \in \mathcal{L}(1, \llbracket \Gamma \rrbracket)$ using the categorical constructs of \mathcal{L} in a straightforward way, see [6]. Then one proves that if π and π' are proofs of $\vdash \Gamma$ and π reduces to π' by the cut-elimination rules, then $\llbracket \pi \rrbracket = \llbracket \pi' \rrbracket$. This is done by an inspection of the various cut-elimination rules. In the case of $(\mu\text{-fold})/(\nu\text{-rec})$ cut-elimination, we need the following lemma.

¹⁵Again the fact that the formulas of the context bear a $?_$ is absolutely necessary to make this definition possible.

$$\begin{array}{c}
\vdots F[\rho'/\zeta] \qquad \qquad \qquad \vdots \pi \qquad \qquad \qquad \vdots \lambda \qquad \qquad \qquad \vdots \rho \\
\vdash ?\Gamma, F[A/\zeta], (F[\mu\zeta F/\zeta])^\perp \quad \vdash \Delta, F[\mu\zeta F/\zeta] \quad (\text{cut}) \quad \vdash \Delta, A^\perp \quad \vdash ?\Gamma, A, (F[A/\zeta])^\perp \quad (\text{cut}) \\
\hline
\vdash \Delta, ?\Gamma, F[A/\zeta] \qquad \qquad \qquad \vdash \Delta, ?\Gamma, (F[A/\zeta])^\perp \quad (\text{cut}) \\
\hline
\vdash \Delta, \Delta, ?\Gamma, ?\Gamma \\
\hline
\vdash \Delta, \Delta, ?\Gamma \\
\hline
\vdash \Delta, \Delta, ?\Gamma \quad (\text{c}) \\
\hline
\vdash \Delta, \Delta, ?\Gamma
\end{array}$$

Fig. 2. Reduction (μ -fold)/(ν -rec)

Lemma 8. Let $\Gamma = (D_1, \dots, D_n)$ be a sequence of closed formulas and let $P = ![D_1] \otimes \dots \otimes ![D_n]$, considered as an object of \mathcal{L}^\dagger . Let F be a formula and $\zeta, \xi_1, \dots, \xi_k$ be a repetition-free list of variables containing all the free variables of F so that $\mathbb{F} = \llbracket F \rrbracket_{\zeta, \vec{\xi}}$ is a strong functor $\mathcal{L}^{k+1} \rightarrow \mathcal{L}$ which belongs to \mathcal{L}_k . As shown in Section II-D this strong functor lifts to a functor $\mathbb{F}[P] : \mathcal{L}[P]^{k+1} \rightarrow \mathcal{L}[P]$. Let π be a proof of $\vdash ?\Gamma, A^\perp, B$, so that $\llbracket \pi \rrbracket \in \mathcal{L}[P](\llbracket A \rrbracket, \llbracket B \rrbracket)$. Let $\vec{C} = (C_1, \dots, C_k)$ be a list of closed formulas. Then

$$\begin{aligned}
\llbracket F \left[\pi/\zeta, \vec{C}/\vec{\xi} \right] \rrbracket &= \mathbb{F}[P](\llbracket \pi \rrbracket, \llbracket C_1 \rrbracket, \dots, \llbracket C_k \rrbracket) \\
&\in \mathcal{L}[P](\mathbb{F}(\llbracket A \rrbracket, \llbracket C_1 \rrbracket, \dots, \llbracket C_k \rrbracket), \mathbb{F}(\llbracket B \rrbracket, \llbracket C_1 \rrbracket, \dots, \llbracket C_k \rrbracket)).
\end{aligned}$$

The proof of the lemma is a simple verification. Notice that we use the fact that the objects of $\mathcal{L}[P]$ are the same as those of \mathcal{L} .

III. SETS AND RELATIONS

The category \mathbf{Rel} has sets as objects, and given sets E and F , $\mathbf{Rel}(E, F) = \mathcal{P}(E \times F)$. Identity is the diagonal relation and composition is the usual composition of relations, denoted by simple juxtaposition. If $t \in \mathbf{Rel}(E, F)$ and $u \subseteq E$ then $t \cdot u = \{b \in F \mid \exists a \in u (a, b) \in t\}$.

A. Rel as a model of LL

This category is a well-known model of LL in which $1 = \perp = \{*\}$, $E \otimes F = (E \multimap F) = E \wp F = E \times F$ so that $E^\perp = E$. As to the additives, $0 = \top = \emptyset$ and $\&_{i \in I} E_i = \oplus_{i \in I} E_i = \bigcup_{i \in I} \{i\} \times E_i$. The exponentials are given by $!E = ?E = \mathcal{M}_{\text{fin}}(E)$ (finite multisets of elements of E which are functions $m : E \rightarrow \mathbb{N}$ such that $m(a) \neq 0$ for finitely many a 's; addition of multisets is defined in the obvious pointwise way, and $[a_1, \dots, a_k]$ is the multiset which maps a to the number of i 's such that $a_i = a$). For the additives and multiplicatives, the operations on morphisms are defined in the obvious way. Let us be more specific about the exponentials. Given $s \in \mathbf{Rel}(E, F)$, $!s \in \mathbf{Rel}(!E, !F)$ is $!s = \{([a_1, \dots, a_n], [b_1, \dots, b_n]) \mid n \in \mathbb{N} \text{ and } \forall i (a_i, b_i) \in s\}$, $\text{der}_E \in \mathbf{Rel}(!E, E)$ is given by $\text{der}_E = \{([a], a) \mid a \in E\}$ and $\text{dig}_E \in \mathbf{Rel}(!E, !!E)$ is given by $\text{dig}_E = \{(m_1 + \dots + m_n, [m_1, \dots, m_n]) \mid \forall i m_i \in \mathcal{M}_{\text{fin}}(E)\}$. Last $m^0 \in \mathbf{Rel}(1, !\top)$ is $m^0 = \{(*, [])\}$ and $m_{E,F}^2 \in \mathbf{Rel}(!E \otimes !F, !(E \& F))$ is given by

$$\begin{aligned}
m_{E,F}^2 &= \{([a_1, \dots, a_k], [b_1, \dots, b_l]), \\
&\quad [(1, a_1), \dots, (1, a_k), (2, b_1), \dots, (2, b_l)] \mid \\
&\quad a_1, \dots, a_k \in E \text{ and } b_1, \dots, b_l \in F\}.
\end{aligned}$$

Weakening $w_E \in \mathbf{Rel}(!E, 1)$ and $\text{contr}_E \in \mathbf{Rel}(!E, !E \otimes !E)$ are given by $w_E = \{([], *)\}$ and $\text{contr}_E = \{(m_1 + m_2, (m_1, m_2)) \mid m_i \in \mathcal{M}_{\text{fin}}(E) \text{ for } i = 1, 2\}$.

B. Hom-continuous functors on Rel

A functor $\mathbb{F} : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ is hom-continuous¹⁶ if, for all $\vec{E}, \vec{F} \in \mathbf{Rel}^n$ and all directed set $D \subseteq \mathbf{Rel}^n(\vec{E}, \vec{F})$, one has $\mathbb{F}(\bigcup D) = \bigcup \{\mathbb{F}(\vec{s}) \mid \vec{s} \in D\}$. This implies in particular that if $\vec{s} \subseteq \vec{t}$, one has $\mathbb{F}(\vec{s}) \subseteq \mathbb{F}(\vec{t})$ (taking $D = \{\vec{s}, \vec{t}\}$). To simplify notations assume that $n = 1$ (but what follows holds for all values of n).

Lemma 9. Let E and F be sets and let $s \in \mathbf{Rel}(E, F)$ and $t \in \mathbf{Rel}(F, E)$. Assume that $ts = \text{Id}_E$ and that $st \subseteq \text{Id}_F$. Then s is (the graph of) an injective function and $t = \{(b, a) \in F \times E \mid (a, b) \in s\}$.

Lemma 10. Let $\mathbb{F} : \mathbf{Rel} \rightarrow \mathbf{Rel}$ be a hom-continuous functor. Assume that $E \subseteq F$ and let $\eta_{E,F}^+ = \{(a, a) \mid a \in E\} \in \mathbf{Rel}(E, F)$ and $\eta_{E,F}^- = \{(a, a) \mid a \in E\} \in \mathbf{Rel}(F, E)$. Then $\mathbb{F}(\eta_{E,F}^+) \in \mathbf{Rel}(\mathbb{F}(E), \mathbb{F}(F))$ is an injective function.

Proof: We have $\eta_{E,F}^- \eta_{E,F}^+ = \text{Id}_E$ and $\eta_{E,F}^+ \eta_{E,F}^- \subseteq \text{Id}_F$ and hence $\mathbb{F}(\eta_{E,F}^-) \mathbb{F}(\eta_{E,F}^+) = \text{Id}$ by functoriality and $\mathbb{F}(\eta_{E,F}^+) \mathbb{F}(\eta_{E,F}^-) \subseteq \text{Id}$ by hom-continuity. The announced property results from Lemma 9. ■

Let \mathbf{Rel}^\subseteq be the category whose objects are sets and morphisms are set inclusions (so that $\mathbf{Rel}^\subseteq(E, F)$ has $\eta_{E,F}^+$ as unique element if $E \subseteq F$ and is empty otherwise). Then η^+ can be thought of as the ‘‘inclusion functor’’ $\mathbf{Rel}^\subseteq \rightarrow \mathbf{Rel}$, acting as the identity on objects. Obviously, \mathbf{Rel}^\subseteq is cocomplete¹⁷.

Proposition 11. If $\mathbb{F} : \mathbf{Rel} \rightarrow \mathbf{Rel}$ is hom-continuous then $\mathbb{F}\eta^+ : \mathbf{Rel}^\subseteq \rightarrow \mathbf{Rel}$ is directed-cocontinuous (that is, preserves the colimits of directed sets of sets).

Proof: Let \mathcal{D} be a directed set of sets and let H be a set. For each $E \in \mathcal{D}$ let $s_E \in \mathbf{Rel}(\mathbb{F}(E), H)$ so that $(s_E)_{E \in \mathcal{D}}$ defines a cocone, that is, for each $E, F \in \mathcal{D}$ such that $E \subseteq F$, one has $s_E = s_F \mathbb{F}(\eta_{E,F}^+)$. Let $L = \bigcup \mathcal{D}$. Let $s \in \mathbf{Rel}(\mathbb{F}(L), H)$ be given by $s = \bigcup_{E \in \mathcal{D}} s_E \mathbb{F}(\eta_{E,L}^-)$. Let $E \in \mathcal{D}$, we have $s \mathbb{F}(\eta_{E,L}^+) = \bigcup_{F \in \mathcal{D}} s_F \mathbb{F}(\eta_{F,L}^- \eta_{E,L}^+)$ so that $s_E \subseteq s \mathbb{F}(\eta_{E,L}^+)$ (since $s_F \mathbb{F}(\eta_{F,L}^- \eta_{E,L}^+) = s_E$ when $F = E$).

¹⁶Hom-set of \mathbf{Rel}^n s are considered as equipped with the inclusion order on relations. The right setting to express hom-continuity would be that of *cpo-enriched categories* but we don't really need this general concept here.

¹⁷Notice that it is not complete, for instance it has no final object.

We prove the converse inclusion. Let $F \in \mathcal{D}$ and let $G \in \mathcal{D}$ be such that $E, F \subseteq G$ (remember that \mathcal{D} is directed). We have

$$\begin{aligned} s_F \mathbb{F}(\eta_{F,L}^- \eta_{E,L}^+) &= s_F \mathbb{F}(\eta_{F,G}^- \eta_{G,L}^- \eta_{G,L}^+ \eta_{E,G}^+) \\ &= s_F \mathbb{F}(\eta_{F,G}^- \eta_{E,G}^+) = s_G \mathbb{F}(\eta_{F,G}^+) \mathbb{F}(\eta_{F,G}^- \eta_{E,G}^+) \\ &\subseteq s_G \mathbb{F}(\eta_{E,G}^+) = s_E \end{aligned}$$

where we have used the fact that $\eta_{F,G}^- \eta_{F,G}^+ \subseteq \text{Id}_G$ and hence $\mathbb{F}(\eta_{F,G}^- \eta_{F,G}^+) \subseteq \text{Id}_{\mathbb{F}(G)}$ by hom-continuity of \mathbb{F} .

So $s_F \mathbb{F}(\eta_{F,L}^- \eta_{E,L}^+) \subseteq s_E$ for all $F \in \mathcal{D}$ and hence $s \mathbb{F}(\eta_{E,L}^+) \subseteq s_E$ as contended.

Let now $s' \in \mathbf{Rel}(\mathbb{F}(L), H)$ be such that $s' \mathbb{F}(\eta_{E,L}^+) = s_E$ for each $E \in \mathcal{D}$, we show that $s' = s$ thus proving the uniqueness part of the universal property. For $E \in \mathcal{D}$, let $\theta_E = \eta_{E,L}^- \eta_{E,L}^+ \in \mathbf{Rel}(L, L)$. Then $(\theta_E)_{E \in \mathcal{D}}$ is a directed family (for \subseteq) and $\bigcup_{E \in \mathcal{D}} \theta_E = \text{Id}_L$. By hom-continuity of \mathbb{F} , we have

$$\begin{aligned} s' &= s' \text{Id}_{\mathbb{F}(L)} = s' \bigcup_{E \in \mathcal{D}} \mathbb{F}(\theta_E) \\ &= \bigcup_{E \in \mathcal{D}} s' \mathbb{F}(\eta_{E,L}^+) \mathbb{F}(\eta_{E,L}^-) = \bigcup_{E \in \mathcal{D}} s_E \mathbb{F}(\eta_{E,L}^-) = s \end{aligned}$$

by our assumption on s' and by definition of s . This shows that the cocone $(\mathbb{F}(\eta_{E,L}^+))_{E \in \mathcal{D}}$ on $\mathbb{F} \eta^+$ is colimiting, thus proving that $\mathbb{F} \eta^+$ is directed cocontinuous. ■

We know that a hom-continuous functor \mathbb{F} maps inclusions to injections, we shall say that \mathbb{F} is *strict* if it maps inclusions to inclusions, that is, if $E \subseteq F$ then $\mathbb{F}(E) \subseteq \mathbb{F}(F)$ and $\mathbb{F}(\eta_{E,F}^+) = \eta_{\mathbb{F}(E), \mathbb{F}(F)}^+$ (which implies $\mathbb{F}(\eta_{E,F}^-) = \eta_{\mathbb{F}(E), \mathbb{F}(F)}^-$). As a direct consequence of Proposition 11, we get:

Lemma 12. *If \mathbb{F} is strict hom-continuous then, for any directed set of sets \mathcal{D} , one has $\mathbb{F}(\bigcup \mathcal{D}) = \bigcup_{E \in \mathcal{D}} \mathbb{F}(E)$.*

C. Variable sets and basic constructions on them

Definition 13. An n -ary variable set is a strong functor $\mathbb{V} : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ such that $\overline{\mathbb{V}}$ is hom-continuous and strict.

By the general considerations of Section II-D, there is a constant strong functor $\mathbf{Rel}^n \rightarrow \mathbf{Rel}$ with value E for each set E . There are projection strong functors $\mathbf{Rel}^n \rightarrow \mathbf{Rel}$, \times (that is \otimes) and $+$ (that is \oplus) define strong functors $\mathbf{Rel}^2 \rightarrow \mathbf{Rel}$, $\mathcal{M}_{\text{fin}}(_)$ (that is $_!$) defines a strong functor $\mathbf{Rel} \rightarrow \mathbf{Rel}$. Strong functors on \mathbf{Rel} are stable under composition, and if \mathbb{V} is a strong functor $\mathbf{Rel}^n \rightarrow \mathbf{Rel}$ then there is a “dual” strong functor \mathbb{V}^\perp (which is actually identical to \mathbb{V} in this very simple model). To check that these strong functors \mathbb{V} are variable sets we have only to check that the underlying functors $\overline{\mathbb{V}}$ are strict hom-continuous.

We deal with $_!$ and composition, the other cases are similar. We already defined the functor¹⁸ $_!$ in Section III-A. If $s \subseteq t \in \mathbf{Rel}(E, F)$, it follows from the definition that $!s \subseteq !t$. Let $D \subseteq \mathbf{Rel}(E, F)$ be directed, we prove $!(\bigcup D) \subseteq \bigcup_{s \in D} !s$:

¹⁸We use the same notation for the strong functor and its underlying functor, this slight abuse of notations should not lead to confusions.

an element of $!(\bigcup D)$ is a pair $([a_1, \dots, a_k], [b_1, \dots, b_k])$ with $(a_i, b_i) \in \bigcup D$ for $i = 1, \dots, k$. Since D is directed, there is an $s \in D$ such that $(a_i, b_i) \in s$ for $i = 1, \dots, k$ and the inclusion follows. Strictness is obvious.

Let $\mathbb{V}_i : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ be variable sets for $i = 1, \dots, k$ and let $\mathbb{W} : \mathbf{Rel}^k \rightarrow \mathbf{Rel}$ be a variable set. Then the functor $\overline{\mathbb{W}} \circ \overline{\mathbb{V}} : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ is clearly strict hom-continuous (since these conditions are preservation properties) from which it follows that the strong functor $\mathbb{U} = \mathbb{W} \circ \overline{\mathbb{V}}$ is a variable type.

1) *Fixed point of a variable set:* Let $\mathbb{F} : \mathbf{Rel} \rightarrow \mathbf{Rel}$ be a strict hom-continuous functor. Since $\emptyset \subseteq \mathbb{F}(\emptyset)$ we have $\mathbb{F}^n(\emptyset) \subseteq \mathbb{F}^{n+1}(\emptyset)$ for all $n \in \mathbb{N}$, by induction on n and hence $\mathbb{F}(\bigcup_{n=0}^\infty \mathbb{F}^n(\emptyset)) = \bigcup \mathbb{F}^n(\emptyset)$ by Lemma 12 since $\{\mathbb{F}^n(\emptyset) \mid n \in \mathbb{N}\}$ is directed. Let $\sigma \mathbb{F} = \bigcup_{n=0}^\infty \mathbb{F}^n(\emptyset)$, so that $(\sigma \mathbb{F}, \text{Id}_{\sigma \mathbb{F}})$ is an \mathbb{F} -coalgebra.

Lemma 14. *The coalgebra $(\sigma \mathbb{F}, \text{Id})$ is final in $\mathbf{Coalg}_{\mathbf{Rel}}(\mathbb{F})$.*

Notice that $(\sigma \mathbb{F}, \text{Id})$ is also an initial object in $\mathbf{Alg}_{\mathbf{Rel}}(\mathbb{F})$. When we insist on considering $\sigma \mathbb{F}$ as a final coalgebra, we denote it as $\nu \mathbb{F}$.

Lemma 15. *Let $\mathbb{F} : \mathbf{Rel}^{n+1} \rightarrow \mathbf{Rel}$ be a strict hom-continuous functor. The functor $\nu \mathbb{F} : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ is strict hom-continuous.*

Let $\mathbb{V} : \mathbf{Rel}^{n+1} \rightarrow \mathbf{Rel}$ be a variable set, by Lemma 5, there is a unique strong functor $\nu \mathbb{V} : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ which is characterized by: $\overline{\nu \mathbb{V}}(\vec{E}) = \nu \overline{\mathbb{V}}_{\vec{E}}$, for each $\vec{s} \in \mathbf{Rel}^n(\vec{E}, \vec{F})$, $\overline{\nu \mathbb{V}}(\vec{s}) = \overline{\mathbb{V}}(\vec{s}, \nu \overline{\mathbb{V}}(\vec{s}))$ and last $\overline{\mathbb{V}}(!E \otimes \vec{F}, \hat{\mathbb{V}}_{E, \vec{F}}) = \hat{\mathbb{V}}_{E, \vec{F}}$.

Lemma 16. *The functor $\nu \mathbb{V}$ is a variable set.*

Proof: By the conditions above satisfied by $\nu \mathbb{V}$ we have that $\overline{\nu \mathbb{V}} = \nu \overline{\mathbb{V}}$ and hence $\overline{\nu \mathbb{V}}$ is strict hom-continuous by Lemma 15. ■

2) *A model of μLL based on variable sets:* Let \mathbf{Rel}_n be the class of all n -ary variable sets, so that $\mathbf{Rel}_0 = \text{Obj}(\mathbf{Rel})$. The fact that $(\mathbf{Rel}, (\mathbf{Rel}_n)_{n \in \mathbb{N}})$ is a Seely model of μLL in the sense of Definition 7 results mainly from the fact that we take *all* variable sets in the \mathbf{Rel}_n 's so that there is essentially nothing to check. More explicitly: (1) holds by Section III-A, (2) holds by construction, (3) holds by the fact that variable sets compose as explained in Section III-C (notice that this condition refers to the general composition of strong functors defined at the beginning of Section II-D1), (4) holds by Section III-C and by the fact that the De Morgan dual of a strong functor is strong, see Section II-D1 and (5) holds by Lemma 16.

IV. NON-UNIFORM TOTALITY SPACES

We enrich the model of Section III with a notion of totality, we use notations from that section for operations on sets and relations.

A. *Basic definitions.*

Let E be a set and let $\mathcal{T} \subseteq \mathcal{P}(E)$. We define $\mathcal{T}^\perp = \{u' \subseteq E \mid \forall u \in \mathcal{T} u \cap u' \neq \emptyset\}$. If $\mathcal{S} \subseteq \mathcal{T} \subseteq \mathcal{P}(E)$ then $\mathcal{T}^\perp \subseteq \mathcal{S}^\perp$.

We also have $\mathcal{T} \subseteq \mathcal{T}^{\perp\perp}$ and therefore $\mathcal{T}^{\perp\perp\perp} = \mathcal{T}^{\perp}$. The biorthogonal closure has a nice and simple characterization.

Lemma 17. *Let $\mathcal{T} \subseteq \mathcal{P}(E)$, then $\mathcal{T}^{\perp\perp} = \uparrow\mathcal{T} = \{v \subseteq E \mid \exists u \in \mathcal{T} \ u \subseteq v\}$.*

Proof: The \supseteq direction is obvious, the converse is not difficult either: let $u \in \mathcal{T}^{\perp\perp}$. Then $E \setminus u \notin \mathcal{T}^{\perp}$, so there is $v \in \mathcal{T}$ such that $v \cap (E \setminus u) = \emptyset$, that is $v \subseteq u$. Hence $u \in \uparrow\mathcal{T}$. ■

A *non-uniform totality space* (NUTS) is a pair $X = (|X|, \mathcal{T}(X))$ where $|X|$ is a set and $\mathcal{T}(X) \subseteq \mathcal{P}(|X|)$ satisfies $\mathcal{T}(X) = \mathcal{T}(X)^{\perp\perp}$, that is $\mathcal{T}(X) = \uparrow\mathcal{T}(X)$. Of course we set $X^{\perp} = (|X|, \mathcal{T}(X)^{\perp})$.

Example 18. Let $X = (\mathbb{N}, \mathcal{T}(X))$ where $\mathcal{T}(X)$ is the set of all infinite subsets of \mathbb{N} . It is a NUTS because a superset of an infinite set is infinite. Then $|X^{\perp}| = \mathbb{N}$ and $\mathcal{T}(X^{\perp})$ is the set of all cofinite subsets of \mathbb{N} (the subsets u of \mathbb{N} such that $\mathbb{N} \setminus u$ is finite). If, with the same web \mathbb{N} , we take $\mathcal{T}(X) = \{u \subseteq \mathbb{N} \mid u \neq \emptyset\}$ (again $\mathcal{T}(X) = \uparrow\mathcal{T}(X)$ obviously), then $\mathcal{T}(X^{\perp}) = \{\mathbb{N}\}$.

We define four basic NUTS: $0 = (\emptyset, \emptyset)$, $\top = (\emptyset, \{\emptyset\})$ and $1 = \perp = (\{*\}, \{\{*\}\})$. Given NUTS X_1 and X_2 we define a NUTS $X_1 \otimes X_2$ by $|X_1 \otimes X_2| = |X_1| \times |X_2|$ and $\mathcal{T}(X_1 \otimes X_2) = \uparrow\{u_1 \times u_2 \mid u_i \in \mathcal{T}(X_i) \text{ for } i = 1, 2\}$. And then we define $X \multimap Y = (X \otimes Y^{\perp})^{\perp}$.

Lemma 19. $t \in \mathcal{T}(X \multimap Y) \Leftrightarrow \forall u \in \mathcal{T}(X) \ t \cdot u \in \mathcal{T}(Y)$.

We define the category **Nuts** whose objects are the NUTS and $\mathbf{Nuts}(X, Y) = \mathcal{T}(X \multimap Y)$, composition being defined as the usual composition in **Rel** (relational composition) and identities as the diagonal relations. Lemma 19 shows that we have indeed defined a category.

1) *Multiplicative structure:*

Lemma 20. *Let X and Y be NUTS and $t \in \mathbf{Nuts}(X, Y)$. Then t is an iso in **Nuts** iff t is (the graph of) a bijection $|X| \rightarrow |Y|$ such that $\forall u \subseteq |X| \ u \in \mathcal{T}(X) \Leftrightarrow t(u) \in \mathcal{T}(Y)$.*

Lemma 21. *Let $t \subseteq |X| \times |Y|$. One has $t \in \mathbf{Nuts}(X, Y)$ iff $t^{\perp} = \{(b, a) \mid (a, b) \in t\} \in \mathbf{Nuts}(Y^{\perp}, X^{\perp})$.*

Proof: This is an obvious consequence of Lemma 19 and of the fact that $(X \multimap Y)^{\perp} = (X \otimes Y^{\perp})^{\perp}$ and $(Y^{\perp} \multimap X^{\perp}) = (Y^{\perp} \otimes X)^{\perp}$. ■

Lemma 22. *Let $t \subseteq |X_1 \otimes X_2 \multimap Y|$. One has $t \in \mathbf{Nuts}(X_1 \otimes X_2, Y)$ iff for all $u_1 \in \mathcal{T}(X_1)$ and $u_2 \in \mathcal{T}(X_2)$ one has $t \cdot (u_1 \otimes u_2) \in \mathcal{T}(Y)$.*

Lemma 23. *The bijection $\alpha_{|X_1|, |X_2|, |Y|}$ is an isomorphism from $(X_1 \otimes X_2) \multimap Y$ to $X_1 \multimap (X_2 \multimap Y)$.*

We turn now \otimes into a functor, its action on morphisms being defined as in **Rel**. Let $t_i \in \mathbf{Nuts}(X_i, Y_i)$ for $i = 1, 2$, we have $t_1 \otimes t_2 \in \mathbf{Nuts}(X_1 \otimes X_2, Y_1 \otimes Y_2)$ by Lemma 22 and by the equation $(t_1 \otimes t_2) \cdot (u_1 \otimes u_2) = (t_1 \cdot u_1) \otimes (t_2 \cdot u_2)$. This functor is monoidal, with unit 1 and symmetric monoidality

isomorphisms λ, ρ, γ and α defined as in **Rel**. The only non-trivial thing to check is that α is indeed a morphism, namely $\alpha_{|X_1|, |X_2|, |X_3|} \in \mathbf{Nuts}((X_1 \otimes X_2) \otimes X_3, X_1 \otimes (X_2 \otimes X_3))$. This results from Lemma 23 and from the observation that $((X_1 \otimes X_2) \otimes X_3)^{\perp} = ((X_1 \otimes X_2) \multimap X_3^{\perp})^{\perp}$ and $(X_1 \otimes (X_2 \otimes X_3))^{\perp} = (X_1 \multimap (X_2 \multimap X_3^{\perp}))^{\perp}$.

The SMC category **Nuts** is closed, with $X \multimap Y$ as internal hom object from X to Y , and evaluation morphism $\text{ev} = \{((a, b), a), b \mid a \in |X| \text{ and } b \in |Y|\}$ which indeed belongs to $\mathbf{Nuts}((X \multimap Y) \otimes X, Y)$ by Lemma 22 since, for all $t \in \mathcal{T}(X \multimap Y)$ and $u \in \mathcal{T}(X)$ we have $\text{ev}(t \otimes u) = t \cdot u \in \mathcal{T}(Y)$. This category **Nuts** is also $*$ -autonomous with dualizing object $\perp = 1$.

2) *Additive structure:* Let $(X_i)_{i \in I}$ be an at most countable family of objects of **Nuts**. We define $X = \&_{i \in I} X_i$ by: $|X| = \bigcup_{i \in I} \{i\} \times |X_i|$ and $\mathcal{T}(X) = \{u \subseteq |X| \mid \forall i \in I \ \pi_i \cdot u \in \mathcal{T}(X_i)\}$. It is clear that $\mathcal{T}(X) = \uparrow\mathcal{T}(X)$ and hence X is an object of **Nuts**. By definition of X and by Lemma 19 we have $\forall i \in I \ \pi_i \in \mathbf{Nuts}(X, X_i)$. Given $\vec{t} = (t_i)_{i \in I}$ with $\forall i \in I \ t_i \in \mathbf{Nuts}(Y, X_i)$, we have $\langle \vec{t} \rangle \in \mathbf{Nuts}(Y, X)$ as easily checked (using Lemma 19 again). It follows that $(\&_{i \in I} X_i, (\pi_i)_{i \in I})$ is the cartesian product of the X_i 's in **Nuts**. This shows that the category **Nuts** has all countable products and hence is cartesian. Since it is $*$ -autonomous, the category **Nuts** is also cocartesian, coproduct being given by $\oplus_{i \in I} X_i = (\&_{i \in I} X_i^{\perp})^{\perp}$. It follows that $X = \oplus_{i \in I} X_i = (\&_{i \in I} X_i^{\perp})^{\perp}$ satisfies $|X| = \bigcup_{i \in I} \{i\} \times |X_i|$ and

$$\mathcal{T}(X) = \{v \subseteq \bigcup_{i \in I} \{i\} \times |X_i| \mid \exists i \in I \exists u \in \mathcal{T}(X_i) \ \{i\} \times u \subseteq v\}$$

as easily checked. Notice that the final object is $\top = (\emptyset, \{\emptyset\})$ and that $0 = \top^{\perp} = (\emptyset, \emptyset)$.

3) *Exponential:* We extend the exponential of **Rel** with totality. If $u \in \mathcal{P}(|X|)$ we set $u^{(1)} = \mathcal{M}_{\text{fin}}(u) \in \mathcal{P}(|!X|)$. Then we set $!|X| = \mathcal{M}_{\text{fin}}(|X|)$ and $\mathcal{T}(!X) = \{u^{(1)} \mid u \in \mathcal{T}(X)\}^{\perp\perp} = \uparrow\{u^{(1)} \mid u \in \mathcal{T}(X)\}$.

Lemma 24. *Let $t \subseteq \mathcal{M}_{\text{fin}}(|X|) \times |Y|$. One has $t \in \mathbf{Nuts}(!X, Y)$ iff for all $u \in \mathcal{T}(X)$ one has $t \cdot u^{(1)} \in \mathcal{T}(Y)$.*

Lemma 25. *Let $t \subseteq \mathcal{M}_{\text{fin}}(|X_1|) \times \dots \times \mathcal{M}_{\text{fin}}(|X_k|) \times |Y|$. One has $t \in \mathbf{Nuts}(!X_1 \otimes \dots \otimes !X_k, Y)$ iff for all $\vec{u} \in \prod_{i=1}^k \mathcal{T}(X_i)$ one has $t \cdot (u_1^{(1)} \otimes \dots \otimes u_k^{(1)}) \in \mathcal{T}(Y)$.*

Lemma 26. *If $t \in \mathbf{Nuts}(X, Y)$ then $!t \in \mathbf{Nuts}(!X, !Y)$.*

Proof: By Lemma 24 and the fact that $!t \cdot u^{(1)} = (t \cdot u)^{(1)}$. ■

To prove that **Nuts** is a categorical model of LL, it suffices to show that the various relational morphisms defining the strong symmetric monoidal monadic structure of $\underline{!}$ in **Rel** (see Section III-A) are actually morphisms in **Nuts**. This is essentially straightforward and based on Lemma 24.

Lemma 27. *Equipped with der , dig , m^0 and m^2 defined as in **Rel**, $\underline{!}$ is a symmetric monoidal comonad which turns **Nuts** into a Seely model of LL.*

B. Variable non-uniform totality spaces (VNUTS)

Let E be a set, we use $\text{Tot}(E)$ for the set of all *totality candidates* on E , that is, of all subsets \mathcal{T} of $\mathcal{P}(E)$ such that $\mathcal{T} = \mathcal{T}^{\perp\perp}$ (remember that $\mathcal{T}^{\perp} = \{u' \subseteq E \mid \forall u \in \mathcal{T} \ u \cap u' \neq \emptyset\}$). In other words $\mathcal{T} \in \text{Tot}(E)$ means that $\mathcal{T} = \uparrow\mathcal{T}$ by Lemma 17. Ordered by \subseteq , this set $\text{Tot}(E)$ is a complete lattice.

Definition 28. Let $n \in \mathbb{N}$, an n -ary VNUTS is a pair $\mathbb{X} = (|\mathbb{X}|, \mathcal{T}(\mathbb{X}))$ where $|\mathbb{X}| : \mathbf{Rel}^n \rightarrow \mathbf{Rel}$ is a variable set $|\mathbb{X}| = (|\overline{\mathbb{X}}|, \widehat{|\mathbb{X}}|)$ (see Section III-C2) and $\mathcal{T}(\mathbb{X})$ is an operation which with each n -tuple \vec{X} of objects of \mathbf{Nuts} associates an element $\mathcal{T}(\mathbb{X})(\vec{X})$ of $\text{Tot}(|\overline{\mathbb{X}}|(|\vec{X}|))$ in such a way that

- 1 for any $\vec{t} \in \mathbf{Nuts}^n(\vec{X}, \vec{Y})$, the element $|\overline{\mathbb{X}}|(\vec{t})$ of $\mathbf{Rel}(|\overline{\mathbb{X}}|(|\vec{X}|), |\overline{\mathbb{X}}|(|\vec{Y}|))$ belongs to $\mathbf{Nuts}(\overline{\mathbb{X}}(\vec{X}), \overline{\mathbb{X}}(\vec{Y}))$ (where $\overline{\mathbb{X}}(\vec{X})$ denotes the NUTS $(|\overline{\mathbb{X}}|(|\vec{X}|), \mathcal{T}(\mathbb{X})(\vec{X}))$)
- 2 and for any $\vec{Y} \in \text{Obj}(\mathbf{Nuts}^n)$ and any $X \in \text{Obj}(\mathbf{Nuts})$ one has $|\overline{\mathbb{X}}|_{|X|, |\vec{Y}|} \in \mathbf{Nuts}(!X \otimes \overline{\mathbb{X}}(\vec{Y}), \overline{\mathbb{X}}(!X \otimes \vec{Y}))$. In other words, for an $u \in \mathcal{T}(X)$ and $v \in \text{Tot}(\mathbb{X})(\vec{Y})$, one has $|\overline{\mathbb{X}}|_{|X|, |\vec{Y}|} \cdot (u^{(1)} \otimes v) \in \text{Tot}(\mathbb{X})(!X \otimes \vec{Y})$.

Lemma 29. Any VNUTS $\mathbb{X} : \mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$ induces a strong functor $\mathcal{X} : \mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$ which satisfies

- $|\mathcal{X}(\vec{X})| = |\overline{\mathbb{X}}|(|\vec{X}|)$,
 - $\mathcal{T}(\mathcal{X}(\vec{X})) = \mathcal{T}(\mathbb{X})(\vec{X})$,
 - $\mathcal{X}(\vec{t}) = |\overline{\mathbb{X}}|(\vec{t}) \in \mathbf{Nuts}(\overline{\mathbb{X}}(\vec{X}), \overline{\mathbb{X}}(\vec{Y}))$ for $\vec{t} \in \mathbf{Nuts}(\vec{X}, \vec{Y})$,
 - and $\mathcal{X}_{X, \vec{Y}} = |\overline{\mathbb{X}}|_{|X|, |\vec{Y}|}$
- and \mathbb{X} can be retrieved from \mathcal{X} .

Proof: It is clear that \mathcal{X} so defined is a strong functor. Let us check that \mathbb{X} can be retrieved from \mathcal{X} . Given a set E , $(E, \mathcal{P}(E))$ is a NUTS that we denote as $\mathfrak{p}(E)$. Notice that \mathfrak{p} can be extended into a functor $\mathbf{Rel} \rightarrow \mathbf{Nuts}$ which acts as the identity on morphisms. There is also a forgetful functor $\mathfrak{u} : \mathbf{Nuts} \rightarrow \mathbf{Rel}$ which maps X to $|X|$ and acts as the identity on morphisms (btw. \mathfrak{p} is right adjoint to \mathfrak{u}). Let \mathbb{X} be a unary VNUTS and let $\mathcal{X} : \mathbf{Nuts} \rightarrow \mathbf{Nuts}$ be the associated strong functor. Then we have $|\overline{\mathbb{X}}| = \mathfrak{u} \circ \mathcal{X} \circ \mathfrak{p}$ and $|\overline{\mathbb{X}}|_{E, F} = \widehat{\mathcal{X}}_{\mathfrak{p}(E), \mathfrak{p}(F)}$ for any sets E and F . Last, given a NUTS X , we have that $\mathcal{T}(\mathbb{X})(X)$ is just the totality component of the NUTS $\mathcal{X}(X)$. This shows that \mathbb{X} is determined by \mathcal{X} as contended. ■

For this reason we use \mathbb{X} to denote the functor \mathcal{X} . Given $n \in \mathbb{N}$ let \mathbf{Vnuts}_n be the class of strong n -ary VNUTS. We identify \mathbf{Vnuts}_0 with the class of objects of the Seely category \mathbf{Nuts} . The following refers to Definition 7

Theorem 30. $(\mathbf{Nuts}, (\mathbf{Vnuts}_n)_{n \in \mathbb{N}})$ is a Seely model of μLL .

Partial proof: We deal with Condition (5).

Let first $\mathbb{X} = (|\mathbb{X}|, \mathcal{T}(\mathbb{X}))$ be a unary VNUTS. Let $E = \sigma|\overline{\mathbb{X}}|$ which is the least set such that $|\overline{\mathbb{X}}|(E) = E$, that is $E = \bigcup_{n=0}^{\infty} |\overline{\mathbb{X}}|^n(\emptyset)$. Let $\Phi : \text{Tot}(E) \rightarrow \text{Tot}(E)$ be defined as follows: given $S \in \text{Tot}(E)$, then (E, S) is a NUTS, and we set $\Phi(S) = \mathcal{T}(\mathbb{X})(E, S) \in \text{Tot}(|\overline{\mathbb{X}}|(E)) = \text{Tot}(E)$. This function Φ is monotone. Let indeed $S_1, S_2 \in \text{Tot}(E)$ with $S_1 \subseteq S_2$.

Then we have $\text{Id} \in \mathbf{Nuts}((E, S_1), (E, S_2))$ and therefore, by Condition (1) satisfied by \mathbb{X} , we have $\text{Id} = |\overline{\mathbb{X}}|(\text{Id}) \in \mathbf{Nuts}(\overline{\mathbb{X}}(E, S_1), \overline{\mathbb{X}}(E, S_2)) = \mathbf{Nuts}((E, \Phi(S_1)), (E, \Phi(S_2)))$ which means that $\Phi(S_1) \subseteq \Phi(S_2)$. By the Knaster Tarski Theorem (remember that $\text{Tot}(E)$ is a complete lattice), Φ has a greatest fixpoint \mathcal{T} that we can describe as follows. Let $(\mathcal{T}_\alpha)_{\alpha \in \mathbb{O}}$, where \mathbb{O} is the class of ordinals, be defined by: $\mathcal{T}_0 = \mathcal{P}(E)$ (the largest possible notion of totality on E), $\mathcal{T}_{\alpha+1} = \Phi(\mathcal{T}_\alpha)$ and $\mathcal{T}_\lambda = \bigcap_{\alpha < \lambda} \mathcal{T}_\alpha$ when λ is a limit ordinal. This sequence is decreasing (easy induction on ordinals using the monotonicity of Φ) and there is an ordinal θ such that $\mathcal{T}_{\theta+1} = \mathcal{T}_\theta$ (by a cardinality argument; we can assume that θ is the least such ordinal). The greatest fixpoint of Φ is then \mathcal{T}_θ as easily checked.

By construction $((E, \mathcal{T}_\theta), \text{Id})$ is an object of $\mathbf{Coalg}_{\mathbf{Nuts}}(\overline{\mathbb{X}})$, we prove that it is the final object. So let (Y, t) be another object of the same category. Since $(|Y|, t)$ is an object of $\mathbf{Coalg}_{\mathbf{Rel}}(|\overline{\mathbb{X}}|)$ and since (E, Id) is the final object in that category, we know by Lemma 14 that there is exactly one $e \in \mathbf{Rel}(|Y|, E)$ such that $|\overline{\mathbb{X}}|(e) t = e$. We prove that actually $e \in \mathbf{Nuts}(Y, (E, \mathcal{T}_\theta))$ so let $v \in \mathcal{T}(Y)$. We prove by induction on the ordinal α that $e \cdot v \in \mathcal{T}_\alpha$. For $\alpha = 0$ it is obvious since $\mathcal{T}_0 = \mathcal{P}(E)$. Assume that the property holds for α and let us prove it for $\alpha + 1$. We have $t \cdot v \in \mathcal{T}(\mathbb{X})(Y) = \mathcal{T}(\overline{\mathbb{X}}(Y))$ since $t \in \mathbf{Nuts}(Y, \overline{\mathbb{X}}(Y))$. Since $\overline{\mathbb{X}}(e) \in \mathbf{Nuts}(\overline{\mathbb{X}}(Y), \overline{\mathbb{X}}(E, \mathcal{T}_\alpha))$ and $\overline{\mathbb{X}}(E, \mathcal{T}_\alpha) = (E, \mathcal{T}_{\alpha+1})$ we have $(\overline{\mathbb{X}}(e) t) \cdot v \in \mathcal{T}_{\alpha+1}$, that is $e \cdot v \in \mathcal{T}_{\alpha+1}$. Last if λ is a limit ordinal and if we assume $\forall \alpha < \lambda \ e \cdot v \in \mathcal{T}_\alpha$ we have $e \cdot v \in \bigcap_{\alpha < \lambda} \mathcal{T}_\alpha = \mathcal{T}_\lambda$. Therefore $e \cdot v \in \mathcal{T}_\theta$. We use $\nu\overline{\mathbb{X}}$ to denote this final coalgebra (E, \mathcal{T}_θ) (its definition depends only on $\overline{\mathbb{X}}$ and does not involve the strength $\widehat{\mathbb{X}}$).

So we have proven the first part of Condition (5) in the definition of a Seely model of μLL (see Section 7). As to the second part, let \mathbb{X} be an $n + 1$ -ary VNUTS. We know by the general Lemma 5 how to build a strong functor $\nu\mathbb{X} : \mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$ with suitable properties. To end the proof, it suffices to exhibit an n -ary VNUTS $\mathbb{Y} = (|\mathbb{Y}|, \mathcal{T}(\mathbb{Y}))$ whose associated strong functor coincides with $\nu\mathbb{X}$. The construction of \mathbb{Y} is essentially straightforward, using the constructions available in \mathbf{Rel} (see the Appendix). ■

Remark 7. For any closed formula A , the web of its interpretation $\llbracket A \rrbracket^{\mathbf{Nuts}}$ in \mathbf{Nuts} coincides with its interpretation $\llbracket A \rrbracket^{\mathbf{Rel}}$ in \mathbf{Rel} . It is also easy to check that for any proof π of $\vdash A$, one has $\llbracket \pi \rrbracket^{\mathbf{Nuts}} = \llbracket \pi \rrbracket^{\mathbf{Rel}}$ (this can be formalized using the functor $\mathfrak{u} : \mathbf{Nuts} \rightarrow \mathbf{Rel}$ introduced in the proof of Lemma 29, which acts trivially on morphisms).

Remark 8. We have recently been informed that the definition above of the functorial objects VNUTS based on variable sets and the category \mathbf{Nuts} might be an instance of a more general construction based on topological functor. We will develop this idea in further work.

C. Examples of data-types

1) *Integers:* The type of “flat integers” is defined by $\iota = \mu\zeta(1 \oplus \zeta)$. In \mathbf{Rel} , $1 \oplus \zeta$ is interpreted as the unary variable

set $\llbracket 1 \oplus \zeta \rrbracket_{\zeta}^{\mathbf{Rel}} : \mathbf{Rel} \rightarrow \mathbf{Rel}$ which maps a set E to $1 \oplus E = \{(1, *)\} \cup (\{2\} \times E)$. Hence $\llbracket \iota \rrbracket^{\mathbf{Rel}}$ is the least set such that $\llbracket \iota \rrbracket = \{(1, *)\} \cup (\{2\} \times \llbracket \iota \rrbracket)$ that is, the set of all tuples $\bar{n} = (2, (2, (\dots (1, *) \dots)))$ where n is the number of occurrence of 2, that is $\llbracket \iota \rrbracket^{\mathbf{Rel}} = \mathbb{N}$ up to renaming. We have $|\llbracket \iota \rrbracket^{\mathbf{Nuts}}| = |\llbracket \iota \rrbracket^{\mathbf{Rel}}| = \mathbb{N}$ and we compute $\mathcal{T}(\llbracket \iota \rrbracket^{\mathbf{Nuts}})$ dually wrt. the proof of Theorem 30: it is the least fixed point of the operator $\Phi : \text{Tot}(\mathbb{N}) \rightarrow \text{Tot}(\mathbb{N})$ (remember that $\text{Tot}(\mathbb{N})$ is just the set of all \subseteq -upwards-closed subsets of \mathbb{N}) such that, if $\mathcal{T} \in \text{Tot}(\mathbb{N})$ then $\Phi(\mathcal{T}) = \{u \subseteq \mathbb{N} \mid 0 \in u \text{ or } \{n \in \mathbb{N} \mid n+1 \in u\} \in \mathcal{T}\}$. Therefore $\text{Tot}(\llbracket \iota \rrbracket^{\mathbf{Nuts}}) = \{u \subseteq \mathbb{N} \mid u \neq \emptyset\}$.

Theorem 31. *If π is a proof of $\vdash \iota$ then $\llbracket \pi \rrbracket^{\mathbf{Nuts}} = \llbracket \pi \rrbracket^{\mathbf{Rel}}$ is a non-empty subset of \mathbb{N} .*

Indeed we know that $\llbracket \pi \rrbracket^{\mathbf{Rel}} = \llbracket \pi \rrbracket^{\mathbf{Nuts}} \in \mathcal{T}(\llbracket \iota \rrbracket^{\mathbf{Nuts}})$. Using an additional notion of coherence (which can be fully compatible with \mathbf{Rel} as in the non-uniform coherence space models of [12], [13]) we can also prove that $\llbracket \pi \rrbracket^{\mathbf{Rel}}$ has at most one element, and hence is a singleton $\{n\}$. This is a denotational version of normalization expressing that indeed π “has a value” (and actually exactly one, which expresses a weak form of confluence).

2) *Binary trees with integer leaves:* This type can be defined as $\tau = \mu\zeta(\iota \oplus (\zeta \otimes \zeta))$. Then an element of $\llbracket \tau \rrbracket^{\mathbf{Rel}} = |\llbracket \tau \rrbracket^{\mathbf{Nuts}}|$ is an element of the set described by the following syntax: $\alpha, \beta, \dots := \langle n \rangle \mid \langle \alpha, \beta \rangle$. A computation similar to the previous one shows that $\text{Tot}(\llbracket \tau \rrbracket^{\mathbf{Nuts}}) = \{u \subseteq \llbracket \tau \rrbracket^{\mathbf{Rel}} \mid u \neq \emptyset\}$.

3) *An empty type of streams of integers:* After reading [5], one could be tempted to define the type of streams of integers as $\sigma_0 = \nu\zeta(\iota \otimes \zeta)$. The variable set $\llbracket \iota \otimes \zeta \rrbracket_{\zeta}^{\mathbf{Rel}} : \mathbf{Rel} \rightarrow \mathbf{Rel}$ maps a set E to $\mathbb{N} \times E$. The least fixed point of this operation on sets is \emptyset and hence $|\llbracket \sigma_0 \rrbracket^{\mathbf{Nuts}}| = \emptyset$ and notice that $\text{Tot}(\emptyset) = \{\emptyset, \{\emptyset\}\}$. In that case, the operation $\Phi : \text{Tot}(\emptyset) \rightarrow \text{Tot}(\emptyset)$ maps \mathcal{T} to $\{u \times v \mid v \in \mathcal{T} \text{ and } u \in \mathcal{P}(\mathbb{N}) \setminus \{\emptyset\}\}$ and hence $\{\emptyset\}$ to itself. It follows that $\mathcal{T}(\llbracket \sigma_0 \rrbracket^{\mathbf{Nuts}}) = \{\emptyset\}$ that is $\llbracket \sigma_0 \rrbracket^{\mathbf{Nuts}} = \top$, the final object of \mathbf{Nuts} . What is the meaning of this trivial interpretation? It simply reflects that, though σ_0 has a lot of non trivial proofs in μLL , it is impossible to extract any finite information from these proofs within μLL , and accordingly all these proofs are interpreted as \emptyset .

Theorem 32. *In μLL , there is no proof of $\vdash \sigma_0^{\perp}, \iota$.*

In other words there is no proof of $\vdash \sigma_0 \multimap \iota$ in μLL ; typically a function extracting the first element of a stream would be a proof of this type... if it would exist! Here is the argument: if π were a proof of $\vdash \sigma_0^{\perp}, \iota$, we would have $\llbracket \pi \rrbracket \in \mathbf{Nuts}(\llbracket \sigma_0 \rrbracket^{\mathbf{Nuts}}, \llbracket \iota \rrbracket^{\mathbf{Nuts}})$ and hence $\llbracket \pi \rrbracket \cdot \emptyset \in \mathcal{T}(\llbracket \iota \rrbracket^{\mathbf{Nuts}})$ which is not the case since $\llbracket \pi \rrbracket \cdot \emptyset = \emptyset$. If types like σ_0 are meaningful in a proof-search perspective, their relevance as data-types in a Curry-Howard approach to μLL is dubious.

4) *A non-empty type of streams of integers:* We set now $\sigma = \nu\zeta(1 \& (\iota \otimes \zeta))$. This type looks like the previous one, but the type 1 leaves space for *partial* empty streams. Warning: it is *not* a type of finite or infinite streams; the $\&$ means that this empty stream will not be a total element: it will have

to be complemented by some total element from the right argument of the $\&$. More precisely $\llbracket 1 \& (\mathbb{N} \otimes \zeta) \rrbracket_{\zeta}^{\mathbf{Rel}} : \mathbf{Rel} \rightarrow \mathbf{Rel}$ is the variable set which maps a set E to $\{(1, *)\} \cup \{2\} \times \mathbb{N} \times E$ so that up to renaming $|\llbracket \sigma \rrbracket^{\mathbf{Nuts}}| = \mathbb{N}^{<\omega}$ (all *finite sequences* of integers). In this case, the operator $\Phi : \text{Tot}(\mathbb{N}^{<\omega}) \rightarrow \text{Tot}(\mathbb{N}^{<\omega})$ maps \mathcal{T} to

$$\{v \subseteq \mathbb{N}^{<\omega} \mid () \in v \text{ and } \exists n \in \mathbb{N}, u \in \mathcal{T} \{n\} \times u \subseteq v\}$$

where we use $()$ for the empty sequence. So for instance

$$\begin{aligned} \Phi^0(\mathcal{P}(\mathbb{N}^{<\omega})) &= \mathcal{P}(\mathbb{N}^{<\omega}) & \Phi^1(\mathcal{P}(\mathbb{N}^{<\omega})) &= \{u \in \mathcal{P}(\mathbb{N}^{<\omega}) \mid () \in u\} \\ \Phi^3(\mathcal{P}(\mathbb{N}^{<\omega})) &= \{u \in \mathcal{P}(\mathbb{N}^{<\omega}) \mid \exists n_1, n_2 () , (n_1), (n_1, n_2) \in u\} . \end{aligned}$$

The greatest fixed point is reached in ω steps:

$$\begin{aligned} \text{Tot}(\llbracket \sigma \rrbracket^{\mathbf{Nuts}}) &= \bigcap_{n < \omega} \Phi^n(\mathcal{P}(\mathbb{N}^{<\omega})) \\ &= \{u \subseteq \mathbb{N}^{<\omega} \mid \exists f \in \mathbb{N}^{\omega} \forall k < \omega (f(1), \dots, f(k)) \in u\} . \end{aligned}$$

So a total subset of $|\llbracket \sigma \rrbracket^{\mathbf{Nuts}}|$ must contain (at least) an infinite stream of integer. For this type of streams σ it is easy to build a proof of $\vdash \sigma^{\perp}, \iota$ extracting the first element of a stream, interpreted as $\{(n), n \mid n \in \mathbb{N}\}$.

V. CONCLUSION AND FURTHER WORK

We will study next the semantics of infinite proofs of μLL (whose definition extends that of [5] for μMALL). A crucial step is to prove that these infinite proofs can be interpreted as total sets in \mathbf{Nuts} , this will be presented in a further paper. This interpretation of proofs is based on the interpretation of their finite approximations in \mathbf{Rel} (remember that the interpretations of a μLL proof in \mathbf{Nuts} and in \mathbf{Rel} are *exactly the same set*).

Our models will also serve as guidelines for the design of a functional language based on μLL , generalizing Gödel’s System T in the spirit of [21] though, as explained in the Introduction, Loader’s syntax is not fully compatible with LL as it is based on *cocartesian* cartesian closed categories. Our system will primarily implement Park’s rule, but we will also consider other options based on polymorphism in the spirit of [36], [18] or [37], or general recursion with guardedness restrictions as in [24], [25], [26].

Its syntax will be based on the idea of representing data-types as *positive* formulas of μLL interpreted in $\mathcal{L}^!$ and therefore equipped with morphisms implementing weakening, contraction and promotion: as noticed in [4], $\mu\zeta_$ is a positive operation whereas $\nu\zeta_$ is negative. In $\mathcal{L}^!$, the \oplus of LL is a coproduct and the \otimes is a cartesian product as expected (and \otimes distributes over \oplus). The targeted calculus will feature a notion of values (positive terms) accounting for the morphisms of $\mathcal{L}^!$, substitution in terms being allowed only for values because only them can safely be discarded and duplicated. Thanks to this choice of design, weakening and contraction will remain implicit operations as in the usual λ -calculus. Our calculus will have explicit promotion and dereliction operations, allowing to implement both CBN and CBV in the same setting, just as in Levy’s Call-by-push-value [38], [39].

REFERENCES

- [1] J.-Y. Girard, "Linear logic," *Theoretical Computer Science*, vol. 50, pp. 1–102, 1987.
- [2] J.-Y. Girard, Y. Lafont, and P. Taylor, *Proofs and types*, ser. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 1989, vol. 7.
- [3] J.-Y. Girard, "Fixpoint theorem in linear logic," 1992, an email posting to the mailing list linear@cs.stanford.edu.
- [4] D. Baelde, "Least and Greatest Fixed Points in Linear Logic," *ACM Trans. Comput. Log.*, vol. 13, no. 1, pp. 2:1–2:44, 2012. [Online]. Available: <https://doi.org/10.1145/2071368.2071370>
- [5] D. Baelde, A. Doumane, and A. Saurin, "Infinitary Proof Theory: the Multiplicative Additive Case," in *25th EACSL Annual Conference on Computer Science Logic, CSL 2016, August 29 - September 1, 2016, Marseille, France*, ser. LIPIcs, J. Talbot and L. Regnier, Eds., vol. 62. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016, pp. 42:1–42:17. [Online]. Available: <https://doi.org/10.4230/LIPIcs.CSL.2016.42>
- [6] P.-A. Melliès, "Categorical semantics of linear logic," *Panoramas et Synthèses*, vol. 27, 2009.
- [7] D. de Carvalho and L. Tortora de Falco, "The relational model is injective for multiplicative exponential linear logic (without weakenings)," *Annals of Pure and Applied Logic*, vol. 163, no. 9, pp. 1210–1236, 2012. [Online]. Available: <https://doi.org/10.1016/j.apal.2012.01.004>
- [8] D. de Carvalho, "The Relational Model Is Injective for Multiplicative Exponential Linear Logic," in *25th EACSL Annual Conference on Computer Science Logic, CSL 2016, August 29 - September 1, 2016, Marseille, France*, ser. LIPIcs, J. Talbot and L. Regnier, Eds., vol. 62. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016, pp. 41:1–41:19. [Online]. Available: <https://doi.org/10.4230/LIPIcs.CSL.2016.41>
- [9] M. Hyland and A. Schalk, "Glueing and orthogonality for models of linear logic," *Theoretical Computer Science*, vol. 294, no. 1/2, pp. 183–231, 2003. [Online]. Available: [https://doi.org/10.1016/S0304-3975\(01\)00241-9](https://doi.org/10.1016/S0304-3975(01)00241-9)
- [10] J.-Y. Girard, "The system F of variable types, fifteen years later," *Theoretical Computer Science*, vol. 45, pp. 159–192, 1986.
- [11] R. Loader, "Linear logic, totality and full completeness," in *Proceedings of the 9th Annual IEEE Symposium on Logic in Computer Science*. IEEE Computer Society, 1994.
- [12] A. Bucciarelli and T. Ehrhard, "On phase semantics and denotational semantics: the exponentials," *Annals of Pure and Applied Logic*, vol. 109, no. 3, pp. 205–241, 2001.
- [13] P. Boudes, "Non-uniform (hyper/multi)coherence spaces," *Mathematical Structures in Computer Science*, vol. 21, no. 1, pp. 1–40, 2011. [Online]. Available: <https://doi.org/10.1017/S0960129510000320>
- [14] A. Doumane, "On the infinitary proof theory of logics with fixed points," Ph.D. dissertation, Université de Paris, 2017.
- [15] J. Brotherston and A. Simpson, "Sequent calculi for induction and infinite descent," *Journal of Logic and Computation*, vol. 21, no. 6, pp. 1177–1216, 2011. [Online]. Available: <https://doi.org/10.1093/logcom/exq052>
- [16] A. Tiu and A. Momigliano, "Cut elimination for a logic with induction and co-induction," *J. Appl. Log.*, vol. 10, no. 4, pp. 330–367, 2012. [Online]. Available: <https://doi.org/10.1016/j.jal.2012.07.007>
- [17] R. McDowell and D. Miller, "Cut-elimination for a logic with definitions and induction," *Theoretical Computer Science*, vol. 232, no. 1-2, pp. 91–119, 2000. [Online]. Available: [https://doi.org/10.1016/S0304-3975\(99\)00171-1](https://doi.org/10.1016/S0304-3975(99)00171-1)
- [18] M. D. Campos and M. Fiore, "Classical logic with Mendler induction," *Journal of Logic and Computation*, vol. 30, no. 1, pp. 77–106, 2020. [Online]. Available: <https://doi.org/10.1093/logcom/exaa004>
- [19] L. Santocanale, "A Calculus of Circular Proofs and Its Categorical Semantics," in *Foundations of Software Science and Computation Structures, 5th International Conference, FOSSACS 2002. Held as Part of the Joint European Conferences on Theory and Practice of Software, ETAPS 2002 Grenoble, France, April 8-12, 2002, Proceedings*, ser. Lecture Notes in Computer Science, M. Nielsen and U. Engberg, Eds., vol. 2303. Springer-Verlag, 2002, pp. 357–371. [Online]. Available: https://doi.org/10.1007/3-540-45931-6_25
- [20] J. Fortier and L. Santocanale, "Cuts for circular proofs: semantics and cut-elimination," in *Computer Science Logic 2013 (CSL 2013), CSL 2013, September 2-5, 2013, Torino, Italy*, ser. LIPIcs, S. R. D. Rocca, Ed., vol. 23. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2013, pp. 248–262. [Online]. Available: <https://doi.org/10.4230/LIPIcs.CSL.2013.248>
- [21] R. Loader, "Equational Theories for Inductive Types," *Annals of Pure and Applied Logic*, vol. 84, no. 2, pp. 175–217, 1997. [Online]. Available: [https://doi.org/10.1016/S0168-0072\(96\)00021-8](https://doi.org/10.1016/S0168-0072(96)00021-8)
- [22] H. Geuvers, "Inductive and Coinductive types with Iteration and Recursion," Tech. Rep., 1992, proceedings of the 1992 Workshop on Types for Proofs and Programs, Basta.
- [23] A. Bucciarelli, T. Ehrhard, and G. Manzonetto, "A relational semantics of parallelism and non-determinism in a functional setting," *Annals of Pure and Applied Logic*, vol. 163, no. 7, pp. 918–934, 2012.
- [24] T. Coquand, "Infinite Objects in Type Theory," in *Types for Proofs and Programs, International Workshop TYPES'93, Nijmegen, The Netherlands, May 24-28, 1993, Selected Papers*, ser. Lecture Notes in Computer Science, H. Barendregt and T. Nipkow, Eds., vol. 806. Springer-Verlag, 1993, pp. 62–78. [Online]. Available: https://doi.org/10.1007/3-540-58085-9_72
- [25] C. Paulin-Mohring, "Inductive Definitions in the system Coq - Rules and Properties," in *Typed Lambda Calculi and Applications, International Conference on Typed Lambda Calculi and Applications, TLCA '93, Utrecht, The Netherlands, March 16-18, 1993, Proceedings*, ser. Lecture Notes in Computer Science, M. Bezem and J. F. Groote, Eds., vol. 664. Springer-Verlag, 1993, pp. 328–345. [Online]. Available: <https://doi.org/10.1007/BFb0037116>
- [26] E. Giménez, "Structural Recursive Definitions in Type Theory," in *Automata, Languages and Programming, 25th International Colloquium, ICALP'98, Aalborg, Denmark, July 13-17, 1998, Proceedings*, ser. Lecture Notes in Computer Science, K. G. Larsen, S. Skyum, and G. Winskel, Eds., vol. 1443. Springer-Verlag, 1998, pp. 397–408. [Online]. Available: <https://doi.org/10.1007/BFb0055070>
- [27] P. Clairambault, "Least and greatest fixpoints in game semantics," in *Foundations of Software Science and Computational Structures, 12th International Conference, FOSSACS 2009, Held as Part of the Joint European Conferences on Theory and Practice of Software, ETAPS 2009, York, UK, March 22-29, 2009, Proceedings*, ser. Lecture Notes in Computer Science, L. de Alfaró, Ed., vol. 5504. Springer, 2009, pp. 16–31. [Online]. Available: https://doi.org/10.1007/978-3-642-00596-1_1
- [28] —, "Strong functors and interleaving fixpoints in game semantics," *rtia*, vol. 47, no. 1, pp. 25–68, 2013.
- [29] J.-Y. Girard, "On denotational completeness," *Theoretical Computer Science*, vol. 227, pp. 249–273, 1999.
- [30] P. Curien, M. P. Fiore, and G. Munch-Maccagnoni, "A theory of effects and resources: adjunction models and polarised calculi," in *Proceedings of the 43rd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages, POPL 2016, St. Petersburg, FL, USA, January 20 - 22, 2016*, R. Bodík and R. Majumdar, Eds. ACM, 2016, pp. 44–56. [Online]. Available: <https://doi.org/10.1145/2837614.2837652>
- [31] T. Uustalu and V. Vene, "Comonadic Notions of Computation," in *Proceedings of the Ninth Workshop on Coalgebraic Methods in Computer Science, CMCS 2008, Budapest, Hungary, April 4-6, 2008*, ser. Electronic Notes in Theoretical Computer Science, J. Adámek and C. Kupke, Eds., vol. 203, no. 5. Elsevier, 2008, pp. 263–284. [Online]. Available: <https://doi.org/10.1016/j.entcs.2008.05.029>
- [32] J. Power and E. Robinson, "Premonoidal Categories and Notions of Computation," *Mathematical Structures in Computer Science*, vol. 7, no. 5, pp. 453–468, 1997. [Online]. Available: <https://doi.org/10.1017/S0960129597002375>
- [33] E. Moggi, "Computational lambda-calculus and monads," in *Proceedings of the 4th Annual IEEE Symposium on Logic in Computer Science*. IEEE Computer Society, 1989.
- [34] R. Hasegawa, "Two applications of analytic functors," *Theoretical Computer Science*, vol. 272, no. 1-2, pp. 113–175, 2002.
- [35] T. Ehrhard, "An introduction to differential linear logic: proof-nets, models and antiderivatives," *Mathematical Structures in Computer Science*, vol. 28, no. 7, pp. 995–1060, 2018.
- [36] N. P. Mendler, "Inductive Types and Type Constraints in the Second-Order lambda Calculus," *Annals of Pure and Applied Logic*, vol. 51, no. 1-2, pp. 159–172, 1991. [Online]. Available: [https://doi.org/10.1016/0168-0072\(91\)90069-X](https://doi.org/10.1016/0168-0072(91)90069-X)
- [37] R. Matthes, "Monotone Fixed-Point Types and Strong Normalization," in *Computer Science Logic, 12th International Workshop, CSL '98, Annual Conference of the EACSL, Brno, Czech Republic, August 24-28, 1998, Proceedings*, ser. Lecture Notes in Computer Science, G. Gottlob,

E. Grandjean, and K. Seyr, Eds., vol. 1584. Springer, 1998, pp. 298–312. [Online]. Available: https://doi.org/10.1007/10703163_20

- [38] P. B. Levy, “Call-by-push-value: Decomposing call-by-value and call-by-name,” *Higher-Order and Symbolic Computation*, vol. 19, no. 4, pp. 377–414, 2006. [Online]. Available: <http://dx.doi.org/10.1007/s10990-006-0480-6>
- [39] T. Ehrhard, “Call-By-Push-Value from a Linear Logic Point of View,” in *Programming Languages and Systems - 25th European Symposium on Programming, ESOP 2016, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2016, Eindhoven, The Netherlands, April 2-8, 2016, Proceedings*, ser. Lecture Notes in Computer Science, P. Thiemann, Ed., vol. 9632. Springer-Verlag, 2016, pp. 202–228. [Online]. Available: <https://doi.org/10.1007/978-3-662-49498-1>

VI. APPENDIX

A. Proof of Lemma 4

Proof: We have $\mathcal{F}(g, \nu\mathcal{F}(B)) \in \mathcal{A}(\nu\mathcal{F}(B), \mathcal{F}(B', \nu\mathcal{F}(B)))$ thus defining an $\mathcal{F}_{B'}$ -coalgebra structure on $\nu\mathcal{F}(B)$ and hence there exists a unique morphism $\nu\mathcal{F}(g)$ such that

$$\mathcal{F}(B', \nu\mathcal{F}(g)) \mathcal{F}(g, \nu\mathcal{F}(B)) = \nu\mathcal{F}(g),$$

that is $\mathcal{F}(g, \nu\mathcal{F}(g)) = \nu\mathcal{F}(g)$.

Functoriality follows: consider also $g' \in \mathcal{B}(B', B'')$, then we know that $h = \nu\mathcal{F}(g' g)$ satisfies $\mathcal{F}(g' g, h) = h$ by the definition above. Now $h' = \nu\mathcal{F}(g') \nu\mathcal{F}(g)$ satisfies the same equation by functoriality of \mathcal{F} and because $\mathcal{F}(g, \nu\mathcal{F}(g)) = \nu\mathcal{F}(g)$ and $\mathcal{F}(g', \nu\mathcal{F}(g')) = \nu\mathcal{F}(g')$, and hence $h' = h$ by Lemma 3, taking $l = \mathcal{F}(g' g, \nu\mathcal{F}(B))$. In the same way one proves that $\nu\mathcal{F}(\text{Id}) = \text{Id}$. ■

B. Proof of Lemma 5

Proof: The part of the statement which concerns the functor $\widehat{\nu\mathbb{F}}$ is a direct application of Lemma 4 so we only have to deal with the strength. Let us prove naturality so let $\vec{f} \in \mathcal{L}^n(\vec{X}, \vec{X}')$ and $g \in \mathcal{L}(Y, Y')$, we must prove that the following diagram commutes

$$\begin{array}{ccc} !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X}) & \xrightarrow{\widehat{\nu\mathbb{F}}_{Y, \vec{X}}} & \widehat{\nu\mathbb{F}}(!Y \otimes \vec{X}) \\ !g \otimes \widehat{\nu\mathbb{F}}(\vec{f}) \downarrow & & \downarrow \widehat{\nu\mathbb{F}}(!g \otimes \vec{f}) \\ !Y' \otimes \widehat{\nu\mathbb{F}}(\vec{X}') & \xrightarrow{\widehat{\nu\mathbb{F}}_{Y', \vec{X}'}} & \widehat{\nu\mathbb{F}}(!Y' \otimes \vec{X}') \end{array}$$

Let $h_1 = \widehat{\nu\mathbb{F}}_{Y, \vec{X}}(!g \otimes \widehat{\nu\mathbb{F}}(\vec{f}))$ and $h_2 = \widehat{\nu\mathbb{F}}(!g \otimes \vec{f}) \widehat{\nu\mathbb{F}}_{Y, \vec{X}}$ be the two morphisms we must prove equal. We use Lemma 3, taking the following morphism l .

$$\begin{array}{c} !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X}) = !Y \otimes \widehat{\mathbb{F}}(\vec{X}, \widehat{\nu\mathbb{F}}(\vec{X})) \\ \downarrow \widehat{\mathbb{F}}_{Y, (\vec{X}, \widehat{\nu\mathbb{F}}(\vec{X}))} \\ \widehat{\mathbb{F}}(!Y \otimes \vec{X}, !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X})) \\ \downarrow \widehat{\mathbb{F}}(!g \otimes \vec{f}, \text{Id}) \\ \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X})) \end{array}$$

With these notations we have

$$\begin{aligned} & \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', h_1) l \\ &= \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', \widehat{\nu\mathbb{F}}_{Y', \vec{X}'} \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', !g \otimes \widehat{\nu\mathbb{F}}(\vec{f}))) \\ & \quad \widehat{\mathbb{F}}(!g \otimes \vec{f}, !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X})) \widehat{\mathbb{F}}_{Y, (\vec{X}, \widehat{\nu\mathbb{F}}(\vec{X}))} \\ &= \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', \widehat{\nu\mathbb{F}}_{Y', \vec{X}'} \widehat{\mathbb{F}}(!g \otimes \vec{f}, !g \otimes \widehat{\nu\mathbb{F}}(\vec{f}))) \widehat{\mathbb{F}}_{Y, (\vec{X}, \widehat{\nu\mathbb{F}}(\vec{X}))} \\ &= \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', \widehat{\nu\mathbb{F}}_{Y', \vec{X}'} \widehat{\mathbb{F}}_{Y', (\vec{X}', \widehat{\nu\mathbb{F}}(\vec{X}'))} (!g \otimes \widehat{\mathbb{F}}(\vec{f}, \widehat{\nu\mathbb{F}}(\vec{f})))) \\ & \quad \text{by naturality of } \widehat{\mathbb{F}} \\ &= \widehat{\nu\mathbb{F}}_{Y', \vec{X}'} (!g \otimes \widehat{\mathbb{F}}(\vec{f}, \widehat{\nu\mathbb{F}}(\vec{f}))) \text{ by Lemma 5} \\ &= \widehat{\nu\mathbb{F}}_{Y', \vec{X}'} (!g \otimes \widehat{\nu\mathbb{F}}(\vec{f})) \text{ by Lemma 4} \end{aligned}$$

so that $\widehat{\mathbb{F}}(!Y' \otimes \vec{X}', h_1) l = h_1$ as required. On the other hand we have

$$\begin{aligned} & \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', h_2) l \\ &= \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', \widehat{\nu\mathbb{F}}(!g \otimes \vec{f})) \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', \widehat{\nu\mathbb{F}}_{Y', \vec{X}'} \widehat{\mathbb{F}}_{Y, (\vec{X}, \widehat{\nu\mathbb{F}}(\vec{X}))} \\ & \quad \widehat{\mathbb{F}}(!g \otimes \vec{f}, !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X}))) \\ &= \widehat{\mathbb{F}}(!Y' \otimes \vec{X}', \widehat{\nu\mathbb{F}}(!g \otimes \vec{f})) \widehat{\mathbb{F}}(!g \otimes \vec{f}, !Y \otimes \widehat{\nu\mathbb{F}}(\vec{X})) \\ & \quad \widehat{\mathbb{F}}(!Y \otimes \vec{X}, \widehat{\nu\mathbb{F}}_{Y, \vec{X}} \widehat{\mathbb{F}}_{Y, (\vec{X}, \widehat{\nu\mathbb{F}}(\vec{X}))} \\ &= \widehat{\mathbb{F}}(!g \otimes \vec{f}, \widehat{\nu\mathbb{F}}(!g \otimes \vec{f})) \widehat{\nu\mathbb{F}}_{Y, \vec{X}} \text{ by Lemma 5} \\ &= \widehat{\nu\mathbb{F}}(!g \otimes \vec{f}) \widehat{\nu\mathbb{F}}_{Y, \vec{X}} \text{ by Lemma 4} \end{aligned}$$

so that $\widehat{\mathbb{F}}(!Y' \otimes \vec{X}', h_2) l = h_2$ which proves our contention. The commutation of the diagrams of Figure 1 for $\widehat{\nu\mathbb{F}}$ is proven similarly. ■

C. Proof of Lemma 9

Proof: Let $a \in E$, since $(a, a) \in \text{Id}_E = ts$, there must exist $b \in F$ such that $(a, b) \in s$ and $(b, a) \in t$. If $(a, b') \in s$ then $(b, b') \in st \subseteq \text{Id}_F$ and hence $b' = b$. It follows that s is a total function $E \rightarrow F$. Let $(a, b) \in s$ (that is $a \in E$ and $b = s(a)$). Since $ts = \text{Id}_E$, we must have $(b, a) \in t$. Conversely let $(b, a) \in t$, we have $(b, s(a)) \in st$ and hence $b = s(a)$. We have proven that $t = \{(s(a), a) \mid a \in E\}$. If $a, a' \in E$ satisfy $s(a) = s(a')$ we have therefore $(a, a') \in ts = \text{Id}_E$ and hence $a = a'$; this shows that s is injective. ■

D. Proof of Lemma 14

Proof: Let (E, t) be an \mathbb{F} -coalgebra. We define a sequence $e_n \in \mathbf{Rel}(E, \sigma\mathbb{F})$ as follows: $e_0 = \emptyset$ and $e_{n+1} = \mathbb{F}(e_n) t$. Then $e_n \subseteq e_{n+1}$ for all n by an easy induction, using the fact that \mathbb{F} is hom-continuous. Let $e = \bigcup_{n=0}^{\infty} e_n \in \mathbf{Rel}(E, \sigma\mathbb{F})$, by hom-continuity of \mathbb{F} we have $\mathbb{F}(e) t = (\bigcup_{n=0}^{\infty} \mathbb{F}(e_n)) t = \bigcup_{n=0}^{\infty} (\mathbb{F}(e_n) t) = \bigcup_{n=0}^{\infty} e_{n+1} = e$ which means that

$$e \in \mathbf{Coalg}_{\mathbf{Rel}}(\mathbb{F})((E, t), (\sigma\mathbb{F}, \text{Id})).$$

We end the proof by showing that e is the unique such morphism, so let $e' \in \mathbf{Coalg}_{\mathbf{Rel}}(\mathbb{F})((E, t), (\sigma\mathbb{F}, \text{Id}))$ which means that $e' \in \mathbf{Rel}(E, \sigma\mathbb{F})$ and $\mathbb{F}(e') t = e'$.

Let $i_n \in \mathbf{Rel}(\sigma\mathbb{F}, \sigma\mathbb{F})$ be defined by induction by $i_0 = \emptyset$ and $i_{n+1} = \mathbb{F}(i_n)$. Then $(i_n)_{n \in \mathbb{N}}$ is monotone and $\bigcup_{n=0}^{\infty} i_n = \text{Id}$ by definition of $\sigma\mathbb{F}$. We prove by induction on n that $\forall n \in \mathbb{N} \ i_n e' = i_n e$. Clearly $i_0 e' = i_0 e = \emptyset$. Next

$$\begin{aligned} i_{n+1} e' &= \mathbb{F}(i_n) \mathbb{F}(e') t \\ &= \mathbb{F}(i_n e') t \\ &= \mathbb{F}(i_n e) t \quad \text{by inductive hypothesis} \\ &= i_{n+1} e. \end{aligned}$$

Therefore $e' = (\bigcup_{n \in \mathbb{N}} i_n) e' = \bigcup_{n \in \mathbb{N}} (i_n e') = \bigcup_{n \in \mathbb{N}} (i_n e) = e$. \blacksquare

E. Proof of Lemma 15

Proof: As usual we assume that $n = 1$ to increase readability. We need to prove first that $\nu\mathbb{F}$ is monotone on morphisms, so let $s, t \in \mathbf{Rel}(E, F)$ with $s \subseteq t$. We have $\nu\mathbb{F}(s) = \bigcup_{n \in \mathbb{N}} s_n$ and $\nu\mathbb{F}(t) = \bigcup_{n \in \mathbb{N}} t_n$ with $s_0 = t_0 = \emptyset$, $s_{n+1} = \mathbb{F}(s, s_n)$ and $t_{n+1} = \mathbb{F}(t, t_n)$ (we use the action of $\nu\mathbb{F}$ on morphisms resulting from Lemma 4 and from the characterization of the morphisms to the final object given in the proof of Lemma 14). By induction and hom-monotonicity of \mathbb{F} we have $\forall n \in \mathbb{N} \ s_n \subseteq t_n$ and hence $\nu\mathbb{F}(s) \subseteq \nu\mathbb{F}(t)$. Let us prove now hom-continuity so let $D \subseteq \mathbf{Rel}(E, F)$ be directed and let $t = \bigcup D$, we prove that $\nu\mathbb{F}(t) = \bigcup_{s \in D} \nu\mathbb{F}(s) \in \mathbf{Rel}(\nu\mathbb{F}(E), \nu\mathbb{F}(F))$ using Lemma 3 (with the notations of that lemma, we take $l = \mathbb{F}(t, \nu\mathbb{F}(E))$). We have

$$\mathbb{F}_F(\nu\mathbb{F}(t)) \mathbb{F}(t, \nu\mathbb{F}(E)) = \nu\mathbb{F}(t)$$

by definition of the functor $\nu\mathbb{F}$ and

$$\begin{aligned} &\mathbb{F}_F\left(\bigcup_{s \in D} \nu\mathbb{F}(s)\right) \mathbb{F}(t, \nu\mathbb{F}(E)) \\ &= \bigcup_{s \in D} \mathbb{F}(F, \nu\mathbb{F}(s)) \bigcup_{s \in D} \mathbb{F}(s, \nu\mathbb{F}(E)) \quad \text{by hom-cont.} \\ &= \bigcup_{s \in D} \mathbb{F}(s, \nu\mathbb{F}(s)) = \bigcup_{s \in D} \nu\mathbb{F}(s). \end{aligned}$$

In the second equation, we used the facts that D is directed and the monotonicity of \mathbb{F} and $\nu\mathbb{F}$ on morphisms.

Let $E \subseteq F$, we prove that $\nu\mathbb{F}(E) \subseteq \nu\mathbb{F}(F)$. This results from the observation that if $E' \subseteq F'$, then $\mathbb{F}_E(E') \subseteq \mathbb{F}_F(F')$ and hence $\forall n \in \mathbb{N} \ \mathbb{F}_E^n(\emptyset) \subseteq \mathbb{F}_F^n(\emptyset)$. Let us check that $\nu\mathbb{F}(\eta_{E,F}^+) = \eta_{\nu\mathbb{F}(E), \nu\mathbb{F}(F)}^+ \in \mathbf{Rel}(\nu\mathbb{F}(E), \nu\mathbb{F}(F))$. We have

$$\begin{aligned} &\mathbb{F}(F, \nu\mathbb{F}(\eta_{E,F}^+)) \mathbb{F}(\eta_{E,F}^+, \nu\mathbb{F}(E)) \\ &= \mathbb{F}(\eta_{E,F}^+, \nu\mathbb{F}(\eta_{E,F}^+)) = \nu\mathbb{F}(\eta_{E,F}^+) \end{aligned}$$

by definition of the functor $\nu\mathbb{F}$ and

$$\begin{aligned} &\mathbb{F}(F, \eta_{\nu\mathbb{F}(E), \nu\mathbb{F}(F)}^+) \mathbb{F}(\eta_{E,F}^+, \nu\mathbb{F}(E)) = \eta_{\mathbb{F}(E, \nu\mathbb{F}(E)), \mathbb{F}(F, \nu\mathbb{F}(F))}^+ \\ &= \eta_{\nu\mathbb{F}(E), \nu\mathbb{F}(F)}^+ \end{aligned}$$

by strictness of \mathbb{F} . The equation follows by Lemma 3, so that the functor $\nu\mathbb{F}$ is strict. \blacksquare

F. Proof of Lemma 19

Proof: Let $t \in \mathcal{T}(X \multimap Y)$ and let $u \in \mathcal{T}(X)$. Let $v' \in \mathcal{T}(Y)^\perp$, since $u \times v' \in \mathcal{T}(X \otimes Y^\perp)$ we have $t \cap (u \times v') \neq \emptyset$ and hence $(t \cdot u) \cap v' \neq \emptyset$. Therefore $t \cdot u \in \mathcal{T}(Y)^{\perp\perp} = \mathcal{T}(Y)$. Conversely assume that $\forall u \in \mathcal{T}(X) \ t \cdot u \in \mathcal{T}(Y)$. Let $u \in \mathcal{T}(X)$ and $v' \in \mathcal{T}(Y^\perp) = \mathcal{T}(Y)^\perp$. Since $t \cdot u \in \mathcal{T}(Y)$ we have $(t \cdot u) \cap v' \neq \emptyset$ and hence $t \cap (u \times v') \neq \emptyset$ and this shows that $t \in \mathcal{T}(X \multimap Y)$. \blacksquare

G. Proof of Lemma 20

Proof: Assume that t is an iso in \mathbf{Nuts} so that there is $t' \in \mathbf{Nuts}(Y, X)$ such that $t't = \text{Id}_{|X|}$ and $tt' = \text{Id}_{|Y|}$ and since we know that the isos in \mathbf{Rel} are the bijections we know that t is a bijection. The fact that $\forall u \subseteq |X| \ u \in \mathcal{T}(X) \Leftrightarrow t(u) \in \mathcal{T}(Y)$ results from the fact that both t and $t' = t^{-1}$ are morphisms in \mathbf{Nuts} .

The converse implication is obvious. \blacksquare

H. Proof of Lemma 22

Proof: The condition is obviously necessary, let us prove that it is sufficient so assume that t fulfills it and let us prove that $t \in \mathcal{T}(X_1 \otimes X_2 \multimap Y)$. To this end it suffices to prove that $t^\perp \in \mathcal{T}(Y^\perp \multimap (X_1 \otimes X_2)^\perp)$. So let $v' \in \mathcal{T}(Y^\perp)$ and let us prove that $t^\perp \cdot v' \in \mathcal{T}((X_1 \otimes X_2)^\perp) = \{u_1 \otimes u_2 \mid u_1 \in \mathcal{T}(X_1) \text{ and } u_2 \in \mathcal{T}(X_2)\}^\perp$. So let $u_i \in \mathcal{T}(X_i)$ for $i = 1, 2$. We know that $t \cdot (u_1 \otimes u_2) \in \mathcal{T}(Y)$ and hence $(t \cdot (u_1 \otimes u_2)) \cap v' \neq \emptyset$, that is $(u_1 \otimes u_2) \cap (t^\perp \cdot v') \neq \emptyset$, proving our contention. \blacksquare

I. Proof of Lemma 23

Proof: Let $t \in \mathcal{T}((X_1 \otimes X_2) \multimap Y)$ and let us prove that $s = \alpha \cdot t \in \mathcal{T}(X_1 \multimap (X_2 \multimap Y))$. Given $u_i \in \mathcal{T}(X_i)$ it suffices to prove that $(t' \cdot u_1) \cdot u_2 \in \mathcal{T}(Y)$ which results from the fact that $(s \cdot u_1) \cdot u_2 = t \cdot (u_1 \otimes u_2)$. Conversely let $s \in \mathcal{T}(X_1 \multimap (X_2 \multimap Y))$ and let us prove that $t = \alpha^{-1} \cdot s \in \mathcal{T}((X_1 \otimes X_2) \multimap Y)$. This results from lemma 22 and from the equation $(s \cdot u_1) \cdot u_2 = t \cdot (u_1 \otimes u_2)$. \blacksquare

J. Proof of Lemma 24

Proof: The condition is obviously necessary, so let us assume that it holds. By Lemma 21, it suffices to prove that $t^\perp \in \mathbf{Nuts}(Y^\perp, (!X)^\perp)$. Let $v' \in \mathcal{T}(Y^\perp)$, we prove that $t^\perp \cdot v' \in \mathcal{T}(!Y)^\perp$. So let $u \in \mathcal{T}(X)$, since $t \cdot u^{(!)} \in \mathcal{T}(Y)$ and hence $(t \cdot u^{(!)}) \cap v' \neq \emptyset$, that is $(t^\perp \cdot v') \cap u^{(!)} \neq \emptyset$. \blacksquare

K. Proof of Lemma 25

Proof: We deal with the case $k = 2$. The condition is necessary since, if $u_1 \in \mathcal{T}(X_1)$ and $u_2 \in \mathcal{T}(X_2)$, then $u_1^{(!)} \otimes u_2^{(!)} \in \mathcal{T}(!X_1 \otimes !X_2)$. So assume that it holds. Let $t' = \text{cur}(t) \in \mathbf{Rel}(|X_1| \multimap (|X_2| \multimap |Y|))$. Let $u_1 \in \mathcal{T}(X_1)$, we have $t' \cdot u_1^{(!)} \in \mathcal{P}(!X_2 \multimap Y)$. Let $u_2 \in \mathcal{T}(X_2)$, we have $(t' \cdot u_1^{(!)}) \cdot u_2^{(!)} = t \cdot (u_1^{(!)} \otimes u_2^{(!)}) \in \mathcal{T}(Y)$ by our assumption. It follows by Lemma 24 that $t' \cdot u_1^{(!)} \in \mathcal{T}(!X_2 \multimap Y)$ and since this holds for any $u_1 \in \mathcal{T}(X_1)$ we actually have $t' \in \mathbf{Nuts}(!X_1, !X_2 \multimap Y)$. It follows that $t = \text{cur}^{-1}(t') \in \mathbf{Nuts}(!X_1 \otimes !X_2, Y)$ as contended. \blacksquare

L. Proof of Lemma 27

Proof: Given an object X of \mathbf{Nuts} , we set $\text{der}_X = \text{der}_{|X|} \in \mathbf{Rel}(|X|, |X|)$ and $\text{dig}_X = \text{dig}_{|X|} \in \mathbf{Rel}(|X|, |!X|)$. Given $u \in \mathcal{T}(X)$, we have $\text{der}_X \cdot u^{(!)} = u \in \mathcal{T}(X)$ and $\text{dig}_X \cdot u^{(!)} = u^{(!)} \in \mathcal{T}(!X)$. It follows by Lemma 24 that $\text{der}_X \in \mathbf{Nuts}(!X, X)$ and $\text{dig}_X \in \mathbf{Nuts}(!X, !X)$.

Naturality and monadicity trivially hold because they hold in \mathbf{Rel} : we have an obvious faithful forgetful functor $\mathbf{Nuts} \rightarrow \mathbf{Rel}$ which commutes with all LL categorical constructs.

We are left with defining the strong monoidality structure of $!_-$ (Seely isomorphisms), for $m^0 \in \mathbf{Nuts}(1, !\top)$ we take the same morphism as in \mathbf{Rel} . And we set $m_{X_1, X_2}^2 = m_{|X_1|, |X_2|}^2 \in \mathbf{Rel}(!X_1 \otimes !X_2, !(X_1 \& X_2))$. Let $u_i \in \mathcal{T}(X_i)$ for $i = 1, 2$. We have $m_{X_1, X_2}^2 \cdot (u_1^{(!)} \otimes u_2^{(!)}) = (u_1 \& u_2)^{(!)} \in \mathcal{T}!(X_1 \& X_2)$ since $u_1 \& u_2 \in \mathcal{T}(X_1 \& X_2)$, and hence by Lemma 25 we have $m_{X_1, X_2}^2 \in \mathbf{Nuts}(!X_1 \otimes !X_2, !(X_1 \& X_2))$. Any element w of $\mathcal{T}(X_1 \& X_2)$ is of shape $w = u_1 \& u_2$ with $u_i \in \mathcal{T}(X_i)$, namely $u_i = \pi_i \cdot w$. We have $(m_{X_1, X_2}^2)^{-1} \cdot w^{(!)} = u_1^{(!)} \otimes u_2^{(!)} \in \mathcal{T}(!X_1 \otimes !X_2)$ and hence by Lemma 24 we have $(m_{X_1, X_2}^2)^{-1} \in \mathbf{Nuts}!(X_1 \& X_2, !X_1 \otimes !X_2)$. This ends the proof that \mathbf{Nuts} is a model of classical Linear Logic since the required commutations obviously hold because they hold in \mathbf{Rel} . ■

M. Full proof of Theorem 30

Proof: Concerning Condition (3), let $(\mathbb{X}_i)_{i=1}^k$ be elements of \mathbf{Vnuts}_n and let $\mathbb{X} \in \mathbf{Vnuts}_k$. Considering \mathbb{X} and the \mathbb{X}_i 's as strong functors, we know that $\mathbb{X} \circ \overline{\mathbb{X}}$ is a strong functor $\mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$. We simply have to exhibit a VNUTS whose associated strong functor is $\mathbb{X} \circ \overline{\mathbb{X}}$. Let $\mathbb{F} = |\mathbb{X}| \circ |\overline{\mathbb{X}}|$ (composition of variable sets, Section III-C2). Let $\vec{X} \in \mathbf{Nuts}^n$, each $\overline{\mathbb{X}_i}(\vec{X})$ is an object of \mathbf{Nuts} and hence $(\overline{\mathbb{F}}(|\vec{X}|), \mathcal{T}(\mathbb{X})(\overline{\mathbb{X}_1}(\vec{X}), \dots, \overline{\mathbb{X}_k}(\vec{X})))$ is a NUTS. Moreover given $\vec{t} \in \mathbf{Nuts}^n(\vec{X}, \vec{Y})$, we know that for each $i = 1, \dots, k$, one has $\overline{\mathbb{X}_i}(\vec{t}) \in \mathbf{Nuts}(\overline{\mathbb{X}_i}(\vec{X}), \overline{\mathbb{X}_i}(\vec{Y}))$ since \mathbb{X}_i is a VNUTS. Since \mathbb{X} is a VNUTS we have

$$\begin{aligned} & \overline{\mathbb{F}}(\vec{t}) \\ & \in \mathbf{Nuts}(\overline{\mathbb{X}}(\overline{\mathbb{X}_1}(\vec{X}), \dots, \overline{\mathbb{X}_k}(\vec{X})), \overline{\mathbb{X}}(\overline{\mathbb{X}_1}(\vec{Y}), \dots, \overline{\mathbb{X}_k}(\vec{Y}))). \end{aligned}$$

Let $X \in \text{Obj}(\mathbf{Nuts})$ and $\vec{Y} \in \text{Obj}(\mathbf{Nuts}^k)$. For $i = 1, \dots, k$ we know that $\overline{\mathbb{X}_i} \in \mathbf{Nuts}(!X \otimes \overline{\mathbb{X}_i}(\vec{Y}), \overline{\mathbb{X}_i}(!X \otimes \vec{Y}))$. Therefore

$$\begin{aligned} & \overline{\mathbb{X}}((\overline{\mathbb{X}_i} \otimes \overline{\mathbb{Y}})_{i=1}^k) \\ & \in \mathbf{Nuts}(\overline{\mathbb{X}}((!X \otimes \overline{\mathbb{X}_i}(\vec{Y}))_{i=1}^k), \overline{\mathbb{X}}((\overline{\mathbb{X}_i}(!X \otimes \vec{Y}))_{i=1}^k)) \end{aligned}$$

and hence

$$\begin{aligned} & \overline{\mathbb{X}}((\overline{\mathbb{X}_i} \otimes \overline{\mathbb{Y}})_{i=1}^k) \widehat{\mathbb{X}}_{X, (\overline{\mathbb{X}_i}(\vec{Y}))_{i=1}^k} \\ & \in \mathbf{Nuts}(!X \otimes \overline{\mathbb{X}}((\overline{\mathbb{X}_i}(\vec{Y}))_{i=1}^k), \overline{\mathbb{X}}((\overline{\mathbb{X}_i}(!X \otimes \vec{Y}))_{i=1}^k)). \end{aligned}$$

Moreover we have

$$\begin{aligned} \widehat{\mathbb{F}}_{|X|, |\vec{Y}|} &= \overline{\mathbb{X}}(|\overline{\mathbb{X}}|(|\overline{\mathbb{X}_i}|_{|X|, |\vec{Y}|})_{i=1}^k) \widehat{\mathbb{X}}_{|X|, (|\overline{\mathbb{X}_i}(\vec{Y})|)_{i=1}^k} \\ & \text{by definition of } \overline{\mathbb{F}} \\ &= \overline{\mathbb{X}}(|\overline{\mathbb{X}}|(|\overline{\mathbb{X}_i}|_{|X|, |\vec{Y}|})_{i=1}^k) \widehat{\mathbb{X}}_{|X|, (|\overline{\mathbb{X}_i}(\vec{Y})|)_{i=1}^k} \\ &= \overline{\mathbb{X}}((\widehat{\mathbb{X}_i}_{X, \vec{Y}})_{i=1}^k) \widehat{\mathbb{X}}_{X, (\overline{\mathbb{X}_i}(\vec{Y}))_{i=1}^k} \end{aligned}$$

using again the fact that \mathbb{X} and the \mathbb{X}_i 's are VNUTS. This shows that the pair $\mathbb{Y} = (|\mathbb{Y}|, \mathcal{T}(\mathbb{Y}))$ given by $|\mathbb{Y}| = \mathbb{F}$ and $\mathcal{T}(\mathbb{Y})(\vec{X}) = \mathcal{T}(\mathbb{X})(\overline{\mathbb{X}_1}(\vec{X}), \dots, \overline{\mathbb{X}_k}(\vec{X}))$ is a VNUTS whose associated strong functor is $\mathbb{X} \circ \overline{\mathbb{X}}$ thus proving our contention.

Concerning Condition (4), let us deal only with the case of $!_-$, the others being similar. We have to exhibit a unary VNUTS \mathbb{X} whose associated strong functor $\mathbf{Nuts} \rightarrow \mathbf{Nuts}$ coincides with $!_-$ (which is known to be a strong functor $\mathbf{Nuts} \rightarrow \mathbf{Nuts}$ by Section IV-A3 and by the general considerations of Section II-D1). For $|\mathbb{X}|$, which has to be a variable set $\mathbf{Rel} \rightarrow \mathbf{Rel}$, we take the interpretation \mathbb{E} of $!_-$ in the model \mathbf{Rel} (Section III-C2) which is an element of \mathbf{Rel}_1 , that is, a unary variable set. Next, given $X \in \text{Obj}(\mathbf{Nuts})$, we take $\mathcal{T}(\mathbb{X})(X) = \mathcal{T}(!X)$. Condition (1) in the definition of VNUTS holds by functoriality of $!_-$ on \mathbf{Nuts} . Condition (2) holds by the definition of $\widehat{\mathbb{F}}_{|X|, |Y|}$ as described in Section II-D1 which coincides with $\mu^2(\text{dig}_X \otimes !Y) \in \mathbf{Nuts}(!X \otimes !Y, !(X \otimes Y))$.

Let us now turn to Condition (5) which is a bit more challenging.

1) *Fixed Points of VNUTS:* Let first $\mathbb{X} = (|\mathbb{X}|, \mathcal{T}(\mathbb{X}))$ be a unary VNUTS. Let $E = \sigma|\overline{\mathbb{X}}|$ which is the least set such that $|\overline{\mathbb{X}}|(E) = E$, that is $E = \bigcup_{n=0}^{\infty} |\overline{\mathbb{X}}|^n(\emptyset)$. Let $\Phi : \text{Tot}(E) \rightarrow \text{Tot}(E)$ be defined as follows: given $S \in \text{Tot}(E)$, then (E, S) is a NUTS, and we set $\Phi(S) = \mathcal{T}(\mathbb{X})(E, S) \in \text{Tot}(|\overline{\mathbb{X}}|(E)) = \text{Tot}(E)$. This function Φ is monotone. Let indeed $S_1, S_2 \in \text{Tot}(E)$ with $S_1 \subseteq S_2$. Then we have $\text{ld} \in \mathbf{Nuts}((E, S_1), (E, S_2))$ and therefore, by Condition (1) satisfied by \mathbb{X} , we have

$$\begin{aligned} \text{ld} &= |\overline{\mathbb{X}}|(\text{ld}) \in \mathbf{Nuts}(\overline{\mathbb{X}}(E, S_1), \overline{\mathbb{X}}(E, S_2)) \\ &= \mathbf{Nuts}((E, \Phi(S_1)), (E, \Phi(S_2))) \end{aligned}$$

which means that $\Phi(S_1) \subseteq \Phi(S_2)$. By the Knaster Tarski Theorem (remember that $\text{Tot}(E)$ is a complete lattice), Φ has a greatest fixpoint \mathcal{T} that we can describe as follows. Let $(\mathcal{T}_\alpha)_{\alpha \in \mathbb{O}}$, where \mathbb{O} is the class of ordinals, be defined by: $\mathcal{T}_0 = \mathcal{P}(E)$ (the largest possible notion of totality on E), $\mathcal{T}_{\alpha+1} = \Phi(\mathcal{T}_\alpha)$ and $\mathcal{T}_\lambda = \bigcap_{\alpha < \lambda} \mathcal{T}_\alpha$ when λ is a limit ordinal. This sequence is decreasing (easy induction on ordinals using the monotonicity of Φ) and there is an ordinal θ such that $\mathcal{T}_{\theta+1} = \mathcal{T}_\theta$ (by a cardinality argument; we can assume that θ is the least such ordinal). The greatest fixpoint of Φ is then \mathcal{T}_θ as easily checked.

By construction $((E, \mathcal{T}_\theta), \text{ld})$ is an object of $\mathbf{Coalg}_{\mathbf{Nuts}}(\overline{\mathbb{X}})$, we prove that it is the final object. So let (Y, t) be another object of the same category. Since $(|Y|, t)$ is an object of $\mathbf{Coalg}_{\mathbf{Rel}}(|\overline{\mathbb{X}}|)$ and since (E, ld) is

the final object in that category, we know by Lemma 14 that there is exactly one $e \in \mathbf{Rel}(|Y|, E)$ such that $\overline{X}(e)t = e$. We prove that actually $e \in \mathbf{Nuts}(Y, (E, \mathcal{T}_\theta))$ so let $v \in \mathcal{T}(Y)$. We prove by induction on the ordinal α that $e \cdot v \in \mathcal{T}_\alpha$. For $\alpha = 0$ it is obvious since $\mathcal{T}_0 = \mathcal{P}(E)$. Assume that the property holds for α and let us prove it for $\alpha + 1$. We have $t \cdot v \in \mathcal{T}(\overline{X})(Y) = \mathcal{T}(\overline{X}(Y))$ since $t \in \mathbf{Nuts}(Y, \overline{X}(Y))$. Since $\overline{X}(e) \in \mathbf{Nuts}(\overline{X}(Y), \overline{X}(E, \mathcal{T}_\alpha))$ and since $\overline{X}(E, \mathcal{T}_\alpha) = (E, \mathcal{T}_{\alpha+1})$ we have $(\overline{X}(e)t) \cdot v \in \mathcal{T}_{\alpha+1}$, that is $e \cdot v \in \mathcal{T}_{\alpha+1}$. Last if λ is a limit ordinal and if we assume $\forall \alpha < \lambda \ e \cdot v \in \mathcal{T}_\alpha$ we have $e \cdot v \in \bigcap_{\alpha < \lambda} \mathcal{T}_\alpha = \mathcal{T}_\lambda$. Therefore $e \cdot v \in \mathcal{T}_\theta$. We use $\nu\overline{X}$ to denote this final coalgebra (E, \mathcal{T}_θ) (its definition depends only on \overline{X} and does not involve the strength \widehat{X}).

So we have proven the first part of Condition (5) in the definition of a Seely model of μLL (see Section 7). As to the second part, let \overline{X} be an $n + 1$ -ary VNUTS. We know by the general Lemma 5 that there is a uniquely defined strong functor $\nu\overline{X} : \mathbf{Nuts}^n \rightarrow \mathbf{Nuts}$ such that

- $\nu\overline{X}(\overrightarrow{X}) = \nu(\overline{X}_{\overrightarrow{X}})$, so that $\overline{X}(\overrightarrow{X}, \nu\overline{X}(\overrightarrow{X})) = \overline{X}(\overrightarrow{X})$, for all $\overrightarrow{X} \in \text{Obj}(\mathbf{Nuts}^n)$,
- $\overline{X}(\overrightarrow{t}, \nu\overline{X}(\overrightarrow{t})) = \nu\overline{X}(\overrightarrow{t})$ for all $\overrightarrow{t} \in \mathbf{Nuts}(\overrightarrow{X}, \overrightarrow{Y})$
- and $\overline{X}(Y \otimes \overrightarrow{X}, \widehat{\nu\overline{X}}_{Y, \overrightarrow{X}}) \widehat{X}_{Y, (\overrightarrow{X}, \nu\overline{X}(\overrightarrow{X}))} = \widehat{\nu\overline{X}}_{Y, \overrightarrow{X}}$ for all $Y \in \text{Obj}(\mathbf{Nuts})$ and $\overrightarrow{X} \in \text{Obj}(\mathbf{Nuts}^n)$.

To end the proof, it will be enough to exhibit an n -ary VNUTS $\mathbb{Y} = (|\mathbb{Y}|, \mathcal{T}(\mathbb{Y}))$ whose associated strong functor coincides with $\nu\overline{X}$. We know that $|\overline{X}|$ is a variable set $\mathbf{Rel}^{n+1} \rightarrow \mathbf{Rel}$ so let $\mathbb{F} = \nu|\overline{X}| = \sigma|\overline{X}|$ which is a variable set $\mathbf{Rel}^n \rightarrow \mathbf{Rel}$ (see Section III-C1). Let $\overrightarrow{X} \in \text{Obj}(\mathbf{Nuts}^n)$, we have $|\nu\overline{X}(\overrightarrow{X})| = |\nu(\overline{X}_{\overrightarrow{X}})| = \bigcup_{n=0}^{\infty} |\overline{X}_{\overrightarrow{X}}|^n(\emptyset) = \mathbb{F}(|\overrightarrow{X}|)$. Let $\overrightarrow{t} \in \mathbf{Nuts}^n(\overrightarrow{X}, \overrightarrow{Y})$, then $\nu\overline{X}(\overrightarrow{t})$ is the unique element s of $\mathbf{Nuts}(\nu\overline{X}(\overrightarrow{X}), \nu\overline{X}(\overrightarrow{Y}))$ (this hom-set is a subset of $\mathbf{Rel}(\mathbb{F}(|\overrightarrow{X}|), \mathbb{F}(|\overrightarrow{Y}|))$) which satisfies $\overline{X}(\overrightarrow{t}, s) = s$, that is $|\overline{X}|(\overrightarrow{t}, s) = s$. This means that $\nu\overline{X}(\overrightarrow{t}) = s = \mathbb{F}(\overrightarrow{t})$. By a completely similar uniqueness argument we have $\widehat{\nu\overline{X}}_{X, \overrightarrow{Y}} = \widehat{\mathbb{F}}_{|X|, |\overrightarrow{Y}|}$ for all $X \in \text{Obj}(\mathbf{Nuts})$ and $\overrightarrow{Y} \in \text{Obj}(\mathbf{Nuts}^n)$. So we set $|\mathbb{Y}| = \mathbb{F}$.

Next, given $\overrightarrow{X} \in \text{Obj}(\mathbf{Nuts}^n)$ we set

$$\begin{aligned} \mathcal{T}(\mathbb{Y})(\overrightarrow{X}) &= \mathcal{T}(\nu\overline{X}(\overrightarrow{X})) \\ &\in \text{Tot}(|\nu\overline{X}(\overrightarrow{X})|) = \text{Tot}(\mathbb{F}(|\overrightarrow{X}|)) \end{aligned}$$

Given $\overrightarrow{t} \in \mathbf{Nuts}(\overrightarrow{X}, \overrightarrow{Y})$ we have

$$\begin{aligned} \mathbb{F}(\overrightarrow{t}) &= \nu\overline{X}(\overrightarrow{t}) \\ &\in \mathbf{Nuts}((\mathbb{F}(|\overrightarrow{X}|), \mathcal{T}(\mathbb{Y})(\overrightarrow{X})), (\mathbb{F}(|\overrightarrow{Y}|), \mathcal{T}(\mathbb{Y})(\overrightarrow{Y}))) \end{aligned}$$

since $(\mathbb{F}(|\overrightarrow{X}|), \mathcal{T}(\mathbb{Y})(\overrightarrow{X})) = \nu\overline{X}(\overrightarrow{X})$ and similarly for \overrightarrow{Y} . Last since $\widehat{\mathbb{F}}_{|X|, |\overrightarrow{Y}|} = \widehat{\nu\overline{X}}_{X, \overrightarrow{Y}} \in \mathbf{Nuts}(!X \otimes \nu\overline{X}(\overrightarrow{Y}), \nu\overline{X}(X \otimes \overrightarrow{Y}))$ we know that $\mathbb{Y} = (|\mathbb{Y}|, \mathcal{T}(\mathbb{Y}))$ is a VNUTS whose associated strong functor is $\nu\overline{X}$. This ends the proof that $(\mathbf{Nuts}, (\mathbf{Vnuts}_n)_{n \in \mathbb{N}})$ is a Seely model of μLL . \blacksquare