

THEMED ISSUE: CANNABINOIDS

REVIEW

CB₂: a cannabinoid receptor with an identity crisis

Brady K Atwood^{1,2} and Ken Mackie¹

¹The Gill Center and the Department of Psychological & Brain Sciences, Indiana University, Bloomington, IN, USA, and

²University of Washington, Program of Neurobiology and Behavior, Seattle, WA, USA

CB₂ was first considered to be the 'peripheral cannabinoid receptor'. This title was bestowed based on its abundant expression in the immune system and presumed absence from the central nervous system. However, multiple recent reports question the absence of CB₂ from the central nervous system. For example, it is now well accepted that CB₂ is expressed in brain microglia during neuroinflammation. However, the extent of CB₂ expression in neurons has remained controversial. There have been studies claiming either extreme-its complete absence to its widespread expression-as well as everything in between. This review will discuss the reported tissue distribution of CB₂ with a focus on CB₂ in neurons, particularly those in the central nervous system as well as the implications of that presence. As CB₂ is an attractive therapeutic target for pain management and immune system modulation without overt psychoactivity, defining the extent of its presence in neurons will have a significant impact on drug discovery. Our recommendation is to encourage cautious interpretation of data that have been presented for and against CB₂'s presence in neurons and to encourage continued rigorous study.

British Journal of Pharmacology (2010) **160**, 467–479; doi:10.1111/j.1476-5381.2010.00729.x

This article is part of a themed issue on Cannabinoids. To view the editorial for this themed issue visit <http://dx.doi.org/10.1111/j.1476-5381.2010.00831.x>

Keywords: GPCR tissue distribution; microglia; sensory neurons; pain; immunocytochemistry

Abbreviations: 2-AG, 2-arachidonylglycerol; CB_{1/2}, type1/2 cannabinoid receptor; DMNX, dorsal motor nucleus of the vagus; DRG, dorsal root ganglion; FACS, fluorescence activated cell sorting; GFAP, glial fibrillary acidic protein; ICC, immunocytochemistry; IHC, immunohistochemistry; IPSC, inhibitory post synaptic current; ISH, in situ hybridization; MAPK, mitogen-activated protein kinases; NB, northern blot; PGE₂, prostaglandin E₂; PI3K, phosphoinositide 3-kinase; RLB, radioligand binding; RT-PCR, reverse transcriptase polymerase chain reaction; SB, Southern blot; THC, Δ⁹-tetrahydrocannabinol; TRPV1, transient receptor potential cation channel V1; WB, western blot

The therapeutic potential of cannabis as well as its psychoactive effects has been known for thousands of years (Adams and Martin, 1996). However, it was not until the discovery of cannabinoid binding sites in brain (Devane *et al.*, 1988; Herkenham *et al.*, 1990; Herkenham *et al.*, 1991; Matsuda *et al.*, 1993) and the subsequent cloning of the CB₁ receptor (Matsuda *et al.*, 1990; Alexander *et al.*, 2008) that cellular mechanisms for these effects began to be elucidated. A second cannabinoid receptor (CB₂) was identified and first cloned from HL60 cells by Munro *et al.* in 1993 (Munro *et al.*, 1993). CB₂ was dubbed the 'peripheral cannabinoid receptor' as a

result of *in situ* hybridization analysis that revealed high levels of CB₂ mRNA in spleen and its absence from brain. CB₂ receptors were cloned from mouse and rat in later years (Shire *et al.*, 1996; Griffin *et al.*, 2000; Brown *et al.*, 2002).

CB₂ and cellular signalling

Around the time that CB₁ and CB₂ were cloned, anandamide and 2-arachidonylglycerol (2-AG) were identified as endogenous cannabinoid agonists (Devane *et al.*, 1992; Felder *et al.*, 1993; Sugiura *et al.*, 1995; Hanus *et al.*, 2001). 2-AG is a high efficacy agonist at CB₂ (Lynn and Herkenham, 1994; Slipetz *et al.*, 1995; Gonsiorek *et al.*, 2000; Sugiura *et al.*, 2000; Shoemaker *et al.*, 2005b). However, anandamide has a low efficacy at CB₂, often functioning as a weak partial agonist (Showalter *et al.*, 1996; Gonsiorek *et al.*, 2000; Sugiura *et al.*, 2000).

Correspondence: Ken Mackie, Department of Psychological and Brain Sciences, Indiana University, 1101 E. 10th Street, Bloomington, IN 47405, USA. E-mail: kmackie@indiana.edu

Received 21 December 2009; revised 8 February 2010; accepted 16 February 2010

Similar to CB₁, CB₂ is a G_{i/o} coupled G protein coupled receptor and as such inhibits adenylyl cyclase (Bayewitch *et al.*, 1995; Felder *et al.*, 1995; Slipetz *et al.*, 1995; Gonsiorek *et al.*, 2000; Sugiura *et al.*, 2000; Shoemaker *et al.*, 2005b). Furthermore, it can also promote MAPK activation (p38 and p42/44), PI3K, ceramide production and gene transcription (Bouaboula *et al.*, 1996; Bouaboula *et al.*, 1999a; Bouaboula *et al.*, 1999b; Howlett, 2002; Howlett *et al.*, 2002; Herrera *et al.*, 2005; Herrera *et al.*, 2006; Grimaldi *et al.*, 2009; Romero-Sandoval *et al.*, 2009). A key difference, however, is that unlike CB₁, CB₂ appears to poorly modulate calcium channels or inwardly rectifying potassium channels (Felder *et al.*, 1995). Studies using SR144528, a CB₂ antagonist/inverse agonist, have revealed that the receptor possesses a high degree of constitutive activity in expression systems (Bouaboula *et al.*, 1999a; Bouaboula *et al.*, 1999b). Of further interest is that CB₂ receptors from different species often have different pharmacological responses to identical drugs, complicating the drug discovery process (Mukherjee *et al.*, 2004; Yao *et al.*, 2006; Bingham *et al.*, 2007). Therefore, despite coupling to the same family of G proteins and sharing some ligands, CB₁ and CB₂ appear to differ significantly from one another in their signalling.

CB₂ as a therapeutic target

CB₂ is an attractive therapeutic target. The abundant CB₂ expression in immune cells presents a plausible explanation for cannabinoid immunomodulatory activity (Lynn and Herkenham, 1994; Berdyshev, 2000; Howlett, 2002; Costa, 2007). Indeed, CB₂ activation affects a myriad of immune responses from inflammation to neuroprotection (Cabral and Griffin-Thomas, 2009). Additionally, numerous reports indicate that CB₂ activation is analgesic and CB₂ agonists suppress responses in many animal models of pain, from acute to neuropathic (Anand *et al.*, 2009), although these effects may involve CB₁ activation as well. CB₁ is abundant within the brain, where it appears responsible for mediating the psychoactive effects of cannabis (Mackie, 2005). Thus, the scarcity of central nervous system (CNS) CB₂ receptors makes CB₂ selective drugs attractive as therapeutics as they would presumably lack psychoactivity. In support of this notion, mice with 'knockout' of CB₂ had typical behavioural responses to Δ^9 -tetrahydrocannabinol (THC) but lost their normal immune responsiveness to THC (Buckley *et al.*, 2000). CB₂ levels can also be increased under certain conditions and disease states further adding to its attractiveness as a therapeutic target (Zhang *et al.*, 2003; Wotherspoon *et al.*, 2005; Yiangou *et al.*, 2006).

CB₂ agonists and antagonists

As CB₂ is such an attractive therapeutic target, much effort has been made to synthesize selective CB₂ agonists and antagonists. Some of the early cannabinoid agonists such as CP55940, WIN55212-2 and HU210 demonstrate high affinity at CB₂ and are considered full agonists, but are not selective for CB₂ over CB₁ [see Miller and Stella (2008) for a summary of

the binding data]. JWH015 was one of the first potential CB₂ selective agonists (Showalter *et al.*, 1996; Griffin *et al.*, 2000), but has since been shown to also be an agonist at GPR55 (Ryberg *et al.*, 2007; Lauckner *et al.*, 2008). Numerous other compounds have been synthesized with the aim of making CB₂ selective agonists. AM1241 (Malan *et al.*, 2001) and JWH133 (Huffman *et al.*, 1999) are two of the most commonly used 'selective' CB₂ agonists. Other early ones include HU308 (Hanus *et al.*, 1999) and GW405833 (L-768242) (Gallant *et al.*, 1996; Valenzano *et al.*, 2005). More recently synthesized compounds include GW833972A (Belvisi *et al.*, 2008), MDA7 (Naguib *et al.*, 2008), A-796260 (Yao *et al.*, 2008) and A-836339 (Yao *et al.*, 2009). Cannabinalones have also been suggested as potential CB₂ selective compounds (Khanolkar *et al.*, 2007). For the interested reader, a review by Whiteside *et al.* contains detailed analysis of many of these compounds as well as numerous others (Whiteside *et al.*, 2007). SR144528 (Rinaldi-Carmona *et al.*, 1998) and AM630 (Pertwee *et al.*, 1995; Ross *et al.*, 1999) are the two of the most commonly used CB₂ selective antagonists and have been frequently used to demonstrate specificity of many of these other CB₂ selective agonists. However, a fundamental problem with designing selective agonists and antagonists is possible interactions with other unforeseen targets. These compounds may exhibit a strong preference for CB₂ over CB₁, but as evidenced by JWH015, other non-CB₁/CB₂ binding sites may still exist. It is extremely difficult to conclusively establish an agonist (or antagonist) is specific for CB₂ and no other receptors. This caveat must be kept in mind when evaluating studies that solely use a pharmacological approach and allege CB₂ involvement in a process. Furthermore, as demonstrated for AM1241, CB₂ agonists may produce very different effects at CB₂ receptors from different species (Bingham *et al.*, 2007). Here, a racemic mixture of AM1241 was an agonist at human CB₂ but functioned as an inverse agonist at rodent CB₂. R-AM1241 has a higher affinity for CB₂ than S-AM1241, but functions similar to the racemate. On the other hand S-AM1241 was an efficacious agonist at both human and rodent CB₂. Furthermore naloxone, a μ opioid receptor antagonist, can block the analgesic effects of AM1241, but this appears not to be the case for other CB₂ agonists (Ibrahim *et al.*, 2005; Yao *et al.*, 2009). It also important to bear in mind that selective agonists and antagonists for a particular receptor may differentially alter coupling to distinct signalling pathways, a concept known as functional selectivity (Urban *et al.*, 2007). Thus, CB₂ agonists that share identical binding characteristics may have different potencies in activating different signalling pathways and evoke substantially different physiological responses. For example, with CB₂ expressed in Chinese hamster ovary cells, 2-AG, CP55940 and noladin ether had different rank orders of potency depending on the signalling pathway analysed: inhibition of adenylyl cyclase, MAPK activation and stimulation of calcium transients (Shoemaker *et al.*, 2005a). This concept of functional selectivity mandates caution in comparing pharmacological studies that employ different agonists and antagonists for CB₂. This may explain the differences previously mentioned between AM1241 and other CB₂ agonists and their ability to produce analgesia and may even extend to other agonist-specific effects obtained with other compounds. Thus, it is important

that functional evidence obtained using CB₂ 'selective' agonists and antagonists be balanced with careful controls and supported by additional genetic and anatomical analyses to assure that some other unanticipated target is not the true site of action.

CB₂: the 'peripheral' cannabinoid receptor?

CB₂ and evidence for absence from CNS

In addition to the data obtained during the CB₂ knockout mouse characterization, earlier reports also arrived at the conclusion that CB₂ is absent from the CNS. (Of course, 'absence' just means below the level of detection of the particular assay being employed.) When CB₂ was first cloned, *in situ* hybridization demonstrated a lack of CB₂ mRNA signal in rat brain (Munro *et al.*, 1993). While characterizing CB₁ and CB₂ in immune cells, Schatz *et al.* performed Northern blots on mouse brain and rat cerebellums and could not detect the presence of CB₂ in these tissues (Schatz *et al.*, 1997). However, RT-PCR demonstrated the presence of CB₂ mRNA, at levels too low to be quantified. Northern blot analysis performed by Galiegue *et al.* is in agreement with the Schatz *et al.* data (Galiegue *et al.*, 1995). However, in RT-PCR experiments performed as a part of this study, CB₂ was undetectable in human cortex, cerebellum, whole brain and pituitary gland. McCoy *et al.* also did not detect CB₂ mRNA in mouse brain using RT-PCR/Southern blot analysis (McCoy *et al.*, 1999). As part of the characterization of SR144528 as a CB₂ antagonist, rat brain radioligand binding and GTP γ S binding analyses were performed (Griffin *et al.*, 1999). The authors found little SR144528 binding in whole brain and cerebellum and the results of their GTP γ S binding analysis supported this. Furthermore, Northern blotting did not detect CB₂ mRNA in cerebellum, cortex or spinal cord. Rat cortex has also been reported to lack CB₂ mRNA (Beltramo *et al.*, 2006). Included in the review by Howlett *et al.* (Howlett *et al.* 2002), Herkenham and Hohmann replicated the *in situ* hybridization results of Munro and colleagues. Derbenev and colleagues did not detect CB₂ mRNA or protein in rat brainstem (Derbenev *et al.*, 2004). As part of an initial characterization of cannabinoid receptors in dorsal root ganglia (DRG), *in situ* hybridization revealed the presence of CB₁ but not CB₂ receptors (Hohmann and Herkenham, 1999a,b). Price *et al.* could also not detect CB₂ mRNA in rat trigeminal ganglia (Price *et al.*, 2003).

Based on all these data, CB₂ was informally referred to as the peripheral cannabinoid receptor. However, a number of more recent reports have suggested that, in contrast to these previous claims of its absence, CB₂ may in fact be expressed in the CNS (see below). This finding has had a significant impact on drug discovery and our understanding of the biology of the endocannabinoid system. This review will focus on reports of CB₂ in neurons and in the brain and the implications of that presence.

CB₂ and the immune system

CB₂ research continues to have a large focus on its role in the immune system. Analysis of the presence and function of CB₂ in the brain necessitates a discussion concerning CB₂ in

immune cells. CB₂ mRNA has been identified in many immune tissues (Munro *et al.*, 1993; Lynn and Herkenham, 1994; Galiegue *et al.*, 1995; Schatz *et al.*, 1997; Berdyshev, 2000; Buckley *et al.*, 2000). Of specific immune cell types, the highest levels of CB₂ are found in macrophages, CD4+ T cells, CD8+ T cells, B cells, natural killer cells, monocytes and polymorphonuclear neutrophils (Derocq *et al.*, 1995; Galiegue *et al.*, 1995; Schatz *et al.*, 1997; Carayon *et al.*, 1998; McCoy *et al.*, 1999; Buckley *et al.*, 2000; Carlisle *et al.*, 2002; Maresz *et al.*, 2007; Dittel, 2008). Of particular relevance for the role of CB₂ in the CNS, CB₂ mRNA and protein have been found in microglia (Carlisle *et al.*, 2002; Klegeris *et al.*, 2003; Walter *et al.*, 2003; Beltramo *et al.*, 2006; Maresz *et al.*, 2007). Microglia are derived from macrophages and can be viewed as the resident immune cells of the brain where they monitor the brain for pathological damage. In response to specific signals within the brain they transition between different states of activity (Ashton and Glass, 2007; Hanisch and Kettenmann, 2007). The expression levels of CB₂ in microglia vary depending on the activation state of the cell (Carlisle *et al.*, 2002; Walter *et al.*, 2003; Stella, 2004; Maresz *et al.*, 2007; Cabral *et al.*, 2008; Pietr *et al.*, 2009). CB₂ modulates microglial migration and infiltration into brain areas with active neuroinflammation and degeneration (Walter *et al.*, 2003; Ashton *et al.*, 2007; Fernandez-Ruiz *et al.*, 2008; Miller and Stella, 2008; Price *et al.*, 2009). In healthy brain microglia do not appear to express CB₂ (Stella, 2004). However, in Alzheimer's brain tissue, CB₂ can be detected in neuritic plaque-associated microglia (Benito *et al.*, 2003). Similarly, in models of neuropathic pain (but not inflammatory pain) CB₂ mRNA increases in association with activated microglia in the spinal cord (Zhang *et al.*, 2003). In addition during amyotrophic lateral sclerosis and multiple sclerosis, CB₂ microglial expression increases in the spinal cord (Yiangou *et al.*, 2006). According to this evidence it is clear that under certain conditions brain microglia are capable of expressing CB₂.

CB₂ and tissue distribution

Despite being initially described as an immune cell cannabinoid receptor, CB₂ has been identified molecularly and pharmacologically in numerous other cell types. Evidence for the presence of CB₂ receptors has been found in pulmonary endothelial cells (Zoratti *et al.*, 2003). In these cells, CB₂ activation by anandamide results in phospholipase C-mediated calcium release from smooth ER with subsequent increases in mitochondrial calcium. CB₂ can also be found in bone (in osteocytes, osteoblasts and osteoclasts) where it modulates bone formation and turnover (Ofek *et al.*, 2006). The gastrointestinal system appears to express CB₂ receptors as well (Storr *et al.*, 2002; Hillsley *et al.*, 2007; Duncan *et al.*, 2008). 2-AG affects meiosis in spermatogonia via CB₂ (Grimaldi *et al.*, 2009) as well as a number of other aspects of reproductive function (Maccarrone, 2008; Grimaldi *et al.*, 2009). Keratinocytes release beta-endorphin in response to CB₂ selective agonist stimulation (Ibrahim *et al.*, 2005), although this result is controversial (Whiteside *et al.*, 2007; Anand *et al.*, 2008; Yao *et al.*, 2008; Yao *et al.*, 2009). These cells have also been reported to have CB₂ immunoreactivity. In the eye, trabecular

meshwork cells have been shown to have functional CB₂ receptors (Zhong *et al.*, 2005; He and Song, 2007). Mature and precursor adipocytes express functional CB₂ receptors that are negatively coupled to adenylyl cyclase (Roche *et al.*, 2006). In cirrhotic liver, CB₂ receptors are expressed in hepatic myofibroblasts, but are absent in normal liver (Julien *et al.*, 2005). THC protects cardiomyocytes from hypoxic damage by acting at CB₂ receptors resulting in nitric oxide production (Shmist *et al.*, 2006).

CB₂ and nociception

To better understand the possible presence of CB₂ in neurons it is helpful to consider the role of CB₂ in nociception. Cannabinoids have long been known to possess analgesic activity, but evidence for CB₂ having a role in analgesia was not presented until 1998 (Calignano *et al.*, 1998). Shortly thereafter, HU308, a CB₂ selective agonist, was shown to have analgesic activity without typical cannabinoid CNS side effects (Hanus *et al.*, 1999). AM1241, another CB₂ selective agonist was also shown to promote analgesia when injected peripherally and this did not produce CNS side effects, suggesting that CB₂ receptors modulate nociception (Malan *et al.*, 2001; Ibrahim *et al.*, 2003; Malan *et al.*, 2003; Ibrahim *et al.*, 2006). It has since been shown that a number of different CB₂ agonists can modulate many types of pain: acute, inflammatory, neuropathic, post-surgical and cancer pain (Khanolkar *et al.*, 2007; Whiteside *et al.*, 2007; Jhaveri *et al.*, 2008; Naguib *et al.*, 2008; Ohta *et al.*, 2008; Yao *et al.*, 2008; Anand *et al.*, 2009; Yao *et al.*, 2009). It is still unclear as to where these CB₂ agonists exert their analgesic activity. The site could be microglia, astrocytes, neurons, another cell type or a combination of these. Furthermore, as discussed above, the specificity of these CB₂ 'selective' compounds may not be as specific as previously thought. Additional work must be performed to state with confidence that these agonists produce analgesia solely via activation of CB₂ receptors.

CB₂ and peripheral neurons

The first step to determine if CB₂ activation has a direct effect on neural mechanisms is to determine whether CB₂ is expressed in neurons. The existence of functional CB₂ receptors in peripheral neurons has been suggested by a number of studies. The first evidence for CB₂ function in peripheral neurons came in 1997 when CB₂ mRNA was identified in mouse vas deferens tissue (Griffin *et al.*, 1997). In support of a functional role for CB₂, JWH015 and JWH051 (agonists preferring CB₂ over CB₁) produced concentration dependent inhibition of evoked contractions presumably via a prejunctional site. However, a submicromolar concentration of AM630, a CB₂ antagonist, could not block this effect. Further, JWH015 is also an agonist for GPR55 (Ryberg *et al.*, 2007; Lauckner *et al.*, 2008), so the involvement of CB₂ cannot be unequivocally asserted.

CB₂ and sensory neurons

Functional studies have hinted at the presence of CB₂ receptors on sensory neurons. In these studies, it is necessary to

consider the possible involvement of CB₂-expressing immune cells as microglia can affect synaptic properties (Cullheim and Thams, 2007; Abbadie *et al.*, 2009). Patel and colleagues provided some of the first functional evidence of CB₂ in sensory neurons (Patel *et al.*, 2003). Using isolated guinea pig and human vagus nerve preparations, they demonstrated that the CB₂ agonist JWH133 inhibited nerve depolarizations in response to capsaicin, PGE₂ and hypertonic saline. These three treatments activate vagal C and/or A δ fibres. SR144528 blocked the effects of JWH133. A follow-up study with another putative CB₂ agonist, GW833972A, produced similar results (Belvisi *et al.*, 2008). Neither study was designed to determine a specific site or mechanism of action. While CB₂ does not appear to play a role in myenteric contractions, it does seem to play a role in activation of mesenteric sensory nerves. AM1241 administered intravenously inhibits bradykinin induced activation of isolated mesenteric afferents in mice (Hillsley *et al.*, 2007). This effect was blocked by AM630 and absent in CB₂ knockout mice. Interestingly, while CB₂ agonists do not affect normal enteric contractility, JWH133 can prevent lipopolysaccharide (LPS) induced increases in evoked contractions (Mathison *et al.*, 2004; Duncan *et al.*, 2008). JWH133 also blocks LPS stimulation of Fos expression in enteric neurons. AM630 antagonizes these effects. CB₂ receptors in myenteric neurons were identified as the most likely target of this drug effect (see below). CB₂ mRNA has also been identified in rat and mouse retina using RT-PCR as well as within specific layers of the retina (ganglion, inner nuclear and photoreceptor inner layers) using *in situ* hybridization (Lu *et al.*, 2000). Additionally, Burdyga *et al.* identified low, barely detectable levels of CB₂ mRNA in rat nodose ganglion, but were unable to detect CB₂ in the human vagal nerve trunk (Burdyga *et al.*, 2004).

CB₂ and nociceptive neurons

Further functional studies point to a role for CB₂ in sensory neuron function, particularly nociceptive neurons. A study was performed to address AM1241's ability to prevent windup of wide dynamic range (WDR) neurons in spinal cord (Nackley *et al.*, 2004). Here, AM1241 administered locally or systemically reduced the activity of C-fibres synapsing onto WDR neurons and this was reversed by SR144528, but not SR141716A. Significantly, suppression occurred in the presence and absence of inflammation. This, combined with the time course observed suggests long-term changes in presynaptic facilitation, makes the effects less likely to be due to CB₂ targeting immune cells. The authors speculate a direct effect of CB₂ activation on C-fibre neurons. Elmes *et al.* performed a similar study using JWH133 as a CB₂ agonist to test WDR spinal neuron responses in models of inflammatory and neuropathic pain as well as in non-inflammatory and sham-operated conditions (Elmes *et al.*, 2004). Like the Nackley study, they also found that peripherally administered CB₂ agonist inhibits WDR activity in both naïve and inflammatory conditions as well as following neuropathic injury. Once again, the data are suggestive of a non-immune function of CB₂, possibly in peripheral neurons. A follow-up study analysed JWH133's ability to inhibit capsaicin-induced calcium increases in DRG neurons cultured from sham-operated and

neuropathic rats (Sagar *et al.*, 2005). JWH133 slightly inhibited calcium increases in DRG cultured from both neuropathic and sham rats in a SR144528-sensitive fashion, consistent with the presence of functional CB₂ receptors in peripheral neurons. However, spinally administered JWH133 inhibited mechanically evoked responses of dorsal horn neurons from laminae V and VI only in neuropathic rats, but not in sham-operated animals. This points to an up-regulation of CB₂ in intrinsic spinal cord neurons in pain states, although does not provide evidence of the site of up-regulation. AM1241 and L768242 (another CB₂ agonist) can also decrease capsaicin-induced calcitonin gene-related peptide release, a pain biomarker, from neurons in spinal cord slices and this can be blocked by SR144528 (Beltramo *et al.*, 2006). These studies are most consistent with CB₂ participating in neural mechanisms rather than via immune cells, but do not directly answer the question of whether or not CB₂ is expressed in neurons.

The initial support for the presence of CB₂ protein in neurons came from Ross and colleagues. Using fluorescence-activated cell sorting analysis, they determined that DRG cultures and F-11 cells (DRG neuron × neuroblastoma hybrid) express both CB₁ and CB₂, but could not conclude that CB₂ was functionally expressed in DRG neurons (Ross *et al.*, 2001). To further address the site of CB₂ expression in DRG, Wotherspoon and colleagues used immunohistochemistry on DRG and spinal cord of naive and nerve damaged rats and mice (Wotherspoon *et al.*, 2005). No CB₂ could be detected in normal rat or mouse spinal cord and DRG neurons. However, upon nerve sectioning or ligation, CB₂ immunoreactivity could be detected in the ipsilateral dorsal horn. This immunoreactivity was strongly reduced in CB₂ knockout mice and was blocked by incubation with the immunizing peptide, suggesting specificity of the primary antibody used. Of great interest, and in contrast to what would be expected based on Zhang *et al.*'s (2003) study, was that the CB₂ signal did not co-localize with markers of astrocytes (GFAP) or microglia (OX-42). Instead it co-localized with markers of damaged sensory neuron terminals (GAP-43 and galanin). CB₂ immunoreactivity also accumulated in axons proximal to the ligation site. They could not identify CB₂ in cell bodies in tissue sections, but were able to identify CB₂ in isolated DRG neurons grown in culture from lesioned mice. They also did not observe CB₂ immunoreactivity in skin, in contrast to other studies (Stander *et al.*, 2005; Kress and Kuner, 2009) and Ibrahim *et al.* who found it in keratinocytes (Ibrahim *et al.*, 2005). A few studies have detected CB₂ mRNA in DRG and spinal cord using quantitative RT-PCR. Here levels increased following nerve ligation, but this does not necessarily implicate a neuronal source (Zhang *et al.*, 2003; Beltramo *et al.*, 2006).

A more recent study by Anand and colleagues is consistent with the above findings (Anand *et al.*, 2008). Specifically, they found CB₂ positive, small diameter neurons in human DRG and peripheral nerves using three different CB₂ antibodies. The specificity of the antibodies was assessed using peptide block. CB₂ levels increased following nerve injury. They further extended the analysis by demonstrating CB₂ colocalization with neuronal (GAP-43), axonal (neurofilament) and nociceptive neuronal markers (TRPV1). Similar staining was

observed in mouse, rat and guinea pig DRG. This study also replicated the functional data reported by Sagar *et al.* in that a CB₂ agonist (GW833972) inhibited capsaicin-induced calcium increases in DRG sensory neurons. They further sought to identify a mechanism for this activity and determined that CB₂-mediated cAMP depletion attenuated TRPV1 activation. This presumably decreased PKA-mediated phosphorylation of TRPV1, analogous to the effects of μ opioid receptor activation.

CB₂ and the enteric nervous system

Despite earlier findings (Griffin *et al.*, 1997), several studies suggest CB₂ is expressed in the enteric nervous system. Duncan *et al.* and Storr *et al.* found CB₂ mRNA in the enteric nervous system (Storr *et al.*, 2002; Duncan *et al.*, 2008). The site of action of JWH133 in preventing LPS-induced increases in ileum contractility was addressed using RT-PCR and immunohistochemistry (Duncan *et al.*, 2008). CB₂ mRNA was detected in the full-wall thickness ileum, ileal muscle, submucosal and mucosal layers in normal rats. LPS treatment had no effect on the levels of expression. A number of different antibodies and knockout tissue were used for controls. CB₂ protein was detected in all the same tissues in which CB₂ mRNA was found. CB₂ colocalized with markers of enteric ganglia, pan-neuronal markers and synaptic terminals suggesting a strong presence in myenteric neurons. CB₂ immunoreactivity did not colocalize with glial markers.

CB₂: another central cannabinoid receptor?

CB₂ in the cerebellum

One of the earliest reports of the presence of CB₂ in the CNS came from a study performed by Skaper and colleagues (Skaper *et al.*, 1996). *In situ* hybridization revealed the presence of CB₂ mRNA in cultured granule cells. In addition *in situ* hybridization localized CB₂ to the granule and Purkinje cell layers of mouse cerebellum. Radioligand binding analysis of cerebellar membranes revealed the presence of two WIN55212 binding sites. The affinities of WIN55212 at these sites were reported to be close to those of CB₁ and CB₂, although the exact identity of the binding sites could not be specifically determined. RT-PCR analysis has identified CB₂ mRNA in the rat cerebellum and Western blotting has revealed expressed CB₂ protein in rat and ferret cerebellum as well (Van Sickle *et al.*, 2005). Peptide block was used as a control for those Western blots.

Additional studies have also attempted to localize CB₂ protein in the cerebellum (Ashton *et al.*, 2006; Baek *et al.*, 2008). Using an antibody directed against the C-terminus of CB₂, with peptide block control, they identified CB₂ protein expression in the granule, Purkinje and white matter layers of the rat cerebellum. The signal did not overlap with astrocytes markers and the staining pattern in the Purkinje layer and parts of the other layers appeared to be capillary endothelial in nature. There were fine fibres in the white matter and granule cell layers that were CB₂ positive but their origin remains to be determined. These could possibly arise from microglia or neurons. Onaivi *et al.* have also reported CB₂

expression in the Purkinje and molecular layers of the cerebellum using Western blot, immunohistochemistry and *in situ* hybridization techniques (Gong *et al.*, 2006; Onaivi *et al.*, 2008b).

CB₂ in the brainstem

CB₂ has been identified within the brainstem as well. A thorough analysis was performed that investigated CB₂ expression in brain, focusing on mRNA, protein and functional expression within the brainstem (Van Sickle *et al.*, 2005). Quantitative RT-PCR showed that the rat brainstem contains CB₂ mRNA at significantly lower levels than spleen (1.5% of spleen levels). Western blotting confirmed this expression for rat as well as for ferret. Immunocytochemistry identified the dorsal motor nucleus of the vagus nerve (DMNX) of the mouse, rat and ferret as a brainstem nucleus containing CB₂ protein. The CB₂ knockout mouse did not show any immunostaining in the DMNX. The DMNX immunoreactivity colocalized with neuronal markers, but in contrast to what Ashton *et al.* found in the cerebellum (Ashton *et al.*, 2006), the signal did not overlap with glial or blood vessel markers. The authors acknowledged the differences in results between their study and that of Derbenev *et al.* that did not find CB₂ in similar regions (Derbenev *et al.*, 2004) and state that in the latter study a faint signal could be observed in Western blots consistent with low levels of expression. The authors also demonstrated that AM630 blocked the anti-emetic actions of 2-AG treatment in ferrets, suggesting CB₂ receptor involvement. Furthermore a sub-eficacious concentration of anandamide combined with AM1241 treatment produced anti-emetic effects. Another, more superficial study of the brainstem using immunohistochemistry was later performed to look for CB₂ in other brainstem nuclei (Baek *et al.*, 2008). CB₂ immunoreactivity was found in the medial vestibular nucleus as well as the dorsal and ventral cochlear nuclei, but no attempts were made to identify cell types. Peptide block and secondary antibody controls were used to determine CB₂ antibody specificity. Viscomi *et al.* did not find CB₂ protein and only low levels of CB₂ mRNA in inferior olive and pontine nuclei using immunohistochemistry and quantitative PCR in normal rats (Viscomi *et al.*, 2009). However, following a hemispherectomy, CB₂ expression dramatically increased in both mRNA and protein levels in these nuclei. The CB₂ immunoreactivity colocalized with neuronal markers but not with microglial or astrocytic ones. Further JWH015 had a neuroprotective effect, preventing cell death due to the hemispherectomy. This was likely operating through CB₂, although they did not report block of the neuroprotective effect with a CB₂ antagonist. Gong *et al.* reported the presence of CB₂ in many nuclei of the brainstem using RT-PCR and immunohistochemistry (Gong *et al.*, 2006).

CB₂ and the hippocampal formation

Using several approaches, Onaivi and his collaborators have reported finding CB₂ immunoreactivity in many areas of the hippocampal formation (Gong *et al.*, 2006; Onaivi, 2006; Onaivi *et al.*, 2006; 2008a,b; Brusco *et al.*, 2008). They report

a predominately postsynaptic expression and an association with rough endoplasmic reticulum and Golgi structures. They have also demonstrated CB₂ staining in hippocampal cultures. In contrast to their immunohistochemical results, they have had mixed results in finding CB₂ mRNA in the hippocampus (Gong *et al.*, 2006; Onaivi *et al.*, 2008b).

Functional evidence for CB₂ expression in the cortex comes from recording spontaneous inhibitory postsynaptic currents (sIPSCs) in layers II and V of the medial entorhinal cortex. Here, 2-AG mediated suppression of sIPSCs was not blocked by LY320135, a CB₁ antagonist/inverse agonist, whereas they were blocked by AM630 and JTE907 (a structurally distinct CB₂ antagonist) (Morgan *et al.*, 2009). Further JWH133 suppressed sIPSCs in a CB₂ antagonist sensitive fashion. The site of CB₂ agonist action remains to be conclusively demonstrated.

CB₂ and other brain regions

Evidence exists for CB₂ expression in other brain regions. While recording from the ventral posterior nucleus of the thalamus, Jhaveri *et al.* found that after spinal nerve ligation, JWH133 reduced spontaneous and evoked responses in a SR144528-sensitive fashion, but that this effect was absent in sham operated rats (Jhaveri *et al.*, 2008). Gong *et al.* have also reported CB₂ immunoreactivity in many thalamic nuclei, but could not detect CB₂ mRNA using RT-PCR (Gong *et al.*, 2006). Furthermore, this group has reported finding CB₂ mRNA in striatum and hypothalamus, but not in olfactory bulb, cortex and spinal cord and mixed results in midbrain (Gong *et al.*, 2006; Onaivi *et al.*, 2008b). Additionally they report CB₂ immunoreactivity in olfactory bulb, cortex, midbrain as well as the other areas already mentioned (Gong *et al.*, 2006; Onaivi, 2006; Onaivi *et al.*, 2006; 2008b).

CB₂ and neurogenesis

CB₂ also appears to play a role in neurogenesis. Both CB₁ and CB₂ are expressed in stem cells (Jiang *et al.*, 2007; Molina-Holgado *et al.*, 2007). More specifically, RT-PCR, Western blot and immunohistochemical analyses have all revealed the presence of CB₂ in embryonic and adult neural progenitor cells (Palazuelos *et al.*, 2006; Molina-Holgado *et al.*, 2007). CB₂ blockade or genetic disruption impairs neurosphere formation and prevents progenitor cell proliferation, whereas CB₂ agonists promote these activities via ERK and Akt signalling (Palazuelos *et al.*, 2006; Molina-Holgado *et al.*, 2007). However, CB₂ expression seems to diminish as the cells differentiate, being nearly absent by the time neuronal and astrocytic markers appear (Palazuelos *et al.*, 2006). Further, CB₁ agonists and antagonists have similar effects in neurosphere formation and in COR-1 neural stem cell cultures (Molina-Holgado *et al.*, 2007; Goncalves *et al.*, 2008) suggesting either functional interactions or redundant signalling. In contrast to these data, CB₁ agonists and antagonists had no effect on neurogenesis in the subventricular zone (SVZ) of either young or adult mice (Goncalves *et al.*, 2008). On the

other hand, JWH133 and WIN55212 stimulated SVZ neurogenesis, whereas AM630 and JTE907 decreased it (Goncalves *et al.*, 2008).

CB₂: where is its real home and why do we care?

We feel careful analysis of the studies reviewed above allows us to reach the following conclusions: CB₂ is expressed by microglia, with levels increasing as they are activated, and CB₂ is present at detectable and functionally relevant levels in a subset of neurons, with increasing levels following injury. We care where CB₂ is expressed primarily for understanding pathology that involves CB₂ and to develop therapies that target difficult to treat conditions. To this end it is important to have a rigorous understanding of where and under what conditions CB₂ is expressed in the CNS.

Approaches aimed at identifying CB₂ receptor expression in the brain can be divided into functional (pharmacological), biochemical and anatomical techniques. All three have their strengths and weaknesses. The most convincing studies will incorporate a combination of these techniques. Table 1 summarizes the studies presented here, detailing the brain region analysed and whether or not CB₂ was detected and the techniques(s) used to detect it. Pharmacological studies rely on the specificity of the drugs used. When interpreting these studies it is necessary to recall that specificity is never absolute – at sufficiently high concentrations any drug will interact with additional targets. Thus, it is important to relate the concentration of the drug being used to the binding affinity of the CB₂ receptor for that drug. The second consideration is that drugs considered to be 'specific' or 'selective' based on our current understanding may soon be found to interact with other receptors. Examples of this in the cannabinoid system include AM251, often used as a 'selective' CB₁ receptor antagonist, but it is also a GPR55 agonist (Henstridge *et al.*, 2009; Kapur *et al.*, 2009) and JWH015, sometimes used as a 'selective' CB₂ agonist, but it, too, is a GPR55 agonist (Ryberg *et al.*, 2007; Lauckner *et al.*, 2008). Approaches to circumvent this issue include using several structurally diverse agonists and antagonists (presumably decreasing the likelihood of having the same 'off-target' actions) and knockout or 'knock-down' controls, when appropriate.

Biochemical studies include Western blotting and PCR-based approaches. For Western blotting, the key limitations are the sensitivity and specificity of the antibody used. At a minimum, blots from knockout (assuming the antibody is recognizing an epitope present in mouse CB₂) and positive control tissues (e.g. spleen) should be shown. Blindly trusting an antibody to 'work' without concurrent controls is unacceptable. Block with the immunizing antibody is desirable, but will not rule out a fortuitous interaction of the antibody with an unintended epitope on another protein. The sensitivity of Western blotting will depend on the abundance of CB₂ as well as the affinity of the antibody. The lack of detection of CB₂ on the blot can only be interpreted as that the level of CB₂ in the brain is below a certain level. (This level, relative to a CB₂-expressing tissue like spleen, can be determined by serial dilution.) PCR-approached tissues are the most sensitive. Their high sensitivity makes their interpreta-

tion subject to several considerations (Suzuki *et al.*, 2000; Lion, 2001). These include amplification of CB₂ mRNA from immune cells trapped in the cerebral vasculature and amplification of CB₂ mRNA from a very small subset of activated microglia. In order to rationally interpret results from PCR-based experiments it is necessary that they be performed in a quantitative fashion, preferably calculating copy number, to facilitate comparisons.

Anatomical studies need to be conducted and interpreted with a similarly critical approach. These studies fall into three categories: autoradiography, *in situ* hybridization and immunocytochemistry. As above the issue of sensitivity needs consideration – it is possible to show CB₂ is present, but it is very hard to conclusively demonstrate that it is not present, just that it is present at a level below the limit of detection. However, this information, coupled with a lack of functional response, can be very valuable in sorting out the role of CB₂ receptors in a particular physiological response. The caveats of autoradiography include the pharmacological considerations discussed above as well as specific technical issues (Frey and Albin, 2001). As this technique has not been widely applied to directly identifying CB₂ receptors in the brain, it will not be further discussed here. *In situ* hybridization studies can yield useful information on which cells express CB₂ and thus can complement PCR-based studies. However, the lower sensitivity of *in situ* hybridization may make this difficult. A necessary control for *in situ* hybridization includes lack of hybridization in knockout tissues (when possible).

Immunocytochemistry studies have the powerful potential to identify the precise localization of CB₂. However, for meaningful information to be drawn from them it is essential that proper controls are followed [as reviewed by Bussolati and Leonardo (2008); Lorincz and Nusser (2008); Saper and Sawchenko (2003)]. Briefly, some of the controls are: adsorption with the immunizing peptide, parallel, blinded staining of wild-type and knockout tissue, the use of two or more antibodies raised against distinct epitopes, antibody titration, omitting the primary antibody from the staining procedure and supporting these findings with RT-PCR, *in situ* hybridization and other such detection methods. Using just one control for one experimental setup is usually insufficient proof of specificity. These basic controls must be remembered when interpreting the data presented from any study cited in this review and future studies as well.

In conclusion, despite originally being thought of as the 'peripheral' cannabinoid receptor, considerable functional and anatomical evidence suggests that CB₂ is expressed in the nervous system – certainly in activated microglia and very likely in some neurons. In addition, this raises the point that any report that identifies CB₂ in neurons of the nervous system must incorporate careful controls to ensure that the CB₂ signal found originates from neurons and not from microglia or immune cells associated with brain blood vessels. Given the importance of determining the functional role of CB₂ in the CNS, under what conditions it is up regulated, and the potential therapeutic applications of CB₂ agonists it is vital to understand where in the CNS CB₂ receptors are expressed. We encourage those working in the field and those reviewing manuscripts to conduct and review these studies in a careful, thoughtful and rigorous fashion.

Table 1 CB₂ distribution in the peripheral and central nervous systems

<i>Location</i>	<i>Presence</i>	<i>Detection method</i>	<i>Species</i>	<i>Reference</i>
Whole brain				
Whole brain	Absent	ISH	Mouse	Munro et al., 1993
Whole brain	Absent	NB/RT-PCR	Human	Galiegue et al., 1995
Whole brain	Absent	NB/RT-PCR	Mouse	Schatz et al., 1997
Whole brain	Absent	SB	Mouse	McCoy et al., 1999
Whole brain	Absent	RLB/GTP γ S	Rat	Griffin et al., 1999
Whole brain	Absent	ISH	Mouse	Howlett et al., 2002
Whole brain	Present	WB	Rat	Gong et al., 2006
Whole brain	Present	WB/RT-PCR	Mouse	Onaivi, 2006; Onaivi et al., 2008b
Brainstem				
DMNX	Present	RT-PCR/WB/IHC	Rat	Van Sickle et al., 2005
DMNX	Present	ICC	Mouse	Van Sickle et al., 2005
DMNX	Present	WB/ICC/functional	Ferret	Van Sickle et al., 2005
Cochlear nuclei	Present	IHC	Rat	Baek et al., 2008
Medial vestibular nuclei	Present	IHC	Rat	Baek et al., 2008
Inferior olive and pontine nuclei	Present	IHC/ICC/RT-PCR	Rat	Viscomi et al., 2009
Brainstem	Present	RT-PCR/IHC	Rat	Gong et al., 2006
Brainstem	Present	RT-PCR	Mouse	Onaivi et al., 2006; Liu et al., 2009
Cerebellum				
Cerebellum	Absent	NB/RT-PCR	Human	Galiegue et al., 1995
Granule and Purkinje cell layers	Present	ISH/RLB	Mouse	Skaper et al., 1996
Cerebellum	Absent	NB/RT-PCR	Rat	Schatz et al., 1997
Cerebellum	Absent	NB/RLB/GTP γ S	Rat	Griffin et al., 1999
Cerebellum	Present	RT-PCR/WB	Rat	Van Sickle et al., 2005
Cerebellum	Present	WB	Ferret	Van Sickle et al., 2005
Cerebellum	Present	ICC	Rat	Ashton et al., 2006
Cerebellum	Present	IHC	Rat	Baek et al., 2008
Cerebellum	Present	IHC/ISH	Rat/mouse	Onaivi, 2006; Onaivi et al., 2008b
Cerebellum	Present	RT-PCR	Human	Liu et al., 2009
Cortex				
Cortex	Absent	RT-PCR/NB	Human	Galiegue et al., 1995
Cortex	Absent	NB	Rat	Griffin et al., 1999
Cortex	Absent	RT-PCR	Rat	Beltramo et al., 2006
Cortex	Present/absent	IHC/RT-PCR	Rat	Gong et al., 2006
Cortex	Present	IHC	Mouse	Onaivi, 2006; Onaivi et al., 2006; 2008b
Cortex	Present	RT-PCR	Mouse/human	Liu et al., 2009
Hippocampus				
Hippocampus	Present/absent	IHC/RT-PCR	Rat	Gong et al., 2006; Brusco et al., 2008
Hippocampus	Present	IHC	Mouse	Onaivi, 2006; Onaivi et al., 2006; 2008b
Hippocampus	Present	IHC	Rat	Onaivi et al., 2008b
Hippocampus	Present	RT-PCR	Human	Liu et al., 2009
Other brain regions				
Thalamus	Present/absent	IHC/RT-PCR	Rat	Gong et al., 2006
Hypothalamus	Present	RT-PCR	Rat	Gong et al., 2006
Midbrain	Present/absent	IHC/RT-PCR	Rat	Gong et al., 2006
Olfactory bulb	Present/absent	IHC/RT-PCR	Rat	Gong et al., 2006
VPN of thalamus	Present	Functional	Rat	Jhaveri et al., 2008
Entorhinal cortex	Present	Functional	Rat	Morgan et al., 2009
Peripheral neurons/spinal cord				
Vas deferens	Present	Functional	Mouse	Griffin et al., 1997
Spinal cord	Absent	NB	Rat	Griffin et al., 1999
DRG neurons	Absent	ISH	Rat	Hohmann and Herkenham, 1999b
Retina	Present	ISH/RT-PCR	Mouse	Lu et al., 2000
DRG neurons	Present	FACS	Rat	Ross et al., 2001
Enteric system	Present	RT-PCR	Rat	Storr et al., 2002
Trigeminal ganglia	Absent	ISH	Rat	Price et al., 2003
Spinal cord	Present	ISH	Rat	Zhang et al., 2003
Vagus nerve	Present	Functional	Guinea pig/human	Patel et al., 2003
DRG neurons	Present	Functional	Rat	Nackley et al., 2004
DRG neurons	Present	Functional	Rat	Elmes et al., 2004
Nodose ganglion	Present	RT-PCR	Rat	Burdyga et al., 2004
Vagus nerve trunk	Absent	RT-PCR	Human	Burdyga et al., 2004
DRG neurons	Present	Functional	Rat	Sagar et al., 2005
DRG neurons/spinal cord	Inducible	ICC	Rat/mouse	Wotherspoon et al., 2005
Skin sensory neurons	Present	IHC	Human	Stander et al., 2005
DRG neurons/spinal cord	Present	Functional/RT-PCR	Rat	Beltramo et al., 2006
Mesenteric sensory neurons	Present	Functional	Mouse	Hillsley et al., 2007
Vagus nerve	Present	Functional	Guinea pig	Belvisi et al., 2008
Myenteric neurons	Present	Functional/ICC/RT-PCR	Rat	Duncan et al., 2008
DRG neurons	Present	IHC/WB/functional	Gp/human/rat/mouse	Anand et al., 2008

Table 1 Continued

Location	Presence	Detection method	Species	Reference
Neural progenitor cells				
Hippocampal neural progenitor cells	Present	RT-PCR/ICC/functional	Mouse	Palazuelos et al., 2006
Neurospheres	Present	WB/ICC/functional	Mouse	Molina-Holgado et al., 2007
SVZ neural progenitor cells	Present	WB/ICC/functional	Mouse	Goncalves et al., 2008

Summary of studies investigating CB₂ expression in the nervous system. Details the location studied, whether CB₂ was detected or not, the method(s) used to detect it, the species analysed and the reference(s) for the studies.

FACS, fluorescence-activated cell sorting; ICC, immunocytochemistry; IHC, immunohistochemistry; ISH, *in situ* hybridization; NB, Northern blot; RLB, radioligand binding; RT-PCR, reverse transcriptase polymerase chain reaction; SB, Southern blot; WB, Western blot.

Acknowledgements

Grant no. NIH DA21696 RR025761.

Conflicts of interest

None.

References

- Abbadie C, Bhargoo S, De Koninck Y, Malcangio M, Melik-Parsadaniantz S, White FA (2009). Chemokines and pain mechanisms. *Brain Res Rev* **60**: 125–134.
- Adams IB, Martin BR (1996). Cannabis: pharmacology and toxicology in animals and humans. *Addiction* **91**: 1585–1614.
- Alexander SP, Mathie A, Peters JA (2008). Guide to Receptors and Channels (GRAC), 3rd edition. *Br J Pharmacol* **153**: S1–209.
- Anand U, Otto WR, Sanchez-Herrera D, Facer P, Yiangou Y, Korchev Y *et al.* (2008). Cannabinoid receptor CB₂ localisation and agonist-mediated inhibition of capsaicin responses in human sensory neurons. *Pain* **138**: 667–680.
- Anand P, Whiteside G, Fowler CJ, Hohmann AG (2009). Targeting CB₂ receptors and the endocannabinoid system for the treatment of pain. *Brain Res Rev* **60**: 255–266.
- Ashton JC, Friberg D, Darlington CL, Smith PF (2006). Expression of the cannabinoid CB₂ receptor in the rat cerebellum: an immunohistochemical study. *Neurosci Lett* **396**: 113–116.
- Ashton JC, Glass M (2007). The Cannabinoid CB₂ Receptor as a Target for Inflammation-Dependent Neurodegeneration. *Curr Neuroparmacol* **5**: 73–80.
- Ashton JC, Rahman RM, Nair SM, Sutherland BA, Glass M, Appleton I (2007). Cerebral hypoxia-ischemia and middle cerebral artery occlusion induce expression of the cannabinoid CB₂ receptor in the brain. *Neurosci Lett* **412**: 114–117.
- Baek JH, Zheng Y, Darlington CL, Smith PF (2008). Cannabinoid CB₂ receptor expression in the rat brainstem cochlear and vestibular nuclei. *Acta Otolaryngol* **128**: 961–967.
- Bayewitch M, Avidor-Reiss T, Levy R, Barg J, Mechoulam R, Vogel Z (1995). The peripheral cannabinoid receptor: adenylate cyclase inhibition and G protein coupling. *FEBS Lett* **375**: 143–147.
- Beltramo M, Bernardini N, Bertorelli R, Campanella M, Nicolussi E, Fredduzzi S *et al.* (2006). CB₂ receptor-mediated antihyperalgesia: possible direct involvement of neural mechanisms. *Eur J Neurosci* **23**: 1530–1538.
- Belvisi MG, Patel HJ, Freund-Michel V, Hele DJ, Crispino N, Birrell MA (2008). Inhibitory activity of the novel CB₂ receptor agonist, GW833972A, on guinea-pig and human sensory nerve function in the airways. *Br J Pharmacol* **155**: 547–557.
- Benito C, Nunez E, Tolon RM, Carrier EJ, Rabano A, Hillard CJ *et al.* (2003). Cannabinoid CB₂ receptors and fatty acid amide hydrolase are selectively overexpressed in neuritic plaque-associated glia in Alzheimer's disease brains. *J Neurosci* **23**: 11136–11141.
- Berdyshev EV (2000). Cannabinoid receptors and the regulation of immune response. *Chem Phys Lipids* **108**: 169–190.
- Bingham B, Jones PG, Uveges AJ, Kotnis S, Lu P, Smith VA *et al.* (2007). Species-specific *in vitro* pharmacological effects of the cannabinoid receptor 2 (CB₂) selective ligand AM1241 and its resolved enantiomers. *Br J Pharmacol* **151**: 1061–1070.
- Bouaboula M, Desnoyer N, Carayon P, Combes T, Casellas P (1999a). Gi protein modulation induced by a selective inverse agonist for the peripheral cannabinoid receptor CB₂: implication for intracellular signalization cross-regulation. *Mol Pharmacol* **55**: 473–480.
- Bouaboula M, Dussosoy D, Casellas P (1999b). Regulation of peripheral cannabinoid receptor CB₂ phosphorylation by the inverse agonist SR 144528. Implications for receptor biological responses. *J Biol Chem* **274**: 20397–20405.
- Bouaboula M, Poinot-Chazel C, Marchand J, Canat X, Bourrie B, Rinaldi-Carmona M *et al.* (1996). Signaling pathway associated with stimulation of CB₂ peripheral cannabinoid receptor. Involvement of both mitogen-activated protein kinase and induction of Krox-24 expression. *Eur J Biochem* **237**: 704–711.
- Brown SM, Wager-Miller J, Mackie K (2002). Cloning and molecular characterization of the rat CB₂ cannabinoid receptor. *Biochim Biophys Acta* **1576**: 255–264.
- Brusco A, Tagliaferro P, Saez T, Onaivi ES (2008). Postsynaptic localization of CB₂ cannabinoid receptors in the rat hippocampus. *Synapse* **62**: 944–949.
- Buckley NE, McCoy KL, Mezey E, Bonner T, Zimmer A, Felder CC *et al.* (2000). Immunomodulation by cannabinoids is absent in mice deficient for the cannabinoid CB₂ receptor. *Eur J Pharmacol* **396**: 141–149.
- Burdyshev G, Lal S, Varro A, Dimaline R, Thompson DG, Dockray GJ (2004). Expression of cannabinoid CB₁ receptors by vagal afferent neurons is inhibited by cholecystokinin. *J Neurosci* **24**: 2708–2715.
- Bussolati G, Leonardo E (2008). Technical pitfalls potentially affecting diagnoses in immunohistochemistry. *J Clin Pathol* **61**: 1184–1192.
- Cabral GA, Griffin-Thomas L (2009). Emerging role of the cannabinoid receptor CB₂ in immune regulation: therapeutic prospects for neuroinflammation. *Expert Rev Mol Med* **11**: e3.
- Cabral GA, Raborn ES, Griffin L, Dennis J, Marciano-Cabral F (2008). CB₂ receptors in the brain: role in central immune function. *Br J Pharmacol* **153**: 240–251.
- Calignano A, La Rana G, Giuffrida A, Piomelli D (1998). Control of pain initiation by endogenous cannabinoids. *Nature* **394**: 277–281.
- Carayon P, Marchand J, Dussosoy D, Derocq JM, Jbilou O, Bord A *et al.* (1998). Modulation and functional involvement of CB₂ peripheral cannabinoid receptors during B-cell differentiation. *Blood* **92**: 3605–3615.
- Carlisle SJ, Marciano-Cabral F, Staab A, Ludwick C, Cabral GA (2002). Differential expression of the CB₂ cannabinoid receptor by rodent

- macrophages and macrophage-like cells in relation to cell activation. *Int Immunopharmacol* 2: 69–82.
- Costa B (2007). On the pharmacological properties of Delta9-tetrahydrocannabinol (THC). *Chem Biodivers* 4: 1664–1677.
- Cullheim S, Thams S (2007). The microglial networks of the brain and their role in neuronal network plasticity after lesion. *Brain Res Rev* 55: 89–96.
- Derbenev AV, Stuart TC, Smith BN (2004). Cannabinoids suppress synaptic input to neurones of the rat dorsal motor nucleus of the vagus nerve. *J Physiol* 559: 923–938.
- Derocq JM, Segui M, Marchand J, Le Fur G, Casellas P (1995). Cannabinoids enhance human B-cell growth at low nanomolar concentrations. *FEBS Lett* 369: 177–182.
- Devane WA, Dysarz FA, 3rd, Johnson MR, Melvin LS, Howlett AC (1988). Determination and characterization of a cannabinoid receptor in rat brain. *Mol Pharmacol* 34: 605–613.
- Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, Griffin G *et al.* (1992). Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* 258: 1946–1949.
- Dittel BN (2008). Direct suppression of autoreactive lymphocytes in the central nervous system via the CB2 receptor. *Br J Pharmacol* 153: 271–276.
- Duncan M, Mouihate A, Mackie K, Keenan CM, Buckley NE, Davison JS *et al.* (2008). Cannabinoid CB2 receptors in the enteric nervous system modulate gastrointestinal contractility in lipopolysaccharide-treated rats. *Am J Physiol Gastrointest Liver Physiol* 295: G78–G87.
- Elmes SJ, Jhaveri MD, Smart D, Kendall DA, Chapman V (2004). Cannabinoid CB2 receptor activation inhibits mechanically evoked responses of wide dynamic range dorsal horn neurons in naive rats and in rat models of inflammatory and neuropathic pain. *Eur J Neurosci* 20: 2311–2320.
- Felder CC, Briley EM, Axelrod J, Simpson JT, Mackie K, Devane WA (1993). Anandamide, an endogenous cannabimimetic eicosanoid, binds to the cloned human cannabinoid receptor and stimulates receptor-mediated signal transduction. *Proc Natl Acad Sci U S A* 90: 7656–7660.
- Felder CC, Joyce KE, Briley EM, Mansouri J, Mackie K, Blond O *et al.* (1995). Comparison of the pharmacology and signal transduction of the human cannabinoid CB1 and CB2 receptors. *Mol Pharmacol* 48: 443–450.
- Fernandez-Ruiz J, Pazos MR, Garcia-Arencibia M, Sagredo O, Ramos JA (2008). Role of CB2 receptors in neuroprotective effects of cannabinoids. *Mol Cell Endocrinol* 286: S91–S96.
- Frey KA, Albin RL (2001). Receptor binding techniques. *Curr Protoc Neurosci* 14: 1–14. Chapter 1: Unit1 4.
- Galiegue S, Mary S, Marchand J, Dussosoy D, Carriere D, Carayon P *et al.* (1995). Expression of central and peripheral cannabinoid receptors in human immune tissues and leukocyte subpopulations. *Eur J Biochem* 232: 54–61.
- Gallant M, Dufresne C, Gareau Y, Guay D, Leblanc Y, Prasit P *et al.* (1996). New class of potent ligands for the human peripheral cannabinoid receptor. *Bioorganic & Medicinal Chemistry Lett* 6: 2263–2268.
- Goncalves MB, Suetterlin P, Yip P, Molina-Holgado F, Walker DJ, Oudin MJ *et al.* (2008). A diacylglycerol lipase-CB2 cannabinoid pathway regulates adult subventricular zone neurogenesis in an age-dependent manner. *Mol Cell Neurosci* 38: 526–536.
- Gong JP, Onaivi ES, Ishiguro H, Liu QR, Tagliaferro PA, Brusco A *et al.* (2006). Cannabinoid CB2 receptors: immunohistochemical localization in rat brain. *Brain Res* 1071: 10–23.
- Gonsiorek W, Lunn C, Fan X, Narula S, Lundell D, Hipkin RW (2000). Endocannabinoid 2-arachidonyl glycerol is a full agonist through human type 2 cannabinoid receptor: antagonism by anandamide. *Mol Pharmacol* 57: 1045–1050.
- Griffin G, Fernando SR, Ross RA, McKay NG, Ashford ML, Shire D *et al.* (1997). Evidence for the presence of CB2-like cannabinoid receptors on peripheral nerve terminals. *Eur J Pharmacol* 339: 53–61.
- Griffin G, Tao Q, Abood ME (2000). Cloning and pharmacological characterization of the rat CB(2) cannabinoid receptor. *J Pharmacol Exp Ther* 292: 886–894.
- Griffin G, Wray EJ, Tao Q, McAllister SD, Rorrer WK, Aung MM *et al.* (1999). Evaluation of the cannabinoid CB2 receptor-selective antagonist, SR144528: further evidence for cannabinoid CB2 receptor absence in the rat central nervous system. *Eur J Pharmacol* 377: 117–125.
- Grimaldi P, Orlando P, Di Siena S, Lolicato F, Petrosino S, Bisogno T *et al.* (2009). The endocannabinoid system and pivotal role of the CB2 receptor in mouse spermatogenesis. *Proc Natl Acad Sci U S A* 106: 11131–11136.
- Hanisch UK, Kettenmann H (2007). Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 10: 1387–1394.
- Hanus L, Abu-Lafi S, Fride E, Breuer A, Vogel Z, Shalev DE *et al.* (2001). 2-arachidonyl glyceryl ether, an endogenous agonist of the cannabinoid CB1 receptor. *Proc Natl Acad Sci U S A* 98: 3662–3665.
- Hanus L, Breuer A, Tchilibon S, Shiloah S, Goldenberg D, Horowitz M *et al.* (1999). HU-308: a specific agonist for CB(2), a peripheral cannabinoid receptor. *Proc Natl Acad Sci U S A* 96: 14228–14233.
- He F, Song ZH (2007). Molecular and cellular changes induced by the activation of CB2 cannabinoid receptors in trabecular meshwork cells. *Mol Vis* 13: 1348–1356.
- Henstridge CM, Balenga NA, Ford LA, Ross RA, Waldhoer M, Irving AJ (2009). The GPR55 ligand L-alpha-lysophosphatidylinositol promotes RhoA-dependent Ca²⁺ signaling and NFAT activation. *FASEB J* 23: 183–193.
- Herkenham M, Lynn AB, Johnson MR, Melvin LS, de Costa BR, Rice KC (1991). Characterization and localization of cannabinoid receptors in rat brain: a quantitative in vitro autoradiographic study. *J Neurosci* 11: 563–583.
- Herkenham M, Lynn AB, Little MD, Johnson MR, Melvin LS, de Costa BR *et al.* (1990). Cannabinoid receptor localization in brain. *Proc Natl Acad Sci U S A* 87: 1932–1936.
- Herrera B, Carracedo A, Diez-Zaera M, Gomez del Pulgar T, Guzman M, Velasco G (2006). The CB2 cannabinoid receptor signals apoptosis via ceramide-dependent activation of the mitochondrial intrinsic pathway. *Exp Cell Res* 312: 2121–2131.
- Herrera B, Carracedo A, Diez-Zaera M, Guzman M, Velasco G (2005). p38 MAPK is involved in CB2 receptor-induced apoptosis of human leukaemia cells. *FEBS Lett* 579: 5084–5088.
- Hillsley K, McCaul C, Aerssens J, Peeters PJ, Gijzen H, Moechars D *et al.* (2007). Activation of the cannabinoid 2 (CB2) receptor inhibits murine mesenteric afferent nerve activity. *Neurogastroenterol Motil* 19: 769–777.
- Hohmann AG, Herkenham M (1999a). Cannabinoid receptors undergo axonal flow in sensory nerves. *Neuroscience* 92: 1171–1175.
- Hohmann AG, Herkenham M (1999b). Localization of central cannabinoid CB1 receptor messenger RNA in neuronal subpopulations of rat dorsal root ganglia: a double-label in situ hybridization study. *Neuroscience* 90: 923–931.
- Howlett AC (2002). The cannabinoid receptors. *Prostaglandins Other Lipid Mediat* 68–69: 619–631.
- Howlett AC, Barth F, Bonner TI, Cabral G, Casellas P, Devane WA *et al.* (2002). International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol Rev* 54: 161–202.
- Huffman JW, Liddle J, Yu S, Aung MM, Abood ME, Wiley JL *et al.* (1999). 3-(1',1'-Dimethylbutyl)-1-deoxy-delta8-THC and related compounds: synthesis of selective ligands for the CB2 receptor. *Bioorg Med Chem* 7: 2905–2914.
- Ibrahim MM, Deng H, Zvonok A, Cockayne DA, Kwan J, Mata HP *et al.*, (2003). Activation of CB2 cannabinoid receptors by AM1241 inhibits experimental neuropathic pain: pain inhibition by recep-

- tors not present in the CNS. *Proc Natl Acad Sci U S A* **100**: 10529–10533.
- Ibrahim MM, Porreca F, Lai J, Albrecht PJ, Rice FL, Khodorova A *et al.*, (2005). CB₂ cannabinoid receptor activation produces antinociception by stimulating peripheral release of endogenous opioids. *Proc Natl Acad Sci U S A* **102**: 3093–3098.
- Ibrahim MM, Rude ML, Stagg NJ, Mata HP, Lai J, Vanderah TW *et al.*, (2006). CB₂ cannabinoid receptor mediation of antinociception. *Pain* **122**: 36–42.
- Jhaveri MD, Elmes SJ, Richardson D, Barrett DA, Kendall DA, Mason R *et al.* (2008). Evidence for a novel functional role of cannabinoid CB₂ receptors in the thalamus of neuropathic rats. *Eur J Neurosci* **27**: 1722–1730.
- Jiang S, Fu Y, Williams J, Wood J, Pandarinathan L, Avraham S *et al.* (2007). Expression and function of cannabinoid receptors CB₁ and CB₂ and their cognate cannabinoid ligands in murine embryonic stem cells. *Plos One* **2**: e641.
- Julien B, Grenard P, Teixeira-Clerc F, Van Nhieu JT, Li L, Karsak M *et al.* (2005). Antifibrogenic role of the cannabinoid receptor CB₂ in the liver. *Gastroenterology* **128**: 742–755.
- Kapur A, Zhao P, Sharir H, Bai Y, Caron MG, Barak LS *et al.* (2009). Atypical responsiveness of the orphan receptor GPR55 to cannabinoid ligands. *J Biol Chem* **284**: 29817–29827.
- Khanolkar AD, Lu D, Ibrahim M, Duclos RI, Jr, Thakur GA, Malan TP *et al.* (2007). Cannabillactones: a novel class of CB₂ selective agonists with peripheral analgesic activity. *J Med Chem* **50**: 6493–6500.
- Klegeris A, Bissonnette CJ, McGeer PL (2003). Reduction of human monocytic cell neurotoxicity and cytokine secretion by ligands of the cannabinoid-type CB₂ receptor. *Br J Pharmacol* **139**: 775–786.
- Kress M, Kuner R (2009). Mode of action of cannabinoids on nociceptive nerve endings. *Exp Brain Res* **196**: 79–88.
- Lauckner JE, Jensen JB, Chen HY, Lu HC, Hille B, Mackie K (2008). GPR55 is a cannabinoid receptor that increases intracellular calcium and inhibits M current. *Proc Natl Acad Sci U S A* **105**: 2699–2704.
- Lion T (2001). Current recommendations for positive controls in RT-PCR assays. *Leukemia* **15**: 1033–1037.
- Liu QR, Pan CH, Hishimoto A, Li CY, Xi ZX, Llorente-Berzal A *et al.* (2009). Species differences in cannabinoid receptor 2 (CB₂ gene): identification of novel human and rodent CB₂ isoforms, differential tissue expression and regulation by cannabinoid receptor ligands. *Genes Brain Behav* **8**: 519–530.
- Lorincz A, Nusser Z (2008). Specificity of immunoreactions: the importance of testing specificity in each method. *J Neurosci* **28**: 9083–9086.
- Lu Q, Straiker A, Maguire G (2000). Expression of CB₂ cannabinoid receptor mRNA in adult rat retina. *Vis Neurosci* **17**: 91–95.
- Lynn AB, Herkenham M (1994). Localization of cannabinoid receptors and nonsaturable high-density cannabinoid binding sites in peripheral tissues of the rat: implications for receptor-mediated immune modulation by cannabinoids. *J Pharmacol Exp Ther* **268**: 1612–1623.
- Maccarrone M (2008). CB₂ receptors in reproduction. *Br J Pharmacol* **153**: 189–198.
- Mackie K (2005). Distribution of cannabinoid receptors in the central and peripheral nervous system. *Handb Exp Pharmacol* **168**: 299–325.
- Malan TP, Jr, Ibrahim MM, Deng H, Liu Q, Mata HP, Vanderah T *et al.* (2001). CB₂ cannabinoid receptor-mediated peripheral antinociception. *Pain* **93**: 239–245.
- Malan TP, Jr, Ibrahim MM, Lai J, Vanderah TW, Makriyannis A, Porreca F (2003). CB₂ cannabinoid receptor agonists: pain relief without psychoactive effects? *Curr Opin Pharmacol* **3**: 62–67.
- Maresz K, Pryce G, Ponomarev ED, Marsicano G, Croxford JL, Shriver LP *et al.* (2007). Direct suppression of CNS autoimmune inflammation via the cannabinoid receptor CB₁ on neurons and CB₂ on autoreactive T cells. *Nat Med* **13**: 492–497.
- Mathison R, Ho W, Pittman QJ, Davison JS, Sharkey KA (2004). Effects of cannabinoid receptor-2 activation on accelerated gastrointestinal transit in lipopolysaccharide-treated rats. *Br J Pharmacol* **142**: 1247–1254.
- Matsuda LA, Bonner TI, Lolait SJ (1993). Localization of cannabinoid receptor mRNA in rat brain. *J Comp Neurol* **327**: 535–550.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI (1990). Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* **346**: 561–564.
- McCoy KL, Matveyeva M, Carlisle SJ, Cabral GA (1999). Cannabinoid inhibition of the processing of intact lysozyme by macrophages: evidence for CB₂ receptor participation. *J Pharmacol Exp Ther* **289**: 1620–1625.
- Miller AM, Stella N (2008). CB₂ receptor-mediated migration of immune cells: it can go either way. *Br J Pharmacol* **153**: 299–308.
- Molina-Holgado F, Rubio-Araiz A, Garcia-Ovejero D, Williams RJ, Moore JD, Arevalo-Martin A *et al.* (2007). CB₂ cannabinoid receptors promote mouse neural stem cell proliferation. *Eur J Neurosci* **25**: 629–634.
- Morgan NH, Stanford IM, Woodhall GL (2009). Functional CB₂ type cannabinoid receptors at CNS synapses. *Neuropharmacology* **57**: 356–368.
- Mukherjee S, Adams M, Whiteaker K, Daza A, Kage K, Cassar S *et al.* (2004). Species comparison and pharmacological characterization of rat and human CB₂ cannabinoid receptors. *Eur J Pharmacol* **505**: 1–9.
- Munro S, Thomas KL, Abu-Shaar M (1993). Molecular characterization of a peripheral receptor for cannabinoids. *Nature* **365**: 61–65.
- Nackley AG, Zvonok AM, Makriyannis A, Hohmann AG (2004). Activation of cannabinoid CB₂ receptors suppresses C-fiber responses and windup in spinal wide dynamic range neurons in the absence and presence of inflammation. *J Neurophysiol* **92**: 3562–3574.
- Naguib M, Diaz P, Xu JJ, Astruc-Diaz F, Craig S, Vivas-Mejia P *et al.* (2008). MDA7: a novel selective agonist for CB₂ receptors that prevents allodynia in rat neuropathic pain models. *Br J Pharmacol* **155**: 1104–1116.
- Ofek O, Karsak M, Leclerc N, Fogel M, Frenkel B, Wright K *et al.* (2006). Peripheral cannabinoid receptor, CB₂, regulates bone mass. *Proc Natl Acad Sci U S A* **103**: 696–701.
- Ohta H, Ishizaka T, Tatsuzuki M, Yoshinaga M, Iida I, Yamaguchi T *et al.* (2008). Imine derivatives as new potent and selective CB₂ cannabinoid receptor agonists with an analgesic action. *Bioorg Med Chem* **16**: 1111–1124.
- Onaivi ES (2006). Neuropsychobiological evidence for the functional presence and expression of cannabinoid CB₂ receptors in the brain. *Neuropsychobiology* **54**: 231–246.
- Onaivi ES, Carpio O, Ishiguro H, Schanz N, Uhl GR, Benno R (2008a). Behavioral effects of CB₂ cannabinoid receptor activation and its influence on food and alcohol consumption. *Ann N Y Acad Sci* **1139**: 426–433.
- Onaivi ES, Ishiguro H, Gong JP, Patel S, Meozzi PA, Myers L *et al.* (2008b). Brain neuronal CB₂ cannabinoid receptors in drug abuse and depression: from mice to human subjects. *Plos One* **3**: e1640.
- Onaivi ES, Ishiguro H, Gong JP, Patel S, Perchuk A, Meozzi PA *et al.* (2006). Discovery of the presence and functional expression of cannabinoid CB₂ receptors in brain. *Ann N Y Acad Sci* **1074**: 514–536.
- Palazuelos J, Aguado T, Egia A, Mechoulam R, Guzman M, Galve-Roperh I (2006). Non-psychoactive CB₂ cannabinoid agonists stimulate neural progenitor proliferation. *FASEB J* **20**: 2405–2407.
- Patel HJ, Birrell MA, Crispino N, Hele DJ, Venkatesan P, Barnes PJ *et al.* (2003). Inhibition of guinea-pig and human sensory nerve activity and the cough reflex in guinea-pigs by cannabinoid (CB₂) receptor activation. *Br J Pharmacol* **140**: 261–268.
- Pertwee R, Griffin G, Fernando S, Li X, Hill A, Makriyannis A (1995). AM630, a competitive cannabinoid receptor antagonist. *Life Sci* **56**: 1949–1955.
- Pietr M, Kozela E, Levy R, Rimmerman N, Lin YH, Stella N *et al.* (2009).

- Differential changes in GPR55 during microglial cell activation. *FEBS Lett* **583**: 2071–2076.
- Price TJ, Helesic G, Parghi D, Hargreaves KM, Flores CM (2003). The neuronal distribution of cannabinoid receptor type 1 in the trigeminal ganglion of the rat. *Neuroscience* **120**: 155–162.
- Price DA, Martinez AA, Seillier A, Koek W, Acosta Y, Fernandez E *et al.* (2009). WIN55,212-2, a cannabinoid receptor agonist, protects against nigrostriatal cell loss in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease. *Eur J Neurosci* **29**: 2177–2186.
- Rinaldi-Carmona M, Barth F, Millan J, Derocq JM, Casellas P, Congy C *et al.* (1998). SR 144528, the first potent and selective antagonist of the CB₂ cannabinoid receptor. *J Pharmacol Exp Ther* **284**: 644–650.
- Roche R, Hoareau L, Bes-Houtmann S, Gonthier MP, Laborde C, Baron JF *et al.* (2006). Presence of the cannabinoid receptors, CB₁ and CB₂, in human omental and subcutaneous adipocytes. *Histochem Cell Biol* **126**: 177–187.
- Romero-Sandoval EA, Horvath R, Landry RP, DeLeo JA (2009). Cannabinoid receptor type 2 activation induces a microglial anti-inflammatory phenotype and reduces migration via MKP induction and ERK dephosphorylation. *Mol Pain* **5**: 25.
- Ross RA, Brockie HC, Stevenson LA, Murphy VL, Templeton F, Makriyannis A *et al.* (1999). Agonist-inverse agonist characterization at CB₁ and CB₂ cannabinoid receptors of L759633, L759656, and AM630. *Br J Pharmacol* **126**: 665–672.
- Ross RA, Coutts AA, McFarlane SM, Anavi-Goffer S, Irving AJ, Pertwee RG *et al.* (2001). Actions of cannabinoid receptor ligands on rat cultured sensory neurones: implications for antinociception. *Neuropharmacology* **40**: 221–232.
- Ryberg E, Larsson N, Sjogren S, Hjorth S, Hermansson NO, Leonova J *et al.* (2007). The orphan receptor GPR55 is a novel cannabinoid receptor. *Br J Pharmacol* **152**: 1092–1101.
- Sagar DR, Kelly S, Millns PJ, O'Shaughnessy CT, Kendall DA, Chapman V (2005). Inhibitory effects of CB₁ and CB₂ receptor agonists on responses of DRG neurons and dorsal horn neurons in neuropathic rats. *Eur J Neurosci* **22**: 371–379.
- Saper CB, Sawchenko PE (2003). Magic peptides, magic antibodies: guidelines for appropriate controls for immunohistochemistry. *J Comp Neurol* **465**: 161–163.
- Schatz AR, Lee M, Condie RB, Pulaski JT, Kaminski NE (1997). Cannabinoid receptors CB₁ and CB₂: a characterization of expression and adenylate cyclase modulation within the immune system. *Toxicol Appl Pharmacol* **142**: 278–287.
- Shire D, Calandra B, Rinaldi-Carmona M, Oustric D, Pesseque B, Bonnin-Cabanne O *et al.* (1996). Molecular cloning, expression and function of the murine CB₂ peripheral cannabinoid receptor. *Biochim Biophys Acta* **1307**: 132–136.
- Shmist YA, Goncharov I, Eichler M, Shneyvays V, Isaac A, Vogel Z *et al.* (2006). Delta-9-tetrahydrocannabinol protects cardiac cells from hypoxia via CB₂ receptor activation and nitric oxide production. *Mol Cell Biochem* **283**: 75–83.
- Shoemaker JL, Joseph BK, Ruckle MB, Mayeux PR, Prather PL (2005a). The endocannabinoid noladin ether acts as a full agonist at human CB₂ cannabinoid receptors. *J Pharmacol Exp Ther* **314**: 868–875.
- Shoemaker JL, Ruckle MB, Mayeux PR, Prather PL (2005b). Agonist-directed trafficking of response by endocannabinoids acting at CB₂ receptors. *J Pharmacol Exp Ther* **315**: 828–838.
- Showalter VM, Compton DR, Martin BR, Abood ME (1996). Evaluation of binding in a transfected cell line expressing a peripheral cannabinoid receptor (CB₂): identification of cannabinoid receptor subtype selective ligands. *J Pharmacol Exp Ther* **278**: 989–999.
- Skaper SD, Buriani A, Dal Toso R, Petrelli L, Romanello S, Facci L *et al.* (1996). The ALLamide palmitoylethanolamide and cannabinoids, but not anandamide, are protective in a delayed postglutamate paradigm of excitotoxic death in cerebellar granule neurons. *Proc Natl Acad Sci U S A* **93**: 3984–3989.
- Slipetz DM, O'Neill GP, Favreau L, Dufresne C, Gallant M, Gareau Y *et al.* (1995). Activation of the human peripheral cannabinoid receptor results in inhibition of adenylyl cyclase. *Mol Pharmacol* **48**: 352–361.
- Stander S, Schmelz M, Metzger D, Luger T, Rukwied R (2005). Distribution of cannabinoid receptor 1 (CB₁) and 2 (CB₂) on sensory nerve fibers and adnexal structures in human skin. *J Dermatol Sci* **38**: 177–188.
- Stella N (2004). Cannabinoid signaling in glial cells. *Glia* **48**: 267–277.
- Storr M, Gaffal E, Saur D, Schusdziarra V, Allescher HD (2002). Effect of cannabinoids on neural transmission in rat gastric fundus. *Can J Physiol Pharmacol* **80**: 67–76.
- Sugiura T, Kondo S, Kishimoto S, Miyashita T, Nakane S, Kodaka T *et al.* (2000). Evidence that 2-arachidonoylglycerol but not N-palmitoylethanolamine or anandamide is the physiological ligand for the cannabinoid CB₂ receptor. Comparison of the agonistic activities of various cannabinoid receptor ligands in HL-60 cells. *J Biol Chem* **275**: 605–612.
- Sugiura T, Kondo S, Sukagawa A, Nakane S, Shinoda A, Itoh K *et al.* (1995). 2-Arachidonoylglycerol: a possible endogenous cannabinoid receptor ligand in brain. *Biochem Biophys Res Commun* **215**: 89–97.
- Suzuki T, Higgins PJ, Crawford DR (2000). Control selection for RNA quantitation. *Biotechniques* **29**: 332–337.
- Urban JD, Clarke WP, von Zastrow M, Nichols DE, Kobilka B, Weinstein H *et al.* (2007). Functional selectivity and classical concepts of quantitative pharmacology. *J Pharmacol Exp Ther* **320**: 1–13.
- Valenzano KJ, Tafesse L, Lee G, Harrison JE, Boulet JM, Gottshall SL *et al.* (2005). Pharmacological and pharmacokinetic characterization of the cannabinoid receptor 2 agonist, GW405833, utilizing rodent models of acute and chronic pain, anxiety, ataxia and catalepsy. *Neuropharmacology* **48**: 658–672.
- Van Sickle MD, Duncan M, Kingsley PJ, Mouihate A, Urbani P, Mackie K *et al.* (2005). Identification and functional characterization of brainstem cannabinoid CB₂ receptors. *Science* **310**: 329–332.
- Viscomi MT, Oddi S, Latini L, Pasquariello N, Florenzano F, Bernardi G *et al.* (2009). Selective CB₂ receptor agonism protects central neurons from remote axotomy-induced apoptosis through the PI3K/Akt pathway. *J Neurosci* **29**: 4564–4570.
- Walter L, Franklin A, Witting A, Wade C, Xie Y, Kunos G *et al.* (2003). Nonpsychotropic cannabinoid receptors regulate microglial cell migration. *J Neurosci* **23**: 1398–1405.
- Whiteside GT, Lee GP, Valenzano KJ (2007). The role of the cannabinoid CB₂ receptor in pain transmission and therapeutic potential of small molecule CB₂ receptor agonists. *Curr Med Chem* **14**: 917–936.
- Wotherspoon G, Fox A, McIntyre P, Colley S, Bevan S, Winter J (2005). Peripheral nerve injury induces cannabinoid receptor 2 protein expression in rat sensory neurons. *Neuroscience* **135**: 235–245.
- Yao BB, Hsieh G, Daza AV, Fan Y, Grayson GK, Garrison TR *et al.* (2009). Characterization of a cannabinoid CB₂ receptor-selective agonist, A-836339 [2,2,3,3-tetramethyl-cyclopropanecarboxylic acid [3-(2-methoxy-ethyl)-4,5-dimethyl-3H-thiazol-(2Z)-ylidene]-amide], using in vitro pharmacological assays, in vivo pain models, and pharmacological magnetic resonance imaging. *J Pharmacol Exp Ther* **328**: 141–151.
- Yao BB, Hsieh GC, Frost JM, Fan Y, Garrison TR, Daza AV *et al.* (2008). In vitro and in vivo characterization of A-796260: a selective cannabinoid CB₂ receptor agonist exhibiting analgesic activity in rodent pain models. *Br J Pharmacol* **153**: 390–401.
- Yao BB, Mukherjee S, Fan Y, Garrison TR, Daza AV, Grayson GK *et al.* (2006). In vitro pharmacological characterization of AM1241: a protean agonist at the cannabinoid CB₂ receptor? *Br J Pharmacol* **149**: 145–154.
- Yiangou Y, Facer P, Durrenberger P, Chessell IP, Naylor A, Bountra C *et al.* (2006). COX-2, CB₂ and P2X₇-immunoreactivities are increased in activated microglial cells/macrophages of multiple

- sclerosis and amyotrophic lateral sclerosis spinal cord. *BMC Neurol* **6**: 12.
- Zhang J, Hoffert C, Vu HK, Groblewski T, Ahmad S, O'Donnell D (2003). Induction of CB2 receptor expression in the rat spinal cord of neuropathic but not inflammatory chronic pain models. *Eur J Neurosci* **17**: 2750–2754.
- Zhong L, Geng L, Njie Y, Feng W, Song ZH (2005). CB2 cannabinoid receptors in trabecular meshwork cells mediate JWH015-induced enhancement of aqueous humor outflow facility. *Invest Ophthalmol Vis Sci* **46**: 1988–1992.
- Zoratti C, Kipmen-Korgun D, Osibow K, Malli R, Graier WF (2003). Anandamide initiates Ca(2+) signaling via CB2 receptor linked to phospholipase C in calf pulmonary endothelial cells. *Br J Pharmacol* **140**: 1351–1362.