



## REVIEW

# Celebrating 20 Years of Genetic Discoveries in Legume Nodulation and Symbiotic Nitrogen Fixation<sup>[OPEN]</sup>

Sonali Roy,<sup>a,1</sup> Wei Liu,<sup>a,1</sup> Raja Sekhar Nandety,<sup>a</sup> Ashley Crook,<sup>b</sup> Kirankumar S. Mysore,<sup>a</sup> Catalina I. Pislariu,<sup>c</sup> Julia Frugoli,<sup>b</sup> Rebecca Dickstein,<sup>d</sup> and Michael K. Udvardi<sup>a,2</sup>

<sup>a</sup> Noble Research Institute, Ardmore, Oklahoma 73401

<sup>b</sup> College of Science, Clemson University, Clemson, South Carolina 29634

<sup>c</sup> Department of Biology, Texas Woman's University, Denton Texas 76204

<sup>d</sup> Department of Biological Sciences and BioDiscovery Institute, University of North Texas, Denton Texas 76203

ORCID IDs: 0000-0002-8114-8300 (S.R.); 0000-0003-4653-0448 (W.L.); 0000-0002-1129-0790 (R.S.N.); 0000-0002-9805-5741 (K.S.M.); 0000-0001-7957-6341 (C.I.P.); 0000-0003-1642-5911 (J.F.); 0000-0002-5401-426X (R.D.); 0000-0001-9850-0828 (M.K.U.).

Since 1999, various forward- and reverse-genetic approaches have uncovered nearly 200 genes required for symbiotic nitrogen fixation (SNF) in legumes. These discoveries advanced our understanding of the evolution of SNF in plants and its relationship to other beneficial endosymbioses, signaling between plants and microbes, the control of microbial infection of plant cells, the control of plant cell division leading to nodule development, autoregulation of nodulation, intracellular accommodation of bacteria, nodule oxygen homeostasis, the control of bacteroid differentiation, metabolism and transport supporting symbiosis, and the control of nodule senescence. This review catalogs and contextualizes all of the plant genes currently known to be required for SNF in two model legume species, *Medicago truncatula* and *Lotus japonicus*, and two crop species, *Glycine max* (soybean) and *Phaseolus vulgaris* (common bean). We also briefly consider the future of SNF genetics in the era of pan-genomics and genome editing.

## INTRODUCTION

Fabaceae (legumes) is the third largest family of flowering plants, with members spread around the globe (Doyle and Luckow, 2003). Part of their success is due to N-fixing symbiosis with soil bacteria called rhizobia, which enables legumes to grow on atmospheric nitrogen (N<sub>2</sub>) in soils lacking mineral and organic N. Their natural ability to inject fixed N into soils make them keystone species for natural and agricultural ecosystems. Although once the primary source of N for agricultural systems, symbiotic nitrogen fixation (SNF) in legumes (approximately 50 million tons per annum) now contributes less than half that provided by chemical fertilizer N (Canfield et al., 2010). Unfortunately, manufactured fertilizer N is costly both economically and environmentally (Sutton et al., 2011). By contrast, SNF in legumes is seen as a sustainable source of N for agriculture. Research on the genetics of SNF, especially over the last 20 years, has revealed the great complexity of this process in plants as well as aspects of its evolution in different plant lineages. This knowledge, reviewed here, will underpin future efforts to improve SNF in legumes via conventional plant breeding and is already guiding more audacious efforts to engineer SNF in nonlegumes (Charpentier and Oldroyd, 2010).

During the 1980s, it was thought that a few dozen genes might be required for SNF in legumes based on the number of nodule-

specific genes, or nodulin genes, known at the time. With the genomic revolution over the past 20 years, it has become clear that thousands of genes are expressed at relatively high levels in legume nodules (Benedito et al., 2008; Mun et al., 2016). Over the same period, genetic approaches have revealed many genes with large effects on SNF, with mutations in these genes causing defects in nodule development and/or plant growth in the absence of soil N (Figure 1). Reverse genetics, which begins with a gene of interest and ends with a phenotype, in contrast to classical/forward genetics, has uncovered both large- and moderate-effect genes for SNF. Together, forward and reverse genetics have revolutionized our understanding of the molecular and cellular processes underpinning SNF, through the discovery of at least 196 genes required for effective symbiosis (Supplemental Data Set; <https://nobleapps.noble.org/legumegenetics>).


## Genetic Resources Developed for Legume Research

Mutations in genes are the “bread and butter” of genetics, and natural mutations in genes of *Pisum sativum* (garden pea) were instrumental in the development of genetics as a scientific discipline (Mendel, 1866). However, natural rates of mutation are low and generally mitigated by cellular DNA repair mechanisms. To accelerate genetic discovery, model species have been subjected to chemical and physical mutagens, including EMS and/or fast neutron bombardment, in *Lotus japonicus*, *Medicago truncatula*, soybean (*Glycine max*), and common bean (*Phaseolus vulgaris*; Szczyglowski et al., 1998; Penmetza and Cook, 2000; Perry et al., 2003; Bolon et al., 2011; Espina et al., 2018). The resulting mutant populations have been instrumental

<sup>1</sup> These authors contributed equally to this work.

<sup>2</sup> Address correspondence to [mudvardi@noble.org](mailto:mudvardi@noble.org).

<sup>[OPEN]</sup>Articles can be viewed without a subscription.  
[www.plantcell.org/cgi/doi/10.1105/tpc.19.00279](http://www.plantcell.org/cgi/doi/10.1105/tpc.19.00279)



**Figure 1.** Historical Timeline of Gene Discovery in *M. truncatula* and *L. japonicus*.

Mutant resources developed using a variety of mutagens have been utilized since 1995 in both species. Mutagens are indicated in boldface. The inset shows the number of SNF genes discovered by mutational analyses each year over the past 20 years. Refer to the text and the Supplemental Data Set for more information on individual genes.

in the discovery of genes required for SNF (Figures 1 and 2; Supplemental Data Set). Over time, mutant populations that were easier to work with were developed using transposons as mutagens. These include an *L. japonicus* mutant population resulting from T-DNA insertions (Schauer et al., 1998), an *M. truncatula* mutant population containing nearly a million genomic insertions of the tobacco (*Nicotiana tabacum*) retrotransposon *Tnt1* (Tadege et al., 2008; Pislariu et al., 2012), and an *L. japonicus* mutant population resulting from mobilization of the endogenous retrotransposon 1 (*LORE1*; Fukai et al., 2013; Malolepszy et al., 2016). The genomic DNA sequences adjacent to sites of transposon insertion, called flanking sequence tags, can be used to identify the causative mutation for both forward and reverse genetics. These strategies are particularly potent in diploid in-breeding

species such as *M. truncatula* ( $2n = 16$ ), *L. japonicus* ( $2n = 12$ ), and *P. vulgaris* ( $2n = 22$ ) but less so in paleopolyploid species such as soybean ( $2n = 40$ ).

### Overview of SNF

SNF is the culmination of a complex series of chemical and physical interactions between legumes and compatible rhizobia, including signaling processes that trigger changes in gene expression in both partners, shape partner selection, and suppress plant defenses. These signals also provide bacterial access to plant epidermal and cortical cells, induce root cell division and nodule meristem formation, and eventually produce thousands of specialized cellular organelles called “symbiosomes,” each



containing one or more nitrogen-fixing bacteroids. Along the way, differentiation of plant tissues in nodules provides an environment rich in nutrients for the bacteria, with low levels of free oxygen compatible with nitrogen reduction by the oxygen-labile nitrogenase enzyme complex yet sufficient for respiration to energize the process. In essence, SNF is a metabolic symbiosis built upon the trading of reduced carbon from the plant for reduced nitrogen from the bacterial symbionts. The following sections describe and contextualize the genetic discoveries (Figure 2) that have transformed our understanding of how legume-rhizobia symbioses develop.

## EARLY SIGNALING AND PARTNER SELECTION

Legume-rhizobia interactions typically begin in N-limited soils with legumes secreting a class of metabolites called flavonoids (Peters et al., 1986; Redmond et al., 1986). Rhizobia recognize these as signals, which trigger the synthesis and release of lipochitooligosaccharides or nodulation (Nod) factors via rhizobial *nod* genes/proteins (Dénarié and Cullimore, 1993). Nod factors serve as a “calling card” to inform the plant of a potentially beneficial symbiont. Subsequent perception of Nod factors by plant receptors leads to recognition of the symbiont, triggering a series of plant responses that roll out a welcome mat to the bacteria (D’Haeze and Holsters, 2002).

### Flavonoids as Signals to Rhizobia

Flavonoids in legume root exudates act as chemotactic signals to rhizobia under low-N conditions (reviewed in Liu and Murray, 2016). Together with isoflavonoids, they confer host specificity. For example, the isoflavones genistein and diadzein found in root exudates of *G. max* and *P. vulgaris* induce *nod* genes in their compatible rhizobial symbionts, *Bradyrhizobium japonicum* and *Rhizobium leguminosarum* bv *phaseoli*, respectively (Bolaños-Vásquez and Werner, 1997). Host-specific flavonoids are thought to interact with rhizobial NodD protein, a transcription factor (TF) that induces the expression of “common” nod(ulation) genes, leading to the production of lipochitooligosaccharide Nod factors with a conserved core structure (Peters et al., 1986; Lerouge et al., 1990). A variety of side groups produced by enzymes encoded by “host-specific” *nod* genes further contribute to partner selection (Long, 1996).

Suppression of plant flavonoid production by RNA interference (RNAi) of *MtCHS* (*CHALCONE SYNTHASE*) substantially decreases nodulation in *M. truncatula* (Wasson et al., 2006). Knockdown of other flavonoid biosynthetic genes, including *MtCHR* (*CHALCONE REDUCTASE*) and *MtFNS* (*FLAVONE SYNTHASE*) but not *MtIFS* (*ISOFLAVONE SYNTHASE*), inhibits nodulation to varying degrees (Zhang et al., 2009). However, *GmIFS* is required for nodulation in soybean, highlighting the differing requirements of isoflavonoids between legumes (Subramanian et al., 2006).


### Nod Factor Signaling and the Common Symbiotic Pathway

Rhizobial Nod factor is necessary and sufficient to induce nodule development on legume roots, even in the absence of rhizobia (Truchet et al., 1991). Plant symbiotic signaling genes, including Nod factor receptor genes, were among the first SNF genes to be isolated, in part because of their dramatic phenotypes: the

absence of nodulation (Nod<sup>−</sup>) and nitrogen fixation (Fix<sup>−</sup>), poor growth on low-N soils, and defects in Nod factor responses. Analysis of plant Nod<sup>−</sup> mutants that are unable to perceive Nod factors enabled the discovery of Nod factor receptor genes, including the orthologous pairs *LjNFR1* (*NOD FACTOR RECEPTOR*) / *MtLYK3* (*LysM RECEPTOR KINASE*) and *LjNFR5* / *MtNFP* (*NOD FACTOR PERCEPTION*; Limpens et al., 2003; Madsen et al., 2003; Radutoiu et al., 2003; Arrighi et al., 2006; Smit et al., 2007). These genes encode receptor kinases with three extracellular lysin motif (LysM) domains that form homomeric and heteromeric complexes at the plasma membrane and the infection thread (IT) membrane (Figure 3), which bind Nod factors (Haney et al., 2011; Broghammer et al., 2012; Moling et al., 2014). *M. truncatula* transformed with *LjNFR1* and *LjNFR5* can interact with *Mesorhizobium loti*, a natural symbiont of *L. japonicus*, and activate early nodulin gene expression as well as nodule organogenesis, in addition to its natural partner *Sinorhizobium meliloti*, indicating that these receptors are crucial for partner selection/host range determination (Radutoiu et al., 2007). *LjNFR5* also forms heteromers with a plasma membrane leucine-rich repeat receptor-like kinase (LRR-RLK), the symbiotic receptor-like kinase *LjSYMRK* (Antolín-Llovera et al., 2014). *LjSYMRK* / *MtDMI2* were the first “common symbiotic” genes isolated and found to be required for both rhizobial and mycorrhizal symbioses (Endre et al., 2002; Stracke et al., 2002). *Ljsymrk* mutants form no ITs or nodules but have excessive root hair branching responses to Nod factor. Interestingly, *Ljsymrk-14*, a mutant with defects in an essential GPDC tetra-amino acid motif in the extracellular domain of the receptor, supports nodule organogenesis but not infection (Kosuta et al., 2011). Recently, epidermal Nod Factor Receptor (*LjNFR*), another LysM-RLK, was found to phosphorylate *LjNFR5* and support calcium spiking, nodule initiation, and activation of the TF NIN (Schauser et al., 1999; Murakami et al., 2018).

Downstream of these receptors is a signaling pathway that shares several components with the arbuscular mycorrhizal (AM) signaling pathway. This “common symbiosis” signaling pathway has been reviewed extensively (Oldroyd et al., 2011; Oldroyd, 2013; Geurts et al., 2016). Common symbiotic signaling components include a Nod factor-induced E3 ubiquitin ligase called *MtPUB1* (PLANT U-BOX PROTEIN1) that is phosphorylated by the *LjSYMRK* ortholog *MtDMI2* as well as the entry receptor *MtLYK3* (Mbengue et al., 2010; Vernié et al., 2016). *Mtpub1-1* mutants have increased numbers of nodules and infection events, indicating that *MtPUB1* is a negative regulator of nodulation (Vernié et al., 2016). Silencing of *MtPUB1* allowed efficient nodulation by an *S. meliloti* *nodFnodL* mutant strain that produced altered Nod factors and poorly nodulated wild-type plants, suggesting that *MtPUB1* plays a role in Nod factor discrimination (Mbengue et al., 2010). Mutants affecting three other SYMRK INTERACTING PROTEINS (SIPs) that presumably act in early signaling, *LjSIP1*, a mitogen-activated protein kinase kinase, *LjSIP2*, an ARID domain-containing protein, and SYMRK INTERACTING E3 UBIQUITIN LIGASE (*LjSIE3*), are defective in nodule organogenesis (Zhu et al., 2008; Chen et al., 2012; Yuan et al., 2012; Wang et al., 2013).

The perception of Nod factors leads to depolarization of cell membranes and changes in ion fluxes. These changes include oscillations in calcium concentrations called “calcium spiking” in the nuclei of epidermal root hair cells, which appear to drive



**Figure 3.** Genes and Processes Involved in Early Signaling during Nodulation.

(Iso)flavonoids produced under low soil N (1) trigger the production of bacterial Nod factors (2) that, together with other signals, are perceived by receptors at the plasma membrane of epidermal cells (3). This triggers biochemical and physiological responses (4–6) that lead to changes in nuclear gene expression (7). See main text for key to gene/protein names and further explanation of early signaling pathways and components. *M. truncatula* protein names are provided unless otherwise specified.

changes in gene expression associated with nodule development and infection (Figure 3; Charpentier and Oldroyd, 2013). This is followed by deformation of root hairs, early nodulin gene expression, and, subsequently, the formation of nodule primordia. Nod- mutants defective in calcium spiking were used to identify various nuclear envelope proteins that help generate this signal, including the calcium channels LjCASTOR and LjPOLLUX/MtDMI1 (Ané, 2004; Imaizumi-Anraku, 2005; Charpentier et al., 2008; Kim et al., 2019); nucleoporin subunits LjNUCLEOPORIN85 (LjNUP85) and LjNUP133 (Kanamori et al., 2006; Saito et al., 2007); and calcium channels MtCNGC (CYCLIC NUCLEOTIDE GATED CHANNELS) a/b/c (Figure 3; Charpentier et al., 2016). Interestingly, MtDMI1 seems to serve the purposes of both LjCASTOR and LjPOLLUX, showing how evolution has come up with different solutions to the same problem in different species (Supplemental Data Set; Ané et al., 2004; Charpentier et al., 2008). Mutations in the nucleoporin-localized protein LjNENA lead to defects in infection by both rhizobial and mycorrhizal symbionts. *Ljnen*a mutants do not exhibit calcium spiking in response to Nod factor and show clear defects in rhizobial infections without any changes

in nodule organogenesis (Groth et al., 2010). However, nodules that form on *Ljnen*a are empty, with all infections blocked at the microcolony stage. An unknown calcium pump and the known calcium channels (CNGCa/b/c) are believed to work in concert to generate nuclear calcium spiking (Charpentier et al., 2016). A gain-of-function mutation in *LjBRUSH*, another CNGC gene, impairs rhizobial infection but only at high temperatures (Maekawa-Yoshikawa et al., 2009; Chiasson et al., 2017). Analysis of other Nod- mutants revealed that the nuclear calcium-calmodulin kinase MtDMI3/CCaMK acts as an intermediary between Nod factor perception and nodule development (Gleason et al., 2006; Tirichine et al., 2006), presumably as a sensor of the signal encoded in nuclear calcium spiking. The TF LjCYCLOPS/INTERACTING PROTEIN OF DMI3 (MtIPD3) regulates gene expression, leading to nodulation (Yano et al., 2008; Singh et al., 2014). Importantly, this TF can be phosphorylated by CCaMK, apparently tying calcium spiking to gene transcription (Figure 3; Singh et al., 2014). Two other TFs, MtNSP1 and MtNSP2 (NODULATION SIGNALING PATHWAY), of the GIBBERELIC ACID INSENSITIVE (GAI), REPRESSOR OF GAI, and SCARECROW (GRAS) family were also discovered from


forward genetics analysis of Nod<sup>-</sup> mutants (Kaló et al., 2005; Smit et al., 2005). NSP1 and NSP2 interact to form heterodimers that are required for binding to and activating Nod factor-responsive genes (Hirsch et al., 2009).

Downstream of the Nod factor signaling pathway, paralogous Nod factor hydrolases MtNFH1 (NOD FACTOR HYDROLASE1) and LjCHIT5 (CHITINASE5) cleave the lipochitooligosaccharidic backbone of Nod factors produced by their respective symbionts (Figure 3). This degradation of Nod factors appears to be crucial for rhizobial infection, colonization, and nodule formation (Cai et al., 2018; Malolepszy et al., 2018). Interestingly, delays in nodule formation by aberrant Nod factor-producing strains were abolished in *Ljchit5* mutants, implicating Nod factor hydrolysis in partner recognition (Malolepszy et al., 2018).

### Other Signaling Pathways

In addition to Nod factor, bacterial surface exopolysaccharides (EPS) are recognized by plant plasma membrane receptors (Figure 3). The exopolysaccharide receptor LJEP3 (EXOPOLYSACCHARIDE RECEPTOR) is a LysM domain-containing entry receptor that is dispensable for early responses to Nod factor but is required for infection progression beyond the microcolony stage (Kawaharada et al., 2015). *LjEP3* is differentially regulated in the epidermis and the nodule primordia (Kawaharada et al., 2017b). An *Ljep3* LORE1-insertion mutant identified in a suppressor screen with the EPS-defective *M. loti* *exoU* mutant overcame the strain incompatibility and formed infrequent but mature pink nodules in addition to some small, uncolonized nodule primordia like those that form on wild-type plants (Kawaharada et al., 2015). Further studies are required to uncover components downstream of this receptor and understand how EPS signaling complements Nod factor signaling in rhizobial partner selection.

Several hormone signaling pathways crosstalk with Nod factor signaling to control nodulation and SNF at various stages (López-Ráez et al., 2017; Liu et al., 2018). *LjCCD7* (CAROTENOID CLEAVAGE DIOXYGENASE7), encoding an enzyme required for strigolactone biosynthesis, and *MtBRI1* (BRASSINOSTEROID INSENSITIVE1), encoding an LRR-RLK that perceives the brassinosteroid hormone brassinolide, both act early during nodulation (Liu et al., 2013; Cheng et al., 2017). Silencing of either gene has a mild negative effect on the number of nodules formed. Analysis of *Tnt1* insertion mutants in *MtDELLA1*, *MtDELLA2*, and *MtDELLA3*, encoding negative regulators of gibberellic acid signaling, revealed that these DELLA proteins control Nod factor-induced gene expression and rhizobial infection in the epidermis as well as nodule number and density (Fonouni-Farde et al., 2016; Liu and Murray, 2016). DELLAs appear to do this by interacting directly with the TFs MtIPD3 and MtNSP2 (Fonouni-Farde et al., 2016; Jin et al., 2016). Cytokinin orchestrates nodule initiation and development together with auxin, which also appears to play roles in the progression of rhizobial infection (Figure 4; see Organogenesis). Lastly, the knockdown of *MtHMGR1* (3-HYDROXY-3-METHYLGLUTARYL COA REDUCTASE1), encoding an enzyme involved in the production of isoprenoid compounds in the mevalonate pathway that likely interacts with MtDMI2, suppressed nodule formation (Kevei et al., 2007). It remains unclear how this fits into early signaling pathways leading to nodulation.



**Figure 4.** Rhizobial Infection.

The entry of rhizobia into the plant cell via tubular ITs triggers a transient ROS burst (1). This is accompanied by cell wall degradation (2), membrane remodeling (3), and cytoskeletal rearrangements (4). The IT is led by a complex at its tip called the infectosome (5). In parallel, several other transcriptional and posttranslational changes (6) and the autophagy pathway (7) ensure accommodation of the symbiont within the plant host.

### PLANT IMMUNITY AND HOST RANGE RESTRICTION

In parallel with Nod factor-dependent partner selection, the plant immune system helps exclude other soil microorganisms from legume roots (Zipfel and Oldroyd, 2017). Partner selection is a continuous process. The host legume employs multiple checkpoints throughout this process to discriminate between symbionts and pathogens. Defensive receptor kinase complexes including LRR-RLKs and LysM-RLKs recognize microbial molecules at the surfaces of plant cells, while microbial “effectors” injected into cells to disarm plant defenses are recognized and neutralized by NBS-LRR (nucleotide binding site leucine-rich repeat) R (Resistance) proteins (Cao et al., 2017). Consequently, the immune system limits the number of rhizobial strains that a legume can associate with. Genes for “host-range restriction” include the NBS-LRR-encoding *Rj2/Rfg1* (*Rhizobium japonicum2/Rhizobium fast-growing1*) and *Rj4*, encoding a member of the pathogenesis-related protein family 5 (Yang et al., 2010; Tang et al., 2016). *Rj2*-carrying cultivars of *G. max* form active N-fixing symbioses with many strains of *Bradyrhizobium* but not with *B. japonicum* USDA 122. *Rfg1* restricts symbiosis with certain *Sinorhizobium fredii* strains (Fan et al., 2017). *Rj2* and *Rfg1* are

alleles of a TIR-NBS-LRR gene, with variant nucleotides affecting seven amino acids (Yang et al., 2010). Rj2 recognizes the effector protein NopP secreted by the incompatible strain *B. japonicum* USDA 122 type III secretion system, which triggers the plant immune system and blocks rhizobial infection (Sugawara et al., 2018). Rj4 is a dominant allele encoding a thaumatin-like protein, which restricts nodulation with ineffective strains of *B. japonicum* and *Bradyrhizobium elkanii* (Tang et al., 2016). Soybean cultivars with the alternate allele, *rj4*, are nodulated by many ineffective strains of *B. elkanii* (Sadovsky and Cregan, 1992). The restriction of host range helps ensure effective SNF. Editing of Rj4 using CRISPR/Cas9 led to interactions with the incompatible rhizobial strain, *B. japonicum* Is-34 (Hayashi et al., 2014; Tang et al., 2016). Both Rj2 and Rj4 plants have fewer infections with incompatible rhizobial strains compared with *rj2* and *rj4* plants. The nodules that occasionally form are developmentally arrested, with only a few cortical divisions and signs of defense responses (Yang et al., 2010; Hayashi et al., 2014). It remains unknown which bacterial molecule is recognized by Rj4.

Nodule cysteine-rich (NCR) peptides also affect partner selection. Editing of two NCR genes in *M. truncatula* ecotype Jemalong A17 allowed colonization by *Sinorhizobium meliloti* strain Rm41, normally a poor nodulator of A17. NCR peptide-encoding loci *MtNFS1* (NITROGEN FIXATION SPECIFICITY) and *MtNFS2* were identified in a recombinant inbred line population derived from ecotypes A17 and DZA315 (Wang et al., 2017; Yang et al., 2017). These functionally dominant peptides ensure that A17 plants are colonized preferentially by efficient N-fixing symbionts.

Interestingly, OsCERK1 (CHITIN ELICITOR RECEPTOR KINASE1), a rice (*Oryza sativa*) LysM-RLK that recognizes chitin oligosaccharides of fungal pathogens, also perceives signals from beneficial AM fungi (Miyata et al., 2014; Zhang et al., 2015), suggesting that some symbiotic receptors might participate in receptor complexes that recognize pathogens. This is an exciting, emerging area of research.

## RHIZOBIAL INFECTION

Attachment of rhizobia to root hair cells leads to root hair curling, which “traps” rhizobia in so-called infection pockets that eventually contain microcolonies of dividing rhizobia. The most common route of rhizobial entry into root cells in the four legumes under discussion is via tubular ITs in epidermal root hair cells, although rhizobia are also known to enter via cracks on the root surface (Sprent and James, 2007). Both modes of entry are Nod factor dependent. However, rhizobia can also enter legume roots via intercellular spaces between cells in a Nod factor-independent manner (reviewed by Oldroyd et al., 2011). ITs originate at infection pockets as the plant cell wall is deconstructed, and the adjacent plasma membrane invaginates to allow the entry of multiplying rhizobia. Simultaneous plant cell divisions in the root cortex and pericycle trigger nodule organogenesis in cells undergoing successful infections (Xiao et al., 2014).

## Hormonal Regulation

One of the first plant mutants identified with defective rhizobial infection was *Mtskl* (*sickle*). This mutant is impaired in EIN2

(ETHYLENE INSENSITIVE2), an endoplasmic reticulum-tethered protein crucial for ethylene signaling (Penmetsa and Cook, 1997; Penmetsa et al., 2003). The hormone ethylene coordinates the biosynthesis and signaling of several other phytohormones at multiple stages of SNF while also modulating defense pathways (Larrainzar et al., 2015). This central role for ethylene is exemplified by the pleiotropic phenotypes caused by mutated EIN2 in *Mtskl* plants. *Mtskl* exhibits massive proliferation of ITs and nodule primordia, although these rarely develop into mature nodules. This role of EIN2 is conserved in *L. japonicus* but requires two copies of *EIN2* such that the double mutant *ein2a ein2b* does not fix nitrogen (Miyata et al., 2013; Reid et al., 2018). Not only is *Mtskl* hypersensitive to beneficial bacteria but it is also hypersusceptible to infection by fungal pathogens, suggesting that ethylene plays a crucial role in fine-tuning interactions with microbes (Penmetsa et al., 2008).

Auxin positively regulates rhizobial infection while gibberellic acid appears to be a negative regulator of this process. A mutation in *MtARF16a* (*AUXIN RESPONSE FACTOR*), which is required for IT elongation but not organogenesis, implicates auxin in the infection process (Breakspear et al., 2014). Mutations in *MtARF16a* reduced the number of infection events, although a few productive ITs resulted in nodule colonization (Breakspear et al., 2014). The biosynthesis and degradation of gibberellic acid play distinct roles in SNF in addition to signaling mediated by DELLA proteins (see Early Signaling and Partner Selection). New research indicates that the catabolism of gibberellic acid by MtGA2ox10 (GIBBERELIC ACID OXIDASE10) plays a positive role in infection (Kim et al., 2019), which is in line with the negative correlation between both gibberellic acid and DELLA-mediated signaling and rhizobial infection (Fonouni-Farde et al., 2016; McAdam et al., 2018). Targeted editing that disrupted *MtGA2ox10* reduced the number of infections and the number and size of nodules formed on hairy roots. Interestingly, *MtGA2ox10* is not only induced in the maturation zone of the root in response to infection by rhizobia but also in the meristem, infection, and fixation zones of mature nodules, suggesting that it plays a role in later stages of nodule functioning (Kim et al., 2019).

## Role of Plant Cytoskeleton

Many components of the cytoskeleton are required for reorientation of root hair tip growth and IT development (Figure 4; Timmers, 2008). Mutations in *LjNap1* (*Nck-associated protein1*) and *LjPir1* (*121F-specific p53 inducible RNA*), encoding components of the SCAR/WAVE complex, affect rhizobial infection but not nodule organogenesis (Yokota et al., 2009). SCAR/WAVE proteins drive actin assembly and consequently control cell growth by initiating the formation of actin complexes from which filaments can extend. *Ljnap1* and *Ljpir1* mutants initiate fewer infections, which are also delayed compared with the wild type (Yokota et al., 2009). Like the *Ljnap1* and *Ljpir1* mutants, most ITs that form on *Ljscarn* (*scar-nodulation*) mutants abort prematurely and release bacteria within the root hair (Qiu et al., 2015). Epidermally expressed *LjSCARN* is required for actin nucleation without which small, white nodules generally form (Qiu et al., 2015). SCAR proteins bind to and activate the Actin-Related Protein2/3 (ARP2/3) complex, a subunit of which, *LjARPC1* (ACTIN-RELATED PROTEIN COMPONENT1), is also required for infection. *Ljarp1* mutants produce few microcolonies and ITs,

which abort/disintegrate prematurely (Hossain et al., 2012; Qiu et al., 2015), resulting in mostly uncolonized nodules. Interestingly, in addition to abnormalities in root hair growth due to actin rearrangements, a knockdown of *MtCDPK1* (*CALCIUM DEPENDENT PROTEIN KINASE1*) resulted in constitutive root hair bending and branching, even in the absence of rhizobia. Only 50% of all incipient infections on *CDPK* RNAi lines were successful, and nodules rarely formed (Ivashuta et al., 2005).

### Cell Wall and Cell Membrane Components

IT formation and extension require the plant cell wall to degrade to allow entry of rhizobia (van Spronsen et al., 1994). Consistent with this notion, the mutation of the pectate lyase-encoding gene *LjNPL* (*NODULE PECTATE LYASE*) resulted in a drastic reduction in the number of ITs in *L. japonicus* roots (Xie et al., 2012). Purified *LjNPL* protein degrades both polygalacturonic acids and pectin. Like other mutants defective in rhizobial infection, *Ljnp1* mostly produced small, white, uninfected nodules (Xie et al., 2012; Liu et al., 2019b).

Membrane trafficking proteins contribute to the development of the IT as the plant plasma membrane invaginates and extends inward (Figure 4). Studies on other integral membrane proteins, including *MtFLOTILLIN2* (*MtFLOT2*), *MtFLOT4*, and *MtSYMREM1* (*SYMBIOTIC REMORIN1*), demonstrated the importance of membrane processes during infection progression (Haney and Long, 2010; Lefebvre et al., 2010). *FLOTILLINS* are lipid raft markers present in membrane microdomains (Glebov et al., 2006). *MtFLOT4* is a nodulin that accumulates at the tips of root hairs upon inoculation with rhizobia (Winzer et al., 1999). *Mtflot4* RNAi resulted in fewer initiating and elongating ITs, which were more likely to abort in root hairs compared with the wild type (Haney and Long, 2010). Like *Mtflot4*, knockdown of *Mtflot2* by RNAi resulted in fewer ITs and nodules, while *Mtflot2* mutations are lethal in the homozygous state, indicating that *Mtflot2* plays a broader, essential role in the plant (Liang et al., 2018). Together, *FLOTILLINS* are thought to play a role in invagination of ITs into root hairs, with *MtFLOT4* playing a specialized role in IT elongation (Haney and Long, 2010). *MtFLOT4* colocalizes to membrane nanodomains within the root hair cells with the symbiotic remorin protein, *MtSYMREM1*. Upon rhizobial inoculation, *MtSYMREM1* and *MtFLOT4* interact with the plant-encoded bacterial entry receptor *MtLYK3* to mediate endocytosis of the receptor protein and allow infection progression (Lefebvre et al., 2010; Haney et al., 2011). The density of *MtSYMREM1* on root epidermal cells is reduced sixfold in *Mtflot4* mutants, suggesting that *MtFLOT4* is required for the recruitment of *MtSYMREM1*. Like *Mtflot4*, *Mtsymrem1* mutants have increased numbers of infection foci but very few elongating ITs (Haney and Long, 2010; Lefebvre et al., 2010; Liang et al., 2018). Reduced expression levels of *MtFLOT4* in *MtN5* (*NODULIN5*) RNAi lines could explain the infection defects observed by silencing this nonsulfated lipid transfer protein gene (Pii et al., 2012).

### The Infectosome

Recent research points to the existence of an exocyst complex at the tip of the IT, termed the “infectosome,” which ensures polar growth of the IT toward the nodule cortex (Figure 4; Liu et al., 2019a). The infectosome comprises the common symbiotic

protein *MtVpy* (*MtVAPYRIN*) and its two interacting partners, *MtLIN* (*LUMPY INFECTION*) and the exocyst complex subunit *MtEXO70 H4* (*EXOCYST* subunit H4; (Pumplin et al., 2010; Murray, 2011; Liu et al., 2019a). Mutating any of the three components causes defects in IT elongation (Liu et al., 2019c). Both *Mtvp1* and *Mtlin* mutants produce few nodules, which are small, white, and uninfected, while *Mtexo70 h4* mutants form normal, functional nodules, possibly reflecting genetic redundancy (Murray, 2011; Liu et al., 2019a).

### The Autophagy Pathway

Components of the autophagy pathway play positive roles in IT formation, such as a common symbiosis kinase of the phosphatidylinositol 3-kinase (PI3K) family identified using RNAi in *P. vulgaris*. *PvPI3K* is expressed in root hairs, and its expression is further enhanced by infection with *Rhizobium tropici* (Estrada-Navarrete et al., 2016). Downregulating *PvPI3K* decreased root hair length and the frequency with which root hairs responded to rhizobia. Most infections were arrested at the bases of root hairs. The resulting nodules were devoid of intracellular infections and unable to fix nitrogen. Knockdown of *PvPI3K* suppressed multiple autophagy genes, including *PvBECLIN*. Independent knockdown of *PvBECLIN* and another autophagy-related kinase gene, *PvTOR* (*TARGET OF RAPAMYCIN*), phenocopied the *PvPI3K*-silenced lines and led to a drastic reduction in nodule number (Estrada-Navarrete et al., 2016; Nanjareddy et al., 2016).

### Reactive Oxygen Species

Nod factor-triggered reactive oxygen species (ROS) production is correlated with the induction of the nodulin gene *RIP1* (*RHIZOBIUM INDUCED PEROXIDASE1*) in *M. truncatula* and is dependent on *DMI1* (Ramu et al., 2002). RNAi uncovered a positive role for the transient ROS burst during rhizobial infection. Silencing of the protein kinase gene *MtSPK1* (*SYMBIOTIC PROTEIN KINASE1*), *LjROP6* (*Rho family of small GTPases in plants*), and two non-orthologous NADPH oxidase genes, *PvRBOHa* and *MtRBOHa* (for *RESPIRATORY BURST OXIDASE HOMOLOGS*), affected SNF to varying degrees. *MtSPK1* expression is induced by both Nod factor and H<sub>2</sub>O<sub>2</sub>, primarily in infected root hair cells and the infection zones of nodules (Andrio et al., 2013). Artificial microRNA *Mtspk1* lines had fewer nodules than the wild type. *L. japonicus* *ROP6* RNAi lines had fewer ITs in the root epidermis, cortex, and nodule primordia and fewer nodules than wild-type plants (Ke et al., 2012). Interfering with NADPH oxidase-catalyzed ROS production by silencing either *MtRBOHa* or *PvRBOHa* strongly suppressed SNF, underlining a positive role for ROS in SNF (Marino et al., 2011; Arthikala et al., 2017).

### The Roles of Transcription and Posttranslational Machinery in Rhizobial Infection

Components of the general transcriptional machinery have been implicated in the infection process, including *LjLAN* (*LACK OF SYMBIONT ACCOMMODATION*), encoding a putative component of a mediator complex required as a cofactor for the recruitment of RNA polymerase II (Suzaki et al., 2019). The EMS-induced *Ljlan*



mutant was Nod<sup>-</sup> 2 weeks after inoculation, although intracellular crack entry of rhizobia led to nodulation during later stages. Analysis of nodule sections confirmed the presence of bacterial clusters in the absence of any cortical ITs. Curiously, LORE1-insertion mutants and CRISPR/Cas9 mutations in the C-terminal domain of *LjLAN* did not reproduce this phenotype, indicating that this domain is dispensable for its function (Suzaki et al., 2019). Silencing *PvAGO5*, encoding a scaffolding Argonaute protein of the RNA-silencing machinery, reduced root hair bending in response to rhizobia and resulted in fewer, smaller nodules (Reyero-Saavedra et al., 2017).

### Ancillary Genes

Several genes that support rhizobial infection have been identified via genetics but cannot be placed into any particular process/pathway. Defects in *MtRPG* (*RHIZOBIUM DIRECTED POLAR GROWTH*) result in abnormally thick cell IT walls and impaired infection, although early Nod factor responses are normal (Arrighi et al., 2008). By contrast, a *Tnt1* insertion in *MtCBS1* (*Cystathionine-β-Synthase-like Domain-Containing*) resulted in more microcolonies and fewer ITs but normal IT wall thickness (Sinharoy et al., 2016).

Infection and organogenesis are interdependent but genetically separable processes; this often complicates the identification of mutants affecting either process alone. For example, the expression of *LjnsRING* (*nodule-specific REALLY INTERESTING NEW GENE*), a component of the ubiquitination pathway, correlates with bacterial infection into nodule cells but ceases as nodule development progresses (Shimomura et al., 2006). *LjnsRING* transcript levels are significantly reduced in nonfunctional nodules induced by a *Fix-ΔnifH M. loti* mutant. Yet, not only do a limited number of ITs form on *LjnsRING* RNAi lines, but nodule development is also severely affected, with some transgenic roots infrequently forming small bumps (Shimomura et al., 2006). Genetics has uncovered many other genes required for infection; see Figure 2 and the Supplemental Data Set for more information.

### ORGANOGENESIS

Nodule organogenesis is driven by three distinctive processes that occur simultaneously: bacterial colonization, nodule inception, and autoregulation of nodule number. Legumes such as *M. truncatula* have persistent nodule meristems, which generate elongated, sometimes multi-lobed “indeterminate” nodules, whereas “determinate,” mostly spherical nodules are formed by transiently active meristems in the nodules of species such as *L. japonicus*, *P. vulgaris*, and *G. max*.

#### Cell Division and Nodule Inception


As root hairs curl, entrap compatible rhizobia, and develop ITs for bacterial transport into root tissues, root pericycle and cortical cells reenter the cell cycle and start dividing, which initiates nodule primordia formation and vascularization (Oldroyd et al., 2011; Guan et al., 2013). Although organogenesis of both indeterminate and determinate types of nodules requires the initiation of cell division in the pericycle, indeterminate nodules emerge from the inner cortex, while determinate nodules emerge from the middle and outer cortex (Hadri et al., 1998). Nodule tissues consist of

outer cortex, cortical endodermis, vascular bundles, and inner cortex, which encase the central, mostly infected region. In mature indeterminate nodules, this region is organized into developmental zones: the meristem (zone I), infection zone (zone II), nitrogen fixation zone (zone III), and senescence zone (zone IV). In mature determinate nodules, the central region consists only of nitrogen fixation and, eventually, senescence zones (Xiao et al., 2014). Development of the various tissues and zones of nodules results from successive anticlinal and periclinal divisions in the pericycle, endodermis, and C3 to C5 cortical layers (Xiao et al., 2014). For functional nodules, it is important that some meristematic cells differentiate into large, polyploid cells, which will accommodate the symbionts. Endoreduplication involves DNA replication without cell division, a process facilitated by ubiquitin-dependent proteolysis of mitotic cyclins. *MtCCS52A*, encoding an anaphase-promoting complex activator, controls endoreduplication during infected cell differentiation in indeterminate nodules. *Mtccs52a* mutant nodule cells remain small with low ploidy, leading to symbiont and host cell death (Cebolla et al., 1999; Vinardell et al., 2003). The fate of nodule meristem cells was explored by downregulating *MtPLETHORA* (*MtPLT*) genes using an RNAi approach. Distinct domains were identified in which *MtPLT1* to *MtPLT4* genes control differentiation into either cells of the central region (*MtPLT3* and *MtPLT4*) or cells of the nodule periphery (*MtPLT1* and *MtPLT2*; Franssen et al., 2015). Nodule morphogenesis is promoted by class 2 KNOX homeobox KNAT3/4/5-like TFs. RNAi silencing of *MtKNAT3/4/5-like* genes resulted in fused nodules, although a number of nodules and infection events were not affected (Di Giacomo et al., 2017). Although infection and nodule organogenesis are distinct, genetically separate programs, proper nodule development is dependent on rhizobial infection (Voroshilova et al., 2009). Analysis of *lin-4*, a weak mutant allele of the *MtLIN* gene, revealed that normal nodule development requires the initiation of ITs. Moreover, *lin-4* nodules that develop 2 to 3 months postinoculation have centrally located vascular bundles. This is reminiscent of nodule formation on N-fixing actinorrhizal plants, which nodulate in association with actinobacteria called *Frankia* (Kuppusamy et al., 2004; Kiss et al., 2009; Guan et al., 2013). Maintenance of root nodule identity is controlled by the *M. truncatula* *NODULE ROOT* (*MtNOOT*) gene, an ortholog of the Arabidopsis (*Arabidopsis thaliana*) *BLADE-ON-PETIOLE* transcriptional activator gene (Couzigou et al., 2012). In *noot* mutants, roots emerge from the nodule meristem, suggesting that these genes were recruited to repress root identity in nodules.

#### Transcriptional Control of Nodule Organogenesis

The very first symbiotic gene to be discovered, named *Nodule Inception* (*NIN*), was predicted to be a TF gene in *L. japonicus* (Schauer et al., 1999). A transposon-tagged *nin* mutant was defective in bacterial recognition, displaying excessive root hair curling without infection or nodulation (Schauer et al., 1999). *NIN*, which encodes an RWP-RK-containing TF, is a master regulator of multiple genes/processes that have positive or negative effects on nodule organogenesis, depending on the developmental time point, including IT initiation, nodule organogenesis, and the control of nodule number (Figure 4; Yano, 2008; Soyano et al., 2013, 2014; Liu, 2019b).

*NIN* triggers IT initiation by inducing the *LjNPL* (see Cell Wall and Cell Membrane Components) and *nuclear transcription factor Y*



**Figure 5.** TFs Involved in Nodule Organogenesis.

Cell divisions opposite protoxylem poles initiate nodules. Cytokinin signaling downstream of the CRE1 receptor activates the GRAS domain TFs NSP1 and NSP2, leading to the activation of NIN. LjCYCLOPS/IPD3 bind to the promoter of *NIN*, leading to the activation of downstream TFs NF-YA1, ERN1, and ERN2. Medicago gene names are displayed unless otherwise indicated.

*subunit A-1 (MtNF-YA1)* genes (Figure 5; Xie et al., 2012; Laporte et al., 2014). In response to  $Ca^{2+}$  oscillations triggered by Nod factor signaling, the calcium/calmodulin-dependent protein kinase LjCCaMK/MtDMI3 phosphorylates LjCYCLOPS/MtIPD3, which activates *NIN* expression, leading to IT formation and nodule inception (Yano et al., 2008; Figures 1 and 5). A functionally redundant *IPD3-like (MtIPD3L)* gene was recently found to act downstream of *MtDMI3* to control nodule organogenesis (Jin et al., 2018). *NIN* expression is also induced by the NSP1 GRAS-type TF, as part of a heterodimer with NSP2 (Hirsch et al., 2009).

IT formation and nodule organogenesis in Medicago are dependent on *Ethylene Response Factor Required for Nodulation1 (ERN1)*, which encodes an APETALA2/Ethylene Responsive Factor (AP2/ERF) TF. *Mtern1* mutants form infection pockets with limited cortical cell divisions that produce empty nodule “bumps” (Middleton et al., 2007). Similar to *NIN*, *ERN1* is induced by the CcCaMK/CYCLOPS complex (Cerri et al., 2017; Kawaharada et al., 2017a) and controls the expression of cell wall-associated *ENOD11* and *ENOD12*, which are critical for IT development (Andriankaja et al., 2007). When *MtERN2*, a close homolog of *MtERN1*, is disrupted, functional nodules develop, but they senesce early (Cerri et al., 2016). Since an *ern1-1 ern2-1* double mutant was found to be devoid of both infection and nodule organogenesis, it was postulated that, although *ERN1* and *ERN2* work together in the root epidermis, only *ERN1* is required for nodule organogenesis. The induction of *LjERN1* by Nod factor is dependent on the Nod factor receptor LjNFR1 and the TFs LjNSP2 and LjCYCLOPS, but not LjNIN or LjNF-YA1, in the epidermis


(Kawaharada et al., 2017b). Interestingly, MtNIN controls the extent of infection by limiting the expression of *MtENOD11 (EARLY NODULIN1)* in the root epidermis, thus acting as an *ERN1* antagonist (Vernié et al., 2015). Analysis of a weak *nin* allele termed *daphne-like*, with an insertion 7 kb upstream of the NIN start site, further suggested that LjNIN plays a negative role in rhizobial infection but not organogenesis (Yoro et al., 2014).

Like NIN, the Nuclear Factor Y complex (a CCAAT-box binding heterotrimeric TF complex consisting of NF-YA, NF-YB, and NF-YC) controls multiple steps of nodule organogenesis. MtNF-YA1 is critical for nodule meristem initiation and persistence (Comber et al., 2006, 2008). Furthermore, LjNF-YA1 and LjNF-YB1 promote cortical cell division in an LjNIN-dependent manner (Soyano et al., 2013). In *P. vulgaris*, the NF-Y heterotrimer of PvNF-YAa, PvNF-YB7, and PvNF-YC1 is required not only for infection and organogenesis, but it also plays a role in symbiotic partner selection for improved nitrogen fixation efficiency (Zanetti et al., 2010; Rípodas et al., 2019). The GRAS domain TF PvSIN1 (Scarecrow-like13 Involved in Nodulation) interacts with PvNF-YC1 to control nodule number, size, and IT formation (Figure 5; Battaglia et al., 2014).

### Hormonal Regulation

Hormones have long been known to control nodule organogenesis (Figure 6; Grunewald et al., 2009). The concentrations and ratios of auxins and cytokinins determine where and when cells divide. Cytokinins regulate cell division leading to nodule primordium formation, and their signals are mainly perceived by the cytokinin receptor, MtCRE1/LjLHK1 (Gonzalez-Rizzo et al., 2006; Murray et al., 2007; Tirichine et al., 2007). The cytokinin receptors LjLHK1A and LjLHK3 share partially redundant functions with LjLHK1 during nodule development (Held et al., 2014). Direct targets of cytokinin signaling include *MtNSP2* and the basic helix-loop-helix TF gene *MtbHLH476*, both encoding positive regulators of nodule organogenesis (Ariel et al., 2012). Genes for cytokinin biosynthesis and homeostasis are required for normal nodule development, including *LjIPT3*, *LjCKX3*, and *MtLOG1*. Cytokinin biosynthesis is mediated by isopentenyl transferase (encoded by *LjIPT3*), and cytokinin homeostasis during nodule development is maintained by cytokinin oxidase/dehydrogenase3 (encoded by *LjCKX3*) in *L. japonicus* and the cytokinin-activating enzyme riboside 5'-monophosphate phosphoribohydrolase (encoded by the LONELY GUY gene *MtLOG1*) in *M. truncatula* (Chen et al., 2014; Mortier et al., 2014; Reid et al., 2016). The involvement of the key TF NIN in cytokinin signaling during nodule organogenesis was recently demonstrated by Liu et al., 2019c. A careful analysis of the *NIN* promoter revealed that a 2.2-kb segment can reverse the excessive root hair curling and restore infection pocket formation, a 5-kb segment can restore IT formation, but only the presence of 10 cytokinin-responsive promoter elements 18 kb upstream of the NIN translation site can restore nodule organogenesis.

Although auxin response genes are induced at the site of initiation of both determinate and indeterminate nodules, the inhibition of polar auxin transport is crucial for indeterminate but not determinate nodule development (Figure 6; Ng and Mathesius, 2018). Medicago PINFORMED (PIN) auxin exporters MtPIN2, MtPIN3, and MtPIN4 and the auxin influx carrier MtLAX2 (LIKE AUXIN2) promote nodule organogenesis (Huo et al., 2006; Roy



**Figure 6.** Hormonal Control of Symbiotic Nitrogen Fixation.

Ethylene (ET) represses calcium spiking induced upon recognition of a compatible Nod factor (1). Rhizobial infection is countered by the hormones ET, jasmonic acid (JA), and gibberellic acid (GA). By contrast, the hormone cytokinin (CK; whose biosynthesis is mediated by IPT3) and auxin (AUX) signaling (via ARF16a) positively regulate IT development (2 and 3). Multiple hormones control cell divisions at the site of infection that lead to the formation of the nodule primordium (4). As the primordium develops into a mature nodule, hormone requirements differ between indeterminate and determinate nodules (5 and 6). Positive and negative roles are shown in green and magenta, respectively. Medicago gene names are shown unless otherwise indicated.

et al., 2017). Auxin-mediated organogenesis of determinate nodules also involves the microRNA *miR167* and the auxin response factors GmARF8a and GmARF8b in soybean. *miR167* acts upstream of NIN, NSP1, NF-YA1, NF-YA2, and ENOD40 and positively regulates nodule formation by inhibiting *GmARF8a* and *GmARF8b* expression (Wang et al., 2015). Another microRNA, *miR160*, plays a crucial role in maintaining the auxin/cytokinin balance during soybean nodule inception and maturation (Nizampatnam et al., 2015). Auxin-induced transcriptional regulation of nodule development is also regulated by the basic helix-loop-helix TF MtbHLH1 (Godiard et al., 2011). Finally, *M. truncatula* CELL DIVISION CYCLE16 (MtCDC16), a presumed component of the anaphase-promoting complex, participates in auxin-dependent nodule organogenesis. The *cdc16* mutant has fewer lateral roots and slightly more nodules than the wild type but shows an auxin-resistant root elongation phenotype (Kuppusamy et al., 2009).

Carotenoid cleavage products, including two plant hormones, strigolactone and abscisic acid, also play roles in nodule organogenesis. The  $\beta$ -carotene hydroxylases GmBCH1 and GmBCH2 catalyze the conversion of  $\beta$ -carotene to  $\beta$ -zeaxanthin in soybean. RNAi silencing of *GmBCH1/2* impaired nodule development and SNF (Kim et al., 2013). Silencing of *P. vulgaris* BYPASS1 (*PvBPS1*) resulted in defective nodule development, which was partially rescued by the carotenoid biosynthesis inhibitor fluridone (Arthikala et al., 2018).

#### Additional Genes Required for Nodule Organogenesis

Defects in *MtNPF1.7* of the Nitrate and Peptide Transporter Family account for the mutant phenotypes of *Mtnip-1*, *Mtnip-3*, and *Mtlatd* (numerous infections and polyphenolics/lateral root-organ defective; Yendrek et al., 2010). These mutants have nodules and lateral roots that lack a persistent meristem, with the more severe


*Mtnip-1* and *Mtlatd* mutants showing rhizobia trapped within ITs (Veereshlingam et al., 2004; Teillet et al., 2008). Although MtNPF1.7 transported nitrate in *Xenopus laevis* oocytes at high affinity (Bagchi et al., 2012), the *in vivo* substrate and precise role of MtNPF1.7 remain unclear, given that some NPFs also transport hormones (Zhang et al., 2014).

Nodule organogenesis requires other genes involved in cellular events and processes in addition to those described above. Refer to Figure 2 and the Supplemental Data Set for more information on these genes.

#### AUTOREGULATION OF NODULATION

Autoregulation of nodulation (AON) is a systemic, long-distance signaling pathway involving signals from developing nodules that are transported to the shoot, where they trigger retrograde signaling from the shoot to control (suppress) further nodulation (Caetano-Anollés and Gresshoff, 1991; Krusell et al., 2002). Nodule numbers are also controlled by local feedback confined to roots, as described below (Figure 7).

Shoot-root grafting of hypermodulating mutant and wild-type soybean showed that the root supernodulation of the *nts382* (nitrate tolerant symbiosis382) soybean mutant is determined by the shoot genotype (Delves et al., 1986), implicating a long-distance signaling pathway. The causative mutation in *nts382* is present in the *GmNARK* (NODULE AUTOREGULATION RECEPTOR KINASE) gene, which is orthologous to the *LjHAR1* (HYPERNODULATION AND ABERRANT ROOT) and *MtSUNN* genes (Krusell et al., 2002; Searle et al., 2003; Schnabel et al., 2005). *LjHAR1/GmNARK/MtSUNN*, which are homologous to Arabidopsis *CLAVATA1* (*CLV1*), a gene involved in meristem maintenance, encode class XI LRR-RLKs containing a short N-terminal plasma membrane-targeting signal, LRRs, a transmembrane domain, and a cytoplasmic Ser/Thr



**Figure 7.** Autoregulation of Nodulation.

Upon perception of rhizobia, the activation of cytokinin signaling results in increased expression of the genes encoding CLE12 and CLE13 peptides (1), which are then modified with arabinose residues by the RDN1 enzyme (2) and transported through the xylem to the shoot (3). Perception of the peptides in the shoot requires the receptor kinase SUNN, CLV2 receptor-like protein, and CRN pseudo-kinase (4). A second pathway involving the CRA2 receptor also controls nodule number by responding to a CEP peptide. Receptors in the shoot increase auxin and cytokinin transport to the root and the transport of *miR2111* (5), which affects *TML* expression (6).

kinase domain (Figure 7; Krusell et al., 2002; Nishimura et al., 2002; Schnabel et al., 2005).

Constitutive expression of the nodulation-induced peptide-encoding gene *MtCLE12/LjCLE-RS1* or *MtCLE13/LjCLE-RS2* eliminates nodulation (Okamoto et al., 2009; Mortier et al., 2010). However, overexpression of *MtCLE12* or *MtCLE13* in the *sun* mutant background reduces but does not eliminate nodulation, indicating that SUNN is required for responses to these two peptides (Mortier et al., 2012). *LjCLE-RS2* binds to *LjHAR1*, and the application of the peptide through vascular “feeding” showed that *LjCLE-RS2* is sufficient to halt nodulation in wild-type *L. japonicus* but not in a *har1* mutant, indicating that the *LjHAR1* receptor kinase is required for the control of the AON pathway by *LjCLE* (Okamoto et al., 2013). These data point to CLE peptides as

the root-to-shoot signal in the AON pathway. Counterparts of these CLEs include the rhizobia-induced *GmRIC1* and *GmRIC2* in *G. max* (Lim et al., 2011; Reid et al., 2011; Ferguson et al., 2014). Some CLE peptides in *Arabidopsis* must be arabinosylated at Hyp residues in order to function (Ogawa-Ohnishi et al., 2013; Imin et al., 2018). The *Mtrdn1/Ljplenty* mutants, carrying a disrupted Hyp arabinosyl transferase enzyme, also hypernodulate, but unlike *MtSUNN*, *MtRDN1* exerts its effect in roots (Yoshida et al., 2010; Schnabel et al., 2011; Yoro et al., 2019). Wild-type plants exhibit reduced nodule number in the presence of abundant available nitrogen (e.g., 10 mM  $\text{KNO}_3$ ), but most hypernodulation mutants have nitrate-resistant nodulation, implying that nitrogen status functions as an input to AON. While *LjCLE-RS1* and *LjCLE-RS2* genes were first identified as induced by rhizobia, *LjCLE-RS2* is also induced by nitrate (Okamoto et al., 2009), connecting nodulation, CLEs, and nitrate sensing. The nitrate-induced expression of *LjCLE-RS1* and *LjCLE-RS2* is dependent on NITRATE UNRESPONSIVE SYMBIOSIS1 (*LjNRSYM1*), a NIN-like protein (Nishida et al., 2018). IT formation and nodulation are insensitive to the application of high nitrate in *Ljnrsym1* mutants. However, analysis of the *Ljhar1-7 Ljnrsym1-1* double mutant indicated that while *LjHAR1* may be involved in regulating nodule number under N-deficient conditions, the control of IT number, nodule cell size, and N fixation is independent of *LjHAR1* (Nishida et al., 2018). Thus, the control of SNF by *LjNRSYM1* under high-N conditions likely occurs in parallel to AON.

Given the findings described above, legume *CLV2* homologs were also pursued for a role in AON. RNAi and EMS mutagenesis of *LjCLV2* produced a hypernodulation phenotype (Krusell et al., 2011). *MtCLV2* interacts with the *MtSUNN* receptor kinase and *MtCORYNE* (CRN), a pseudokinase that causes hypernodulation when mutated (Crook et al., 2016). The *Mtcm* mutant behaves like a *Mtsunn* mutant in grafting experiments: the effect is shoot derived (Crook et al., 2016), and *Mtcm* mutants do not respond to overexpression of *MtCLE12* or *MtCLE13* (Nowak et al., 2019). Together, mutational and physical evidence suggests crosstalk between AON receptors in the shoot (Figure 7, top).

Mutants in other receptors affecting nodule number yielded information about more potential interactions for AON receptors. The *Ljklavier* (*Ljklv*) mutant is a shoot-controlled hypernodulation mutant (Oka-Kira et al., 2005). *LjKLV* encodes a receptor-like kinase placed genetically in the AON pathway alongside *LjHAR1*, based upon mutant responses to *CLE-RS1* and *CLE-RS2* overexpression and physical interaction between the *KLV* and *HAR1* proteins (Miyazawa et al., 2010). Interestingly, *M. truncatula* has two *MtKLV* sequence homologs (Supplemental Data Set), which remain to be tested for roles in AON. Another receptor kinase identified by mutation, *MtCRA2*, regulates nodule number systemically from the shoot, independently of *MtSUNN*, to limit nodule development in the presence of nitrate (Huault et al., 2014; Mohd-Radzman et al., 2016). The C-terminally encoded peptide (CEP) family act as nitrogen “hunger” signals in plants and are transported systemically from root to shoot (Okamoto et al., 2016). Overexpression of *MtCEP1* enhances nodulation (Imin et al., 2013), and *CRA2* is required for CEP activity (Laffont et al., 2019), but analysis of a *cra2 sunn* double mutant suggested that these nodule regulatory pathways are independent (Laffont et al., 2019; Nowak et al., 2019).

Exciting progress has also been made on genes/proteins that act in roots in response to the shoot-derived inhibitor and on this mobile inhibitor itself. *TOO MUCH LOVE (TML)*, a gene identified via analysis of a hypernodulating mutant of *L. japonicus*, regulates nodule number from the root and is active after the signal is sent from the shoot (Magori and Kawaguchi, 2009). *LjTML* encodes a Kelch-repeat F-box protein possibly involved in proteasomal degradation of target proteins as a late step in AON (Takahara et al., 2013). Medicago has two *TML* genes, *MtTML1* and *MtTML2*. Knockdown of these genes by RNAi increased relative nodule number, implicating them in AON (Gautrat et al., 2019). Interestingly, homologous sequences in *Arabidopsis* and *G. max* are targeted for degradation by *miR2111*, a conserved microRNA that may be transported systemically (Ferguson et al., 2019). In fact, the expression of *miR2111* regulates *TML* in *L. japonicus* in a shoot-controlled manner dependent on *har1* (Tsikou et al., 2018), which would be expected of the AON shoot-derived inhibitor.

*NODULE NUMBER CONTROL1 (GmNNC1)* gene transcripts, encoding an AP2 TF, are cleaved by *miR172c*, which enables nodule development in soybean (Figure 7). Expression of *miR172c* is increased in *Gmnrk* nodule primordia, making *miR172c* another potential player in AON (Wang et al., 2014). *GmNNC1* physically interacts with the TF BES1/BZR1 HOMOLOG LIKE1 (*GmBEHL1*), a soybean homolog of *Arabidopsis* BES1/BZR1 HOMOLOG1 (*BEH1*) involved in brassinosteroid signaling (Yan et al., 2018). Knockdown of *GmBEHL1* expression doubled the number of nodules formed, concomitant with the induction of *GmmiR172c* expression.

### Cytokinin and AON-Independent Control of Nodulation

*Ljlhk1* cytokinin receptor mutants exhibit suppressed AON, leading to increased nodulation in *L. japonicus* (Murray et al., 2007). By contrast, mutations in *MtCRE1*, the Medicago ortholog of *LjLHK1*, reduce nodule number, which in an *Mtsunn* mutant background brings nodule number closer to wild-type levels, indicating that two separate pathways control nodule number (Gonzalez-Rizzo et al., 2006). *CRE1* is one of several genes that interact with AON signals to affect nodule number locally. In *M. truncatula*, the induction of *MtCLE12* and *MtCLE13* in the nodule meristem requires an active *MtCRE1* cytokinin receptor (Mortier et al., 2012). Similarly, *LjCLE-RS1* and *LjCLE-RS3* accumulation during infection is reduced in *lhk1-1* mutant roots (Tsikou et al., 2018), suggesting that early cytokinin signaling is required to send the AON signal. Evidence from tissue-specific expression of *LjLHK1* supports the notion that cortical cell cytokinin signaling functions as a local link to the systemic AON pathway (Miri et al., 2019). Other genes affecting nodule number are also tied to cytokinin signaling, but less directly to AON. Please see Figure 2 and the Supplemental Data Set for more information.

### BACTERIAL RELEASE AND SYMBIOSOME FORMATION

When the IT reaches the incipient nodule cortex, it branches and releases bacteria into plant cells via unwalling infection “droplets,” resulting in a unique “organelle” called the symbiosome (Figure 8). Bacteria within symbiosomes grow and divide, along with the plant-produced symbiosome membrane (SM), until the plant cell


is more or less packed with thousands of symbiosomes, each containing one or a few bacteria (Roth and Stacey, 1989).

### Role of the Exocytotic Pathway

Although early studies proposed that symbiosome formation occurs through an endocytic process (Mellor, 1989; Verma, 1992), recent genetic evidence implicates the secretory/exocytotic pathway, which delivers membrane vesicles to the plasma membrane, in this process (Limpens et al., 2009). Membrane fusion during exocytosis is mediated by SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) complexes, which are of two types: vesicle membrane-localized SNAREs (v-SNAREs) and target membrane-localized SNAREs (t-SNAREs). Downregulation of two v-SNARE genes in *M. truncatula*, *MtVAMP721d* and *MtVAMP721e* (*VESICLE-ASSOCIATED MEMBRANE PROTEIN*), blocked bacterial release from ITs and symbiosome formation without affecting nodule development (Ivanov et al., 2012). Similar results were found in *G. max* and *L. japonicus* (Gavrin et al., 2016; Sogawa et al., 2019). *MtVAMP721d* and *MtVAMP721e* are localized to the site of bacterial release and on the SM (Ivanov et al., 2012; Gavrin et al., 2016). Interestingly, reducing *MtVAMP721d* and *MtVAMP721e* expression also blocked arbuscule formation but not fungal colonization during AM symbiosis, indicating that these genes have conserved functions in intracellular symbioses in plants (Ivanov et al., 2012). As described above, *LjNPL* is required for the degradation of plant cell walls during rhizobium infection (Xie et al., 2012). The encapsulation of its ortholog, *GmNPL1*, in *GmVAMP721d*-positive vesicles indicates that exocytosis also plays a role in loosening cell walls prior to bacterial release into host cells (Gavrin et al., 2016). Another member of the *M. truncatula* VAMP72 family, *MtVAMP721a*, which may be involved in plant defense, is repressed by the nodule-enhanced C2H2-type TF, *MtRSD* (*REGULATOR OF SYMBIOSOME DEVELOPMENT*), and *MtRsd* mutants are defective in symbiosome development and nitrogen fixation (Sinharoy et al., 2013). Whether VAMP721a interferes or competes with VAMP721d/e-mediated vesicle trafficking and/or directs different types of vesicles (e.g., those containing defense molecules) to sites of bacterial entry in *MtRsd* mutants remains to be determined.

The exocytotic pathway involves the t-SNARE protein *MtSYP132* (*SYNTAXIN OF PLANTS132*), which is localized to the IT membrane, infection droplets, and SM in *M. truncatula* (Catalano et al., 2007; Limpens et al., 2009). RNAi of *MtSYP132A* impairs symbiosome differentiation and SNF (Pan et al., 2016). *MtSYP132A* is also required for arbuscule development during AM symbiosis, indicating that it plays conserved roles in plant-microbe interactions (Pan et al., 2016). *LjSYP71* has also been implicated in symbiosome differentiation, with *Ljsyp71* mutants exhibiting enlarged symbiosomes and a Fix<sup>-</sup> phenotype (Hakoyama et al., 2012b).

Synaptotagmins bind to t-SNAREs and v-SNAREs and facilitate the fusion of vesicles to target membranes. *MtSYT1*, *MtSYT2*, and *MtSYT3* are nodule synaptotagmins that are expressed from the meristematic to fixation zones. RNAi of the corresponding genes delayed rhizobium release, blocked symbiosome differentiation, and resulted in Fix<sup>-</sup> nodules (Gavrin et al., 2017).



**Figure 8.** Transport and Metabolism during Symbiotic Nitrogen Fixation.

Bacteria are released into the host cell in infection droplets (1) derived from Golgi deposits at the site of bacterial release (6), where they divide (2) and differentiate (3). Mature symbiosomes have multiple transporters on the symbiosome membrane and other membrane proteins that are essential for symbiosome membrane biogenesis (4). Other transporters on the cell membrane and from the vasculature (7) also import elements required for bacterial metabolism and optimal N fixation before undergoing senescence (8).

### Role of the Endocytic Pathway

Although exocytosis plays a pivotal role in rhizobium release and symbiosome formation, two endocytic pathway proteins, MtFLOT2/4 and MtSYMREM1, are found in the SM (Haney and Long, 2010; Lefebvre et al., 2010). RNAi of *MtFLOT2/4* caused the formation of small, white nodules with reduced SNF capacity, whereas an *Mtsymrem1* mutant was defective in bacterial release. Thus, MtFLOT2/4, MtSYMREM1, and endocytosis may participate in signal transduction and/or membrane trafficking associated with bacterial release and/or symbiosome development. Symbiosome maturation is also impaired upon knockdown of the late endosomal marker GTPase MtRab7 (Ras-related protein in brain), resulting in early nodule senescence (Limpens et al., 2009).

### Additional Components Involved in Symbiosome Differentiation

The cytoskeleton, including microtubules and actin filaments, plays a crucial role in intracellular transport (Petrásek and Schwarzerová, 2009). Actin undergoes dynamic rearrangement during bacterial release and symbiosome differentiation in nodules (Zhang et al., 2019). RNAi of *MtARP3* (*ACTIN RELATED PROTEIN*), encoding an actin nucleation complex component required for actin filament branching, impaired symbiosome development, resulting in small, nonfunctional symbiosomes (Gavrín

et al., 2015). TFs also affect symbiosome differentiation. Silencing a Kruppel-like zinc finger protein gene (*MtZPT2-1*) impaired early bacterial differentiation and SNF (Frugier et al., 2000). Additionally, the aquaporin MtTIP1g, which is required for water transport, is redirected from the vacuolar membrane to the SM in the nodule fixation zone. Repression of *MtTIP1g* arrested symbiosome elongation prior to maturation (Gavrín et al., 2014).

### HOST CONTROL OF BACTERIAL MATURATION

Rhizobia within symbiosomes differentiate into N-fixing bacteroids that release ammonia into the host cell in exchange for reduced carbon and other nutrients from the plant, with exchange mediated by transporters in the SM (Udvardi and Poole, 2013). In certain legumes, primarily galeoid (temperate) legumes including *M. truncatula*, bacterial differentiation involves changes in volume and shape (e.g., elongation), DNA content (endoreduplication), and loss of viability as a free-living organism (Mergaert et al., 2006). Bacteroid differentiation is accompanied (and in part controlled) by a decrease in free oxygen concentration in infected cells, which prevents the inactivation of oxygen-labile nitrogenase. Microaerobic conditions within nodules are achieved by restricted gaseous diffusion into nodules and high rates of respiration facilitated by oxygen-binding and oxygen-transporting leg(ume) hemoglobin in the cytoplasm of plant cells (Ott et al. 2005).

### Activation of Bacterial Nitrogenase

The induction of nitrogenase genes in rhizobia is controlled by oxygen. Free-oxygen concentrations in nodule cells are as low as  $10^{-8}$  M in the fixation zone, permitting nitrogenase to function (Tjepkema and Yocum, 1974). However, plant mitochondria and bacteroids require high oxygen flux to sustain respiration for ATP supply to nitrogenase. Leghemoglobins are essential for buffering oxygen in nodules at low levels while transporting it rapidly to rhizobia. RNAi and CRISPR/Cas9 of *LjLB1*, *LjLB2*, and *LjLB3* resulted in nodules with elevated oxygen concentrations but reduced ATP:ADP ratios and defective SNF (Ott et al., 2005; Wang et al., 2019). Importantly, *nif* genes (*nifH*) were not induced in these nodules (Ott et al., 2009).

Interestingly, bacterial nitrogenase activity is also dependent on plant supply of the cofactor homocitrate, at least in rhizobia that lack *nifE* and cannot make homocitrate. Mutations in *LjFEN1*, which encodes homocitrate synthase, impair SNF, but this can partially be overcome by adding homocitrate (Hakoyama et al., 2009).

### Terminal Differentiation

Indeterminate nodule-forming legumes belonging to the inverted repeat-lacking clade, including *M. truncatula*, require their symbionts to undergo terminal differentiation, including loss of motility, elongation, and endoreduplication, before they fix nitrogen. This process is controlled by the Cys-rich NCR peptides, which are cleaved into their biologically active forms from prepropeptides by peptidases (Figure 8). Mutations in two enzymes controlling NCR peptide maturation cause severe defects in nitrogen fixation. *DNF1* (*DEFECTIVE IN NITROGEN FIXATION1*) encodes a signal peptidase enzyme required for cleaving the N-terminal signal peptide directing NCR peptides to the symbiosome. Unprocessed prepropeptide NCRs in *dnf1* mutants do not induce terminal differentiation of bacteria, blocking SNF (Van de Velde et al., 2010; Wang et al., 2010). Nonetheless, infection and nodule organogenesis of *dnf1* are normal. A DNA methylase required for maturation of at least 82 NCR peptides, encoded by *DEMETER* (*MtDME*; Satgé et al., 2016), is also required for bacteroid endoreduplication, elongation, and SNF.

Mature 35- to 70-amino acid NCR peptides are targeted to the symbiosome, where they induce bacteroid differentiation. The NCR peptide family has at least 700 members in *M. truncatula* (de Bang et al., 2017), making it difficult to study them using conventional genetic tools. NCR coding sequences are short, making it hard to retrieve insertion mutants by reverse genetics. Furthermore, functional redundancy is expected to stymie genetic studies. Surprisingly, therefore, single mutations in *MtNCR211* (*dnf4*) or *MtNCR169* (*dnf7*) impair SNF (Horváth et al., 2015; Kim et al., 2015). Thus, their roles appear to be unique. Irregularly shaped undifferentiated bacteroids in *dnf7* mutant lines appear to be incompletely surrounded by the SM, while bacteroid-containing cells in *dnf4* mutants appear degenerated (Horváth et al., 2015; Kim et al., 2015). These observations demonstrate the importance of the NCR family in establishing effective symbiosis in *M. truncatula*.

Although determinate nodule-forming legumes do not possess NCR peptides, the Fix<sup>-</sup> phenotype of an *Ljapn1* (*aspartic peptidase nodule induced1*) mutant hints at the idea that other

peptides control bacterial persistence within nodules (Yamaya-Ito et al., 2018). Further studies of the roles of plant peptides during SNF are clearly warranted.

### SYMBIOTIC METABOLISM AND TRANSPORT

SNF is a metabolic symbiosis based upon the exchange of reduced C derived from plant photosynthesis for reduced N from rhizobial nitrogen fixation. Genetic studies have shown that effective SNF requires the coordination of multiple metabolic and transport pathways in both the plant and the microbe (Figure 8).

#### Carbon Metabolism and Transport

Sucrose derived from photosynthesis is transported through the vascular system to nodules, where it is hydrolyzed by Suc synthases or invertases that are typically nodule-enhanced (Morell and Copeland, 1984; Hohnjec et al., 2003; Fletmetakis et al., 2006). The reduced Suc synthase activity in *P. sativum* *Pssus1* and *L. japonicus* *Ljsus1* *Ljsus3* double mutants impairs SNF, as does RNAi silencing of *MtSucS* (Gordon et al., 1999; Baier et al., 2007; Horst et al., 2007). Built-in metabolic redundancy seems to be a theme in nodule carbon metabolism and transport. Both a *Tnt1*-insertion mutant of the nodule-enhanced Suc transporter gene *MtSWEET11* and a null mutation of *LjSUS3*, encoding the major nodule-induced isoform of Suc synthase, are uncompromised in SNF (Horst et al., 2007; Kryvoruchko et al., 2016).

#### Redox Metabolism

High rates of respiration result in the production of ROS, which are harmful to nodule metabolism (Chang et al., 2009). Antioxidants, including ascorbate, reduced GSH, and homoglutathione, can scavenge ROS and protect cells from oxidative damage. In *M. truncatula*,  $\gamma$ -glutamylcysteine synthetase (*Mt $\gamma$ ECS*) is required for GSH biosynthesis; silencing *Mt $\gamma$ ECS* impairs nodule growth and SNF (El Msehli et al., 2011).

#### Nitrogen Metabolism and Transport

Plant ammonia transport and assimilation and organic N export are central aspects of nodule function. The Gln synthetase-glutamate synthase (GOGAT) cycle catalyzes the assimilation of ammonium into amino acids. RNAi of *MtNADH-GOGAT* reduced nodule amino acid content (primarily Glu, Gln, and Ala) and impaired SNF in alfalfa (*Medicago sativa*; Cordoba et al. 2003). Amino acids, especially Gln and Asn, are exported from nodules of temperate legumes, including *M. truncatula* and *L. japonicus*, whereas ureides, especially allantoin and allantoic acid, are synthesized and transported from nodules to shoots in tropical legumes, including *G. max* and *P. vulgaris* (Smith and Atkins, 2002). In plants, ureides are synthesized by purine oxidation. Glutamine phosphoribosyl pyrophosphate amidotransferase (PRAT) catalyzes the first step of de novo purine synthesis. Silencing of nodule-enhanced *PvPRAT3* severely reduced ureide production and slightly reduced SNF (Coletto et al., 2016).

Several transporters of nitrogen-containing compounds are required for effective SNF. These include *LjAMT1;1*, *GmUPS1-1*

and GmUPS1-2, LjNPF8.6, and MtNPF1.7. RNAi of two nodule cortical and vascular plasma membrane-localized ureide permease genes, *GmUPS1-1* and *GmUPS1-2*, resulted in increased nodule ureide levels but reduced plant growth, reflecting decreased translocation of N from nodules to shoots (Collier and Tegeder, 2012). *LjAMT1;1* encodes a plasma membrane-localized high-affinity ammonium transporter. RNAi of *LjAMT1;1* resulted in less effective SNF, suggesting that this ammonium transporter might function in recovering ammonium that diffuses out of nodule cells before it is incorporated into amino acids for export to the shoot (Rogato et al., 2008). LjNPF8.6, a nodule-induced member of the NPF, showed dual-affinity nitrate transport activity in *X. laevis* oocytes, and SNF was reduced in an LjNPF8.6 mutant (Valkov et al., 2017).

### Phosphate and Sulfate Transport

The macronutrient phosphate is required by developing and functional nodules to sustain growth and metabolism. Low soil phosphate is an important limiting factor of nodule initiation and growth (Le Roux et al., 2009). PHOSPHATE STARVATION RESPONSE25 (GmPHR25)-regulated phosphate transporters GmPT5 (PHOSPHATE TRANSPORTER5) and GmPT7, which likely control phosphorus uptake into nodules, control nodule number and size (Qin et al., 2012; Xue et al., 2017; Chen et al., 2019). Sulfate is also important for SNF. Mutants of *L. japonicus* Symbiotic Sulfate Transporter1 (LjSST1) are defective in SNF (Krusell et al., 2005). A proteomics study placed LjSST1 on the SM (Wienkoop and Saalbach, 2003), indicating that it plays a role in sulfate transport from infected cells to rhizobia (Krusell et al., 2005).

### Metal Ion Transport

Metal ions play crucial roles in SNF as cofactors of metalloenzymes, including (Fe)-hemoglobins, (Fe-Mo)- and (Fe-S)-nitrogenases, and (Cu-Fe)-cytochromes, as well as TFs, including zinc finger TFs (Figure 8; Rubio et al., 2007; González-Guerrero et al., 2014).

Iron is delivered by the vascular system to nodules, where it is released to the apoplast before being taken up by plant cells and delivered to bacteroids (Rodríguez-Haas et al., 2013). The *M. truncatula* plasma membrane iron transporter, Natural Resistance-Associated Macrophage Protein1 (MtNRAMP1), transports iron from the apoplast into infected nodule cells where it is essential for SNF, as shown using *Mtnramp1 Tnt1*-insertion mutants (Tejada-Jiménez et al., 2015). Iron is often chelated to citrate to ensure its solubility and movement throughout the plant. The plasma membrane-localized, iron-activated citrate efflux transporter MtMATE67 is required for iron uptake from the apoplast of nodule cells and for effective SNF (Kryvoruchko et al., 2018). Similarly, RNAi of *Ljmate1* altered iron distribution in nodules and impaired SNF in *L. japonicus* (Takanashi et al., 2013).

LjSEN1 is required for rhizobial differentiation into N-fixing bacteroids in *L. japonicus* nodules (Suganuma et al., 2003). *LjSEN1* is homologous to Arabidopsis *VACUOLAR IRON TRANSPORTER1* and is expressed specifically in nodule-infected cells. LjSEN1 is thought to transport iron, but this remains to be demonstrated (Hakoyama et al., 2012a).

Molybdenum is crucial for iron-molybdenum nitrogenase activity. Two molybdenum transporters, MtMOT1.2 and MtMOT1.3, are essential for SNF in *M. truncatula*. MtMOT1.2 is located on the plasma membrane of endodermal cells surrounding nodule vascular vessels and mediates molybdate uptake and distribution into nodule cells (Gil-Díez et al., 2019). MtMOT1.3, which is located on the plasma membrane of infected cells, mediates molybdate uptake into these cells (Tejada-Jiménez et al., 2017). The mutation of either *MtMOT1.2* or *MtMOT1.3* impairs SNF (Tejada-Jiménez et al., 2017; Gil-Díez et al., 2019). How molybdate is transported across the SM remains to be determined.

Zinc is a cofactor of superoxide dismutases that detoxify ROS and is present in zinc-finger motif TFs (Rubio et al., 2007; González-Guerrero et al., 2014). Two zinc transporters, MtZIP6 (Zinc-Iron Permease6) and MtMTP2, are essential for SNF. RNAi of *MtZIP6*, encoding an enzyme in the plasma membrane of infected cells responsible for zinc uptake from the apoplast, impaired SNF (Abreu et al., 2017). *M. truncatula* Metal Tolerance Protein2 (MtMTP2), which is localized to the endomembrane system, is a zinc efflux protein (León-Mediavilla et al., 2018). Mutation of *Mtmtp2* resulted in aberrant bacteroid differentiation and early senescence, suggesting a defect in intracellular zinc allocation (León-Mediavilla et al., 2018).

Copper is a cofactor of several enzymes, including cytochrome oxidases. *M. truncatula* Copper Transporter1 (MtCOPT1), which is localized to the plasma membrane, imports copper from the apoplast into nodule cells. *Mtcopt1* mutants displayed lower SNF and lower copper-dependent cytochrome oxidase activity than wild-type plants (Senovilla et al., 2018).

### SENESCENCE AND DEFENSE

Nodule senescence is a controlled process that enables nutrient recycling and can be triggered by developmental and environmental cues such as drought and high nitrate levels. During age-related senescence, bacteroids lyse, forming numerous vesicles within the cytosol of infected cells, followed by senescence of infected cells (Van de Velde et al., 2006). Typically, senescence is mediated by TFs and involves proteases and lipases that degrade proteins and lipids, respectively, as well as transporters that export breakdown products (Schippers et al., 2015). The TF gene *MtNAC969* (*NAM/ATAF/CUC-encoding969*) is induced in the nitrogen fixation zone upon nitrate treatment (de Zélicourt et al., 2012). RNAi of *MtNAC969* did not alter nodule formation but it activated senescence genes, including two senescence-induced cysteine proteinase (CP) genes, and caused premature nodule aging (de Zélicourt et al., 2012). Therefore, MtNAC969 appears to act as a suppressor of nodule senescence that is executed, at least in part, by CPs. RNAi of two CP genes expressed in the nodule senescence zone, *MtCP6*, encoding a papain-type proteinase, and *MtVPE* (*VACUOLAR PROCESSING ENZYME*), encoding a protein of the legumain proteinase subclass, delayed nodule senescence and prolonged nodule growth and nitrogen fixation (Pérez Guerra et al., 2010; Pierre et al., 2014). MtCP6 and MtVPE are localized to the SM, suggesting that they are involved in symbiosome degradation. Conversely, knockdown of the CP inhibitor gene *MtSERPIN6* accelerated nodule aging (Dhanushkodi et al., 2018). Interestingly, mutations in the *STAY-GREEN* gene *MtSGR*



led to aberrant shoot senescence and delayed nodule senescence (Zhou et al., 2011).

Iron liberated from leghemoglobin, especially during senescence, is a potential source of ROS, which can interfere with nodule metabolism. Iron storage proteins that scavenge free iron include ferritin multimeric complexes (Theil, 2003). Silencing of two ferritin genes, *MtFer2* and *MtFer3*, increased free iron levels in the nodule and induced early nodule senescence (Dhanushkodi et al., 2018). Therefore, preventing leghemoglobin degradation might delay nodule senescence. Accordingly, nodules of the *Mtcas31* (*cold acclimation specific31*) mutant, which is defective in a dehydrin protein that interacts with the leghemoglobin MtLB120-1, had elevated transcript levels of senescence genes and reduced nitrogenase activity under drought stress (Li et al., 2018). Although nodule senescence enables plants to recover part of their investment in nodules via the remobilization of nutrients, premature nodule senescence compromises SNF and N supply to the plant. Therefore, it would be interesting to identify key signaling components and regulators of senescence as a step toward improving SNF by delaying nodule senescence.

Defense phenotypes often overlap with those exhibited by senescent nodules. For example, knockdown of *MtSNARP2* (*SMALL NODULIN ACIDIC RNA BINDING PROTEIN2*) resulted in defense responses such as the development of a thick nodule outer cortex and a closed endodermal layer (Laporte et al., 2010). On the other hand, *Mtsnarp2* mutants senesce prematurely without accumulating brown pigments often associated with defense. In *nodule with activated defenses1* (*Mtnad1*) mutant nodules, cells of both symbiotic partners are disrupted and the defense marker gene *PR10* is upregulated (Wang et al., 2016; Domonkos et al., 2017). Bacteria in *Mtnad1* mutants have an unorganized bacteroid cell wall showing degradation. Consequently, SNF is arrested and does not progress to the N-fixation stage.

Interfering with SNF progression at any stage also elicits defense-like responses. *Fix-* mutants of the *DOES NOT FIX2* (*MtDNF2*) gene, encoding a phosphatidylinositol phospholipase C-like protein, and the coregulated *SYMBIOTIC CYSTEINE RICH RECEPTOR-LIKE KINASE* (*MtSymCRK*), which do not belong to known defense pathways, exhibit defense-like phenotypes such as brown-pigmented, necrotic nodules, defense marker gene induction, a block in bacterial terminal differentiation, and no N-fixing activity (Bourcy et al., 2013). Interestingly, these defense-like reactions in the *dnf2* mutant manifest only when grown on agar containing impurities, further supporting the idea that defense responses triggered by an external elicitor can impair SNF progression (Berrabah et al., 2014). Additional research is warranted to understand whether such molecular players are direct repressors of defense pathways or if the associated mutant phenotypes are secondary responses to interrupted SNF.

## CONCLUSIONS AND FUTURE DIRECTIONS

The last 20 years have seen amazing progress in the discovery of genes and mechanisms underlying SNF. While most SNF genes have homologs in nonlegumes, many of these genes have evolved roles in SNF via gene family expansion and/or neofunctionalization, often resulting in nodule-specific expression. Mycorrhizal root symbiosis evolved in land plants hundreds

of millions of years before legume-rhizobia SNF, and work over the last 20 years has uncovered several common symbiotic genes that are required for SNF that first evolved for mycorrhizal symbiosis. Genetic studies in legumes have focused on one or two genotypes per species, but genome sequencing of multiple genotypes per species has revealed that the pan-genome, which encompasses sequences of all individuals of a species, is much larger than any single genome. This raises questions about the functions of additional (or missing) sequences in specific genotypes and the extent to which they might contribute to SNF. New genetic tools, especially genome editing (Curtin et al., 2017) that enables targeted genetic changes in novel genotypes, should help to answer such questions. Genome editing will also be useful in addressing the functions of multigene families, whose roles in SNF have thus far been shrouded by genetic redundancy, as well as the roles of small genes that are less frequently mutated by conventional approaches. While genomics is beginning to illuminate the sequence diversity present within individual plant species, plant breeders have been using such diversity through phenotypic selection for improved plant traits for many years. However, breeding programs have rarely if ever selected directly for SNF effectiveness in crop or forage legumes. Doing so now with the benefit of the genome sequences of hundreds or thousands of diverse genotypes for a given species opens up the possibility of identifying genes and alleles of those genes that contribute to SNF effectiveness. This, in turn, may facilitate breeding to enhance SNF in legumes via marker-assisted or genomic-selection approaches, for instance.

Despite the remarkable progress of the last 20 years, much remains to be learned about the genetics and molecular biology of SNF in legumes and how the various pieces fit together. The next 20 years will likely to be as exciting as the past, especially as we endeavor to translate genetic knowledge into plant breeding programs that optimize SNF in legumes and possibly extend it into nonlegumes of agricultural importance.

## Supplemental Data

**Supplemental Data Set.** List of functionally validated SNF genes in legumes.

## ACKNOWLEDGMENTS

We thank Marcus Griffiths and Josh Meo for help with figure preparation and Yinbing Ge and Jody Beard for developing the accompanying webpage (<https://nobleapps.noble.org/legumegenetics>), Nick Krom for sequence information, and the three anonymous reviewers for thorough and constructive feedback. We thank the creators of the Reactome icon library, which we used extensively (Sidiropoulos et al., 2017). This work was funded by the National Science Foundation Plant Genome Research Program (grants DBI-1127155 and DBI-1733470) and by the Noble Research Institute.

## AUTHOR CONTRIBUTIONS

S.R. and W.L. curated gene information; R.S.N. performed database searches; S.R. prepared all figures with input from W.L.; S.R., W.L.,

R.S.N., A.C., C.I.P., K.S.M., J.F., and R.D. contributed to the writing of the article; M.K.U wrote, edited and supervised all aspects of the article.

Received April 19, 2019; revised September 17, 2019; accepted October 24, 2019; published October 24, 2019.

## REFERENCES

- Abreu, I., Saéz, Á., Castro-Rodríguez, R., Escudero, V., Rodríguez-Haas, B., Senovilla, M., Larue, C., Grolimund, D., Tejada-Jiménez, M., Imperial, J., and González-Guerrero, M. (2017). *Medicago truncatula* Zinc-Iron Permease6 provides zinc to rhizobia-infected nodule cells. *Plant Cell Environ.* **40**: 2706–2719.
- Andriankaja, A., Boisson-Dernier, A., Frances, L., Sauviac, L., Jauneau, A., Barker, D.G., and de Carvalho-Niebel, F. (2007). AP2-ERF transcription factors mediate Nod factor dependent *Mt ENOD11* activation in root hairs via a novel *cis*-regulatory motif. *Plant Cell* **19**: 2866–2885.
- Andrio, E., Marino, D., Marmeys, A., de Segonzac, M.D., Damiani, I., Genre, A., Huguet, S., Frendo, P., Puppo, A., and Pauly, N. (2013). Hydrogen peroxide-regulated genes in the *Medicago truncatula*-*Sinorhizobium meliloti* symbiosis. *New Phytol.* **198**: 179–189.
- Ané, J.-M., et al. (2004). *Medicago truncatula* DMI1 required for bacterial and fungal symbioses in legumes. *Science* **303**: 1364–1367.
- Antolín-Llovera, M., Ried, M.K., and Parniske, M. (2014). Cleavage of the SYMBIOSIS RECEPTOR-LIKE KINASE ectodomain promotes complex formation with Nod factor receptor 5. *Curr. Biol.* **24**: 422–427.
- Ariel, F., et al. (2012). Two direct targets of cytokinin signaling regulate symbiotic nodulation in *Medicago truncatula*. *Plant Cell* **24**: 3838–3852.
- Arrighi, J.-F., et al. (2006). The *Medicago truncatula* lysin [corrected] motif-receptor-like kinase gene family includes NFP and new nodule-expressed genes. *Plant Physiol.* **142**: 265–279.
- Arrighi, J.-F., Godfroy, O., de Billy, F., Saurat, O., Jauneau, A., and Gough, C. (2008). The *RPG* gene of *Medicago truncatula* controls Rhizobium-directed polar growth during infection. *Proc. Natl. Acad. Sci. USA* **105**: 9817–9822.
- Arthikala, M.-K., Montiel, J., Sánchez-López, R., Nava, N., Cárdenas, L., and Quinto, C. (2017). *Respiratory Burst Oxidase Homolog Gene A* is crucial for rhizobium infection and nodule maturation and function in common bean. *Front. Plant Sci.* **8**: 2003.
- Arthikala, M.-K., Nanjareddy, K., and Lara, M. (2018). In *BPS1* downregulated roots, the *BYPASS1* signal disrupts the induction of cortical cell divisions in bean-Rhizobium symbiosis. *Genes (Basel)* **9**: 11.
- Bagchi, R., Salehin, M., Adeyemo, O.S., Salazar, C., Shulaev, V., Sherrier, D.J., and Dickstein, R. (2012). Functional assessment of the *Medicago truncatula* NIP/LATD protein demonstrates that it is a high-affinity nitrate transporter. *Plant Physiol.* **160**: 906–916.
- Baier, M.C., Barsch, A., Küster, H., and Hohnjec, N. (2007). Antisense repression of the *Medicago truncatula* nodule-enhanced sucrose synthase leads to a handicapped nitrogen fixation mirrored by specific alterations in the symbiotic transcriptome and metabolome. *Plant Physiol.* **145**: 1600–1618.
- Battaglia, M., Ripodas, C., Clúa, J., Baudin, M., Aguilar, O.M., Niebel, A., Zanetti, M.E., and Blanco, F.A. (2014). A nuclear factor Y interacting protein of the GRAS family is required for nodule organogenesis, infection thread progression, and lateral root growth. *Plant Physiol.* **164**: 1430–1442.
- Benedito, V.A., et al. (2008). A gene expression atlas of the model legume *Medicago truncatula*. *Plant J.* **55**: 504–513.
- Berrabah, F., Bourcy, M., Eschstruth, A., Cayrel, A., Guefrachi, I., Mergaert, P., Wen, J., Jean, V., Mysore, K.S., Gourion, B., and Ratet, P. (2014). A nonRD receptor-like kinase prevents nodule early senescence and defense-like reactions during symbiosis. *New Phytol.* **203**: 1305–1314.
- Bolaños-Vásquez, M.C., and Werner, D. (1997). Effects of *Rhizobium tropici*, *R. etli*, and *R. leguminosarum* bv. *phaseoli* on nod gene-inducing flavonoids in root exudates of *Phaseolus vulgaris*. *Mol. Plant Microbe Interact.* **10**: 339–346.
- Bolon, Y.-T., et al. (2011). Phenotypic and genomic analyses of a fast neutron mutant population resource in soybean. *Plant Physiol.* **156**: 240–253.
- Bourcy, M., Brocard, L., Pislariu, C.I., Cosson, V., Mergaert, P., Tadege, M., Mysore, K.S., Udvardi, M.K., Gourion, B., and Ratet, P. (2013). *Medicago truncatula* DNF2 is a PI-PLC-XD-containing protein required for bacteroid persistence and prevention of nodule early senescence and defense-like reactions. *New Phytol.* **197**: 1250–1261.
- Breakspear, A., Liu, C., Roy, S., Stacey, N., Rogers, C., Trick, M., Morieri, G., Mysore, K.S., Wen, J., Oldroyd, G.E., Downie, J.A., and Murray, J.D. (2014). The root hair “infectome” of *Medicago truncatula* uncovers changes in cell cycle genes and reveals a requirement for auxin signaling in rhizobial infection. *Plant Cell* **26**: 4680–4701.
- Brogghammer, A., et al. (2012). Legume receptors perceive the rhizobial lipochitin oligosaccharide signal molecules by direct binding. *Proc. Natl. Acad. Sci. USA* **109**: 13859–13864.
- Caetano-Anollés, G., and Gresshoff, P.M. (1991). Plant genetic control of nodulation. *Annu. Rev. Microbiol.* **45**: 345–382.
- Cai, J., et al. (2018). Role of the Nod factor hydrolase MtNHF1 in regulating Nod factor levels during rhizobial infection and in mature nodules of *Medicago truncatula*. *Plant Cell* **30**: 397–414.
- Canfield, D.E., Glazer, A.N., and Falkowski, P.G. (2010). The evolution and future of Earth’s nitrogen cycle. *Science* **330**: 192–196.
- Cao, Y., Halane, M.K., Gassmann, W., and Stacey, G. (2017). The role of plant innate immunity in the legume-rhizobium symbiosis. *Annu. Rev. Plant Biol.* **68**: 535–561.
- Catalano, C.M., Czymmek, K.J., Gann, J.G., and Sherrier, D.J. (2007). *Medicago truncatula* syntaxin SYP132 defines the symbiosome membrane and infection droplet membrane in root nodules. *Planta* **225**: 541–550.
- Cebolla, A., Vinardell, J.M., Kiss, E., Oláh, B., Roudier, F., Kondorosi, A., and Kondorosi, E. (1999). The mitotic inhibitor *ccs52* is required for endoreduplication and ploidy-dependent cell enlargement in plants. *EMBO J.* **18**: 4476–4484.
- Cerri, M.R., Frances, L., Kelner, A., Fournier, J., Middleton, P.H., Auriac, M.-C., Mysore, K.S., Wen, J., Erard, M., Barker, D.G., Oldroyd, G.E., and de Carvalho-Niebel, F. (2016). The symbiosis-related ERN transcription factors act in concert to coordinate rhizobial host root infection. *Plant Physiol.* **171**: 1037–1054.
- Cerri, M.R., et al. (2017). The *ERN1* transcription factor gene is a target of the CCaMK/CYCLOPS complex and controls rhizobial infection in *Lotus japonicus*. *New Phytol.* **215**: 323–337.
- Chang, C., Damiani, I., Puppo, A., and Frendo, P. (2009). Redox changes during the legume-rhizobium symbiosis. *Mol. Plant* **2**: 370–377.
- Charpentier, M., Bredemeier, R., Wanner, G., Takeda, N., Schleiff, E., and Parniske, M. (2008). *Lotus japonicus* CASTOR and POLLUX are ion channels essential for perinuclear calcium spiking in legume root endosymbiosis. *Plant Cell* **20**: 3467–3479.
- Charpentier, M., and Oldroyd, G. (2010). How close are we to nitrogen-fixing cereals? *Curr. Opin. Plant Biol.* **13**: 556–564.

- Charpentier, M., and Oldroyd, G.E. (2013). Nuclear calcium signaling in plants. *Plant Physiol.* **163**: 496–503.
- Charpentier, M., Sun, J., Vaz Martins, T., Radhakrishnan, G.V., Findlay, K., Soumpourou, E., Thouin, J., Véry, A.-A., Sanders, D., Morris, R.J., and Oldroyd, G.E. (2016). Nuclear-localized cyclic nucleotide-gated channels mediate symbiotic calcium oscillations. *Science* **352**: 1102–1105.
- Chen, L., et al. (2019). A nodule-localized phosphate transporter GmPT7 plays an important role in enhancing symbiotic N<sub>2</sub> fixation and yield in soybean. *New Phytol.* **221**: 2013–2025.
- Chen, T., Zhu, H., Ke, D., Cai, K., Wang, C., Gou, H., Hong, Z., and Zhang, Z. (2012). A MAP kinase kinase interacts with SymRK and regulates nodule organogenesis in *Lotus japonicus*. *Plant Cell* **24**: 823–838.
- Chen, Y., Chen, W., Li, X., Jiang, H., Wu, P., Xia, K., Yang, Y., and Wu, G. (2014). Knockdown of *LjIPT3* influences nodule development in *Lotus japonicus*. *Plant Cell Physiol.* **55**: 183–193.
- Cheng, X., Gou, X., Yin, H., Mysore, K.S., Li, J., and Wen, J. (2017). Functional characterisation of brassinosteroid receptor *MtBRI1* in *Medicago truncatula*. *Sci. Rep.* **7**: 9327.
- Chiasson, D.M., Haage, K., Sollweck, K., Brachmann, A., Dietrich, P., and Parniske, M. (2017). A quantitative hypermorphic *CNGC* allele confers ectopic calcium flux and impairs cellular development. *eLife* **6**: e25012.
- Coletto, I., Trenas, A.T., Erban, A., Kopka, J., Pineda, M., and Alamillo, J.M. (2016). Functional specialization of one copy of *glutamine phosphoribosyl pyrophosphate amidotransferase* in ureide production from symbiotically fixed nitrogen in *Phaseolus vulgaris*. *Plant Cell Environ.* **39**: 1767–1779.
- Collier, R., and Tegeder, M. (2012). Soybean ureide transporters play a critical role in nodule development, function and nitrogen export. *Plant J.* **72**: 355–367.
- Combiér, J.P., de Billy, F., Gamas, P., Niebel, A., and Rivas, S. (2008). Trans-regulation of the expression of the transcription factor *MtHAP2-1* by a uORF controls root nodule development. *Genes Dev.* **22**: 1549–1559.
- Combiér, J.-P., Frugier, F., de Billy, F., Boualem, A., El-Yahyaoui, F., Moreau, S., Vernié, T., Ott, T., Gamas, P., Crespi, M., and Niebel, A. (2006). *MtHAP2-1* is a key transcriptional regulator of symbiotic nodule development regulated by microRNA169 in *Medicago truncatula*. *Genes Dev.* **20**: 3084–3088.
- Cordoba, E., Shishkova, S., Vance, C.P., and Hernández, G. (2003). Antisense inhibition of NADH glutamate synthase impairs carbon/nitrogen assimilation in nodules of alfalfa (*Medicago sativa* L.). *Plant J.* **33**: 1037–1049.
- Couzigou, J.-M., et al. (2012). *NODULE ROOT* and *COCHLEATA* maintain nodule development and are legume orthologs of Arabidopsis *BLADE-ON-PETIOLE* genes. *Plant Cell* **24**: 4498–4510.
- Crook, A.D., Schnabel, E.L., and Frugoli, J.A. (2016). The systemic nodule number regulation kinase SUNN in *Medicago truncatula* interacts with MtCLV2 and MtCRN. *Plant J.* **88**: 108–119.
- Curtin, S.J., Tiffin, P., Guhlin, J., Trujillo, D.I., Burghart, L.T., Atkins, P., Baltes, N.J., Denny, R., Voytas, D.F., Stupar, R.M., and Young, N.D. (2017). Validating genome-wide association candidates controlling quantitative variation in nodulation. *Plant Physiol.* **173**: 921–931.
- de Bang, T.C., et al. (2017). Genome-wide identification of *Medicago* peptides involved in macronutrient responses and nodulation. *Plant Physiol.* **175**: 1669–1689.
- Delves, A.C., Mathews, A., Day, D.A., Carter, A.S., Carroll, B.J., and Gresshoff, P.M. (1986). Regulation of the soybean-Rhizobium nodule symbiosis by shoot and root factors. *Plant Physiol.* **82**: 588–590.
- Dénarié, J., and Cullimore, J. (1993). Lipo-oligosaccharide nodulation factors: A new class of signaling molecules mediating recognition and morphogenesis. *Cell* **74**: 951–954.
- de Zélicourt, A., Diet, A., Marion, J., Laffont, C., Ariel, F., Moison, M., Zahaf, O., Crespi, M., Gruber, V., and Frugier, F. (2012). Dual involvement of a *Medicago truncatula* NAC transcription factor in root abiotic stress response and symbiotic nodule senescence. *Plant J.* **70**: 220–230.
- D’Haeze, W., and Holsters, M. (2002). Nod factor structures, responses, and perception during initiation of nodule development. *Glycobiology* **12**: 79R–105R.
- Dhanushkodi, R., Matthew, C., McManus, M.T., and Dijkwel, P.P. (2018). Drought-induced senescence of *Medicago truncatula* nodules involves serpin and ferritin to control proteolytic activity and iron levels. *New Phytol.* **220**: 196–208.
- Di Giacomo, E., Laffont, C., Sciarra, F., Iannelli, M.A., Frugier, F., and Frugis, G. (2017). KNAT3/4/5-like class 2 KNOX transcription factors are involved in *Medicago truncatula* symbiotic nodule organ development. *New Phytol.* **213**: 822–837.
- Domonkos, Á., et al. (2017). NAD1 controls defense-like responses in *Medicago truncatula* symbiotic nitrogen fixing nodules following rhizobial colonization in a BacA-independent manner. *Genes (Basel)* **8**: 387.
- Doyle, J.J., and Luckow, M.A. (2003). The rest of the iceberg: Legume diversity and evolution in a phylogenetic context. *Plant Physiol.* **131**: 900–910.
- El Msehli, S., Lambert, A., Baldacci-Cresp, F., Hopkins, J., Boncompagni, E., Smiti, S.A., Hérouart, D., and Frendo, P. (2011). Crucial role of (homo)glutathione in nitrogen fixation in *Medicago truncatula* nodules. *New Phytol.* **192**: 496–506.
- Endre, G., Kereszt, A., Kevei, Z., Mihacea, S., Kaló, P., and Kiss, G.B. (2002). A receptor kinase gene regulating symbiotic nodule development. *Nature* **417**: 962–966.
- Espina, M.J., Ahmed, C.M.S., Bernardini, A., Adeleke, E., Yadegari, Z., Arelli, P., Pantalone, V., and Taheri, A. (2018). Development and phenotypic screening of an ethyl methane sulfonate mutant population in soybean. *Front. Plant Sci.* **9**: 394.
- Estrada-Navarrete, G., Cruz-Mireles, N., Lascano, R., Alvarado-Affantranger, X., Hernández-Barrera, A., Barraza, A., Olivares, J.E., Arthikala, M.-K., Cárdenas, L., Quinto, C., and Sanchez, F. (2016). An autophagy-related kinase is essential for the symbiotic relationship between *Phaseolus vulgaris* and both rhizobia and arbuscular mycorrhizal fungi. *Plant Cell* **28**: 2326–2341.
- Fan, Y., Liu, J., Lyu, S., Wang, Q., Yang, S., and Zhu, H. (2017). The soybean *Rfg1* gene restricts nodulation by *Sinorhizobium fredii* USDA193. *Front. Plant Sci.* **8**: 1548.
- Ferguson, B.J., Li, D., Hastwell, A.H., Reid, D.E., Li, Y., Jackson, S.A., and Gresshoff, P.M. (2014). The soybean (*Glycine max*) nodulation-suppressive CLE peptide, GmRIC1, functions interspecifically in common white bean (*Phaseolus vulgaris*), but not in a supernodulating line mutated in the receptor PvNARK. *Plant Biotechnol. J.* **12**: 1085–1097.
- Ferguson, B.J., Mens, C., Hastwell, A.H., Zhang, M., Su, H., Jones, C.H., Chu, X., and Gresshoff, P.M. (2019). Legume nodulation: The host controls the party. *Plant Cell Environ.* **42**: 41–51.
- Flemetakis, E., Efroze, R.C., Ott, T., Stedel, C., Aivalakis, G., Udvardi, M.K., and Katinakis, P. (2006). Spatial and temporal organization of sucrose metabolism in *Lotus japonicus* nitrogen-fixing nodules suggests a role for the elusive alkaline/neutral invertase. *Plant Mol. Biol.* **62**: 53–69.
- Fonouni-Farde, C., Tan, S., Baudin, M., Brault, M., Wen, J., Mysore, K.S., Niebel, A., Frugier, F., and Diet, A. (2016). DELLA-mediated gibberellin signalling regulates Nod factor signalling and rhizobial infection. *Nat. Commun.* **7**: 12636.

- Franssen, H.J., Xiao, T.T., Kulikova, O., Wan, X., Bisseling, T., Scheres, B., and Heidstra, R. (2015). Root developmental programs shape the *Medicago truncatula* nodule meristem. *Development* **142**: 2941–2950.
- Frugier, F., Poirier, S., Satiat-Jeunemaitre, B., Kondorosi, A., and Crespi, M. (2000). A Krüppel-like zinc finger protein is involved in nitrogen-fixing root nodule organogenesis. *Genes Dev.* **14**: 475–482.
- Fukai, E., Stougaard, J., and Hayashi, M. (2013). Activation of an endogenous retrotransposon associated with epigenetic changes in *Lotus japonicus*: A tool for functional genomics in legumes. *Plant Genome* **6**.
- Gautrat, P., Mortier, V., Laffont, C., De Keyser, A., Fromentin, J., Frugier, F., and Goormachtig, S. (2019). Unraveling new molecular players involved in the autoregulation of nodulation in *Medicago truncatula*. *J. Exp. Bot.* **70**: 1407–1417.
- Gavrin, A., Chiasson, D., Ovchinnikova, E., Kaiser, B.N., Bisseling, T., and Fedorova, E.E. (2016). VAMP721a and VAMP721d are important for pectin dynamics and release of bacteria in soybean nodules. *New Phytol.* **210**: 1011–1021.
- Gavrin, A., Jansen, V., Ivanov, S., Bisseling, T., and Fedorova, E. (2015). ARP2/3-mediated actin nucleation associated with symbiosome membrane is essential for the development of symbiosomes in infected cells of *Medicago truncatula* root nodules. *Mol. Plant Microbe Interact.* **28**: 605–614.
- Gavrin, A., Kaiser, B.N., Geiger, D., Tyerman, S.D., Wen, Z., Bisseling, T., and Fedorova, E.E. (2014). Adjustment of host cells for accommodation of symbiotic bacteria: Vacuole de-functionalization, HOPS suppression, and TIP1g re-targeting in *Medicago*. *Plant Cell* **26**: 3809–3822.
- Gavrin, A., Kulikova, O., Bisseling, T., and Fedorova, E.E. (2017). Interface symbiotic membrane formation in root nodules of *Medicago truncatula*: The role of synaptotagmins *MtSyt1*, *MtSyt2* and *MtSyt3*. *Front. Plant Sci.* **8**: 201.
- Geurts, R., Xiao, T.T., and Reinhold-Hurek, B. (2016). What does it take to evolve a nitrogen-fixing endosymbiosis? *Trends Plant Sci.* **21**: 199–208.
- Gil-Diez, P., Tejada-Jiménez, M., León-Mediavilla, J., Wen, J., Mysore, K.S., Imperial, J., and González-Guerrero, M. (2019). MtMOT1.2 is responsible for molybdate supply to *Medicago truncatula* nodules. *Plant Cell Environ.* **42**: 310–320.
- Gleason, C., Chaudhuri, S., Yang, T., Muñoz, A., Poovaiah, B.W., and Oldroyd, G.E. (2006). Nodulation independent of rhizobia induced by a calcium-activated kinase lacking autoinhibition. *Nature* **441**: 1149–1152.
- Glebov, O.O., Bright, N.A., and Nichols, B.J. (2006). Flotillin-1 defines a clathrin-independent endocytic pathway in mammalian cells. *Nat. Cell Biol.* **8**: 46–54.
- Godiard, L., Lepage, A., Moreau, S., Laporte, D., Verdenaud, M., Timmers, T., and Gamas, P. (2011). MtbHLH1, a bHLH transcription factor involved in *Medicago truncatula* nodule vascular patterning and nodule to plant metabolic exchanges. *New Phytol.* **191**: 391–404.
- González-Guerrero, M., Matthiadis, A., Sáez, Á., and Long, T.A. (2014). Fixating on metals: New insights into the role of metals in nodulation and symbiotic nitrogen fixation. *Front. Plant Sci.* **5**: 45.
- Gonzalez-Rizzo, S., Crespi, M., and Frugier, F. (2006). The *Medicago truncatula* CRE1 cytokinin receptor regulates lateral root development and early symbiotic interaction with *Sinorhizobium meliloti*. *Plant Cell* **18**: 2680–2693.
- Gordon, A.J., Minchin, F.R., James, C.L., and Komina, O. (1999). Sucrose synthase in legume nodules is essential for nitrogen fixation. *Plant Physiol.* **120**: 867–878.
- Groth, M., Takeda, N., Perry, J., Uchida, H., Dräxl, S., Brachmann, A., Sato, S., Tabata, S., Kawaguchi, M., Wang, T.L., and Parniske, M. (2010). *NENA*, a *Lotus japonicus* homolog of *Sec13*, is required for rhizodermal infection by arbuscular mycorrhiza fungi and rhizobia but dispensable for cortical endosymbiotic development. *Plant Cell* **22**: 2509–2526.
- Grunewald, W., van Noorden, G., Van Isterdael, G., Beeckman, T., Gheysen, G., and Mathesius, U. (2009). Manipulation of auxin transport in plant roots during *Rhizobium* symbiosis and nematode parasitism. *Plant Cell* **21**: 2553–2562.
- Guan, D., et al. (2013). Rhizobial infection is associated with the development of peripheral vasculature in nodules of *Medicago truncatula*. *Plant Physiol.* **162**: 107–115.
- Hadri, A.-E., Spaik, H.P., Bisseling, T., and Brewin, N.J. (1998). Diversity of root nodulation and rhizobial infection processes. In *The Rhizobiaceae*, H.P. Spaik, A. Kondorosi, and P.J.J. Hooykaas, eds (Dordrecht, The Netherlands: Springer), pp. 347–360.
- Hakoyama, T., et al. (2009). Host plant genome overcomes the lack of a bacterial gene for symbiotic nitrogen fixation. *Nature* **462**: 514–517.
- Hakoyama, T., et al. (2012a). The integral membrane protein SEN1 is required for symbiotic nitrogen fixation in *Lotus japonicus* nodules. *Plant Cell Physiol.* **53**: 225–236.
- Hakoyama, T., et al. (2012b). The SNARE protein SYP71 expressed in vascular tissues is involved in symbiotic nitrogen fixation in *Lotus japonicus* nodules. *Plant Physiol.* **160**: 897–905.
- Haney, C.H., and Long, S.R. (2010). Plant flotillins are required for infection by nitrogen-fixing bacteria. *Proc. Natl. Acad. Sci. USA* **107**: 478–483.
- Haney, C.H., Riely, B.K., Tricoli, D.M., Cook, D.R., Ehrhardt, D.W., and Long, S.R. (2011). Symbiotic rhizobia bacteria trigger a change in localization and dynamics of the *Medicago truncatula* receptor kinase LYK3. *Plant Cell* **23**: 2774–2787.
- Hayashi, M., et al. (2014). A thaumatin-like protein, *Rj4*, controls nodule symbiotic specificity in soybean. *Plant Cell Physiol.* **55**: 1679–1689.
- Held, M., Hou, H., Miri, M., Huynh, C., Ross, L., Hossain, M.S., Sato, S., Tabata, S., Perry, J., Wang, T.L., and Szczyglowski, K. (2014). *Lotus japonicus* cytokinin receptors work partially redundantly to mediate nodule formation. *Plant Cell* **26**: 678–694.
- Hirsch, S., Kim, J., Muñoz, A., Heckmann, A.B., Downie, J.A., and Oldroyd, G.E. (2009). GRAS proteins form a DNA binding complex to induce gene expression during nodulation signaling in *Medicago truncatula*. *Plant Cell* **21**: 545–557.
- Hohnjec, N., Perlick, A.M., Pühler, A., and Küster, H. (2003). The *Medicago truncatula* sucrose synthase gene *MtSucS1* is activated both in the infected region of root nodules and in the cortex of roots colonized by arbuscular mycorrhizal fungi. *Mol. Plant Microbe Interact.* **16**: 903–915.
- Horst, I., Welham, T., Kelly, S., Kaneko, T., Sato, S., Tabata, S., Parniske, M., and Wang, T.L. (2007). TILLING mutants of *Lotus japonicus* reveal that nitrogen assimilation and fixation can occur in the absence of nodule-enhanced sucrose synthase. *Plant Physiol.* **144**: 806–820.
- Horváth, B., et al. (2015). Loss of the nodule-specific cysteine rich peptide, *NCR169*, abolishes symbiotic nitrogen fixation in the *Medicago truncatula dnf7* mutant. *Proc. Natl. Acad. Sci. USA* **112**: 15232–15237.
- Hossain, M.S., Liao, J., James, E.K., Sato, S., Tabata, S., Jurkiewicz, A., Madsen, L.H., Stougaard, J., Ross, L., and Szczyglowski, K. (2012). *Lotus japonicus ARPC1* is required for rhizobial infection. *Plant Physiol.* **160**: 917–928.
- Huault, E., Laffont, C., Wen, J., Mysore, K.S., Ratet, P., Duc, G., and Frugier, F. (2014). Local and systemic regulation of plant root

- system architecture and symbiotic nodulation by a receptor-like kinase. *PLoS Genet.* **10**: e1004891.
- Huo, X., Schnabel, E., Hughes, K., and Frugoli, J.** (2006). RNAi phenotypes and the localization of a protein:GUS fusion imply a role for *Medicago truncatula* PIN genes in nodulation. *J. Plant Growth Regul.* **25**: 156–165.
- Imaizumi-Anraku, H., et al.** (2005). Plastid proteins crucial for symbiotic fungal and bacterial entry into plant roots. *Nature* **433**: 527–531.
- Imin, N., Mohd-Radzman, N.A., Ogilvie, H.A., and Djordjevic, M.A.** (2013). The peptide-encoding CEP1 gene modulates lateral root and nodule numbers in *Medicago truncatula*. *J. Exp. Bot.* **64**: 5395–5409.
- Imin, N., Patel, N., Corcilius, L., Payne, R.J., and Djordjevic, M.A.** (2018). CLE peptide tri-arabinylation and peptide domain sequence composition are essential for SUNN-dependent autoregulation of nodulation in *Medicago truncatula*. *New Phytol.* **218**: 73–80.
- Ivanov, S., Fedorova, E.E., Limpens, E., De Mita, S., Genre, A., Bonfante, P., and Bisseling, T.** (2012). Rhizobium-legume symbiosis shares an exocytotic pathway required for arbuscule formation. *Proc. Natl. Acad. Sci. USA* **109**: 8316–8321.
- Ivashuta, S., Liu, J., Liu, J., Lohar, D.P., Haridas, S., Bucciarelli, B., VandenBosch, K.A., Vance, C.P., Harrison, M.J., and Gantt, J.S.** (2005). RNA interference identifies a calcium-dependent protein kinase involved in *Medicago truncatula* root development. *Plant Cell* **17**: 2911–2921.
- Jin, Y., Chen, Z., Yang, J., Mysore, K.S., Wen, J., Huang, J., Yu, N., and Wang, E.** (2018). IPD3 and IPD3L function redundantly in rhizobial and mycorrhizal symbioses. *Front. Plant Sci.* **9**: 267.
- Jin, Y., Liu, H., Luo, D., Yu, N., Dong, W., Wang, C., Zhang, X., Dai, H., Yang, J., and Wang, E.** (2016). DELLA proteins are common components of symbiotic rhizobial and mycorrhizal signalling pathways. *Nat. Commun.* **7**: 12433.
- Kaló, P., et al.** (2005). Nodulation signaling in legumes requires NSP2, a member of the GRAS family of transcriptional regulators. *Science* **308**: 1786–1789.
- Kanamori, N., et al.** (2006). A nucleoporin is required for induction of Ca<sup>2+</sup> spiking in legume nodule development and essential for rhizobial and fungal symbiosis. *Proc. Natl. Acad. Sci. USA* **103**: 359–364.
- Kawaharada, Y., James, E.K., Kelly, S., Sandal, N., and Stougaard, J.** (2017a). The ethylene responsive factor required for nodulation 1 (ERN1) transcription factor is required for infection-thread formation in *Lotus japonicus*. *Mol. Plant Microbe Interact.* **30**: 194–204.
- Kawaharada, Y., et al.** (2015). Receptor-mediated exopolysaccharide perception controls bacterial infection. *Nature* **523**: 308–312.
- Kawaharada, Y., Nielsen, M.W., Kelly, S., James, E.K., Andersen, K.R., Rasmussen, S.R., Füchtbauer, W., Madsen, L.H., Heckmann, A.B., Radutoiu, S., and Stougaard, J.** (2017b). Differential regulation of the *Epr3* receptor coordinates membrane-restricted rhizobial colonization of root nodule primordia. *Nat. Commun.* **8**: 14534.
- Ke, D., Fang, Q., Chen, C., Zhu, H., Chen, T., Chang, X., Yuan, S., Kang, H., Ma, L., Hong, Z., and Zhang, Z.** (2012). The small GTPase ROP6 interacts with NFR5 and is involved in nodule formation in *Lotus japonicus*. *Plant Physiol.* **159**: 131–143.
- Kevei, Z., et al.** (2007). 3-Hydroxy-3-methylglutaryl coenzyme A reductase 1 interacts with NORL and is crucial for nodulation in *Medicago truncatula*. *Plant Cell* **19**: 3974–3989.
- Kim, G.-B., Son, S.-U., Yu, H.-J., and Mun, J.-H.** (2019). *MtGA2ox10* encoding C20-GA2-oxidase regulates rhizobial infection and nodule development in *Medicago truncatula*. *Sci. Rep.* **9**: 5952.
- Kim, M., Chen, Y., Xi, J., Waters, C., Chen, R., and Wang, D.** (2015). An antimicrobial peptide essential for bacterial survival in the nitrogen-fixing symbiosis. *Proc. Natl. Acad. Sci. USA* **112**: 15238–15243.
- Kim, Y.-K., et al.** (2013). Functional implication of  $\beta$ -carotene hydroxylases in soybean nodulation. *Plant Physiol.* **162**: 1420–1433.
- Kiss, E., Oláh, B., Kaló, P., Morales, M., Heckmann, A.B., Borbola, A., Lózsa, A., Kontár, K., Middleton, P., Downie, J.A., Oldroyd, G.E., and Andre, G.** (2009). LIN, a novel type of U-box/WD40 protein, controls early infection by rhizobia in legumes. *Plant Physiol.* **151**: 1239–1249.
- Kosuta, S., Held, M., Hossain, M.S., Morieri, G., Macgillivray, A., Johansen, C., Antolín-Llovera, M., Parniske, M., Oldroyd, G.E., Downie, A.J., Karas, B., and Szczyglowski, K.** (2011). *Lotus japonicus symRK-14* uncouples the cortical and epidermal symbiotic program. *Plant J.* **67**: 929–940.
- Krusell, L., et al.** (2005). The sulfate transporter SST1 is crucial for symbiotic nitrogen fixation in *Lotus japonicus* root nodules. *Plant Cell* **17**: 1625–1636.
- Krusell, L., et al.** (2002). Shoot control of root development and nodulation is mediated by a receptor-like kinase. *Nature* **420**: 422–426.
- Krusell, L., et al.** (2011). The *Clavata2* genes of pea and *Lotus japonicus* affect autoregulation of nodulation. *Plant J.* **65**: 861–871.
- Kryvoruchko, I.S., Routray, P., Sinharoy, S., Torres-Jerez, I., Tejada-Jiménez, M., Finney, L.A., Nakashima, J., Pislariu, C.I., Benedito, V.A., González-Guerrero, M., Roberts, D.M., and Udvardi, M.K.** (2018). An iron-activated citrate transporter, MtMATE67, is required for symbiotic nitrogen fixation. *Plant Physiol.* **176**: 2315–2329.
- Kryvoruchko, I.S., Sinharoy, S., Torres-Jerez, I., Sosso, D., Pislariu, C.I., Guan, D., Murray, J., Benedito, V.A., Frommer, W.B., and Udvardi, M.K.** (2016). MtSWEET11, a nodule-specific sucrose transporter of *Medicago truncatula*. *Plant Physiol.* **171**: 554–565.
- Kuppusamy, K.T., Andre, G., Prabhu, R., Penmetsa, R.V., Veereshlingam, H., Cook, D.R., Dickstein, R., and Vandenbosch, K.A.** (2004). *LIN*, a *Medicago truncatula* gene required for nodule differentiation and persistence of rhizobial infections. *Plant Physiol.* **136**: 3682–3691.
- Kuppusamy, K.T., Ivashuta, S., Bucciarelli, B., Vance, C.P., Gantt, J.S., and Vandenbosch, K.A.** (2009). Knockdown of *CELL DIVISION CYCLE16* reveals an inverse relationship between lateral root and nodule numbers and a link to auxin in *Medicago truncatula*. *Plant Physiol.* **151**: 1155–1166.
- Laffont, C., Huault, E., Gautrat, P., Andre, G., Kalo, P., Bourion, V., Duc, G., and Frugier, F.** (2019). Independent regulation of symbiotic nodulation by the SUNN negative and CRA2 positive systemic pathways. In *Plant Physiol.*, Volume **180**, pp. 559–570.
- Laporte, P., Lepage, A., Fournier, J., Catrice, O., Moreau, S., Jardinaud, M.-F., Mun, J.-H., Larrainzar, E., Cook, D.R., Gamas, P., and Niebel, A.** (2014). The CCAAT box-binding transcription factor NF-YA1 controls rhizobial infection. *J. Exp. Bot.* **65**: 481–494.
- Laporte, P., Satiat-Jeunemaitre, B., Velasco, I., Csorba, T., Van de Velde, W., Campalans, A., Burgyan, J., Arevalo-Rodriguez, M., and Crespi, M.** (2010). A novel RNA-binding peptide regulates the establishment of the *Medicago truncatula*-*Sinorhizobium meliloti* nitrogen-fixing symbiosis. *Plant J.* **62**: 24–38.
- Larrainzar, E., et al.** (2015). Deep sequencing of the *Medicago truncatula* root transcriptome reveals a massive and early interaction between nodulation factor and ethylene signals. *Plant Physiol.* **169**: 233–265.
- Lefebvre, B., et al.** (2010). A remorin protein interacts with symbiotic receptors and regulates bacterial infection. *Proc. Natl. Acad. Sci. USA* **107**: 2343–2348.

- León-Mediavilla, J., Senovilla, M., Montiel, J., Gil-Díez, P., Saez, Á., Kryvoruchko, I.S., Reguera, M., Udvardi, M.K., Imperial, J., and González-Guerrero, M. (2018). MtMTP2-facilitated zinc transport into intracellular compartments is essential for nodule development in *Medicago truncatula*. *Front. Plant Sci.* **9**: 990.
- Lerouge, P., Roche, P., Faucher, C., Maillet, F., Truchet, G., Promé, J.C., and Dénarié, J. (1990). Symbiotic host-specificity of *Rhizobium meliloti* is determined by a sulphated and acylated glucosamine oligosaccharide signal. *Nature* **344**: 781–784.
- Le Roux, M., Khan, S., and Valentine, A. (2009). Nitrogen and carbon costs of soybean and lupin root systems during phosphate starvation. *Symbiosis* **48**: 102–109.
- Li, X., Feng, H., Wen, J., Dong, J., and Wang, T. (2018). MtCAS31 aids symbiotic nitrogen fixation by protecting the leghemoglobin MtLb120-1 under drought stress in *Medicago truncatula*. *Front. Plant Sci.* **9**: 633.
- Liang, P., Stratil, T.F., Popp, C., Marín, M., Folgmann, J., Mysore, K.S., Wen, J., and Ott, T. (2018). Symbiotic root infections in *Medicago truncatula* require remorin-mediated receptor stabilization in membrane nanodomains. *Proc. Natl. Acad. Sci. USA* **115**: 5289–5294.
- Lim, C.W., Lee, Y.W., and Hwang, C.H. (2011). Soybean nodule-enhanced CLE peptides in roots act as signals in GmNARK-mediated nodulation suppression. *Plant Cell Physiol.* **52**: 1613–1627.
- Limpens, E., Franken, C., Smit, P., Willemse, J., Bisseling, T., and Geurts, R. (2003). LysM domain receptor kinases regulating rhizobial Nod factor-induced infection. *Science* **302**: 630–633.
- Limpens, E., Ivanov, S., van Esse, W., Voets, G., Fedorova, E., and Bisseling, T. (2009). *Medicago* N<sub>2</sub>-fixing symbiosomes acquire the endocytic identity marker *Rab7* but delay the acquisition of vacuolar identity. *Plant Cell* **21**: 2811–2828.
- Liu, C.-W., et al. (2019a). A protein complex required for polar growth of rhizobial infection threads. *Nat Commun* **10**: 2848.
- Liu, C.-W., et al. (2019b). NIN acts as a network hub controlling a growth module required for rhizobial infection. *Plant Physiol.* **179**: 1704–1722.
- Liu, C.-W., and Murray, J.D. (2016). The role of flavonoids in nodulation host-range specificity: An update. *Plants (Basel)* **5**: 33.
- Liu, H., Zhang, C., Yang, J., Yu, N., and Wang, E. (2018). Hormone modulation of legume-rhizobial symbiosis. *J. Integr. Plant Biol.* **60**: 632–648.
- Liu, J., Novero, M., Charnikhova, T., Ferrandino, A., Schubert, A., Ruyter-Spira, C., Bonfante, P., Lovisolo, C., Bouwmeester, H.J., and Cardinale, F. (2013). Carotenoid cleavage dioxygenase 7 modulates plant growth, reproduction, senescence, and determinate nodulation in the model legume *Lotus japonicus*. *J. Exp. Bot.* **64**: 1967–1981.
- Liu, J., Rutten, L., Limpens, E., van der Molen, T., van Velzen, R., Chen, R., Chen, Y., Geurts, R., Kohlen, W., Kulikova, O., and Bisseling, T. (2019c). A remote cis-regulatory region is required for NIN expression in the pericycle to initiate nodule primordium formation in *Medicago truncatula*. *Plant Cell* **31**: 68–83.
- Long, S.R. (1996). Rhizobium symbiosis: Nod factors in perspective. *Plant Cell* **8**: 1885–1898.
- López-Ráez, J.A., Shirasu, K., and Foo, E. (2017). Strigolactones in plant interactions with beneficial and detrimental organisms: The yin and yang. *Trends Plant Sci.* **22**: 527–537.
- Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M., Szczygłowski, K., Sato, S., Kaneko, T., Tabata, S., Sandal, N., and Stougaard, J. (2003). A receptor kinase gene of the LysM type is involved in legume perception of rhizobial signals. *Nature* **425**: 637–640.
- Maekawa-Yoshikawa, M., Müller, J., Takeda, N., Maekawa, T., Sato, S., Tabata, S., Perry, J., Wang, T.L., Groth, M., Brachmann, A., and Parniske, M. (2009). The temperature-sensitive *brush* mutant of the legume *Lotus japonicus* reveals a link between root development and nodule infection by rhizobia. *Plant Physiol.* **149**: 1785–1796.
- Magori, S., and Kawaguchi, M. (2009). Long-distance control of nodulation: Molecules and models. *Mol. Cells* **27**: 129–134.
- Malolepszy, A., et al. (2018). A plant chitinase controls cortical infection thread progression and nitrogen-fixing symbiosis. *eLife* **7**: e38874.
- Malolepszy, A., et al. (2016). The LORE1 insertion mutant resource. *Plant J.* **88**: 306–317.
- Marino, D., Andrio, E., Danchin, E.G.J., Oger, E., Gucciardo, S., Lambert, A., Puppo, A., and Pauly, N. (2011). A *Medicago truncatula* NADPH oxidase is involved in symbiotic nodule functioning. *New Phytol.* **189**: 580–592.
- Mbengue, M., Camut, S., de Carvalho-Niebel, F., Deslandes, L., Froidure, S., Klaus-Heisen, D., Moreau, S., Rivas, S., Timmers, T., Hervé, C., Cullimore, J., and Lefebvre, B. (2010). The *Medicago truncatula* E3 ubiquitin ligase PUB1 interacts with the LYK3 symbiotic receptor and negatively regulates infection and nodulation. *Plant Cell* **22**: 3474–3488.
- McAdam, E.L., Reid, J.B., and Foo, E. (2018). Gibberellins promote nodule organogenesis but inhibit the infection stages of nodulation. *J. Exp. Bot.* **69**: 2117–2130.
- Mellor, R.B. (1989). Bacteroids in the Rhizobium-legume symbiosis inhabit a plant internal lytic compartment: implications for other microbial endosymbioses. *J. Exp. Bot.* **40**: 831–839.
- Mendel, G. (1866). Experiments in plant hybridization. In *Proceedings of the Natural History Society of Brno, Volume IV (Biodiversity Heritage Library)*, pp. 3–47.
- Mergaert, P., Uchiumi, T., Alunni, B., Evanno, G., Cheron, A., Catrice, O., Mausset, A.-E., Barloy-Hubler, F., Galibert, F., Kondorosi, A., and Kondorosi, E. (2006). Eukaryotic control on bacterial cell cycle and differentiation in the Rhizobium-legume symbiosis. *Proc. Natl. Acad. Sci. USA* **103**: 5230–5235.
- Middleton, P.H., et al. (2007). An ERF transcription factor in *Medicago truncatula* that is essential for Nod factor signal transduction. *Plant Cell* **19**: 1221–1234.
- Miri, M., Janakirama, P., Huebert, T., Ross, L., McDowell, T., Orosz, K., Markmann, K., and Szczygłowski, K. (2019). Inside out: Root cortex-localized LHK1 cytokinin receptor limits epidermal infection of *Lotus japonicus* roots by *Mesorhizobium loti*. *New Phytol.* **222**: 1523–1537.
- Miyata, K., Kawaguchi, M., and Nakagawa, T. (2013). Two distinct *EIN2* genes cooperatively regulate ethylene signaling in *Lotus japonicus*. *Plant Cell Physiol.* **54**: 1469–1477.
- Miyata, K., et al. (2014). The bifunctional plant receptor, OsCERK1, regulates both chitin-triggered immunity and arbuscular mycorrhizal symbiosis in rice. *Plant Cell Physiol.* **55**: 1864–1872.
- Miyazawa, H., et al. (2010). The receptor-like kinase KLAVER mediates systemic regulation of nodulation and non-symbiotic shoot development in *Lotus japonicus*. *Development* **137**: 4317–4325.
- Mohd-Radzman, N.A., Laffont, C., Ivanovici, A., Patel, N., Reid, D., Stougaard, J., Frugier, F., Imin, N., and Djordjevic, M.A. (2016). Different pathways act downstream of the CEP peptide receptor CRA2 to regulate lateral root and nodule development. *Plant Physiol.* **171**: 2536–2548.
- Moling, S., Pietraszewska-Bogiel, A., Postma, M., Fedorova, E., Hink, M.A., Limpens, E., Gadella, T.W., and Bisseling, T. (2014). Nod factor receptors form heteromeric complexes and are essential for intracellular infection in *Medicago* nodules. *Plant Cell* **26**: 4188–4199.
- Morell, M., and Copeland, L. (1984). Enzymes of sucrose breakdown in soybean nodules: Alkaline invertase. *Plant Physiol.* **74**: 1030–1034.

- Mortier, V., Den Herder, G., Whitford, R., Van de Velde, W., Rombauts, S., D'Haeseleer, K., Holsters, M., and Goormachtig, S. (2010). CLE peptides control *Medicago truncatula* nodulation locally and systemically. *Plant Physiol.* **153**: 222–237.
- Mortier, V., De Wever, E., Vuylsteke, M., Holsters, M., and Goormachtig, S. (2012). Nodule numbers are governed by interaction between CLE peptides and cytokinin signaling. *Plant J.* **70**: 367–376.
- Mortier, V., Wasson, A., Jaworek, P., De Keyser, A., Decroos, M., Holsters, M., Tarkowski, P., Mathesius, U., and Goormachtig, S. (2014). Role of *LONELY GUY* genes in indeterminate nodulation on *Medicago truncatula*. *New Phytol.* **202**: 582–593.
- Mun, T., Bachmann, A., Gupta, V., Stougaard, J., and Andersen, S.U. (2016). Lotus Base: An integrated information portal for the model legume *Lotus japonicus*. *Sci. Rep.* **6**: 39447.
- Murakami, E., et al. (2018). Epidermal LysM receptor ensures robust symbiotic signalling in *Lotus japonicus*. *eLife* **7**: e33506.
- Murray, J.D., Karas, B.J., Sato, S., Tabata, S., Amyot, L., and Szczyglowski, K. (2007). A cytokinin perception mutant colonized by *Rhizobium* in the absence of nodule organogenesis. *Science* **315**: 101–104.
- Murray, J.D., et al. (2011). *Vapyrin*, a gene essential for intracellular progression of arbuscular mycorrhizal symbiosis, is also essential for infection by rhizobia in the nodule symbiosis of *Medicago truncatula*. *Plant J.* **65**: 244–252.
- Nanjareddy, K., Blanco, L., Arthikala, M.-K., Alvarado-Affantranger, X., Quinto, C., Sánchez, F., and Lara, M. (2016). A legume TOR protein kinase regulates rhizobium symbiosis and is essential for infection and nodule development. *Plant Physiol.* **172**: 2002–2020.
- Ng, J.L.P., and Mathesius, U. (2018). Acropetal auxin transport inhibition is involved in indeterminate but not determinate nodule formation. *Front. Plant Sci.* **9**: 169.
- Nishida, H., Tanaka, S., Handa, Y., Ito, M., Sakamoto, Y., Matsunaga, S., Betsuyaku, S., Miura, K., Soyano, T., Kawaguchi, M., and Suzaki, T. (2018). A NIN-LIKE PROTEIN mediates nitrate-induced control of root nodule symbiosis in *Lotus japonicus*. *Nat. Commun.* **9**: 499.
- Nishimura, R., Hayashi, M., Wu, G.-J., Kouchi, H., Imaizumi-Anraku, H., Murakami, Y., Kawasaki, S., Akao, S., Ohmori, M., Nagasawa, M., Harada, K., and Kawaguchi, M. (2002). HAR1 mediates systemic regulation of symbiotic organ development. *Nature* **420**: 426–429.
- Nizampatnam, N.R., Schreier, S.J., Damodaran, S., Adhikari, S., and Subramanian, S. (2015). microRNA160 dictates stage-specific auxin and cytokinin sensitivities and directs soybean nodule development. *Plant J.* **84**: 140–153.
- Nowak, S., Schnabel, E., and Frugoli, J. (2019). The *Medicago truncatula* CLAVATA3-LIKE CLE12/13 signaling peptides regulate nodule number depending on the CORYNE but not the COMPACT ROOT ARCHITECTURE2 receptor. *Plant Signal. Behav.* **14**: 1598730.
- Ogawa-Ohnishi, M., Matsushita, W., and Matsubayashi, Y. (2013). Identification of three hydroxyproline O-arabinosyltransferases in *Arabidopsis thaliana*. *Nat. Chem. Biol.* **9**: 726–730.
- Oka-Kira, E., et al. (2005). *klavier (klv)*, a novel hypernodulation mutant of *Lotus japonicus* affected in vascular tissue organization and floral induction. *Plant J.* **44**: 505–515.
- Okamoto, S., Ohnishi, E., Sato, S., Takahashi, H., Nakazono, M., Tabata, S., and Kawaguchi, M. (2009). Nod factor/nitrate-induced CLE genes that drive HAR1-mediated systemic regulation of nodulation. *Plant Cell Physiol.* **50**: 67–77.
- Okamoto, S., Shinohara, H., Mori, T., Matsubayashi, Y., and Kawaguchi, M. (2013). Root-derived CLE glycopeptides control nodulation by direct binding to HAR1 receptor kinase. *Nat. Commun.* **4**: 2191.
- Okamoto, S., Tabata, R., and Matsubayashi, Y. (2016). Long-distance peptide signaling essential for nutrient homeostasis in plants. *Curr. Opin. Plant Biol.* **34**: 35–40.
- Oldroyd, G.E. (2013). Speak, friend, and enter: Signalling systems that promote beneficial symbiotic associations in plants. *Nat. Rev. Microbiol.* **11**: 252–263.
- Oldroyd, G.E., Murray, J.D., Poole, P.S., and Downie, J.A. (2011). The rules of engagement in the legume-rhizobial symbiosis. *Annu. Rev. Genet.* **45**: 119–144.
- Ott, T., Sullivan, J., James, E.K., Flemetakis, E., Günther, C., Gibon, Y., Ronson, C., and Udvardi, M. (2009). Absence of symbiotic leghemoglobins alters bacteroid and plant cell differentiation during development of *Lotus japonicus* root nodules. *Mol. Plant Microbe Interact.* **22**: 800–808.
- Ott, T., van Dongen, J.T., Günther, C., Krusell, L., Desbrosses, G., Vigeolas, H., Bock, V., Czechowski, T., Geigenberger, P., and Udvardi, M.K. (2005). Symbiotic leghemoglobins are crucial for nitrogen fixation in legume root nodules but not for general plant growth and development. *Curr. Biol.* **15**: 531–535.
- Pan, H., Oztas, O., Zhang, X., Wu, X., Stonoha, C., Wang, E., Wang, B., and Wang, D. (2016). A symbiotic SNARE protein generated by alternative termination of transcription. *Nat. Plants* **2**: 15197.
- Penmetsa, R.V., and Cook, D.R. (1997). A legume ethylene-insensitive mutant hyperinfected by its rhizobial symbiont. *Science* **275**: 527–530.
- Penmetsa, R.V., and Cook, D.R. (2000). Production and characterization of diverse developmental mutants of *Medicago truncatula*. *Plant Physiol.* **123**: 1387–1398.
- Penmetsa, R.V., Frugoli, J.A., Smith, L.S., Long, S.R., and Cook, D.R. (2003). Dual genetic pathways controlling nodule number in *Medicago truncatula*. *Plant Physiol.* **131**: 998–1008.
- Penmetsa, R.V., et al. (2008). The *Medicago truncatula* ortholog of Arabidopsis EIN2, *sickle*, is a negative regulator of symbiotic and pathogenic microbial associations. *Plant J.* **55**: 580–595.
- Pérez Guerra, J.C., Coussens, G., De Keyser, A., De Rycke, R., De Bodt, S., Van De Velde, W., Goormachtig, S., and Holsters, M. (2010). Comparison of developmental and stress-induced nodule senescence in *Medicago truncatula*. *Plant Physiol.* **152**: 1574–1584.
- Perry, J.A., Wang, T.L., Welham, T.J., Gardner, S., Pike, J.M., Yoshida, S., and Parniske, M. (2003). A TILLING reverse genetics tool and a web-accessible collection of mutants of the legume *Lotus japonicus*. *Plant Physiol.* **131**: 866–871.
- Peters, N.K., Frost, J.W., and Long, S.R. (1986). A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. *Science* **233**: 977–980.
- Petrásek, J., and Schwarzerová, K. (2009). Actin and microtubule cytoskeleton interactions. *Curr. Opin. Plant Biol.* **12**: 728–734.
- Pierre, O., Hopkins, J., Combiér, M., Baldacci, F., Engler, G., Brouquisse, R., Hérouart, D., and Boncompagni, E. (2014). Involvement of papain and legumain proteinase in the senescence process of *Medicago truncatula* nodules. *New Phytol.* **202**: 849–863.
- Pii, Y., Molesini, B., Masiero, S., and Pandolfini, T. (2012). The non-specific lipid transfer protein N5 of *Medicago truncatula* is implicated in epidermal stages of rhizobium-host interaction. *BMC Plant Biol.* **12**: 233.
- Pislariu, C.I., et al. (2012). A *Medicago truncatula* tobacco retrotransposon insertion mutant collection with defects in nodule development and symbiotic nitrogen fixation. *Plant Physiol.* **159**: 1686–1699.
- Pumplin, N., Mondo, S.J., Topp, S., Starker, C.G., Gantt, J.S., and Harrison, M.J. (2010). *Medicago truncatula* Vapyrin is a novel protein required for arbuscular mycorrhizal symbiosis. *Plant J.* **61**: 482–494.

- Qin, L., Zhao, J., Tian, J., Chen, L., Sun, Z., Guo, Y., Lu, X., Gu, M., Xu, G., and Liao, H. (2012). The high-affinity phosphate transporter GmPT5 regulates phosphate transport to nodules and nodulation in soybean. *Plant Physiol.* **159**: 1634–1643.
- Qiu, L., Lin, J.S., Xu, J., Sato, S., Parniske, M., Wang, T.L., Downie, J.A., and Xie, F. (2015). SCARN a novel class of SCAR protein that is required for root-hair infection during legume nodulation. *PLoS Genet.* **11**: e1005623.
- Radutoiu, S., Madsen, L.H., Madsen, E.B., Felle, H.H., Umehara, Y., Grønlund, M., Sato, S., Nakamura, Y., Tabata, S., Sandal, N., and Stougaard, J. (2003). Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature* **425**: 585–592.
- Radutoiu, S., Madsen, L.H., Madsen, E.B., Jurkiewicz, A., Fukai, E., Quistgaard, E.M., Albrechtsen, A.S., James, E.K., Thirup, S., and Stougaard, J. (2007). LysM domains mediate lipochitin-oligosaccharide recognition and *Nfr* genes extend the symbiotic host range. *EMBO J.* **26**: 3923–3935.
- Ramu, S.K., Peng, H.-M., and Cook, D.R. (2002). Nod factor induction of reactive oxygen species production is correlated with expression of the early nodulin gene rip1 in *Medicago truncatula*. *Mol. Plant Microbe Interact.* **15**: 522–528.
- Redmond, J.W., Batley, M., Djordjevic, M.A., Innes, R.W., Kuempel, P.L., and Rolfe, B.G. (1986). Flavones induce expression of nodulation genes in *Rhizobium*. *Nature* **323**: 632–635.
- Reid, D., Liu, H., Kelly, S., Kawaharada, Y., Mun, T., Andersen, S.U., Desbrosses, G., and Stougaard, J. (2018). Dynamics of ethylene production in response to compatible Nod factor. *Plant Physiol.* **176**: 1764–1772.
- Reid, D.E., Ferguson, B.J., Hayashi, S., Lin, Y.-H., and Gresshoff, P.M. (2011). Molecular mechanisms controlling legume autoregulation of nodulation. *Ann. Bot.* **108**: 789–795.
- Reid, D.E., Heckmann, A.B., Novák, O., Kelly, S., and Stougaard, J. (2016). CYTOKININ OXIDASE/DEHYDROGENASE3 maintains cytokinin homeostasis during root and nodule development in *Lotus japonicus*. *Plant Physiol.* **170**: 1060–1074.
- Reyero-Saavedra, M.D.R., Qiao, Z., Sánchez-Correa, M.D.S., Díaz-Pineda, M.E., Reyes, J.L., Covarrubias, A.A., Libault, M., and Valdés-López, O. (2017). Gene silencing of *argonaute5* negatively affects the establishment of the legume-rhizobia symbiosis. *Genes (Basel)* **8**: 352.
- Ripodas, C., Castaingts, M., Clúa, J., Villafañe, J., Blanco, F.A., and Zanetti, M.E. (2019). The PvNF-YA1 and PvNF-YB7 subunits of the heterotrimeric NF-Y transcription factor influence strain preference in the *Phaseolus vulgaris*-*Rhizobium etli* symbiosis. *Front. Plant Sci.* **10**: 221.
- Rodríguez-Haas, B., Finney, L., Vogt, S., González-Melendi, P., Imperial, J., and González-Guerrero, M. (2013). Iron distribution through the developmental stages of *Medicago truncatula* nodules. *Metallomics* **5**: 1247–1253.
- Rogato, A., D'Apuzzo, E., Barbulova, A., Omrane, S., Stedel, C., Simon-Rosin, U., Katinakis, P., Flemetakis, M., Udvardi, M., and Chiurazzi, M. (2008). Tissue-specific down-regulation of *LjAMT1*;1 compromises nodule function and enhances nodulation in *Lotus japonicus*. *Plant Mol. Biol.* **68**: 585–595.
- Roth, L.E., and Stacey, G. (1989). Bacterium release into host cells of nitrogen-fixing soybean nodules: The symbiosome membrane comes from three sources. *Eur. J. Cell Biol.* **49**: 13–23.
- Roy, S., et al. (2017). *MtLAX2*, a functional homologue of the Arabidopsis auxin influx transporter AUX1, is required for nodule organogenesis. *Plant Physiol.* **174**: 326–338.
- Rubio, M.C., Becana, M., Sato, S., James, E.K., Tabata, S., and Spaink, H.P. (2007). Characterization of genomic clones and expression analysis of the three types of superoxide dismutases during nodule development in *Lotus japonicus*. *Mol. Plant Microbe Interact.* **20**: 262–275.
- Sadowsky, M.J., and Cregan, P.B. (1992). The soybean *Rj4* allele restricts nodulation by *Bradyrhizobium japonicum* serogroup 123 strains. *Appl. Environ. Microbiol.* **58**: 720–723.
- Saito, K., et al. (2007). NUCLEOPORIN85 is required for calcium spiking, fungal and bacterial symbioses, and seed production in *Lotus japonicus*. *Plant Cell* **19**: 610–624.
- Satgé, C., Moreau, S., Sallet, E., Lefort, G., Auriac, M.-C., Remblière, C., Cottret, L., Gallardo, K., Noiro, C., Jardinaud, M.-F., and Gamas, P. (2016). Reprogramming of DNA methylation is critical for nodule development in *Medicago truncatula*. *Nat. Plants* **2**: 16166.
- Schauser, L., Handberg, K., Sandal, N., Stiller, J., Thykjaer, T., Pajuelo, E., Nielsen, A., and Stougaard, J. (1998). Symbiotic mutants deficient in nodule establishment identified after T-DNA transformation of *Lotus japonicus*. *Mol. Gen. Genet.* **259**: 414–423.
- Schauser, L., Roussis, A., Stiller, J., and Stougaard, J. (1999). A plant regulator controlling development of symbiotic root nodules. *Nature* **402**: 191–195.
- Schippers, J.H., Schmidt, R., Wagstaff, C., and Jing, H.-C. (2015). Living to die and dying to live: The survival strategy behind leaf senescence. *Plant Physiol.* **169**: 914–930.
- Schnabel, E., Journet, E.-P., de Carvalho-Niebel, F., Duc, G., and Frugoli, J. (2005). The *Medicago truncatula* *SUNN* gene encodes a CLV1-like leucine-rich repeat receptor kinase that regulates nodule number and root length. *Plant Mol. Biol.* **58**: 809–822.
- Schnabel, E.L., Kassaw, T.K., Smith, L.S., Marsh, J.F., Oldroyd, G.E., Long, S.R., and Frugoli, J.A. (2011). The *ROOT DETERMINED NODULATION1* gene regulates nodule number in roots of *Medicago truncatula* and defines a highly conserved, uncharacterized plant gene family. *Plant Physiol.* **157**: 328–340.
- Searle, I.R., Men, A.E., Laniya, T.S., Buzas, D.M., Iturbe-Ormaetxe, I., Carroll, B.J., and Gresshoff, P.M. (2003). Long-distance signaling in nodulation directed by a CLAVATA1-like receptor kinase. *Science* **299**: 109–112.
- Senovilla, M., Castro-Rodríguez, R., Abreu, I., Escudero, V., Kryvoruchko, I., Udvardi, M.K., Imperial, J., and González-Guerrero, M. (2018). *Medicago truncatula* copper transporter 1 (*MtCOPT1*) delivers copper for symbiotic nitrogen fixation. *New Phytol.* **218**: 696–709.
- Shimomura, K., Nomura, M., Tajima, S., and Kouchi, H. (2006). LjnsRING, a novel RING finger protein, is required for symbiotic interactions between *Mesorhizobium loti* and *Lotus japonicus*. *Plant Cell Physiol.* **47**: 1572–1581.
- Sidiropoulos, K., et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics* **33**: 3461–3467.
- Singh, S., Katzer, K., Lambert, J., Cerri, M., and Parniske, M. (2014). CYCLOPS, a DNA-binding transcriptional activator, orchestrates symbiotic root nodule development. *Cell Host Microbe* **15**: 139–152.
- Sinharoy, S., Liu, C., Breakspear, A., Guan, D., Shailes, S., Nakashima, J., Zhang, S., Wen, J., Torres-Jerez, I., Oldroyd, G., Murray, J.D., and Udvardi, M.K. (2016). A *Medicago truncatula* cystathionine- $\beta$ -synthase-like domain-containing protein is required for rhizobial infection and symbiotic nitrogen fixation. *Plant Physiol.* **170**: 2204–2217.
- Sinharoy, S., Torres-Jerez, I., Bandyopadhyay, K., Kereszt, A., Pislariu, C.I., Nakashima, J., Benedito, V.A., Kondorosi, E., and Udvardi, M.K. (2013). The C2H2 transcription factor regulator of symbiosome differentiation represses transcription of the secretory pathway gene *VAMP721a* and promotes symbiosome development in *Medicago truncatula*. *Plant Cell* **25**: 3584–3601.



- Smit, P., Limpens, E., Geurts, R., Fedorova, E., Dolgikh, E., Gough, C., and Bisseling, T. (2007). Medicago LYK3, an entry receptor in rhizobial nodulation factor signaling. *Plant Physiol.* **145**: 183–191.
- Smit, P., Raedts, J., Portyanko, V., Debellé, F., Gough, C., Bisseling, T., and Geurts, R. (2005). NSP1 of the GRAS protein family is essential for rhizobial Nod factor-induced transcription. *Science* **308**: 1789–1791.
- Smith, P.M., and Atkins, C.A. (2002). Purine biosynthesis: Big in cell division, even bigger in nitrogen assimilation. *Plant Physiol.* **128**: 793–802.
- Sogawa, A., Yamazaki, A., Yamasaki, H., Komi, M., Manabe, T., Tajima, S., Hayashi, M., and Nomura, M. (2019). SNARE proteins LjVAMP72a and LjVAMP72b are required for root symbiosis and root hair formation in *Lotus japonicus*. *Front. Plant Sci.* **9**: 1992.
- Soyano, T., Hirakawa, H., Sato, S., Hayashi, M., and Kawaguchi, M. (2014). Nodule inception creates a long-distance negative feedback loop involved in homeostatic regulation of nodule organ production. *Proc. Natl. Acad. Sci. USA* **111**: 14607–14612.
- Soyano, T., Kouchi, H., Hirota, A., and Hayashi, M. (2013). NODULE INCEPTION directly targets NF-Y subunit genes to regulate essential processes of root nodule development in *Lotus japonicus*. *PLoS Genet.* **9**: e1003352.
- Sprent, J.I., and James, E.K. (2007). Legume evolution: Where do nodules and mycorrhizas fit in? *Plant Physiol.* **144**: 575–581.
- Stracke, S., Kistner, C., Yoshida, S., Mulder, L., Sato, S., Kaneko, T., Tabata, S., Sandal, N., Stougaard, J., Szczyglowski, K., and Parniske, M. (2002). A plant receptor-like kinase required for both bacterial and fungal symbiosis. *Nature* **417**: 959–962.
- Subramanian, S., Stacey, G., and Yu, O. (2006). Endogenous isoflavones are essential for the establishment of symbiosis between soybean and *Bradyrhizobium japonicum*. *Plant J.* **48**: 261–273.
- Suganuma, N., Nakamura, Y., Yamamoto, M., Ohta, T., Koiwa, H., Akao, S., and Kawaguchi, M. (2003). The *Lotus japonicus Sen1* gene controls rhizobial differentiation into nitrogen-fixing bacteroids in nodules. *Mol. Genet. Genomics* **269**: 312–320.
- Sugawara, M., Takahashi, S., Umehara, Y., Iwano, H., Tsurumaru, H., Odake, H., Suzuki, Y., Kondo, H., Konno, Y., and Yamakawa, T. (2018). Variation in bradyrhizobial NopP effector determines symbiotic incompatibility with Rj2-soybeans via effector-triggered immunity. *Nat. Commun.* **9**: 3139.
- Sutton, M.A., Oenema, O., Erisman, J.W., Leip, A., van Grinsven, H., and Winiwarter, W. (2011). Too much of a good thing. *Nature* **472**: 159–161.
- Suzaki, T., Takeda, N., Nishida, H., Hoshino, M., Ito, M., Misawa, F., Handa, Y., Miura, K., and Kawaguchi, M. (2019). LACK OF SYMBIONT ACCOMMODATION controls intracellular symbiont accommodation in root nodule and arbuscular mycorrhizal symbiosis in *Lotus japonicus*. *PLoS Genet.* **15**: e1007865.
- Szczyglowski, K., Shaw, R.S., Wopereis, J., Copeland, S., Hamburger, D., Kasiborski, B., Dazzo, F.B., and de Bruijn, F.J. (1998). Nodule organogenesis and symbiotic mutants of the model legume *Lotus japonicus*. *Mol. Plant Microbe Interact.* **11**: 684–697.
- Tadege, M., Wen, J., He, J., Tu, H., Kwak, Y., Eschstruth, A., Cayrel, A., Andre, G., Zhao, P.X., Chabaud, M., Ratet, P., and Mysore, K.S. (2008). Large-scale insertional mutagenesis using the Tnt1 retrotransposon in the model legume *Medicago truncatula*. *Plant J.* **54**: 335–347.
- Takahara, M., et al. (2013). TOO MUCH LOVE, a novel Kelch repeat-containing F-box protein, functions in the long-distance regulation of the legume-Rhizobium symbiosis. *Plant Cell Physiol.* **54**: 433–447.
- Takanashi, K., Yokosho, K., Saeki, K., Sugiyama, A., Sato, S., Tabata, S., Ma, J.F., and Yazaki, K. (2013). LjMATE1: A citrate transporter responsible for iron supply to the nodule infection zone of *Lotus japonicus*. *Plant Cell Physiol.* **54**: 585–594.
- Tang, F., Yang, S., Liu, J., and Zhu, H. (2016). *Rj4*, a gene controlling nodulation specificity in soybeans, encodes a thaumatin-like protein but not the one previously reported. *Plant Physiol.* **170**: 26–32.
- Teillet, A., Garcia, J., de Billy, F., Gherardi, M., Huguet, T., Barker, D.G., de Carvalho-Niebel, F., and Journet, E.-P. (2008). *api*, a novel *Medicago truncatula* symbiotic mutant impaired in nodule primordium invasion. *Mol. Plant Microbe Interact.* **21**: 535–546.
- Tejada-Jiménez, M., Castro-Rodríguez, R., Kryvoruchko, I., Lucas, M.M., Udvardi, M., Imperial, J., and González-Guerrero, M. (2015). *Medicago truncatula* Natural Resistance-Associated Macrophage Protein1 is required for iron uptake by rhizobia-infected nodule cells. *Plant Physiol.* **168**: 258–272.
- Tejada-Jiménez, M., Gil-Díez, P., León-Mediavilla, J., Wen, J., Mysore, K.S., Imperial, J., and González-Guerrero, M. (2017). *Medicago truncatula* Molybdate Transporter type 1 (MtMOT1.3) is a plasma membrane molybdenum transporter required for nitrogenase activity in root nodules under molybdenum deficiency. *New Phytol.* **216**: 1223–1235.
- Theil, E.C. (2003). Ferritin: At the crossroads of iron and oxygen metabolism. *J. Nutr.* **133** (suppl. 1): 1549S–1553S.
- Timmer, A.C. (2008). The role of the plant cytoskeleton in the interaction between legumes and rhizobia. *J. Microsc.* **231**: 247–256.
- Tirichine, L., et al. (2006). Dereglulation of a Ca<sup>2+</sup>/calmodulin-dependent kinase leads to spontaneous nodule development. *Nature* **441**: 1153–1156.
- Tirichine, L., Sandal, N., Madsen, L.H., Radutoiu, S., Albrechtsen, A.S., Sato, S., Asamizu, E., Tabata, S., and Stougaard, J. (2007). A gain-of-function mutation in a cytokinin receptor triggers spontaneous root nodule organogenesis. *Science* **315**: 104–107.
- Tjepkema, J.D., and Yocum, C.S. (1974). Measurement of oxygen partial pressure within soybean nodules by oxygen microelectrodes. *Planta* **119**: 351–360.
- Truchet, G., Roche, P., Lerouge, P., Vasse, J., Camut, S., de Billy, F., Promé, J.-C., and Dénarié, J. (1991). Sulphated lipooligosaccharide signals of *Rhizobium meliloti* elicit root nodule organogenesis in alfalfa. *Nature* **351**: 670–673.
- Tsikou, D., Yan, Z., Holt, D.B., Abel, N.B., Reid, D.E., Madsen, L.H., Bhasin, H., Sexauer, M., Stougaard, J., and Markmann, K. (2018). Systemic control of legume susceptibility to rhizobial infection by a mobile microRNA. *Science* **362**: 233–236.
- Udvardi, M., and Poole, P.S. (2013). Transport and metabolism in legume-rhizobia symbioses. *Annu. Rev. Plant Biol.* **64**: 781–805.
- Valkov, V.T., Rogato, A., Alves, L.M., Sol, S., Noguero, M., Lérans, S., Lacombe, B., and Chiurazzi, M. (2017). The nitrate transporter family protein LjNPF8.6 controls the N-fixing nodule activity. *Plant Physiol.* **175**: 1269–1282.
- Van de Velde, W., Guerra, J.C., De Keyser, A., De Rycke, R., Rombauts, S., Maunoury, N., Mergaert, P., Kondorosi, E., Holsters, M., and Goormachtig, S. (2006). Aging in legume symbiosis: A molecular view on nodule senescence in *Medicago truncatula*. *Plant Physiol.* **141**: 711–720.
- Van de Velde, W., et al. (2010). Plant peptides govern terminal differentiation of bacteria in symbiosis. *Science* **327**: 1122–1126.
- van Spronsen, P.C., Bakhuizen, R., van Brussel, A.A., and Kijne, J.W. (1994). Cell wall degradation during infection thread formation by the root nodule bacterium *Rhizobium leguminosarum* is a two-step process. *Eur. J. Cell Biol.* **64**: 88–94.
- Veereshlingam, H., Haynes, J.G., Penmetsa, R.V., Cook, D.R., Sherrier, D.J., and Dickstein, R. (2004). *nip*, a symbiotic *Medicago truncatula* mutant that forms root nodules with aberrant infection threads and plant defense-like response. *Plant Physiol.* **136**: 3692–3702.
- Verma, D. (1992). Signals in root nodule organogenesis and endocytosis of Rhizobium. *Plant Cell* **4**: 373–382.

- Vernié, T., et al. (2016). PUB1 interacts with the receptor kinase DMI2 and negatively regulates rhizobial and arbuscular mycorrhizal symbioses through its ubiquitination activity in *Medicago truncatula*. *Plant Physiol.* **170**: 2312–2324.
- Vernié, T., Kim, J., Frances, L., Ding, Y., Sun, J., Guan, D., Niebel, A., Gifford, M.L., de Carvalho-Niebel, F., and Oldroyd, G.E. (2015). The NIN transcription factor coordinates diverse nodulation programs in different tissues of the *Medicago truncatula* root. *Plant Cell* **27**: 3410–3424.
- Vinardell, J.M., Fedorova, E., Cebolla, A., Kevei, Z., Horvath, G., Kelemen, Z., Tarayre, S., Roudier, F., Mergaert, P., Kondorosi, A., and Kondorosi, E. (2003). Endoreduplication mediated by the anaphase-promoting complex activator *CCS52A* is required for symbiotic cell differentiation in *Medicago truncatula* nodules. *Plant Cell* **15**: 2093–2105.
- Voroshilova, V.A., Demchenko, K.N., Brewin, N.J., Borisov, A.Y., and Tikhonovich, I.A. (2009). Initiation of a legume nodule with an indeterminate meristem involves proliferating host cells that harbour infection threads. *New Phytol.* **181**: 913–923.
- Wang, C., et al. (2016). NODULES WITH ACTIVATED DEFENSE 1 is required for maintenance of rhizobial endosymbiosis in *Medicago truncatula*. *New Phytol.* **212**: 176–191.
- Wang, C., Zhu, H., Jin, L., Chen, T., Wang, L., Kang, H., Hong, Z., and Zhang, Z. (2013). Splice variants of the *SIP1* transcripts play a role in nodule organogenesis in *Lotus japonicus*. *Plant Mol. Biol.* **82**: 97–111.
- Wang, D., Griffiths, J., Starker, C., Fedorova, E., Limpens, E., Ivanov, S., Bisseling, T., and Long, S. (2010). A nodule-specific protein secretory pathway required for nitrogen-fixing symbiosis. *Science* **327**: 1126–1129.
- Wang, L., Rubio, M.C., Xin, X., Zhang, B., Fan, Q., Wang, Q., Ning, G., Becana, M., and Duanmu, D. (2019). CRISPR/Cas9 knockout of leghemoglobin genes in *Lotus japonicus* uncovers their synergistic roles in symbiotic nitrogen fixation. *New Phytol.* **224**: 818–832.
- Wang, Q., et al. (2017). Host-secreted antimicrobial peptide enforces symbiotic selectivity in *Medicago truncatula*. *Proc. Natl. Acad. Sci. USA* **114**: 6854–6859.
- Wang, Y., Li, K., Chen, L., Zou, Y., Liu, H., Tian, Y., Li, D., Wang, R., Zhao, F., Ferguson, B.J., Gresshoff, P.M., and Li, X. (2015). MicroRNA167-directed regulation of the auxin response factors *GmARF8a* and *GmARF8b* is required for soybean nodulation and lateral root development. *Plant Physiol.* **168**: 984–999.
- Wang, Y., Wang, L., Zou, Y., Chen, L., Cai, Z., Zhang, S., Zhao, F., Tian, Y., Jiang, Q., Ferguson, B.J., Gresshoff, P.M., and Li, X. (2014). Soybean miR172c targets the repressive AP2 transcription factor *NNC1* to activate *ENOD40* expression and regulate nodule initiation. *Plant Cell* **26**: 4782–4801.
- Wasson, A.P., Pellerone, F.I., and Mathesius, U. (2006). Silencing the flavonoid pathway in *Medicago truncatula* inhibits root nodule formation and prevents auxin transport regulation by rhizobia. *Plant Cell* **18**: 1617–1629.
- Wienkoop, S., and Saalbach, G. (2003). Proteome analysis: Novel proteins identified at the peribacteroid membrane from *Lotus japonicus* root nodules. *Plant Physiol.* **131**: 1080–1090.
- Winzer, T., Bairl, A., Linder, M., Linder, D., Werner, D., and Müller, P. (1999). A novel 53-kDa nodulin of the symbiosome membrane of soybean nodules, controlled by *Bradyrhizobium japonicum*. *Mol. Plant Microbe Interact.* **12**: 218–226.
- Xiao, T.T., Schilderink, S., Moling, S., Deinum, E.E., Kondorosi, E., Franssen, H., Kulikova, O., Niebel, A., and Bisseling, T. (2014). Fate map of *Medicago truncatula* root nodules. *Development* **141**: 3517–3528.
- Xie, F., Murray, J.D., Kim, J., Heckmann, A.B., Edwards, A., Oldroyd, G.E., and Downie, J.A. (2012). Legume pectate lyase required for root infection by rhizobia. *Proc. Natl. Acad. Sci. USA* **109**: 633–638.
- Xue, Y.-B., Xiao, B.-X., Zhu, S.-N., Mo, X.-H., Liang, C.-Y., Tian, J., Liao, H., and Miriam, G. (2017). *GmPHR25*, a GmPHR member up-regulated by phosphate starvation, controls phosphate homeostasis in soybean. *J. Exp. Bot.* **68**: 4951–4967.
- Yamaya-Ito, H., Shimoda, Y., Hakoyama, T., Sato, S., Kaneko, T., Hossain, M.S., Shibata, S., Kawaguchi, M., Hayashi, M., Kouchi, H., and Umehara, Y. (2018). Loss-of-function of ASPARTIC PEPTIDASE NODULE-INDUCED 1 (*APN1*) in *Lotus japonicus* restricts efficient nitrogen-fixing symbiosis with specific *Mesorhizobium loti* strains. *Plant J.* **93**: 5–16.
- Yan, Q., Wang, L., and Li, X. (2018). GmBEHL1, a BES1/BZR1 family protein, negatively regulates soybean nodulation. *Sci. Rep.* **8**: 7614.
- Yang, S., Tang, F., Gao, M., Krishnan, H.B., and Zhu, H. (2010). *R* gene-controlled host specificity in the legume-rhizobia symbiosis. *Proc. Natl. Acad. Sci. USA* **107**: 18735–18740.
- Yang, S., Wang, Q., Fedorova, E., Liu, J., Qin, Q., Zheng, Q., Price, P.A., Pan, H., Wang, D., Griffiths, J.S., Bisseling, T., and Zhu, H. (2017). Microsymbiont discrimination mediated by a host-secreted peptide in *Medicago truncatula*. *Proc. Natl. Acad. Sci. USA* **114**: 6848–6853.
- Yano, K., et al. (2008). CYCLOPS, a mediator of symbiotic intracellular accommodation. *Proc. Natl. Acad. Sci. USA* **105**: 20540–20545.
- Yendrek, C.R., et al. (2010). A putative transporter is essential for integrating nutrient and hormone signaling with lateral root growth and nodule development in *Medicago truncatula*. *Plant J.* **62**: 100–112.
- Yokota, K., et al. (2009). Rearrangement of actin cytoskeleton mediates invasion of *Lotus japonicus* roots by *Mesorhizobium loti*. *Plant Cell* **21**: 267–284.
- Yoro, E., Nishida, H., Ogawa-Ohnishi, M., Yoshida, C., Suzaki, T., Matsubayashi, Y., and Kawaguchi, M. (2019). PLENTY, a hydroxyproline O-arabinoxyltransferase, negatively regulates root nodule symbiosis in *Lotus japonicus*. *J. Exp. Bot.* **70**: 507–517.
- Yoro, E., Suzaki, T., Toyokura, K., Miyazawa, H., Fukaki, H., and Kawaguchi, M. (2014). A positive regulator of nodule organogenesis, NODULE INCEPTION, acts as a negative regulator of rhizobial infection in *Lotus japonicus*. *Plant Physiol.* **165**: 747–758.
- Yoshida, C., Funayama-Noguchi, S., and Kawaguchi, M. (2010). *plenty*, a novel hypernodulation mutant in *Lotus japonicus*. *Plant Cell Physiol.* **51**: 1425–1435.
- Yuan, S., Zhu, H., Gou, H., Fu, W., Liu, L., Chen, T., Ke, D., Kang, H., Xie, Q., Hong, Z., and Zhang, Z. (2012). A ubiquitin ligase of symbiosis receptor kinase involved in nodule organogenesis. *Plant Physiol.* **160**: 106–117.
- Zanetti, M.E., Blanco, F.A., Beker, M.P., Battaglia, M., and Aguilar, O.M. (2010). A C subunit of the plant nuclear factor NF-Y required for rhizobial infection and nodule development affects partner selection in the common bean-*Rhizobium etli* symbiosis. *Plant Cell* **22**: 4142–4157.
- Zhang, C., Bousquet, A., and Harris, J.M. (2014). Abscisic acid and LATERAL ROOT ORGAN DEFECTIVE/NUMEROUS INFECTIONS AND POLYPHENOLICS modulate root elongation via reactive oxygen species in *Medicago truncatula*. *Plant Physiol.* **166**: 644–658.
- Zhang, J., Subramanian, S., Stacey, G., and Yu, O. (2009). Flavones and flavonols play distinct critical roles during nodulation of *Medicago truncatula* by *Sinorhizobium meliloti*. *Plant J.* **57**: 171–183.

- Zhang, X., Dong, W., Sun, J., Feng, F., Deng, Y., He, Z., Oldroyd, G.E., and Wang, E.** (2015). The receptor kinase CERK1 has dual functions in symbiosis and immunity signalling. *Plant J.* **81**: 258–267.
- Zhang, X., Han, L., Wang, Q., Zhang, C., Yu, Y., Tian, J., and Kong, Z.** (2019). The host actin cytoskeleton channels rhizobia release and facilitates symbiosome accommodation during nodulation in *Medicago truncatula*. *New Phytol.* **221**: 1049–1059.
- Zhou, C., et al.** (2011). From model to crop: Functional analysis of a *STAY-GREEN* gene in the model legume *Medicago truncatula* and effective use of the gene for alfalfa improvement. *Plant Physiol.* **157**: 1483–1496.
- Zhu, H., Chen, T., Zhu, M., Fang, Q., Kang, H., Hong, Z., and Zhang, Z.** (2008). A novel ARID DNA-binding protein interacts with SymRK and is expressed during early nodule development in *Lotus japonicus*. *Plant Physiol.* **148**: 337–347.
- Zipfel, C., and Oldroyd, G.E.** (2017). Plant signalling in symbiosis and immunity. *Nature* **543**: 328–336.