

Cell-free expression and characterization of multivalent rhamnose-binding lectins using bilayer interferometry

Katherine F. Warfel^{a,b,c}, Eugénie Laigre^d, Sarah E. Sobol^{a,b,c}, Emilie Gillon^e, Annabelle Varrot^e, Olivier Renaudet^d, Jerome Dejeu^{d,f}, Michael C. Jewett^{a,b,c,g,h,*}, Anne Imberty^{e,*}

^aDepartment of Chemical and Biological Engineering, Northwestern University, 2145 Sheridan Road, Technological Institute E136, Evanston, IL 60208, USA

^bChemistry of Life Processes Institute, Northwestern University, 2170 Campus Drive, Evanston, IL 60208, USA

^cCenter for Synthetic Biology, Northwestern University, 2145 Sheridan Road, Technological Institute E136, Evanston, IL 60208, USA

^dUniv. Grenoble Alpes, CNRS, DCM, UMR 5250, 570 Rue de la Chimie, 38000 Grenoble, France

^eUniv. Grenoble Alpes, CNRS, CERMAV, UPR5301, 601 Rue de la Chimie, 38000 Grenoble, France

^fFEMTO-ST Institute, CNRS UMR-6174, Université de Bourgogne Franche-Comté, F-25000 Besançon, France

^gRobert H. Lurie Comprehensive Cancer Center, Northwestern University, 676 North Saint Clair Street, Suite 1200, Chicago, IL 60611, USA

^hSimpson Querrey Institute, Northwestern University, 303 East Superior Street, Suite 11-131, Chicago, IL 60611, USA

*Address correspondence to:

1) Anne Imberty: anne.imberty@cermav.cnrs.fr and 2) Michael Jewett: m-jewett@northwestern.edu

Abstract

Lectins are important biological tools for binding glycans, but recombinant protein expression poses challenges for some lectin classes, limiting the pace of discovery and characterization. To discover and engineer lectins with new functions, workflows amenable to rapid expression and subsequent characterization are needed. Here, we present bacterial cell-free protein synthesis as a means for efficient, small-scale expression of multivalent, disulfide bond-rich, rhamnose-binding lectins. Furthermore, we demonstrate that the cell-free expressed lectins can be directly coupled with BLI analysis, either in solution or immobilized on the sensor, to measure interaction with carbohydrate ligands without purification. This workflow enables determination of lectin substrate specificity and estimation of binding affinity. Overall, we believe that this method will enable high-throughput expression, screening, and characterization of new and engineered multivalent lectins for applications in synthetic glycobiology.

Introduction

Lectins are proteins that recognize and bind specific glycans, making them interesting scaffolds for therapeutics, diagnostics and quality control reagents for glycoprotein products (Arnaud et al. 2013; Fernandez-Poza et al. 2021). However, recombinant expression of lectins can be challenging due to properties such as toxicity, size or presence of disulfide bonds (Martínez-Alarcón et al. 2018). One strategy to overcome these challenges is cell-free expression (CFE), which harnesses biological machinery to enable high-yielding transcription and translation outside of the living cell (Carlson et al. 2012; Silverman et al. 2020). The modular and open CFE reaction environment allows for manipulation of expression conditions, enabling production of complex products including proteins containing disulfide bonds (Goerke and Swartz 2008; Dopp and Reuel 2020), membrane proteins (Matthies et al. 2011; Kruyer et al. 2021), and glycosylated proteins (Kightlinger et al. 2019; Hershewe et al. 2021; Stark et al. 2021; Jaroentomeechai et al. 2022). CFE is also scalable from the nanoliter to liter scale (Zawada et al. 2011), allowing for small-scale parallel expression of many proteins simply by switching out the template DNA added to the reaction, and accelerating protein screening (Hunt et al. 2021; Hunt et al. 2022).

To take advantage of CFE for rapid lectin screening, it would be ideal to assess the functionality of produced lectins directly in the reaction without purification. Biolayer interferometry (BLI) is a technique that has been recently used to characterize protein binding parameters without a purification step, indicating possible compatibility with CFE reactions (Pogoutse et al. 2016). In addition, BLI has been used to study multivalent lectin-carbohydrate interactions in purified systems (Laigre et al. 2018; Picault et al. 2022). Notably, this technique requires less material than surface plasmon resonance (SPR) or isothermal titration calorimetry (ITC), is label free, and can be run in a plate-based format, making it more compatible with small-scale cell-free expression (Concepcion et al. 2009).

To demonstrate our cell-free workflow, we selected two model eukaryotic lectins, SUL-I (PDB-ID 5H4S) from *Toxopneustes pileolus* (sea urchin) venom and CSL3 (PDB-ID 2ZX2) from

Oncorhynchus keta (chum salmon) eggs, that are characterized, but are difficult to isolate naturally or to produce in reasonable quantities in *Escherichia coli* (Shiina et al. 2002; Hatakeyama et al. 2015). These dimeric lectins are of interest due to their binding affinity to the glycosphingolipid globotriaosylceramide (Gb3), a tumor associated glycolipid, as well as to rhamnose-containing bacterial O-antigens (Shiina et al. 2002; Siukstaite et al. 2021). Here, we show that these rhamnose-binding lectins are soluble and active when expressed in a bacterial cell-free system. Further, we show that cell-free expressed lectins are compatible with BLI binding assays in crude reactions without purification, with lectins either immobilized or in solution. We believe that this demonstration will enable future high-throughput screening and characterization of the specificity and affinity of predicted and engineered lectins.

Results

Cell-free expressed multivalent lectins are compatible with direct in-solution BLI analysis

Each SUL-I monomer presents 3 binding sites with 13 disulfide bonds, while each CSL3 monomer has 2 binding sites with 8 disulfide bonds and both of them are dimeric in solution (**Figure S1A-B**) (Shirai et al. 2009; Hatakeyama et al. 2017). To support proper protein folding, SUL-I and CSL3 were produced in an *E. coli* cell-free expression system using an extract derived from T7 SHuffle cells (Dopp and Reuel 2020) and additives optimized for disulfide bond expression (**Figure 1A, Table S1**). We produced 8.2 μM (274 $\mu\text{g}/\text{mL}$) and 3.9 μM (94 $\mu\text{g}/\text{mL}$) full-length, soluble SUL-I and CSL3 respectively, as determined by radiolabeled amino acid (^{14}C -Leucine) incorporation and quantification (**Figure 1B, S2A-C**). Importantly, lowering the expression temperature to 16 °C was necessary to increase the solubility and yields of both lectins (**Figure S2A-B**).

We first confirmed that cell-free expressed SUL-I and CSL3 were active by isolating them using either D-lactose or D-galactose affinity resin respectively (**Figure S2D**). Next, we

determined if BLI analysis was directly compatible with unpurified lectins in cell-free reactions. We loaded biotinylated L-rhamnose monosaccharide (L-Rha α -sp3-biot) onto a streptavidin BLI tip. We then accounted for background interaction from the cell-free reaction mixture by adding a “blocking” step in a negative cell-free reaction (containing water instead of plasmid DNA encoding the lectin) following rhamnose substrate immobilization on the tip and prior to the interaction assay (**Figure 1C, Figure S3**). We observe a rapid and strong association phase for both lectins, with almost no dissociation phase, when cell-free reactions expressing either SUL-I or CSL3 are present in solution (**Figure 1D**). No interaction occurs in reactions that did not contain lectin DNA expression template (no DNA), confirming the specificity of the method (**Figure 1D**). The signal variation was different for the two lectins at increased dilution, indicating better binding from SUL-I. This may be related to the higher expression levels for SUL-I in the cell-free reaction and by the larger number of binding sites available in SUL-I compared to CSL3 (**Figure S4**). For SUL-I, the signal level remains unchanged with a 50-fold dilution, indicating that the dissociation constant (k_D) is lower than the lectin concentration at this dilution (160 nM). Whereas for CSL3, we observe a more rapid decrease of the signal with increasing dilution (80 nM), yielding an approximation of k_D that is higher than the concentration of CSL3 at this dilution (between 80 and 400 nM).




Figure 1. Cell-free expressed rhamnose-binding lectins are directly compatible with BLI analysis. A) Schematic representation of cell-free expression system for disulfide bond containing lectins. B) Average total and soluble yields of SUL-I and CSL3 expressed in cell-free at 16°C by radiolabeled amino acid (^{14}C -Leucine) incorporation. Error bars represent the standard deviation for $n=3$ CFE reactions. C) Schematic of BLI assay where a biotinylated rhamnose monosaccharide is immobilized on the tip, followed by blocking with a cell-free reaction with no DNA template, and then measuring the interaction with a lectin synthesized in the cell-free reaction. D) BLI sensorgram of the interaction between a rhamnose monosaccharide immobilized on the tip and a 10-fold dilution in PBS of either negative control (no DNA template), SUL-I (~800 nM), or CSL3 (~400 nM) enriched cell-free reaction.

Cell-free expression coupled with BLI enables determination of SUL-I binding specificity and affinity

To further validate this experimental set up, we tested the specificity of SUL-I for different rhamnose substrate architectures immobilized on the BLI tip (**Figure 2A, Figure S4**). In addition to the monosaccharide, we also assessed two synthesized multivalent compounds, a tetravalent (Rha4) and a hexadecavalent (Rha16) cyclopeptide functionalized with L-rhamnose by triazole linkages, using architectures that have previously been used to assay multivalent lectin/substrate

interactions (Picault et al. 2022). We also examined a commercial multivalent polyacrylamide polymer functionalized with rhamnose (PAA-rhamnose). We observed higher signal as the valency of the substrate on the tip increased, with the smallest change in wavelength for the monosaccharide and the largest change in wavelength (~ 8-fold higher) for the Rha16 substrate, indicating that more SUL-I was able to bind to the additional L-rhamnose immobilized on the tip (**Figure 2A**). Very little dissociation was observed in all cases.

Additionally, we employed this strategy to determine the specificity of SUL-I by screening the ability of different mono- and oligo-saccharides (rhamnose, Gb3, isoGb3, lactose, galactose, glucose, fucose, and mannose) to inhibit the binding interaction with the rhamnose monosaccharide on the tip. We determined that a 50-fold dilution of SUL-I CFPS (~ 160 nM) still resulted in robust signal (**Figure S4**) and incubated the diluted cell-free reaction with sugar competitors at varying concentrations for one hour before the interaction step. For all experiments where the sugar acted as an inhibitor, the signal decreased with increased concentration of inhibitor (**Figure S5**). Subsequently, we compared the equilibrium response R_{eq} (maximum equilibrium change in wavelength), as determined by fitting a 1:1 association curve for each condition, to estimate relative IC₅₀ values for each sugar (**Figure 2B, Figure S5**) (Orthwein et al. 2021). Rhamnose had the lowest IC₅₀ for this system (0.07 mM), followed by Gb3 (0.15 mM), while galactose had the highest observed IC₅₀ (1.26 mM) (**Figure 2B, Figure S5A, Table S2**). Fucose, glucose, and mannose did not fully inhibit binding at the concentrations tested (**Figure 2B**).

Finally, we demonstrated the utility of this method for estimating binding kinetics and affinity (**Figure 2C**). As previously described for multivalent lectin-ligand systems with weak dissociation, we estimated the kinetic parameters k_{on} , k_{off} , and k_D from the linear relationship between k_{obs} , fit by a 1:1 association model, and the lectin concentration, determined by ¹⁴C-Leucine incorporation (**Figure 2C-D, Figure S6**) (Picault et al. 2022). Values of k_{on} were not affected much by the architecture of the substrates, with variations from $3.5 \times 10^{-5} \text{ M}^{-1} \text{ s}^{-1}$ to $1.9 \times$

$10^{-5} \text{ M}^{-1} \text{ s}^{-1}$ (**Table S3**). We obtained the highest k_D of $\sim 80 \text{ nM}$ for the monosaccharide and slightly higher affinity k_D ($\sim 30\text{-}40 \text{ nM}$) for the Rha4, Rha16 and PAA-Rha on the tip (**Figure S6, Table S3**). When the rhamnose ligands are immobilized, all sensors present rhamnose multivalently regardless of individual ligand structure, resulting in little difference in binding behavior between the architectures.

We also reversed the experiment and immobilized the lectins with N-terminal 6xHis tag from a crude cell-free reaction using Ni-NTA tips, to measure interaction with rhamnose ligands in solution (**Figure 2E, Figure S7A**). This system did not follow the same kinetics as when the rhamnose ligand was on the tip, possibly due to different affinity when the lectin is immobilized, preventing a 1:1 binding analysis. However, using steady state analysis we were able to estimate a k_D of $\sim 100 \text{ nM}$ for Rha4 and $\sim 50 \text{ nM}$ for Rha16 with SUL-I on the tip (**Figure S8, Table S4**). Importantly, these k_D values are the same order of magnitude as our estimates for SUL-I binding affinity when the rhamnose substrate was immobilized, although the Rha16 has higher affinity than Rha4 when SUL-I is immobilized.

While we could detect interaction with the Rha4 and Rha16 ligands in solution, this experimental set up is limited by the detectable ligand size, and we could not detect binding with the rhamnose monosaccharide in solution (**Figure S7B**) as previously observed for other monosaccharides (Laignre et al. 2018). However, this method is also of interest for detection of glycans in crude samples. Towards this goal, we show that SUL-I enables detection of the *Shigella flexneri* O-antigen glycan, which contains terminal rhamnose residues (Anderson et al. 2016), in crude cell-free glycoprotein synthesis (CFGpS) extracts (Jaroentomeechai et al. 2018; Kightlinger et al. 2019; Warfel et al. 2022) without purification (**Figure 2F**). In these experiments, we accounted for background binding to the Ni-NTA tip with a reference sensor that was loaded with cell-free reaction that did not contain the DNA expression template for the lectin (**Figure S7, S8**), shown without subtraction in **Figure 2F**. Nevertheless, as previously described for protein loading onto BLI tips from crude cell lysate, a stronger, more specific interaction such as biotin-

streptavidin may result in more uniform and saturated binding of the lectin to the tip (Pogoutse et al. 2016).




Figure 2. Cell-free expression coupled with biolayer interferometry enables determination of lectin binding specificity and affinity. A) Sensorgram of the interaction between different rhamnose substrate architectures (Rha16, PAA-Rha, Rha4, Rha monosaccharide) immobilized on the SA BLI tip with 50-fold diluted SUL-I (~160 nM) cell-free reaction in solution. B) Impact of different sugar inhibitors on the interaction between the rhamnose monosaccharide tip and SUL-

I lectin in solution. The normalized equilibrium binding signal (R_{eq}) as determined by a 1:1 association fit, is plotted as a function of the log (inhibitor concentration) for each sugar inhibitor. Nonlinear curve fit using Graphpad PRISM log(inhibitor) vs normalized response is shown in the corresponding color for each inhibitor. C) Interaction of Rha4 substrate immobilized on the SA tip, with varying dilutions of cell-free reaction containing SUL-I (~16 nM - 320 nM) in solution. The 1:1 model fit used to determine k_{obs} is shown in a black dotted line for each concentration and estimated concentration of SUL-I in solution is indicated to the right of the corresponding curve. D) k_{obs} values plotted as a function of SUL-I concentration and fit with a simple linear regression in Graphpad PRISM for Rha4. E) Interaction of SUL-I substrate immobilized on the Ni-NTA tip from a cell-free reaction, with varying dilutions of Rha4 substrate (~25nM - 200 nM) in solution. F) Detection of 100-fold dilution of crude extract enriched with *Shigella flexneri* 2a O-antigen in solution with SUL-I in cell-free reaction (blue) or control cell-free reaction with no DNA (red) immobilized on the Ni-NTA tip. The monomer structure of the O-antigen glycan is indicated above the sensorgram. All data is representative of at least two independent experiments.

Discussion

This work couples a bacterial cell-free expression platform directly with biolayer interferometry (BLI) to create a workflow compatible with small-scale expression and screening of lectins in high-throughput. As a proof of concept, we expressed two multivalent rhamnose-binding lectins from marine organisms and demonstrated that lectins in crude cell-free reactions are compatible with BLI interaction assays when tips are functionalized with glycan ligands. Importantly, we demonstrated this method with two commercially available biotinylated rhamnose ligands as well as two synthesized rhamnose substrates of varying sizes, enabling wide use of this technique. This approach does not require immobilization of the lectin and enables in-solution determination of binding specificity as well as affinity. Further, the ability to quantify the cell-free expressed protein in the crude mixture enables kinetics measurements using this method.

We also demonstrated that SUL-I expressed in a crude cell-free reaction can be immobilized on the BLI tip and used for qualitative detection of glycan epitopes in solution. While immobilization of the lectin has some drawbacks, such as size of glycoconjugate that can be detected or the impact of immobilization on binding activity, it can enable detection of glycan substrates from crude mixtures. Here, we demonstrate detection of the *S. flexneri* 2a O-antigen, of interest for vaccine development, directly in cell extract, aligning with previous reports that rhamnose binding lectins can bind smooth LPS from other *Shigella* serotypes (Shiina et al. 2002).

In total, we have shown that unpurified lectins in cell-free protein synthesis reactions can be used directly in BLI experiments, either in solution or immobilized on the sensor tip, for qualitative detection of carbohydrate-lectin interactions. This work advances efforts to characterize binding interactions without a purification step, which could expedite discovery and screening of new binders (Khavrutskii et al. 2013; Pogoutse et al. 2016). We expect that this workflow can be coupled to tools like LectomeExplore (Bonnardel et al. 2021) to express and screen uncharacterized lectins, and will advance the field of synthetic glycobiology for lectin discovery and application.

Materials and Methods

Cell-free Protein Synthesis (CFPS)

Cell-free protein synthesis and ^{14}C -Leucine quantification were performed as described previously using a modified PANOx-SP (PEP) formulation in an extract prepared from SHuffle T7 Express (NEB) cells (Warfel et al. 2022). Extract was pre-incubated with 50 μM IAM at room temperature for 30 minutes. Reactions were supplemented with 10 μM DsbC, 1 mM GSH, and 4 mM GSSG and incubated at 16 °C for 20 hours for lectin expression. Detailed methods can be found in the Supplementary Information.

Bio-layer interferometry (BLI)

All BLI experiments were performed using an Octet RED96 Instrument with data collected with ForteBio DataAcquisition9, analyzed and fit with ForteBio DataAnalysis9, and plotted with Graphpad PRISM. Detailed methods can be found in the Supplementary Information.

Acknowledgements

The authors would like to thank Derek Wong, Dr. Simona Notova, Maddie DeWinter, and Dr. Andrew Hunt for helpful scientific discussions, and Dr. Ashty Karim for help with manuscript preparation. The authors also thank Hugues Bonnet for assistance with the BLI equipment. K.F.W. thanks the Chateaubriand Fellowship program: this material is based upon research supported by the Chateaubriand Fellowship of the Office for Science & Technology of the Embassy of France in the United States. A.I. acknowledges support from Glyco@Alps (ANR-15-IDEX-0002) and Labex Arcane/CBH-EUR-GS (ANR-17-EURE-0003). M.C.J. acknowledges DTRA (HDTRA1-21-1-0038) and the National Science Foundation (MCB 1936789). The authors also acknowledge support from ICMG UAR 2607 for BLI facilities.

References

- Anderson M, Sansonetti PJ, Marteyn BS. 2016. Shigella diversity and changing landscape: Insights for the twenty-first century. *Frontiers in Cellular and Infection Microbiology*. 6:45. doi:10.3389/fcimb.2016.00045.
- Arnaud J, Audfray A, Imberty A. 2013. Binding sugars: From natural lectins to synthetic receptors and engineered neolectins. *Chem Soc Rev*. 42:4798–4813. doi:10.1039/c2cs35435g.
- Bonnardel F, Mariethoz J, Pérez S, Imberty A, Lisacek F. 2021. LectomeXplore, an update of UniLectin for the discovery of carbohydrate-binding proteins based on a new lectin classification. *Nucleic Acids Res*. 49(D1):D1548–D1554. doi:10.1093/nar/gkaa1019.
- Carlson ED, Gan R, Hodgman CE, Jewett MC. 2012. Cell-free protein synthesis: applications come of age. *Biotechnol Adv*. 30(5):1185–1194. doi:10.1016/j.biotechadv.2011.09.016.
- Concepcion J, Witte K, Wartchow C, Choo S, Yao D, Persson H, Wei J, Li P, Heidecker B, Ma W, et al. 2009. Label-free detection of biomolecular interactions using BioLayer interferometry for kinetic characterization. *Comb Chem High Throughput Screen*. 12(8):791–800. doi:10.2174/138620709789104915.
- Dopp JL, Reuel NF. 2020. Simple, functional, inexpensive cell extract for in vitro prototyping of proteins with disulfide bonds. *Biochem Eng J*. 164:107790. doi:10.1016/j.bej.2020.107790.
- Fernandez-Poza S, Padros A, Thompson R, Butler L, Islam M, Mosely JA, Scrivens JH, F Rehman M, Akram MS. 2021. Tailor-made recombinant prokaryotic lectins for characterisation of glycoproteins. *Anal Chim Acta*. 1155:338352. doi:10.1016/j.aca.2021.338352.
- Goerke AR, Swartz JR. 2008. Development of cell-free protein synthesis platforms for disulfide bonded proteins. *Biotechnol Bioeng*. 99(2):351–367. doi:10.1002/bit.21567.

Hatakeyama T, Ichise A, Unno H, Goda S, Oda T, Tateno H, Hirabayashi J, Sakai H, Nakagawa H. 2017. Carbohydrate recognition by the rhamnose-binding lectin SUL-I with a novel three-domain structure isolated from the venom of globiferous pedicellariae of the flower sea urchin *Toxopneustes pileolus*. *Protein Sci.* 26(8):1574. doi:10.1002/PRO.3185.

Hatakeyama T, Ichise A, Yonekura T, Unno H, Goda S, Nakagawa H. 2015. CDNA cloning and characterization of a rhamnose-binding lectin SUL-I from the toxopneustid sea urchin *Toxopneustes pileolus* venom. *Toxicon.* 94:8–15. doi:10.1016/J.TOXICON.2014.11.236.

Hershewe JM, Warfel KF, Iyer SM, Peruzzi JA, Sullivan CJ, Roth EW, DeLisa MP, Kamat NP, Jewett MC. 2021. Improving cell-free glycoprotein synthesis by characterizing and enriching native membrane vesicles. *Nat Commun.* 12(1):2363. doi:10.1038/s41467-021-22329-3.

Hunt AC, Case JB, Park Y-J, Cao L, Wu K, Walls AC, Liu Z, Bowen JE, Yeh H-W, Saini S, et al. 2022. Multivalent designed proteins neutralize SARS-CoV-2 variants of concern and confer protection against infection in mice. *Sci Transl Med.* 14(646):eabn1252. doi:10.1126/scitranslmed.abn1252.

Hunt AC, Vögeli B, Kightlinger WK, Yoesep DJ. 2021. A high-throughput, automated, cell-free expression and screening platform for antibody discovery. *bioRxiv.* <https://www.biorxiv.org/content/10.1101/2021.11.04.467378.abstract>.

Jaroentomeechai T, Kwon YH, Liu Y, Young O, Bhawal R, Wilson JD, Li M, Chapla DG, Moremen KW, Jewett MC, et al. 2022. A universal glycoenzyme biosynthesis pipeline that enables efficient cell-free remodeling of glycans. *Nat Commun.* 13(1):6325. doi:10.1038/s41467-022-34029-7.

Jaroentomeechai T, Stark JC, Natarajan A, Glasscock CJ, Yates LE, Hsu KJ, Mrksich M, Jewett MC, DeLisa MP. 2018. Single-pot glycoprotein biosynthesis using a cell-free transcription-translation system enriched with glycosylation machinery. *Nat Commun.* 9(1):2686. doi:10.1038/s41467-018-05110-x.

Khavrutskii L, Yeh J, Timofeeva O, Tarasov SG, Pritt S, Stefanisko K, Tarasova N. 2013. Protein purification-free method of binding affinity determination by microscale thermophoresis. *J Vis Exp.*(78):50541. doi:10.3791/50541.

Kightlinger W, Duncker KE, Ramesh A, Thames AH, Natarajan A, Stark JC, Yang A, Lin L, Mrksich M, DeLisa MP, et al. 2019. A cell-free biosynthesis platform for modular construction of protein glycosylation pathways. *Nat Commun.* 10(1):5404. doi:10.1038/s41467-019-12024-9.

Kruyer NS, Sugianto W, Tickman BI, Alba Burbano D, Noireaux V, Carothers JM, Peralta-Yahya P. 2021. Membrane Augmented Cell-Free Systems: A New Frontier in Biotechnology. *ACS Synth Biol.* 10(4):670–681. doi:10.1021/acssynbio.0c00625.

Laigre E, Goyard D, Tiertant C, Dejeu J, Renaudet O. 2018. The study of multivalent carbohydrate-protein interactions by bio-layer interferometry. *Org Biomol Chem.* 16(46):8899–8903. doi:10.1039/c8ob01664j.

Martínez-Alarcón D, Blanco-Labra A, García-Gasca T. 2018. Expression of lectins in heterologous systems. *Int J Mol Sci.* 19(2):616. doi:10.3390/ijms19020616.

Matthies D, Haberstock S, Joos F, Dötsch V, Vonck J, Bernhard F, Meier T. 2011. Cell-free expression and assembly of ATP synthase. *J Mol Biol.* 413(3):593–603. doi:10.1016/j.jmb.2011.08.055.

Orthwein T, Huergo LF, Forchhammer K, Selim KA. 2021. Kinetic Analysis of a Protein-protein Complex to Determine its Dissociation Constant (KD) and the Effective Concentration (EC50) of an Interplaying Effector Molecule Using Bio-layer Interferometry. *Bio Protoc.* 11(17):e4152. doi:10.21769/BioProtoc.4152.

Picault L, Laigre E, Gillon E, Tiertant C, Renaudet O, Imberty A, Goyard D, Dejeu J. 2022. Characterization of the interaction of multivalent glycosylated ligands with bacterial lectins by biolayer interferometry. *Glycobiology.* 32(10):886–896. doi:10.1093/glycob/cwac047.

Pogoutse AK, Lai CC-L, Ostan N, Yu R-H, Schryvers AB, Moraes TF. 2016. A method for measuring binding constants using unpurified in vivo biotinylated ligands. *Anal Biochem.* 501:35–43. doi:10.1016/j.ab.2016.02.001.

Shiina N, Tateno H, Ogawa T, Muramoto K, Saneyoshi M, Kamiya H. 2002. Isolation and characterization of L-rhamnose-binding lectins from chum salmon (*Oncorhynchus keta*) eggs. *Fish Sci.* 68(6):1352–1366. doi:10.1046/J.1444-2906.2002.00575.X.

Shirai T, Watanabe Y, Lee MS, Ogawa T, Muramoto K. 2009. Structure of Rhamnose-binding Lectin CSL3: Unique Pseudo-tetrameric Architecture of a Pattern Recognition Protein. *J Mol Biol.* 391(2):390–403. doi:10.1016/J.JMB.2009.06.027.

Silverman AD, Karim AS, Jewett MC. 2020. Cell-free gene expression: an expanded repertoire of applications. *Nat Rev Genet.* 21(3):151–170. doi:10.1038/s41576-019-0186-3.

Siukstaite L, Imberty A, Römer W. 2021. Structural Diversities of Lectins Binding to the Glycosphingolipid Gb3. *Frontiers Mol Biosci.* 8:696. doi:10.3389/FMOLB.2021.704685.

Stark JC, Jaroentomeechai T, Moeller TD, Hershewe JM, Warfel KF, Moricz BS, Martini AM, Dubner RS, Hsu KJ, Stevenson TC, et al. 2021. On-demand biomanufacturing of protective conjugate vaccines. *Sci Adv.* 7(6): eabe9444. doi:10.1126/sciadv.abe9444.

Warfel KF, Williams A, Wong DA, Sobol SE, Desai P, Li J, Chang Y-F, DeLisa MP, Karim AS, Jewett MC. 2022. A Low-Cost, Thermostable, Cell-Free Protein Synthesis Platform for On-Demand Production of Conjugate Vaccines. *ACS Synth Biol.* in press. doi:10.1021/acssynbio.2c00392.

Zawada JF, Yin G, Steiner AR, Yang J, Naresh A, Roy SM, Gold DS, Heinsohn HG, Murray CJ. 2011. Microscale to manufacturing scale-up of cell-free cytokine production—a new approach for shortening protein production development timelines. *Biotechnol Bioeng.* 108(7):1570–1578. doi:10.1002/bit.23103.