

REVIEW

Cellular mechanisms of potassium transport in plants

Dev T. Britto and Herbert J. Kronzucker*

Department of Biological Sciences, University of Toronto, 1265 Military Trail, Toronto, Ontario, Canada M1C 1A4

Correspondence*Corresponding author,
e-mail: herbertk@utsc.utoronto.caReceived 19 September 2007;
revised 7 January 2008

doi: 10.1111/j.1399-3054.2008.01067.x

Potassium (K^+) is the most abundant ion in the plant cell and is required for a wide array of functions, ranging from the maintenance of electrical potential gradients across cell membranes, to the generation of turgor, to the activation of numerous enzymes. The majority of these functions depend more or less directly upon the activities and regulation of membrane-bound K^+ transport proteins, operating over a wide range of K^+ concentrations. Here, we review the physiological aspects of potassium transport systems in the plasma membrane, re-examining fundamental problems in the field such as the distinctions between high- and low-affinity transport systems, the interactions between K^+ and other ions such as NH_4^+ and Na^+ , the regulation of cellular K^+ pools, the generation of electrical potentials and the problems involved in measurement of unidirectional K^+ fluxes. We place these discussions in the context of recent discoveries in the molecular biology of K^+ acquisition and produce an overview of gene families encoding K^+ transporters.

Introduction

2007 marks the 200th anniversary of the discovery of potassium by the English chemist Sir Humphrey Davy. Since then, much intensive and insightful research has been conducted on this element and its function in the living world. A recent biological highlight of this work was the awarding of the 2003 Nobel Prize in chemistry to R. MacKinnon for elucidation of the structure and function of the *Streptomyces lividans* K^+ channel at the atomic scale (Doyle et al. 1998). Potassium is one of the most abundant elements in plant tissues, comprising about 1–10% of dry matter (Epstein and Bloom 2005). Globally, over 25 MT of potash fertilizer is applied to cropland every year (Minerals Mining and Sustainable Development 2002: http://www.iied.org/mmsd/mmsd_pdfs/065_ifa.pdf), and improved efficiency of potassium acquisition by plants is an important concern among plant scientists (Lynch 2007).

Although sodium can, under some circumstances, partially replace potassium for relatively non-specific functions such as osmoregulation (Marschner et al. 1981,

Subbarao et al. 1999), potassium is classified as an essential macronutrient for all plants. Many of the diverse roles of K^+ in plant cells depend on the transport of K^+ through K^+ -specific membrane-bound transport proteins. Such functions include the short-term maintenance of electrical potentials across membranes (Cheeseman and Hanson 1979) and turgor-related phenomena such as cell expansion (Dolan and Davies 2004), plant movements (Moran 2007, Philippar et al. 1999), pollen tube development (Mouline et al. 2002) and stomatal opening and closing (Dietrich et al. 2001, Humble and Hsiao 1969). Other well-characterized biological functions of potassium include the activation of numerous enzymes (Suelter 1970), long-distance transport of nitrate (Ben-Zioni et al. 1971) and sucrose (Cakmak et al. 1994) and the charge stabilization of anions within the cell (Clarkson and Hanson 1980). In this review, we focus on the key topics of membrane transport and compartmentation of potassium in plant cells, summarizing the state of knowledge and indicating new directions in potassium research.

Abbreviations – CNG, cyclic-nucleotide-gated channel; HATS, high-affinity transport system; LATS, low-affinity transport system.

HATS and LATS: the two-mechanism model

Research on potassium transport in plant roots includes some of the first instances of the utilization of novel radiotracer technologies in the 1940s. Of particular note were the pioneering studies of Epstein and co-workers, who established that, at low external concentrations (under 1 mM), the unidirectional uptake kinetics of K^+ can be mathematically analyzed using a Michaelis–Menten model (Epstein et al. 1963). This led to the assignment of maximal velocity (V_{max}) and half-saturation concentration (K_M) terms to what was initially referred to as ‘Mechanism 1’ and later as the high-affinity transport system (HATS) for potassium. At external concentrations higher than 1 mM, K^+ transport patterns become dominated by a kinetically distinct system, which shows little if any saturation. This linear component of K^+ transport was first termed ‘Mechanism 2’ by Epstein and later the low-affinity transport system (LATS). The two-mechanism model is depicted in Fig. 1.

The distinct kinetics, energetics and regulatory aspects of HATS and LATS were subsequently worked out in

greater detail (Glass and Dunlop 1978, Kochian and Lucas 1982, Kochian et al. 1985, Maathuis and Sanders 1995). Apart from the different external K^+ concentrations at which they operate and the shapes of their associated concentration dependence curves, each system has several unique characteristics. Most importantly, there appear to be distinct mechanisms by which HATS and LATS transporters catalyze the flux of K^+ across the plasma membrane. In the case of HATS, it is likely that K^+ enters the cell via symport with H^+ , with a proposed (although possibly variable) 1:1 stoichiometry (Kochian et al. 1989, Maathuis and Sanders 1994, Maathuis et al. 1997), in an energy-dependent process involving the transmembrane proton motive force (Fig. 2). The H^+ and electrical gradients underlying this force are maintained by membrane-bound, ATP-hydrolyzing transporters that pump H^+ out of the cytosol and into the external medium (Cheeseman and Hanson 1980, Cheeseman et al. 1980, Kochian et al. 1989, Maathuis and Sanders 1994, Palmgren 2001, Pardo et al. 2006). The regulation of HATS is accomplished both by limitations to substrate binding with the carrier molecule and by feedback on




Fig. 1. General schematic of K^+ influx kinetics in plant roots. Isotherms for HATS (open squares) and LATS (open triangles) are indicated, in addition to the combined flux (solid line). Arrows and dashed lines indicate up- and downregulation of HATS, in response to plant K^+ status. Inset: Effect of accompanying anions on low-affinity K^+ fluxes.




Fig. 2. General mechanisms proposed for K⁺ influx into plant cells, via the HATS (upper diagram) and the LATS (lower diagram). In the HATS mechanism, the thermodynamically uphill flux of K⁺ is driven by the downhill flux of H⁺; charge balance is achieved by the outward pumping of two H⁺ by the plasma membrane proton ATPase. In the LATS mechanism, by contrast, an electrogenic uniport of K⁺ is electrically balanced by the ATP-driven efflux of one H⁺.

gene transcription (Gierth et al. 2005). In the LATS, K⁺ is thought to be transported via ion channels (K⁺ specific as well as non-selective), which, by forming pores that cross the membrane bilayer, can catalyze thermodynamically downhill fluxes that are at least three orders of magnitude higher than those catalyzed by pumps and carriers (Tester 1990). Thermodynamically, channel activity relies on electrochemical potential gradients for K⁺ to drive transport and is regulated by a wide range of agents: membrane voltage (Gassmann and Schroeder 1994), pH (Hoth and Hedrich 1999), cyclic nucleotides (Talke et al. 2003), CO₂-dependent light activation (Deeken et al. 2000), reactive oxygen species (Cakmak 2005) and K⁺ itself (Johansson et al. 2006, Liu et al. 2006). The thermodynamic distinction between these two basic types of transporters forms the basis for their designation, in the recently devised transporter classification (TC) system, as class 1 (channel/pore type) for LATS transport, and class 2 (electrochemical potential-driven transporter type) for HATS transport (Busch and Saier 2002).

K⁺ acquisition from low external concentrations is usually considered to be an energy-demanding process, while that from high concentrations is energetically passive. On the surface, this view is supported by analyses of the electrochemical potential gradient for K⁺ transport into the plant cell, which is defined primarily by the differences in K⁺ concentration and electrical potential on either side of the plasma membrane (Cheeseman and Hanson 1980, Szczerba et al. 2006a). K⁺ is usually the most abundant cation in the cytosol, with concentrations typically ranging from 40 to 200 mM (Kronzucker et al. 2003, Leigh and Wyn Jones 1984, Walker et al. 1996), a condition that could present an energetic obstacle to K⁺ entry even under abundant potassium supply, were it not for the electrical charge separation across the membrane (inside negative), which greatly enhances the uptake of cations. However, in very dilute solutions, this electrical pull is insufficient to drive K⁺ influx, and an active transport mechanism is thus postulated under such conditions (Cheeseman and Hanson 1980, Maathuis and Sanders 1994).

Nevertheless, if, at higher external concentrations, thermodynamic conditions favor an energetically downhill influx of K⁺, a more comprehensive view indicates that this process must still entail cellular energy consumption. This is because for every K⁺ ion that enters the cell and is retained, a slight depolarization of the membrane will occur, in turn requiring a subsequent, energy-dependent electrical compensation via the plasma membrane H⁺ ATPase. In addition, it has been amply demonstrated that, as potassium provision goes up, increased influx of K⁺ is accompanied by a substantial increase in K⁺ efflux. Except for the special case in which the thermodynamic gradient for K⁺ transport across the membrane is zero (a Nernstian condition – see, e.g. Maathuis and Sanders 1994), a passive influx of K⁺ entails that the efflux of K⁺ must be directly energy requiring (Szczerba et al. 2006a). The idea that energy demand for K⁺ acquisition can occur over a broad range of K⁺ supply (1–50 mM) is supported by the strong correlations between K⁺ uptake and ATP hydrolysis by roots of numerous plant species (Fisher et al. 1970).

Responses to plant K⁺ status also distinguish HATS from LATS (Fig. 1). HATS is strongly downregulated under K⁺-replete conditions and, conversely, strongly upregulated under K⁺ starvation (Glass 1978, Kochian and Lucas 1982). By contrast, influx in the LATS range appears, at least in some studies, to be insensitive to K⁺ status (Kochian and Lucas 1982), although, in at least one instance, low-affinity K⁺ influx was enhanced by low external K⁺ (Maathuis and Sanders 1995). In the latter case, however, the hyperpolarized state of the plasma membrane under low K⁺, rather than a specific regulatory

mechanism, may have enhanced the passive, inward, channel-mediated current. In addition, the apparent insensitivity of LATS to plant K^+ status in other work (Kochian and Lucas 1982) may be somewhat misleading because flux underestimates are more prone to occur in the LATS range (Szczerba et al. 2006b); thus, it is possible that LATS-range K^+ influx is generally upregulated with increased K^+ status, a situation similar to the increase in low-affinity NH_4^+ transport that has been repeatedly seen with higher plant N status (Cerezo et al. 2001, Rawat et al. 1999; see below). Armengaud et al. (2004) and Gierth et al. (2005), in transcriptome analyses of *Arabidopsis*, showed that the response of putative K^+ transporter expression to K^+ starvation was largely limited to the strong upregulation of a single high-affinity transport gene, *athak5*, which may be responsible for the majority of high-affinity uptake. In another study examining gene expression in roots of wheat and barley (Wang et al. 1998), the gene encoding the putative K^+ transporter HKT1 was rapidly and strongly upregulated upon K^+ starvation, although the relevance of this gene product to K^+ acquisition has been called into question (Haro et al. 2005, Hayes et al. 2001).

Another notable difference between the two systems is seen in the interaction between K^+ uptake and the presence of other ions. HATS transport can be very sensitive to competitors or toxicants such as NH_4^+ and Na^+ , while LATS transport appears to be relatively resistant to their influence (Kronzucker et al. 2003, Maathuis and Sanders 1994, Maathuis et al. 1996, Nieves-Cordones et al. 2007, Pardo and Quintero 2002, Santa-Maria et al. 2000, Spalding et al. 1999). Indeed, the sensitivity of HATS to NH_4^+ is the basis of the surprising finding that channel mediation can contribute to K^+ transport in the high-affinity range (Hirsch et al. 1998; see below). We have recently shown (Szczerba et al. 2008) that the reverse effect is not seen; that, rather, low-affinity NH_4^+ transport in barley roots is substantially reduced when the K^+ supply is increased, offering a partial explanation for the alleviation of NH_4^+ toxicity by elevated external $[K^+]$.

Although the suppression of high-affinity K^+ transport by NH_4^+ is well established, evidence for the interactions between Na^+ and K^+ is more equivocal: for instance, Maathuis et al. (1996) found that, in a wide range of terrestrial plants (including *Arabidopsis*), the addition of Na^+ either inhibited or had no effect on high-affinity K^+ absorption; by contrast, Spalding et al. (1999) showed that the non-channel-mediated uptake of K^+ into *Arabidopsis* roots (see discussion of AKT1 below), at an external $[K^+]$ of 10 μM , was in fact stimulated by Na^+ . The latter finding may indicate a role, if limited, of a proposed Na^+/K^+ exchange mechanism of the HKT1

transporter (Rubio et al. 1995). Our recent work (Kronzucker et al. 2006) shows that, in barley roots, Na^+ suppresses K^+ influx at both 0.1 and 1.5 mM external $[K^+]$, while a reciprocal effect of K^+ on Na^+ influx is not observed; in still more recent experiments (under review), we have found evidence that Na^+ inhibits K^+ influx well into the low-affinity range (up to 40 mM), while such high concentrations of K^+ had no effect on Na^+ transport or compartmentation. A similar lack of effect of K^+ on Na^+ uptake was observed by Wang et al. (2007) in *Sueda maritima*, but strong reciprocal interactions of the two ions were seen in a classic study on barley roots (Rains and Epstein 1967). That the latter study shows results, in the same species, contradictory to ours is likely an indication of the non-steady-state conditions used in that study. This, however, is not to discount such an approach; it is essential to consider both short- and long-term effects of these ionic interactions, particularly with regard to the potentially important role that enriched soil K^+ may play in the amelioration of Na^+ toxicity in glycophytes.

A further aspect of differential ion interaction in the two transport ranges is the effect on the LATS of the accompanying anion (Fig. 1, inset). Epstein et al. (1963), for instance, showed that K^+ (Rb^+) influx in the LATS range was much higher with $RbCl$ than with Rb_2SO_4 . Similarly, Kochian et al. (1985) found that the velocity of LATS-range K^+ transport was 60% higher when the counterion was Cl^- rather than SO_4^{2-} , NO_3^- or $H_2PO_4^-$. In neither study was this counterion effect observed in the HATS range.

One final distinction between HATS and LATS transport to be mentioned here is that, under low-affinity conditions, the plasma membrane is much more permeable to K^+ , in both the influx and the efflux directions (Pettersson and Kasimir-Klemedtsson 1990, Szczerba et al. 2006a). This increased 'leakiness' results in very high ratios of efflux to influx as $[K^+]_{ext}$ rises through the millimolar range, a phenomenon common to a wide array of nutrient ions (Britto and Kronzucker 2006). One aspect of this high permeability is that K^+ ions can be lost from roots as a result of physical disturbance, possibly because of the engagement of mechanically or stretch-activated channels (Britto et al. 2006, Shabala et al. 2000). Clearly, this ion loss from the plant system can pose a problem for the flux researcher, and its possibility must be taken into account.

In the decades prior to the identification of genes encoding K^+ transport proteins, there was a vigorous debate about whether fluxes in the high- and low-affinity ranges were each catalyzed by specialized transport proteins or, alternatively, by proteins that undergo state changes in response to varying nutritional conditions,

resulting in modified substrate affinities (Borstlap 1981, Nissen 1980). The eventual identification of genetically distinct molecular entities operating in specific parts of the transport range appeared at first to have resolved this issue, until the publication of several reports indicating that individual transporters may have both high- and low-affinity activity. Such dual-affinity behavior has also been reported for the transport of nitrate (Liu et al. 1999) and phosphate (Shin et al. 2004). In the case of potassium, at least three transporters from *Arabidopsis thaliana* appear to have dual (or very broad spectrum) affinities: the carrier AtKUP1 (Fu and Luan 1998, Kim et al. 1998) and the inward-rectifying channels AKT1 (Hirsch et al. 1998) and KAT1 (Brüggemann et al. 1999). AtKUP1, when expressed in yeast cells, shows two half-saturation constants (K_M): one in the range of 22–44 μM external $[\text{K}^+]$ and another near 11 mM (Fu and Luan 1998, Kim et al. 1998). The mechanism by which the switch between affinities occurs may involve phosphorylation of the transport protein, as has been demonstrated for the switch between affinities of the nitrate transporter CHL1 (Liu and Tsay 2003), and for the change in gating characteristics of the voltage-gated channel AKT2 (Michard et al. 2005). The AKT1 channel was shown to have a dual- (or broad-spectrum) affinity character by allowing growth of *A. thaliana* under conditions where high-affinity K^+ transport was blocked by NH_4^+ . In this situation, an inward channel-mediated flux of K^+ was calculated to be thermodynamically feasible even with external $[\text{K}^+]$ as low as 10 μM , given a sufficiently hyperpolarized plasma membrane (Hirsch et al. 1998). A recent report by Li et al. (2006) indicates that AKT1 is upregulated at low external $[\text{K}^+]$ via a calcium-dependent phosphorylation event, providing further evidence for the role of this channel in 'high-affinity' K^+ influx. In addition, the guard cell channel KAT1 was shown to mediate K^+ uptake from an equally low $[\text{K}^+]$, when expressed both in *Arabidopsis* and in yeast cells (Brüggemann et al. 1999).

The physiological significance of these findings, however, remains unclear. In the case of KAT1, some evidence indicates that the channel inactivates at submillimolar external $[\text{K}^+]$, although in a heterologous expression system (mammalian HEK cells; Hertel et al. 2005), although this effect was not observed in *Arabidopsis* itself (Brüggemann et al. 1999). In the case of AKT1, the K_M of Rb^+ uptake by *Arabidopsis* roots (measured as the difference in uptake between wild-type plants and *akt1-1* mutants) was found to be 0.9 mM (Gierth et al. 2005), suggesting that AKT1 does in fact normally operate in the low-affinity range. In addition, K^+ uptake by channels relies on the membrane electrical potential as a driving force; the measured potential in the study by Hirsch et al. (1998) was only sufficiently negative in 20% of wild-type

cells to drive passive uptake through AKT1. Nevertheless, it has been reported elsewhere that AKT1 may be responsible for 55–63% of K^+ influx in the submillimolar range, again based on differences between wild-type and mutant plants (Spalding et al. 1999). In both examples, however, it is difficult to rule out pleiotropic effects in mutant lines, and thus the conclusions must be appreciated with some caution. Still, it seems clear that considerable kinetic overlap exists among genetically distinct groups of transport proteins.

Moreover, it is beyond question that plant roots have phenotypically plastic systems in place that allow potassium to be extracted from soil solutions that vary considerably in $[\text{K}^+]$, reflecting edaphically realistic concentration ranges of 0.1–6 mM or even higher in fertilized soils (Adams 1971, Kochian and Lucas 1982, Reisenauer 1966), which can change rapidly in the field, given the substantial impact of soil drying and rehydration.

In retrospect, given the multiplicity of distinct K^+ transport proteins found in plant cells and the number of different cell types found within the root, it is perhaps surprising that the HATS-range concentration-dependence isotherms of K^+ transport into roots fit so well with the Michaelis–Menten model of enzyme kinetics, a model that was initially developed for purified enzymes in solution. Interestingly, protoplasts isolated from maize roots exhibited kinetic patterns of K^+ uptake that were essentially identical to those measured in whole roots (Kochian and Lucas 1983), indicating that root K^+ transport in this range may be dominated by a single protein (Gierth et al. 2005) or by a group of closely related proteins that share similar kinetic characteristics.

Another problematic aspect of the interpretation of influx isotherms is that they are based on measurements that are underestimated because of the simultaneous occurrence of K^+ efflux (Britto and Kronzucker 2001). In the steady state (i.e. when plant growth and experimentation are conducted under the same physicochemical conditions), the degree to which efflux can result in an underestimate of influx depends greatly on external $[\text{K}^+]$ concentration. Because of the higher ratios of efflux to influx seen in the LATS range (see above), 'direct', steady-state influx measurements in this range are inherently more prone to inaccuracies than in the HATS range and can lead to substantial underestimates of K^+ influx (Szczerba et al. 2006b). This problem of accurate flux measurements in the LATS is not unique to K^+ because increased efflux at higher external concentrations is seen with several other nutrient ions (Britto and Kronzucker 2006).

This limitation of the direct influx method can be circumvented under some conditions by means of

compartmental analysis, which monitors both efflux and retention of potassium by the plant, in addition to providing information about intracellular pool sizes of K^+ (see below). However, a crucial assumption underlying such analyses is that the plant system is examined under steady-state conditions. Influx isotherms, on the other hand, depict influxes over a range of external concentrations, only one of which represents the steady-state provision. Thus, compartmental analysis cannot by itself resolve the distortions present in such isotherms as a result of simultaneous efflux. While the extent of efflux under non-steady-state conditions has yet to be investigated, preliminary results in our laboratory, from a combination of compartmental and direct influx analyses, indicate that steady-state K^+ efflux is conserved by the plant in the short term, when $[K^+]_{\text{ext}}$ is shifted, and that a new steady state requires several hours to be achieved. This information will be used to develop influx isotherms that more accurately reflect plant capacity to transport K^+ and other ions.

Another aspect of K^+ transport research that needs to be approached with caution is experimental work that employs rubidium-86 as a radiotracer for potassium. Justification of the widespread use of $^{86}\text{Rb}^+$ as a potassium analogue stems from several reports suggesting that, at least in some cases, there is little discrimination between the two ions (Kochian and Lucas 1982, Läuchli and Epstein 1970, Polley and Hopkins 1979). However, numerous other studies have shown that, on the contrary, $^{86}\text{Rb}^+$ is a poor tracer for K^+ in higher plant systems (Behl and Jeschke 1982, Cline and Hungate 1960, de Agazio et al. 1983, Jacoby 1975, Jacoby and Nissen 1977, Kannan and Ramani 1973, Maas and Leggett 1968, Marschner and Schimanski 1968, Schachtman et al. 1992); substantial discrimination between the two elements has also been observed in *Nitellopsis* (MacRobbie and Dainty 1958), *Chlamydomonas* (Malhotra and Glass 1995, Polley and Doctor 1985), *Chara* (Keifer and Spanswick 1978), *Ulva* and *Chaetomorpha* (West and Pitman 1967), as well as in *Escherichia coli* (Rhoads et al. 1977). Moreover, the discrimination between these elements is not uniform across experimental conditions. For instance, two studies have shown that it increases in *Chlamydomonas* with K^+ deprivation. In addition, de Agazio et al. (1983) concluded that, in maize roots, discrimination against Rb^+ is particularly strong in the high-affinity, energy-dependent range of K^+ uptake. A similar conclusion was reached with *E. coli* (Rhoads et al. 1977), in which a 1000-fold discrimination against Rb^+ was seen in the high-affinity K^+ uptake range and a 10- to 25-fold discrimination in two of three low-affinity systems. In the third low-affinity system, K^+ was actually discriminated against, in favor of Rb^+ .

Unfortunately, the majority of K^+ transport studies using radiotracers have used $^{86}\text{Rb}^+$ instead of $^{42}\text{K}^+$. This is largely because of the convenience of using $^{86}\text{Rb}^+$, which has a much longer half-life than $^{42}\text{K}^+$ (18.65 days vs 12.36 h). For the same reason, studies using $^{86}\text{Rb}^+$ rarely involve corroboration of results by use of $^{42}\text{K}^+$. Given the numerous instances of Rb/K discrimination, some of the conclusions concerning K^+ transport, arrived at using $^{86}\text{Rb}^+$ exclusively, are likely to be incorrect. This problem is compounded by the issue of simultaneous influx and efflux, mentioned above, that may give rise to inaccurate isotherms. A further complication in the measurement of K^+ fluxes is the rarely reported phenomenon of flux alterations caused by physical disturbance of experimental plants, a problem that is particularly pressing in the LATS range (Britto et al. 2006; see above).

Electrical aspects of K^+ transport

Potassium transport is a key component underlying the homeostatic regulation of the plasma membrane electrical potential difference ($\Delta\Psi$) in plant cells. Interestingly, however, a range of set points for $\Delta\Psi$ can be achieved by the cell, depending on factors such as the external concentration of K^+ itself (Cheeseman and Hanson 1979, Etherton and Higinbotham 1960, Hayes et al. 2001, Maathuis and Sanders 1993). Membranes depolarize rapidly in response to increases in external K^+ and hyperpolarize with decreasing $[K^+]_{\text{ext}}$. This can be explained by the passive, channel-mediated, electrogenic flux of K^+ either into the cell (causing depolarization) or out of it (causing hyperpolarization). The direction of the flux is determined by the direction of the potassium diffusion potential, which is a function of $\Delta\Psi$ and of the ratio of $[K^+]$ on either side of the membrane. $\Delta\Psi$ shows a roughly linear response to the log of $[K^+]_{\text{ext}}$ over several orders of magnitude (e.g. three orders in the study by Etherton and Higinbotham 1960). Thus, plant cells behave very similarly to potassium electrodes within this response range (Fig. 3).

Conversely, plants engage electrogenic K^+ transport that rectifies alterations in $\Delta\Psi$ brought about by other means. For instance, sudden exposure of roots to cations such as Rb^+ , Cs^+ and NH_4^+ causes membrane depolarization in root cells of maize, and the subsequent efflux of K^+ counteracts this change (Nocito et al. 2002). This efflux of K^+ is mediated by outward-rectifying K^+ channels; similarly, the influx of K^+ can counteract membrane hyperpolarization via the activity of inward-rectifying channels. Clearly, both types of channels are voltage regulated.

It should be pointed out, however, that the channel-mediated flux of K^+ along its diffusion potential is not




Fig. 3. Potassium-induced plasma membrane depolarization. Plant cells behave like potassium electrodes, adjusting their membrane electrical potentials to variable set points as external $[K^+]$ changes.

a primary generator of $\Delta\Psi$, as it involves merely passive transport. The active transport of K^+ , mediated by HATS, also does not contribute to the primary generation of $\Delta\Psi$ but is contingent upon an electrochemical proton gradient to furnish the required energy. Ultimately, the establishment of this gradient by the activity of the plasma membrane H^+ -pumping ATPase is the means by which plant cells become electrically polarized (Palmgren 2001), in contrast to animal cells, which generate $\Delta\Psi$ by the ATP-driven counter-exchange of three Na^+ ions and two K^+ ions.

Cytosolic K^+ pools

The cytosolic concentration of potassium is under fairly strict homeostatic control and is maintained by fluxes to and from the soil solution, as well as to and from the storage pool in the vacuole; these fluxes are catalyzed by transporters in the plasma membrane and tonoplast membrane, respectively. The vacuolar K^+ pool is much more dynamic than that of the cytosol, gaining and losing potassium as required to maintain cytosolic K^+ homeostasis (Leigh 2001, Walker et al. 1996). Typically, cytosolic $[K^+]$ is held between 80 and 200 mM, although declines in this value have been observed under conditions of K^+ starvation (Walker et al. 1996), salt stress (Hajibagheri et al. 1987, 1988, Kronzucker et al. 2006, Speer and Kaiser 1991), ammonium toxicity (Kronzucker et al. 2003) and aluminum toxicity (Lindberg and Strid 1997). Surprisingly, in our own laboratory, a substantial decline of 40–50% in cytosolic $[K^+]$ was also observed

with an increase in external $[K^+]$ from an HATS-range value of 0.1 mM to an LATS-range value of 1.5 mM (Kronzucker et al. 2003; Szczerba et al. 2006a; at higher $[K^+]_{ext}$ this pool was seen to recover). This counter-intuitive finding, determined under steady-state conditions, might be explained by a link between transmembrane K^+ distribution and the shift in K^+ transport from the high-affinity to the low-affinity condition, represented by these two external concentrations of K^+ . A fundamental distinction between these conditions is the active transport of K^+ in the HATS range and its passive transport in the LATS. In the latter situation, an electrochemically neutral distribution of K^+ across the membrane may result (Maathuis and Sanders 1993), and this, in conjunction with the potassium-dependent drop in membrane potential, could dramatically alter the distribution of K^+ across the plasma membrane.

Table 1 shows values for cytosolic $[K^+]$, obtained from a variety of plant systems and using several analytical methods. Ranges of values, where given, represent the response of the cell to K^+ availability; in many cases, this range is fairly narrow, reflecting the homeostasis of the K^+ pool. Interestingly, the use of widely differing laboratory techniques to determine cytosolic $[K^+]$, including tracer analysis (Pitman and Saddler 1967), longitudinal ion profiling (Jeschke and Stelter 1976), X-ray microanalysis (Flowers and Hajibagheri 2001), the K^+ -sensitive fluorescent dye PBFI (potassium-binding benzofuran isophthalate; Halperin and Lynch 2003), K^+ -specific microelectrodes (Walker et al. 1996) and cell fractionation (Speer and Kaiser 1991), nevertheless results in reasonably close agreement. This is in contrast to large differences in values obtained using a similar range of methods for other ions such as sodium (Carden et al. 2003, Kronzucker et al. 2006), ammonium (Britto et al. 2001, Lee and Ratcliffe 1991), and nitrate (Britto and Kronzucker 2003, Siddiqi and Glass 2002). The reasons for inconsistent agreement is unclear; the general agreement seen among methods in the case of potassium may be indicative of the homeostatic control of this pool, which may render it more resistant to physical disturbances that may be imposed during experimentation, for example with microelectrode impalement or with tracer elution protocols.

K^+ transport families

In the past decade, there has been substantial progress in the identification of genes encoding K^+ transporters in plants, as can be seen in the plethora of excellent recent reviews on the subject (Ashley et al. 2006, Cherel 2004, Gierth and Mäser 2007, Grabov 2007, Hedrich and Marten 2006, Lebaudy et al. 2007, Rodriguez-Navarro

Table 1. Cytosolic concentrations or activities of potassium, as determined by a variety of methods in several plant systems.

| Method | Plant material | [K ⁺] _{cytosol} (mM) | Reference |
|--|---|---|---------------------------------|
| Efflux analysis | Barley root | 92 | Pitman and Saddler (1967) |
| | | 236 | Bange (1979) |
| | | 316–320 | Behl and Jeschke (1982) |
| | | 127–184 | Memon et al. (1985) |
| | | 40–130 | Kronzucker et al. (2003) |
| | | 25–200 | Szczerba et al. (2006a) |
| Longitudinal ion profiling | Onion root | 184 | Macklon (1975) |
| | Maize root | 99–108 | Davis and Higinbotham (1976) |
| X-ray microanalysis | Barley root | 110 | Jeschke and Stelter (1976) |
| | Maize root | 129–162 | Hajibagheri et al. (1987, 1988) |
| Fluorescent dye | Barley protoplasts | 71–119 | Pitman et al. (1981) |
| | <i>Arabidopsis thaliana</i> root hairs | 65–70 | Lindberg (1995) |
| K ⁺ -selective microelectrode | <i>Acer pseudoplatanus</i> (suspension cells) | 55–60 | Halperin and Lynch (2003) |
| | <i>A. thaliana</i> root | 126 | Rona et al. (1982) |
| | Barley root | 83 | Maathuis and Sanders (1993) |
| | Barley root (± salt stress) | 45–83 | Walker et al. (1996) |
| Cell fractionation | Pea leaf | 39–63 | Carden et al. (2003) |
| | Spinach leaf | 53 | Speer and Kaiser (1991) |
| | | 147 | Speer and Kaiser (1991) |

and Rubio 2006, Shabala 2003, Véry and Sentenac 2003). This progress has been made despite a somewhat false start with the discovery of the first putative high-affinity K⁺ transporter HKT1 (Schachtman and Schroeder 1994), which is now thought to belong to a group of transporters, the HKT/Trk family, that are clearly implicated in the transport and internal recirculation of Na⁺ but appear to be of relatively minor significance to K⁺ transport. This revision is an important example of the problems associated with heterologous expression systems (Gierth and Mäser 2007, Haro et al. 2005, Lebaudy et al. 2007), which are used extensively in genetic screening projects. Potentially more promising is the development of plant-based expression systems for physiological studies on specific gene products (Bei and Luan 1998). Nevertheless, the discovery and characterization of HKT proteins have some relevance to K⁺ transport, in that, despite being implicated in high-affinity cation transport, their molecular architecture is similar to that of bona fide low-affinity K⁺ transporters (Shaker and KcsA-type channels). In both types of transporter, membrane-pore-membrane (MPM) motifs are found, with conserved residues within the pore loop area functioning as an ion selectivity filter (Mäser et al. 2002; see below).

At present, it appears clear that the high-affinity range of transport in plants is dominated by the activity of transporters from the KT/HAK/KUP family, which were identified based on sequence homologies to transporters from bacteria (Schleyer and Bakker 1993) and fungi (Bañuelos et al. 1995), as well as through mutant complementation studies. These transporters have 10 to

14 transmembrane regions (Gierth and Mäser 2007) and probably catalyze K⁺ transport by engaging a potassium/proton symport mechanism, as has been shown for the *Neurospora crassa* HAK1 gene product (Haro et al. 1999). In *Arabidopsis*, microarray analysis has shown that several members of this family are upregulated by K⁺ starvation, conforming to the phenotype of HATS upregulation; of these, AtHAK5 is particularly strongly upregulated and may be the most important contributor to high-affinity K⁺ uptake (Gierth et al. 2005). In addition to K⁺ uptake, there appear to be several other roles for transporters in this group, specifically in the distribution of auxin (Vicente-Agullo et al. 2004) and in cell expansion (Elumalai et al. 2002), two processes that, clearly, are intimately connected (Jones et al. 1998). The relationship between potassium transport and these other functions requires further study.

In addition to the major contribution to high-affinity K⁺ transport made by KT/HAK/KUP transporters and the minor possible involvement of HKT/Trk transporters, there is growing evidence that two large, related families of electroneutral cation-proton antiporters, CPA1 and CPA2, are also responsible for non-channel-mediated K⁺ transport (Brett et al. 2005, Gierth and Mäser 2007). So far, however, the functional attributions of these proteins have been largely for the transport of Na⁺, one prominent example being the tonoplast transporter NHX1 (a CPA1-type protein), which can confer salinity tolerance, most likely by enhancing the Na⁺ flux to the vacuole (Apse et al. 1999). Another example from the CPA1 family is the SOS1 protein, which also plays an important role in salt

tolerance by catalyzing Na^+ efflux from the cell (Shi et al. 2000). Nevertheless, NHX1 (and similar proton antiporters) has been shown to be responsible for vacuolar K^+ loading, as well as for Na^+ transport across the tonoplast, although it appears to have lower specificity for K^+ (Zhang and Blumwald 2001); the tandem-pore TPK1 channel may be more important for the trans-tonoplast movement of K^+ , particularly in its release from guard cell vacuoles (Gobert et al. 2007). At least two CHX proteins from the CPA2 family appear to be involved in K^+ homeostasis and pH regulation (Gierth and Mäser 2007), properties that are tied together mechanistically via the counter-exchange of K^+ and H^+ (Britto and Kronzucker 2005).

Potassium-selective channels have been identified at the molecular genetic level as multimers composed of α -subunits belonging to three major families: the Shaker, TPK (tandem-pore) and Kir (inward-rectifying)-like families (Lebaudy et al. 2007). Of these, the best characterized, and possibly the most functionally important, is the family of Shaker-type channels. The tetrameric channel is composed of four α -subunits arranged around a central pore (Zimmermann and Sentenac 1999). Each subunit consists of six hydrophobic, transmembrane segments, with a region of positively charged amino acids on the fourth segment acting as a voltage sensor, and a pore domain between the fifth and sixth segment containing the highly conserved GYGD (glycine-tyrosine-glycine-aspartate) motif, a key characteristic of potassium selectivity filters (Véry and Sentenac 2003).

In addition to the above-mentioned families of K^+ -specific channels, K^+ currents can also be conducted via non-selective cation channels (Demidchik and Maathuis 2007), a heterogeneous group that includes the Shaker-like, cyclic-nucleotide-gated channels (CNGs). Twenty CNGs have been found in *Arabidopsis* (Talke et al. 2003), one of which, the putative K^+ channel AtCNGC10, was recently found to profoundly influence growth and starch accumulation (Borsics et al. 2007). By contrast, only nine Shaker-type channel genes have been identified in *Arabidopsis*, but, because Shaker subunits can form heteromers, the structural–functional diversity of these proteins may be vast (Lebaudy et al. 2007). A vivid demonstration of this was seen in the case of trans-formation of *Xenopus* oocytes with five Shaker genes, the resulting gene products being shown to polymerize indiscriminately, suggesting the presence of a major source of K^+ channel diversity in plants (Dreyer et al. 1997, Xicluna et al. 2007).

Apart from their primary role in mediating potassium uptake from the soil, particularly under K^+ -replete conditions, potassium channels have a wide range of functions within the plant. One major role, mentioned

above, is the short-term maintenance of electrical potential gradients across cell membranes, in which K^+ channels operate in concert with electrogenic pumps (including the plasma membrane H^+ ATPase) and chloride transporters (Gradmann and Hoffstadt 1998). Because K^+ channels are electrogenic in mechanism, all can contribute to the polarization state of the cell, but the voltage-gated channels have regulatory components that render them more specifically suited to this function. These channels are generally of the Shaker type, for both inward- and outward-rectifying fluxes (Lebaudy et al. 2007). Shaker channels have also been shown to function in the xylem loading of K^+ (SKOR channels; Gaymard et al. 1998), as well as in phloem loading and unloading (AKT2/3; Deeken et al. 2000, 2002, Lacombe et al. 2000).

To conclude this review on cellular K^+ transport in plants, we wish to highlight three very recent discoveries in the literature, which illustrate the utility of molecular biology to unravel physiological phenomena of broad interest to plant biologists. In one example (Yu et al. 2006), the activity of the AKT2 channel homologue SPICK2 was shown to be a likely participant in the turgor-related movements of *Samanea* motor cells, acting differentially in extensor and flexor halves of the leaf pulvinus, depending on the phosphorylation state of the transporter. In another example (Sano et al. 2007), K^+ -dependent cell turgor influencing different phases of the cell cycle appears to be reciprocally controlled by the inward-rectifying K^+ channel NKT1 during the G1 phase, and the outward-rectifying K^+ channel NTORK1 during the S phase. Finally, Gazzarrini et al. (2006) showed that the *Chlorella* virus MT325 encodes both water channels and K^+ channels that appear to work synergistically in the host, providing both an osmotic driving force (cellular K^+ accumulation) and a pathway for water flow into the cell. These examples indicate the ongoing progress in demonstrating critical functions for potassium pools and transporters in plant systems.

Acknowledgement – We wish to thank the Natural Sciences and Engineering Council of Canada (NSERC) for funding this work.

References

- Armengaud P, Breitling R, Amtmann A (2004) The potassium-dependent transcriptome of *Arabidopsis* reveals a prominent role of jasmonic acid in nutrient signaling. *Plant Physiol* 136: 2556–2576
- Adams F (1971) Soil solution. In: Carson EW (ed) *The Plant Root and Its Environment*. University Press of Virginia, Charlottesville, VA, pp 441–481

- Apse MP, Aharon GS, Snedden WA, Blumwald E (1999) Salt tolerance conferred by overexpression of a vacuolar Na⁺/H⁺ antiport in *Arabidopsis*. *Science* 285: 1256–1258
- Ashley MK, Grant M, Grabov A (2006) Plant responses to potassium deficiencies: a role for potassium transport proteins. *J Exp Bot* 57: 425–436
- Bange G (1979) Comparison of compartmental analysis for Rb⁺ and Na⁺ in low-salt and high-salt barley roots. *Physiol Plant* 46: 179–183
- Bañuelos MA, Klein RD, Alexander-Bowman SJ, Rodríguez-Navarro A (1995) A potassium transporter of the yeast *Schwanniomyces occidentalis* homologous to the Kup system of *Escherichia coli* has a high concentrative capacity. *EMBO J* 14: 3021–3027
- Behl R, Jeschke WD (1982) Potassium fluxes in excised barley roots. *J Exp Bot* 33: 584–600
- Bei Q, Luan S (1998) Functional expression and characterization of a plant K⁺ channel gene in a plant cell model. *Plant J* 13: 857–865.
- Ben-Zioni A, Vaadia Y, Lips SH (1971) Nitrate uptake by roots as regulated by nitrate reduction products of the shoot. *Physiol Plant* 24: 288–290
- Borsics T, Webb D, Andeme-Ondzighi C, Staehelin LA, Christopher DA (2007) The cyclic nucleotide-gated calmodulin-binding channel AtCNGC10 localizes to the plasma membrane and influences numerous growth responses and starch accumulation in *Arabidopsis thaliana*. *Planta* 225: 563–573
- Borstlap AC (1981) Invalidity of the multiphasic concept of ion absorption in plants. *Plant Cell Environ* 4: 189–195
- Brett CL, Donowitz M, Rao R (2005) Evolutionary origins of eukaryotic sodium/proton exchangers. *Am J Physiol Cell Physiol* 288: C223–C239
- Britto DT, Kronzucker HJ (2001) Can unidirectional influx be measured in higher plants? A mathematical approach using parameters from efflux analysis. *New Phytol* 150: 37–47
- Britto DT, Kronzucker HJ (2003) The case for cytosolic NO₃⁻ heterostasis: a critique of a recently proposed model. *Plant Cell Environ* 26: 183–188
- Britto DT, Kronzucker HJ (2005) Nitrogen acquisition, PEP carboxylase, and cellular pH homeostasis: new views on old paradigms. *Plant Cell Environ* 28: 1396–1409
- Britto DT, Kronzucker HJ (2006) Futile cycling at the plasma membrane: a hallmark of low-affinity nutrient transport. *Trends Plant Sci* 11: 529–534
- Britto DT, Siddiqi MY, Glass ADM, Kronzucker HJ (2001) Futile transmembrane NH₄⁺ cycling: a cellular hypothesis to explain ammonium toxicity in plants. *Proc Natl Acad Sci USA* 98: 4255–4258
- Britto DT, Szczerba MW, Kronzucker HJ (2006) A new, non-perturbing, sampling procedure in tracer exchange measurements. *J Exp Bot* 6: 1309–1314
- Brüggemann L, Dietrich P, Becker D, Dreyer I, Palme K, Hedrich R (1999) Channel-mediated high-affinity K⁺ uptake into guard cells from *Arabidopsis*. *Proc Natl Acad Sci USA* 96: 3298–3302
- Busch W, Saier MH (2002) The transporter classification (TC) system. *Crit Rev Biochem Mol Biol* 37:287–337
- Cakmak I (2005) The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *J Plant Nutr Soil Sci* 168: 521–530
- Cakmak I, Hengeler C, Marschner H (1994) Changes in phloem export of sucrose in leaves in response to phosphorus, potassium and magnesium deficiency in bean plants. *J Exp Bot* 45: 1251–1257
- Carden DE, Walker DJ, Flowers TJ, Miller AJ (2003) Single-cell measurements of the contributions of cytosolic Na⁺ and K⁺ to salt tolerance. *Plant Physiol* 131: 676–683
- Cerezo M, Tillard P, Gojon A, Primo-Millo E, Garcia-Agustin P (2001) Characterization and regulation of ammonium transport systems in *Citrus* plants. *Planta* 214: 97–105
- Cheeseman JM, Hanson JB (1979) Mathematical analysis of the dependence of cell potential on external potassium in corn roots *Plant Physiol* 63: 1–4
- Cheeseman JM, Hanson JB (1980) Does active K⁺ influx to roots occur? *Plant Sci Lett* 18: 81–84
- Cheeseman JM, Lafayette PR, Gronewald JW, Hanson JB (1980) Effect of ATPase inhibitors on cell potential and K⁺ influx in corn roots. *Plant Physiol* 65: 1139–1145
- Cherel I (2004) Regulation of K⁺ channel activities in plants: from physiological to molecular aspects. *J Exp Bot* 55: 337–351
- Clarkson DT, Hanson JB (1980) The mineral nutrition of higher-plants. *Annu Rev Plant Physiol Plant Mol Biol* 31: 239–298
- Cline JF, Hungate FP (1960) Accumulation of potassium, cesium¹³⁷ and rubidium⁸⁶ in bean plants grown in nutrient solutions. *Plant Physiol* 35: 826–829.
- Davis RF, Higinbotham N (1976) Electrochemical gradients and K⁺ and Cl⁻ fluxes in excised corn roots. *Plant Physiol* 57: 129–136
- de Agazio M, Giardina MC, Federico R (1983) ⁸⁶Rb influx in maize roots at different external concentrations of RbCl or KCl. *Plant Cell Physiol* 24: 453–455
- Deeken R, Sanders C, Ache P, Hedrich R (2000) Developmental and light-dependent regulation of a phloem-localised K⁺ channel of *Arabidopsis thaliana*. *Plant J* 23: 285–290
- Deeken R, Geiger D, Fromm J, Koroleva O, Ache P, Langenfeld-Heyser R, Sauer N, May ST, Hedrich R (2002) Loss of the AKT2/3 potassium channel affects sugar loading into the phloem of *Arabidopsis*. *Planta* 216: 334–344
- Demidchik V, Maathuis JM (2007) Physiological roles of nonselective cation channels in plants: from salt stress to signaling and development. *New Phytol* 175: 387–404
- Dietrich P, Sanders D, Hedrich R (2001) The role of ion channels in light-dependent stomatal opening. *J Exp Bot* 52: 1959–1967

- Dolan L, Davies J (2004) Cell expansion in roots. *Curr Opin Plant Biol* 7: 33–39
- Doyle DA, Cabral JM, Pfuetzner RA, Kuo AL, Gulbis JM, Cohen SL, Chait BT, MacKinnon R (1998) The structure of the potassium channel: molecular basis of K⁺ conduction and selectivity. *Science* 280: 69–77
- Dreyer I, Antunes S, Hoshi T, MullerRober B, Palme K, Pongs O, Reintanz B, Hedrich R (1997) Plant K⁺ channel alpha-subunits assemble indiscriminately. *Biophys J* 72: 2143–2150
- Elumalai RP, Nagpal P, Reed JW (2002) A mutation in the *Arabidopsis* KT2/KUP2 potassium transporter gene affects shoot cell expansion. *Plant Cell* 14: 119–131
- Epstein E, Bloom AJ (2005) *Mineral Nutrition of Plants: Principles and Perspectives*, 2nd Edn. Sinauer Associates Inc., Sunderland MA
- Epstein E, Rains DW, Elzam OE (1963) Resolution of dual mechanisms of potassium absorption by barley roots. *Proc Natl Acad Sci USA* 49: 684–692
- Etherton B, Higinbotham N (1960) Transmembrane potential measurements of cells of higher plants as related to salt uptake. *Science* 131: 409–410
- Fisher JD, Hausen D, Hodges TK (1970) Correlation between ion fluxes and ion stimulated adenosine triphosphatase activity of plant roots. *Plant Physiol* 46: 812–814
- Flowers TJ, Hajibagheri MA (2001) Salinity tolerance in *Hordeum vulgare*: ion concentrations in root cells of cultivars differing in salt tolerance. *Plant Soil* 231: 1–9
- Fu HH, Luan S (1998) AtKUP1: a dual-affinity K⁺ transporter from *Arabidopsis*. *Plant Cell* 10: 63–73
- Gassmann W, Schroeder JI (1994) Inward-rectifying K⁺ channels in root hairs of wheat – a mechanism for aluminum-sensitive low-affinity K⁺ uptake and membrane potential control. *Plant Physiol* 105: 1399–1408
- Gaymard F, Pilot G, Lacombe B, Bouchez D, Bruneau D, Boucherez J, Michaux-Ferriere N, Thibaud JB, Sentenac H (1998) Identification and disruption of a plant shaker-like outward channel involved in K⁺ release into the xylem sap. *Cell* 94: 647–655
- Gazzarrini S, Kang M, Epimashko S, Van Etten JL, Dainty J, Thiel G, Moroni A (2006) *Chlorella* virus MT325 encodes water and potassium channels that interact synergistically. *Proc Natl Acad Sci USA* 103: 5355–5360
- Gierth M, Mäser P (2007) Potassium transporters in plants – involvement in K⁺ acquisition, redistribution and homeostasis. *FEBS Lett* 581: 2348–2356
- Gierth M, Mäser P, Schroeder JI (2005) The potassium transporter AtHAK5 functions in K⁺ deprivation-induced high-affinity K⁺ uptake and AKT1 K⁺ channel contribution to K⁺ uptake kinetics in *Arabidopsis* roots. *Plant Physiol* 137: 1105–1114
- Glass ADM (1978) The regulation of K⁺ influx into intact roots of barley (*Hordeum vulgare* (L) cv. Conquest) by internal K⁺. *Can J Bot* 56: 1759–1764
- Glass ADM, Dunlop J (1978) Influence of potassium content on kinetics of potassium influx into excised ryegrass and barley roots. *Planta* 141: 117–119
- Goerttler A, Isayenkova S, Voelker C, Czempinski K, Maathuis FJM (2007) The two-pore channel *TPK1* gene encodes the vacuolar K⁺ conductance and plays a role in K⁺ homeostasis. *Proc Natl Acad Sci USA* 104: 10726–10731
- Grabov A (2007) Plant KT/KUP/HAK potassium transporters: single family – multiple functions. *Ann Bot* 99: 1035–1041
- Gradmann D, Hoffstadt J (1998) Electrocoupling of ion transporters in plants: interaction with internal ion concentrations. *J Membr Biol* 166: 51–59
- Hajibagheri MA, Harvey DMR, Flowers TJ (1987) Quantitative ion distribution within root-cells of salt-sensitive and salt-tolerant maize varieties. *New Phytol* 105: 367–379
- Hajibagheri MA, Flowers TJ, Collins JC, Yeo AR (1988) A comparison of the methods of X-ray microanalysis, compartmental analysis and longitudinal ion profiles to estimate cytoplasmic ion concentrations in 2 maize varieties. *J Exp Bot* 39: 279–290
- Halperin SJ, Lynch JP (2003) Effects of salinity on cytosolic Na⁺ and K⁺ in root hairs of *Arabidopsis thaliana*: in vivo measurements using the fluorescent dyes SBFI and PBF1. *J Exp Bot* 54: 2035–2043
- Haro R, Bañuelos MA, Senn MAE, Barrero-Gil J, Rodriguez-Navarro A (2005) HKT1 mediates sodium uniprot in roots. Pitfalls in the expression of HKT1 in yeast. *Plant Physiol* 139: 1495–1506
- Haro R, Sainz L, Rubio F, Rodriguez-Navarro A (1999) Cloning of two genes encoding potassium transporters in *Neurospora crassa* and expression of the corresponding cDNAs in *Saccharomyces cerevisiae*. *Mol Microbiol* 31: 511–520
- Hayes DE, Smith FA, Walker NA (2001) High-affinity potassium transport into wheat roots involves sodium – a role for HKT1? *Aust J Plant Physiol* 28: 643–652
- Hedrich R, Marten I (2006) 30-year progress of membrane transport in plants. *Planta* 224: 725–739
- Hertel B, Horvath F, Wodala B, Hurst A, Moroni A, Thiel G (2005) KAT1 inactivates at sub-millimolar concentrations of external potassium. *J Exp Bot* 56: 3103–3110
- Hirsch RE, Lewis BD, Spalding EP, Sussman MR (1998) A role for the AKT1 potassium channel in plant nutrition. *Science* 280: 918–921
- Hoth S, Hedrich R (1999) Distinct molecular bases for pH sensitivity of the guard cell K⁺ channels KST1 and KAT1. *J Biol Chem* 274: 11599–11603
- Humble GD, Hsiao TC (1969) Specific requirement of potassium for light-activated opening of stomata in epidermal strips. *Plant Physiol* 44: 230–234
- Jacoby B (1975) Light sensitivity of ²²Na, ⁸⁶Rb, and K⁴² absorption by different tissues of bean-leaves. *Plant Physiol* 55: 978–981

- Jacoby B, Nissen P (1977) Potassium and rubidium interaction in their absorption by bean leaf slices. *Physiol Plant* 40: 42–44
- Jeschke WD, Stelzer W (1976) Measurement of longitudinal ion profiles in single roots of *Hordeum* and *Atriplex* by use of flameless atomic-absorption spectroscopy. *Planta* 128: 107–112
- Johansson I, Wulfetange K, Poree F, Michard E, Gajdanowicz P, Lacombe B, Sentenac H, Thibaud JB, Mueller-Roeber B, Blatt MR, Dreyer I (2006) External K⁺ modulates the activity of the *Arabidopsis* potassium channel SKOR via an unusual mechanism. *Plant J* 46: 269–281
- Jones AM, Im KH, Savka MA, Wu MJ, DeWitt NG, Shillito R, Binns AN (1998) Auxin-dependent cell expansion mediated by overexpressed auxin-binding protein 1. *Science* 282: 1114–1117
- Kannan S, Ramani S (1973) Discrimination between Rb and K ions in their absorption by rice roots. *Int J Appl Rad Iso* 24: 431–432
- Keifer DW, Spanswick RM (1978) Activity of electrogenic pump in *Chara corallina* as inferred from measurements of membrane potential, conductance, and potassium permeability. *Plant Physiol* 62: 653–661
- Kim EJ, Kwak JM, Uozumi N, Schroeder JI (1998) *AtKUP1*: an *Arabidopsis* gene encoding high-affinity potassium transport activity. *Plant Cell* 10: 51–62
- Kochian LV, Lucas WJ (1982) Potassium-transport in corn roots. I. Resolution of kinetics into a saturable and linear component. *Plant Physiol* 70: 1723–1731
- Kochian LV, Lucas WJ (1983) Potassium-transport in corn roots. II. The significance of the root periphery. *Plant Physiol* 73: 208–215
- Kochian LV, Jiao XZ, Lucas WJ (1985) Potassium-transport in corn roots. IV. Characterization of the linear component. *Plant Physiol* 79: 771–776
- Kochian LV, Shaff JE, Lucas WJ (1989) High-affinity K⁺ uptake in maize roots – a lack of coupling with H⁺ efflux. *Plant Physiol* 91: 1202–1211
- Kronzucker HJ, Szczerba MW, Britto DT (2003) Cytosolic potassium homeostasis revisited: ⁴²K-tracer analysis in *Hordeum vulgare* L. reveals set-point variations in [K⁺]. *Planta* 217: 540–546
- Kronzucker HJ, Szczerba MW, Moazami-Goudarzi M, Britto DT (2006) The cytosolic Na⁺: K⁺ ratio does not explain salinity-induced growth impairment in barley: a dual tracer study using ⁴²K⁺ and ²⁴Na⁺. *Plant Cell Environ* 29: 2228–2237
- Lacombe B, Pilot G, Michard E, Gaymard F, Sentenac H, Thibaud JB (2000) A shaker-like K⁺ channel with weak rectification is expressed in both source and sink phloem tissues of *Arabidopsis*. *Plant Cell* 12: 837–851
- Läuchli A, Epstein E (1970) Transport of potassium and rubidium in plant roots – significance of calcium. *Plant Physiol* 45: 639–641
- Lebaudy A, Very AA, Sentenac H (2007) K⁺ channel activity in plants: genes, regulations and functions. *FEBS Lett* 581: 2357–2366
- Lee RB, Ratcliffe RG (1991) Observations on the subcellular distribution of the ammonium ion in maize root-tissue using *in-vivo* ¹⁴N-nuclear magnetic resonance spectroscopy. *Planta* 183: 359–367
- Leigh RA (2001) Potassium homeostasis and membrane transport. *J Plant Nutr Soil Sci* 164: 193–198
- Leigh RA, Wyn Jones RG (1984) A hypothesis relating critical potassium concentrations for growth to the distribution and functions of this ion in the plant cell. *New Phytol* 97: 1–13
- Li LG, Kim BG, Cheong YH (2006) A Ca²⁺ signaling pathway regulates a K⁺ channel for low-K response in *Arabidopsis*. *Proc Natl Acad Sci USA* 103: 12625–12630
- Lindberg S (1995) *In situ* determination of intracellular concentrations of K⁺ in barley (*Hordeum vulgare* L. cv. Kara) using the K⁺-binding fluorescent dye benzofuran isophthalate. *Planta* 195: 525–529
- Lindberg S, Strid H (1997) Aluminium induces rapid changes in cytosolic pH and free calcium and potassium concentrations in root protoplasts of wheat (*Triticum aestivum*). *Physiol Plant* 99: 405–414
- Liu KH, Tsay YF (2003) Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *EMBO J* 22: 1005–1013
- Liu KH, Huang CY, Tsay YF (1999) CHL1 is a dual-affinity nitrate transporter of *Arabidopsis* involved in multiple phases of nitrate uptake. *Plant Cell* 11: 865–874
- Liu K, Li LG, Luan S (2006) Intracellular K⁺ sensing of SKOR, a Shaker-type K⁺ channel from *Arabidopsis*. *Plant J* 46: 260–268
- Lynch JP (2007) Roots of the second green revolution. *Aust J Bot* 55: 493–512
- Maas EV, Leggett JE (1968) Uptake of ⁸⁶Rb and K by excised maize roots. *Plant Physiol* 43: 2054–2056
- Maathuis FJM, Sanders D (1993) Energization of potassium uptake in *Arabidopsis thaliana*. *Planta* 191: 302–307
- Maathuis FJM, Sanders D (1994) Mechanism of high-affinity potassium uptake in roots of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 91: 9272–9276
- Maathuis FJM, Sanders D (1995) Contrasting roles in ion-transport of 2 K⁺-channel types in root cells of *Arabidopsis thaliana*. *Planta* 197: 456–464
- Maathuis FJM, Verlin D, Smith FA, Sanders D, Fernandez JA, Walker NA (1996) The physiological relevance of Na⁺-coupled K⁺-transport. *Plant Physiol* 112: 1609–1616
- Maathuis FJM, Sanders D, Gradmann D (1997) Kinetics of high-affinity K⁺ uptake in plants, derived from K⁺-induced changes in current-voltage relationships – a modelling approach to the analysis of carrier-mediated transport. *Planta* 203: 229–236

- Macklon AES (1975) Cortical cell fluxes and transport to stele in excised root segments of *Allium cepa* L. 1. Potassium, sodium and chloride. *Planta* 122: 109–130
- MacRobbie EAC, Dainty J (1958) Ion transport in *Nitellopsis obtusa*. *J Gen Physiol* 42: 335–353
- Malhotra B, Glass ADM (1995) Potassium fluxes in *Chlamydomonas reinhardtii*. I. Kinetics and electrical potentials. *Plant Physiol* 108: 1527–1536
- Marschner H, Schimanski C (1968) Differential uptake of potassium and rubidium by barley. *Naturwissenschaften* 55: 499
- Marschner H, Kuiper PJC, Kylin A (1981) Genotypic differences in the response of sugar-beet plants to replacement of potassium by sodium. *Physiol Plant* 51: 239–244
- Mäser P, Hosoo Y, Goshima S, Horie T, Eckelman B, Yamada K, Yoshida K, Bakker EP, Shinmyo A, Oiki S, Schroeder JI, Uozumi N (2002) Glycine residues in potassium channel-like selectivity filters determine potassium selectivity in four-loop-per-subunit HKT transporters from plants. *Proc Natl Acad Sci USA* 99: 6428–6433
- Memon AR, Saccomani M, Glass ADM (1985) Efficiency of potassium utilization by barley varieties - the role of subcellular compartmentation. *J Exp Bot* 36: 1860–1876
- Michard E, Lacombe B, Poree F, Mueller-Roeber B, Sentenac H, Thibaud JB, Dreyer I (2005) A unique voltage sensor sensitizes the potassium channel AKT2 to phosphoregulation. *J Gen Physiol* 126: 605–617
- Moran N (2007) Osmoregulation of leaf motor cells. *FEBS Lett* 581: 2337–2347
- Mouline K, Very AA, Gaymard F, Boucherez J, Pilot G, Devic M, Bouchez D, Thibaud JB, Sentenac H (2002) Pollen tube development and competitive ability are impaired by disruption of a Shaker K⁺ channel in *Arabidopsis*. *Genes Dev* 16: 339–350
- Nieves-Cordones M, Martinez-Cordero MA, Martinez V, Rubio F (2007) An NH₄⁺-sensitive component dominates high-affinity K⁺ uptake in tomato plants. *Plant Sci* 172: 273–280
- Nissen P (1980) Multiphasic uptake of potassium by barley roots of low and high potassium content – separate sites for uptake and transitions. *Physiol Plant* 48: 193–200
- Nocito FF, Sacchi GA, Cocucci M (2002) Membrane depolarization induces K⁺ efflux from subapical maize root segments. *New Phytol* 154: 45–51
- Palmgren MG (2001) Plant plasma membrane H⁺-ATPases: powerhouses for nutrient uptake. *Annu Rev Plant Physiol Plant Mol Biol* 52: 817–845
- Pardo JM, Quintero FJ (2002) Plants and sodium ions: keeping company with the enemy. *Genome Biol* 3: 10171–10174
- Pardo JM, Cubero B, Leidi EO, Quintero FJ (2006) Alkali cation exchangers: roles in cellular homeostasis and stress tolerance. *J Exp Bot* 57: 1181–1199
- Pettersson S, Kasimir-Klemedtsson A (1990) Influx and efflux of K⁺ in sunflower roots after transfer between solutions with different K⁺ concentrations. *Physiol Plant* 79: 686–692
- Philippart K, Fuchs I, Luthen H, Hoth S, Bauer CS, Haga K, Thiel G, Ljung K, Sandberg G, Bottger M, Becker D, Hedrich R (1999) Auxin-induced K⁺ channel expression represents an essential step in coleoptile growth and gravitropism. *Proc Natl Acad Sci USA* 96: 12186–12191
- Pitman MG, Saddler HDW (1967) Active sodium and potassium transport in cells of barley roots. *Proc Natl Acad Sci USA* 57: 44–49
- Pitman MG, Lauchli A, Stelzer R (1981) Ion distribution in roots of barley seedlings measured by electron-probe X-ray microanalysis. *Plant Physiol* 68: 673–679
- Polley LD, Hopkins JW (1979) Rubidium (potassium) uptake by *Arabidopsis*. *Plant Physiol* 64: 374–378
- Polley LD, Doctor DD (1985) Potassium transport in *Chlamydomonas reinhardtii* – isolation and characterization of transport deficient mutant strains. *Planta* 163: 208–213
- Rains DW, Epstein E (1967) Sodium absorption by barley roots: its mediation by mechanism 2 of alkali cation transport. *Plant Physiol* 42: 319–323
- Rawat SR, Silim SN, Kronzucker HJ, Siddiqi MY, Glass ADM (1999) *AtAMT1* gene expression and NH₄⁺ uptake in roots of *Arabidopsis thaliana*: evidence for regulation by root glutamine levels. *Plant J* 19: 143–152
- Reisenauer HM (1966) Mineral nutrients in soil solution. In: Altman PL, Ditter DS (eds) *Environmental Biology*. Federation of American Societies for Experimental Biology, Bethesda, MD, pp 507–508
- Rhoads DB, Woo A, Epstein W (1977) Discrimination between Rb⁺ and K⁺ by *Escherichia coli*. *Biochim Biophys Acta* 469: 45–51
- Rodriguez-Navarro A, Rubio F (2006) High-affinity potassium and sodium transport systems in plants. *J Exp Bot* 57: 1149–1160
- Rona JP, Cornel D, Grignon C, Heller R (1982) The electrical potential difference across the tonoplast of *Acer pseudoplatanus* cells. *Physiol Veg* 20: 459–463
- Rubio F, Gassmann W, Schroeder JI (1995) Sodium-driven potassium uptake by the plant potassium transporter HKT1 and mutations conferring salt tolerance. *Science* 270: 1660–1663
- Sano T, Becker D, Ivashikina N, Wegner LH, Zimmermann U, Roelfsema MR, Nagata T, Hedrich R (2007) Plant cells must pass a K⁺ threshold to re-enter the cell cycle. *Plant J* 50: 401–413
- Santa-Maria GE, Danna CH, Czibener C (2000) High-affinity potassium transport in barley roots. Ammonium-sensitive and -insensitive pathways. *Plant Physiol* 123: 297–306
- Schachtman DP, Schroeder JI (1994) Structure and transport mechanism of a high-affinity potassium uptake transporter from higher-plants. *Nature* 370: 655–658

- Schachtman DP, Schroeder JI, Lucas WJ, Anderson JA, Gaber RF (1992) Expression of an inward-rectifying potassium channel by the *Arabidopsis* *KAT1* cDNA. *Science* 258: 1654–1658
- Schleyer M, Bakker EP (1993) Nucleotide-sequence and 3'-end deletion studies indicate that the K⁺-uptake protein Kup from *Escherichia coli* is composed of a hydrophobic core linked to a large and partially essential hydrophilic C terminus. *J Bacteriol* 175: 6925–6931
- Shabala S (2003) Regulation of potassium transport in leaves: from molecular to tissue level. *Ann Bot* 92: 627–634
- Shabala S, Babourina O, Newman I (2000) Ion-specific mechanisms of osmoregulation in bean mesophyll cells. *J Exp Bot* 51: 1243–1253
- Shi HZ, Ishitani M, Kim CS, Zhu JK (2000) The *Arabidopsis thaliana* salt tolerance gene *SOS1* encodes a putative Na⁺/H⁺ antiporter. *Proc Natl Acad Sci USA* 97: 6896–6901
- Shin H, Shin HS, Dewbre GR, Harrison MJ (2004) Phosphate transport in *Arabidopsis*: Pht1;1 and Pht1;4 play a major role in phosphate acquisition from both low- and high-phosphate environments. *Plant J* 39: 629–642
- Siddiqi MY, Glass ADM (2002) An evaluation of the evidence for, and implications of, cytoplasmic nitrate homeostasis. *Plant Cell Environ* 25: 1211–1217
- Spalding EP, Hirsch RE, Lewis DR, Qi Z, Sussman MR, Lewis BD (1999) Potassium uptake supporting plant growth in the absence of AKT1 channel activity – inhibition by ammonium and stimulation by sodium. *J Gen Physiol* 113: 909–918
- Speer M, Kaiser WM (1991) Ion relations of symplastic and apoplastic space in leaves from *Spinacia oleracea* L. and *Pisum sativum* L. under salinity. *Plant Physiol* 97: 990–997
- Subbarao GV, Wheeler RM, Stutte GW, Levine LH (1999) How far can sodium substitute for potassium in red beet? *J Plant Nutr* 22: 1745–1761
- Suelter CH (1970) Enzymes activated by monovalent cations. *Science* 168: 789–795
- Szczerba MW, Britto DT, Kronzucker HJ (2006a) Rapid, futile K⁺ cycling and pool-size dynamics define low-affinity potassium transport in barley. *Plant Physiol* 141: 1494–1507
- Szczerba MW, Britto DT, Kronzucker HJ (2006b) The face value of ion fluxes: the challenge of determining influx in the low-affinity transport range. *J Exp Bot* 57: 3293–3300
- Szczerba MW, Britto DT, Balkos KD, Kronzucker HJ (2008) Alleviation of rapid, futile ammonium cycling at the plasma membrane by potassium reveals K⁺-sensitive and -insensitive components of NH₄⁺ transport. *J Exp Bot*, doi:10.1093/jxb/erm309
- Talke IN, Blaudez D, Maathuis FJM, Sanders D (2003) CNGCs: prime targets of plant cyclic nucleotide signalling? *Trends Plant Sci* 8: 286–293
- Tester M (1990) Plant ion channels – whole-cell and single-channel studies. *New Phytol* 114: 305–340
- Véry AA, Sentenac H (2003) Molecular mechanisms and regulation of K⁺ transport in higher plants. *Annu Rev Plant Biol* 54: 575–603
- Vicente-Agullo F, Rigas S, Desbrosses G, Dolan L, Hatzopoulos P, Grabov A (2004) Potassium carrier TRH1 is required for auxin transport in *Arabidopsis* roots. *Plant J* 40: 523–535
- Walker DJ, Leigh RA, Miller AJ (1996) Potassium homeostasis in vacuolate plant cells. *Proc Natl Acad Sci USA* 93: 10510–10514
- Wang TB, Gassmann W, Rubio F, Schroeder JI, Glass ADM (1998) Rapid up-regulation of *HKT1*, a high-affinity potassium transporter gene, in roots of barley and wheat following withdrawal of potassium. *Plant Physiol* 118: 651–659
- Wang S-M, Zhang J-L, Flowers TJ (2007) Low-affinity Na⁺ uptake in the halophyte *Suaeda maritima*. *Plant Physiol* 145: 559–571
- West KR, Pitman MG (1967) Rubidium as a tracer for potassium in the marine algae *Ulva lactuca* L. and *Chaetomorpha darwinii* (Hooker) Kuetzing. *Nature* 214: 1262–1263
- Xicluna J, Lacombe B, Dreyer I, Alcon C, Jeanguenin L, Sentenac H, Thibaud JB, Cherel I (2007) Increased functional diversity of plant K⁺ channels by preferential heteromerization of the Shaker-like subunits AKT2 and KAT2. *J Biol Chem* 282: 486–494
- Yu L, Becker D, Levi H, Moshelion M, Hedrich R, Lotan I, Moran A, Pick U, Naveh L, Libal Y, Moran N (2006) Phosphorylation of SPICK2, an AKT2 channel homologue from *Samanea* motor cells. *J Exp Bot* 57: 3583–3594
- Zhang HX, Blumwald E (2001) Transgenic salt-tolerant tomato plants accumulate salt in foliage but not in fruit. *Nat Biotech* 19: 765–768
- Zimmermann S, Sentenac H (1999) Plant ion channels: from molecular structures to physiological functions. *Curr Opin Plant Biol* 2: 477–482